

# Semi-reflexive polytopes

Tiago Royer

## Abstract

The Ehrhart function  $L_P(t)$  of a polytope  $P$  is usually defined only for integer dilation arguments  $t$ . By allowing arbitrary real numbers as arguments we may also detect integer points entering (or leaving) the polytope in fractional dilations of  $P$ , thus giving more information about the polytope. Nevertheless, there are some polytopes that only gain new integer points for integer values of  $t$ ; that is, these polytopes satisfy  $L_P(t) = L_P(\lfloor t \rfloor)$ . We call those polytopes *semi-reflexive*. In this paper, we give a characterization of these polytopes in terms of their hyperplane description, and we use this characterization to show that a polytope is reflexive if and only if both it and its dual are semi-reflexive.

## 1 Introduction

Given a polytope  $P \subseteq \mathbb{R}^d$ , the classical Ehrhart lattice point enumerator  $L_P(t)$  is defined as

$$L_P(t) = \#(tP \cap \mathbb{Z}^d), \quad \text{integer } t > 0.$$

Here,  $\#(A)$  is the number of elements in  $A$  and  $tP = \{tx \mid x \in P\}$  is the dilation of  $P$  by  $t$ . The above definition may be extended to allow arbitrary real numbers as dilation parameters; we will assume this extension in this paper. Moreover, we will agree that  $L_P(0) = 1$ .

To minimize confusion, we will denote real dilation parameters with the letter  $s$ , so that  $L_P(t)$  denotes the classical Ehrhart function and  $L_P(s)$  denotes the extension considered in this paper. So, for example,  $L_P(t)$  is just the restriction of  $L_P(s)$  to integer arguments.

Every polytope  $P \subset \mathbb{R}^d$  may be written as

$$P = \bigcap_{i=1}^n \{x \in \mathbb{R}^d \mid \langle a_i, x \rangle \leq b_i\}, \quad (1)$$

where each  $a_i$  is a vector in  $\mathbb{R}^d$  and each  $b_i$  is a real number.

We define a polytope  $P$  to be *semi-reflexive* if  $P$  is rational and satisfies  $L_P(s) = L_P(\lfloor s \rfloor)$  for all  $s \geq 0$ . We have the following characterization.

**Theorem 1.** *Let  $P$  be a rational polytope. Then  $P$  is a semi-reflexive polytope if and only if  $P$  may be written as in (1), with all the  $a_i$  being integers and all  $b_i$  being either 0 or 1.*

Section 2 contains some examples of semi-reflexive polytopes. The characterization above is proven in Section 3. A similar proof yields another, similar, characterization of semi-reflexive polytopes in terms of their relative interiors; this is discussed in Section 4. Finally, in Section 5 we show how semi-reflexive polytopes relate with reflexive polytopes.

## 2 Examples of semi-reflexive polytopes

Here, we will use the “if” part of the characterization to provide some examples of semi-reflexive polytopes.

- The unit cube  $P = [0, 1]^d$ .  $P$  may be represented by the inequalities  $x_i \geq 0$  and  $x_i \leq 1$ , for all  $i$ .
- The standard simplex, which is defined by the inequalities  $x_i \geq 0$  for all  $i$ , and  $x_1 + \dots + x_d \leq 1$ .
- The cross-polytope. This polytope is defined by  $|x_1| + \dots + |x_d| \leq 1$ . Each of the  $2^d$  vectors  $(\alpha_1, \dots, \alpha_d) \in \{-1, 1\}^d$  gives a bounding inequality of the form  $\alpha_1 x_1 + \dots + \alpha_d x_d \leq 1$ , all of which satisfy the characterization.
- Order polytopes [3]. Let  $\prec$  be a partial order over the set  $\{1, \dots, d\}$ . The order polytope for the partial order  $\prec$  is the set of points  $(x_1, \dots, x_d) \in \mathbb{R}^d$  which satisfy  $0 \leq x_i \leq 1$  for all  $i$ , and  $x_i \leq x_j$  whenever  $i \prec j$ .
- Chain polytopes [3]. Let  $\prec$  be a partial order over the set  $\{1, \dots, d\}$ . The chain polytope for the partial order  $\prec$  is the set of points  $(x_1, \dots, x_d) \in \mathbb{R}^d$  which satisfy  $0 \leq x_i$  for all  $i$ , and  $x_{i_1} + \dots + x_{i_k} \leq 1$  for all chains  $i_1 \prec i_2 \prec \dots \prec i_k$ .
- Quasi-metric polytopes [2]. Let  $G$  be a graph such that every vertex has degree 1 or 3. Identify its edge set with  $\{1, \dots, d\}$ . The quasi-metric polytope  $\mathcal{P}_G$  is defined to be the set of points  $(x_1, \dots, x_d) \in \mathbb{R}^d$  satisfying all inequalities of the form  $x_i \leq x_j + x_k$  and  $x_i + x_j + x_k \leq 1$ , whenever  $i, j$  and  $k$  are the edges incident to a degree-3 vertex in  $G$ .
- And, as we will see in Section 5, all reflexive polytopes are also semi-reflexive.

## 3 Characterizing semi-reflexive polytopes in terms of their hyperplane description

(We will use the Iverson bracket in this section.)

In this section, mostly deal with full-dimensional polytopes. If  $P$  is full-dimensional, then in the representation (1) the number  $n$  may be chosen to be the number of facets in  $P$ . In this case, each hyperplane  $\{x \mid \langle a_i, x \rangle = b_i\}$

intersects  $P$  in a facet, so that there is no redundant hyperplanes; that is, such representation is minimal. We'll assume such representations are always minimal for full-dimensional polytopes.

For full-dimensional polytopes, there is a slight generalization of the hyperplane description of Theorem 1. Instead of demanding that  $a_i$  is an integer for all  $i$ , we will only require that  $a_i$  is an integer if the corresponding  $b_i$  is 1. Thus, if  $b_i = 0$ , then  $a_i$  may be an arbitrary vector. Observe that, for rational polytopes,  $a_i$  will necessarily be a rational vector, so when  $b_i = 0$  we may multiply both sides by an appropriate integer constant (the lcm of all denominators of the coordinates of  $a_i$ ) to get the same condition of Theorem 1.

One direction is easy.

**Theorem 2.** *Let  $P$  be as in (1). If all  $b_i$  is either 0 or 1, and  $a_i$  is integral whenever  $b_i = 1$ , then  $L_P(s) = L_P(\lfloor s \rfloor)$ .*

*Proof.*

$$L_P(s) = \sum_{x \in \mathbb{Z}^d} [x \in sP] = \sum_{x \in \mathbb{Z}^d} \prod_{i=1}^n [\langle a_i, x \rangle \leq sb_i].$$

If  $b_i = 0$ , the term  $[\langle a_i, x \rangle \leq sb_i]$  reduces to  $[\langle a_i, x \rangle \leq 0]$ , which is constant for all  $s$ ; thus  $[\langle a_i, x \rangle \leq sb_i] = [\langle a_i, x \rangle \leq \lfloor s \rfloor b_i]$ .

If  $b_i = 1$ , as  $x$  and  $a_i$  are integral, the number  $\langle a_i, x \rangle$  is an integer, thus  $[\langle a_i, x \rangle \leq sb_i] = [\langle a_i, x \rangle \leq \lfloor s \rfloor b_i]$  again.

Therefore,

$$\begin{aligned} L_P(s) &= \sum_{x \in \mathbb{Z}^d} \prod_{i=1}^n [\langle a_i, x \rangle \leq sb_i] \\ &= \sum_{x \in \mathbb{Z}^d} \prod_{i=1}^n [\langle a_i, x \rangle \leq \lfloor s \rfloor b_i] \\ &= \sum_{x \in \mathbb{Z}^d} [x \in \lfloor s \rfloor P] \\ &= L_P(\lfloor s \rfloor). \end{aligned} \quad \square$$

For the other direction we need a lemma.

Denote the open ball with radius  $\delta$  centered at  $x$  by  $B_\delta(x)$ ; that is, if  $x \in \mathbb{R}^d$ , we have

$$B_\delta(x) = \{y \in \mathbb{R}^d \mid \|y - x\| < \delta\}.$$

**Lemma 3.** *Let  $K$  be a full-dimensional cone with apex 0, and let  $\delta > 0$  be any value. Then there are infinitely many integer points  $x \in K$  such that  $B_\delta(x) \subset K$ .*

That is, there are many points which are “very inside”  $K$ .

*Proof.* Choose  $x$  to be any rational point in the interior of the cone. By definition, there is some  $\varepsilon > 0$  with  $B_\varepsilon(x) \subseteq K$ . For any  $\lambda > 0$ , we have

$$\lambda B_\varepsilon(x) = B_{\lambda\varepsilon}(\lambda x) \subseteq K.$$




Figure 1: The polytope  $P$ , when dilated, loses the point  $x_0$ , but if the dilation is small enough then it does not gain any new integral point. Therefore,  $L_P(s)$  will decrease from  $s_0$  to  $s_0 + \varepsilon$ .

For all sufficiently large  $\lambda$ , we have  $\lambda\varepsilon > \delta$ , so we just need to take the infinitely many integer  $\lambda$  such that  $\lambda x$  is an integer vector.  $\square$

We'll need the fact that these integral points are distant from the boundary only in the proof of Theorem 5. For the next theorem, existence of infinitely many such points would be enough.

**Proposition 4.** *Let  $P$  be a full-dimensional polytope. If  $0 \notin P$ , then  $L_P(s)$  is not a nondecreasing function. In fact,  $L_P(s)$  has infinitely many “drops”; that is, there are infinitely many points  $s_0$  such that  $L_P(s_0) > L_P(s_0 + \varepsilon)$  for sufficiently small  $\varepsilon > 0$ .*

*Proof.* Writing  $P$  as in (1), we conclude at least one of the  $b_i$  must be negative (because the vector  $0$  satisfies every linear restriction where  $b_i \geq 0$ ). Equivalently (dividing both sides by  $b_i$ ), there is a half-space of the form  $\{x \mid \langle u, x \rangle \geq 1\}$  such that some facet  $F$  of  $P$  is contained in  $\{x \mid \langle u, x \rangle = 1\}$ .

Now, consider the cone  $\bigcup_{\lambda > 0} \lambda F$ . The previous lemma says there is an integral point  $x_0$  in this cone, and so by its definition we have  $x_0 \in s_0 F$  for some  $s_0 > 0$ ; thus,  $x_0 \in s_0 P$ . We'll argue that, for all sufficiently small  $\varepsilon > 0$ , the polytopes  $(s_0 + \varepsilon)P$  do not contain any integral point which is not present in  $s_0 P$  (Figure 1).

For small  $\varepsilon$  (say,  $\varepsilon < 1$ ), all these polytopes are “uniformly bounded”; that is, there is some  $N$  such that  $(s_0 + \varepsilon)P \subseteq [-N, N]^d$  for all  $0 \leq \varepsilon < 1$ . Each integral  $x$  in  $[-N, N]^d$  which is not in  $s_0 P$  must violate a linear restriction of the form  $\langle a_i, x \rangle \leq s_0 b_i$ ; that is,  $\langle a_i, x \rangle > s_0 b_i$  for some  $i$ . As we're dealing with real variables, we also have  $\langle a_i, x \rangle > (s_0 + \varepsilon)b_i$  for all sufficiently small  $\varepsilon$ , say, for all  $\varepsilon < \delta_x$  for some  $\delta_x > 0$ . Now, as there is a finite number of such relevant integral  $x$ , we can take  $\delta$  to be the smallest of all such  $\delta_x$ ; then if  $0 < \varepsilon < \delta$  every integral point of  $(s_0 + \varepsilon)P$  also appears in  $s_0 P$ .

But the special restriction  $\langle u, x \rangle \geq 1$ , considered above, “dilates” to the linear restriction  $\langle u, x \rangle \geq s_0 + \varepsilon$  in  $(s_0 + \varepsilon)P$ . Since  $x_0$  satisfy this restriction with equality for  $\varepsilon = 0$ , for any  $\varepsilon > 0$  we’ll have  $x_0 \notin (s_0 + \varepsilon)P$ . Therefore, not only the dilates  $(s_0 + \varepsilon)P$  do not contain new integral points (for small enough  $\varepsilon$ ), but actually these dilates lose the point  $x_0$  if  $\varepsilon > 0$ . Thus,  $L_P(s_0) > L_P(s_0 + \varepsilon)$  for all sufficiently small  $\varepsilon > 0$ .

Finally, there are infinitely many integral  $x_0$  in the cone  $\bigcup_{\lambda > 0} \lambda F$ ; thus we have infinitely many different values of  $\|x_0\|$  and hence of  $s_0$ , and the reasoning above shows  $L_P(s)$  “drops” in every such  $s_0$ .  $\square$

A simple consequence of this proposition is that if  $s_0$  is a point where  $L_P(s)$  “drops”, then in the interval  $[[s_0], [s_0] + 1)$  the function  $L_P(s)$  will not be a constant function, so we cannot have  $L_P(s) = L_P(\lfloor s \rfloor)$  for all  $s$  if  $P$  does not contain the origin.

We’re now able to show that, for full-dimensional polytopes, the converse of Theorem 2 holds.

**Theorem 5.** *Let  $P$  be a full-dimensional polytope. If  $L_P(s) = L_P(\lfloor s \rfloor)$ , then the polytope  $P$  can be written as in (1) where each  $b_i$  is either 0 or 1, and when  $b_i = 1$  the vector  $a_i$  must be integral.*

*Proof.* Since  $0 \in P$ , by the previous proposition, we know we can write  $P$  as in (1), with each  $b_i$  being nonnegative. By dividing each inequality by the corresponding  $b_i$  (if  $b_i \neq 0$ ), we may assume each  $b_i$  is either 0 or 1. Now we must show that when  $b_i = 1$ , the vector  $a_i$  will be integral.

Let  $\langle u, x \rangle \leq 1$  be one of the linear restrictions where  $b_i = 1$ . Using an approach similar to the proof of the previous proposition, we’ll show  $u$  must be an integral vector.

Let  $F$  be the facet of  $P$  which is contained in  $\{\langle u, x \rangle = 1\}$ , and again define  $K = \bigcup_{\lambda \geq 0} \lambda F$ . Suppose  $u$  has a non-integer coordinate; first, we’ll find an integral point  $x_0$  and a non-integer  $s_0 > 0$  such that  $x_0 \in s_0 F$ .

By Lemma 3 (using  $\delta = \frac{3}{2}$ ), there is some integral  $y \in K$  such that  $B_{\frac{3}{2}}(y) \subset K$ . If  $\langle u, y \rangle$  is not an integer, we may let  $x_0 = y$  and  $s_0 = \langle u, y \rangle$ ; then  $x_0$  is an integral point which is in the relative interior of  $s_0 F$ . If  $\langle u, y \rangle$  is an integer, as  $u$  is not an integral vector, some of its coordinates is not an integer, say the  $j$ th; then  $\langle u, y + e_j \rangle$  will not be an integer, so (as  $y + e_j \in B_{\frac{3}{2}}(y) \subset K$ ) we may let  $x_0 = y + e_j$  and  $s_0 = \langle u, y + e_j \rangle$  to obtain our desired integral point which is in a non-integral dilate of  $F$ .

We have  $sP \subset s_0 P$  for  $s < s_0$ , but  $x_0 \notin sP$  for any  $s < s_0$ , so  $L_P(s) < L_P(s_0)$  for all  $s < s_0$ . As  $s_0$  is not an integer (by construction), this shows that  $L_P(\lfloor s_0 \rfloor) < L_P(s_0)$ , a contradiction.  $\square$

Finally, using unimodular transforms, we may show the characterization for rational polytopes.

**Theorem 1.** *Let  $P$  be a rational polytope written as in (1). Then  $L_P(s) = L_P(\lfloor s \rfloor)$  if and only if all  $a_i$  are integers and all  $b_i$  are either 0 or 1.*

*Proof.* The “if” part is Theorem 2, so assume that  $L_P(s) = L_P(\lfloor s \rfloor)$ . By Proposition 4, we must have  $0 \in P$ . If  $P$  is full-dimensional, we just need to apply Theorem 5: if any of the  $a_i$  is non-integer, then it must be rational (because  $P$  is rational) and the corresponding  $b_i$  must be zero, so we may just multiply the inequality by the lcm of the denominators of the coordinates of  $a_i$ . Thus, assume that  $P$  is not full-dimensional.

Let  $L = \text{aff } P$ , the affine hull of  $P$ . Since  $P$  contains the origin,  $L$  is a vector space; since  $P$  is rational,  $L$  is spanned by integer points. Let  $\dim P$  be the dimension of  $P$  (and of  $L$ ) and let  $d$  be the dimension of the ambient space (so that  $P \subseteq \mathbb{R}^d$ ). Then there is a unimodular transform  $M$  which maps  $L$  to  $\mathbb{R}^{\dim P} \times \{0\}^{d-\dim P}$ .

If  $P$  is contained in the half-space

$$\{x \in \mathbb{R}^d \mid \langle a, x \rangle \leq b\},$$

then  $MP$  is contained in the half-space

$$\{x \in \mathbb{R}^d \mid \langle M^{-t}a, x \rangle \leq b\},$$

so applying unimodular transforms don’t change neither the hypothesis nor the conclusions. Thus, let  $Q$  be the projection of  $MP$  to  $\mathbb{R}^{\dim P}$ ; then  $Q$  is a full-dimensional polytope which satisfies  $L_Q(s) = L_P(s)$ , so we may apply Theorem 5 to conclude the proof.  $\square$

We finish this section by remarking that the hypothesis of  $P$  being either full-dimensional or rational is indeed necessary. For example, if  $H$  is the set of vectors which are orthogonal to  $(\ln 2, \ln 3, \ln 5, \ln 7, \dots, \ln p_d)$  (where  $p_d$  is the  $d$ th prime number) and  $x = (x_1, \dots, x_d)$  is an integral vector,  $x \in H$  if and only if

$$x_1 \log 2 + \dots + x_d \log p_d = 0,$$

which (by applying  $e^x$  to both sides) we may rewrite as

$$2^{x_1} 3^{x_2} \dots p_d^{x_d} = 1,$$

which is only possible if  $x_1 = \dots = x_d = 0$ . That is, the only integral point in  $H$  is the origin. So, if  $P \subseteq H$ , then  $L_P(s)$  will be a constant function, regardless of any other assumption over  $P$ .

## 4 Interiors of polytopes

It is interesting to note there are results similar to Theorems 1 and 5, but for interiors of polytopes. More precisely:

**Theorem 6.** *Let  $P$  be a full-dimensional polytope written as in (1). Then  $L_{P^\circ}(s) = L_{P^\circ}(\lceil s \rceil)$  for all  $s \geq 0$  if and only if all  $b_i$  is either 0 or 1, and  $a_i$  is integral whenever  $b_i = 1$ .*

In other words,  $L_P(s) = L_P(\lfloor s \rfloor)$  if and only if  $L_{P^\circ}(s) = L_{P^\circ}(\lceil s \rceil)$ .

*Proof.* The proof is very similar to the proofs of the theorems 2, 4, and 5, so only the needed changes will be stated.

For Theorem 2, the case  $b_i = 0$  is left unchanged, and when  $b_i = 1$  we need to use  $[n < x] = [n < \lceil x \rceil]$  to conclude that, in all cases,  $[\langle a_i, x \rangle < sb_i] = [\langle a_i, x \rangle < \lceil s \rceil b_i]$ .

For Proposition 4, we can use the same  $u$ ,  $s_0$  and  $x_0$ , but now we will show  $L_{P^\circ}(s_0) < L_{P^\circ}(s_0 - \varepsilon)$ . A point  $x$  which is in the interior of  $s_0 P^\circ$  satisfy all linear restrictions of the form  $\langle x, a_i \rangle < s_0 b_i$ . For all sufficiently small  $\varepsilon > 0$  we have  $\langle x, a_i \rangle < (s_0 - \varepsilon) b_i$ , and so every integer point in  $s_0 P^\circ$  will also be present on  $(s_0 - \varepsilon) P^\circ$ . Now the linear restriction  $\langle u, x \rangle > 1$  “shrinks” to  $\langle u, x \rangle > 1 - \varepsilon$ , and so the point  $x_0$  (which satisfy  $\langle u, x_0 \rangle = 1$ ) will be contained in  $(s_0 - \varepsilon) P^\circ$ , and thus  $L_{P^\circ}(s_0 - \varepsilon) > L_{P^\circ}(s_0)$ .

And for Theorem 5, we again can use the same  $s_0$  and  $x_0$  and also the face  $F$ , but now we will dilate  $P^\circ$  instead of shrinking. As  $sP \subseteq s'P$  for all  $s \leq s'$ , we also have  $sP^\circ \subseteq s'P^\circ$  for all  $s \leq s'$ . Now  $x_0 \in s_0 F$ , so for all  $\varepsilon > 0$  we will have  $x_0 \in (s_0 + \varepsilon) P^\circ$ , which thus show  $L_P(s_0) < L_P(s_0 + \varepsilon)$ .  $\square$

Thus, using essentially the same proof as of Theorem 1, we have the following characterization of semi-reflexive polytopes.

**Theorem 7.** *Let  $P$  be a rational polytope. Then  $P$  is semi-reflexive if and only if  $L_{P^\circ}(s) = L_{P^\circ}(\lceil s \rceil)$  for all real  $s \geq 0$ .*

## 5 Relation with reflexive polytopes

The hyperplane representation of Theorem 1 may be rewritten in matricial form as follows:

$$P = \{x \in \mathbb{R}^d \mid A_1 x \leq \mathbf{1}, A_2 x \leq 0\}, \quad (2)$$

where  $\mathbf{1} = (1, \dots, 1)$  is the all-ones vector, and  $A_1$  and  $A_2$  are integer matrices. We will use this representation to relate semi-reflexive and reflexive polytopes.

There are several equivalent definitions of reflexive polytopes (see e.g. [1, p. 97]). We mention two of them.

First,  $P$  is a reflexive polytope if it is an integer polytope which may be written as

$$P = \{x \in \mathbb{R}^d \mid Ax \leq \mathbf{1}\},$$

where  $\mathbf{1} = (1, \dots, 1)$  is the all-ones vector and  $A$  is an integer matrix. This definition make it obvious that every reflexive polytope is also semi-reflexive.

The second definition uses the dual  $P^*$  of a polytope  $P$ , defined by

$$P^* = \{x \in \mathbb{R}^d \mid \langle x, y \rangle \leq 1 \text{ for all } y \in P\}.$$

Then  $P$  is reflexive if and only if both  $P$  and  $P^*$  are integer polytopes.

We have the following relation between reflexive and semi-reflexive polytopes.

**Theorem 8.**  *$P$  is a reflexive polytope if and only if both  $P$  and  $P^*$  are semi-reflexive polytopes.*

*Proof.* If  $P$  is reflexive, then  $P^*$  is also reflexive, and thus both are semi-reflexive.

Now, suppose that  $P$  and  $P^*$  are semi-reflexive polytopes. This contains the implicit assumption that  $P^*$  is bounded, and thus  $P$  must contain the origin in its interior. Therefore, by Theorem 1, we must have

$$P = \{x \in \mathbb{R}^d \mid Ax \leq \mathbf{1}\}$$

for some integer matrix  $A$ . (The fact that 0 is in the interior of  $P$  allowed us to ignore  $A_2$  in the representation (2).)

So, we just need to show that  $P$  has integer vertices. Since  $(P^*)^* = P$ , we may apply the same reasoning to  $P^*$  to write

$$P^* = \bigcap_{i=1}^n \{x \in \mathbb{R}^d \mid \langle a_i, x \rangle \leq 1\}$$

for certain integers  $a_1, \dots, a_n$ . But these  $a_i$  are precisely the vertices of  $P$ , being, thus, an integral polytope.  $\square$

## References

- [1] Matthias Beck and Sinai Robins. *Computing the Continuous Discretely: Integer-Point Enumeration in Polyhedra*. Undergraduate Texts in Mathematics. Springer, New York, second edition, 2015.
- [2] Cristina Gomes Fernandes, José Coelho de Pina, Jorge Luis Ramírez Alfonso, and Sinai Robins. Polytopes attached to cubic graphs, and their Ehrhart quasi-polynomials. In preparation.
- [3] Richard Peter Stanley. Two poset polytopes. *Discrete & Computational Geometry*, 1(1):9–23, 1986.