

# The Binomial Spin Glass

Mohammad-Sadegh Vaezi,<sup>1</sup> Gerardo Ortiz,<sup>2,3</sup> Martin Weigel,<sup>4</sup> and Zohar Nussinov<sup>1,\*</sup>

<sup>1</sup>*Department of Physics, Washington University, St. Louis, MO 63160, USA*

<sup>2</sup>*Department of Physics, Indiana University, Bloomington, IN 47405, USA*

<sup>3</sup>*Department of Physics, University of Illinois, 1110 W. Green Street, Urbana, Illinois 61801, USA*

<sup>4</sup>*Applied Mathematics Research Centre, Coventry University, Coventry CV1 5FB, UK*

To establish a unified framework for studying both discrete and continuous coupling distributions, we introduce the *binomial* spin glass, a class of models where the couplings are sums of  $m$  identically distributed Bernoulli random variables. In the continuum limit  $m \rightarrow \infty$ , the class reduces to one with Gaussian couplings, while  $m = 1$  corresponds to the  $\pm J$  spin glass. We demonstrate that for short-range Ising models on  $d$ -dimensional hypercubic lattices the ground-state entropy density for  $N$  spins is bounded from above by  $(\sqrt{d/2m} + 1/N) \ln 2$ , and further show that the actual entropies follow the scaling behavior implied by this bound. We thus uncover a fundamental non-commutativity of the thermodynamic and continuous coupling limits that leads to the presence or absence of degeneracies depending on the precise way the limits are taken. Exact calculations of defect energies reveal a crossover length scale  $L^*(m) \sim L^\kappa$  below which the binomial spin glass is indistinguishable from the Gaussian system. Since  $\kappa = -1/(2\theta)$ , where  $\theta$  is the spin-stiffness exponent, discrete couplings become irrelevant at large scales for systems with a finite-temperature spin-glass phase.

PACS numbers: 05.50.+q, 64.60.De, 75.10.Hk

Spin glasses are extremely rich systems that have continued to surprise for many decades [1–13]. They represent paradigmatic realizations of complexity that are abundant in nature and numerous combinatorial optimization problems [14]. Abstractions of spin-glass physics have led to new optimization algorithms and new insight into computational complexity [15–18], shed light on protein folding [19], and provided models of neural networks [20]. Notwithstanding this success, several fundamental questions still linger. These include [21] the character of the low-lying states and whether there are many incongruent [22] ground states. It has long been known that spin-glass systems with discrete couplings may rigorously exhibit an extensive degeneracy [23, 24], but these results do not extend to continuous coupling distributions [25–29]. The possibility of vanishing spectral gaps mandates the distinction of localized and extended excitations, and only the latter can give rise to a multitude of states.

In this paper, we connect the  $\pm J$  and the Gaussian spin glass models by interpolating them via the *binomial* spin glass that has a tunable control parameter  $m$ . We establish bounds of the spectral degeneracy of the Ising system on bipartite graphs, which includes the usual Edwards-Anderson (EA) model with  $\pm J$  ( $m = 1$ ) and Gaussian ( $m \rightarrow \infty$ ) couplings [5–18, 30]. We thus show that discrete (finite  $m$ ) spin-glass samples exhibit an extensive ground-state degeneracy, while continuous ones ( $m \rightarrow \infty$ ) become two-fold degenerate, while more generally the degeneracy depends on the precise way the non-commuting limits  $N \rightarrow \infty$  and  $m \rightarrow \infty$  are taken.

We define the *binomial Ising spin glass* on a graph of

$N$  sites [45] by the Hamiltonian

$$H_m = - \sum_{\langle xy \rangle} \mathcal{J}_{xy}^m s_x s_y \equiv - \sum_{\alpha=1}^{\mathcal{L}} \mathcal{J}_\alpha^m z_\alpha. \quad (1)$$

Here, the sum is over sites  $x$  and  $y$ , defining a link  $\alpha = \langle xy \rangle$ ,  $\mathcal{L}$  denotes the total number of links, and  $s_x = \pm 1$ . The *binomial coupling* for each link  $\alpha$ ,  $\mathcal{J}_\alpha^m \equiv \frac{1}{\sqrt{m}} \sum_{k=1}^m J_\alpha^{(k)}$ , is a sum of  $m$  copies (or “layers”) of binary couplings  $J_\alpha^{(k)} = \pm 1$ , each with probability  $p$  of being  $+1$ . The probability distribution of  $\mathcal{J}_\alpha^m$ ,

$$\tilde{P}(\mathcal{J}_\alpha^m) = \sum_{j=0}^m \binom{m}{j} p^{m-j} (1-p)^j \delta\left(\mathcal{J}_\alpha^m - \frac{m-2j}{\sqrt{m}}\right), \quad (2)$$

is a binomial. In the large- $m$  limit, the distribution (2) approaches a Gaussian of mean  $\sqrt{m}(2p-1)$  and variance  $\sigma^2 = 4p(1-p)$ . In particular, for  $p = 1/2$ , the distribution  $\tilde{P}(\mathcal{J}_\alpha^m)$  approaches the standard normal distribution usually considered for the EA model [10].

To understand the degeneracies in the spectrum, we study the entropy density of the  $\ell$ -th energy level,

$$\mathcal{S}_\ell \equiv \frac{\sum_{\{\mathcal{J}_\alpha^m\}} P(\{\mathcal{J}_\alpha^m\}) \ln D_\ell(\{\mathcal{J}_\alpha^m\})}{N}, \quad (3)$$

where  $D_\ell$  is the degeneracy of the  $\ell$ -th energy level [23].  $P(\{\mathcal{J}_\alpha^m\}) = \prod_{\alpha=1}^{\mathcal{L}} \tilde{P}(\mathcal{J}_\alpha^m)$  is the probability of the coupling configuration.

We first embark on the derivation of an upper bound on the ground state entropy density  $\mathcal{S}_0$ . We restrict ourselves to bipartite graphs, where any closed loop encompasses an even number of links  $\alpha$ . Consider two spin

\* zohar@wuphys.wustl.edu

configurations  $|\mathfrak{s}\rangle \neq |\mathfrak{s}'\rangle$  and evaluate their energy difference  $\Delta E = E(\mathfrak{s}) - E(\mathfrak{s}')$ . From Eq. (1),

$$\Delta E = - \sum_{\alpha=1}^{\mathcal{L}} \mathcal{J}_{\alpha}^m \left( z_{\alpha}(\mathfrak{s}) - z_{\alpha}(\mathfrak{s}') \right) = -2 \sum_{\alpha=1}^{\mathcal{L}} \mathcal{J}_{\alpha}^m n_{\alpha}, \quad (4)$$

with integers  $n_{\alpha} = 0, \pm 1$  defined by  $n_{\alpha} \equiv [z_{\alpha}(\mathfrak{s}) - z_{\alpha}(\mathfrak{s}')]/2$ , where  $z_{\alpha}(\mathfrak{s}) = s_x s_y$ . If  $|\mathfrak{s}\rangle$  and  $|\mathfrak{s}'\rangle$  are degenerate then  $\Delta E = 0$ . A degeneracy only occurs for some realizations  $\{\mathcal{J}_{\alpha}^m\}$  of the couplings, and Eq. (4) can be understood as a set of conditions for the couplings to ensure this.

Consider an arbitrary reference configuration  $|\mathfrak{s}\rangle$  of energy  $E(\mathfrak{s})$  and examine its viable degeneracy with the contending  $2^N - 1$  other configurations  $|\mathfrak{s}'\rangle$ . Each of these leads to a particular set of integers  $\mathbf{C}_j = \{n_{\alpha}\}_j$ , which form the set  $\{\mathbf{C}_j\}_{j=1, 2^N-1}^{|\mathfrak{s}\rangle}$ . A subset of those,  $\text{Sat}_{|\mathfrak{s}\rangle} = \{\mathbf{C}_{j_1}, \mathbf{C}_{j_2}, \dots, \mathbf{C}_{j_N}\}$ , will satisfy the degeneracy condition  $\Delta E = 0$  in Eq. (4) for some coupling realizations. There are two types of solutions to the equation  $\Delta E = 0$ : (i)  $n_{\alpha} = 0, \forall \alpha$ , or (ii)  $n_{\alpha} \neq 0$ , for at least one link  $\alpha$ . It is straightforward to demonstrate that there is a single configuration  $|\mathfrak{s}'\rangle (\neq |\mathfrak{s}\rangle)$  for which (i)  $n_{\alpha} = 0, \forall \alpha$  [46]. This is the degenerate configuration  $|\mathfrak{s}'\rangle$  obtained by inverting all of the spins in  $|\mathfrak{s}\rangle$ . To determine whether the degeneracy may be larger than two, we need to compute the probability  $\mathcal{P}$  that constraints of type (ii) may be satisfied. While we cannot exactly calculate this probability for general  $N$  and  $m$ , bounds that we will derive suggest that  $\lim_{N \rightarrow \infty} \lim_{m \rightarrow \infty} \mathcal{S}_{\ell} = 0$ . As we will emphasize, different large  $m$  and  $N$  limits may yield incompatible results.

Constraints  $\mathbf{C}_j \in \text{Sat}_{|\mathfrak{s}\rangle}$  are in a *one-to-one correspondence* with zero-energy interfaces [47], whose *size* is equal to the number  $g_j$  of non-zero integers in the set  $\{n_{\alpha}\}_j$ . That is, given a fixed reference configuration  $|\mathfrak{s}\rangle$  and a degenerate one  $|\mathfrak{s}'\rangle$ , all type (ii) solutions to Eq. (4) are associated with configurations where the product  $s_x s'_x$  is equal to  $-1$  in a non-empty set of sites  $x \in R$ . To avoid the trivial redundancy due to global spin inversion, consider the states  $|\mathfrak{s}\rangle$  and  $|\mathfrak{s}'\rangle$  for which the spin at an arbitrarily chosen ‘‘origin’’ of the lattice assumes the value  $+1$ . These states are related via  $|\mathfrak{s}'\rangle = U_{\mathfrak{s}'\mathfrak{s}} |\mathfrak{s}\rangle$ , where the *domain-wall operator*  $U_{\mathfrak{s}'\mathfrak{s}}$  is the product of Pauli matrices that flip the sign of the spins  $s'_x$  at the sites  $x$  where  $|\mathfrak{s}\rangle$  and  $|\mathfrak{s}'\rangle$  differ. Regions  $R$  are bounded by zero-energy domain walls that are interfaces dual to the links with  $n_{\alpha} = \pm 1$ , i.e., surrounding the areas  $R$  where the spins in  $|\mathfrak{s}\rangle$  and  $|\mathfrak{s}'\rangle$  have opposite orientation. Each satisfied constraint  $\mathbf{C}_j \in \text{Sat}_{|\mathfrak{s}\rangle}$  is associated with a state  $|\mathfrak{s}'\rangle = U_{\mathfrak{s}'\mathfrak{s}} |\mathfrak{s}\rangle$  that is degenerate with  $|\mathfrak{s}\rangle$  for some coupling realization(s).

We next formalize the counting of *independent domain walls* or clusters of free spins to arrive at an asymptotic bound on their number [Eq. (9)]. This will, in turn, provide a bound on the degeneracy. We define a *complete* set of *independent* constraints  $\overline{\text{Sat}}_{|\mathfrak{s}\rangle} \subset \text{Sat}_{|\mathfrak{s}\rangle}$ , of cardinality  $\mathcal{M}$ , to be composed of *all* constraints  $\mathbf{C}_j \in \text{Sat}_{|\mathfrak{s}\rangle}$  that lead to *linearly independent equations* of the form

of Eq. (4),  $\Delta E = E(\mathfrak{s}) - E(\mathfrak{s}_j) = 0$ , on the coupling constants  $\{\mathcal{J}_{\alpha}^m\}$  [47]. All constraints in  $\text{Sat}_{|\mathfrak{s}\rangle}$  are a consequence of the linearly independent subset of constraints  $\overline{\text{Sat}}_{|\mathfrak{s}\rangle}$ . Each constraint  $\mathbf{C}_j \in \text{Sat}_{|\mathfrak{s}\rangle}$  is associated with a domain wall operator  $U_{\mathfrak{s}_j\mathfrak{s}}$  that generates a degenerate state  $|\mathfrak{s}_j\rangle = U_{\mathfrak{s}_j\mathfrak{s}} |\mathfrak{s}\rangle$ . If for a given coupling realization  $\{\mathcal{J}_{\alpha}^m\}$  there are  $M(\{\mathcal{J}_{\alpha}^m\}) \leq \mathcal{M}$  such independently satisfied constraints, then the states

$$|\bar{n}_1 \bar{n}_2 \dots \bar{n}_M\rangle \equiv U_{\mathfrak{s}_1\mathfrak{s}}^{\bar{n}_1} U_{\mathfrak{s}_2\mathfrak{s}}^{\bar{n}_2} \dots U_{\mathfrak{s}_M\mathfrak{s}}^{\bar{n}_M} |\mathfrak{s}\rangle, \quad (5)$$

( $\bar{n}_i = 0, 1$ ) will include all of the spin configurations degenerate with  $|\mathfrak{s}\rangle$ . Taking global spin inversion into account, the degeneracy of  $|\mathfrak{s}\rangle$  is

$$D_{\ell(|\mathfrak{s}\rangle, \{\mathcal{J}_{\alpha}^m\})} \leq 2^{M(\{\mathcal{J}_{\alpha}^m\})+1}, \quad (6)$$

where, for a system defined by the coupling constants  $\{\mathcal{J}_{\alpha}^m\}$ , the index  $\ell(|\mathfrak{s}\rangle, \{\mathcal{J}_{\alpha}^m\})$  denotes the level  $\ell$  the state  $|\mathfrak{s}\rangle$  belongs to. The set  $\{|\bar{n}_1 \bar{n}_2 \dots \bar{n}_M\rangle\}$  may contain additional states not degenerate with  $|\mathfrak{s}\rangle$  [48].

After averaging over disorder, the expected number of the linearly independent satisfied constraints  $\text{Sat}_{|\mathfrak{s}\rangle}$  is

$$\langle M \rangle_m \equiv \sum_{\{\mathcal{J}_{\alpha}^m\}} \sum_{\mathbf{C}_j \in \overline{\text{Sat}}_{|\mathfrak{s}\rangle}} P(\{\mathcal{J}_{\alpha}^m\}) \delta^{\{\mathcal{J}_{\alpha}^m\}}(\mathbf{C}_j) \equiv \sum_{\mathbf{C}_j \in \overline{\text{Sat}}_{|\mathfrak{s}\rangle}} \mathcal{P}(\mathbf{C}_j). \quad (7)$$

Here,  $\mathcal{P}(\mathbf{C}_j)$  is the probability that a linearly independent constraint  $\mathbf{C}_j$  is satisfied. The Kronecker  $\delta^{\{\mathcal{J}_{\alpha}^m\}}(\mathbf{C}_j)$  equals 1 if  $\mathbf{C}_j$  is satisfied for the couplings  $\{\mathcal{J}_{\alpha}^m\}$  and is zero otherwise. Let us bound the probability  $\mathcal{P}(\mathbf{C}_j)$  by taking the form (2) of the coupling distribution into account. From the definition of the couplings  $\{\mathcal{J}_{\alpha}^m\}$ , the sum in Eq. (4) can effectively be read as including a sum over layers  $k = 1, \dots, m$ , which hence includes  $g_j m$  non-zero terms. For general  $m \geq 1$ , and even  $g_j m$ , the probability that half of the nonzero integers  $n_{\alpha} \mathcal{J}_{\alpha}^{(k)}$  in Eq. (4) are  $+1$  and the remainder are  $-1$  is

$$\mathcal{P}(\mathbf{C}_j) = \binom{g_j m}{\frac{g_j m}{2}} \frac{1}{2^{g_j m}} < \frac{1}{\sqrt{g_j m}}. \quad (8)$$

(Eq. (4) cannot be satisfied for odd  $g_j m$ .) From asymptotic analysis [49] and Eq. (8), the probability  $\mathcal{P}(\mathbf{C}_j)$  scales (for large  $m$ ) as (and, for any  $m$ , is bounded by)  $1/\sqrt{g_j m}$ . Denoting by  $g_{\min}$  the smallest possible value of  $g_j$  for the graph/lattice at hand,

$$\langle M \rangle_m \leq \frac{\mathcal{M}}{\sqrt{g_{\min} m}}. \quad (9)$$

On a general graph, the number  $\mathcal{M}$  of linearly *independent* constraints  $\mathbf{C}_j$  on the coupling constants  $\{\mathcal{J}_{\alpha}^m\}$  cannot be larger than their total number,  $\mathcal{M} \leq \mathcal{L}$ , i.e., the number of links  $\mathcal{L}$  on this graph. Putting all of the pieces together, Eqs. (6) and (9) imply

$$\sum_{\{\mathcal{J}_{\alpha}^m\}} P(\{\mathcal{J}_{\alpha}^m\}) \ln D_{\ell(|\mathfrak{s}\rangle, \{\mathcal{J}_{\alpha}^m\})} \leq \left(1 + \frac{\mathcal{L}}{\sqrt{g_{\min} m}}\right) \ln 2. \quad (10)$$




FIG. 1. Ground-state entropy  $\mathcal{S}_0 N$  of the binomial Ising spin glass with  $m$  layers, cf. Eq. (1), on square lattices of  $N = L^2$  spins from exact ground-state calculations (from the bottom:  $L = 8, 16, 20, 24$ , and  $32$ ). Lines are fits of the form of (13) to the data for sufficiently large  $m$ . The inset shows the linear scaling of the amplitude  $A(N)$ . The top line indicates the constraint imposed by the upper bound (11).

Trying to evaluate the l.h.s. of Eq. (10) we must take into account that whatever  $|s\rangle$  we pick might be a ground state for some coupling configurations, but will be an excited state for others. Hence we cannot directly infer a bound to the average entropy  $\mathcal{S}_\ell$  from (10). Since the inverse temperature  $1/(k_B T) = \partial \ln D / \partial E$ , however, the system's ground-state degeneracy for couplings  $\{\mathcal{J}_\alpha^m\}$  is typically lower than (or equal to) that of any other level  $\ell$  [50], i.e.,  $D_0 \leq D_\ell$ . This monotonicity of  $D(E)$  implies that, typically,  $\mathcal{S}_0 N = \sum_{\{\mathcal{J}_\alpha^m\}} P(\{\mathcal{J}_\alpha^m\}) \ln D_0(\{\mathcal{J}_\alpha^m\}) \leq \sum_{\{\mathcal{J}_\alpha^m\}} P(\{\mathcal{J}_\alpha^m\}) \ln D_\ell(|s\rangle, \{\mathcal{J}_\alpha^m\})$ . Then, Eq. (10) yields

$$\mathcal{S}_0 \leq \left( \frac{\mathcal{L}}{N \sqrt{g_{\min} m}} + \frac{1}{N} \right) \ln 2. \quad (11)$$

This is the promised rigorous bound. For  $p \neq 1/2$  one has a lower entropy density than that of  $p = 1/2$ . Thus, Eq. (11) constitutes a generous upper bound on  $\mathcal{S}_0$  for general  $p$ . To study higher energy levels, consider the average of Eq. (10) over all possible  $2^N$  reference spin configurations  $|s\rangle$ . Performing this average and invoking the monotonicity of  $D(E)$  suggests that the entropy density  $\mathcal{S}_\ell$  of Eq. (3) of low-lying excited levels  $\ell > 0$  is, typically, also bounded by the r.h.s of Eq. (11). For  $d$ -dimensional hypercubic lattices with periodic boundary conditions, the ratio  $\mathcal{L}/N = d$  while  $g_{\min} = 2d$ . Thus,  $\mathcal{S}_0 \leq (\sqrt{d/2m} + 1/N) \ln 2$ . Eq. (11) further suggests that, in the thermodynamic ( $N \rightarrow \infty$ ) limit [51],

$$\mathcal{S}_0(m') \sim \sqrt{\frac{m}{m'}} \mathcal{S}_0(m) \quad \text{for finite } m, m' \gg 1. \quad (12)$$

We now study the exact  $m$  dependence of the ground state entropies of the binomial model on the square lattice with periodic boundaries and  $N = L^2$ . To this end,

we employed an implementation of the Pfaffian technique of counting dimer coverings of the lattice as discussed in Ref. [52], which is a generalization of earlier methods [53, 54] to fully periodic lattices. In Fig. 1, we present the results for the ground-state entropy, averaged over 1000 coupling realizations for each lattice size. The data are well described by

$$\mathcal{S}_0 N = \left( \frac{A(N)}{\sqrt{m}} + 1 \right) \ln 2. \quad (13)$$

Linear fits in  $1/\sqrt{m}$  for fixed  $N$  work well for sufficiently large  $m$ , as is illustrated by the straight lines in Fig. 1. Thus, for any finite  $N$ , as  $m \rightarrow \infty$  the ground-state entropy is equal to  $\ln 2$ , implying a single degenerate ground-state pair. The slope  $A(N)$  shown in the inset follows a linear behavior,  $A(N) = aN + b$ , and we find  $a = 0.0858(4)$  and  $b = 1.09(12)$ . For not too small  $m$ , our data are hence fully consistent with

$$\mathcal{S}_0 = \left( \frac{a}{\sqrt{m}} + \frac{1}{N} + \frac{b}{N\sqrt{m}} \right) \ln 2. \quad (14)$$

When  $N \gg \sqrt{m} \gg 1$ , Eq. (14) is consistent with the physically inspired [51] scaling of Eq. (12). For large  $N$ , the bound of Eq. (11) would have been asymptotically saturated if  $a \simeq 1$ , far larger than the actual value of  $a$ . The behavior in the double limit  $m, N \rightarrow \infty$  is subtle: (1) for  $m \rightarrow \infty$ ,  $N$  finite, we have a single ground-state pair; (2) for  $N \rightarrow \infty$ ,  $m$  finite, there is a finite ground-state entropy  $\sim \ln 2/\sqrt{m}$ ; (3) for  $N \rightarrow \infty$ ,  $m \rightarrow \infty$ ,  $\kappa = N/\sqrt{m}$  fixed, there is a finite number  $2^{a\kappa}$  of ground-state pairs. Thus clearly the continuum and thermodynamic limits are not commutative in general. Note further that according to the bound  $\mathcal{S}_0 \leq (\sqrt{d/2m} + 1/N) \ln 2$  for hypercubic lattices additional rich behavior is expected if the limit of high dimensions is correlated with that of large  $m$ .

Let us turn to the study of excitations. By construction, cf. Eq. (4), for finite  $m$  the energy is ‘‘quantized’’ in multiples of  $1/\sqrt{m}$ . It is therefore natural to expect a closing of the spectral gap as  $m \rightarrow \infty$ . That this is indeed the case can be shown rigorously for the one-dimensional binomial spin glass in its thermodynamic limit, with different behaviors for odd and even  $m$ , see the discussion in the Supplemental Material [55]. The closing of the gap is a consequence of the existence of (rare) local excitations, i.e., finite-size clusters of almost free spins [56]. Whether gapless *non-local* excitations exist and which form they take in the thermodynamic limit is a long-standing question [57]. One possible approach of investigating such excitations consists of subjecting individual samples to a system spanning perturbation by a change of boundary condition and studying how this affects the energy and configuration of the ground state. Such defect energy calculations [58] enable us to extract a scaling  $\langle |\Delta E| \rangle \sim L^\theta$  of the defect energies with the spin stiffness exponent  $\theta$ . Generalizing Peierls' argument [59–62] for the stability of the ordered phase, one should




FIG. 2. Effective spin stiffness exponents  $\theta = \theta(m)$  resulting from fits of the power law  $\langle |\Delta E| \rangle = BL^\theta$  to the defect energies for the binomial model of  $m$  layers (inset, from the top:  $m = 1, 5, 11, 51, 201$ , and  $1001$ ), averaged over 10 000 disorder samples. The solid line of the inset corresponds to the Gaussian model.

find  $\theta > 0$  for cases where there is a finite-temperature spin-glass phase, and  $\theta \leq 0$  otherwise. The latter case is expected for dimensions  $d = 1$  and  $d = 2$ , whereas  $\theta$  is positive for  $d \geq 3$  [63, 64]. We employed techniques based on minimum-weight perfect matching [65, 66] to perform such calculations for the binomial model on the square lattice. The resulting disorder-averaged defect energies from *exact* ground-state calculations for samples with periodic and antiperiodic boundaries are shown in the inset of Fig. 2. As  $m$  increases, the decay of defect energies as a function of  $L$  becomes steeper and the data approach the behavior of the Gaussian EA model. The effective spin stiffness exponents  $\theta$  extracted from fits of the type  $\langle |\Delta E| \rangle = BL^\theta$  are shown in the main panel of Fig. 2. These exponents appear to interpolate smoothly between the limiting cases of the Gaussian model with  $\theta = -0.2793(3)$  and the  $\pm J$  system with  $\theta = 0$  [63, 66]. Asymptotically, however, we expect that  $\theta(m) = 0$  for any finite value of  $m$  when  $L \gtrsim L^*(m)$ . The scaling of the crossover length  $L^*(m) \sim m^\kappa$  follows by considering the model with the unscaled couplings  $\sqrt{m}\mathcal{J}_\alpha^m$ , for which the energy gap  $\Delta$  is independent of  $m$ . The discreteness of the spectrum becomes apparent once the corresponding defect energies  $\sqrt{m}\langle |\Delta E| \rangle \sim L^\theta$  have decayed below the size of the gap, i.e., for

$$L \geq L^*(m) \sim m^{-1/(2\theta)},$$

such that  $\kappa = -1/(2\theta)$ . For the  $d = 2$  system we have  $\theta = -0.2793(3)$  [66], such that  $\kappa = 1.790(2)$ , which is in excellent agreement with the actual defect energies for our system shown in Fig. 3.

It is clear that if  $\theta < 0$ , as is the case for the Gaussian spin glass in two dimensions, excitations of a divergent length scale may entail a vanishing energy penalty. At zero temperature, the discreteness of the spectrum is




FIG. 3. Scaling collapse of the defect energies of the binomial model for system sizes rescaled with the crossover length scale  $L^*(m) \sim m^\kappa$  with  $\kappa = 1.79$ .

then always seen at large scales  $L \gtrsim L^*(m)$ . On the other hand, for  $\theta \geq 0$  (i.e.,  $d \geq 3$ ), the above arguments imply that the discreteness does not matter at large scales. Also, in this case one should inspect the full probability distribution of domain wall energies and the weight it carries in the limit  $\Delta E \rightarrow 0$  [56]. In how far such excitations correspond to incongruent states, however, one might only be able to infer by inspecting the configurations themselves.

In summary, we introduced and discussed the *binomial spin glass*. This class of models affords controlled access to the enigmatic continuous ( $m \rightarrow \infty$ ) finite dimensional EA model. Its  $m = 1$  realization is the quintessential discrete spin glass, the  $\pm J$  model. We derived bounds on the spectral degeneracy of the binomial Ising spin glass on general graphs and suggested an asymptotic scaling that is fully supported by exact two-dimensional calculations. The behavior of defect energies suggests the existence of a crossover length  $L^*(m) \sim L^{-1/2\theta}$  below which the binomial model behaves like the Gaussian system. Our results show that the existence of degeneracies depends on the particular way of taking the thermodynamic ( $N \rightarrow \infty$ ) and continuous coupling ( $m \rightarrow \infty$ ) limits, and limiting states with and without degeneracies can be reached by corresponding correlated limiting processes, thus accommodating theories that postulate degeneracies as well as pictures stipulating a unique ground-state pair. An intriguing prediction regards an effectively negative crossover scaling exponent in three dimensions, where hence discreteness of the spectrum is expected not to matter at large scales.

The physics of spin-glass models and, in particular, the role of degeneracies has also recently attracted attention from another side. In the context of quantum annealing [67] as implemented in the devices by D-Wave and similar machines that are being developed by competing consortia, degeneracies are not a desired feature as the quantum annealing process does not sample such

states uniformly [68]. On the other hand, continuous coupling distributions may also be undesired because of increased susceptibility to external noise implied by chaos in spin glasses [69–72]. Our binomial glasses may allow for realizations that suffer the least from these combined problems. While the present system is already a generalization of the usually considered spin-glass models,

we believe that the approach of decomposing continuous couplings into discrete layers and the intriguing consequences it allowed us to uncover in terms of the general non-commutativity of the thermodynamic and continuous coupling limits is promising and we expect exciting applications to models in other fields.

*Acknowledgements.* This research was partially supported by the NSF CMMT under grant number 1411229.

- 
- [1] D. L. Stein and C. M. Newman, *Spin Glasses and Complexity* (Princeton University Press, 2013).
- [2] K. H. Fischer and J. A. Hertz, *Spin Glasses* (Cambridge University Press, Cambridge, 1991).
- [3] D. L. Stein and C. M. Newman, *Complex Systems* **20**, 115 (2011).
- [4] J. A. Mydosh, *Spin Glasses: An Experimental Introduction* (Taylor and Francis, London, Washington D. C., 1993).
- [5] V. Cannella and J. A. Mydosh *Phys. Rev. B* **6**, 4220 (1972).
- [6] H. Nishimori, *Statistical Physics of Spin Glasses and Information Processing: An Introduction* (Oxford University Press, Oxford, 2011).
- [7] H. Nishimori and G. Ortiz *Elements of Phase Transitions and Critical Phenomena* (Oxford University Press, Oxford, 2011).
- [8] P. W. Anderson, *Physics Today* **41**(1), 9 (1988); **41**(3), 9 (1988); **41**(6), 9 (1988); **41**(9), 9 (1988); *Physics Today* **42**(7), 9 (1989); **42**(9), 9 (1989); **43**(3), 9 (1990).
- [9] J. Lukic, A. Galluccio, E. Marinari, Olivier C. Martin, and G. Rinaldi, *Phys. Rev. Lett.* **92**, 117202 (2004).
- [10] S. Edwards and P. W. Anderson, *J. Phys. F* **5**, 965 (1975).
- [11] D. Sherrington and S. Kirkpatrick *Phys. Rev. Lett.* **35**, 1792 (1975).
- [12] K. Binder and A. P. Young, *Rev. Mod. Phys.*, **58**, 80 (1986).
- [13] G. Parisi, *Phys. Rev. Lett.* **43**, 1754 (1979); G. Parisi, *J. Phys. A* **13**, L115 (1980); G. Parisi, *J. Phys. A* **13**, 1101 (1980); G. Parisi, *J. Phys. A* **13**, 1887 (1980); G. Parisi, *Phys. Rev. Lett.* **50**, 1946 (1983).
- [14] F. Barahona, *J. Phys. A* **15**, 3241 (1982).
- [15] M. Mezard, G. Parisi, and M. A. Virasoro, *Europhys. Lett.* **1**, 77 (1985).
- [16] M. Mezard, G. Parisi, and M. A. Virasoro, *Spin Glass Theory and Beyond* (World Scientific, Singapore, 1987).
- [17] M. Mezard, G. Parisi, and R. Zecchina, *Science* **297**, 812 (2002).
- [18] A. Braunstein, M. Mezard, and R. Zecchina, *Random Structures and Algorithms* **27**, 201 (2005).
- [19] J. D. Bryngelson and P. G. Wolynes, *Proceedings of the Natl. Acad. of Science (USA)* **84**, 7524 (1987).
- [20] J. J. Hopfield, *Proc. Natl. Acad. Sci. USA* **79**, 2554 (1982); D. J. Amit, H. Gutfreund, and H. Sompolinsky, *Phys. Rev. Lett.* **55**, 1530 (1985); J. J. Hopfield and D. W. Tank, *Science* **233**, 625 (1986); H. Sompolinsky, *Physics Today* **41** (12), 70 (1988).
- [21] C. M. Newman and D. L. Stein, *Comm. Math. Phys.* **224**, 205 (2001).
- [22] Given any Ising spin configuration one may inspect the sign of each of the links  $z_\alpha$  on the lattice. If there is an extensive (volume proportional) number of links  $z_\alpha$  that are of different signs in two different Ising spin configurations  $|s\rangle$  and  $|s'\rangle$  then the two states are said to be “incongruent” relative to one another [3].
- [23] J. E. Avron, G. Roepstorff, and L. S. Schulman, *J. Stat. Phys.* **26**, 25 (1981).
- [24] M. Loebl and J. Vondrak, *Discrete Mathematics* **271** (1–3), 179 (2003).
- [25] J. W. Landry and S. N. Coppersmith, *Phys. Rev. B* **65**, 134404 (2002).
- [26] E. Marinari and G. Parisi, *Phys. Rev. B* **62**, 11677 (2000).
- [27] H. Rieger, *Frustrated Systems: Ground State Properties via Combinatorial Optimization*, Lecture Notes in Physics Vol. 501 (Springer-Verlag, Heidelberg, 1998).
- [28] J. Houdayer and O. Martin, *Phys. Rev. Lett.* **83**, 1030 (1999).
- [29] R. N. Bhatt and A. P. Young, *Phys. Rev. B* **37**, 5606 (1988).
- [30] See Section A of the Supplemental Material for a brief overview that further discusses Refs. [5, 6, 8–18].
- [31] F. Guerra and F. L. Toninelli, *Comm. Math. Phys.* **230**, 71 (2002);
- [32] M. Talagrand, *Spin Glasses: A Challenge to Mathematicians*, Springer-Verlag (2003).
- [33] M. Mezard, G. Parisi, N. Sourlas, G. Toulouse, and M. Virasoro, *Phys. Rev. Lett.*, **52**, 1156 (1984).
- [34] G. Parisi, *J. Phys.* **45**, 843 (1984).
- [35] R. Rammal, G. Toulouse, and M. A. Virasoro, *Rev. Mod. Phys.* **58**, 765 (1986).
- [36] S. Franz, M. Mezard, G. Parisi, and L. Peliti, *Phys. Rev. Lett.* **81**, 1758 (1998).
- [37] W. L. MacMillan *Phys. Rev. B* **31**, 340 (1985).
- [38] A. J. Bray and M. A. Moore, *Phys. Rev. B* **31**, 631 (1985).
- [39] R. G. Calfish and J. R. Banavar, *Phys. Rev. B* **32**, 7617 (1985).
- [40] M. A. Moore and A. J. Bray, *J. Phys C: Solid State Phys.* **18**, L699 (1985).
- [41] D. S. Fisher and D. A. Huse, *J. Phys. A: Math. Gen.* **20**, L1005 (1987).
- [42] D. A. Huse and D. S. Fisher, *J. Phys. A: Math. Gen.* **20**, L997 (1987).
- [43] C. M. Newman and D. L. Stein, *Phys. Rev. Lett.* **84**, 3966 (2000).
- [44] C. M. Newman and D. L. Stein, *Commun. Math. Phys.* **224**, 205 (2001).
- [45] We exclude classical systems with topological degeneracy, M.-S. Vaezi, G. Ortiz, and Z. Nussinov, *Phys. Rev. B* **93**, 205112 (2016).
- [46] Supplemental Material; see Section B.
- [47] Supplemental Material; see Section C.

- [48] Supplemental Material; see Section D.  
 [49] Notice that one can write the asymptotic form

$$\mathcal{P}(C_j) \sim \sqrt{\frac{2}{\pi g_j m}}, \quad (15)$$

for large  $m$  after applying Stirling's approximation.

- [50] Supplemental Material; see Section E.  
 [51] Supplemental Material; see Section F.  
 [52] A. Galluccio, M. Loebl, and J. Vondrak, *Phys. Rev. Lett.* **84**, 5924 (2000).  
 [53] J. A. Blackman, J. R. Goncalves, and J. Poulter, *Phys. Rev. E* **58**, 1502 (1998).  
 [54] L. Saul and M. Kardar, *Phys. Rev. E* **48**, R3221 (1993).  
 [55] Supplemental Material; see Section G.  
 [56] Supplemental Material; see Section H.  
 [57] N. Kawashima and H. Rieger, in *Frustrated Spin Systems*, edited by H. T. Diep (World Scientific, Singapore, 2005), chap. 9, p. 491.  
 [58] M. Cieplak and J. R. Banavar, *Phys. Rev. B* **27**, 293 (1983).  
 [59] R. Peierls, *Proc. Camb. Phil. Soc.* **32**, 477 (1936).  
 [60] R. B. Griffiths, *Phys. Rev.* **136**, A437 (1964).  
 [61] G. Ortiz, E. Cobanera, and Z. Nussinov, *Nuclear Phys. B* **854**, 780 (2011); see Appendix E in particular.  
 [62] C. Bonati, *European Journal of Physics B* **35**, 035002 (2014).  
 [63] A. K. Hartmann and A. P. Young, *Phys. Rev. B* **64**, 180404 (2001).  
 [64] S. Boettcher, *European Physics Journal B* **38**, 83 (2004).  
 [65] I. Bieche, R. Maynard, R. Rammal, and J. P. Uhry, *J. Phys. A* **13**, 2553 (1980).  
 [66] H. Khoshbakht and M. Weigel, *Phys. Rev. B* **97**, 064410 (2018).  
 [67] A. B. Finnila, M. A. Gomez, C. Sebenik, C. Stenson, and J. D. Doll, *Chem. Phys. Lett.* **219**, 343 (1994).  
 [68] S. Mandrà, Z. Zhu, and H. G. Katzgraber, *Phys. Rev. Lett.* **118**, 070502 (2017).  
 [69] A. J. Bray and M. A. Moore, *Phys. Rev. Lett.* **58**, 57 (1987).  
 [70] C. K. Thomas, D. A. Huse, and A. Middleton, *Phys. Rev. Lett.* **107**, 047203 (2011).  
 [71] D. Hu, P. Ronhovde, and Z. Nussinov, *Philosophical Magazine* **92**, 406 (2012).  
 [72] Z. Zhu, A. J. Ochoa, S. Schnabel, F. Hamze, and H. G. Katzgraber, *Phys. Rev. A* **93**, 012317 (2016).

## Supplemental Material for The Binomial Spin Glass

M.-S. Vaezi,<sup>1</sup> G. Ortiz,<sup>2,3</sup> M. Weigel,<sup>4</sup> and Z. Nussinov<sup>1,\*</sup>

<sup>1</sup> *Department of Physics, Washington University, St. Louis, MO 63160, USA*

<sup>2</sup> *Department of Physics, Indiana University, Bloomington, IN 47405, USA*

<sup>3</sup> *Department of Physics, University of Illinois, 1110 W. Green Street, Urbana, Illinois 61801, USA*

<sup>4</sup> *Applied Mathematics Research Centre, Coventry University, Coventry CV1 5FB, UK*

Below, we further provide a lightning overview of the problem that prompted the current investigation (Section A). We then elaborate on several aspects that were alluded to in the main text (Sections (B-H)).

### A. General Background and Motivation

The quintessential short-range Ising spin glass system is the Edwards-Anderson (EA) model, where at each lattice site  $\mathbf{x}$  lies a classical spin  $s_{\mathbf{x}} = \pm 1$ , that interacts with nearest-neighbor spins only [1]. In the *discrete* binary version, the random couplings may assume only the two values  $\pm J$ . Conversely, the couplings are *continuous* random Gaussian variables in the *continuous* EA model. While the extensive ground state degeneracy is well established for various binary distributions, the situation for the continuous EA model has been mired by controversy. Parisi’s tour de force solution [2] led to insights concerning the extensive nature of the ground state entropy of the infinite-range Sherrington-Kirkpatrick (SK) model [3]. The latter harbors a plethora of distinct thermodynamic states [4–6]. A measure of similarity between disparate thermodynamic states is provided by the well-known “overlap function” [4, 7, 8]  $q_{rr'} = \frac{1}{N} \sum_{\mathbf{x}} \langle s_{\mathbf{x}} \rangle_r \langle s_{\mathbf{x}} \rangle_{r'}$ , where  $N$  is the total number of lattice sites, and its average over the probabilities  $W_r$  and  $W_{r'}$  of the realizations of the different pairs of states  $r$  and  $r'$  (the “overlap distribution function”),  $P(q) = \sum_{r,r'} q_{rr'} W_r W_{r'}$ . The SK model displays a cascade of different overlaps (an ultrametric structure [9]) and replica symmetry breaking wherein  $P(q)$  becomes nontrivial [10]. Standard ordered systems typically display a small number of symmetry related thermodynamic states (and zero temperature ground states) associated with a distribution  $P(q)$  that is a sum of simple delta functions. While the Parisi solution and various related (effective infinite dimension or infinite range) mean-field treatments raise the possibility of an exponentially large number of ground states, other considerations [4, 11–18] suggest that (similar to ferromagnets) in typical short-range spin glasses, there are only two symmetry related ground states. The understanding of this problem underlies our work. This question is not merely of academic importance; the behavior of real finite dimensional magnetic spin glass systems has long been of direct experimental pertinence, e.g., [19, 20].

We now explicitly define the standard EA model. Consider a general bipartite lattice (in any finite number of

dimensions  $d$ ) of size  $N$ , endowed with periodic boundaries, with an Ising spin  $s_{\mathbf{x}}$  at each lattice site  $\mathbf{x}$ . The EA spin glass Hamiltonian is given by

$$H = - \sum_{\langle \mathbf{xy} \rangle} J_{\mathbf{xy}} s_{\mathbf{x}} s_{\mathbf{y}} \equiv - \sum_{\alpha=1}^{\mathcal{L}} J_{\alpha} z_{\alpha}. \quad (\text{S1})$$

The summation in Eq. (S1) is over nearest-neighbor spins at sites  $\mathbf{x}$  and  $\mathbf{y}$  sharing the link  $\alpha = \langle \mathbf{xy} \rangle$ ,  $z_{\alpha} = \pm 1$ , and the total number of these links is  $\mathcal{L} = d \times N$ . In various standard Ising spin glass models, the spin couplings  $\{J_{\alpha}\}$  in Eq. (S1) are customarily drawn from one of several well studied distributions. For instance, in the “binary Ising spin glass model” [21], the couplings  $\{J_{\alpha}\}$  are random variables that assume the two values  $\pm 1$  with probabilities  $P(J_{\alpha} = 1) = p$ ,  $P(J_{\alpha} = -1) = 1 - p$  (i.e., a Bernoulli distribution). In the continuous EA model the couplings  $\{J_{\alpha}\}$  are drawn from a Gaussian distribution of vanishing mean and variance equals to unity.

### B. The trivial ground state pair given an assignment of link variables

Given the definition of the link variable  $z_{\alpha} \equiv s_{\mathbf{x}} s_{\mathbf{y}}$ , a moment’s reflection reveals that

$$s_{\mathbf{y}} = s_{\mathbf{x}} \prod_{\alpha \in \Gamma_{\mathbf{xy}}} z_{\alpha}, \quad (\text{S2})$$

where  $\Gamma_{\mathbf{xy}}$  is any path on the lattice, composed of nearest-neighbor links, joining site  $\mathbf{x}$  to site  $\mathbf{y}$ . Thus, with  $s_{\mathbf{y}}|_{\mathbf{s}}$  denoting the value of the spin at site  $\mathbf{y}$  in configuration  $|\mathbf{s}\rangle$ , we have that

$$s_{\mathbf{y}}|_{\mathbf{s}} = s_{\mathbf{x}}|_{\mathbf{s}} \prod_{\alpha \in \Gamma_{\mathbf{xy}}} z_{\alpha}|_{\mathbf{s}}, \quad s_{\mathbf{y}}|_{\mathbf{s}'} = s_{\mathbf{x}}|_{\mathbf{s}'} \prod_{\alpha \in \Gamma_{\mathbf{xy}}} z_{\alpha}|_{\mathbf{s}'}. \quad (\text{S3})$$

Now, if for all links  $\alpha$ , the values of  $z_{\alpha}$  are the same in both configurations  $|\mathbf{s}\rangle$  and  $|\mathbf{s}'\rangle$  (i.e., if  $\{z_{\alpha}\}|_{\mathbf{s}} = \{z_{\alpha}\}|_{\mathbf{s}'}$ ) then, trivially,

$$\prod_{\alpha \in \Gamma_{\mathbf{xy}}} z_{\alpha}|_{\mathbf{s}} = \prod_{\alpha \in \Gamma_{\mathbf{xy}}} z_{\alpha}|_{\mathbf{s}'}. \quad (\text{S4})$$

Taken together, Eqs. (S3) and (S4) imply that if, at a particular site  $\mathbf{x}$ , the spin configurations  $|\mathbf{s}\rangle$  and  $|\mathbf{s}'\rangle$  share the same value of the spin,  $s_{\mathbf{x}}|_{\mathbf{s}'} = s_{\mathbf{x}}|_{\mathbf{s}}$ , then the spins must be identical at all other lattices sites  $\mathbf{y}$ ,  $s_{\mathbf{y}}|_{\mathbf{s}'} = s_{\mathbf{y}}|_{\mathbf{s}}$ . This, however, leads to a contradiction as  $|\mathbf{s}'\rangle \neq |\mathbf{s}\rangle$ . Therefore, if two distinct spin configurations satisfy

condition (i) it must be that the respective spin values at any lattice site  $x$  are different,  $s_x|_{|s'\rangle} = -s_x|_{|s\rangle}$ . That is,

$$s_y|_{|s'\rangle} = -s_y|_{|s\rangle}, \quad \forall y. \quad (\text{S5})$$

Hence, if  $n_\alpha = 0, \forall \alpha$  in Eq. (4) of the main text, then there are, trivially, only two degenerate configurations ( $|s'\rangle \neq |s\rangle$ ) related by a global spin inversion. The above simple proof applies for arbitrary energy levels. Replicating, *mutatis mutandis*, the above argument to a general set of (non-necessarily vanishing) integers  $\{n_\alpha\}$  over all lattice links  $\alpha$ , illustrates that any set  $\{n_\alpha\}$  may correspond to exactly two unique spin configurations.

### C. Graphical Representation of the Constraints

In the main text we defined  $\text{Sat}_{|s\rangle}$  to be the set composed of *all* constraints  $C_j$  satisfying the relation  $\Delta E = E(s) - E(s') = 0$ , in Eq. (4) of the main text. We also defined the subset  $\overline{\text{Sat}}_{|s\rangle} \subset \text{Sat}_{|s\rangle}$ , comprising *all linearly independent* constraints. Here, we further introduce a restricted subset of constraints, that of *geometrically disjoint* and independent zero energy domain walls,  $\text{Sat}_{|s\rangle}^g \subset \text{Sat}_{|s\rangle}$ . The subset  $\text{Sat}_{|s\rangle}^g$  is defined by having no pair of different constraints on the coupling constants that involve links associated with the same lattice sites  $x$ .

In what follows, we provide a few simple examples illuminating the above definitions. To this end, we consider a  $5 \times 5$  square lattice with binomial couplings  $\{\mathcal{J}_\alpha^m\}$  (Fig. S1). We start with a random spin configuration  $|s\rangle$  (panel (a)). Panels (b) through (e), represent spin configurations  $|s'\rangle$  for which one or more spins are being flipped with respect to panel (a). The energy difference in each case can be easily calculated. For example,

$$\begin{aligned} \Delta E_{a,b} = E_a - E_b = & -2(\mathcal{J}_{19,14}^m n_{19,14} \\ & + \mathcal{J}_{19,18}^m n_{19,18} + \mathcal{J}_{19,20}^m n_{19,20} + \mathcal{J}_{19,24}^m n_{19,24}), \end{aligned} \quad (\text{S6})$$

gives the energy difference between spin configurations in panel (a) and (b). It is easy to see that  $n_{19,18} = n_{19,20} = n_{19,24} = 1$ , and  $n_{19,14} = -1$ . Following the same procedure we end up with,

$$\begin{aligned} \Delta E_{a,b} &= -2(-\mathcal{J}_{19,14}^m + \mathcal{J}_{19,18}^m + \mathcal{J}_{19,20}^m + \mathcal{J}_{19,24}^m), \\ \Delta E_{a,c} &= -2(\mathcal{J}_{8,3}^m + \mathcal{J}_{8,7}^m + \mathcal{J}_{8,9}^m + \mathcal{J}_{8,13}^m), \\ \Delta E_{a,d} &= -2(-\mathcal{J}_{7,2}^m + \mathcal{J}_{7,6}^m + \mathcal{J}_{8,3}^m + \mathcal{J}_{8,9}^m + \mathcal{J}_{8,13}^m \\ &\quad - \mathcal{J}_{12,11}^m - \mathcal{J}_{12,13}^m - \mathcal{J}_{12,17}^m), \\ \Delta E_{a,e} &= \Delta E_{a,b} + \Delta E_{a,d} \\ &= -2(-\mathcal{J}_{19,14}^m + \mathcal{J}_{19,18}^m + \mathcal{J}_{19,20}^m + \mathcal{J}_{19,24}^m \\ &\quad - \mathcal{J}_{7,2}^m + \mathcal{J}_{7,6}^m + \mathcal{J}_{8,3}^m + \mathcal{J}_{8,9}^m + \mathcal{J}_{8,13}^m \\ &\quad - \mathcal{J}_{12,11}^m - \mathcal{J}_{12,13}^m - \mathcal{J}_{12,17}^m). \end{aligned} \quad (\text{S7})$$

Now, assume  $C_1, C_2, C_3$ , and  $C_4$  are constraints associated with  $\Delta E_{a,b}, \Delta E_{a,c}, \Delta E_{a,d}, \Delta E_{a,e}$ , respectively. If these

constraints are satisfied, i.e.,  $\Delta E_{a,b} = \Delta E_{a,c} = \Delta E_{a,d} = \Delta E_{a,e} = 0$ , for certain coupling realizations, then they belong to the set  $\text{Sat}_{|s\rangle}$ . That is,  $C_1, C_2, C_3, C_4 \in \text{Sat}_{|s\rangle}$ .

To understand this better, consider the case  $m = 4$ . Since  $\mathcal{J}_\alpha^m \equiv \frac{1}{\sqrt{m}} \sum_{k=1}^m J_\alpha^{(k)}$ , the couplings  $\mathcal{J}_\alpha^4$  may acquire the values  $-2, -1, 0, 1, 2$ . In Fig. S2, we provide three examples of random coupling realizations. The spin configuration is the same as in panel (a) of Fig. S1. From Eq. (S7) and Fig. S2, we can see that, only  $C_3$  in panel (a), none in panel (b), and only  $C_1$  in panel (c) are satisfied.

In order to create the subset  $\overline{\text{Sat}}_{|s\rangle}$ , we should note that it is not necessarily unique, since we may have different linearly independent constraints that span the same set of conditions in  $\text{Sat}_{|s\rangle}$ . In addition to that, the satisfaction of constraints depends on the coupling realizations as well. For instance, if for a given realization,  $C_1, C_2$  and  $C_3$  are satisfied, trivially from Eq. (S7) (i.e.,  $\Delta E_{a,e} = \Delta E_{a,b} + \Delta E_{a,d}$ ),  $C_4$  is automatically satisfied. Therefore, for such cases,  $C_4$  is a linear combination of  $C_1$  and  $C_3$ , and one may define the subset  $\overline{\text{Sat}}_{|s\rangle}^{(I)}$  for which  $C_1, C_2, C_3 \in \overline{\text{Sat}}_{|s\rangle}^{(I)}$ , but  $C_4 \notin \overline{\text{Sat}}_{|s\rangle}^{(I)}$ . On the other hand, there exist some realizations for which  $\Delta E_{a,b} = -\Delta E_{a,d} \neq 0$ , but  $\Delta E_{a,e} = 0$ . Meaning,  $C_4$  is satisfied, however,  $C_1$  and  $C_3$  are not.

The geometrically disjoint constraints may also give rise to different subsets. For instance, from Fig. S1, one can trivially show that the pairs  $C_1, C_2$  and  $C_1, C_3$  are each geometrically disjoint, however,  $C_2$  and  $C_3$  are not. Therefore, we could define two different subsets  $\overline{\text{Sat}}_{|s\rangle}^{(I)}$  and  $\overline{\text{Sat}}_{|s\rangle}^{(II)}$  so that  $C_1, C_2 \in \overline{\text{Sat}}_{|s\rangle}^{(I)}$  and  $C_1, C_3 \in \overline{\text{Sat}}_{|s\rangle}^{(II)}$ .

These examples further illustrate the difference between  $\mathcal{M}$  and  $M(\{\mathcal{J}_\alpha^m\})$  in the main text, where  $\mathcal{M}$  is associated with the maximum number of linearly independent satisfied constraints, i.e., the cardinality of  $\overline{\text{Sat}}_{|s\rangle}$ , while  $M(\{\mathcal{J}_\alpha^m\})$  denotes the number of constraints satisfied for a particular realization of coupling constants. Trivially,  $M(\{\mathcal{J}_\alpha^m\}) \leq \mathcal{M}$ .






FIG. S1. Graphical representations of the constraints. Panel (a) represents a random spin configuration. Blue solid circles and red diamonds denote spin up and down, respectively. Flipping one or more spins at different sites of panel (a) would result in new spin configurations such as in panels (b) through (e) (e.g., the spin configuration of panel (b) is obtained from flipping the spin at site 19 of panel (a)). The dashed yellow dotted lines represent the links that contribute to the energy difference. The green dashed lines crossing such links correspond to a domain wall.

#### D. The meaning of Equation (5) of the main text

In Eq. (5) of the main text, we mentioned that the set  $\{|\bar{n}_1\bar{n}_2\cdots\bar{n}_M\rangle\}$  includes all of the spin configurations degenerate with  $|\mathbf{s}\rangle$ . We also pointed out that it may contain additional states not degenerate with  $|\mathbf{s}\rangle$ . The latter point is usually associated with the domain walls that are not geometrically disjoint (see section C). To accentuate this consider, e.g., a  $5 \times 5$  lattice with a given random spin configuration and coupling constants (see panel (a) of Fig. S3), in which  $U_{ba}$ ,  $U_{ca}$ , and  $U_{da}$  are spin flip operators leading, respectively, to zero energy domain walls around the sites 7, 18 and 19 (corresponding to panels (b),(c) and (d)).

From Fig. S3, the domain walls in panel (c) and (d) are not geometrically disjoint, where  $U_{ca}$  and  $U_{da}$  act on the *nearest neighbor* sites 18 and 19 such that the sign of the link connecting them, is altered by both op-

erators. In such a case, even though the two states  $U_{ca}|\mathbf{a}\rangle \equiv |\mathbf{c}\rangle$  and  $U_{da}|\mathbf{a}\rangle \equiv |\mathbf{d}\rangle$  are degenerate with  $|\mathbf{a}\rangle$ , the state  $U_{da}U_{ca}|\mathbf{a}\rangle \equiv |\mathbf{e}\rangle$  (i.e., from panel (e),  $U_{ea} = U_{da}U_{ca}$ ) is not degenerate with  $|\mathbf{a}\rangle$ . One should note that in general this might not be true. That is, for some coupling realizations the state  $|\mathbf{e}\rangle$  can be degenerate with  $|\mathbf{a}\rangle$ .

By contrast, the two spin flip operators  $U_{ba}$  and  $U_{da}$  associated with the geometrically disjoint domain walls in panel (b) and (d), respectively, do not alter the signs of any common links. Therefore, the state  $U_{da}U_{ba}|\mathbf{a}\rangle \equiv |\mathbf{f}\rangle$  (i.e., from panel (f),  $U_{fa} = U_{da}U_{ba}$ ) is degenerate with  $|\mathbf{a}\rangle$ .




FIG. S2. Three examples of coupling realizations for the binomial model with  $m = 4$  (i.e.,  $\mathcal{J}_\alpha^4 = -2, -1, 0, 1, 2$ ). The numbers in green (brown) color provide the values of horizontal (vertical) coupling constants.

### E. The ground state entropy is bounded by the entropy of a random energy level

In deriving the bound of Eq. (11) of the main text, we assumed that no information other than the probability distribution  $P(\{\mathcal{J}_\alpha^m\})$  is provided. The configuration  $|\mathfrak{s}\rangle$  that we considered in the main text was an arbitrary random state. We next consider a more sophisticated problem. Suppose that the coupling constants are drawn from a binomial distribution and that once chosen a *ground state configuration*  $|\mathfrak{s}\rangle$  is given (i.e., the values of the spins  $s_x$  at all sites  $x$  in this ground state are provided). We then calculate the average of Eq. (7) of the main text with the condition that the (otherwise random binomial) coupling constants admit the particular configuration  $|\mathfrak{s}\rangle$  as a *ground state*. When applicable, the fact that  $|\mathfrak{s}\rangle$  is a ground state may generally yield nontrivial constraints on the coupling constants  $\{J_\alpha^{(k)}\}$  (recall that  $\mathcal{J}_\alpha^m \equiv \frac{1}{\sqrt{m}} \sum_{k=1}^m J_\alpha^{(k)}$ ). In such a situation, given the configuration  $|\mathfrak{s}\rangle$ , we may not simply use the initial binomial distribution for the coupling constants.

We now trivially demonstrate that if the energy density associated with the high temperature limit is unique then Eq. (11) of the main text will constitute an upper bound on the average ground state entropy density even if such information was provided for each realization of  $\{J_\alpha^{(k)}\}$ . This assertion follows as the entropy  $S_\ell(\{J_\alpha^{(k)}\})$  associated with any energy  $E = E_\ell$  is typically larger than the ground state entropy,

$$S_0 \leq S_\ell. \quad (\text{S8})$$

The proof of Eq. (S8) is rather elementary and relies on a trivial symmetry of the spectrum. Let us denote the two sublattices forming the large bipartite lattice by  $A$  and  $B$ . If we flip all spins in sublattice  $A$  (i.e.,  $s_{x \in A} \rightarrow -s_{x \in A}$ ) and do not alter those in sublattice  $B$  ( $s_{y \in B} \rightarrow s_{y \in B}$ ), then all

nearest-neighbor links (i.e., the products  $s_x s_y$  for nearest neighbor sites  $x$  and  $y$ ) on the original lattice change their sign,  $z_\alpha \rightarrow -z_\alpha$ . This single sublattice spin inversion constitutes a one-to-one mapping of the Ising spin states, that changes the sign of the total energy ( $E \rightarrow -E$ ). We may thus conclude that as a function of the energy  $E$ , the entropy density  $\mathcal{S} = S(\{J_\alpha^{(k)}\})/N$  for a system with fixed couplings  $\{J_\alpha^{(k)}\}$  satisfies the simple relation  $\mathcal{S}(E_\ell) = \mathcal{S}(-E_\ell)$  where  $E_\ell$  is the energy of the  $\ell$ -th level. It follows that the energy  $E = 0$  is an extremum of the entropy density  $\mathcal{S}(E) \equiv \mathcal{S}(E_\ell)$ . Consequently, for any fixed couplings  $\{J_\alpha^{(k)}\}$ ,

$$\frac{1}{T} = N \frac{\partial \mathcal{S}}{\partial E} \geq 0. \quad (\text{S9})$$

(The factor of  $N$  appears in the above equation since  $\mathcal{S}$  is the entropy density). Thus,  $E \leq 0$  for any positive temperature  $T$ . In what follows we discuss what occurs if there is a *unique* high temperature limit for each set of coupling constants. In such a case, the entropy density  $\mathcal{S}(E)$  (averaged over all realization of the coupling constants) is maximal at  $E = 0$ . The semi-positive definite nature of the derivative in Eq. (S9) implies (as in all common systems satisfying the third law of thermodynamics) that the entropy is lowest at  $T = 0$ . Since the state  $|\mathfrak{s}\rangle$  for which we performed the analysis was arbitrary (and corresponds to an energy  $E_{|\mathfrak{s}\rangle}$  for which the entropy density is greater than or equal to that of the ground state), we see that Eq. (S9) must hold even if information is provided as to the explicit ground state configuration  $|\mathfrak{s}\rangle$  for each particular realization of the couplings  $\{J_\alpha^{(k)}\}$ . We thus observe that even if given such additional information, the ground state entropy density must satisfy the bound of Eq. (11) of the main text.




FIG. S3. Panel (a) represents a random spin configuration with some given coupling constants. Blue solid circles and red diamonds denote spin up and down, respectively. The numbers in green (brown) color provide the values of horizontal (vertical) coupling constants. Flipping one or more spins at different sites of panel (a) would result in new spin configurations such as in panels (b) through (f). The dashed yellow dotted lines represent the links that contribute to the energy difference. The green dashed lines crossing such links correspond to a domain wall. Please note that the values associate with different links in each panel is the same as in panel (a).

### F. Asymptotic Scaling of the Entropy Density

We now motivate a scaling that suggests that the rigorous bound of Eq. (11) of the main text leads to Eq. (12) as an approximate asymptotic relation for large  $N$  and  $m$ . In Section C of this supplemental material, we defined the subset  $\overline{\text{Sat}}_{|\mathbf{s}}^{\mathbf{g}} \subset \text{Sat}_{|\mathbf{s}}$  composed of geometrically disjoint constraints. If there are  $n_g$  such constraints (or associated zero energy domain walls when these constraints are satisfied) then the degeneracy will be trivially bounded from below by  $2^{n_g}$ . This bound is established by noting that, since no spin is common to two domain walls, all of the spins in each of these  $n_g$  domain walls may be flipped independently of all others. When applied to domain walls in  $\overline{\text{Sat}}_{|\mathbf{s}}^{\mathbf{g}}$  then, in the notation of Eq. (5) of the main text, each binary string of length  $n_g$  will correspond to a different configuration that is degenerate with the reference state  $|\mathbf{s}\rangle$ . This is to be contrasted with the set of zero energy domain walls  $\text{Sat}_{|\mathbf{s}}$  for which

various binary strings of the form of Eq. (5) may correspond to states that are *not* degenerate with  $|\mathbf{s}\rangle$ . As  $m$  grows, by Eq. (8) of the main text, both the number of satisfied constraints and the number of independent zero energy domain walls may diminish as  $1/\sqrt{m}$ . When fewer walls appear in  $\overline{\text{Sat}}_{|\mathbf{s}}$ , it may become increasingly rare for different walls in this subset to share the same lattice sites. If this occurs then, for large  $m$ , we will have the asymptotic relation  $\overline{\text{Sat}}_{|\mathbf{s}}^{\mathbf{g}} \sim \text{Sat}_{|\mathbf{s}}$ . In such a case, in the large  $N$  limit,  $\mathcal{S} \sim n_g/N \ln 2$ . The number  $n_g$  and the probability of these zero energy domain walls decay, for  $m \gg 1$ , as  $1/\sqrt{m}$  (or  $1/\sqrt{m'}$  for  $m' \gg 1$ ). Similarly, if a finite fraction of the  $M$  domain walls in  $\overline{\text{Sat}}_{|\mathbf{s}}$  does not remain geometrically disjoint such that, asymptotically, one may only generate  $q^M$  (with  $q < 2$ ) degenerate states (Eq. (5)) given  $M$  independent domain walls, then  $\mathcal{S} \sim \frac{M}{N} \ln q$ . Either way, we anticipate that, in the thermodynamic limit, Eq. (12) of the main text will hold.

### G. One-dimensional Binomial Spin Glass

Let us start with the simplest one-dimensional binomial spin glass system (which by a simple change of variables ( $s_x \rightarrow s'_x \equiv s_x \prod_{u < x} \text{sign}(\mathcal{J}_{u,u+1}^m)$ ) may be transformed onto a random Ising ferromagnet with couplings  $|\mathcal{J}_{x',x'+1}^m|$ ). Here, the ground state energy  $E_0 = -\sum_x |\mathcal{J}_{x,x+1}^m|$ . In an open chain of  $N$  sites, the lowest excitation consists of identifying the weakest link,  $|\mathcal{J}_{x',x'+1}^m| \equiv \min_x \{|\mathcal{J}_{x,x+1}^m|\}$  and flipping all spins  $s_x \rightarrow -s_x$  for which  $x > x'$  (or consistently doing the same thing and only flipping all spins to the left of  $x'$ ); this generates a state that has an energy  $E_0 + \Delta E_{\min}$  with  $\Delta E_{\min} = 2|\mathcal{J}_{x',x'+1}^m|$ . (On a periodic chain, we may similarly identify the two weakest links and flip all spins lying between those two links leading to an energy cost  $\Delta E_{\min}$  that is twice the sum of the moduli of these two weakest links.) Calculations of the density of states and all ensuing thermodynamic properties are trivial [22]. For instance, the disorder averaged entropy in the low temperature,  $T \ll 1$ , limit of the binary model is  $[S_{m=1}(T)] \sim k_B(\ln 2 + (N-1)(1+2\beta)e^{-2\beta})$ , with  $\beta = 1/(k_B T)$ . The exponential suppression becomes  $e^{-2\beta/\sqrt{m}}$  and  $e^{-4\beta/\sqrt{m}}$  for odd and even  $m$ , respectively. Thus the excitation gap scales as  $m^{-1/2}$  (yet differently for odd and even  $m$ ). By contrast, the low- $T$  entropy of the continuum model is  $[S_{m \rightarrow \infty}(T)] \sim k_B(\ln 2 + \frac{N-1}{\sqrt{2\pi}}(k_B T - \frac{(k_B T)^3}{8}))$ , indicating the vanishing of the spectral gap in the thermodynamic limit. In that limit, these lowest excitations differ, relative to the ground state, by an extensive number of flipped spins.

### H. Distribution of excitations

Given any *ground state* configuration on a hypercubic lattice in  $d$  dimensions, one may compute the probability distribution for excitations of energy  $\Delta E_x = |\Delta E_x| = 2 \sum_y n_{xy} \mathcal{J}_\alpha^m$  generated by flipping a single spin  $x$ . Here, the sum is over all sites  $y$  that are nearest neighbor of site  $x$  and  $n_{xy} = -\text{sign}(s_x s_y) = \pm 1$ . Given the probability distribution for the links  $\{\mathcal{J}_\alpha^m\}$ , one may compute the probability distribution associated with a finite sum of these links  $2 \sum_y n_{xy} \mathcal{J}_\alpha^m$  in the ground state. The latter sum is that over a finite number of links (with bounded mean and variance) and thus for any  $\epsilon > 0$  (no matter how small), the probability that  $|\Delta E_x| < \epsilon$  is strictly smaller than unity. In order for the system to have a spectral gap that is greater than  $\epsilon$ , it must be that for each of the  $N$  lattice sites  $x$ , the energy penalty  $|\Delta E_x| > \epsilon$ . Given that the condition  $|\Delta E_x| > \epsilon$  must, in the thermodynamic limit, be satisfied an infinite number of times, while for any single  $x$  the probability that this condition is satisfied is strictly smaller than one, it is essentially impossible to have a gap larger than any arbitrary positive number  $\epsilon$ . From this, it follows that the gapless local excitations must be appear. If the local energy penal-




FIG. S4. Distribution of defect energies  $|\Delta E|$  for the 2D Gaussian system and a number of different system sizes. The curves collapse if rescaled by  $L^{-\theta}$  with the value  $\theta \approx -0.28$  of the stiffness exponent.

ties in the ground state were independent of one another then the probability that all local flips result in an energy penalty larger than  $\epsilon$  would be the product of the probabilities of having  $|\Delta E_x| > \epsilon$  for all sites  $x$ . Although the local flips are not independent of one another (since they all relate to flips relative to the same special state—the ground state), it seems highly unlikely  $|\Delta E_x| > \epsilon$  for all  $x$  when the probability of having a local energy penalty larger than  $\epsilon$  for any single  $x$  is strictly smaller than one.

We now explicitly discuss a measure that, in general dimensions, may provide physical insight – the *distribution* of such individual defect energies (i.e., the distribution of domain wall energies in our binomial Ising spin system). In Fig. S4, we plot this distribution in the continuous  $m = \infty$  Gaussian limit. If  $f(\epsilon, \tilde{l})$  denotes the cumulative probability that the energy penalty of a domain wall (of size  $\tilde{l}$ ) is smaller than  $\epsilon$ , then the probability that amongst  $\mathcal{N}_{\tilde{l}}$  independent domain walls, no single domain wall entails an energy cost lower than  $\epsilon$  will be bounded from above by  $e^{-f(\epsilon, \tilde{l})\mathcal{N}_{\tilde{l}}}$  as we briefly elaborate on now. Since, by definition,  $f(\epsilon, \tilde{l})$  is the cumulative probability that the energy cost of a random wall of size  $\tilde{l}$  is smaller than  $\epsilon$  (i.e.,  $\text{Prob.}(|\Delta E| \leq \epsilon) = f(\epsilon, \tilde{l})$ ), the probability that amongst  $\mathcal{N}_{\tilde{l}}$  independent domain walls, we explicitly have that the probability that no single domain wall has an energy cost larger than  $\epsilon$  is, trivially,  $[\text{Prob.}(|\Delta E| > \epsilon)]^{\mathcal{N}_{\tilde{l}}} = (1 - f(\epsilon, \tilde{l}))^{\mathcal{N}_{\tilde{l}}} \leq e^{-\mathcal{N}_{\tilde{l}}f(\epsilon, \tilde{l})}$  (where we invoked  $e^{-f} \geq (1 - f)$  for all  $f \geq 0$ ). For small  $f \rightarrow 0^+$  (associated with  $\epsilon \rightarrow 0^+$  in  $d \geq 3$ ), this general inequality is replaced by an equality (i.e.,  $[\text{Prob.}(|\Delta E| > \epsilon)]^{\mathcal{N}_{\tilde{l}}} = e^{-\mathcal{N}_{\tilde{l}}f(\epsilon, \tilde{l})}$ ).

Thus, if the area ( $d = 2$ ) or volume ( $d = 3$ ) of the entire lattice is  $\|\Lambda\|$ , then whenever the sum

$$\lim_{\epsilon \rightarrow 0^+} \lim_{\tilde{l}_0 \rightarrow \infty} \lim_{N \rightarrow \infty} \sum_{\|\Lambda\|^{1/d} - \tilde{l}_0 \geq \tilde{l} \geq \tilde{l}_0} f(\epsilon, \tilde{l}) \mathcal{N}_{\tilde{l}} = \infty \quad (\text{S10})$$

then gapless (or degenerate) states of diverging  $\tilde{l}$  may

appear. This is so because flipping all of the spins links one ground state to its conjugate. The inequality  $||\Lambda||^{1/d} - \tilde{l}_0 \geq \tilde{l} \geq \tilde{l}_0$  in Eq. (S10) means that the an extensive number of spin flips is needed to connect a given spin configuration to either of the two members of the degenerate ground state pair.

Since  $\theta_{d=2} < 0$  then (as is further underscored in the full distribution of Fig. S4), in two dimensions nearly all large domain walls entail a *vanishing* energy penalty. In  $d = 2$ ,  $\lim_{\epsilon \rightarrow 0^+} \lim_{\tilde{l} \rightarrow \infty} f(\epsilon, \tilde{l}) = 1$  and the probability of obtaining, in the thermodynamic limit, degenerate states that differ by an extensive number of flipped spins is unity. The existence of gapless states in  $d = 2$  is hardly surprising; such gapless states may be trivially constructed by the insertion of random domain walls of

divergent size into a ground state. Indeed, in  $d = 2$  (where the typical energy cost  $\mathcal{O}(\tilde{l}^\theta)$  vanishes as  $\tilde{l} \rightarrow \infty$ ), knowledge of the detailed distribution of the energy cost as a function of the domain wall size  $\tilde{l}$  is unnecessary for establishing gapless states. However, in  $d \geq 3$  (where  $\theta_d > 0$ ), the lowest energy states are related to *the asymptotic low energy limit* of the domain wall energy distribution (a distribution that, in these higher dimensions, is associated with a divergent average energy  $\mathcal{O}(\tilde{l}^{\theta_d})$  when  $\tilde{l} \rightarrow \infty$ ). A gap (for states that differ from one another by an extensive number of flipped spins) is potentially possible if the sum of Eq. (S10) vanishes. Thus, we stress that in  $d \geq 3$ , knowledge of the cumulative probability distribution  $f(\epsilon, \tilde{l})$  can be of paramount importance. We reserve the analysis of the  $d = 3$  domain wall energy distribution for future work.

- 
- [1] S. Edwards and P. W. Anderson, J. Phys. F **5**, 965 (1975).
  - [2] G. Parisi, Phys. Rev. Lett. **43**, 1754 (1979); G. Parisi, J. Phys. A **13**, L115 (1980); G. Parisi, J. Phys. A **13**, 1101 (1980); G. Parisi, J. Phys. A **13**, 1887 (1980); G. Parisi, Phys. Rev. Lett. **50**, 1946 (1983).
  - [3] D. Sherrington and S. Kirkpatrick Phys. Rev. Lett. **35**, 1792 (1975).
  - [4] D. L. Stein and C. M. Newman, *Spin Glasses and Complexity* (Princeton University Press, 2013).
  - [5] F. Guerra and F. L. Toninelli, Comm. Math. Phys. **230**, 71 (2002);
  - [6] M. Talagrand, *Spin Glasses: A Challenge to Mathematicians*, Springer-Verlag (2003).
  - [7] M. Mezard, G. Parisi, N. Sourlas, G. Toulouse, and M. Virasoro, Phys. Rev. Lett., **52**, 1156 (1984).
  - [8] G. Parisi, J. Phys. **45**, 843 (1984).
  - [9] R. Rammal, G. Toulouse, and M. A. Virasoro, Rev. Mod. Phys. **58**, 765 (1986).
  - [10] S. Franz, M. Mezard, G. Parisi, and L. Peliti, Phys. Rev. Lett. **81**, 1758 (1998).
  - [11] W. L. MacMillan Phys. Rev. B **31**, 340 (1985).
  - [12] A. J. Bray and M. A. Moore, Phys. Rev. B **31**, 631 (1985).
  - [13] R. G. Calfish and J. R. Banavar, Phys. Rev. B **32**, 7617 (1985).
  - [14] M. A. Moore and A. J. Bray, J. Phys C: Solid State Phys. **18**, L699 (1985).
  - [15] D. S. Fisher and D. A. Huse, J. Phys. A: Math. Gen. **20**, L1005 (1987).
  - [16] D. A. Huse and D. S. Fisher, J. Phys. A: Math. Gen. **20**, L997 (1987).
  - [17] C. M. Newman and D. L. Stein, Phys. Rev. Lett. **84**, 3966 (2000).
  - [18] C. M. Newman and D. L. Stein, Commun. Math. Phys. **224**, 205 (2001).
  - [19] J. A. Mydosh, *Spin Glasses: An Experimental Introduction* (Taylor and Francis, London, Washington D. C., 1993).
  - [20] V. Cannella and J. A. Mydosh Phys. Rev. B **6**, 4220 (1972).
  - [21] J. Lukic, A. Galluccio, E. Marinari, O. C. Martin, and G. Rinaldi, Phys. Rev. Lett. **92**, 117202 (2004).
  - [22] In order to attain any state, a number of links  $\{\mathcal{J}_\alpha^m\}$  must be flipped relative to a ground state. The probability for obtaining a particular energy amounts to a convolution on the probability distribution of the links. The latter convolution readily becomes a trivial product after Fourier transformation.