

Enumeration of words that contain the pattern 123 exactly once

Mingjia Yang

Abstract: Enumeration problems related to words avoiding patterns as well as permutations that contain the pattern 123 exactly once have been studied in great detail. However, the problem of enumerating words that contain the pattern 123 exactly once is new and will be the focus of this paper. Previously, Doron Zeilberger provided a shortened version of Alexander Burstein’s combinatorial proof of John Noonan’s theorem that the number of permutations with exactly one 321 pattern is equal to $\frac{3}{n} \binom{2n}{n+3}$. Surprisingly, a similar method can be directly adapted to words. We are able to use this method to find a formula enumerating the words with exactly one 123 pattern. Further inspired by Nathaniel Shar and Zeilberger’s paper on generating functions enumerating 123-avoiding words with r occurrences of each letter, we examine the algebraic equations for generating functions for words with r occurrences of each letter and with exactly one 123 pattern.

1. Introduction

Recall that a word $w = w_1 \dots w_k$ is an ordered list of letters on some alphabet. To say a word contains a pattern (a certain permutation of $\{1, \dots, m\}$) σ is to say there exist $1 \leq i_1 < i_2 < \dots < i_m \leq k$ such that the subword $w_{i_1} \dots w_{i_m}$ is *order isomorphic* to σ (for example, 246 is order isomorphic to 123). A word avoids the pattern σ if it does not contain it.

For a lucid history on the study of *forbidden patterns*, readers are welcome to refer to the introduction of Shar and Zeilberger’s paper [SZ]. We also found the organization of [SZ] to be nice and suitable for the development of our paper, and we will be more or less following that.

In the present article, we say that a word w in the alphabet $\{a_1, a_2, \dots, a_n\}$ ($a_1 < a_2 < \dots < a_n$) is *associated* with the list $[l_1, \dots, l_n]$ if w has l_i many a_i ’s in it, for i from 1 to n . For example, 231113233 is a word associated with the list $[3, 2, 4]$, and 223344 is a word associated with the list $[2, 2, 2]$. Without specifying, our default alphabet will be $\{1, \dots, n\}$ for some $n \geq 1$.

In the second section, we will generalize Zeilberger’s bijective proof [Z1] (a shortened version of Alexander Burstein’s elegant combinatorial proof [Bu]) that the number of permutations of $\{1, \dots, n\}$ that contain the pattern 321 exactly once equals $\frac{3}{n} \binom{2n}{n+3}$ and apply it to words. Although no closed form formula was found, we have a summation whose summands are expressions involving enumeration of 123-avoiding words (for details, see Theorem 1).

In the third section, we will study, using ideas from the second section, how to extend Shar and Zeilberger's work [SZ] on generating functions enumerating 123-avoiding words (with r occurrences of each letter) to words (with r occurrences of each letter) having exactly one pattern 123. More precisely, for every positive integer r , Shar and Zeilberger found an algorithm for finding the defining algebraic equation for the ordinary generating function enumerating 123-avoiding words of length rn where each of the n letters of $\{1, 2, \dots, n\}$ occurs exactly r times.

We will present an algorithm for finding an analogue of that, that is, a defining algebraic equation for the ordinary generating function enumerating words of length rn where each of the n letters of $\{1, 2, \dots, n\}$ occurs exactly r times, now with exactly one pattern 123. We used the same (as in Shar and Zeilberger's paper [SZ]) memory-intensive, and exponential time, Buchberger's algorithm for finding Gröbner bases, and our computer (running Maple) found the defining algebraic equation for $r = 2$:

$$\begin{aligned} x^4(x+4)^2 F^4 + 2x^3(x+4)(11x+23)F^3 - 4x(3x^4 - 10x^3 - 97x^2 - 146x + 1)F^2 \\ + (-168x^4 - 840x^3 - 744x^2 + 336x - 24)F + 144x^3(x+2) = 0. \end{aligned}$$

This took about a second. The minimal algebraic equation for $r = 3$ has 12 as the highest power for F and the computation took about 20 seconds. Interested readers can find it on the website accompanying this paper: <http://sites.math.rutgers.edu/~my237/0ne123>. The case when $r = 4$ already took too long to compute (more than a month).

Now, let $a_r(n)$ be the number words of length rn where each of the n letters of $\{1, 2, \dots, n\}$ occurs exactly r times, with exactly one pattern 123. In the last section of the present article, we will use the Maple package **SCHUTZENBERGER** to derive recurrence relations for our sequences. Having obtained the defining algebraic equations of the generating functions for $a_r(n)$ in the cases $r = 2$ and $r = 3$, Manuel Kauers kindly helped us in finding the asymptotics for our sequences $a_2(n)$ and $a_3(n)$ (thanks to Kauers, the constants in front are fully rigorous and were computed via a step by step procedure; for details, please refer to [KP]):

$$\begin{aligned} a_2(n) &= \frac{3(13 - \sqrt{21})}{49} \cdot \frac{1}{\sqrt{\pi}} \cdot 12^n \cdot n^{-3/2} \cdot (1 + O(n^{-1})), \\ a_3(n) &= \frac{-7 + 6\sqrt{7}}{56} \cdot \frac{1}{\sqrt{\pi}} \cdot 32^n \cdot n^{-3/2} \cdot (1 + O(n^{-1})). \end{aligned}$$

We noticed how similar these are to the asymptotics of the sequences enumerating 123-avoiding words with r occurrences of each letter, given on page 8 of [SZ], and we have a similar conjecture as on page 3 of [SZ] that $a_r(n)$ is asymptotically $C_r \cdot ((r+1)2^r)^n \cdot n^{-3/2}$, where C_r is a constant depending on r (possibly $\frac{1}{\sqrt{\pi}}$ times a fraction of expressions involving square roots).

2.1. Zeilberger's shortened version of Burstein's proof on permutations containing 321 exactly once

In a paper published in 2011 [Bu], Burstein gave an elegant combinatorial proof of John Noonan's theorem [N] that the number of permutations of $\{1, \dots, n\}$ that contain the pattern 321 exactly once equals $\frac{3}{n} \binom{2n}{n+3}$. Zeilberger [Z1] was able to shorten Burstein's proof by using a bijection between a permutation with exactly one pattern 321, denoted as $\pi_1 c \pi_2 b \pi_3 a \pi_4$ ($a < b < c$), with the pair $(\pi_1 b \pi_2 a, c \pi_3 b \pi_4)$ where $\pi_1 b \pi_2 a$ is a 321-avoiding permutation of $\{1, \dots, b\}$ and $c \pi_3 b \pi_4$ is a 321-avoiding permutation of $\{b, \dots, n\}$. Readers are encouraged to read Zeilberger's proof as a motivation and warm-up. Below we will see how we can use the same logic and apply it to words.

2.2. Extension to words

Theorem 1. Let $A(l_1, \dots, l_n)$ be the number of 123-avoiding words associated with the list $[l_1, \dots, l_n]$. Then the number of words associated with list $[l_1, \dots, l_n]$ that contain the pattern 123 exactly once is:

$$\sum_{b=2}^{n-1} \sum_{j=0}^{l_b-1} (A(l_1, \dots, l_{b-1}, j+1) - A(l_1, \dots, l_{b-1}, j)) \cdot (A(l_b - j, l_{b+1}, \dots, l_n) - A(l_b - j - 1, l_{b+1}, \dots, l_n)).$$

Before we start the proof, let us first define what a *good* pair of words is. Fix any $2 \leq b \leq n-1$ and a list $[l_1, \dots, l_n]$. For any $0 \leq j \leq l_b - 1$, the pair of words (σ_1, σ_2) is *good* if σ_1 is a 123-avoiding word in $\{1, \dots, b\}$ associated with the list $[l_1, \dots, l_{b-1}, j+1]$ that does not start with b and σ_2 is a 123-avoiding word in $\{b, \dots, n\}$ associated with the list $[l_b - j, l_{b+1}, \dots, l_n]$ that does not end with b . We will also say σ_i ($i = 1, 2$) is good if it belongs to a good pair (σ_1, σ_2) .

Proof. Any word w associated with the list $[l_1, l_2, \dots, l_n]$ with exactly one pattern 123 can be written as $\pi_1 a \pi_2 b \pi_3 c \pi_4$ ($a < b < c$), where abc is the unique 123 pattern. All entries to the left of b , except a , must be greater than or equal to b , and all the entries to the right of b , except for c , must be smaller than or equal to b . Also, π_2 and π_3 must not contain any b 's, otherwise there will be another 123 pattern. Observe that $a \pi_3 b \pi_4$ is a word in $\{1, 2, \dots, b\}$ avoiding pattern 123 and does not start with b and $\pi_1 b \pi_2 c$ is a word in $\{b, b+1, \dots, n\}$ avoiding pattern 123 and does not end with b . Therefore $(a \pi_3 b \pi_4, \pi_1 b \pi_2 c)$ is a good pair.

We now verify that there is indeed a bijection from the set of words having exactly one pattern 123 to the set of good pairs (σ_1, σ_2) (for $2 \leq b \leq n-1$ and $0 \leq j \leq l_b - 1$).

Fix b and j ($2 \leq b \leq n-1$, $0 \leq j \leq l_b-1$). Given a word $\pi_1 a \pi_2 b \pi_3 c \pi_4$ (that has exactly one 123 pattern: abc), we can easily map it to a unique good pair $(a\pi_3 b \pi_4, \pi_1 b \pi_2 c)$ by first finding out what a, c are. This is easy since we have only one 123 pattern. Conversely, given a good pair (σ_1, σ_2) , we take the first letter of σ_1 as “ a ” and the leftmost occurrence of b as “ b ” and get π_3 and π_4 ($\sigma_1 = a\pi_3 b \pi_4$). Similarly, we take the last letter of σ_2 as “ c ” and the rightmost occurrence of b as “ b ” and get π_1 and π_2 ($\sigma_2 = \pi_1 b \pi_2 c$). Putting everything together we get a unique $\pi_1 a \pi_2 b \pi_3 c \pi_4$.

Now, for any b and j , the number of good σ_1 is $A(l_1, \dots, l_{b-1}, j+1) - A(l_1, \dots, l_{b-1}, j)$ and the number of good σ_2 is $A(l_b - j, l_{b+1}, \dots, l_n) - A(l_b - j - 1, l_{b+1}, \dots, l_n)$. Therefore the number of words $\pi_1 a \pi_2 b \pi_3 c \pi_4$ with exactly one pattern 123 is: $(A(l_1, \dots, l_{b-1}, j+1) - A(l_1, \dots, l_{b-1}, j)) \cdot (A(l_b - j, l_{b+1}, \dots, l_n) - A(l_b - j - 1, l_{b+1}, \dots, l_n))$. Summing over all b and j , we get the desired result.

Corollary 1. The number of words associated with list $[l_1, \dots, l_n]$ that contain exactly one pattern 123 is equal to the number of words associated with list $[l_n, \dots, l_1]$ that contain exactly one pattern 123.

Proof. By Theorem 1, the number of words with exactly one pattern 123 and associated with list $[l_1, \dots, l_n]$ is

$$\sum_{b=2}^{n-1} \sum_{j=0}^{l_b-1} (A(l_1, \dots, l_{b-1}, j+1) - A(l_1, \dots, l_{b-1}, j)) \cdot (A(l_b - j, l_{b+1}, \dots, l_n) - A(l_b - j - 1, l_{b+1}, \dots, l_n)) \quad (1)$$

and the number of words with exactly one pattern 123 and associated with list $[l_n, \dots, l_1]$ is

$$\sum_{b=2}^{n-1} \sum_{j=0}^{l_{n-b+1}-1} (A(l_n, \dots, l_{n-b+2}, j+1) - A(l_n, \dots, l_{n-b+2}, j)) \cdot (A(l_{n-b+1} - j, l_{n-b}, \dots, l_1) - A(l_{n-b+1} - j - 1, l_{n-b}, \dots, l_1)). \quad (2)$$

When $b = k$ ($2 \leq k \leq n-1$), the inner sum of (1) becomes

$$\sum_{j=0}^{l_k-1} (A(l_1, \dots, l_{k-1}, j+1) - A(l_1, \dots, l_{k-1}, j)) \cdot (A(l_k - j, l_{k+1}, \dots, l_n) - A(l_k - j - 1, l_{k+1}, \dots, l_n)) \quad (3)$$

while when $b = n - k + 1$ ($2 \leq k \leq n-1$, notice this is the “symmetric counterpart” of $b = k$), the inner sum of (2) becomes

$$\sum_{j=0}^{l_k-1} (A(l_n, \dots, l_{k+1}, j+1) - A(l_n, \dots, l_{k+1}, j)) \cdot (A(l_k - j, l_{k-1}, \dots, l_1) - A(l_k - j - 1, l_{k-1}, \dots, l_1)). \quad (4)$$

We only need to show (3) = (4) in order to show (1) = (2). Now notice that when $j = t$ ($0 \leq t \leq l_k - 1$), the summand of (3) is

$$(A(l_1, \dots, l_{k-1}, t+1) - A(l_1, \dots, l_{k-1}, t)) \cdot (A(l_k - t, l_{k+1}, \dots, l_n) - A(l_k - t - 1, l_{k+1}, \dots, l_n)) \quad (5)$$

and when $j = l_k - 1 - t$ (the “symmetric counterpart” of $j = t$), the summand of (4) is

$$(A(l_n, \dots, l_{k+1}, l_k - t) - A(l_n, \dots, l_{k+1}, l_k - t - 1)) \cdot (A(t+1, l_{k-1}, \dots, l_1) - A(t, l_{k-1}, \dots, l_1)). \quad (6)$$

After a small rearrangement, we can see (5) = (6) because of an important result that $A(l_1, \dots, l_n)$ is symmetric in its arguments (meaning we can re-order the list $[l_1, \dots, l_n]$ in any order we want and get the same number; for details, see [SZ], page 4). Therefore as j ranges from 0 to $l_k - 1$, we have (3) = (4). And as b ranges from 0 to $n - 1$, we have (1) = (2).

Corollary 2. Fix a list $L := [l_1, \dots, l_n]$. The number of words associated with L that contain exactly one pattern 123 is equal to the number of words associated with L that contain exactly one pattern 321.

Proof. Let S_1 be the set of words associated with $[l_1, \dots, l_n]$ that contain exactly one pattern 123 and S_2 be the set of words associated with $[l_n, \dots, l_1]$ that contain exactly one pattern 321. Take any $w_1 \in S_1$, we can map it to a word w_2 associated with $[l_n, \dots, l_1]$ by mapping letter i to letter $n - i + 1$ (for all i from 1 to n). For example, the word 121322 is mapped to 323122. Observe that w_2 must contain exactly one pattern 321, which occurs at the same location in w_2 as the location of the 123 pattern in w_1 . Therefore $w_2 \in S_2$. Clearly this is a bijection from S_1 to S_2 . So $|S_1| = |S_2|$. This along with Corollary 1 gives Corollary 2.

Remark.

One may wonder if the number of words associated with $[l_1, \dots, l_n]$ that contain exactly one pattern 123 is equal to the number of words associated with $[l_1, \dots, l_n]$ that contain exactly one pattern 132. This is not the case (if this were the case, we would have an analogue of the result that the 123-avoiding words associated with $[l_1, \dots, l_n]$ are equinumerous with the 132-avoiding words associated with $[l_1, \dots, l_n]$, see [Z2]). For example, the number of permutations of $\{1, 2, \dots, n\}$ ($n \geq 1$) that contain exactly one pattern 123 is $\frac{3}{n} \binom{2n}{n+3}$ [Z1] while the number of permutations of $\{1, 2, \dots, n\}$ ($n \geq 1$) that contain exactly one pattern 132 is $\binom{2n-3}{n-3}$ [B6].

3.1. Some crucial background and generating functions for words avoiding pattern 123

In the beautiful paper by Shar and Zeilberger [SZ], methods for finding the algebraic equation for the ordinary generating function enumerating 123-avoiding words of length rn , where each of the n letters of $\{1, 2, \dots, n\}$ occurs exactly r times were given. First we present some important definitions and results of that paper here.

For $0 \leq i \leq j \leq r-1$ and $n \geq 0$, let $W_r^{(i,j)}(n)$ be the set of 123-avoiding words of length $rn+i+j$, in the alphabet $\{1, 2, \dots, n, n+1, n+2\}$, with i occurrences of the letter 1, j occurrences of the $n+2$, and exactly r occurrences of the other n letters, and let $W_r^{(i,j)}$ be the union of $W_r^{(i,j)}(n)$ over all $n \geq 0$. Let $g_r^{(i,j)}(x)$ be the *weight enumerator* for $W_r^{(i,j)}$, with respect to the weight $w \rightarrow x^{\text{length}(w)}$. (Note that the $W_r^{(i,j)}$'s have the same weight enumerator if any two letters have i and j occurrences respectively, and the remaining letters each occurs exactly r times. For a detailed explanation, see [SZ], page 3-4.)

Shar and Zeilberger were able to find a system of $\binom{r+1}{2}$ equations for $g_r^{(i,j)}(x)$ ($0 \leq i \leq j \leq r-1$), with the convention that if $s > k$ then $g_r^{(s,k)} = g_r^{(k,s)}$:

$$g_r^{(i,j)}(x) = \delta_{i,0}\delta_{j,0} + x \sum_{t=0}^{r-1} g_r^{(i,t)}(x) g_r^{((r-t) \bmod r, (j-1) \bmod r)}(x) + \sum_{m=0}^{i-1} x^{m+1} g_r^{(i-m, j-1)}(x)$$

For example, in the case when $r = 2$, we would get the following system of equations:

$$g_2^{(0,0)}(x) = 1 + x g_2^{(0,0)}(x) g_2^{(0,1)}(x) + x g_2^{(0,1)}(x) g_2^{(1,1)}(x)$$

$$g_2^{(0,1)}(x) = x g_2^{(0,0)}(x)^2 + x g_2^{(0,1)}(x)^2$$

$$g_2^{(1,1)}(x) = x g_2^{(0,0)}(x) g_2^{(0,1)}(x) + x g_2^{(0,1)}(x) (1 + g_2^{(1,1)}(x))$$

Solving this system of equations in the three unknowns $g_2^{(0,0)}(x)$, $g_2^{(0,1)}(x)$, $g_2^{(1,1)}(x)$, we get the weight enumerators for $W_2^{(0,0)}$, $W_2^{(0,1)}$ and $W_2^{(1,1)}$.

Once we have the weight enumerators, we can easily get the corresponding generating functions by doing a little operation. For example, if we have an explicit expression for $g_2^{(0,0)}(x)$ ($g_2^{(0,0)}(x) = 1 + x^2 + 6x^4 + 43x^6 + 352x^8 + 3114x^{10} + \dots$), the corresponding generating function will be $f_2^{(0,0)}(x) = 1 + x + 6x^2 + 43x^3 + 352x^4 + 3114x^5 + \dots$ (that is, $f_2^{(0,0)}(x) = g_2^{(0,0)}(x^{1/2})$).

3.2. Extension to generating functions for words with exactly r occurrences of each letter, and with exactly one pattern 123

Definitions: Let $V_r(n)$ be the set of words in the alphabet $\{1, \dots, n\}$ with exactly r occurrences of each letter, and with exactly one pattern 123. Let V_r be $\bigcup_{n=0}^{\infty} V_r(n)$.

Let $h_r(x)$ be the weight enumerator for V_r (as always, with weight $w \rightarrow x^{\text{length}(w)}$) and let $f_r(x)$ be the corresponding generating function. We will be following the framework of [SZ], with two warm-up cases leading to the general case.

First warm-up: $r = 1$

Claim: $h_1(x) = (g_1^{(0,0)}(x) - xg_1^{(0,0)}(x) - 1)^2/x$.

Proof. Recall that $g_1^{(0,0)}(x) (= 1 + x + 2x^2 + 5x^3 + 14x^4 + 42x^5 + \dots)$ is the weight enumerator for 123-avoiding permutations on $\{1, \dots, n, \dots\}$. We prove this claim by showing that the coefficient of x^n ($n \geq 0$) on the right hand side is exactly the number of good pairs $(a\pi_3b\pi_4, \pi_1b\pi_2c)$ ($2 \leq b \leq n-1$), which equals to the number of permutations on $\{1, \dots, n\}$ with exactly one pattern 123 (by Zeilberger's proof [Z1]).

For any fixed b ($2 \leq b \leq n-1$), a good $a\pi_3b\pi_4$ would be a 123-avoiding permutation on $\{1, \dots, b\}$ that does not start with b . Similarly, a good $\pi_1b\pi_2c$ would be a 123-avoiding permutation on $\{b, \dots, n\}$ that does not end with b .

Note that the coefficient of x^b in $g_1^{(0,0)}(x) - xg_1^{(0,0)}(x) - 1$ is exactly the number of good $a\pi_3b\pi_4$. (The x in front of $g_1^{(0,0)}(x)$ corresponds to having b in front of a permutation, and the -1 corresponds to an empty permutation. We don't want either of these.)

Similarly, the coefficient of x^{n-b+1} in $g_1^{(0,0)}(x) - xg_1^{(0,0)}(x) - 1$ is the number of good $\pi_1b\pi_2c$. Multiplying the two, we have that the coefficient of x^{n+1} ($= x^b \cdot x^{n-b+1}$) in

$$(g_1^{(0,0)}(x) - xg_1^{(0,0)}(x) - 1)^2$$

is the number of good pairs $(a\pi_3b\pi_4, \pi_1b\pi_2c)$ (b ranges from 2 to $n-1$). Dividing by x , we get the coefficient of x^n in

$$(g_1^{(0,0)}(x) - xg_1^{(0,0)}(x) - 1)^2/x$$

is the number of good pairs $(a\pi_3b\pi_4, \pi_1b\pi_2c)$.

Second warm-up: $r = 2$

Claim: $h_2(x) = 2 \cdot (g_2^{(0,0)}(x) - xg_2^{(0,1)}(x) - 1)(g_2^{(0,1)}(x) - xg_2^{(0,0)}(x))/x$.

Proof. Recall that $g_2^{(0,0)}(x) (= 1 + x^2 + 6x^4 + 43x^6 + \dots)$ is the weight enumerator for 123-avoiding words on $\{1, 1, \dots, n, n, \dots\}$ (or equivalently, 123-avoiding words associated with $[2, 2, \dots]$) and $g_2^{(0,1)}(x) (= x + 3x^3 + 19x^5 + 145x^7 \dots)$ is the weight enumerator for 123-avoiding words on $\{1, 2, 2, 3, 3, \dots, n, n, \dots\}$. As in the first warm-up, we prove this claim by showing that the coefficient of x^{2n} ($n \geq 0$) on the right hand side is exactly the number of good pairs $(a\pi_3 b\pi_4, \pi_1 b\pi_2 c)$ ($2 \leq b \leq n - 1$), which equals to the number of words on $\{1, 1, \dots, n, n\}$ with exactly one pattern 123 (by the proof of Theorem 1).

For any b ($2 \leq b \leq n - 1$), we have the following two cases: either π_4 contains one b and π_1 contains no b or the other way around.

Case 1: π_4 contains one b and π_1 contains no b .

Then a good $a\pi_3 b\pi_4$ would be a 123-avoiding word on $\{1, 1, \dots, b, b\}$ that does not start with b . Similarly, a good $\pi_1 b\pi_2 c$ would be a 123-avoiding word on $\{b, b + 1, b + 1, \dots, n, n\}$ that does not end with b .

Note that the coefficient of x^{2b} in $g_2^{(0,0)}(x) - xg_2^{(0,1)}(x) - 1$ is exactly the number of good $a\pi_3 b\pi_4$. (The x in front of $g_2^{(0,1)}(x)$ corresponds to having b in front of a word, and the -1 corresponds to an empty word. We don't want either of these.)

Similarly, the coefficient of $x^{2(n-b)+1}$ in $g_2^{(0,1)}(x) - xg_2^{(0,0)}(x)$ is the number of good $\pi_1 b\pi_2 c$.

Multiplying the two, and let b range from 2 to $n - 1$, we have that the coefficient of x^{2n+1} ($= x^{2b} \cdot x^{2(n-b)+1}$) in

$$(g_2^{(0,0)}(x) - xg_2^{(0,1)}(x) - 1)(g_2^{(0,1)}(x) - xg_2^{(0,0)}(x))$$

is the number of good pairs $(a\pi_3 b\pi_4, \pi_1 b\pi_2 c)$ if the additional b is in π_4 . Dividing by x , we get the coefficient of x^{2n} in

$$(g_2^{(0,0)}(x) - xg_2^{(0,1)}(x) - 1)(g_2^{(0,1)}(x) - xg_2^{(0,0)}(x))/x$$

is the number of good pairs $(a\pi_3 b\pi_4, \pi_1 b\pi_2 c)$ if the additional b is in π_4 .

Case 2: π_1 contains one b and π_4 contains no b .

Then a good $a\pi_3b\pi_4$ would be a 123-avoiding word on $\{1, 1, \dots, b-1, b-1, b\}$ that does not start with b . A good $\pi_1b\pi_2c$ would be a 123-avoiding word on $\{b, b, \dots, n, n\}$ that does not end with b .

Now, the coefficient of x^{2b-1} in $g_2^{(0,1)}(x) - xg_2^{(0,0)}(x)$ is exactly the number of good $a\pi_3b\pi_4$. Similarly, the coefficient of $x^{2(n-b)+2}$ in $g_2^{(0,0)}(x) - xg_2^{(0,1)}(x) - 1$ is the number of good $\pi_1b\pi_2c$.

Multiplying the two, we have that the coefficient of x^{2n+1} ($= x^{2b-1} \cdot x^{2(n-b)+2}$) in

$$(g_2^{(0,1)}(x) - xg_2^{(0,0)}(x))(g_2^{(0,0)}(x) - xg_2^{(0,1)}(x) - 1)$$

is the number of good pairs $(a\pi_3b\pi_4, \pi_1b\pi_2c)$ (b ranges from 2 to $n-1$) if the additional b is in π_1 . Dividing by x , we get the coefficient of x^{2n} in

$$(g_2^{(0,1)}(x) - xg_2^{(0,0)}(x))(g_2^{(0,0)}(x) - xg_2^{(0,1)}(x) - 1)/x$$

is the number of good pairs $(a\pi_3b\pi_4, \pi_1b\pi_2c)$ if the additional b is in π_1 .

Therefore the coefficient of x^{2n} in $2 \cdot (g_2^{(0,0)}(x) - xg_2^{(0,1)}(x) - 1)(g_2^{(0,1)}(x) - xg_2^{(0,0)}(x))/x$ is the number of good pairs $(a\pi_3b\pi_4, \pi_1b\pi_2c)$, which is equal to the number of words in $\{1, 1, \dots, n, n\}$ that have exactly one pattern 123. That is, we have shown the weight enumerator for V_2 is as claimed to be. To get the generating function $f_2(x)$ for V_2 we simply let $f_2(x) = h_2(x^{1/2})$.

Readers are welcome to compare $h_2(x)$ with the earlier formula in the case when $l_i = 2$ ($1 \leq i \leq n$):

$$\sum_{b=2}^{n-1} \sum_{j=0}^1 \underbrace{(A(j+1, 2, 2, \dots, 2) - A(j, 2, 2, \dots, 2))}_{b-1 \text{ many } 2\text{'s}} \cdot \underbrace{(A(2-j, 2, 2, \dots, 2) - A(1-j, 2, 2, \dots, 2))}_{n-b \text{ many } 2\text{'s}}.$$

The general case

Theorem 2.

$$h_r(x) = \frac{1}{x} \sum_{i=1}^r (g_r^{(0, i \bmod r)} - xg_r^{(0, i-1)} - \delta_{(i \bmod r, 0)}) (g_r^{(0, (r+1-i) \bmod r)} - xg_r^{(0, r-i)} - \delta_{((r+1-i) \bmod r, 0)}).$$

The general case is derived using the exact same logic as for the warm-up cases. Instead of having two cases as in the second warm-up, here we have r cases. Interested readers are welcome to verify the formula for $r = 3$ by themselves, and the general case should be apparent after this verification. As before, to get the generating function for V_r we simply let $f_r(x) = h_r(x^{1/r})$.

4. Using Maple packages

As noted in Shar and Zeilberger's paper ([SZ], page 7), now that we know $f_r(x)$ has the property of being algebraic, the sequence $a_r(n)$ satisfies some homogeneous linear recurrence equation with polynomial coefficients.

Using the *algtoec* procedure in the **SCHUTZENBERGER** package written by Doron Zeilberger, available from:

<http://www.math.rutgers.edu/~zeilberg/tokhniot/SCHUTZENBERGER.txt>

we are able to find (rigorously) recurrences (in operator notation) for our sequences when $r = 1$ and $r = 2$ ($r = 3$ took too long to compute):

For $r = 1$ we get:

$$(2n(2n+1) - (n+4)(n-2)N)a_1(n) = 0$$

(which agrees with the already known formula $a_1(n) = \frac{3}{n} \binom{2n}{n+3}$).

In the case when $r = 2$, *algtoec* returned a operator of degree 8, but it can be reduced to a minimal operator of degree 4 (thanks to Manuel Kauers for pointing it out), that is:

$$\begin{aligned} & (36(1+n)(2+n)(1+2n)(3+2n)(18154800 + 23101940n + 10635771n^2 + 2093616n^3 + 147833n^4) \\ & + 12(2+n)(3+2n)(1283329440 + 3700267618n + 4200957553n^2 + 2408049238n^3 + 735936616n^4 \\ & + 113774584n^5 + 6948151n^6)N + (282564806400 + 1066356868608n + 1704365727480n^2 \\ & + 1511140337906n^3 + 814587362081n^4 + 273775889012n^5 + 56080140110n^6 + 6405068474n^7 \\ & + 312371129n^8)N^2 - 2(4+n)(11939685120 + 40890299130n + 56943840213n^2 + 41794221496n^3 \\ & + 17488032270n^4 + 4183030930n^5 + 531527997n^6 + 27792604n^7)N^3 + 8(1+n)(4+n)(5+n)(11 \\ & + 2n)(3742848 + 7519914n + 5241921n^2 + 1502284n^3 + 147833n^4)N^4)a_2(n) = 0 \end{aligned}$$

Everything in this paper is implemented (with explanation) in the Maple packages **Words123New** and **PW123** and available from: <http://sites.math.rutgers.edu/~my237/One123>, which also includes some sample input and output files.

Acknowledgements

The author thanks Doron Zeilberger for bringing this project into her attention and continued conversation. The author also would like to thank Matthew Russell for offering suggestions that improved the exposition of this paper, Andrew Lohr and Alejandro Ginory for proofreading the draft, and Manuel Kauers for his help in finding the asymptotics.

References

[Bó] M. Bóna, *Permutations with one or two 132-subsequences*, Discrete Math. 181 (1998), 267-274.

<http://www.sciencedirect.com/science/article/pii/S0012365X97000629?via%3Dihub>

[Bu] A. Burstein, *A short proof for the number of permutations containing pattern 321 exactly once*, The Electronic J. of Combinatorics 18(2) (2011), P21.

<http://www.combinatorics.org/ojs/index.php/eljc/article/view/v18i2p21/pdf>

[K] M. Kauers, A Mathematica Package for Computing Asymptotic Expansions of Solutions of P-Finite Recurrence Equations, Technical Report RISC 11-04, Johannes Kepler University, April 2011.

[KP] M. Kauers and P. Paule, *The Concrete Tetrahedron: Symbolic Sums, Recurrence Equations, Generating Functions, Asymptotic Estimates*, Springer (2011), Section 6.5.

[N] J. Noonan, *The number of permutations containing exactly one increasing subsequence of length three*, Discrete Math. 152 (1996), 307-313.

<http://www.sciencedirect.com/science/article/pii/0012365X9500247T>

[SZ] N. Shar and D. Zeilberger, *The (Ordinary) Generating Functions Enumerating 123-Avoiding Words with r occurrences of each of $1, 2, \dots, n$ are Always Algebraic*, Annals of Combinatorics 20 (2016), 387-396.

<http://sites.math.rutgers.edu/~zeilberg/mamarim/mamarimhtml/words123.html>

[Z1] D. Zeilberger, *Alexander Burstein's Lovely Combinatorial Proof of John Noonan's Beautiful Theorem that the number of n -permutations that contain the Pattern 321 Exactly Once Equals $(3/n)(2n!)/((n-3)!(n+3)!)$* , Personal Journal of Shalosh B. Ekhad and Doron Zeilberger, Oct.18,

2011.

<http://sites.math.rutgers.edu/~zeilberg/mamarim/mamarimhtml/burstein.html>

[Z2] D. Zeilberger, *A Snappy Proof That 123-Avoiding Words are Equinumerous With 132-Avoiding Words*, Personal Journal of Shalosh B. Ekhad and Doron Zeilberger, April 11, 2005.

<http://sites.math.rutgers.edu/~zeilberg/mamarim/mamarimhtml/a123.html>

Department of Mathematics, Rutgers University, Piscataway, NJ 08854

E-mail address: **my237@math.rutgers.edu**