

On the cone of f -vectors of cubical polytopes

Ron M. Adin

Department of Mathematics, Bar-Ilan University

radin@math.biu.ac.il

Daniel Kalmanovich

Einstein Institute of Mathematics, The Hebrew University of Jerusalem

daniel.kalmanovich@gmail.com

Eran Nevo

Einstein Institute of Mathematics, The Hebrew University of Jerusalem

nevo.eran@gmail.com

April 23, 2018

Abstract

What is the minimal closed cone containing all f -vectors of cubical d -polytopes? We construct cubical polytopes showing that this cone, expressed in the cubical g -vector coordinates, contains the nonnegative g -orthant, thus verifying one direction of the Cubical Generalized Lower Bound Conjecture of Babson, Billera and Chan. Our polytopes also show that a natural cubical analogue of the simplicial Generalized Lower Bound Theorem does not hold.

1 Introduction

Understanding the possible face numbers of polytopes, and of subfamilies of interest, is a fundamental question, dealt with since antiquity. The celebrated g -theorem, conjectured by McMullen [McM71] and proved by Stanley [Sta80] (necessity) and by Billera and Lee [BL81] (sufficiency), characterizes the f -vectors of simplicial polytopes. Here we consider f -vectors of cubical polytopes; a d -polytope Q is *cubical* if all its facets are combinatorially isomorphic to the $(d - 1)$ -cube. Adin [Adi96] proved analogues of the Dehn–Sommerville relations for cubical polytopes, thus encoding the f -vector of Q by its (long) cubical g -vector

$$g^c(Q) = \left(g_1^c(Q), g_2^c(Q), \dots, g_{\lfloor d/2 \rfloor}^c(Q) \right)$$

(with the constant $g_0^c(Q) = 2^{d-1}$ omitted). The construction of neighborly cubical d -polytopes by Joswig and Ziegler [JZ00], where the number of vertices varies, shows that the linear span of the vectors $g^c(Q)$, over all cubical d -polytopes, is the entire vector space $\mathbb{R}^{\lfloor d/2 \rfloor}$. Adin [Adi96, Question 2] asked whether $g^c(Q)$ is always in the nonnegative orthant, and Babson, Billera and Chan [BBC97, Conjecture 5.2] conjectured further that the minimal closed cone \mathcal{C}_d containing all the vectors $g^c(Q)$ corresponding to cubical d -polytopes is exactly this nonnegative orthant \mathcal{A}_d .

Denote by e_i the i -th unit vector in $\mathbb{R}^{\lfloor d/2 \rfloor}$. Stacked cubical polytopes show that the ray spanned by e_1 is in \mathcal{C}_d , and neighborly cubical polytopes show that the ray spanned by $e_{\lfloor d/2 \rfloor}$ is in \mathcal{C}_d . Our main result is that all the rays spanned by the vectors e_i are in \mathcal{C}_d . Thus

Theorem 1. $\mathcal{A}_d \subseteq \mathcal{C}_d$.

The conjecture of Adin and Babson–Billera–Chan is

Conjecture 2. $\mathcal{A}_d = \mathcal{C}_d$.

An analogue of Theorem 1 was previously known for the much wider class of PL cubical spheres [BBC97, Theorem 5.7]. Also, Conjecture 2 holds for $d \leq 5$, by combining the constructions above with Steve Klee’s result [Kle11, Prop.3.7] asserting that $g_k^c(Q) \geq 0$ for any cubical polytope Q of dimension $2k + 1$.

Sanyal and Ziegler [SZ10] showed how to construct, from any simplicial $(d - 2)$ -polytope P on $n - 1$ vertices and a total order $v_1 < v_2 < \dots < v_{n-1}$ on its vertices, a cubical d -polytope $Q = Q(P, <)$ on 2^n vertices; it is the projection of a deformed n -cube in \mathbb{R}^n onto the last d coordinates. Further, they showed that if P is k -neighborly then the k -skeleton of Q is isomorphic to the k -skeleton of the n -cube. We apply their construction to the case where P_n is the k -neighborly k -stacked $(d - 2)$ -polytope on $n - 1$ vertices constructed by McMullen and Walkup [MW71], with $1 \leq k \leq \lfloor \frac{d-2}{2} \rfloor$, and with a suitable total order $<$. Analyzing the cubical g -vectors of the resulting polytopes $Q(k, d, n) = Q(P_n, <)$, as n tends to infinity, gives Theorem 1. See Theorem 13 and Corollary 14 for the exact values and asymptotic behavior of the cubical g -vectors.

The generalized lower bound theorem for simplicial polytopes (GLBT), conjectured by McMullen and Walkup [MW71] and proved by Murai and Nevo [MN13], asserts that for $1 \leq k < \lfloor \frac{d}{2} \rfloor$, a simplicial d -polytope P is k -stacked if and only if $g_{k+1}(P) = 0$. The polytopes $Q(k, d, n)$ demonstrate that the natural cubical analogue of the GLBT does not hold:

Theorem 3. *For any $k \geq 1$ and $n \geq d \geq 2k + 4$, we have $g_{k+2}^c(Q(k, d, n)) = 0$, and $Q(k, d, n)$ is not cubical $(k + 1)$ -stacked.*

The paper is organized as follows. Basic definitions and notation are given in section 2. In section 3 we define our variant of the McMullen–Walkup polytopes. A sketch of the Sanyal–Ziegler construction is given in section 4. In section 5 we construct the polytopes $Q(k, d, n)$ mentioned above and analyze their cubical g -vector. In section 6 we prove Theorem 3, showing that $Q(k, d, n)$ is not $(k + 1)$ -stacked. The final section 7 concludes with remarks and open questions.

Acknowledgments

The authors thank Michael Joswig and Isabella Novik for helpful comments. The research of Adin was supported by an MIT–Israel MISTI grant. He also thanks the Israel Institute for Advanced Studies, Jerusalem, for its hospitality during part of the work on this paper. The research of Kalmanovich and Nevo was partially supported by Israel Science Foundation grant ISF-1695/15 and by grant 2528/16 of the ISF–NRF Singapore joint research program. This work was also partially supported by the National Science Foundation under Grant No. DMS-1440140, while Nevo was in residence at the Mathematical Sciences Research Institute in Berkeley, California, during the Fall 2017 semester.

An extended abstract version of this paper will appear in the FPSAC 2018 proceedings. We thank the FPSAC 2018 referees for their comments on that extended abstract.

2 Preliminaries

The purpose of this section is mainly to set the notation that we will use throughout the paper. For undefined terminology we refer the reader to [Zie95]. A d -**dimensional polytope** P is the convex hull of a finite set of points in \mathbb{R}^d which affinely span \mathbb{R}^d . A (proper) **face** σ of P is the intersection of P with one of its supporting hyperplanes, the **dimension** $\dim \sigma$ of σ is then the dimension of the affine span of that intersection. The faces of dimensions 0, 1, and $d - 1$ are called **vertices**, **edges**, and **facets**, respectively. The empty set \emptyset and the polytope P itself are called **trivial faces** and have dimensions -1 and d , respectively. We will abbreviate and write d -polytope and i -face to denote dimension. We denote by $\text{Vert}(P)$ the set of vertices of P , and for a vertex $v \in \text{Vert}(P)$, we denote by P/v the **vertex figure** of P at v , that is, P/v is a $(d - 1)$ -polytope obtained as the intersection of P with a hyperplane which strictly separates v from $\text{Vert}(P) \setminus \{v\}$; the face lattice of P/v does not depend on the separating hyperplane chosen.

A **polytopal complex** K is a finite collection of polytopes in \mathbb{R}^d such that

- (i) the empty polytope is in K ,
- (ii) if $P \in K$ then all faces of P are also in K ,
- (iii) if $P, Q \in K$ then $P \cap Q$ is a face of both P and Q .

The **dimension** $\dim K$ of K is the maximum of $\dim P$ over all $P \in K$; we say that K is a $\dim K$ -complex. The elements in K are called **faces**; the faces of dimension $\dim K$ are called **facets**. For $F \in K$ we define the (open) **star** of F and the **antistar** of F , respectively, by

$$\begin{aligned} \text{st}_F(K) &= \{G \in K \mid F \subseteq G\}, \\ \text{ast}_F(K) &= \{G \in K \mid F \not\subseteq G\}. \end{aligned}$$

The number of i -faces in K is denoted by $f_i(K)$, and the f -**vector** of K is $f(K) = (f_0(K), f_1(K), \dots, f_{\dim K}(K))$. The f -**polynomial** of K is defined by

$$f(K, t) = \sum_{i=0}^{\dim K+1} f_{i-1}(K) t^i,$$

where $f_{-1}(K) = 1$ for any nonempty K .

For a polytope P we denote by $\langle P \rangle$ the complex of all faces of P . The **boundary complex** ∂P is the complex formed by all the proper faces of P , that is $\partial P = \langle P \rangle \setminus \{P\}$. We also define the f -vector and f -polynomial of P by $f(P) = f(\partial P)$ and $f(P, t) = f(\partial P, t)$. We use $\text{lk}_v(P)$ to denote the boundary complex $\partial(P/v)$ of the vertex figure of P at v .

2.1 Simplicial complexes and polytopes

A **simplicial complex** is a polytopal complex in which all polytopes are simplices. Let K be a simplicial $(D - 1)$ -complex; the h -**polynomial** of K is defined by

$$\begin{aligned} h(K, t) &= h_0(K) + h_1(K)t + \dots + h_D(K)t^D \\ &:= (1 - t)^D \cdot f\left(K, \frac{t}{1 - t}\right), \end{aligned}$$

and the h -vector of K is $(h_0(K), \dots, h_D(K))$. If $K = \partial P$ for a simplicial D -polytope P then the **Dehn–Sommerville relations** assert that $h_i(K) = h_{D-i}(K)$ for any $0 \leq i \leq D$. The corresponding g -vector $(g_0(K), \dots, g_{\lfloor D/2 \rfloor}(K))$ of K is then defined by

$$\begin{aligned} g_0(K) &= h_0(K) = 1, \\ g_i(K) &= h_i(K) - h_{i-1}(K), \quad \text{for all } 1 \leq i \leq \lfloor D/2 \rfloor. \end{aligned}$$

For two simplicial complexes K and L we define the **join** $K * L$ to be the simplicial complex whose simplices are the disjoint unions of simplices of K and simplices of L .

A polytope is **simplicial** if each of its proper faces is a simplex. For a simplicial polytope P we write $h(P, t)$ to mean $h(\partial P, t)$, and similarly $h_i(P) := h_i(\partial P)$ and $g_i(P) := g_i(\partial P)$.

A simplicial d -polytope P is called **k -stacked** if P has a triangulation in which there are no interior faces of dimension less than $d - k$. A simplicial polytope P is called **k -neighborly** if each subset of at most k vertices forms the vertex set of a face of P . We denote by $C(d, n)$ the **cyclic d -polytope with n vertices**:

$$C(d, n) := \text{conv} \{x(t_1), x(t_2), \dots, x(t_n)\},$$

where $t_1 < t_2 < \dots < t_n$ and $x(t) := (t, t^2, \dots, t^d)$ is the moment curve in \mathbb{R}^d .

2.2 Cubical complexes and polytopes

A **cubical complex** is a polytopal complex in which all polytopes are combinatorially isomorphic to cubes. Let Q be a cubical $(d - 1)$ -complex, the **short cubical h -polynomial** is defined by

$$h^{sc}(Q, t) = \sum_{i=0}^{d-1} h_i^{sc}(Q) t^i = \sum_{j=0}^{d-1} f_j(Q) (2t)^j (1-t)^{d-1-j}.$$

When Q is clear from the context, we may sometimes drop it from the notation, as we do in the following few definitions. The **(long) cubical h -vector** $(h_0^c, h_1^c, \dots, h_d^c)$ is defined by

$$\begin{aligned} h_0^c &= 2^{d-1}, \\ h_i^{sc} &= h_i^c + h_{i+1}^c, \quad \text{for } 0 \leq i \leq d-1, \end{aligned}$$

and the corresponding **(short and long) cubical g -vectors** are defined, as in the simplicial case, by

$$\begin{aligned} g_0^{sc} &= h_0^{sc} = f_0, & g_i^{sc} &= h_i^{sc} - h_{i-1}^{sc} \quad \text{for } 1 \leq i \leq \lfloor (d-1)/2 \rfloor; \\ g_0^c &= h_0^c = 2^{d-1}, & g_i^c &= h_i^c - h_{i-1}^c \quad \text{for } 1 \leq i \leq \lfloor d/2 \rfloor. \end{aligned}$$

A polytope is **cubical** if each of its proper faces is combinatorially a cube. Adin [Adi96] showed that any cubical d -polytope Q satisfies an analogue of the Dehn–Sommerville relations: $h_i^c(Q) = h_{d-i}^c(Q)$ for all $0 \leq i \leq d$.

In analogy with the simplicial case, [BBC97] defined cubical neighborliness and cubical stackedness: a cubical d -polytope is **k -neighborly** if it has the k -skeleton of a cube (of some dimension); it is **k -stacked** if it has a cubical subdivision with no interior faces of dimension less than $d - k$.

Each vertex figure in a cubical d -polytope P is a simplicial $(d - 1)$ -polytope; we finish this section with the relation known as **Hetyei's observation**:

$$(1) \quad h^{sc}(P, t) := h^{sc}(\partial P, t) = \sum_{v \in \text{Vert}(P)} h(\text{lk}_v(P), t).$$

It shows that the cubical Dehn–Sommerville relations follow from the simplicial ones.

3 The McMullen–Walkup polytopes

In section 3 of [MW71], McMullen and Walkup describe the construction of k -neighborly k -stacked simplicial D -polytopes with N vertices, for any set of parameters satisfying $2 \leq 2k \leq D < N$. Their construction takes a k -neighborly $2k$ -polytope C with $N - D + 2k$ vertices (e.g., the cyclic $2k$ -polytope with $N - D + 2k$ vertices) and a $(D - 2k)$ -simplex T , both lying in \mathbb{R}^D in such a way that the relative interior of T intersects the affine hull $\text{Aff}(C)$ in a vertex x of C . Then the convex hull $\text{conv}(C \cup T)$ is the desired polytope. We define a slightly more general construction.

Definition 4. Let $2 \leq K \leq D < N$. Let $C = C(K, N - D + K)$ be the cyclic K -polytope with $N - D + K$ vertices, and let T be a $(D - K)$ -simplex, both lying in \mathbb{R}^D in such a way that the relative interior of T intersects $\text{Aff}(C)$ in a vertex x of C . The polytope $\text{conv}(C \cup T)$ is a D -dimensional simplicial polytope with N vertices, denoted $\text{MW}(K, D, N; x)$.

The faces of $\text{MW}(K, D, N; x)$ are of two types:

- (a) the convex hull of T and a face of C that contains x ,
- (b) the convex hull of a proper face of T and a face of C that does not contain x .

The boundary complex of $\text{MW}(K, D, N; x)$ is thus described by

$$(2) \quad \partial \text{MW}(K, D, N; x) = \langle T \rangle * \text{lk}_x(C) \cup \bigcup_{\partial T * \text{lk}_x(C)} \partial T * \text{ast}_x(C).$$

McMullen and Walkup have shown that $\text{MW}(2k, D, N; x)$ is k -neighborly and k -stacked, thus satisfying

$$(3) \quad g_i(\text{MW}(2k, D, N; x)) = \begin{cases} 0, & i > k; \\ \binom{N-D-1}{i} = \binom{N-D+i-2}{i}, & i \leq k. \end{cases}$$

The proof [MW71, p. 269] that $\text{MW}(2k, D, N; x)$ is k -neighborly and k -stacked actually shows:

Lemma 5. *The polytope $\text{MW}(K, D, N; x)$ is $\lfloor K/2 \rfloor$ -neighborly and $\lfloor K/2 \rfloor$ -stacked. In particular, for $K = 2k - 1$,*

$$(4) \quad g_i(\text{MW}(2k - 1, D, N; x)) = \begin{cases} 0, & i > k - 1; \\ \binom{N-D-1}{i} = \binom{N-D+i-2}{i}, & i \leq k - 1. \end{cases}$$

The vertices of C come with a natural total order $v_1 < v_2 < \dots < v_{N-D+K}$, inherited from the order $t_1 < t_2 < \dots < t_{N-D+K}$ of the parameters in the definition of C . We take x to be the last vertex in that ordering, denoting the resulting polytope simply by $\text{MW}(K, D, N)$. Removing $x = v_{N-D+K}$, we extend the order $v_1 < \dots < v_{N-D+K-1}$ of the remaining vertices of C to an order $v_1 < \dots < v_{N-D+K-1} < v_{N-D+K} < \dots < v_N$ of the vertices of $\text{MW}(K, D, N)$, where v_{N-D+K}, \dots, v_N are the vertices of the $(D - K)$ -simplex T . We will use the following result:

Lemma 6. *The vertex figure $\text{MW}(2k, D, N) / v_1$ is combinatorially isomorphic to $\text{MW}(2k - 1, D - 1, N - 1)$.*

Proof. For $C = C(2k, N - D + 2k)$ denote $C' = C / v_1$, and note that $C' \cong C(2k - 1, N - D + 2k - 1)$. Applying the construction in Definition 4 to C' and a $(D - 2k)$ -simplex T produces an $\text{MW}(2k - 1, D - 1, N - 1)$ with boundary complex

$$(5) \quad \langle T \rangle * \text{lk}_x(C') \cup \bigcup_{\partial T * \text{lk}_x(C')} \partial T * \text{ast}_x(C').$$

We now show that the complex above is equal to $\text{lk}_{v_1}(\text{MW}(2k, D, N))$. Let F be a face of $\text{MW}(2k, D, N)$ containing v_1 . If F contains T then $F \setminus \{v_1\}$ is in $\langle T \rangle * \text{lk}_x(C')$, and if F does not contain T then $F \setminus \{v_1\}$ is in $\partial T * \text{ast}_x(C')$. Thus $\text{lk}_{v_1}(\text{MW}(2k, D, N))$ is contained in the complex (5). Similarly, for the other direction, take a face F' in the complex (5) and observe that $F' \cup \{v_1\}$ is in the boundary complex of $\text{MW}(2k, D, N)$, as described in (2). ■

4 The Sanyal–Ziegler construction

We give a very brief sketch of the construction, focusing on the combinatorial description of vertex figures. The reader is prompted to confer with the paper [SZ10], or with Sanyal's diploma thesis [San05].

Let $(P, <)$ be a simplicial $(d-2)$ -polytope with $n-1$ vertices, totally ordered by $<$. Label the vertices $v_1, \dots, v_{n-1} \in \mathbb{R}^{d-2}$ according to the given order $v_1 < v_2 < \dots < v_{n-1}$, and assume that the vertices are in general position, i.e., no $d-1$ vertices of P lie on a hyperplane. We start by defining the lexicographic diamonds of P .

4.1 Lexicographic diamonds

Let $w_1, \dots, w_{n-1} \in \mathbb{R}$ be a set of **heights**, and denote by $V^w = \{(w_i, v_i) \mid 1 \leq i \leq n-1\} \subset \mathbb{R}^{d-1}$ the set of **lifted vertices**. Let $p = (w_0, v'_0) \in \mathbb{R}^{d-1}$ be an arbitrary point with $w_0 \gg |w_i|$ for every $1 \leq i \leq n-1$, and consider the $(d-1)$ -polytope $D(P, w) = \text{conv}(V^w, p)$. We call $D(P, w)$ the **diamond** over P with heights w , noting that, for w_0 big enough, the combinatorial type of $D(P, w)$ is independent of the choice of the point p . Projecting the lower envelope of $D(P, w)$ onto P shows that $\text{ast}_p(D(P, w))$ is a polytopal subdivision of P .

Of special interest are the subdivisions of P induced by the heights $(w_1, w_2, \dots, w_{n-1}) = (\pm h, 0, \dots, 0)$ with $h > 0$. The subdivision of P induced by $(-h, 0, \dots, 0)$ is obtained by **pulling** v_1 ; it is a triangulation of P , and its facets are the pyramids with apex v_1 over facets in $P \cap P_1$ where $P_1 = \text{conv}(v_2, \dots, v_{n-1})$. The subdivision of P induced by $(h, 0, \dots, 0)$ is obtained by **pushing** v_1 , and its facets are the pyramids with apex v_1 over facets in $P_1 \setminus P$, plus one (possibly non-simplex) facet P_1 . The **a -th lexicographic subdivision** $\text{Lex}_a(P)$ of P is the polytopal subdivision of P obtained by successively pushing the vertices v_1, \dots, v_{a-1} , and then pulling v_a . That is, pushing v_1 creates a subdivision with one non-simplex cell P_1 , which we replace by its subdivision obtained by pushing v_2 , which, in turn, has only one non-simplex cell $P_2 = \text{conv}\{v_3, \dots, v_{n-1}\}$, and so on, until we finally replace $P_{a-1} = \text{conv}\{v_a, \dots, v_{n-1}\}$ by its triangulation obtained by pulling v_a .

The above iterative procedure amounts to choosing a set of heights w_1, \dots, w_{n-1} with

$$w_1 > \dots > w_{a-1} > 0 > w_a, \quad \text{and} \quad w_{a+1} = \dots = w_{n-1} = 0.$$

The resulting diamond (with $w_0 \gg w_1, -w_a$), denoted $D_a = D(P, w)$, is called the **a -th lexicographic diamond**. Its vertices are labeled v_0, v_1, \dots, v_{n-1} , with v_0 corresponding to the apex p ; thus $\text{ast}_{v_0}(D_a) = \text{Lex}_a(P)$.

Remark 7. Note that pushing or pulling a vertex in a simplex has no effect, thus the (possibly) different diamonds are D_a with $1 \leq a \leq n-d+1$.

4.2 The vertex figures of Q

Take a Gale transform $G \in \mathbb{R}^{(n-1) \times (n-d)}$ of P that has the form $G = \begin{bmatrix} I_{n-d} \\ \bar{G} \end{bmatrix}$, where $\bar{G} \in \mathbb{R}^{(d-1) \times (n-d)}$.

Plugging \bar{G} into the **deformed cube template** (see [SZ10, Definition 3.1]) produces a combinatorial n -cube $\bar{C} = C_n(\bar{G})$. The projection $\pi_d(\bar{C})$ of \bar{C} onto the last d coordinates is the cubical polytope $Q = Q(P, <)$ mentioned in the introduction.

The following key result from [SZ10]¹ states that each vertex figure of Q is combinatorially equivalent to some diamond D_a and, moreover, specifies which diamond corresponds to a given vertex v of Q .

Lemma 8 ([SZ10, Theorem 3.7]). *Let v be a vertex of \bar{C} labeled by $\sigma \in \{+, -\}^n$. Then the vertex figure of $\pi_d(v)$ in Q is isomorphic to D_a with*

$$a = \min \left(\{i \in [n] \mid \sigma_i = +\} \cup \{n-d+1\} \right).$$

The isomorphism $D_a \cong Q/v$ is given by: $v_i \in D_a$ corresponds to the neighbor of v obtained by flipping the $(i+1)$ -th coordinate of v ($0 \leq i \leq n-1$).

5 The cubical polytopes $Q(k, d, n)$

Fix positive integers $k \geq 1$ and $n \geq d \geq 2k+2$. We apply the Sanyal–Ziegler construction to the McMullen–Walkup polytope $P = \text{MW}(2k, d-2, n-1)$, with a total order $<$ on its vertices as described after Lemma 5 above. The result is a d -dimensional cubical polytope $Q = Q(k, d, n) = Q(P, <)$ with 2^n vertices. We now compute its cubical g -vector $g^c(Q)$, in stages.

5.1 Computing $g^{sc}(Q(k, d, n))$

By Hetyei's observation (1) we have

$$h_i^{sc}(Q) = \sum_{v \in \text{Vert}(Q)} h_i(\text{lk}_v(Q)) \quad (0 \leq i \leq d-1).$$

Therefore, by Lemma 8, for each $1 \leq i \leq \lfloor \frac{d-1}{2} \rfloor$:

$$(6) \quad g_i^{sc}(Q) = \sum_{v \in \text{Vert}(Q)} g_i(\text{lk}_v(Q)) = \sum_{a=1}^{n-d} 2^{n-a} g_i(D_a) + 2^d g_i(D_{n-d+1}).$$

We now compute the g -vectors of the diamonds D_a at hand, namely for our choice of $(P, <)$. First we compute $g(D_1)$, a computation that is then used to compute $g(D_a)$ for general a .

Proposition 9. *For $a = 1$ and $0 \leq i \leq \lfloor \frac{d-1}{2} \rfloor$:*

$$g_i(D_1) = \begin{cases} 0, & \text{if } i > k+1; \\ \binom{n-d}{k}, & \text{if } i = k+1; \\ \binom{n-d}{i}, & \text{if } i \leq k. \end{cases}$$

¹Theorem 3.7 in [SZ10] actually contains a typo, having $n-d-1$ instead of the correct value $n-d+1$. Their proof, however, does give the correct value. Further, [SZ10, Theorem 3.7] is stated for $Q(P, <)$ where P is neighborly; however, their proof holds verbatim for any simplicial polytope P .

Proof. Let $B_1 = \text{Lex}_1(P)$, namely the simplicial $(d-2)$ -ball triangulating P by starring from v_1 . Then

$$f(D_1, t) = f(B_1, t) + t \cdot f(P, t).$$

Note that, strictly speaking, the simplicial complexes actually considered here are ∂D_1 , B_1 and ∂P , of dimensions $d-2$, $d-2$ and $d-3$, respectively. The corresponding h -polynomial is

$$\begin{aligned} h(D_1, t) &= (1-t)^{d-1} \cdot f\left(D_1, \frac{t}{1-t}\right) \\ &= (1-t)^{d-1} \cdot \left[f\left(B_1, \frac{t}{1-t}\right) + \frac{t}{1-t} \cdot f\left(P, \frac{t}{1-t}\right) \right] \\ &= h(B_1, t) + t \cdot h(P, t). \end{aligned}$$

To compute $h(B_1, t)$, observe that

$$f(B_1, t) = (1+t) \cdot f(\text{ast}_{v_1}(P)) = (1+t) \cdot \left(f(P, t) - t \cdot f(\text{lk}_{v_1}(P), t) \right).$$

The h -polynomial is therefore

$$\begin{aligned} h(B_1, t) &= (1-t)^{d-1} \cdot f\left(B_1, \frac{t}{1-t}\right) \\ &= (1-t)^{d-1} \cdot \left(1 + \frac{t}{1-t} \right) \cdot \left[f\left(P, \frac{t}{1-t}\right) - \frac{t}{1-t} \cdot f\left(\text{lk}_{v_1}(P), \frac{t}{1-t}\right) \right] \\ &= (1-t)^{d-2} \cdot f\left(P, \frac{t}{1-t}\right) - t \cdot (1-t)^{d-3} \cdot f\left(\text{lk}_{v_1}(P), \frac{t}{1-t}\right) \\ &= h(P, t) - t \cdot h(\text{lk}_{v_1}(P), t). \end{aligned}$$

Summing up, we have

$$h(D_1, t) = (1+t) \cdot h(P, t) - t \cdot h(\text{lk}_{v_1}(P), t)$$

which yields, for $1 \leq i \leq \lfloor \frac{d-1}{2} \rfloor$,

$$(7) \quad g_i(D_1) = g_i(P) + g_{i-1}(P) - g_{i-1}(\text{lk}_{v_1}(P)).$$

Using (3), Lemma 6 and Lemma 5, we compute the right-hand side of (7) to obtain the result. \blacksquare

Proposition 10. For each $2 \leq a \leq n-d+1$ and $0 \leq i \leq \lfloor \frac{d-1}{2} \rfloor$:

$$g_i(D_a) = \begin{cases} 0, & \text{if } i > k+1; \\ \binom{n-d-a+1}{k}, & \text{if } i = k+1; \\ \binom{n-d}{i}, & \text{if } i \leq k. \end{cases}$$

Proof. Let v_0 be the apex of the diamond D_a . The link condition $\text{lk}_{v_1 v_0}(D_a) = \text{lk}_{v_1}(D_a) \cap \text{lk}_{v_0}(D_a)$ holds: indeed, $\text{lk}_{v_0}(D_a) = \partial P$ and the faces of $\text{lk}_{v_1}(D_a)$ not containing v_0 form $\text{lk}_{v_1}(\text{Lex}_a(P))$, so

$$\text{lk}_{v_1}(D_a) \cap \text{lk}_{v_0}(D_a) = \text{lk}_{v_1}(\text{Lex}_a(P)) \cap \partial P,$$

and as $a \geq 2$, the right hand side equals $\text{lk}_{v_1}(\partial P)$, for which we have $\text{lk}_{v_1}(\partial P) = \text{lk}_{v_1}(\text{lk}_{v_0}(D_a)) = \text{lk}_{v_1 v_0}(D_a)$.

Denote by K the $(d-2)$ -complex obtained from ∂D_a by contracting the edge $v_1 v_0$. Then, the h -polynomials are related by

$$h(D_a, t) = h(K, t) + t \cdot h(P/v_1, t),$$

see e.g. [Nev07, Eq. (10)].

Note that K is the boundary complex of a polytope, namely of the $(a-1)$ -lexicographic diamond over $P_1 = \text{MW}(2k, d-2, n-2)$; denote this diamond by $Y_{a-1}(n-1)$. Then, by Lemma 6

$$h(D_a, t) = h(Y_{a-1}(n-1), t) + t \cdot h(\text{MW}(2k-1, d-3, n-2), t).$$

We may again contract the edge between the first vertex of P_1 and the apex v_0 in $Y_{a-1}(n-1)$ to obtain $Y_{a-2}(n-2)$. We do these edge contractions $a-1$ times, and get

$$h(D_a, t) = h(Y_1(n-a+1), t) + t \cdot \sum_{j=1}^{a-1} h(\text{MW}(2k-1, d-3, n-1-j), t),$$

and therefore

$$(8) \quad g_i(D_a) = g_i(Y_1(n-a+1)) + \sum_{j=1}^{a-1} g_{i-1}(\text{MW}(2k-1, d-3, n-1-j)).$$

As $Y_1(n-a+1)$ is the 1-lexicographic diamond over $\text{MW}(2k, d-2, n-a)$, the claimed result follows from Proposition 9, Lemma 5, and the identity

$$\begin{aligned} & \binom{n-d-a+1}{i} + \sum_{j=1}^{a-1} \binom{n-d-j+1}{i-1} \\ &= \binom{n-d-a+1}{i} + \sum_{j=1}^{a-1} \left[\binom{n-d-j+1}{i} - \binom{n-d-j}{i} \right] = \binom{n-d}{i}. \quad \blacksquare \end{aligned}$$

Combining (6) with Propositions 9 and 10, and noting that $\binom{0}{k} = 0$ for $k \geq 1$, we conclude

Corollary 11. For each $0 \leq i \leq \lfloor \frac{d-1}{2} \rfloor$,

$$g_i^{sc}(Q) = \begin{cases} 0, & \text{if } i > k+1; \\ \sum_{a=1}^{n-d} 2^{n-a} \binom{n-d-a+1}{i}, & \text{if } i = k+1; \\ 2^n \binom{n-d}{i}, & \text{if } i \leq k. \end{cases}$$

5.2 Computing $g^c(Q(k, d, n))$

In order to compute the cubical g -vector of Q , and in particular $g_{k+2}^c(Q)$, we need the following binomial identity:

Lemma 12. For any integers $k \geq 1$ and $m \geq 0$,

$$\sum_{a=1}^m 2^{m-a} \binom{m-a+1}{k} = (-1)^{k+1} + 2^m \sum_{j=0}^k (-1)^j \binom{m}{k-j}.$$

Proof. Fix $k \geq 1$, and denote

$$\begin{aligned} L_m &:= \sum_{a=1}^m 2^{m-a} \binom{m-a+1}{k} = \sum_{b=0}^{m-1} 2^b \binom{b+1}{k}, \\ R_m &:= (-1)^{k+1} + 2^m \sum_{j=0}^k (-1)^j \binom{m}{k-j}. \end{aligned}$$

We will prove that $L_m = R_m$ for all $m \geq 0$, by induction on m .

For $m = 0$, indeed $L_0 = R_0 = 0$. For the induction step, note that $L_{m+1} = L_m + 2^m \binom{m+1}{k}$. Thus it suffices to prove that $R_{m+1} = R_m + 2^m \binom{m+1}{k}$ as well. Indeed,

$$\begin{aligned} R_{m+1} - R_m &= 2^m \sum_{j=0}^k (-1)^j \left[2 \binom{m+1}{k-j} - \binom{m}{k-j} \right] \\ &= 2^m \sum_{j=0}^k (-1)^j \left[\binom{m+1}{k-j} + \binom{m+1}{k-j-1} \right] \\ &= 2^m \binom{m+1}{k}, \end{aligned}$$

as claimed. ■

Theorem 13. For each $1 \leq i \leq \lfloor d/2 \rfloor$,

$$g_i^c(Q) = \begin{cases} 0, & \text{if } i > k+1; \\ 2^n \sum_{j=1}^i (-1)^{j-1} \binom{n-d}{i-j} + (-1)^i 2^d, & \text{if } i \leq k+1. \end{cases}$$

Proof. From

$$\begin{aligned} g_0^c(Q) &= 2^{d-1}, & g_0^{sc}(Q) &= 2g_0^c(Q) + g_1^c(Q), \\ g_i^{sc}(Q) &= g_i^c(Q) + g_{i+1}^c(Q) & (1 \leq i \leq \lfloor (d-1)/2 \rfloor) \end{aligned}$$

it follows that

$$(9) \quad g_i^c(Q) = \sum_{j=1}^i (-1)^{j-1} g_{i-j}^{sc}(Q) + (-1)^i 2^d \quad (1 \leq i \leq \lfloor d/2 \rfloor).$$

The values of $g_i^c(Q)$ for $i \leq k+1$ now follow easily from Corollary 11. It also follows that

$$g_i^c(Q) + g_{i+1}^c(Q) = 0 \quad (i \geq k+2),$$

and all that remains is to show that $g_{k+2}^c(Q) = 0$. Indeed, by (9) and Corollary 11,

$$\begin{aligned} g_{k+2}^c(Q) &= \sum_{j=1}^{k+2} (-1)^{j-1} g_{k+2-j}^{sc}(Q) + (-1)^{k+2} 2^d \\ &= \sum_{a=1}^{n-d} 2^{n-a} \binom{n-d-a+1}{k} + 2^n \sum_{j=2}^{k+2} (-1)^{j-1} \binom{n-d}{k+2-j} + (-1)^{k+2} 2^d \\ &= \sum_{a=1}^{n-d} 2^{n-a} \binom{n-d-a+1}{k} - \left[2^n \sum_{j=0}^k (-1)^j \binom{n-d}{k-j} + (-1)^{k+1} 2^d \right]. \end{aligned}$$

Using Lemma 12 with $m = n-d$ gives, indeed, $g_{k+2}^c(Q) = 0$ as claimed. ■

Corollary 14. Fix $k \geq 1$ and $d \geq 2k+2$, and let $\{Q_n\}_{n=d}^\infty = \{Q(k, d, n)\}_{n=d}^\infty$. Then

$$\lim_{n \rightarrow \infty} \frac{g_{k+1}^c(Q_n)}{2^n \binom{n-d}{k}} = 1, \quad \text{and} \quad \lim_{n \rightarrow \infty} \frac{g_i^c(Q_n)}{2^n \binom{n-d}{k}} = 0 \quad (\forall i \neq k).$$

Corollary 14 shows that the ray spanned by e_k ($2 \leq k \leq \lfloor d/2 \rfloor$) in $\mathbb{R}^{\lfloor d/2 \rfloor}$ belongs to the closed cone \mathcal{C}_d . Note that this was already known for the ray spanned by $e_{\lfloor d/2 \rfloor}$ because of the existence of neighborly cubical d -polytopes, such as $Q(k, 2k + 2, n)$. The ray spanned by e_1 belongs to this cone because of the existence of stacked cubical d -polytopes. Thus \mathcal{C}_d contains \mathcal{A}_d , and Theorem 1 is proved.

6 No obvious cubical GLBT

In [BBC97], after introducing the definitions of cubical stackedness and cubical neighborliness, the authors show that cubical 1-stacked d -polytopes with at least n vertices exist, for any $n \geq 2^d$ (see [BBC97, Corollary 5.6]). It is also shown (see [BBC97, proof of Proposition 5.5]) that if Q is a cubical k -stacked d -polytope, then $g_i^c(Q) = 0$ for $k < i \leq \lfloor d/2 \rfloor$. The converse claim, namely, that $g_{k+1}^c(Q) = 0$ implies that Q is cubical k -stacked, is false, as our analysis of $Q(k, d, n)$ below shows. This is in apparent contrast with the simplicial GLBT.

Theorem 15. *For any $k \geq 1$ and $n \geq d \geq 2k + 4$, the polytope $Q = Q(k, d, n)$ is cubical k -neighborly but not cubical $(k+1)$ -stacked, although $g_{k+2}^c(Q) = 0$. In fact, it is not j -stacked for any $k+1 \leq j \leq \lfloor d/2 \rfloor - 1$.*

Proof. The polytope Q is cubical k -neighborly by [JZ00, Theorem 16] or [SZ10, Theorem 3.2]; this also follows from our explicit computation of its (short) cubical g -vector. By Theorem 13, $g_{k+2}^c(Q) = 0$. Assume by contradiction that Q is cubical $(k+1)$ -stacked, so Q has some cubulation Q' , namely a subdivision into (combinatorial) cubes, without interior $(d - k - 2)$ -faces. Let C_n be the deformed n -cube that Q is a projection of.

Lemma 16. *All faces of Q' must be faces of C_n .*

Proof. Consider an m -cube F of Q' . Since Q' has no interior 1-faces, the 1-skeleton of F is in Q , and so also in C_n . Now, any 1-dimensional subcomplex of C_n which is isomorphic to the graph of an m -cube, is the 1-skeleton of an m -face of C_n . ■

Next, the cubulation Q' induces a set of *compatible* triangulations of the vertex figures. We extend the notation for the boundary complex of the vertex figure in a polytope, and denote by $\text{lk}_v(B)$ the simplicial complex whose face lattice is the ideal above the vertex v in the face lattice of B , with B a simplicial complex or a cubical complex.

Each $\text{lk}_v(Q')$ is a triangulation of Q/v with no interior $(d - k - 3)$ -faces. Thus the vertex figure of Q at v — isomorphic to some diamond D_a — is $(k+1)$ -stacked, and by the simplicial GLBT (an easy part of it, see [MN13, Thm.2.3(ii)] or [BD13, Thm.2.12]), $\text{lk}_v(Q')$ is the unique triangulation obtained from the diamond D_a by inserting all $(d-1)$ -simplices whose $(d-k-3)$ -skeleton is contained in the boundary of D_a . (We abuse notation and identify ∂D_a with $\text{lk}_v(Q)$ by the isomorphism given in Lemma 8.) This description allows us to determine, for each vertex v of Q , the list of d -cubes in Q' that contain v . The compatibility condition mentioned above is the requirement that if u is a vertex in a d -cube from the list of v , then the list of u must contain this d -cube too. We show, in steps, that the $(k+1)$ -stacked triangulations of the diamonds D_a are incompatible, thus such a cubulation Q' cannot exist.

In order to identify the facets in the unique $(k+1)$ -stacked triangulation $S(D_a)$ of D_a , we first identify the *missing faces* of D_a , namely its minimal non-faces. Kalai [Kal94] and Nagel [Nag08] showed that if $g_i(R) = 0$ for a simplicial j -polytope R , then R has no missing ℓ -faces for $i \leq \ell \leq j - i$. Combined with the simplicial GLBT, a subset A of d vertices of D_a is a facet in $S(D_a)$ iff all its subsets B of size

$\leq k + 2$ are faces of D_a . Since D_a is k -neighborly, it is enough to determine the missing faces of D_a of sizes $k + 1$ and $k + 2$. To analyze these missing faces of D_a we make an essential use of the following observation: let $P = MW(2k, d - 2, m)$, and recall Definition 4, where P is constructed from the cyclic polytope $C = C(2k, m - d + 2 + 2k)$ and the $(d - 2 - 2k)$ -simplex T .

Lemma 17. *The operations MW and Lex commute:*

$$(10) \quad \text{Lex}_a(P) = \langle T \rangle * \text{lk}_x(\text{Lex}_a(C)) \cup_{\partial T * \text{lk}_x(\text{Lex}_a(C))} \partial T * \text{ast}_x(\text{Lex}_a(C)).$$

Proof. By the construction of P , for a subset F of $\text{Vert}(P)$ (so that $x \notin F$), we have:

- F is a face of P , with $\text{Vert}(T) \subseteq \text{Vert}(F)$, iff $(\text{Vert}(F) \setminus \text{Vert}(T)) \cup \{x\}$ is the set of vertices of a face \bar{F} of C , and
- F is a face of P , with $\text{Vert}(T) \not\subseteq \text{Vert}(F)$, iff $\text{Vert}(F) \setminus \text{Vert}(T)$ is the set of vertices of a face \bar{F} of C .

Now, let F be a face added when pulling v_1 in P . Then $F = v_1 * F'$, with F' a face of P , and F' a face of $P_1 = \text{conv}(v_2, \dots, v_m)$. With the notation above, \bar{F}' is a face of both C and C_1 , and thus $\bar{F} = v_1 * \bar{F}'$ is added when pulling v_1 in C . Similarly, when pushing v_1 in P , we add faces F (and $v_1 * F$) of P_1 which are non-faces of P , and so \bar{F} is a face of C_1 , and a non-face of C , and thus is added when pushing v_1 in C . Reversing the argument shows that faces added when pushing/pulling v_1 in C give corresponding added faces in pushing/pulling in P . When constructing $\text{Lex}_a(P)$ and $\text{Lex}_a(C)$ sequentially, we always push/pull v_1 in a suitable MW-polytope and the corresponding suitable cyclic polytope C , hence (10) follows. ■

We need some terminology, to be used in the following two propositions: let G be a subset of the set $\{v_1, \dots, v_{n-d+2k+1}\}$ of vertices of the cyclic polytope C . A **block** $B \subseteq G$ is a maximal (with respect to inclusion) subset of G consisting of consecutive vertices, B is **inner** if $\min B > v_1$ and $\max B < v_{n-d+2k+1} = x$, and B is **even / odd** if its size is. We call a subset of vertices of C **isolated** if no two of them are consecutive. Recall that p denotes the apex of the diamond D_a of $P = MW(2k, d - 2, n - 1)$.

Proposition 18. *For any $1 \leq a \leq n - d + 1$, the missing faces in D_a of size $k + 1$ or $k + 2$ are exactly the sets of the following two types:*

- (1) $F = \{p\} \cup F'$, with F' of size $k + 1$, consisting of isolated vertices of C , $x \notin F'$, and $\min F' = a$; or
- (2) F is of size $k + 1$, consisting of isolated vertices of C , $x \notin F$, $\min F \neq a$.

Proof. Any missing face F of D_a must contain a missing face of P , and is either of the form

- (i) $\{p\} \cup F'$ where F' is an interior face of the ball $\text{Lex}_a(P)$, or
- (ii) a missing face of ∂P which is not a face of $\text{Lex}_a(P)$, or
- (iii) a missing face of $\text{Lex}_a(P)$ containing a missing face F' of ∂P as a strict subset.

Using Lemma 17 we can replace P by C in (i), (ii) and (iii), and F contains a missing face of C not containing x .

By the Gale evenness criterion, a subset of i vertices of C forms a face of C iff it contains at most $2k - i$ inner odd blocks. It follows that the missing faces of C are exactly the sets of $k + 1$ isolated vertices, containing at most one element of $\{v_1, x\}$.

Let F' be a missing face of C , $x \notin F'$, with $\min F' = v_i$, and denote $C_i = \text{conv}\{v_j \mid j \geq i\}$, a cyclic polytope contained in C .

If $i = a$ then F' is a face in $\text{Lex}_a(C_a)$, thus F' is also in $\text{Lex}_a(C)$, yielding a missing face $F = F' \cup \{v_a\}$ of D_a of type (i).

If $i > a$, then F' is a missing face in $\text{Lex}_a(C)$, hence $F = F'$ is a missing face of D_a of type (ii).

If $i < a$, C_{i+1} blocks v_i from seeing $F' \setminus v_i$, so F' is missing in $\text{Lex}_a(C)$, and so $F = F'$ is a missing face of D_a of type (ii).

Now, we will show that missing faces of type (iii) do not exist, and thus the characterization in the statement will follow. Indeed, assume F to be of type (iii), and denote $\{z\} = F \setminus F'$ (indeed this difference set must be a singleton as $|F'| = k + 1 < |F| \leq k + 2$ and $F' \subseteq F$). As $F' \in \text{Lex}_a(C)$, the discussion above shows $\min(F') = v_a$. If $z \neq \min F$, then $\min F = v_a$, and z is adjacent to some vertex of F' (else $F \setminus \{v_a\}$ would be missing in $\text{Lex}_a(C)$). But there is no missing face containing v_a in $\text{Lex}_a(C_a)$ so F is in $\text{Lex}_a(C_a)$, hence also in $\text{Lex}_a(C)$, a contradiction. If $z = \min F$, then $F \setminus \{v_a\}$ is a missing face of D_a of type (ii), a contradiction.

To conclude, note that the items (i) and (ii), with C replacing P , are restatements of (1) and (2), respectively. ■

Proposition 18 and the discussion preceding Lemma 17 imply the following description of the $(d - 1)$ -faces of the $(k + 1)$ -stacked triangulation $S(D_a)$.

Proposition 19. *The sets of d vertices that form the $(d - 1)$ -simplices of the $(k + 1)$ -stacked triangulation $S(D_a)$ of D_a are exactly the sets of the following two types:*

- (I) $\{p\} \cup \{2k\text{-set in } C \setminus \{x\} \text{ consisting of even blocks}\} \cup \text{Vert}(T)$, or
- (II) $\{a\} \cup \{2k\text{-set } F \text{ in } C \setminus \{x\} \text{ consisting of even blocks, } \min F > a\} \cup \text{Vert}(T)$.

Proof. Let Y be a d -set that forms a facet of $S(D_a)$. Then Y cannot contain more than $2k + 1$ vertices of $C \setminus \{x\}$, because then it would contain two disjoint missing faces of C , one of which must be of type (2). Clearly, Y has to contain at least $2k$ vertices of $C \setminus \{x\}$. So, by Proposition 18, Y is either of the form

- $2k$ vertices of $C \setminus \{x\}$, the apex p , and all of the vertices of T , or
- $2k + 1$ vertices of $C \setminus \{x\}$, and all of the vertices of T .

If Y has $2k + 1$ vertices of $C \setminus \{x\}$, then it contains $k + 1$ isolated vertices of $C \setminus \{x\}$, so $\min Y = a$, since Y contains no missing faces of type (2), and $p \notin Y$ since Y contains no missing faces of type (1). These are the d -sets of type (II) in the statement. Otherwise, Y has $2k$ vertices of $C \setminus \{x\}$, and hence must contain p and all vertices of T , and since Y does not contain missing faces of type (1) or (2), it must be of type (I). ■

We are now ready to complete the proof of Theorem 15. Consider a vertex v_σ of Q with $\sigma \in \{+, -\}^n$, and with $a = \min\{i \mid \sigma_i = +\} < n - d + 1$. Lemma 8 says that $Q/v_\sigma \cong D_a$, with the vertex $v_i \in D_a$ corresponding to the vertex in Q/v_σ obtained by flipping the $(i + 1)$ -th coordinate of σ . Consider $v_{\sigma'}$, where σ' is obtained by flipping the a -th coordinate of σ . Then $v_{\sigma'}$ is a neighbor of v_σ with $Q/v_{\sigma'} \cong D_b$, for some $b > a$.

By Proposition 19, there is a $(d - 1)$ -face F in $S(D_a)$ of type (II) (e.g., consisting of a , $\text{Vert}(T)$, and a single $2k$ -block in $C \setminus \{x\}$). It corresponds to a d -cube D in Q' . The same d -set of coordinates F

corresponds to the face of $S(D_b) \cong \text{lk}_{v_{\sigma'}}(Q')$ which corresponds to $D \in Q'$. But F is of neither type (I) nor type (II) in D_b , and so the triangulations are incompatible.

This completes the proof that Q is not $(k+1)$ -stacked. For $k+1 < j \leq \lfloor d/2 \rfloor - 1$, if Q were j -stacked, then the induced j -stacked triangulation of any diamond D_a would again be the same $S(D_a)$ given by the GLBT, and thus the analysis above yields a contradiction as before. ■

7 Concluding remarks

7.1 Cubical k -stacked d -polytopes

As mentioned in Theorem 15, $Q(k, d, n)$ is not cubical j -stacked for any $k+1 \leq j \leq \lfloor d/2 \rfloor - 1$. It is then natural to ask: do cubical k -stacked, and not $(k-1)$ -stacked, d -polytopes actually exist?

The k -elementary cubical d -polytopes C_k^d constructed for any $0 \leq k \leq d$ by Blind and Blind in [BB95] provide a positive answer to this question. For $1 \leq k \leq \lfloor d/2 \rfloor$, C_k^d is k -stacked by construction and its g^c -vector is

$$g_i^c(C_k^d) = \sum_{j=1}^k 2^{d-j} \binom{j-1}{i-1} \quad (1 \leq i \leq \lfloor d/2 \rfloor)$$

so that, in particular, $g_k^c(C_k^d) = 2^{d-k} > 0$ and hence C_k^d is not $(k-1)$ -stacked.

The following question, implicit in [BBC97], asks for a sequence of cubical k -stacked d -polytopes with g^c -vector approaching the ray spanned by e_k . It is still unanswered.

Question 20. *Let $2 \leq k \leq \lfloor d/2 \rfloor - 1$. Does there exist a sequence of cubical k -stacked d -polytopes with dominant k -th coordinate g_k^c of the cubical g -vector?*

7.2 The Cubical Lower Bound Conjecture

Jockusch studied the lower and upper bound problems for cubical polytopes in [Joc93], where he stated a *Cubical Lower Bound Conjecture*, rephrased as follows:

Conjecture 21 (CLBC, [Joc93]). *Let Q be a cubical d -polytope with n vertices. Then*

$$\begin{aligned} f_k(Q) &\geq \left[\binom{d}{k} + \binom{d-1}{k-1} \right] 2^{-k} n - \binom{d-1}{k-1} 2^{d-k} \quad (1 \leq k \leq d-2), \\ f_{d-1}(Q) &\geq (2d-2)2^{-(d-1)}n - (2d-4). \end{aligned}$$

Jockusch remarks that the CLBC for edges is as strong as the general statement. This reduction is a cubical analog of the *MPW-reduction* (see [Bar73, Theorem 1]), and it follows that Conjecture 21 is equivalent to the $k=1$ case, which in turn can be rewritten as

Conjecture 22 (CLBC). *If Q is a cubical d -polytope with n vertices then*

$$g_2^c(Q) \geq 0.$$

References

[Adi96] R. M. Adin. A new cubical h -vector. *Discrete Math.*, 157(1-3):3–14, 1996.

- [Bar73] D. Barnette. A proof of the lower bound conjecture for convex polytopes. *Pacific J. Math.*, 46:349–354, 1973.
- [BB95] G. Blind and R. Blind. The cubical d -polytopes with fewer than 2^{d+1} vertices. *Discrete Comput. Geom.*, 13(3-4):321–345, 1995.
- [BBC97] E. K. Babson, L. J. Billera, and C. S. Chan. Neighborly cubical spheres and a cubical lower bound conjecture. *Israel J. Math.*, 102:297–315, 1997.
- [BD13] B. Bagchi and B. Datta. On k -stellated and k -stacked spheres. *Discrete Math.*, 313(20):2318–2329, 2013.
- [BL81] L. J. Billera and C. W. Lee. A proof of the sufficiency of McMullen’s conditions for f -vectors of simplicial convex polytopes. *J. Combin. Theory Ser. A*, 31(3):237–255, 1981.
- [Joc93] W. Jockusch. The lower and upper bound problems for cubical polytopes. *Discrete Comput. Geom.*, 9(2):159–163, 1993.
- [JZ00] M. Joswig and G. M. Ziegler. Neighborly cubical polytopes. *Discrete Comput. Geom.*, 24(2-3):325–344, 2000. The Branko Grünbaum birthday issue.
- [Kal94] G. Kalai. Some aspects of the combinatorial theory of convex polytopes. In *Polytopes: abstract, convex and computational (Scarborough, ON, 1993)*, volume 440 of *NATO Adv. Sci. Inst. Ser. C Math. Phys. Sci.*, pages 205–229. Kluwer Acad. Publ., Dordrecht, 1994.
- [Kle11] S. Klee. Lower bounds for cubical pseudomanifolds. *Discrete Comput. Geom.*, 46(2):212–222, 2011.
- [McM71] P. McMullen. The numbers of faces of simplicial polytopes. *Israel J. Math.*, 9:559–570, 1971.
- [MN13] S. Murai and E. Nevo. On the generalized lower bound conjecture for polytopes and spheres. *Acta Math.*, 210(1):185–202, 2013.
- [MW71] P. McMullen and D. W. Walkup. A generalized lower-bound conjecture for simplicial polytopes. *Mathematika*, 18:264–273, 1971.
- [Nag08] U. Nagel. Empty simplices of polytopes and graded Betti numbers. *Discrete Comput. Geom.*, 39(1-3):389–410, 2008.
- [Nev07] E. Nevo. Higher minors and Van Kampen’s obstruction. *Math. Scand.*, 101(2):161–176, 2007.
- [San05] R. Sanyal. On the combinatorics of projected deformed products. Diploma Thesis, available online at <http://page.mi.fu-berlin.de/sanyal/DiplomaThesis.pdf>, 2005.
- [Sta80] R. P. Stanley. The number of faces of a simplicial convex polytope. *Adv. in Math.*, 35(3):236–238, 1980.
- [SZ10] R. Sanyal and G. M. Ziegler. Construction and analysis of projected deformed products. *Discrete Comput. Geom.*, 43(2):412–435, 2010.
- [Zie95] G. M. Ziegler. *Lectures on polytopes*, volume 152 of *Graduate Texts in Mathematics*. Springer-Verlag, New York, 1995.