

PARTIAL REGULARITY FOR SOLUTIONS TO SUBELLIPTIC EIKONAL EQUATIONS

PAOLO ALBANO, PIERMARCO CANNARSA, AND TERESA SCARINCI

ABSTRACT. On a bounded domain Ω in euclidean space \mathbb{R}^n , we study the homogeneous Dirichlet problem for the eikonal equation associated with a system of smooth vector fields, which satisfies Hörmander's bracket generating condition. We prove that the solution is smooth in the complement of a closed set of Lebesgue measure zero.

1. INTRODUCTION

Let $\Omega \subset \mathbb{R}^n$ be a bounded open set with boundary Γ , given by a smooth manifold of dimension $n - 1$. Let X_1, \dots, X_N be a system of smooth vector fields defined on some open neighbourhood of Ω , say Ω' . Hereafter, the term *smooth* stands for either C^∞ or C^ω , the latter meaning real analytic functions. We shall assume that *Hörmander's bracket-generating condition* is satisfied, i.e., $\text{Lie}\{X_1, \dots, X_N\}(x) = \mathbb{R}^n$, $\forall x \in \Omega'$, where $\text{Lie}\{X_1, \dots, X_N\}(x)$ denotes the space of all values, at x , of the vector fields of the Lie algebra generated by $\{X_1, \dots, X_N\}$. We point out that we need not suppose such vector fields to be linearly independent, nor that $N < n$.

Under the above assumptions—that will be in force throughout the paper—it is well known that the boundary value problem

$$(1.1) \quad \sum_{j=1}^N (X_j T)^2(x) = 1 \quad \text{in } \Omega, \quad T = 0 \quad \text{on } \Gamma,$$

admits a unique continuous viscosity solution. Moreover, T is Hölder continuous but fails to be more regular, in general.

In [3], we investigated the regularity of T . Building on such results, in this paper we analyse the *singular support* of T .

Date: March 15, 2022.

Key words and phrases. eikonal equation; degenerate equations; sub-Riemannian geometry; semiconcavity.

Definition 1.1. *The singular support of a function $f : \Omega \rightarrow \mathbb{R}$, $\text{Sing supp } f$ in short, is the complement in Ω of the set of all points $x \in \Omega$ that have an open neighbourhood on which f is smooth.*

In a similar way, one can define the $C^{1,1}$ singular support and the Lipschitz singular support of T , which are denoted by $\text{Sing supp}_{C^{1,1}} T$ and $\text{Sing supp}_{Lip} T$, respectively. (It is clear that $\text{Sing supp } T$ is closed in Ω .) We first prove the following result.

Theorem 1.1. $\text{Sing supp } T = \text{Sing supp}_{C^{1,1}} T$.

Moreover, we show that the singular support of T is a negligible set.

Theorem 1.2. $\text{Sing supp } T$ has Lebesgue measure zero.

We note that Theorem 1.2 is related to the so-called *minimizing Sard conjecture* in sub-riemannian geometry (see, e.g., [12, Conjecture 1, p. 158]). One of the formulations of such a conjecture, adapted to the case of a smooth target, claims that the set \mathcal{S}_{min} , which consists of all points lying on a singular minimizing trajectory, should have Lebesgue measure zero. Since, by [3, Theorem 3.2], \mathcal{S}_{min} coincides with the set on which the sub-riemannian distance fails to be Lipschitz, the above conjecture can be proved by appealing to [10], where the almost everywhere differentiability of the sub-riemannian distance to a closed set with the inner ball property is obtained.

A further part of the same conjecture could be rephrased saying that the set of all points, on a neighbourhood of which the sub-riemannian distance is smooth, should have full Lebesgue measure. Since such a set is nothing but the complement of the singular support of the sub-riemannian distance, Theorem 1.2 above shows the conjecture to be true for smooth targets of codimension 1.

2. PROOFS

The proof of theorems 1.1 and 1.2 relies on the fact that the solution T of (1.1) is the value function of a suitable time optimal control problem.

Let $x \in \bar{\Omega}$. For any measurable function $u = (u_1, \dots, u_N) : [0, +\infty[\rightarrow \mathbb{R}^N$ taking values in $\bar{B}_1(0)$, the unit closed ball of \mathbb{R}^N , we denote by $y^{x,u}$ the unique maximal solution of the Cauchy problem

$$(2.2) \quad \begin{cases} y'(t) = \sum_{j=1}^N u_j(t) X_j(y(t)) & (t \geq 0) \\ y(0) = x. \end{cases}$$

The time needed to steer x to Γ along $y^{x,u}$ is given by

$$\tau_\Gamma(x, u) = \inf \{ t \geq 0 : y^{x,u}(t) \in \Gamma \}.$$

Given any $y \in \Omega$, the *Minimum Time Problem* with target Γ is the following:

$$(MTP) \quad \text{minimize } \tau_\Gamma(x, u) \text{ over all controls } u : [0, +\infty[\rightarrow \overline{B}_1(0).$$

The *minimum time function* is defined by

$$T(x) = \inf_{u(\cdot)} \tau_\Gamma(x, u) \quad (x \in \overline{\Omega}).$$

It is well known that T is the unique viscosity solution of the Dirichlet problem (1.1). Moreover, Hörmander's bracket generating condition implies that (2.2) is small time locally controllable, so that T is finite and continuous (see, for instance, [5, Proposition 1.6, Chapter IV]).

We recall that a $u(\cdot)$ is called an *optimal control* at a point $x \in \Omega$ if $T(x) = \tau_\Gamma(x, u)$. The corresponding solution of (2.2), $y^{x,u}$, is called the *time-optimal trajectory* at x associated with u .

We now recall the definition of singular time-optimal trajectories. For any point $z \in \Gamma$, we denote by $\nu(z)$ the outward unit normal to Γ at z .

Definition 2.2. *We say that a time-optimal trajectory $y(\cdot) = y^{x,u}(\cdot)$ at a point $x \in \Omega$ is singular if there exists an absolutely continuous arc $p : [0, T(x)] \rightarrow \mathbb{R}^n \setminus \{0\}$ such that*

$$(2.3) \quad p'_k(t) = \sum_{j=1}^N u_j(t) \langle \partial_{x_k} X_j(y(t)), p(t) \rangle \quad t \in [0, T(x)] \text{ a.e.}$$

$$(k = 1, \dots, N),$$

$$(2.4) \quad \langle X_k(y(t)), p(t) \rangle = 0 \quad \forall t \in [0, T(x)] \quad (k = 1, \dots, N),$$

$$(2.5) \quad \exists \lambda > 0 : p(T(x)) = \lambda \nu(y(T(x))).$$

Notice that (2.4) and (2.5) imply that all the $X_j(y(T(x)))$'s are tangent to Γ , that is,

$$\text{span} \{ X_1(y(T(x))), \dots, X_N(y(T(x))) \} \subset T_\Gamma(y(T(x))).$$

So, $y(T(x))$ is a *characteristic point*.

In order to connect the lack of regularity of T with the presence of singular trajectories, it is useful to look at the Lipschitz singular set of T , i.e.,

$$\text{Sing}_L T = \left\{ x \in \Omega : \limsup_{\Omega \ni y \rightarrow x} \frac{|T(y) - T(x)|}{|y - x|} = \infty \right\}$$

which consists of all points at which T fails to be Lipschitz. Indeed, one can show that:

- (S1) $x \in \text{Sing}_L T$ if and only if x is the initial point of a singular trajectory ([3, Theorem 3.2]);
- (S2) $\text{Sing}_L T$ is closed in Ω ([3, Proposition 4.1]);
- (S3) T is locally semiconcave in $\Omega \setminus \text{Sing}_L T$ ([3, Theorem 4.3]).

We recall that a function is *semiconcave* if it can be locally represented as the sum of a smooth function plus a concave one.

Notice that property (S3) above ensures that $\text{Sing}_L T = \text{Sing supp}_{Lip} T$.

The fact that the existence of singular time-optimal trajectories may destroy the regularity of a solution of a first order Hamilton-Jacobi equation was observed (implicitly) by Sussmann in [13] and (explicitly) by Agrachev in [1]. The regularity these authors considered is subanaliticity of the point-to-point distance function associated with real-analytic distributions. The aforementioned subanaliticity results were extended to solutions of the Dirichlet problem in [14].

We recall that a vector $p \in \mathbb{R}^n$ is a *proximal subgradient* of T at $x \in \Omega$ if $\exists c, \rho > 0$ such that

$$(2.6) \quad T(y) - T(x) - \langle p, y - x \rangle \geq -c|y - x|^2, \quad \forall y \in B(x, \rho) \cap \Omega.$$

The set of all proximal subgradients of T at x is denoted by $\partial_P T(x)$.

The following lemma identifies proximal subdifferentiability as a threshold for local smoothness.

Lemma 2.1. *Let $x_0 \in \Omega$ be such that*

- a):** $\partial_P T(x_0)$ is nonempty,
- b):** T is semiconcave on an open neighbourhood $U_0 \subset \Omega$ of x_0 .

Then, T is of class C^∞ on some open neighbourhood $U \subset U_0$ of x_0 .

Proof. To begin with, we note that a) and b) force T to be differentiable at x_0 . Then, standard arguments based on sensitivity relations guarantee the existence of a unique optimal trajectory, $y_0(\cdot)$, starting from x_0 , and ensure that T stays differentiable along such a trajectory which, therefore, is not singular in view of (S1). So, by (S3), T is semiconcave on a relatively open neighbourhood, W_0 , of $\{y_0(t) : t \in [0, T_0]\}$, where we have set $T_0 = T(x_0)$. Thus, there exists a constant C_1 such that

$$(2.7) \quad \nabla^2 T \leq C_1 I$$

in the sense of distributions on W_0 . Moreover, by the propagation of proximal subdifferentiability (see [7, Theorem 3] or [9, Theorem 2.3]), a) implies that there exists a constant $C_2 \geq 0$ such that, for all $t \in [0, T_0[$ and $h \in \mathbb{R}^n$ sufficiently small,

$$(2.8) \quad T(y_0(t) + h) - T(y_0(t)) - \langle \nabla T(y_0(t)), h \rangle \geq -C_2 |h|^2.$$

The key idea of the proof is to deduce the local smoothness of T along $y_0(\cdot)$, in particular near x_0 , from [11, Theorem 3.1]. For this, we must prove that $\{y_0(t) : t \in [0, T_0]\}$ contains no conjugate points¹. In order to check such an assertion, we identify y_0 as a backward solution of the characteristic system as follows. Since $\xi_0 := y_0(T_0)$ is not a characteristic boundary point, there exists an open neighbourhood $V_0 \subset \Gamma$ of ξ_0 such that $H(\xi, \nu(\xi)) > 0$ for all $\xi \in V_0$, where $H(x, p) = \{\sum_{j=1}^N \langle p, X_j(x) \rangle^2\}^{1/2}$ is the Hamiltonian associated with $\{X_1, \dots, X_N\}$. For any $\xi \in V_0$, denote by $(X(\cdot, \xi), P(\cdot, \xi))$ the solution of

$$(2.9) \quad \begin{cases} -\dot{X} &= \nabla_p H(X, P), & X(0) &= \xi \\ \dot{P} &= \nabla_x H(X, P), & P(0) &= H(\xi, \nu(\xi))^{-1} \nu(\xi), \end{cases} \quad (t \geq 0)$$

defined on some maximal interval $[0, \tau_\xi[$, and by $X_{t,\xi}$ and $P_{t,\xi}$ the Jacobian of the maps X and P composed with a local parametrization of Γ (such matrix-valued functions solve a certain system of ODE's, i.e., the linearization of (2.9)). Observe that $\tau_\xi > T_0$ for all ξ in a suitable relatively open set $V \subset V_0$ because y_0 —coupled with a suitable dual arc p_0 —solves (2.9) backward in time for $\xi = \xi_0$, i.e.,

$$(X(t, \xi_0), P(t, \xi_0)) = (y_0(t - T_0), p_0(t - T_0)) \quad (t \in [0, T_0]).$$

So, proving that $y_0(\cdot)$ contains no conjugate point amounts to showing $\det X_{t,\xi}(t, \xi_0) \neq 0$ for all $t \in [0, T_0]$. If this is not the case, let $t_0 \in]0, T_0]$ be the first time at which $\det X_{t,\xi}(\cdot, \xi_0) = 0$. Then, by the classical method of characteristics, T is smooth at $X(t, \xi_0)$ and $\nabla T(X(t, \xi_0)) = -P(t, \xi_0)$ for all $t \in [0, t_0[$. So,

$$(2.10) \quad \nabla^2 T(X(t, \xi_0)) X_{t,\xi}(t, \xi_0) = -P_{t,\xi}(t, \xi_0), \quad \forall t \in [0, t_0].$$

Since, by well-known properties of solutions to linear systems (see, e.g., [6, p. 155]), $P_{t,\xi}$ can be singular at no point at which $\det X_{t,\xi} = 0$, from (2.10) it follows that

$$(2.11) \quad \lim_{t \nearrow t_0} |\det(\nabla T^2(X(t, \xi_0)))| = \infty.$$

Using the fact that for all $t \in [0, t_0[$ the left-hand side of (2.8) is equal to $\langle \nabla^2 T(X(t, \xi_0))h, h \rangle + o(|h|^2)$, we deduce that $\langle \nabla^2 T(X(t, \xi_0))h, h \rangle \geq -C_2 |h|^2 + o(|h|^2)$. Then, we conclude that there exists $C_2 > 0$ such that $\langle \nabla^2 T(X(t, \xi_0))\eta, \eta \rangle \geq -C_2$ for all $\eta \in S^{n-1}$ and $t \in [0, t_0[$. Finally, the last inequality, together with (2.7), yields that $\nabla^2 T(X(\cdot, \xi_0))$

¹Notice that, in [11], structural assumptions—that are not satisfied in our settings—are imposed. However, such assumptions are not needed for the proof of [11, Theorem 3.1].

is bounded on $[0, t_0[$, in contrast with (2.11), completing the proof. \square

Proof of Theorem 1.1. Let $\Sigma_1(T) = \text{Sing supp}_{C^{1,1}} T$ and $\Sigma(T) = \text{Sing supp } T$. Since $\Sigma_1(T) \subseteq \Sigma(T)$, we just need to show that $\Omega \setminus \Sigma_1(T) \subseteq \Omega \setminus \Sigma(T)$. As mentioned above, T is semiconcave on $\Omega \setminus \Sigma_1(T)$. Moreover, from the very definition (2.6) of proximal subgradients it follows that $\partial_P T(x) \neq \emptyset$ for any $x \in \Omega \setminus \Sigma_1(T)$. Then, the conclusion follows from Lemma 2.1. \square

Proof of Theorem 1.2. We keep the notation $\Sigma_1(T)$ of the previous proof and set $\Sigma_{Lip}(T) = \text{Sing supp}_{Lip} T$. We observe that, by Theorem 1.1, it suffices to show that the $C^{1,1}$ singular support of T has null measure. For this purpose we decompose such a support as $\Sigma_1(T) = \Sigma_{Lip}(T) \cup (\Sigma_1(T) \setminus \Sigma_{Lip}(T))$. By [10, Corollary 3.3], we deduce that $\Sigma_{Lip}(T)$ has measure zero. In order to prove that $\Sigma_1(T) \setminus \Sigma_{Lip}(T)$ has measure zero we use an idea from [2]. Recall that, by [3, Theorem 4.1], T is locally semiconcave in $\Omega \setminus \Sigma_{Lip}(T)$. Then, Alexandroff Theorem (see [4]) guarantees that T has a second order Taylor expansion at a.e. point of $\Omega \setminus \Sigma_{Lip}(T)$. Hence, $\partial_P T(x)$ is nonempty for a.e. $x \in \Omega \setminus \Sigma_{Lip}(T)$. So, thanks to Lemma 2.1 we conclude that there exists a set of full measure in $\Omega \setminus \Sigma_{Lip}(T)$ which lies in the complement of $\Sigma_1(T) \setminus \Sigma_{Lip}(T)$. This proves that the set $\Sigma_1(T) \setminus \Sigma_{Lip}(T)$ has null measure and completes the proof. \square

ACKNOWLEDGEMENTS

The authors are grateful to the referee for her/his careful reading and useful comments.

REFERENCES

- [1] A.AGRACHEV, *Compactness for sub-Riemannian length-minimizers and sub-analyticity*, Rend. Sem. Mat. Univ. Politec. Torino 56 (1998), no.4, 1-12.
- [2] P.ALBANO, *On the cut locus of a closed set*, Nonlinear Anal. 125 (2015), 398-405.
- [3] P.ALBANO, P.CANNARSA AND T.SCARINCI, *Regularity results for the minimum time function with Hörmander vector fields*, J. Differential Equations 264 (2018), no. 5, 3312-3335.
- [4] A.D.ALEXANDROFF, *Almost everywhere existence of the second differential of a convex function and some properties of convex surfaces connected with it* (Russian) Leningrad State Univ. Annals [Uchenye Zapiski] Math. Ser. 6 (1939), 3-35.
- [5] M. BARDI AND I. CAPUZZO DOLCETTA, *Optimal control and viscosity solutions of Hamilton-Jacobi-Bellman equations*. Birkhäuser Boston, Inc., Boston, MA, 1997.

- [6] P.CANNARSA AND C.SINESTRARI, Semiconcave functions, Hamilton-Jacobi equations, and optimal control. *Progress in Nonlinear Differential Equations and their Applications*, 58, Birkhäuser Boston, Inc., Boston, MA, 2004.
- [7] P.CANNARSA AND T.SCARINCI, *Conjugate times and regularity of the minimum time function with differential inclusions*. Analysis and geometry in control theory and its applications, 85–110, Springer INdAM Ser., 11, Springer, Cham, 2015.
- [8] M.DERRIDJ, *Sur un théorème de traces*, Ann. Inst. Fourier (Grenoble) 22 (1972), 73-83.
- [9] H. FRANKOWSKA, L. V. NGUYEN, *Local Regularity of the Minimum Time Function*, J. Optim. Theory Appl. 164 (2015), no.1, 68-91.
- [10] K.T.NGUYEN *Hypographs satisfying an external sphere condition and the regularity of the minimum time function*, J. Math. Anal. Appl. 372 (2010), no. 2, 611-628.
- [11] C. PIGNOTTI *Rectifiability results for singular and conjugate points of optimal exit time problems*, J. Math. Appl. 270 (2002), no. 2, 681-708.
- [12] L.RIFFORD AND E.TRÉLAT, *Morse-Sard type results in sub-Riemannian geometry*, Math. Ann. 332 (2005), no. 1, 145-159.
- [13] H.J.SUSSMANN, *Optimal control theory and piecewise analyticity of the distance function for some real-analytic sub-Riemannian metrics*. Optimization and nonlinear analysis (Haifa, 1990), 298–310, Pitman Res. Notes Math. Ser., 244, Longman Sci. Tech., Harlow, 1992.
- [14] E.TRÉLAT, *Global subanalytic solutions of Hamilton-Jacobi type equations*, Ann. Inst. H. Poincaré Anal. Non Linéaire 23 (2006), no.3, 363-387.

DIPARTIMENTO DI MATEMATICA, UNIVERSITÀ DI BOLOGNA, PIAZZA DI PORTA SAN DONATO 5, 40127 BOLOGNA, ITALY

E-mail address: `paolo.albano@unibo.it`

DIPARTIMENTO DI MATEMATICA, UNIVERSITÀ DI ROMA "TOR VERGATA", VIA DELLA RICERCA SCIENTIFICA 1, 00133 ROMA, ITALY

E-mail address: `cannarsa@mat.uniroma2.it`

DEPARTMENT OF STATISTIC AND OPERATION RESEARCH, UNIVERSITY OF VIENNA, OSKAR-MORGENSTERN-PLATZ 1, 1090 VIENNA, AUSTRIA

E-mail address: `teresa.scarinci@gmail.com`