

# PLANE CURVES POSSESSING TWO OUTER GALOIS POINTS

SATORU FUKASAWA AND PIETRO SPEZIALI

**ABSTRACT.** We classify plane curves  $\mathcal{C}$  possessing two Galois points  $P_1$  and  $P_2 \in \mathbb{P}^2 \setminus \mathcal{C}$  such that the associated Galois groups  $G_{P_1}$  and  $G_{P_2}$  generate the semidirect product  $G_{P_1} \rtimes G_{P_2}$ . New examples of plane curves with two Galois points are also presented.

## 1. INTRODUCTION

A classical problem in Algebraic Geometry is the classification of algebraic varieties. Birational invariants are classical tools for the classification of projective (non-singular) algebraic curves  $\mathcal{C}$  up to birational or projective equivalence. In some cases, the genus  $g$  and the automorphism group  $\text{Aut}(\mathcal{C})$  completely determine a curve; see [1, 6]. More recently, the number of Galois points of  $\mathcal{C}$  has been successfully used for the classification problem ([2, 3, 18, 20]).

Let  $K$  be an algebraically closed field of characteristic  $p \geq 0$ . A point  $P \in \mathbb{P}^2$  is called a Galois point for a plane curve  $\mathcal{C} \subset \mathbb{P}^2$  with function field  $K(\mathcal{C})$ , if the finite field extension  $K(\mathcal{C})/\pi_P^*K(\mathbb{P}^1)$  induced by the projection  $\pi_P$  from  $P$  is a Galois extension ([13, 18]). The associated Galois group of  $P$  is denoted by  $G_P$ . Furthermore, a Galois point  $P$  is said to be outer, if  $P \in \mathbb{P}^2 \setminus \mathcal{C}$ .

In this article, we classify plane curves  $\mathcal{C}$  possessing two outer Galois points  $P_1$  and  $P_2 \in \mathbb{P}^2 \setminus \mathcal{C}$  such that associated groups  $G_{P_1}$  and  $G_{P_2}$  generate  $G_{P_1} \rtimes G_{P_2}$ . Determining the structure of the group generated by  $G_{P_1}$  and  $G_{P_2}$  seems a rather challenging task. On the one hand, by [3, Lemma 7]  $G_{P_1}$  and  $G_{P_2}$  have trivial intersection. On the other hand, in general it is not possible to embed the group  $\langle G_{P_1}, G_{P_2} \rangle$  into  $\text{PGL}(3, K)$ ; see for instance [18, Remark 1] and [11, Proposition 4]. Further, it might be that  $G = \langle G_{P_1}, G_{P_2} \rangle \neq G_{P_1} \rtimes G_{P_2}$  but  $G$  is linear, as the following example shows.

**Example 1.1.** For  $p > 0$ , let  $q = p^h$  be a power of  $p$ . Also, let  $\mathcal{H}$  be the Hermitian curve with affine equation  $x^{q+1} = y^q + y$ . By [7, Theorems 1 and 2], any point in  $\mathbb{P}^2(\mathbb{F}_{q^2}) \setminus \mathcal{H}$  is an outer Galois point for  $\mathcal{H}$  with Galois group isomorphic to a cyclic group  $C_{q+1}$ . Take two points  $P_1, P_2$  lying on the same tangent to  $\mathcal{H}$  at a point  $P$ . Clearly  $\langle G_{P_1}, G_{P_2} \rangle$ , being a subgroup of the stabilizer  $\text{Aut}(\mathcal{H})_P$  of  $P$  in  $\text{Aut}(\mathcal{H}) \cong \text{PGU}(3, q)$ , is linear. Assume that  $\langle G_{P_1}, G_{P_2} \rangle$  is a (semi)direct product. By [6, Theorem A.10],  $|\text{Aut}(\mathcal{H})_P| = q^3(q^2 - 1)$ , a contradiction.

Our main result is the following Theorem.

**Theorem 1.2.** Let  $\mathcal{C}$  be a plane curve of degree  $d \geq 3$  defined over an algebraically closed field  $K$  of characteristic  $p \geq 0$ . Assume that  $p = 0$  or  $d$  is prime to  $p$ . Then  $\mathcal{C}$  possesses two outer Galois

---

2020 *Mathematics Subject Classification.* 14H37, 14H05.

*Key words and phrases.* Algebraic curves; Automorphism groups; Galois points.

The first author was partially supported by JSPS KAKENHI Grant Numbers JP16K05088 and JP19K03438.

The second author was partially supported by FAPESP-Brazil, grant 2017/18776-6.

points  $P_1$  and  $P_2$  such that the set  $G_{P_1}G_{P_2}$  is the semidirect product  $G_{P_1} \rtimes G_{P_2}$ , if and only if  $\mathcal{C}$  is projectively equivalent to the curve defined by one of the following:

- (a)  $x^d + y^d + 1 = 0$ ;
- (b)  $x^d + x^{d/2}(x^{d/2} + 1) = 0$ , where  $d$  is even;
- (c)  $y^d + x^2(x^2 + 1)g(x^2)^2 = 0$ , where  $d$  and  $d/2$  are even, and  $g = \sum a_i x^i \in K[x]$  is a separable polynomial of degree  $d/4 - 1$  determined by the relations

$$a_{d/4-1} = 1, \quad \text{and} \quad \frac{d^2}{2}a_i = 8(i+1)^2a_i + 4(i+1)(2i+3)a_{i+1};$$

- (d)  $y^d + (x^2 + 1)g(x^2)^2 = 0$ , where  $d$  and  $d/2 - 1$  are even, and  $g = \sum a_i x^i \in K[x]$  is a separable polynomial of degree  $(d/2 - 1)/2$  determined by the relations

$$a_{(d-2)/4} = 1, \quad \text{and} \quad \left(\frac{d}{2}\right)^2 a_i = (2i+1)^2a_i + (2i+1)(2i+2)a_{i+1}.$$

For curves in Case (b), (c) or (d), the following hold:

- (1) The genus is  $(d-2)^2/4$ ;
- (2) In Case (b) (resp. Cases (c) and (d)), the number of singular points is one (resp. is  $d/2-1$ );
- (3) If  $d$  is even and  $d/2$  is even (resp.  $d/2$  is odd), then the nonsingular models of curves in Cases (b) and (c) (resp. in Cases (b) and (d)) are isomorphic;
- (4)  $G_{P_1}$  is a dihedral group and  $G_{P_2}$  is a cyclic group.

In particular, if  $d$  is odd, then the Fermat curve is a unique example. On the other hand, the following holds.

**Corollary 1.3.** *Let  $\mathcal{C}$  be a plane curve of degree  $d \geq 3$ . Assume that  $p = 0$  or  $d$  is prime to  $p$ . Then there exist two outer Galois points  $P_1$  and  $P_2$  such that  $G_{P_1}G_{P_2} = G_{P_1} \times G_{P_2}$ , if and only if  $\mathcal{C}$  is projectively equivalent to the Fermat curve  $X^d + Y^d + Z^d = 0$ .*

Plane curves described in Case (c) or in Case (d) in Theorem 1.2 are new examples of plane curves with two outer Galois points (see the Table in [20]).

## 2. PROOFS

The system of homogeneous coordinates of  $\mathbb{P}^2$  is denoted by  $(X : Y : Z)$ , and the system of affine coordinates with  $Z \neq 0$  is denoted by  $(x, y)$ , where  $x = X/Z$ ,  $y = Y/Z$ . The nonsingular model of  $\mathcal{C}$  is denoted by  $\hat{\mathcal{C}}$  and its genus is denoted by  $g$ . For two different points  $R_1$  and  $R_2 \in \mathbb{P}^2$ , the line passing through  $R_1$  and  $R_2$  is denoted by  $\overline{R_1R_2}$ . When  $G_1, G_2$  are subgroups of the full automorphism group  $\text{Aut}(\hat{\mathcal{C}})$ , we define a subgroup

$$C_{G_1}(G_2) := \{\sigma \in G_1 \mid \sigma\tau = \tau\sigma, \text{ for all } \tau \in G_2\}$$

of  $G_1$ .

First, we confirm (some part of) the converse assertion of Theorem 1.2. The following proposition regarding the Fermat curve is known, by Miura and Ohbuchi [12] (see also [1]).

**Proposition 2.1.** *For the Fermat curve  $X^d + Y^d + Z^d = 0$ , points  $P_1 = (1 : 0 : 0)$  and  $P_2 = (0 : 1 : 0)$  are outer Galois points such that  $G_{P_1} \cong C_d$ ,  $G_{P_2} \cong C_d$ , and  $G_{P_1}G_{P_2} = G_{P_1} \rtimes G_{P_2} = G_{P_1} \times G_{P_2}$ , where  $C_d$  is the cyclic group of order  $d$ .*

Another example of a plane curve with two Galois points is the following, which was studied by Takahashi [16].

**Proposition 2.2.** *Let  $\mathcal{C}$  be the curve  $X^d + X^{d/2}Z^{d/2} + Y^d = 0$ , and let  $P_1 = (1 : 0 : 0)$  and  $P_2 = (0 : 1 : 0)$ . Then points  $P_1, P_2$  are outer Galois points such that  $G_{P_1} \cong D_d$ ,  $G_{P_2} \cong C_d$ , and  $G_{P_1}G_{P_2} = G_{P_1} \rtimes G_{P_2}$ , where  $D_d$  is the dihedral group of order  $d$ .*

*Proof.* It is easily verified that  $P_2$  is an outer Galois point with  $G_{P_2} \cong C_d$ , since the linear maps  $(x, y) \mapsto (x, \lambda y)$  with  $\lambda^d = 1$  preserve the curve and they form a cyclic group of order  $d$ . According to a work of Takahashi [16],  $P_1$  is an outer Galois point with  $G_{P_1} \cong D_d$ . In fact, the two rational maps  $(x, y) \mapsto (\eta x, y)$ , where  $\eta$  is a primitive  $d/2$ -th root of unity, and  $(x, y) \mapsto (y^2/x, y)$  act on  $\mathcal{C}$  and preserve the fibers of  $\pi_{P_1}$ . A direct computation shows that  $\langle G_{P_1}, G_{P_2} \rangle = G_{P_1} \rtimes G_{P_2}$ .  $\square$

Takahashi's curve  $X^d + X^{d/2}Z^{d/2} + Y^d = 0$  admits another plane model with two outer Galois points. More precisely:

**Proposition 2.3.** *Let  $d \geq 6$  be even, and let  $\hat{\mathcal{C}}$  be the nonsingular model of the curve  $\mathcal{C} : X^d + X^{d/2}Z^{d/2} + Y^d = 0$ . Then there exist a morphism  $r' : \hat{\mathcal{C}} \rightarrow \mathbb{P}^2$  and points  $P'_1, P'_2 \in \mathbb{P}^2 \setminus r'(\hat{\mathcal{C}})$  such that the following hold:*

- (a) *The morphism  $r' : \hat{\mathcal{C}} \rightarrow \mathcal{C}' := r'(\hat{\mathcal{C}})$  is birational, and  $\deg \mathcal{C}' = d$ .*
- (b) *Points  $P'_1, P'_2$  are outer Galois points.*
- (c)  *$G_{P'_1} \cong D_d$ ,  $G_{P'_2} \cong C_d$ , and  $G_{P'_1}G_{P'_2} = G_{P'_1} \rtimes G_{P'_2}$ .*
- (d)  *$|C_{G_{P'_1}}(G_{P'_2})| = 2$ .*

*Proof.* Let  $r : \hat{\mathcal{C}} \rightarrow \mathcal{C}$  be the normalization,  $P_1 = (1 : 0 : 0) \in \mathbb{P}^2$  for the plane model  $\mathcal{C}$ , and let

$$\sigma(x, y) = (\eta x, y), \quad \tau(x, y) = \left( \frac{y^2}{x}, y \right),$$

where  $\eta$  is a primitive  $d/2$ -th root of unity. As we saw the previous Proposition,  $P_1$  is an outer Galois point with  $G_{P_1} = \langle \sigma, \tau \rangle \cong D_d$ . We take a point  $Q_\mu = (\mu : 1 : 0)$  such that  $\mu \in K$  and  $\mu^d = -1$ . Let

$$\sigma'(x, y) = (\eta x, \eta y), \quad \tau'(x, y) = \left( \frac{y^2}{x}, \frac{y}{\mu^2} \right),$$

and let  $G'_2 = \langle \sigma', \tau' \rangle$ . Since  $\sigma'\tau' = \tau'\sigma'$ , it follows that  $|G'_2| = d$ . Note that the group  $G'_2$  fixes the point  $Q_\mu$ . This implies that  $G'_2 \cong C_d$  (see, for example, [6, Lemma 11.44]). It can be confirmed that  $\sigma'\sigma = \sigma\sigma'$ ,  $\sigma'\tau = \tau\sigma'$ , and  $\tau'\sigma = \sigma^{-1}\tau'$ . Since

$$\tau'\tau(x, y) = \left( x, \frac{y}{\mu^2} \right), \quad \tau\tau' = \left( \frac{x}{\mu^4}, \frac{y}{\mu^2} \right),$$

it follows that  $\tau'\tau = \sigma^m \tau \tau'$ , where  $\eta^m = \mu^4$ . These imply that  $G_{P_1}G'_2 = G_{P_1} \rtimes G'_2$ .

We prove that  $|C_{G_{P_1}}(G_{P'_2})| = 2$ . Assume that  $\sigma^n \in C_{G_{P_1}}(G'_2) \setminus \{1\}$ . Since

$$\sigma^n \tau' = \left( \frac{\eta^n y^2}{x}, \frac{y}{\mu^2} \right), \quad \tau' \sigma^n = \left( \frac{y^2}{\eta^n x}, \frac{y}{\mu^2} \right),$$

it follows that  $\eta^{2n} = 1$ . This implies that  $d/2$  is even and  $n = d/4$ . Assume that  $\sigma^n \tau \in C_{G_{P_1}}(G'_2)$ . Since

$$\sigma^n \tau \tau' = \left( \frac{\eta^n x}{\mu^4}, \frac{y}{\mu^2} \right), \quad \tau' \sigma^n \tau = \left( \frac{x}{\eta^n}, \frac{y}{\mu^2} \right),$$

it follows that  $\eta^{2n} = \mu^4$ . If  $d/2$  is even, then  $1 = (\eta^{2n})^{d/4} = (\mu^4)^{d/4} = -1$ . This implies that  $p = 2$  and  $d$  is divisible by  $p$ . This is a contradiction. Therefore,  $d/2$  is odd. Then  $\eta^n = -\mu^2$ . Whether  $d/2$  is even or odd, it follows that  $|C_{G_{P_1}}(G'_2)| = 2$ .

Since  $G'_2$  fixes  $Q_\mu$  and any element of  $G_{P_1}$  does not fix  $Q_\mu$ , it follows that  $G_{P_1} \cap G'_2 = \{1\}$ . Since  $k(\mathcal{C})^{(\sigma')} = k(y/x) \supset k(\mathcal{C})^{G'_2}$ , it follows from Lüroth's theorem that  $k(\mathcal{C})^{G'_2}$  is rational. Let  $Q \in r^{-1}(\mathcal{C} \cap \{Y = 0\})$ . It can be confirmed that

$$\sum_{\sigma \in G_{P_1}} \sigma(Q) = \sum_{Q' \in r^{-1}(\mathcal{C} \cap \{Y=0\})} Q' = \sum_{\eta \in G'_2} \eta(Q).$$

It follows from a theorem [4, Theorem 1 and Remark 1] of the first author that there exist a birational morphism  $r' : \hat{\mathcal{C}} \rightarrow \mathbb{P}^2$  of degree  $d$  onto its image and points  $P'_1, P'_2 \in \mathbb{P}^2 \setminus r'(\hat{\mathcal{C}})$  such that  $G_{P'_1} = G_{P_1}$  and  $G_{P'_2} = G'_2$ . Conditions (c) and (d) are satisfied for the groups  $G_{P'_1}$  and  $G_{P'_2}$ , according to the discussion above.  $\square$

*Proof of Theorem 1.2.* By Propositions 2.1 and 2.2, the converse assertions in Cases (a) and (b) are obvious. Proposition 2.3 and the former assertion in Theorem 1.2 imply the converse assertions in Cases (c) and (d), since we show later that the projective equivalence class is uniquely determined by the degree  $d$  and the order  $|C_{G_{P_1}}(G_{P_2})|$ . We prove the former assertion in eight steps.

*Step 1:  $G_{P_2}$  is a cyclic group.* The projection  $\pi_{P_1}$  from  $P_1$  induces the covering  $\hat{\mathcal{C}} \rightarrow \mathbb{P}^1 \cong \hat{\mathcal{C}}/G_{P_1}$ . The covering  $\pi_{P_1}(\mathcal{C}) \rightarrow \hat{\mathcal{C}}/G_{P_1}G_{P_2}$  is denoted by  $f_{P_1}$ . By [4, Theorem 1 (c')], there exists a point  $Q \in \hat{\mathcal{C}}$  such that

$$(1) \quad \sum_{\sigma \in G_{P_1}} \sigma(Q) = \sum_{\tau \in G_{P_2}} \tau(Q)$$

as divisors. By this equation, for any  $[\tau] \in G_{P_1}G_{P_2}/G_{P_1}$ ,  $[\tau][Q] = [\tau(Q)] = [Q]$ , where  $[Q] = \pi_{P_1}(Q) \in \hat{\mathcal{C}}/G_{P_1}$  is a class containing  $Q$ . It follows from [15, Theorem 3.8.2] that  $f_{P_1}$  is totally ramified at  $\pi_{P_1}(Q)$ . Since  $f_{P_1}$  is a Galois covering and  $\pi_{P_1}(\mathcal{C}) \cong \mathbb{P}^1$ ,  $(G_{P_1} \rtimes G_{P_2})/G_{P_1} \cong G_{P_2}$  is a cyclic group of order  $d$  (see, for example, [17, Theorem 1]).

*Step 2: The defining equation.* Let  $R \in \pi_{P_1}(\mathcal{C}) \cong \mathbb{P}^1$  be another ramification point and let  $\ell \subset \mathbb{P}^2$  be the line corresponding to  $R$ . For a suitable system of coordinates, we can assume that  $P_1 = (1 : 0 : 0)$ ,  $P_2 = (0 : 1 : 0)$  and  $\ell$  is given by  $Y = 0$ . Note that  $\pi_{P_1}$  and  $\pi_{P_2}$  are represented by  $(X : Y : Z) \mapsto (Y : Z)$  and  $(X : Y : Z) \mapsto (X : Z)$  respectively. Since  $f_{P_1}$  is totally ramified at  $\pi_{P_1}(Q)$  and  $R$ , which correspond to the points  $(1 : 0)$  and  $(0 : 1)$ ,  $f_{P_1} = (aY^d : bZ^d)$  for some  $a, b \in K \setminus \{0\}$ . Since the function field of  $\hat{\mathcal{C}}/(G_{P_1} \rtimes G_{P_2})$  is contained in  $K(\hat{\mathcal{C}}/G_{P_2}) = K(x)$ ,  $y^d = c(x)/d(x)$  for some relatively prime polynomials  $c(x)$  and  $d(x) \in K[x]$ . Since the curve  $\mathcal{C}$  is of degree  $d$ ,  $d(x) \in K$  and the equation  $y^d - c(x) = 0$  is the defining one. The degree of  $c(x)$  is  $d$ , since  $P_1 \in \mathbb{P}^2 \setminus \mathcal{C}$ . The group  $G_{P_2}$  consists of all linear maps  $(x, y) \mapsto (x, \lambda y)$  with  $\lambda^d = 1$ .

*Step 3: The genus  $g$ .* Note that  $\pi_{P_1}$  is not ramified at any point in  $\mathcal{C} \cap \overline{P_1 P_2}$ , since  $\pi_{P_2}$  is not ramified at these points. By the Riemann–Hurwitz formula, there exists a ramification point  $S \in \mathcal{C} - (\overline{P_1 P_2} \cup \ell)$  for the projection  $\pi_{P_1}$ . Let  $m$  be the ramification index of  $S$ . By a property

of Galois coverings [15, Corollary 3.7.2], there exist  $(d/m)$  points with ramification index  $m$  in the line  $\overline{P_1S}$ . Note that any element of  $G_{P_2}$  is the restriction of some linear transformation. By the action of  $G_{P_2}$ , there exist  $d \times (d/m)$  points with index  $m$ . By the Riemann–Hurwitz formula, we get

$$2g - 2 \geq d(-2) + d \times \left( d - \frac{d}{m} \right).$$

If there exists a ramification point not contained in  $\{\eta(S) \mid \eta \in G_{P_1}G_{P_2}\} \cup \ell$ , then  $g \geq (d^2 - 2d + 2)/2$ , but this is a contradiction to  $g \leq (d - 1)(d - 2)/2$ . Therefore,  $\{\eta(S) \mid \eta \in G_{P_1}G_{P_2}\}$  is the set of all ramification points in  $\mathcal{C} \setminus \ell$ .

Let  $n$  be the number of irreducible components of  $c(x)$  and let

$$-c(x) = \gamma \prod_{i=1}^n (x - a_i)^{e_i},$$

where  $\gamma \in K$ ,  $a_1, \dots, a_n \in K$  are pairwise distinct and  $\sum_{i=1}^n e_i = d$ . By [15, Proposition 3.7.3],

$$2g - 2 + 2d = \sum_{i=1}^n (d - r_i),$$

where  $r_i$  is the greatest common divisor of  $d$  and  $e_i$ . Note that  $d - r_i \leq d - 1$ . By the lower bound of the genus  $g$  above,  $n \geq d - d/m + 1$ . In particular, the set  $\mathcal{C} \cap \ell$  consists of  $n > d/2$  points and hence,  $\pi_{P_1}$  is not ramified at each point of  $\mathcal{C} \cap \ell$ , by [15, Corollary 3.7.2]. It follows that

$$2g - 2 + 2d = d \left( d - \frac{d}{m} \right).$$

*Step 4: The number of points on the line  $\ell$  and their multiplicities.* If  $x - a_i$  is a component of  $c(x)$ , then the line defined by  $x - a_i = 0$  intersects  $\mathcal{C}$  at only the point  $(a_i, 0)$ . The fiber of  $(a_i, 0)$  under the normalization consists of exactly  $e_i$  points, since  $\pi_{P_1}$  is unramified at such points. Since  $\pi_{P_2}$  is a Galois covering and  $x - a_i$  gives a fiber of it, it follows that  $e_i$  divides  $d$ . This implies that  $r_i = e_i$ . Furthermore, by the Riemann–Hurwitz formula,

$$2g - 2 + 2d = \sum_{i=1}^n (d - r_i) = nd - \sum_i e_i = nd - d = d(n - 1).$$

Since  $2g - 2 + 2d = d(d - d/m)$  as considered above, it follows that

$$n = d - d/m + 1.$$

Let  $y = \beta$  denote a branch point of  $\pi_{P_1}$ . Then there exists a separable polynomial  $h(x)$  of degree  $d/m$  such that

$$-c(x) + \beta^d = h(x)^m.$$

Then

$$-c(x) = \prod_{k=0}^{m-1} (h(x) - \zeta^k \beta^{\frac{d}{m}}),$$

where  $\zeta$  is a primitive  $m$ -th root of unity. If  $x - a$  is a multiple component of  $c(x)$ , then  $h(a) - \zeta^k \beta^{\frac{d}{m}} = 0$  for some  $k$ , and  $h'(a) = 0$ . This implies that  $e_i \leq d/m$ .

*Step 5: The number of nonsingular points on  $\ell$ .* Recall that

$$C_{G_{P_1}}(G_{P_2}) = \{\sigma \in G_{P_1} \mid \sigma\tau = \tau\sigma, \text{ for all } \tau \in G_{P_2}\}.$$

We prove that for a nonsingular point  $Q \in \mathcal{C} \cap \ell$ ,

$$(C_{G_{P_1}}(G_{P_2}))Q = (\mathcal{C} \setminus \text{Sing}(\mathcal{C})) \cap \ell.$$

Assume by contradiction that there exists  $\sigma \in C_{G_{P_1}}(G_{P_2})$  such that  $\sigma(Q)$  is a singular point. Let  $\tau \in G_{P_2}$  be an automorphism such that  $\tau^{-1}(\sigma(Q)) \neq \sigma(Q)$  (see [15, Theorem 3.7.1]). Since  $\tau(Q) = Q$ ,  $\tau^{-1}\sigma\tau(Q) = \tau^{-1}(\sigma(Q)) \neq \sigma(Q)$ . On the other hand, since  $\sigma \in C_{G_{P_1}}(G_{P_2})$ , it follows that  $\tau^{-1}\sigma\tau(Q) = \sigma(Q)$ . This is a contradiction.

Let  $Q'$  be a nonsingular point on  $\ell$ . It follows from [15, Theorem 3.7.1] that there exists  $\sigma \in G_{P_1}$  such that  $\sigma(Q) = Q'$ . Let  $\tau \in G_{P_2}$ . Since  $Q$  and  $Q'$  are nonsingular points,  $\tau(Q) = Q$  and  $\tau(Q') = Q'$ . Then  $\tau^{-1}\sigma\tau(Q) = \tau^{-1}\sigma(Q) = \tau^{-1}(Q') = Q' = \sigma(Q)$ . Since the stabilizer subgroup of  $G_{P_1}$  of any point on the line  $\ell$  is trivial (see [15, Theorem 3.8.2]), it follows that  $\tau^{-1}\sigma\tau = \sigma$ .

*Step 6:  $m \geq 3$  implies that  $\mathcal{C}$  is the Fermat curve.* Assume that  $m \geq 4$ . For the line  $\ell$ , there exist at least

$$n - \deg h'(x) \geq d - d/m + 1 - (d/m - 1) = d - (2d/m) + 2$$

nonsingular points. Since  $m \geq 4$ , it follows that

$$d - (2d/m) + 2 \geq d/2 + 2.$$

It follows from Step 5 that all points on  $\ell$  are nonsingular and  $G_{P_1}G_{P_2} = G_{P_1} \times G_{P_2}$ . By Steps 2 and 3, it follows that  $m = d$ . It follows from Step 4 that  $\mathcal{C}$  is projectively equivalent to the Fermat curve.

Assume that  $m = 3$ . Then there exist at least  $d/3 + 2$  nonsingular points on the line  $\ell$ . By the discussion above for  $m \geq 4$ ,  $|C_{G_{P_1}}(G_{P_2})| = d/2$  or  $d$ . We prove that any  $\sigma \in C_{G_{P_1}}(G_{P_2})$  is the restriction of some linear transformation of  $\mathbb{P}^2$ . Let  $\Lambda$  be a (base-point-free) linear system of dimension 2 induced by the normalization  $r : \hat{\mathcal{C}} \rightarrow \mathcal{C} \subset \mathbb{P}^2$ . Then  $r^*\overline{P_1P_2}, r^*\ell \in \Lambda$ . Since  $P_1$  is outer Galois, it follows that

$$\sigma^*r^*\overline{P_1P_2} = r^*\overline{P_1P_2}, \sigma^*r^*\ell = r^*\ell.$$

Let  $Q$  be a nonsingular point on  $\ell$ . By Step 5,  $\sigma(Q)$  is also a nonsingular point on  $\ell$ . Let  $T_Q$  and  $T_{\sigma(Q)}$  be tangent lines at  $Q$  and  $\sigma(Q)$  respectively. Then

$$\sigma^*(r^*T_{\sigma(Q)}) = \sigma^*(d\sigma(Q)) = d(\sigma^*\sigma(Q)) = dQ = r^*T_Q.$$

Since  $\Lambda$  is the smallest linear system containing  $r^*\overline{P_1P_2}$ ,  $r^*\ell$  and  $r^*T_{\sigma(Q)}$ , it follows that  $\sigma^*\Lambda = \Lambda$ .

Assume that  $|C_{G_{P_1}}(G_{P_2})| = d/2$ . Since  $C_{G_{P_1}}(G_{P_2})$  acts on all lines passing through  $P_1$ ,  $C_{G_{P_1}}(G_{P_2})$  fixes  $P_1$  and acts on  $\ell$  as a cyclic group of order  $d/2$ . Let  $R$  be a singular point on  $\ell$ . If  $R$  is not a fixed point of  $C_{G_{P_1}}(G_{P_2})$ , then there exist  $d/2$  singular points on  $\ell$ . This is a contradiction. Therefore, on the line  $\ell$ , there exist exactly  $d/2$  nonsingular points and one singular point. This is a contradiction, since there exist exactly

$$d - d/m + 1 = (2d)/3 + 1$$

points on  $\ell$  for  $m = 3$ . Therefore,  $|C_{G_{P_1}}(G_{P_2})| = d$  and  $\mathcal{C}$  is projectively equivalent to the Fermat curve.

*Step 7: The case where  $m = 2$  and  $C_{G_{P_1}}(G_{P_2})$  fixes a point of  $\mathcal{C} \cap \ell$ .* Let  $f = |C_{G_{P_1}}(G_{P_2})| \geq 2$ . As in Step 6, any element of  $C_{G_{P_1}}(G_{P_2})$  is the restriction of some linear transformation of  $\mathbb{P}^2$ . Note

that there exists a line  $\ell' \subset \mathbb{P}^2 \setminus \{P\}$  of which points are fixed by  $C_{G_{P_1}}(G_{P_2})$  (see, for example, [11, 19]). Assume that  $\ell' \cap \ell = \{R\} \subset \mathcal{C}$ . It follows that  $k \times f + 1 = d/2 + 1$  for some  $k$ , by considering the number of points on  $\ell$ . This implies that  $2k = d/f$ .

We can assume that for a suitable system of coordinates,  $R = (0 : 0 : 1)$  and a generator of  $C_{G_{P_1}}(G_{P_2})$  is represented by  $(X : Y : Z) \mapsto (\zeta X : Y : Z)$ , where  $\zeta$  is a primitive  $f$ -th root of unity. It follows that the multiplicity of  $R$  is greater than or equal to  $f$ . Then nonsingular points on  $\ell$  are defined by  $x^f - a^f = 0$  for some  $a \in K$ . We can assume that  $a^f = -1$ . Since  $d/2 + 1 - (f + 1) = d/2 - f$  singular points exist except for  $R$ , and  $-c(x) = h^2 - \beta^d$  has at most  $d - 2f$  components which define singular points on  $\ell$  except for  $R$ , it follows that the multiplicity of singular points except for  $R$  is two and that of  $R$  is equal to  $f$ . Therefore,  $h(x)^2 - \beta^d$  is of the form

$$x^f(x^f + 1) \prod_{i=1}^{d/(2f)-1} (x^f - a_i)^2$$

for some pairwise distinct elements  $a_i \in K$ . It follows that there exists a separable polynomial  $g$  of degree  $d/(2f) - 1$  such that

$$g(x^f) = \prod_{i=1}^{d/(2f)-1} (x^f - a_i).$$

We prove that  $f = 2$  or  $d/2$ . Assume that  $2 < f < d/2$ . For a ramification point  $S$  of the projection  $\pi_{P_1}$ , the orbit of  $S$  under  $G_{P_1}G_{P_2}$  is denoted by  $G \cdot S$ , which coincides with the set of all ramification points for  $\pi_{P_1}$ . It follows from [15, Remark 4.3.7] that

$$(dy) = -2(y)_\infty + \sum_{R \in G \cdot S} R,$$

by considering the projection  $\pi_{P_1}(x, y) = y$ . Let  $D$  be the divisor of degree  $d$  coming from  $\mathcal{C} \cap \{Z = 0\}$ . Then

$$(y)_\infty = D, \quad \text{and} \quad \sum_{R \in G \cdot S} R \sim \frac{d}{2}D.$$

Therefore, the divisor

$$K_{\mathcal{C}} := \frac{d-4}{2}D$$

is a canonical divisor of  $\hat{\mathcal{C}}$ . Note that  $(x)_\infty = D$  and  $(y)_\infty = D$ . Since  $2 < f < d/2$  and  $f$  divides  $d/2$ , it follows that

$$(f-1) + \left(\frac{d}{f} - 2\right) \leq \frac{d-4}{2}$$

and that

$$x^{f-1}y^{d/f-2} \in \mathcal{L}(K_{\mathcal{C}}).$$

For a point  $Q \in \hat{\mathcal{C}}$  with  $r(Q) = (0 : 0 : 1)$ ,

$$\text{ord}_Q(x^{f-1}y^{d/f-2}) = (f-1)\frac{d}{f} + \left(\frac{d}{f} - 2\right) = d-2.$$

Then the number  $d - 1 = (d - 2) + 1$  is a gap number of pole orders at  $Q$ . On the other hand, for a point  $Q_c := (c : 0 : 1)$  with  $c^f = -1$ ,

$$\left(\frac{y}{x-c}\right)_{\infty} = (d-1)Q_c,$$

namely,  $d - 1$  is a non-gap number at  $Q_c$ . This implies that there does not exist an automorphism  $\sigma$  of  $\hat{\mathcal{C}}$  such that  $\sigma(Q_c) = Q$ . This is a contradiction.

If  $f = d/2$ , then we obtain the curve in (b). We can assume that  $f = 2$ . Considering the differentials of both sides of  $h(x)^2 - \beta^d = x^2(x^2 + 1)g(x^2)^2$ , it follows that

$$2h \times h' = g(x^2)x\{(4x^2 + 2)g(x^2) + 4(x^4 + x^2)g'(x^2)\}.$$

On the other hand, since any multiple component of  $-c(x) = h(x)^2 - \beta^d$  is a component of  $h'$ ,

$$h' = (\deg h)g(x^2) \times x.$$

By these two equations, it follows that

$$dh = (4x^2 + 2)g(x^2) + (4x^4 + 4x^2)g'(x^2).$$

By considering the differentials,

$$dh' = x\{8g(x^2) + (24x^2 + 12)g'(x^2) + (8x^4 + 8x^2)g''(x^2)\}.$$

Since  $h' = (d/2)g(x^2) \times x$ , it follows that

$$\frac{d^2}{2}g(x^2) = 8g(x^2) + (24x^2 + 12)g'(x^2) + (8x^4 + 8x^2)g''(x^2).$$

Let  $g(x) = \sum_{i=0}^{d/4-1} a_i x^i$  with  $a_{d/4-1} = 1$ . We have relations

$$(d^2/2)a_i = 8(i+1)^2 a_i + 4(i+1)(2i+3)a_{i+1}$$

for  $1 \leq i \leq d/4 - 1$ , and

$$(d^2/2)a_0 = 8a_0 + 12a_1.$$

The curve  $\mathcal{C}$  is in Case (c).

*Step 8: The case where  $m = 2$  and  $C_{G_{P_1}}(G_{P_2})$  does not fix any point of  $\mathcal{C} \cap \ell$ .* In this case, there exists a point of  $\mathcal{C} \cap (\mathbb{P}^2 \setminus \ell)$  fixed by  $C_{G_{P_1}}(G_{P_2})$ , which is nonsingular. Since the ramification indices of  $\pi_P$  are two, it follows that  $|C_{G_{P_1}}(G_{P_2})| = 2$ . Furthermore, the number of nonsingular points on  $\ell$  is two, and the number of singular points is  $d/2 + 1 - 2 = d/2 - 1$ . It follows from  $\sum_i e_i = d$  that the multiplicity of all singular points is two. Since  $C_{G_{P_1}}(G_{P_2})$  acts on singular points and does not fix any point of  $\mathcal{C} \cap \ell$ , it follows that  $d/2$  is odd. Note that  $h(x)$  is of the form  $\sum_{j: \text{odd}} b_j x^j$ , since each component of  $h(x)$  represents a line passing through  $d$  ramification points of  $\pi_{P_1}$ . Since any irreducible component of  $-c(x) = h^2 - \beta^d$  with multiplicity two is a component of  $h'$ , it follows that

$$(d/2)^2(h^2 - \beta^d) = (h')^2(x^2 + 1)$$

for a suitable system of coordinates. Considering the differentials of both sides, it follows that

$$(d/2)^2 h = h''(x^2 + 1) + xh'.$$



Using these two equations, we have relations

$$b_{d/2} = 1, \quad (d/2)^2 b_j = j^2 b_j + (j+1)(j+2)b_{j+2}, \quad \text{and} \quad \frac{b_1^2}{(d/2)^2} = -\beta^d.$$

Let  $g(x) = \sum a_i x^i$  be a polynomial such that  $(d/2)g(x^2) = h'(x)$ . Then

$$(d/2)a_i = (2i+1)b_{2i+1}$$

for  $i = 0, \dots, (d-2)/4$ . It follows that

$$a_{(d-2)/4}, \quad \text{and} \quad \left(\frac{d}{2}\right)^2 a_i = (2i+1)^2 a_i + (2i+1)(2i+2)a_{i+1}.$$

Since the defining equation is of the form

$$y^d + h^2 - \beta^d = y^d + \left(\frac{h'}{(d/2)}\right)^2 (x^2 + 1) = y^d + (x^2 + 1)g(x^2)^2 = 0,$$

the curve  $\mathcal{C}$  is in Case (d). □

**Example 2.4.** Let  $d = 6$ . We consider Case (d) in Theorem 1.2. The curve  $\mathcal{C}$  is defined by

$$y^6 + (x^2 + 1)(x^2 + a_0)^2 = 0,$$

where  $(6/2)^2 a_0 = (0+1)^2 a_0 + (0+1)(0+2) \times 1$ . Then it follows that  $a_0 = 1/4$ .

Let  $d = 8$ . We consider Case (c) in Theorem 1.2. The curve  $\mathcal{C}$  is defined by

$$y^8 + x^2(x^2 + 1)(x^2 + a_0)^2 = 0,$$

where  $(8^2/2)a_0 = 8(0+1)^2 a_0 + 4(0+1)(0+3) \times 1$ . Then it follows that  $a_0 = 1/2$ .

**Remark 2.5.** There exist inclusions

$$G_{P_2} \hookrightarrow \text{PGL}(3, K) \quad \text{and} \quad C_{G_{P_1}}(G_{P_2}) \hookrightarrow \text{PGL}(3, K).$$

**Remark 2.6.** It is easily seen that Takahashi's curve is isomorphic to a Generalized Fermat curve  $\mathcal{F}_{d/2, d} : z^{d/2} + y^d + 1 = 0$ . According to a result of Kontogeorgis [9], if  $d \geq 6$  and  $d-1$  is not a power of  $p$ , then  $|\text{Aut}(\hat{\mathcal{C}})| = (d/2) \times (2d) = d^2$ . This implies that  $G_{P_1} G_{P_2} = \text{Aut}(\hat{\mathcal{C}})$ .

For Fermat curves, the number of outer Galois points is known ([2, 7, 18]). For Takahashi's curve, the following holds.

**Proposition 2.7.** Let  $d \geq 4$  be an even integer prime to  $p$  and let  $\mathcal{C}$  be the curve  $X^d + X^{d/2} Z^{d/2} + Y^d = 0$ . Then the number of outer Galois points is at most two.

*Proof.* Let  $P_1 = (1 : 0 : 0)$ ,  $P_2 = (0 : 1 : 0)$  and  $Q = (0 : 0 : 1)$ . The normalization is denoted by  $r : \hat{\mathcal{C}} \rightarrow \mathcal{C}$ . Since  $P_1$  is Galois and  $\mathcal{C} \cap \overline{P_1 Q}$  contains a nonsingular point, it follows from a property of Galois coverings [15, Corollary 3.7.2] that  $r^{-1}(Q)$  consists of exactly  $d/2$  points. Let  $P \neq P_2$  be an outer Galois point and let  $G_P$  the Galois group.

Assume that  $d \geq 6$ . We show that  $P \in \{Z = 0\} = \overline{P_1 P_2}$ . Considering information on Weierstrass points,  $G_P$  acts on the set  $\mathcal{C} \cap \{Z = 0\} = \{Q_1, \dots, Q_d\}$ . (see [9, p.129]). If  $G_P$  fixes all points on  $\mathcal{C} \cap \{Z = 0\}$ , then such points are ramification points for the projection  $\pi_P$ , and hence, the lines  $\ell_i = \overline{P Q_i}$  are tangent lines to  $\mathcal{C}$  for every  $i = 1, \dots, d$ . In this case,  $P$  coincides with the singular point  $Q$ . This is a contradiction. Therefore, there exist  $\sigma \in G_P$  and a point  $R \in \mathcal{C} \cap \{Z = 0\}$  such that  $R \neq \sigma(R) \in \{Z = 0\}$ . Then  $P \in \overline{R \sigma(R)} = \{Z = 0\}$ .

Next, we show that there exists  $\sigma \in G_P \setminus \{1\}$  which acts on the set  $r^{-1}(Q)$ . The line given by  $\overline{QP_2}$  is denoted by  $T_Q$ . Note that  $\mathcal{C} \cap T_Q = \{Q\}$ . Let  $G_Q \cong C_{d/2}$  be the Galois group of the cyclic extension  $K(z, y)/K(y)$  with  $1 + z^{d/2} + y^d = 0$ , which is given by the projection  $\pi_Q$  from  $Q$ . Note that  $G_Q$  acts on  $r^{-1}(Q)$  and, for any  $\sigma \in G_P$ ,  $\sigma^{-1}G_Q\sigma = G_Q$  (see [9, p.130]). Let  $R \in (\mathcal{C} \cap \overline{PQ}) \setminus \{Q\}$ . (Since  $\overline{PQ} \neq T_Q$ , such a point exists.) Then there exist  $d/2$  elements  $\sigma_i \in G_P$  such that  $\{\sigma_i(R)\} = r^{-1}(Q)$ . Assume that there exist  $i, j$  such that  $\sigma_i\sigma_j(R) \in r^{-1}(Q)$ . Then there exists  $\tau \in G_Q$  such that  $\tau(\sigma_i\sigma_j(R)) = \sigma_i(R)$ . It follows that  $\sigma_i^{-1}\tau\sigma_i(\sigma_j(R)) = R \notin r^{-1}(Q)$ . This is a contradiction to  $\sigma_i^{-1}G_Q\sigma_i = G_Q$ . Therefore, for each  $i$ ,  $\sigma_i(r^{-1}(Q)) = r^{-1}(\mathcal{C} \cap \overline{PQ} \setminus \{Q\})$  and  $\sigma_i(r^{-1}(\mathcal{C} \cap \overline{PQ} \setminus \{Q\})) = r^{-1}(Q)$ . Since the number of elements of  $r^{-1}(Q)$  is  $d/2 \geq 3$ , there exists a pair  $(i, j)$  with  $\sigma_i\sigma_j \neq 1$  acting on the set  $r^{-1}(Q)$ .

We show that  $\sigma$  is the restriction of some linear transformation on  $\mathbb{P}^2$ . Let  $\Lambda$  be a (base-point-free) linear system of dimension 2 induced by the normalization  $r : \hat{\mathcal{C}} \rightarrow \mathcal{C} \subset \mathbb{P}^2$ . Let  $\Lambda_0 \subset \Lambda$  be the sublinear system of dimension 1 corresponding to the projection  $\pi_P \circ r$ . Since  $P$  is an outer Galois point,  $\sigma^*\Lambda_0 = \Lambda_0$ . Note that  $r^*T_Q \in \Lambda \setminus \Lambda_0$ . Since  $\mathcal{C} \cap T_Q = \{Q\}$  and  $\sigma$  acts on  $r^{-1}(Q)$ , it follows that  $\sigma^*(r^*(T_Q)) = r^*(T_Q)$  as divisors. This implies that  $\sigma^*\Lambda = \Lambda$ . It follows that  $\sigma$  is the restriction of some linear transformation  $\bar{\sigma}$  on  $\mathbb{P}^2$ .

In this case,  $\bar{\sigma}(T_Q) = T_Q$ . Since  $\bar{\sigma}(\{Z = 0\}) = \{Z = 0\}$ ,  $\bar{\sigma}(P_2) = P_2$ . Therefore,  $\bar{\sigma}$  acts on the ramification points for  $\pi_{P_2}$  and hence,  $\bar{\sigma}(\{Y = 0\}) = \{Y = 0\}$ . It follows that  $P = P_1$ . Assertion follows, in the case where  $d \geq 6$ .

Finally, we consider the case where  $d = 4$ . Then  $g = 1$ . It is not difficult to check that the points  $(\lambda : 0 : 1)$  and  $(-\lambda : 0 : 1)$ , with  $\lambda^2 + 1 = 0$ , are all total inflexion points of  $\mathcal{C}$ . If  $G_P$  is a cyclic group of order four (and  $g = 1$ ), then there exist two totally ramified points for  $\pi_P$ . This implies that  $P = P_2$ . We can assume that  $G_P \cong (\mathbb{Z}/2\mathbb{Z})^{\oplus 2}$ . If  $P \in T_Q$ , then there exist five outer Galois points (including  $P_2$ ) on the line  $T_Q$ , since all elements  $\sigma \in G_{P_2}$  are linear transformations and  $\sigma(P) \neq P$  if  $\sigma \neq 1$ . In this case, five involutions fix a common point in the fiber  $r^{-1}(Q)$ . This is a contradiction to the structure of the automorphism group of elliptic curves (see [14, Theorem 10.1 and Appendix A]). Therefore,  $P \notin T_Q$ . In this case, it follows trivially that there exists  $\sigma \in G_P \setminus \{1\}$  which acts on the set  $r^{-1}(Q)$ . Similar to the fourth paragraph,  $\sigma$  is the restriction of some linear transformation  $\bar{\sigma}$  on  $\mathbb{P}^2$ . In this case,  $\bar{\sigma}(Q) = Q$  and  $\bar{\sigma}(T_Q) = T_Q$ . By the former condition,  $\bar{\sigma}(\{Z = 0\}) = \{Z = 0\}$ . Similar to the previous paragraph, it follows that  $P = P_1$ .  $\square$

**Remark 2.8.** For  $d \geq 6$  in Proposition 2.7, the existence of a non-trivial linear subgroup in  $G_P$  can also be proved by a group-theoretical argument as follows. It can be seen that  $G_P$  has an index 2 subgroup  $\Sigma$  that preserves  $r^{-1}(Q)$  acting on it as a sharply transitive permutation group. The linear subgroup  $V$  of  $G_{P_1} \times G_{P_2}$  consists of all maps  $(x, y) \rightarrow (ux, vy)$  with  $u^{d/2} = 1, v^d = 1$ . In particular  $|V| = \frac{1}{2}d^2$ , and  $V$  preserves  $r^{-1}(Q)$ . The center  $U$  of  $\text{Aut}(\hat{\mathcal{C}})$  consists of all maps  $\{\psi_u : (x, y) \rightarrow (ux, uy) \mid u^{d/2} = 1\}$ .

Assume that  $\Sigma \cap V = \{1\}$ . Then the group  $\Delta = \langle \Sigma, V \rangle$  has order at least  $|\Sigma||V| = \frac{1}{4}d^3$ . As  $\frac{1}{4}d^3 > d^2$ , by Remark 2.6,  $d - 1$  is a power of  $p$ . Furthermore,  $\Delta$  preserves  $r^{-1}(Q)$ .

Let  $\Psi$  be the stabilizer of a point  $R \in r^{-1}(Q)$  in  $\Delta$ . As  $U$  is centralized by  $\Delta$  and  $U$  is transitive on  $r^{-1}(Q)$ , any element in  $\Delta$  which fixes  $R$  must fix  $r^{-1}(Q)$  pointwise. Therefore, if  $\Psi$  is the subgroup of  $\Delta$  fixing  $r^{-1}(Q)$  pointwise, then  $|\Delta| = \frac{1}{2}d|\Psi|$ . This implies  $|\Psi| \geq \frac{1}{2}d^2$ . Since  $2g(\hat{\mathcal{C}}) - 2 = \frac{1}{2}d(d - 4)$ , a contradiction is obtained by applying the Riemann–Hurwitz formula to  $\hat{\mathcal{C}} \rightarrow \hat{\mathcal{C}}/\Psi$ , as  $\frac{1}{2}d(d - 4) \geq -d^2 + \frac{1}{2}d(\frac{1}{2}d^2 - 1)$  is satisfied for  $3 - \sqrt{3} \leq d \leq 3 + \sqrt{3} < 6$ .

**Remark 2.9.** When  $K = \mathbb{C}$  and  $d = 4$ , Proposition 2.7 was first proved by Kanazawa and Yoshihara [8].

For another plane model of Takahashi's curve, we have the following:

**Proposition 2.10.** Let  $d \geq 6$ , and let  $\mathcal{C}' \subset \mathbb{P}^2$  be a plane curve in Case (c) or in Case (d) in Theorem 1.2. If  $d - 1$  is not a power of  $p$ , then the number of Galois points is at most two.

*Proof.* Let  $P'_2 = (0 : 1 : 0)$ . Then  $P'_2$  is an outer Galois point and  $G_{P'_2} \cong C_d$ . Let  $P'_1 \in \{Y = 0\}$  be an outer Galois point described in Proposition 2.3. By the assumption on the degree, it follows from a theorem of Kontogeorgis [9] that  $|\text{Aut}(\hat{\mathcal{C}}')| = d^2$ . Therefore,  $G_{P'_1}G_{P'_2} = \text{Aut}(\hat{\mathcal{C}}')$  and  $G_{P'_1}$  is a normal subgroup of  $\text{Aut}(\hat{\mathcal{C}}')$ . Assume that  $P \in \mathbb{P}^2$  be an outer Galois point different from  $P'_1$ . Since  $G_P \subset \text{Aut}(\hat{\mathcal{C}}')$ , it follows that  $G_{P'_1}G_P = G_{P'_1} \times G_P$ . By Theorem 1.2,  $G_P \cong C_d$  and all automorphisms in  $G_P$  are linear. Since  $\text{Sing}(\mathcal{C}')$  is invariant under the action of  $G_P$ , it follows that  $G_P$  fixes the line  $\{Y = 0\}$ . With singular points and nonsingular points on the line  $\{Y = 0\}$  considered, it follows that there exists an automorphism  $\tau \in G_P$  fixing  $(\mathcal{C}' \setminus \text{Sing}(\mathcal{C}')) \cap \{Y = 0\}$  pointwise. Let  $Q_1, Q_2 \in (\mathcal{C}' \setminus \text{Sing}(\mathcal{C}')) \cap \{Y = 0\}$  be different points, which are ramification points of the projection from  $P$ . Then  $P \in T_{Q_1} \cap T_{Q_2}$ , where  $T_{Q_1}$  and  $T_{Q_2}$  are tangent lines. This implies that  $P = P'_2$ .  $\square$

**Remark 2.11.** Assume that  $p > 0$ ,  $d = p^e m$  for some  $e > 0$  and  $m$  is not divisible by  $p$ . Then we can prove that if there exist two outer Galois points  $P_1$  and  $P_2$  such that  $G_{P_1}G_{P_2} = G_{P_1} \times G_{P_2}$ , then  $m$  divides  $p^e - 1$  and  $\mathcal{C}$  is projectively equivalent to the curve defined by

$$\left( \sum_{m \mid p^i - 1} a_i x^{p^i} \right)^m + c_1 \left( \sum_{m \mid p^j - 1} b_j y^{p^j} \right)^m + c_2 = 0,$$

where  $a_e, a_0, b_e, b_0, c_1, c_2 \in K \setminus \{0\}$  and  $a_i, b_j \in K$  for  $i = 1, \dots, e - 1$  and  $j = 1, \dots, e - 1$ . This will be shown in another paper.

### 3. CONCLUDING REMARKS

It is natural to ask how general the situation studied in our paper is. We start by pointing out the following.

**Remark 3.1.** When  $p = 0$ , the Fermat curve, Takahashi's curve and another plane model of Takahashi's curve are only known examples of plane curves satisfying the following conditions (see the Table in [20]):

- (1)  $g > 0$ .
- (2) There exist two outer Galois points.

Based on Remark 3.1, we raise the following Question.

**Question 3.1.** If  $p = 0$  and  $g > 0$ , is it true that any curve with two Galois points is isomorphic to one of the curves described in Theorem 1.2?

Answering to Question 3.1 seems a rather challenging task, to be addressed in a forthcoming paper. A lot of information should be gathered by a careful analysis for the possible structures of  $G = \langle G_{P_1}, G_{P_2} \rangle$ . We end this Section with some results on this direction.

**Proposition 3.2.** *There exists a non-faithful permutation representation  $\varphi : G \rightarrow S_d$ . Further,  $\varphi(G)$  is a transitive subgroup of  $S_d$ .*

*Proof.* Let  $\Omega = \{\sigma(Q) \mid \sigma \in G_{P_1}\} = \{\tau(Q) \mid \tau \in G_{P_2}\}$  (as pointed out before,  $\Omega$  is a  $G$ -short orbit, whence  $G$  acts on  $\Omega$ ). By [5, Lemma 3.4],  $|\Omega| = d$ . This immediately yields the existence of the permutation representation  $\varphi : G \rightarrow S_d$ . We denote by  $N$  its Kernel and by  $n = |N|$ .

Assume that  $\varphi$  is faithful. This is equivalent to  $n = 1$ . Then  $G$  itself can be regarded as a transitive permutation group on  $\Omega$ . Recall that  $|G| \geq d^2$  as it contains  $G_{P_1}G_{P_2}$ . Since  $p = 0$ , then the stabilizer of any point in  $\Omega$  is cyclic. Then by [10, Theorem], we have  $|G| \leq d^2 - d$ , a contradiction.

As both  $G_{P_1}$  and  $G_{P_2}$  act regularly on  $\Omega$ , we have that  $G$  is transitive on  $\Omega$ , whence our second claim.  $\square$

**Remark 3.3.** *As a consequence of the Riemann–Hurwitz formula applied to  $\mathcal{C} \rightarrow \mathcal{C}/N$ , one immediately gets that  $n \leq d$ . If  $n = d$  then the curve  $\mathcal{C}$  is non-singular as it has genus equal to  $(d-1)(d-2)/2$ , and hence it is projectively equivalent to the Fermat curve (see [18]). This happens because for the Fermat curve with  $P_1 = (1 : 0 : 0)$  and  $P_2 = (0 : 1 : 0)$ , the group  $G$  contains the automorphism represented by*

$$\begin{pmatrix} \zeta & & \\ & 1 & \\ & & 1 \end{pmatrix} \times \begin{pmatrix} 1 & & \\ & \zeta & \\ & & 1 \end{pmatrix} = \begin{pmatrix} \zeta & & \\ & \zeta & \\ & & 1 \end{pmatrix} \sim \begin{pmatrix} 1 & & \\ & 1 & \\ & & \zeta^{-1} \end{pmatrix},$$

where  $\zeta$  is a primitive  $d$ -th root of unity.

**Remark 3.4.** *If  $n = d$ , then the group  $G$  is abelian as we have  $G \simeq C_d \times C_d$ . Interestingly, this is the only case when  $G$  is abelian. By contradiction, let  $n < d$  and assume that  $G$  is abelian. Then  $G/N$  is a transitive abelian permutation group. But a transitive abelian permutation group is necessarily regular, hence  $|G/N| = d$ . But then  $|G| = nd < d^2$ , a contradiction.*

**Acknowledgments.** The authors had an opportunity to discuss this article (other than e-mail) during their visit at The University of Pavia, in September, 2017. The authors thank Professor Gian Pietro Pirola for his warm hospitality.

## REFERENCES

- [1] N. Arakelian, P. Speziali, On generalizations of Fermat curves over finite fields and their automorphisms, *Comm. Algebra*, **45** (2017) no. 11, 4926-4938.
- [2] S. Fukasawa, On the number of Galois points for a plane curve in positive characteristic, *Comm. Algebra*, **36** (2008), 29-36; Part II, *Geom. Dedicata*, **127** (2007), 131-137.
- [3] S. Fukasawa, Classification of plane curves with infinitely many Galois points, *J. Math. Soc. Japan*, **63** (2011), 195-209.
- [4] S. Fukasawa, A birational embedding of an algebraic curve into a projective plane with two Galois points, *J. Algebra*, **511** (2018), 95-101.
- [5] S. Fukasawa, On the number of Galois points for a plane curve in characteristic zero, available at arXiv:1604.01907.
- [6] J. W. P. Hirschfeld, G. Korchmáros and F. Torres, Algebraic curves over a finite field, Princeton Univ. Press, Princeton, 2008.
- [7] M. Homma, Galois points for a Hermitian curve, *Comm. Algebra*, **34** (2006), 4503-4511.
- [8] M. Kanazawa, H. Yoshihara, Galois lines for space elliptic curve with  $j = 12^3$ , *Beitr. Algebra Geom.*, **59** (2018), 431-444.

- [9] A. Kontogeorgis, The group of automorphisms of the function field of the curve  $X^n + Y^m + 1 = 0$ , *J. Number Theory*, **72** (1998), 110-136.
- [10] A. Lucchini, On the order of transitive permutation groups with cyclic point stabilizer, *Atti della Accademia Nazionale dei Lincei. Classe di Scienze Fisiche, Matematiche e Naturali. Rendiconti Lincei. Matematica e Applicazioni, Serie 9*, Vol **9**, (1998), n.4, p. 241-243.
- [11] K. Miura, Galois points for plane curves and Cremona transformations, *J. Algebra*, **320** (2008), 987-995.
- [12] K. Miura, A. Ohbuchi, Automorphism group of plane curve computed by Galois points, *Beitr. Algebra Geom.*, **56** (2015), 695-702.
- [13] K. Miura, H. Yoshihara, Field theory for function fields of plane quartic curves, *J. Algebra*, **226** (2000), 283-294.
- [14] J. H. Silverman, The arithmetic of elliptic curves, Graduate Texts in Mathematics **106**, Springer-Verlag, New York, 1986.
- [15] H. Stichtenoth, Algebraic function fields and codes, Graduate Texts in Mathematics **254**, Springer-Verlag, Berlin Heidelberg, 2009.
- [16] T. Takahashi, Galois point for a plane curve with two singular points, Talk in Symposium on Algebraic Geometry in Sado, Niigata, Japan, June 2011.
- [17] R. C. Valentini, M. L. Madan, A Hauptsatz of L. E. Dickson and Artin-Schreier extensions, *J. Reine Angew. Math.*, **318** (1980), 156-177.
- [18] H. Yoshihara, Function field theory of plane curves by dual curves, *J. Algebra*, **239** (2001), 340-355.
- [19] H. Yoshihara, Rational curve with Galois point and extendable Galois automorphism, *J. Algebra*, **321** (2009), 1463-1472.
- [20] H. Yoshihara, S. Fukasawa, List of problems, available at:  
<https://sites.google.com/sci.kj.yamagata-u.ac.jp/fukasawa-lab/open-questions-english>

DEPARTMENT OF MATHEMATICAL SCIENCES, FACULTY OF SCIENCE, YAMAGATA UNIVERSITY, YAMAGATA 990-8560 (JAPAN)

*Email address:* `s.fukasawa@sci.kj.yamagata-u.ac.jp`

INSTITUTO DE MATEMÁTICA, ESTATSTICA E COMPUTAÇÃO CIENTFICA, UNIVERSIDADE DE CAMPINAS, CAMPINAS, SP 13083-859 (BRAZIL)

*Email address:* `speziali@unicamp.br`