

WILDLY COMPATIBLE SYSTEMS AND SIX OPERATIONS

NING GUO

ABSTRACT. For a scheme X separated and of finite type over an excellent regular scheme S , we define wildly compatible systems of constructible sheaves of modules over finite fields on X for certain vector spaces V . The main result is that for $\dim S \leq 1$, wildly compatible systems are preserved by Grothendieck's six operations and Verdier's duality. Finally, for a smooth integral scheme X over a finite field, we prove that all ℓ -adic compatible systems gives wildly compatible systems.

1. INTRODUCTION

For a scheme X separated and of finite type over an excellent scheme S , we consider a family of locally constant étale sheaves $\{\mathcal{F}_i\}_{i \in I}$ of \mathbb{F}_{λ_i} -modules on X . Each sheaf \mathcal{F}_i corresponds to a representation ρ_i of the étale fundamental group of X . The wild ramification of ρ_i is described by Swan conductors in terms of Brauer traces (see [Ser77, 18.1 and 19.2]). To compare wild ramifications of classes $[\mathcal{F}_i]$ (so-called virtual sheaves) in the Grothendieck group $K_c(X, \mathbb{F}_{\lambda_i})$, it is natural to find a subgroup (defined by Brauer traces) of $\prod_{i \in I} K_c(X, \mathbb{F}_{\lambda_i})$ to specify virtual sheaves with the same ramification. In addition, having “the same wild ramification” should be compatible with functorial operations (for instance, Grothendieck's six operations and Verdier's duality). Wild ramifications of sheaves, in turn, have been related to Euler–Poincaré characteristics and cycles and the development is summarized as the following.

- (i) Deligne–Illusie ([Ill81]) defined the notion of “same wild ramification” for an open subvarieties X of a normal proper variety Z over an algebraic closed field of positive characteristic, and they proved that two locally constant sheaves over X with the same rank and the same wild ramification along $Z \setminus X$ have the same Euler–Poincaré characteristic.
- (ii) When S is an excellent Henselian discrete valuation ring or a field with the residue characteristic exponent $p \geq 1$, Vidal ([Vid04], [Vid05]) constructed a subgroup of $K_c(X, \mathbb{F}_\lambda)$ to define the notion “virtual wild ramification 0” and proved its preservation under Grothendieck's six operations.
- (iii) With Vidal's setting, Yatagawa ([Yat17]) defined the subgroup of “the same wild ramification” and proved that it is preserved by four operations and Verdier's duality. Further, Saito and Yatagawa ([SY16]) proved that two constructible complexes with the same wild ramification have the same characteristic cycle.

In this article, we introduce wildly compatible system of (virtual) constructible sheaves as a more general notion, and prove its preservation under Grothendieck's six operations and Verdier's duality. After generalizing Vidal's definition [Vid04, 2.1] of local wild part of $\pi_1(X)$ to excellent base S without restriction on $\dim S$ (Definition 2.4), we establish a formula (Theorem 2.7) connecting global and local wild parts of π_1 to find their functorial properties (see Proposition 2.12, Corollaries 2.14 and 2.15). With these generalities and properties, wildly compatible systems (Definition 3.7) have no restriction on $\dim S$ and rely on certain vector spaces V , which are flexible: by changing V , we obtain Vidal's and Yatagawa's notion as special examples (Remark 3.11). A wildly compatible system is defined by a subgroup $K_c(X/S, I, V) \subset \prod_{i \in I} K_c(X, \mathbb{F}_{\lambda_i})$, and sheaves in $K_c(X/S, I, V)$ can be viewed as having “compatible wild ramifications”: Brauer traces of these sheaves on each stratum are components of some vectors in V . The main result is the preservation of wildly compatible systems as the following.

Theorem 1.1 (Theorem 4.1). *Let S be an excellent regular scheme with $\dim S \leq 1$. Let $f : X \rightarrow Y$ be an S -morphism of schemes over S separated and of finite type. We have:*

DÉPARTEMENT DE MATHÉMATIQUES, UNIVERSITÉ PARIS-SUD, ORSAY CEDEX, FRANCE

E-mail address: ning.guo1@u-psud.fr.

Date: August 15, 2021.

- (i) $f^* : K_c(Y, I) \rightarrow K_c(X, I)$ induces $f^* : K_c(Y/S, I, V) \rightarrow K_c(X/S, I, V)$;
- (ii) $f_* : K_c(X, I) \rightarrow K_c(Y, I)$ induces $f_* : K_c(X/S, I, V) \rightarrow K_c(Y/S, I, V)$;
- (iii) $f_! : K_c(X, I) \rightarrow K_c(Y, I)$ induces $f_! : K_c(X/S, I, V) \rightarrow K_c(Y/S, I, V)$;
- (iv) $f^! : K_c(X, I) \rightarrow K_c(Y, I)$ induces $f^! : K_c(X/S, I, V) \rightarrow K_c(Y/S, I, V)$;
- (v) $R\mathcal{H}om : K_c(X, I) \times K_c(X, I) \rightarrow K_c(X, I)$ induces
 $R\mathcal{H}om : K_c(X/S, I, V) \times K_c(X/S, I, V) \rightarrow K_c(X/S, I, V)$ when V is a subalgebra;
- (vi) $-\otimes - : K_c(X, I) \times K_c(X, I) \rightarrow K_c(X, I)$ induces
 $-\otimes - : K_c(X/S, I, V) \times K_c(X/S, I, V) \rightarrow K_c(X/S, I, V)$ when V is a subalgebra;
- (vii) $D_X : K_c(X, I) \rightarrow K_c(X, I)$ induces $D_X : K_c(X/S, I, V) \rightarrow K_c(X/S, I, V)$.

This generalizes [Vid05, Cor. 0.2] and [Yat17, Cor. 4.1]. Finally, for a smooth integral scheme X over a finite field, compatible ℓ -adic virtual constructible sheaves are related to our wildly compatible virtual constructible sheaves. More precisely, the decomposition map $d : K(X, E) \rightarrow K(X, F)$ sends compatible λ_i -adic virtual constructible sheaves to wildly compatible virtual constructible \mathbb{F}_{λ_i} -sheaves (Theorem 5.1).

1.2. Conventions. By (X/S) we always mean a scheme X separated and of finite type over an excellent scheme S (and Y/R is similar). We use $\bar{s} \rightarrow S$ to denote geometric points of S , and $X_{\bar{s}}$ is the base change of X over the strict Henselization $S_{\bar{s}}$.

Acknowledgements. I thank to my advisor, Weizhe Zheng. With his forward-looking instruction and guidance, this article was written smoothly. He also helped me to correct mistakes and defects with infinite patience. I thank to Lei Fu for important suggestions for revision.

2. THE WILD PART OF π_1

Nagata's theorem predicts that every scheme X separated and of finite type over an excellent scheme S has a compactification \bar{X} . It turns out that wild ramifications of constructible étale \mathbb{F}_{λ_i} -sheaves \mathcal{F}_i over X is considered over the boundary $\bar{X} \setminus X$. By the theory of modular representation, the elements of $\pi_1(X)$ whose super-natural orders are prime to λ_i form a set $\pi_1(X)_{\text{reg}}$, where we take Brauer traces of \mathcal{F}_i (3.1 or see [Ser77, 18.1]). After a recollection of Nagata compactifications (2.1–2.3), we prove a global-local formula (Theorem 2.7) of wild parts of π_1 (Definition 2.4), which corresponds to $\pi_1(X)_{\text{reg}}$ referred above. By this formula, any wildly compatible system is locally wildly compatible (Proposition 3.12 (ii)).

2.1. Nagata compactifications. For a scheme X over a quasi-compact quasi-separated scheme S , we say $f : X \rightarrow S$ is *compactifiable* or has a *compactification*, if f is factorized by an S -scheme \bar{X} as $f : X \xrightarrow{i} \bar{X} \xrightarrow{\bar{f}} S$, where i is an open immersion and \bar{f} is a proper morphism. We denote this compactification of f by (\bar{X}, i, \bar{f}) . By Nagata's compactification theorem ([Con07, Thm. 4.1]), f is compactifiable if and only if it is of finite type and separated. In this case, all compactifications of f form a category \mathcal{C} . The objects of \mathcal{C} are triplets (\bar{X}, i, \bar{f}) . A morphism between two objects $(Z_1, i_1, f_1), (Z_2, i_2, f_2)$ in \mathcal{C} is such a morphism $g : Z_1 \rightarrow Z_2$ so that $g \circ i_1 = i_2$ and $g \circ f_2 = f_1$. In this case, we say that Z_2 is *dominated* by Z_1 , or Z_1 *dominates* Z_2 as compactifications of $X \rightarrow S$.

In fact, \mathcal{C} is a cofiltered category ([Del73, 3.2.6]):

- 1) given $Z_1, Z_2 \in \text{Obj}(\mathcal{C})$, there exists $Z_3 \in \text{Obj}(\mathcal{C})$ and morphisms $Z_1 \xleftarrow{g_1} Z_3 \xrightarrow{g_2} Z_2$;
- 2) given two morphisms $Z_1 \xrightarrow[r]{s} Z_2$ in \mathcal{C} , there exists $Z_3 \in \text{Obj}(\mathcal{C})$ and a morphism $t : Z_3 \rightarrow Z_1$ in \mathcal{C} such that $s \circ t = r \circ t$.

For 1), it suffices to take $Z' := Z_1 \times_S Z_2$ and a compactification Z_3 of $(i_1, i_2) : X \rightarrow Z'$. For 2), we denote by $k : K \rightarrow Z_1$ the equalizer of r and s . By universal property of equalizer, we factorize i_1 as $i_1 = k \circ i$. Even though k is proper and K is proper over S , the morphism i is not necessarily be an open immersion. Therefore, we take a compactification of $i : X \rightarrow K$ as $X \xrightarrow{i_3} Y \xrightarrow{p} K$. The composition $k \circ p : Y \rightarrow Z_1$ is the desired morphism.

In the sequel, we assume that S is an excellent scheme.

2.2. Nagata normal compactifications. For (X/S) with normal connected X and a Nagata compactification $f : X \rightarrow \overline{X} \xrightarrow{\overline{f}} S$, we consider the normalization \overline{X}^ν of \overline{X} .

Claim 2.2.1. The normalization \overline{X}^ν is also a Nagata compactification of $f : X \rightarrow S$.

Proof. By the universal property of normalization, there is a morphism $i : X \rightarrow \overline{X}^\nu$. We show that $X \xrightarrow{i} \overline{X}^\nu \xrightarrow{\overline{f} \circ \nu} S$ is another compactification of $f : X \rightarrow S$. By excellence assumption of schemes over S , the normalization ν is finite so $\overline{f} \circ \nu : \overline{X}^\nu \rightarrow S$ is proper. On the other hand, the base change $Y := \overline{X}^\nu \times_{\overline{X}} X \rightarrow X$ is a finite birational morphism from an integral scheme to a normal connected scheme, hence is an isomorphism. Therefore, $i : X \rightarrow \overline{X}^\nu$ is an open immersion. \square

All normal compactifications of $X \rightarrow S$ form a full subcategory \mathcal{C}' of \mathcal{C} . Further, \mathcal{C}' is co-cofinal:

- 1) for every $x \in \text{Obj}(\mathcal{C})$, there is an object $y \in \text{Obj}(\mathcal{C}')$ and a morphism $y' \rightarrow x$ in $\text{Morph}(\mathcal{C})$;
- 2) given $x \in \text{Obj}(\mathcal{C})$ and $y_1, y_2 \in \text{Obj}(\mathcal{C}')$ with $y_1 \rightarrow x$ and $y_2 \rightarrow x$, there exists $y_3 \in \text{Obj}(\mathcal{C}')$ fitting into the following commutative diagram

$$\begin{array}{ccccc} y_1 & \longleftarrow & y_3 & \longrightarrow & y_2 \\ & \searrow & \downarrow & \swarrow & \\ & & x & & \end{array} .$$

It is clear that 1) and 2) follows directly from the fact that for any two normal compactifications in \mathcal{C}' , there is always a normal compactification dominating them.

2.3. Nagata normal dense compactifications. On the basis of 2.2, we consider normal compactifications \overline{X} such that $X \rightarrow \overline{X}$ are dominant open immersions. Here we assume that X is a normal connected scheme separated and of finite type over an excellent scheme S . In this case, for an arbitrary normal compactification \overline{X} , we can always take the connected component \overline{X}' of \overline{X} that contains X . Hence X is open dense in \overline{X}' , which is also a normal compactification of $X \rightarrow S$ dominating \overline{X} . Subsequently, all normal compactifications of $X \rightarrow S$ such that X is an open subscheme form a cofiltered co-cofinal subcategory of \mathcal{C}' (and of \mathcal{C}).

By the results about co-cofinality above, for convenience, we always assume that X is an open dense subscheme in every normal compactification of $X \rightarrow S$.

Definition 2.4 (Wild part of π_1). For an open subscheme X of a normal connected scheme \overline{X} , we let x be a geometric point of \overline{X} with the characteristic exponent $p \geq 1$. The strict Henselization \overline{X}_x of \overline{X} at x has a fiber product $X_x := \overline{X}_x \times_{\overline{X}} X$. Let v be a geometric point of X_x lying above a geometric generic point η of X . We define some subsets of $\pi_1(X, \overline{\eta})$ as follows:

- (i) $E_{x,v}$ is the union of images of p -Sylow subgroups of $\pi_1(X_x, v)$ in $\pi_1(X, \eta)$;
- (ii) $E_{X,x}$ is the union of conjugates of $E_{x,v}$ by g in $\pi_1(X, \eta)$;
- (iii) $E_{X,\overline{X}}$ is the closure of the union of $E_{X,x}$ for all geometric points x of \overline{X} .

For (X/S) with normal connected X , we define:

- (iv) $E_{X/S}$ is the intersection of $E_{X,\overline{E}}$ for all normal compactifications \overline{X} of $X \rightarrow S$;
- (v) for the image τ of η in X , the subset $E_{\tau/S}$ is the intersection of images of $E_{U/S}$ in $E_{X/S}$ for all open dense subscheme of X .

Claim 2.4.1. Definition 2.4 (ii) is independent of choices of v .

Proof. For another geometric point w above η , there is an isomorphism of fibre functors $\gamma : F_v \simeq F_w$ inducing an isomorphism

$$\phi : \pi_1(X_x, v) = \text{Aut}(F_v) \xrightarrow{\sim} \text{Aut}(F_w) = \pi_1(X_x, w) \quad h \longmapsto \gamma \circ h \circ \gamma^{-1}.$$

Therefore, the image g of γ in $\pi_1(X, \eta)$ induces an inner automorphism such that the following diagram

$$\begin{array}{ccc} \pi_1(X_x, v) & \longrightarrow & \pi_1(X, \eta) \\ \downarrow \phi & & \downarrow (-)^g \\ \pi_1(X_x, w) & \longrightarrow & \pi_1(X, \eta) \end{array}$$

commutes, where $(-)^g$ denotes the inner automorphism induced by g . \square

Remark 2.5. Here we compare our Definition 2.4 with Vidal's notion ([Vid04, §§2.1], [Vid05, §1]).

- Let S be the spectrum of an excellent Henselian discrete valuation ring of mixed characteristic $(0, p)$ (that is, the characteristic of residue field at generic point is 0 and for closed point is p). Let \bar{x} be the geometric point above η , then $E_{X, \bar{x}} = \{1\}$. On the other hand, $E_{X, x}^{\text{Vidal}}$ is not trivial in general. Therefore, $E_{X, \bar{x}} \neq E_{X, x}^{\text{Vidal}}$ in this case.
- Further, we consider $E_{X, \bar{X}}$ and $E_{X, \bar{X}}^{\text{Vidal}}$ for a normal compactification \bar{X} of $X \rightarrow S$. A finite quotient Q of $\pi_1(X)$ corresponds to a Galois cover $Y \rightarrow X$. Let \bar{Y} be the normalization of \bar{X} in Y . From [Vid05, Proof 6.1] we know that $g \in Q$ belongs to $E_{X, \bar{X}}$ if and only if g fixes a point of $(\bar{Y})_s$, where s is the closed point of S . Similarly, $g \in Q$ belongs to $E_{X, \bar{X}}^{\text{Vidal}}$ if and only if g fixes a point of \bar{Y} . But the locus of fixed points \bar{Y}^g in \bar{Y} of g is proper over S , since proper morphism is stable under composition. Consequently, we find that $\bar{Y}^g \cap \bar{Y}_s \neq \emptyset$ and $E_{X, \bar{X}} = E_{X, \bar{X}}^{\text{Vidal}}$.

Remark 2.6. With Definition 2.4 we can also define tameness for Galois covers, which is a classical notion. Let X be a normal connected scheme separated and of finite type over $\text{Spec}(\mathbb{Z})$. A Galois cover $Y \rightarrow X$ is called *tame* if $E_{X/S_{\text{pec}(\mathbb{Z})}}(Q) = \{1\}$, where Q is the finite quotient of $\pi_1(X)$ associated to the Galois cover $Y \rightarrow X$. When Y is regular, this is the notion of numerical tameness in [KS10].

A global-local formula. Given a geometric point \bar{s} of S , the strict Henselization $S_{\bar{s}} \rightarrow S$ at \bar{s} induces a base change of X and \bar{X} . Though the base change $X_{\bar{s}}$ is not connected in general, we can also define its normal compactification as the disjoint union of normal compactifications of its all connected components. All normal compactifications of $X_{\bar{s}}$ also form a cofiltered category. In this sense, the base change $(\bar{X})_{\bar{s}}$ is a normal compactification of $X_{\bar{s}}$ over $S_{\bar{s}}$. Conversely, given a normal compactification $Z_{\bar{s}}$ of $X_{\bar{s}} \rightarrow S_{\bar{s}}$, we want to prove that it can be dominated by the base change of a normal compactification of $X \rightarrow S$ (Proposition 2.8), which leads to a formula relating global and local wild parts of π_1 as the following.

Theorem 2.7. For (X/S) with $\dim S < 3$ and normal connected X , we have

$$E_{X/S} = \overline{\bigcup_{\bar{s} \rightarrow S} \bigcup_{g \in \pi_1(X, \eta)} g^{-1} E'_{X_{\bar{s}}/S_{\bar{s}}} g},$$

where $E'_{X_{\bar{s}}/S_{\bar{s}}}$ are images of $E_{X_{\bar{s}}/S_{\bar{s}}}$ in $\pi_1(X, \eta)$.

Proof. We may assume that S is connected. By Definition 2.4,

$$E_{X, \bar{X}} = \overline{\bigcup_x \bigcup_{g \in \pi_1(X, \eta)} g^{-1} E_{x, v} g}, \quad \text{where } x \text{ goes through geometric points of } \bar{X}.$$

For a geometric point x of \bar{X} , its composition with $\bar{X} \rightarrow S$ is a geometric point s of S . So x gives rise to a geometric point y of $(\bar{X})_{\bar{s}}$. We have the following commutative diagram

$$\begin{array}{ccc} \pi_1((X_{\bar{s}})_y, w) & \longrightarrow & \pi_1(X_{\bar{s}}, \xi) \\ \downarrow & & \downarrow \\ \pi_1(X_x, v) & \longrightarrow & \pi_1(X, \eta), \end{array}$$

where ξ is a geometric generic point above η and w is a geometric point above ξ . The first column is an isomorphism, since $(X_{\bar{s}})_y \simeq X_x$. It follows that the image $E'_{y,w}$ of $E_{y,w}$ in $\pi_1(X, \eta)$ is $E_{x,v}$, and

$$\begin{aligned} E_{X, \bar{X}} &= \overline{\bigcup_{\bar{s}} \bigcup_y \bigcup_{g \in \pi_1(X, \eta)} g^{-1} E_{y,w} g} \\ &= \overline{\bigcup_{\bar{s}} \bigcup_{g \in \pi_1(X, \eta)} g^{-1} E'_{X_{\bar{s}}, (\bar{X})_{\bar{s}}} g}, \end{aligned}$$

where \bar{s} goes through geometric points of S and y goes through geometric points of $(\bar{X})_{\bar{s}}$. For a fixed $\bar{s} \rightarrow S$, by Proposition 2.8 below, the base changes of \bar{X} are co-cofinal in the category of normal compactifications of $X_{\bar{s}} \rightarrow S_{\bar{s}}$. By taking intersections on both sides, we find

$$\begin{aligned} E_{X/S} &= \bigcap_{X \rightarrow \bar{X} \rightarrow S} E_{X, \bar{X}} \stackrel{(1)}{=} \bigcap_{X \rightarrow \bar{X} \rightarrow S} \left(\overline{\bigcup_{\bar{s} \rightarrow S} \bigcup_{g \in \pi_1(X, \eta)} g^{-1} E'_{X_{\bar{s}}, (\bar{X})_{\bar{s}}} g} \right) \\ &\stackrel{(2)}{=} \varprojlim_Q \bigcap_{X \rightarrow \bar{X} \rightarrow S} \left(\left(\bigcup_{\bar{s} \rightarrow S} \bigcup_{g \in \pi_1(X, \eta)} g^{-1} E'_{X_{\bar{s}}, (\bar{X})_{\bar{s}}} g \right) (Q) \right) \\ &\stackrel{(3)}{=} \varprojlim_Q \left(\bigcup_{\bar{s} \rightarrow S} \bigcup_{g \in \pi_1(X, \eta)} g_Q^{-1} \bigcap_{\bar{X}} E'_{X_{\bar{s}}, (\bar{X})_{\bar{s}}} (Q) g_Q \right) \\ &\stackrel{(4)}{=} \varprojlim_Q \left(\left(\bigcup_{\bar{s} \rightarrow S} \bigcup_{g \in \pi_1(X, \eta)} g^{-1} E'_{X_{\bar{s}}/S_{\bar{s}}} g \right) (Q) \right) \\ &\stackrel{(5)}{=} \overline{\bigcup_{\bar{s} \rightarrow S} \bigcup_{g \in \pi_1(X, \eta)} g^{-1} E'_{X_{\bar{s}}/S_{\bar{s}}} g}. \end{aligned}$$

Here Q goes through all finite quotients of $\pi_1(X, \eta)$ and g_Q denotes the image of g in Q . The equalities (2) and (5) come from that a closed subset in a profinite group is the inverse limit of all its finite quotients. The (3) is by the distribution law for intersection and union and the fact that the category of normal compactifications is cofiltered. \square

Proposition 2.8. *We let X and S be as assumed in Theorem 2.7. For any normal compactification $Z_{\bar{s}}$ of $X_{\bar{s}} \rightarrow S_{\bar{s}}$, there exists a normal compactification \bar{X} of $X \rightarrow S$ such that $\bar{X} \times_S S_{\bar{s}}$ dominates $Z_{\bar{s}}$.*

Proof. The base change of any normal compactification of $X \rightarrow S$ to $S_{\bar{s}}$ is a normal compactification of $X_{\bar{s}} \rightarrow S_{\bar{s}}$. To find the desired \bar{X} , we prove in the following steps.

- (i) We have $S_{\bar{s}} = \varprojlim_{i \in I} S_i$ for each affine scheme S_i étale over S . Because $X_{\bar{s}}$ and $Z_{\bar{s}}$ are of finite type over $S_{\bar{s}}$ and $S_{\bar{s}}$ is excellent, a fortiori Noetherian, by [SP, 01TX], $X_{\bar{s}}$ and $Z_{\bar{s}}$ are of finite presentation over $S_{\bar{s}}$. By [EGA IV₃, 8.8.2], there exists $i_0 \in I$ such that the normal compactification $X_{\bar{s}} \rightarrow Z_{\bar{s}}$ descend over $S' := S_{i_0}$ as $X_{S'} \rightarrow Z_{S'}$.
- (ii) Since $S' \rightarrow S$ is étale and affine, it is quasi-finite and separated. By Zariski's main theorem, it is an open immersion followed by a finite morphism as $S' \rightarrow T \rightarrow S$. We may assume that T is connected and replace T by its normalization (as in the proof of Claim 2.4.1) if necessary.
- (iii) We glue X_T and $Z_{S'}$ along the open $X_{S'}$ to obtain $Z := Z_{S'} \cup_{X_{S'}} X_T$. It is clear that $Z \rightarrow T$ is of finite type and we prove its separatedness. We restrict the diagonal morphism $\Delta_{Z/T}: Z \rightarrow Z \times_T Z$ to four open subschemes $Z \times_T Z = (Z_{S'} \times_T Z_{S'}) \cup (Z_{S'} \times_T X_T) \cup (X_T \times_T Z_{S'}) \cup (X_T \times_T X_T)$ to check $\Delta_{Z/T}$ is a closed immersion. Since $Z_{S'}/T$ and X_T/T are separated, by symmetry, it suffices to consider $\Delta_{Z/T}|_{Z_{S'} \times_T X_T}$, which is the closed immersion $\Delta_{X_{S'}/S'}$ by the isomorphism $Z_{S'} \times_T X_T \simeq Z_{S'} \times_{S'} X_{S'}$ and the separatedness of $X_{S'}/S'$. Therefore, by Nagata's theorem, there is a compactification Z_T of $Z \rightarrow T$. Further, we have $(Z_T)_{S'} \simeq Z_{S'}$.
- (iv) Since $T \rightarrow S$ is finite, it follows that $X_T \rightarrow Z_T \rightarrow S$ is also a compactification. Because T is normal and $\dim T < 3$, it is Cohen-Macaulay. Subsequently, the dominant morphism $T \rightarrow S$ is flat and so is $X_T \rightarrow X$. By Lemma 2.9 below, there is a compactification Z'_T of $X_T \rightarrow S$ and a normal compactification \bar{X} of $X \rightarrow S$ with a finite morphism $Z'_T \rightarrow \bar{X}$. The base change of $Z'_T \rightarrow \bar{X}$ to $S_{\bar{s}}$ is a finite morphism between two normal compactifications of $X_{\bar{s}} \rightarrow S_{\bar{s}}$, so it is an isomorphism and $(\bar{X})_{\bar{s}}$ dominates $Z_{\bar{s}}$. \square

Lemma 2.9. *For a finite flat morphism $f : X \rightarrow Y$ between two connected schemes separated and of finite type over S and a compactification \overline{X} of $X \rightarrow S$, there is a compactification \overline{X}' of $X \rightarrow S$ dominating \overline{X} and a compactification \overline{Y}' of $Y \rightarrow S$ fitting into a commutative diagram*

$$\begin{array}{ccc} X & \longrightarrow & \overline{X}' \\ f \downarrow & & \downarrow \overline{f}, \\ Y & \longrightarrow & \overline{Y}' \end{array}, \quad \text{where } \overline{f} \text{ is finite flat.}$$

Proof. This proof comes from O. Gabber's idea and is similar to I. Vidal's proof in [Vid04, Prop. 2.1.1 (ii)].

By Nagata's theorem, there exists \overline{Y} as a compactification of Y . The closure of the graph of f in $\overline{X} \times_S \overline{Y}$ is denoted by Γ . By the separatedness of $\overline{Y} \rightarrow S$, the first projection $p_1 : \Gamma \rightarrow \overline{X}$ is an isomorphism $i : \Gamma_X \simeq X$ when restricted to X . Because f is dominant, the second projection $p_2 : \Gamma \rightarrow \overline{Y}$ when restricted to Y coincides with f , up to a composition with the isomorphism i . Therefore, the morphism p_2 is finite and flat and satisfies the condition in the flattening theorem ([RG71, Thm. 5.2.2]) for $n = 0$: it is flat outside $\overline{Y} \setminus Y$. Then there exists a Y -admissible (that is, the blowing-up locus is in $\overline{Y} \setminus Y$) blowing-up $\overline{Y}' \rightarrow \overline{Y}$ such that the strict transform $\overline{f} : \overline{X}' \rightarrow \overline{Y}'$ is flat. The flatness of \overline{f} implies it is not only finite on $Y \subset \overline{Y}'$ but also on the entire \overline{Y}' . \square

Remark 2.10. If we consider normal compactifications which are algebraic spaces, the assumption in Theorem 2.7 for S can be more general: S can be an arbitrary excellent scheme. The idea is from Chenyang Xu, who suggests to consider equivariant normal compactifications. As in the proof of Theorem 2.7 (ii) let $K(S')$ and $K(S)$ be fraction fields of S' and S respectively. Then we take a Galois extension $K/K(S)$ with Galois group G containing $K(S')$ and the normalization R of S in K . The normalization of S in $K(S')$ is denoted by T . With the similar technique in (iii) before, one can extend $\overline{X}_{S'}$ to a normal compactification \overline{X}_T of $X_T \rightarrow T$ (X_T can be replaced with its normalization if necessary). Then the base change of \overline{X}_T to R can be dominated by a G -equivariant normal compactification of $X_R \rightarrow R$ by the construction in [Zhe09, 3.7]. The normalization of the quotient $\overline{X}_R//G$ is a normal compactification and its base change along $S' \rightarrow S$ dominates $\overline{X}_{S'}$.

2.11. Functoriality of wild parts of π_1 . The wild parts of π_1 have pleasant functorial properties. The following lemma generalizes Vidal's results ([Vid04, Prop. 2.1.1]).

Proposition 2.12. *In this proposition, we fix*

- S and R are excellent scheme;
- $X \rightarrow \overline{X}$ and $Y \rightarrow \overline{Y}$ be two open immersions of normal connected schemes;
- $f : X \rightarrow Y$ is a morphism of schemes;
- $\phi : \pi_1(X, \eta) \rightarrow \pi_1(Y, \xi)$ is the morphism of étale fundamental groups induced by f , where η is a geometric generic point of X and ξ is the image of η in Y .

(i)

$$\begin{array}{ccc} \overline{X} & \longrightarrow & \overline{Y} \\ \uparrow & & \uparrow \\ X & \xrightarrow{f} & Y \end{array} \quad \text{then } \phi(E_{X, \overline{X}}) \subset E_{Y, \overline{Y}}.$$

If there is a commutative square

(ii) Let $X \rightarrow \overline{X} \rightarrow S$ and $Y \rightarrow \overline{Y} \rightarrow R$ be two compactifications.

$$\begin{array}{ccc} X & \xrightarrow{f} & Y \\ \downarrow & & \downarrow \\ S & \longrightarrow & R \end{array} \quad \text{then } \phi(E_{X/S}) \subset E_{Y/R}.$$

If there is a commutative square

(iii) When f is a finite étale, we have $E_{X, \overline{X}} = E_{Y, \overline{Y}} \cap \pi_1(X, \eta)$. If further (X/S) and (Y/S) , then

$$E_{X/S} = E_{Y/S} \cap \pi_1(X, \eta).$$

(iv) With the condition of (ii), if we further assume that $X = Y$ with $f = \text{id}$, then

$$E_{X/S} \subset E_{X/R}.$$

When $S \rightarrow R$ is proper, we have

$$E_{X/S} = E_{X/R}.$$

Proof.

(i) By definition,

$$E_{X, \overline{X}} = \overline{\bigcup_{x \rightarrow \overline{X}} \bigcup_{g \in \pi_1(X, \eta)} g^{-1} E_{x, v} g},$$

where x and v are geometric points of \overline{X} and $X_x = X \times_{\overline{X}} \overline{X}_x$ respectively, and v is above η . It suffices to show that

$$\phi \left(\bigcup_{x \rightarrow \overline{X}} \bigcup_{g \in \pi_1(X, \eta)} g^{-1} E_{x, v} g \right) \subset E_{Y, \overline{Y}},$$

since $E_{Y, \overline{Y}}$ is closed. Any element $\sigma \in \bigcup_{x \rightarrow \overline{X}} \bigcup_{g \in \pi_1(X, \eta)} g^{-1} E_{x, v} g$ can be written as $\gamma_1^{-1} \phi_1 \gamma_1$, where $\phi_1 \in \pi_1(X_x, v)$ and $\gamma_1 \in \pi_1(X, \eta)$. We let y and w be the images of x and v respectively, and consider the following diagram

$$\begin{array}{ccccc} v & \rightarrow & X_x & \longrightarrow & \overline{X}_x \\ & & \downarrow & \searrow & \downarrow \\ & & & & Y_y & \longrightarrow & \overline{Y}_y \\ & & \downarrow & & \downarrow & & \downarrow \\ \eta & \rightarrow & X & \longrightarrow & \overline{X} \\ & & \downarrow & \searrow & \downarrow \\ & & & & Y & \longrightarrow & \overline{Y} \end{array}$$

and a similar diagram for fundamental groups. These diagrams are not necessarily commutative. Let ϕ_2 be the image of ϕ_1 in $\pi_1(Y_y, w)$ and let γ_2 be the image of γ_1 in $\pi_1(Y, \xi)$. Then we have $f(\sigma) = \gamma_2^{-1} \phi_2 \gamma_2 \in E_{Y, \overline{Y}}$ and $f(E_{X, \overline{X}}) \subset E_{Y, \overline{Y}}$.

- (ii) For any normal compactification \overline{Y} of $Y \rightarrow R$, we take a normal compactification of $\overline{Y} \times_R S$ to obtain a normal compactification of $X \rightarrow S$. By the result of (i) the assertion follows.
- (iii) We first prove for the case when \overline{X} is dominated by the normalization of a normal compactification \overline{Y} for $Y \rightarrow S$. Then X_x is a pointed connected component of $Y_y \times_Y X$. To prove that $E_{x, v} = E_{y, w} \cap \pi_1(X, \eta)$, it suffices to show that the following diagram

$$\begin{array}{ccc} \pi_1(X_x, v) & \xrightarrow{\psi_X} & \pi_1(X, \eta) \\ \downarrow & & \downarrow \\ \pi_1(Y_y, w) & \xrightarrow{\psi_Y} & \pi_1(Y, \xi) \end{array}$$

is cartesian. All vertical arrows are injective. Let $U = \psi_Y^{-1}(\pi_1(X, \eta)) \subset \pi_1(Y_y, w)$. This is an open subgroup associated to an intermediate pointed finite étale covering $X_x \rightarrow V \rightarrow Y_y$. The correspondent point is denoted by $t \rightarrow V$. As the map $\pi_1(V, t) \rightarrow \pi_1(Y, \xi)$ is factorized by $\pi_1(X, \eta)$, then by [SGA 1_{new}, Exp. V, 6.4], the natural projection $V \times_Y X \rightarrow V$ admits a pointed section s . Composite the section s with the natural pointed morphism $V \times_Y X \rightarrow Y_y \times_Y X$, we find that the pointed morphism $V \rightarrow Y_y \times_Y X$ must factors through X_x . That means $V \simeq X_x$, hence that diagram is cartesian.

Let $q \in \pi_1(Y, \xi)$ and let (X', η') be a pointed connected finite étale cover of Y corresponding to the open subgroup $\pi_1(X', \eta') = q\pi_1(X, \eta)q^{-1}$. We denote by $\phi : \pi_1(X, \eta) \rightarrow \pi_1(X', \eta')$ the isomorphism induced by conjugation by q in $\pi_1(Y, \xi)$ and $\mu : \pi_1(X', \eta') \rightarrow \pi_1(X, \eta)$ is denoted for its inverse. Then $q^{-1} E_{y, w} q \cap \pi_1(X, \eta) = \mu(E_{y, w} \cap \pi_1(X', \eta')) = \mu(E_{x, v'})$ (for X' it is the same with the above argument). By the result of (i) we find that $\psi(E_{x, v'}) \subset \psi(E_{X', \overline{X'}}) \subset E_{X, \overline{X}}$. Thus

$q^{-1}E_{y,wq} \cap \pi_1(X, \eta) \subset E_{X, \overline{X}}$. As this can be applied to all q and all geometric point $x \rightarrow \overline{X}$, we then use the result of (i) to find that $E_{Y, \overline{Y}} \cap \pi_1(X, \eta) \subset E_{X, \overline{X}}$. By Lemma 2.9 above, we finish the proof.

- (iv) Every normal compactification of $X \rightarrow S$ is also a normal compactification of $X \rightarrow R$. Further, given any normal compactification W of $X \rightarrow R$, there is a normal compactification Z of $X \rightarrow R$ dominating W . Hence Z is also a normal compactification of $X \rightarrow S$ and dominates W . Therefore $E_{X/S} = E_{X/R}$. \square

Definition 2.13.

- (i) Let \mathcal{B} be the (pointed) category whose objects consist of all open immersions $(X \xrightarrow{i} \overline{X})$ of normal connected schemes with fixed geometric generic points. Morphisms in \mathcal{B} consists of all commutative squares as in Proposition 2.12 (i) such that induced morphisms of geometric generic points of $Y, \overline{Y}, X, \overline{X}$ are compatible, where $(X \hookrightarrow \overline{X})$ and $(Y \hookrightarrow \overline{Y})$ are objects in \mathcal{B} .
- (ii) Let \mathcal{D} be the (pointed) category whose objects consist of all (X/S) , where X are normal connected with fixed geometric generic points. Morphisms in \mathcal{D} consists of all commutative squares as in Proposition 2.12 (ii) such that induced morphisms of geometric generic points of X, S, Y, R are compatible, where (X/S) and (Y/R) are objects in \mathcal{D} .

With Definition 2.13 and Proposition 2.12 (i)& (ii) we obtain the following corollary.

Corollary 2.14. *Here we use the same notations as Definition 2.13.*

- (i) *The assignment*

$$b: \mathcal{B} \rightarrow \mathbf{Set} \quad (X \rightarrow \overline{X}) \mapsto E_{X, \overline{X}}$$

is a functor. For each morphism ϕ in \mathcal{B} as in Definition 2.13 (i), b sends it to $b(\phi): E_{X, \overline{X}} \rightarrow E_{Y, \overline{Y}}$ defined by f .

- (ii) *There is a functor $E_{-/-}: \mathcal{D} \rightarrow \mathbf{Set}$ defined as follows. For each $(X/S) \in \text{Obj}(\mathcal{D})$, $E_{-/-}$ sends it to $E_{X/S}$. For each morphism ψ in \mathcal{D} as in Definition 2.13 (ii), $E_{-/-}$ sends it to $E_{-/-}(\psi): E_{X/S} \rightarrow E_{Y/R}$ defined by f .*
- (iii) *Let (Y/R) be an object in \mathcal{D} . Let $f: X \rightarrow Y$ be a morphism separated and of finite type. Then f gives a morphism in \mathcal{D} and hence we have $E_{X/R} \subset E_{Y/R}$.*

Corollary 2.15. *Let S be an excellent scheme. Let \mathcal{D}_S be the category of all normal connected schemes separated and of finite type over S . Let X be a normal connected scheme. Let \mathcal{E}^X be the category consisting of all excellent schemes S with $X \rightarrow S$ separated and of finite type.*

- (i) *There is a covariant functor $E_{-/S}: \mathcal{D}_S \rightarrow \mathbf{Set}$ defined as follows. For each object $X \in \mathcal{D}_S$, $E_{-/S}$ sends it to $E_{-/S}(X) = E_{X/S}$ an object in \mathbf{Set} . For each morphism $f: X \rightarrow Y$ in \mathcal{D}_S , $E_{-/S}$ sends it to $E_{-/S}(f): E_{X/S} \xrightarrow{f_*} E_{Y/S}$ a morphism in \mathbf{Set} .*
- (ii) *There is a covariant functor $E_{X/-}: \mathcal{E}^X \rightarrow \mathbf{Set}$ defined as follows. For each object $S \in \mathcal{E}^X$, $E_{X/-}$ sends it to $E_{X/-}(S) = E_{X/S}$. For each morphism $h: S \rightarrow R$ in \mathcal{E}^X , $E_{X/-}$ sends it to $E_{X/-}(h): E_{X/S} \hookrightarrow E_{X/R}$ a inclusion morphism in \mathbf{Set} .*

Proof. By Corollary 2.14 (ii) the assertion of (i)&(ii) follows. \square

3. WILDLY COMPATIBLE SYSTEMS OF VIRTUAL CONSTRUCTIBLE SHEAVES

Now we consider a family of virtual constructible (resp., locally constant) sheaves $\{\mathcal{F}_i\}_{i \in I}$ on X in (X/S) . Here \mathcal{F}_i are elements in Grothendieck group $K_c(X, \mathbb{F}_{\lambda_i})$ (resp., $K_{\text{coh}}(X, \mathbb{F}_{\lambda_i})$) of constructible (resp., locally constant) sheaves of \mathbb{F}_{λ_i} -modules on X , and each λ_i is a power of prime ℓ_i . In order to compare the wild ramifications of $\{\mathcal{F}_i\}_{i \in I}$, we use Brauer traces.

3.1. Brauer traces. Let G be a profinite group. For a finite field \mathbb{F}_λ , where λ is a power of a prime ℓ , the subset consisting of ℓ -regular elements (i.e., elements of orders prime to ℓ) is denoted by $G_{\ell\text{-reg}}$. For $M \in K.(\mathbb{F}_\lambda[G])$ an element of Grothendieck group of finite dimensional \mathbb{F}_λ -vector spaces with continuous G -actions, the Brauer trace is a central function $\text{Tr}_M^{\text{Br}} : G_{\ell\text{-reg}} \rightarrow W(\mathbb{F}_\lambda)$, where $W(\mathbb{F}_\lambda)$ is the Witt ring of \mathbb{F}_λ . Concretely, each eigenvalue ζ of action of $g \in G_{\ell\text{-reg}}$ on M is in $\overline{\mathbb{F}_\lambda}$ and has a unique lift $[\zeta]$ as a root of unity of order prime to ℓ , and Brauer trace is given by $\text{Tr}_M^{\text{Br}}(g) = \sum[\zeta]$. Brauer trace is an additive function and is multiplicative with respect to the element of $K.(\mathbb{F}_\lambda[G])$.

Note that the Brauer trace, when restricted to $E_{X/S}$, takes value in a CM field $K_{\lambda_i} := \mathbb{Q}_{\lambda_i} \cap \mathbb{Q}(\zeta_{p^\infty})$ considered as a subfield of \mathbb{Q}_{λ_i} , where \mathbb{Q}_{λ_i} are quotient fields of $W(\mathbb{F}_{\lambda_i})$ (i.e., unramified extensions of \mathbb{Q}_{ℓ_i} with residue fields \mathbb{F}_{λ_i}). Let V be a \mathbb{Q} -vector subspace of $\prod_{i \in I} K_{\lambda_i}$. In the sequel we use this notation of K_{λ_i} for convenience and λ_i are all assumed to be invertible on S .

3.2. Wildly compatible locally constant sheaves.

Definition 3.3. Let X be a normal connected in (X/S) . Let V be a fixed subspace of $\prod_{i \in I} K_{\lambda_i}$. We say a system $\{\mathcal{F}_i\}_{i \in I}$ of virtual locally constant \mathbb{F}_{λ_i} -sheaves on X is a *wildly compatible system* for V if $(\text{Tr}_{\mathcal{F}_i}^{\text{Br}}(g))_{i \in I} \in V$ for all $g \in E_{X/S}$. We define a subgroup $K_{\text{coh}}(X/S, I, V)$ of $K_{\text{coh}}(X, I) := \prod_{i \in I} K_{\text{coh}}(X, \mathbb{F}_{\lambda_i})$ to be the subgroup consisting of $(\mathcal{F}_i)_{i \in I}$ in wildly compatible systems.

Remark 3.4 ([Vid04, Rem. 2.2.1]). Let S and X be the same as in Definition 3.3. Assume that $\{\mathcal{F}_i\}_{i \in I}$ are finitely many locally constant virtual sheaves over \mathbb{F}_{λ_i} . Let V be a subspace of $\prod_{i \in I} K_{\lambda_i}$. Then $\{\mathcal{F}_i\}$ form a wildly compatible system for V if and only if there exists a normal compactification \overline{X}_0 of $X \rightarrow S$ such that $\{\mathcal{F}_i\}$ satisfy the following condition: for every $g \in E_{X/S, \overline{X}_0}$, we have $(\text{Tr}_{\mathcal{F}_i}^{\text{Br}}(g))_{i \in I} \in V$. One side is easy to prove by $E_{X/S} \subset E_{X, \overline{X}_0}$. For the other side, we notice that $\pi_1(X)$ acts on \mathcal{F}_i by a finite quotient Q . Since Q is finite and the category \mathcal{C} of normal compactifications of $X \rightarrow S$ is cofiltered, for the projection $p_Q : \pi_1(X, \overline{\eta}) \rightarrow Q$, the finite quotient parts $E_{X/S}(Q) := p_Q(E_{X/S})$ and $E_{X, \overline{X}}(Q) := p_Q(E_{X, \overline{X}})$ satisfy $E_{X/S}(Q) = \bigcap_{\overline{X} \in \mathcal{C}} E_{X, \overline{X}}(Q) = E_{X, \overline{X}_0}(Q)$ for a normal compactification \overline{X}_0 of $X \rightarrow S$ then the assertion follows.

3.5. Wildly compatible constructible sheaves. Given a virtual constructible sheaf \mathcal{F}_ℓ on X , there is a stratification $X = \coprod_{i \in I} U_i$ denoted by P_ℓ , where U_i are normal connected and locally closed subschemes of X such that restrictions of \mathcal{F}_ℓ on U_i are locally constant. Here by ‘‘stratification’’ we mean a ‘‘nice stratification’’, i.e., for each stratum U_i , its closure is a disjoint union of strata $\overline{U}_i = \coprod_{j \in J} U_j$ where $J \subset I$. But X can have another stratification Q_ℓ for \mathcal{F}_ℓ : $X = \coprod_{j \in J} V_j$. We say a stratification P_ℓ is *finer* than Q_ℓ or $P_\ell \geq Q_\ell$, if $V_j = \coprod_{i(j)} U_{i(j)}$ for each j , here $i(j)$ are functions of index sets I and J . Now we consider finitely many virtual constructible sheaves \mathcal{F}_i on X with stratifications P_i . Then there is a common refinement P so that we can compare their wild ramifications in terms of Brauer traces.

Assumption 3.6. In the sequel when considering virtual constructible sheaves, we further assume that $V = \bigcap_J p_J^{-1} p_J(V)$, where J goes through all finite subsets of I and p_J are projections to subspaces.

Definition 3.7. Let V be a fixed \mathbb{Q} -vector subspace of $\prod_{i \in I} K_{\lambda_i}$. We say a system $\{\mathcal{F}_i\}_{i \in I}$ of virtual constructible \mathbb{F}_{λ_i} -sheaves on X in (X/S) is a *wildly compatible system* for V , if for any finite subset $J \subset I$, there is a common stratification $X = \coprod_{\sigma \in \Sigma_J} X_\sigma$ such that $\{\mathcal{F}_j|_{X_\sigma}\}_{j \in J}$ is a wildly compatible system of virtual locally constant sheaves on X_σ for $p_J(V)$ and for each $\sigma \in \Sigma_J$. We define a subgroup $K_c(X/S, I, V)$ of $K_c(X, I) := \prod_{i \in I} K_c(X, \mathbb{F}_{\lambda_i})$ to be the subgroup consisting of such $(\mathcal{F}_i)_{i \in I}$.

Remark 3.8. From Definition 3.7, it is clear that a system $\{\mathcal{F}_i\}_{i \in I}$ of virtual constructible sheaves is a wildly compatible system for V if and only if its restriction to J is a wildly compatible system for $p_J(V)$ for every finite subset $J \subset I$.

Proposition 3.9. For (X/S) with $\dim S \leq 1$, let $\{\mathcal{F}_i\}_{i \in I}$ be finitely many virtual locally constant sheaves on X . We fix a \mathbb{Q} -vector subspace V of $\prod_{i \in I} K_{\lambda_i}$. If there is a common stratification $X = \coprod_{\sigma \in \Sigma} X_\sigma$ for $\{\mathcal{F}_i\}_{i \in I}$ such that for each σ we have $(\text{Tr}_{\mathcal{F}_i|_{X_\sigma}}^{\text{Br}}(g))_{i \in I} \in V$ for all $g \in E_{X_\sigma/S}$, then $(\text{Tr}_{\mathcal{F}_i}^{\text{Br}}(g))_{i \in I} \in V$ for all $g \in E_{X/S}$.

Proof. This proposition and its proof is similar to Vidal’s in [Vid05, 6.2 (ii)]. By the valuative criterion of O. Gabber ([Vid05, 6.1]), $E_{X/S} = \overline{\bigcup_\sigma \text{Im}(E_{X_\sigma/S})}$. Hence for every $g \in E_{X/S}$, we have $(\text{Tr}_{\mathcal{F}_i}^{\text{Br}}(g))_{i \in I} \in V$. \square

Corollary 3.10. *Assumption of X being as in Proposition 3.9, for any index set I and any V , we have*

$$K_{\text{coh}}(X/S, I, V) = K_c(X/S, I, V) \cap K_{\text{coh}}(X, I).$$

Remark 3.11. Here we give three special examples of wildly compatible systems of virtual locally constant sheaves and constructible sheaves.

- (i) When $\#I = 1$ and $V = 0$, a virtual constructible sheaf \mathcal{F} in $K_c(X/S, I, V)$ can be viewed as it “has zero wild ramification”.
- (ii) When $\#I = 2$ and let V be the kernel of the map

$$K_{\lambda_1} \times K_{\lambda_2} \rightarrow \mathbb{Q}, \quad a \mapsto \frac{1}{[K_{\lambda_1} : \mathbb{Q}]} \text{Tr}_{K_{\lambda_1}/\mathbb{Q}}(a) - \frac{1}{[K_{\lambda_2} : \mathbb{Q}]} \text{Tr}_{K_{\lambda_2}/\mathbb{Q}}(a)$$

For $(\mathcal{F}_1, \mathcal{F}_2) \in K_{\text{coh}}(X/S, I, V)$ they “have the same wild ramification” in a weaker sense.

- (iii) Let L be a field of characteristic zero with embeddings $\iota = (\iota_i)_{i \in I} : L \rightarrow \prod_{i \in I} \overline{\mathbb{Q}_{\lambda_i}}$ and let V be $(\prod_{i \in I} K_{\lambda_i}) \cap \text{Image}((\iota_i)_{i \in I})$. If $(\text{Tr}_{\mathcal{F}_i}^{\text{Br}}(g))_{i \in I} \in V$ for all $g \in E_{X/S}$, then $(\mathcal{F}_i)_{i \in I} \in K_{\text{coh}}(X/S, I, V)$. In this case, for any fixed index set I , $(\mathcal{F}_i)_{i \in I} \in K_{\text{coh}}(X/S, I, V)$ if and only if for all subset $J \subset I$ with $\#J \leq 2$, $(\mathcal{F}_j)_{j \in J}$ is a wildly compatible system for $p_J(V)$.
 - (a) For (i), when S is the spectrum of an excellent Henselian discrete valuation ring with residue characteristic exponent $p \geq 1$ or an algebraically closed field k with characteristic p , and X is a normal connected scheme separated and of finite type over S , by Remark 2.5 we obtain $K_c(X/S, I, V) = K_c(X, \mathbb{F}_{\lambda}^0)$ which is Vidal’s notion.
 - (b) For (ii), when S is the spectrum of an excellent Henselian discrete valuation ring of residue characteristic exponent $p \geq 1$ and X is a scheme separated and of finite type over S , we obtain $K_{\text{coh}}(X/S, I, V) = \Delta_{\text{coh}}(X, \mathbb{F}_{\lambda_1}, \mathbb{F}_{\lambda_2})$ which is Yatagawa’s notion. Further with Assumption 3.6 for V , we have $K_c(X/S, I, V) = \Delta_c(X, \mathbb{F}_{\lambda_1}, \mathbb{F}_{\lambda_2})$. In fact, our generalization is the same with Kato’s ([Kat18, Def. 2.2]).
 - (c) Let I, V and S, X be the same with (iii) with extra Assumption 3.6 for V . If for any $J \subset I$ a finite subset, there is a stratification $X = \coprod_{\sigma \in \Sigma_J} X_{\sigma}$ such that for each $g \in E_{X_{\sigma}/S}$ there exists $a \in L$ and $\text{Tr}_{\mathcal{F}_i|X_{\sigma}}^{\text{Br}}(g) = \iota_i(a)$, then $(\mathcal{F}_i)_{i \in I} \in K_c(X/S, I, V)$. In this case, for any fixed index set I , we have $(\mathcal{F}_i)_{i \in I} \in K_c(X/S, I, V)$ if and only if for all subset $J \subset I$ with $\#J \leq 2$, $(\mathcal{F}_j)_{j \in J}$ is a wildly compatible system for V .

Proposition 3.12. *For (X/S) , let $T \xrightarrow{h} S \rightarrow R$ be morphisms of excellent schemes with (S/R) .*

- (i) *If we also denote $X_T \rightarrow X$ by h , then*

$$h^* K_c(X/S, I, V) \subset K_c(X_T/T, I, V).$$

- (ii) *Suppose S is regular with $\dim S < 3$. If $S_{\bar{s}} \rightarrow S$ is denoted by s , then we have*

$$(\mathcal{F}_i)_{i \in I} \in K_c(X/S, I, V) \quad \text{if and only if} \quad (s^* \mathcal{F}_i)_{i \in I} \in K_c(X_{\bar{s}}/S_{\bar{s}}, I, V) \quad \text{for all } \bar{s}.$$

- (iii) *We have $K_c(X/R, I, V) \subset K_c(X/S, I, V)$.*

Proof.

- (i) It suffices to prove for the case when \mathcal{F}_i are locally constant. The assertion follows directly from Corollary 2.12,(ii).
- (ii) For $(\mathcal{F}_i)_{i \in I} \in K_c(X/S, I, V)$, by (i), we conclude that $\{s^*(\mathcal{F}_i)\}_{i \in I}$ is a wildly compatible system.

For the other side, we reduce to the case when $\#I$ is finite and \mathcal{F}_i are locally constant. There exists a finite quotient $E_{X/S}(Q)$ such that for all $g \in E_{X/S}$, $\text{Tr}_{\mathcal{F}_i}^{\text{Br}}(g) = \text{Tr}_{\mathcal{F}_i}^{\text{Br}}(\bar{g})$ for all $i \in I$, where \bar{g} is the image of g in $E_{X/S}(Q)$. By Theorem 2.7, $E_{X/S}(Q) = \bigcup_{\bar{s} \rightarrow S} \bigcup_{h \in \pi_1(X, a)} \bar{h} E'_{X(\bar{s})/S(\bar{s})}(Q) \bar{h}^{-1}$. Thus there is a geometric point $\bar{s}_{\bar{g}} \rightarrow S$ such that $\bar{g} = \bar{h} g' h^{-1}$, where $g' \in E_{X(\bar{s}_{\bar{g}})/S(\bar{s}_{\bar{g}})}(Q)$ and \bar{g}' is the image of g' under $\pi_1(X(\bar{s}), b) \rightarrow \pi_1(X, a) \rightarrow Q$. By assumption, $\{s^*(\mathcal{F}_i)\}_{i \in I}$ are wildly compatible systems for all geometric points $\bar{s} \rightarrow S$. Finally $(\text{Tr}_{\mathcal{F}_i}^{\text{Br}}(g))_{i \in I} = (\text{Tr}_{\mathcal{F}_i}^{\text{Br}}(\bar{g}))_{i \in I} =$

$(\mathrm{Tr}_{\mathcal{F}_i}^{\mathrm{Br}}(\overline{hg'h^{-1}}))_{i \in I} = (\mathrm{Tr}_{\mathcal{F}_i}^{\mathrm{Br}}(\overline{g'}))_{i \in I} = (\mathrm{Tr}_{s^* \mathcal{F}_i}^{\mathrm{Br}}(g'))_{i \in I} \in V$. So $\{\mathcal{F}_i\}_{i \in I}$ is also a wildly compatible system.

(iii) The proof is directly from the result of Corollary 2.15, (ii). \square

4. PRESERVATION BY SIX OPERATIONS

Theorem 4.1. *For (X/S) and (Y/S) with $\dim \leq 1$ and a morphism $f: X \rightarrow Y$, we have:*

- (i) $f^*: K_c(Y, I) \rightarrow K_c(X, I)$ induces $f^*: K_c(Y/S, I, V) \rightarrow K_c(X/S, I, V)$;
- (ii) $f_*: K_c(X, I) \rightarrow K_c(Y, I)$ induces $f_*: K_c(X/S, I, V) \rightarrow K_c(Y/S, I, V)$;
- (iii) $f_!: K_c(X, I) \rightarrow K_c(Y, I)$ induces $f_!: K_c(X/S, I, V) \rightarrow K_c(Y/S, I, V)$;
- (iv) $f^!: K_c(X, I) \rightarrow K_c(Y, I)$ induces $f^!: K_c(X/S, I, V) \rightarrow K_c(Y/S, I, V)$;
- (v) $R\mathcal{H}om: K_c(X, I) \times K_c(X, I) \rightarrow K_c(X, I)$ induces
 $R\mathcal{H}om: K_c(X/S, I, V) \times K_c(X/S, I, V) \rightarrow K_c(X/S, I, V)$ when V is a subalgebra;
- (vi) $-\otimes -: K_c(X, I) \times K_c(X, I) \rightarrow K_c(X, I)$ induces
 $-\otimes -: K_c(X/S, I, V) \times K_c(X/S, I, V) \rightarrow K_c(X/S, I, V)$ when V is a subalgebra;
- (vii) $D_X: K_c(X, I) \rightarrow K_c(X, I)$ induces $D_X: K_c(X/S, I, V) \rightarrow K_c(X/S, I, V)$.

This theorem also implies results for the three cases in Remark 3.11, including Vidal's ([Vid05, Cor. 0.2]) and Yatagawa's ([Yat17, Cor. 4.1]). In 3.11, note that V are subalgebras for (a) and (c) but not (b).

Proof. By Proposition 3.12, it suffices to prove for the case when S is strictly local. Further $f^* = f^!$ by [Zhe15, Cor. 9.5], $f_* = f_!$ by [IZ13, Thm. 3.14] generalizing a theorem of Laumon ([Lau81]). So (i)–(iv) reduces to f^* and $f_!$. By Remark 3.8, we may assume that I is a finite set.

- (i)& (iv) This proof is similar to [Vid04, Prop. 2.3.3] and [Yat17, Cor. 4.1]. For $(\mathcal{F}_i)_{i \in I} \in K_c(Y, I)$, we may assume that they are constructible sheaves. There is a common stratification for $\mathcal{F}_i: Y = \coprod_{\sigma} Y_{\sigma}$ such that $(\mathcal{F}_i)_{i \in I} \in K_{\mathrm{coh}}(Y_{\sigma}/S, I, V)$. Take preimages of Y_{σ} and take their stratifications we have $X = \coprod_{\sigma, \mu} X_{\sigma\mu}$ such that $f^* \mathcal{F}_i|_{X_{\sigma\mu}}$ are locally constant sheaves. By Proposition 2.12 the assertion follows.
- (ii)&(iii) Let $(a_i)_{i \in I} \in K_c(X/S, I, V)$. First, we prove the assertion when f is an immersion. If f is an open immersion, then $Y = X \coprod (Y \setminus X)$ and $f_! a_i|_X = a_i$, $f_! a_i|_{Y \setminus X} = 0$. If f is a closed immersion, then we have $f_* a_i|_X = a_i$, $f_* a_i|_{Y \setminus X} = 0$. By taking a stratification of $Y \setminus X$, we check on each stratum and find that $\{a_i\}_{i \in I}$ for a wildly compatible system.

For general cases, we decompose $X = X_{\eta} \coprod X_s$, $Y = Y_{\eta} \coprod Y_s$ as closed fibre parts and generic fibre parts. Write $f_{\eta}: X_{\eta} \rightarrow Y_{\eta}$ and $f_s: X_s \rightarrow Y_s$. Let $i_{Y_{\eta}}: Y_{\eta} \rightarrow Y$ and $i_{Y_s}: Y_s \rightarrow Y$ be immersions. By $f_! a_i = i_{Y_{\eta}}^! f_{\eta}^!(a_i|_{X_{\eta}}) + i_{Y_s}^! f_s^!(a_i|_{X_s})$ for $i \in I$ and the proof above for immersions, it suffices to prove for f_{η} or f_s . The assertion is local so it suffices to prove when $Y = \mathrm{Spec} A$ is affine. Further, we take $X = \mathrm{Spec} B$ affine and therefore $Y = \mathrm{Spec} B[x_1, \dots, x_r]/I$. The morphism $X \rightarrow Y$ is decomposed as a closed immersion followed by a series of morphisms of relative dimension one.

When f is a closed immersion the proof has been finished above. Now let f be a morphism of relative dimension one. Since a_i are constructible, we may assume that Y is normal connected and $(f_! a_i)_{i \in I} \in K_{\mathrm{coh}}(Y, I)$. Further we may assume that X is normal connected and $(a_i)_{i \in I} \in K_{\mathrm{coh}}(X/S, I, V)$. For any pair of components (a_1, a_2) of an element $(a_i)_{i \in I}$ in $K_{\mathrm{coh}}(X/S, I, V)$ its associated galois étale covering $p: V \rightarrow X$ trivializing them with galois group H . By shrinking Y to an open dense subscheme if necessary, we may assume that images of this pair of components under $(f \circ p)_!$ are in Grothendieck groups of locally constant sheaves. Let Q be a finite quotient of $\pi_1(Y)$ acting on $f_! a_1, f_! a_2, (f \circ p)_! \mathbb{F}_{\lambda_1}$ and $(f \circ p)_! \mathbb{F}_{\lambda_2}$.

By shrinking Y to an open dense subscheme, we may assume that $E_{Y/S}(Q) = (E_{\tau/S}(Y))(Q)$, where τ is the generic point of Y . Let $g \in E_{Y/S}$, then by [Vid05, Cor. 3.0.5 and Thm. 0.1], we express the Brauer traces of a_{ℓ_i} for $i = 1, 2$ as the following

$$\mathrm{Tr}_{f_i a_i}^{\mathrm{Br}}(g) = \frac{1}{|H|} \sum_{h' \in H', p_{P'}(h')=g'} \mathrm{Tr}_{R\Gamma_c(V_{\bar{\tau}}, E_{\lambda_i})}(h') \times \mathrm{Tr}_{a_i}^{\mathrm{Br}}(p_H(h')),$$

where P' is a pro- p -subgroup of $\mathrm{Gal}(\bar{\tau}/\tau)$ with image P under $\mathrm{Gal}(\bar{\tau}/\tau) \rightarrow \pi_1(Y)$ and $H' = P' \times H$, E_{λ_i} is a finite extension of \mathbb{Q}_{ℓ_i} such that its integer ring has the residue field \mathbb{F}_{λ_i} . For all $g \in E_{X/S}$ we have $(\mathrm{Tr}_{a_i}^{\mathrm{Br}}(g))_{i \in I} \in V$. We need to show that for all $g' \in E_{Y/S}$ we have $(\mathrm{Tr}_{f_i a_i}^{\mathrm{Br}}(g'))_{i \in I} \in V$. By [Vid05, Prop. 2.2.1] and [Vid05, Prop. 2.3.1], the traces $\mathrm{Tr}_{R\Gamma_c(V_{\bar{\tau}}, E_{\lambda_i})}(h')$ are integers independent of i . Then by [Vid05, Cor. 3.0.7 and Thm. 0.1], if $h' \in H'$ such that $\mathrm{Tr}_{R\Gamma_c(V_{\bar{\tau}}, E_{\lambda_i})}(h') \neq 0$ then $p_H(h')$ is in the image of $E_{X/S}$ by $\pi_1(X) \rightarrow H$. Thus $(\mathrm{Tr}_{f_i a_i}^{\mathrm{Br}}(g))_{i \in I} \in V$ for every $g \in E_{Y/S}$. So we finish the proof.

- (v) Since $R\mathcal{H}om_X(\mathcal{F}_i, \mathcal{G}_i) \simeq D_X(\mathcal{F}_i \otimes D_X(\mathcal{G}_i))$, the assertion follows from (vi) and (vii).
- (vi) The assertion follows since Brauer trace of a tensor product is a product of Brauer traces.
- (vii) Let $\{\mathcal{F}_i\}_{i \in I} \in K_c(X, I, V)$. Let $j : U \rightarrow X$ be an open immersion such that \mathcal{F}_i are lisse and $U \rightarrow S$ is smooth. Let $i : Y = X \setminus U \rightarrow X$ be the closed immersion. Then $D_X \mathcal{F}_i = D_X j_! j^* \mathcal{F}_i + D_X i_* i^* \mathcal{F}_i$. It suffices to prove that $D_X j_! j^* \mathcal{F}_i$ form a wildly compatible system. By definition we have $D_X j_! j^* \mathcal{F}_i = j_* D_U j^* \mathcal{F}_i = j_* R\mathcal{H}om_U(j^* \mathcal{F}_i, \mathbb{F}_{\lambda_i}(d)[2d])$. By the result of (ii), it suffices to consider $R\mathcal{H}om_U(j^* \mathcal{F}_i, \mathbb{F}_{\lambda_i}(-d)[-2d])$. In fact, for a dual module M^\vee of M , the Brauer trace satisfies $\mathrm{Tr}_{M^\vee}^{\mathrm{Br}}(g) = \mathrm{Tr}_M^{\mathrm{Br}}(g^{-1})$. Note that as a result of definition, $E_{X/S}$ is symmetric: if $g \in E_{X/S}$, then $g^{-1} \in E_{X/S}$. By (i), $j^* \mathcal{F}_i$ form a wildly compatible system. So $(j^* \mathcal{F}_i)^\vee(-d)[-2d]$ also form a wildly compatible system. Thus $D_U j_! j^* \mathcal{F}_i$ form a compatible system and by Noether's induction we finish the proof. \square

5. RELATION WITH COMPATIBLE ℓ -ADIC VIRTUAL CONSTRUCTIBLE SHEAVES

Let X be a smooth curve over \mathbb{F}_{p^f} . Let ℓ_i be primes distinct with p , where $i \in I$. Let E_i/\mathbb{Q}_{ℓ_i} be finite extensions of fields with residue fields \mathbb{F}_{λ_i} and λ_i are integer powers of ℓ_i . We fix a field L of characteristic zero and embeddings $\{\iota_i : L \hookrightarrow E_i \hookrightarrow \overline{\mathbb{Q}_{\ell_i}}\}$. Let K be the function field of X . We recall the definition of (strictly) compatible system of E_i -virtual sheaves on X ([Fuj02, 1.2]). A family of virtual constructible sheaves $\{\mathcal{F}_i\}_{i \in I}$ where \mathcal{F}_i are virtual sheaves over E_i , is called an L -compatible system if for all $x \in |X|$, all geometric points $\bar{x} \rightarrow x$ and all $g \in \mathrm{Gal}(\kappa(\bar{x})/\kappa(x))$ such that g are integer powers of geometric Frobenius as F_0^f , there exist $a_g \in L$ depending on g such that $\mathrm{Tr}(g, (\mathcal{F}_i)_{\bar{x}}) = \iota_i(a_g)$. Now we find a connection between E_i -compatible systems and wildly compatible systems.

Let \mathcal{O} be a complete discrete valuation ring and let its fraction field be E of characteristic zero and its residue field be F of characteristic $\ell > 0$. Let X be a Noetherian and connected scheme. Then the category of lisse ℓ -adic sheaves on X is equivalent to the category of continuous ℓ -adic representations of $\pi_1(X, \bar{x})$, where \bar{x} is a geometric point of X . Let $K(X, \mathcal{O}), K(X, E), K(X, F)$ be the Grothendieck groups of constructible sheaves of \mathcal{O}, E, F -modules over X respectively. Let $j^* : K(X, \mathcal{O}) \rightarrow K(X, E)$ be the ring homomorphism induced by the exact functor

$$\mathrm{Mod}_c(X, \mathcal{O}) \rightarrow \mathrm{Mod}_c(X, E), \quad M \mapsto E \otimes_{\mathcal{O}} M$$

Let $i^* : K(X, \mathcal{O}) \rightarrow K(X, F)$ be the ring homomorphism given by the triangulated functor

$$D_c^b(X, \mathcal{O}) \rightarrow D_c^b(X, F), \quad M \mapsto F \otimes_{\mathcal{O}}^{\mathbb{L}} M$$

By [Zhe15, Prop. 9.4], j^* is an isomorphism. Thus we obtain the decomposition morphism $d := i^* \circ j^*{}^{-1} : K(X, E) \rightarrow K(X, F)$.

Theorem 5.1. *For any L -compatible system $\{\mathcal{F}_i\}_{i \in I}$ of virtual constructible sheaves on a smooth integral scheme X over \mathbb{F}_{p^f} , the image $\{d(\mathcal{F}_i)\}_{i \in I}$ is a wildly compatible system of virtual constructible \mathbb{F}_{λ_i} -sheaves.*

Proof. It suffices to prove for $\#I = 2$ and locally constant sheaves $\mathcal{F}_1, \mathcal{F}_2$ on X . If they are L -compatible, then by definition, for all closed point $x \in |X|$, all $\bar{x} \rightarrow x$ and all $g \in \mathrm{Gal}(\kappa(\bar{x})/\kappa(x))$ an integer power

of F_0^f , there are $a_g \in L$ such that $\mathrm{Tr}(g, (\mathcal{F}_i)_{\bar{x}}) = \iota_i(a_g)$. Note that $(\mathcal{F}_i)_{\bar{\eta}}$ give rise to representations $\rho_i : \pi_1(X, \bar{\eta}) \rightarrow \mathrm{GL}_{n_i}(E_i)$, where $\bar{\eta}$ is the geometric generic point of X . By Chebotarev density theorem ([Ser65, Thm. 7]), the sets consisting of the images of conjugacy classes of $\{\mathrm{Frob}_x^n\}_{n \in \mathbb{Z}}$ under ρ_i for all unramified $x \in X$ are dense subsets in $\rho_i(\pi_1(X, \bar{\eta}))$ respectively. Therefore for all $g \in E_{X/\mathrm{Spec}(\mathbb{F}_{p^f})} \subset \pi_1(X, \bar{\eta})$, we have $\mathrm{Tr}_{d(\mathcal{F}_1)}^{\mathrm{Br}}(g) = \iota_1(a_g)$ and $\mathrm{Tr}_{d(\mathcal{F}_2)}^{\mathrm{Br}}(g) = \iota_2(a_g)$ for $a_g \in L$. This shows that $\{d(\mathcal{F}_i)\}_{i \in I}$ is a wildly compatible system in the sense of Remark 3.11, (c). \square

Remark 5.2. In [Zhe09] Weizhe Zheng has defined L -compatible systems over excellent Henselian discrete valuation rings. Further in [LZ18] Qing Lu and Weizhe Zheng have proved that Theorem 5.1 can be generalized to this case.

REFERENCES

- [EGA IV₃] A. Grothendieck and J. Dieudonné, *Éléments de géométrie algébrique. IV. Étude locale des schémas et des morphismes de schémas. III*, Inst. Hautes Études Sci. Publ. Math. **28** (1966), 255. MR0217086 (36 #178)
- [SGA 1_{new}] *Revêtements étales et groupe fondamental (SGA 1)*, Documents Mathématiques (Paris) [Mathematical Documents (Paris)], 3, Société Mathématique de France, Paris, 2003 (French). Séminaire de géométrie algébrique du Bois Marie 1960–61. [Algebraic Geometry Seminar of Bois Marie 1960–61]; Directed by A. Grothendieck; With two papers by M. Raynaud; Updated and annotated reprint of the 1971 original [Lecture Notes in Math., 224, Springer, Berlin; MR0354651 (50 #7129)]. MR2017446 (2004g:14017)
- [Con07] Brian Conrad, *Deligne’s notes on Nagata compactifications*, Journal-Ramanujan Mathematical Society **22** (2007), no. 3, 205.
- [Del73] Pierre Deligne, *Cohomologie à supports propres*, Théorie des topos et cohomologie étale des schémas, 1973, pp. 250–480.
- [Fuj02] Kazuhiro Fujiwara, *Independence of ℓ for intersection cohomology (after Gabber)*. *Algebraic geometry 2000, Azumino (Hotaka)*, Adv. Stud. Pure Math **36** (2002), 145–151.
- [Ill81] Luc Illusie, *Théorie de Brauer et caractéristique d’Euler-Poincaré (d’après P. Deligne)*, The Euler-Poincaré characteristic (French), Astérisque **82** (1981), 161–172.
- [IZ13] Luc Illusie and Weizhe Zheng, *Odds and ends on finite group actions and traces*, International Mathematics Research Notices **2013** (2013), no. 1, 1–62.
- [Kat18] Hiroki Kato, *Wild ramification and restrictions to curves*, Internat. J. Math. **29** (2018), no. 8, 1850052, 20, DOI 10.1142/S0129167X18500520.
- [KS10] Moritz Kerz and Alexander Schmidt, *On different notions of tameness in arithmetic geometry*, Mathematische Annalen **346** (2010), no. 3, 641.
- [Lau81] Grard Laumon, *Comparaison de caractéristiques d’Euler-Poincaré en cohomologie ℓ -adique*, C. R. Acad. Sci. Paris Sér. I Math **292** (1981), no. 3, 209–212.
- [LZ18] Qing Lu and Weizhe Zheng, *Compatible systems and ramification*, 2018. arXiv preprint: 1801.03221.
- [RG71] Michel Raynaud and Laurent Gruson, *Critères de platitude et de projectivité Techniques de “platification” d’un module*, Inventiones mathematicae **13** (1971), 1–89.
- [SY16] Takehashi Saito and Yuri Yatagawa, *Wild ramification determines the characteristic cycle*, arXiv preprint: 1604.01513 (2016).
- [Ser65] Jean-Pierre Serre, *Zeta and L -functions*, Arithmetic Algebraic Geometry (Proc. Conf. Purdue Univ., 1963), 1965, pp. 82–92.
- [Ser77] ———, *Linear representations of finite groups*, Springer-Verlag, New York-Heidelberg, 1977. Translated from the second French edition by Leonard L. Scott, Graduate Texts in Mathematics, Vol. 42.
- [SP] The Stacks Project Authors, *Stacks Project*, 2018.
- [Vid04] Isabelle Vidal, *Théorie de Brauer et conducteur de Swan*, Journal of Algebraic Geometry **13** (2004), no. 2, 349–391.
- [Vid05] ———, *Courbes nodales et ramification sauvage virtuelle*, Manuscripta Mathematica **118** (2005), no. 1, 43–70.
- [Yat17] Yuri Yatagawa, *Having the same wild ramification is preserved by the direct image*, Manuscripta Mathematica (2017), 1–14.
- [Zhe15] Weizhe Zheng, *Six operations and Lefschetz-Verdier formula for Deligne-Mumford stacks*, Science China Mathematics **58** (2015), no. 3, 565–632. doi: 10.1007/s11425-015-4970-z.
- [Zhe09] ———, *Sur l’indépendance de ℓ en cohomologie ℓ -adique sur les corps locaux*, Ann. Sci. Éc. Norm. Supér. (4) **42** (2009), no. 2, 291–334.