

GENUS ZERO GOPAKUMAR-VAFA TYPE INVARIANTS FOR CALABI-YAU 4-FOLDS II: FANO 3-FOLDS

YALONG CAO

ABSTRACT. In analogy with the Gopakumar-Vafa (GV) conjecture on Calabi-Yau (CY) 3-folds, Klemm and Pandharipande defined GV type invariants on Calabi-Yau 4-folds using Gromov-Witten theory and conjectured their integrality. In a joint work with Maulik and Toda, the author conjectured their genus zero invariants are DT_4 invariants of one dimensional stable sheaves. In this paper, we study this conjecture on the total space of canonical bundle of a Fano 3-fold Y , which reduces to a relation between twisted GW and DT_3 invariants on Y . Examples are computed for both compact and non-compact Fano 3-folds to support our conjecture.

MSC 2010: 14N35, 14J32, 14J45

0. INTRODUCTION

0.1. GV/ DT_4 conjecture on CY 4-folds. Gromov-Witten invariants are rational numbers which virtually count stable maps from complex curves to algebraic varieties (or symplectic manifolds). Because of multiple-cover contributions, they are in general not integers and hence are not honest enumerative invariants. On Calabi-Yau 3-folds, Gopakumar-Vafa [8] conjectured the existence of integral invariants which determine Gromov-Witten invariants. In [13], Klemm-Pandharipande proposed a parallel conjecture on Calabi-Yau 4-folds.

More specifically, let X be a smooth projective Calabi-Yau 4-fold, fix $\beta \in H_2(X, \mathbb{Z})$ and $n \geq 0$. For integral classes $\gamma_i \in H^{m_i}(X, \mathbb{Z})$, $1 \leq i \leq n$, one defines genus zero GW invariants

$$GW_{0,\beta}(X)(\gamma_1, \dots, \gamma_n) := \int_{[\overline{M}_{0,n}(X,\beta)]^{\text{vir}}} \prod_{i=1}^n \text{ev}_i^*(\gamma_i),$$

where $\text{ev}_i: \overline{M}_{0,n}(X, \beta) \rightarrow X$ is the i -th evaluation map.

The *Gopakumar-Vafa type invariants*

$$n_{0,\beta}(X)(\gamma_1, \dots, \gamma_n) \in \mathbb{Q}$$

are defined in terms of the identity

$$\sum_{\beta > 0} GW_{0,\beta}(X)(\gamma_1, \dots, \gamma_n) q^\beta = \sum_{\beta > 0} n_{0,\beta}(X)(\gamma_1, \dots, \gamma_n) \sum_{d=1}^{\infty} d^{n-3} q^{d\beta},$$

and conjectured to be integers [13].

In [5], Cao-Maulik-Toda gave a sheaf-theoretic interpretation of the above GV type invariants in terms of Donaldson-Thomas invariants on CY 4-folds (called DT_4 invariants) introduced by Cao-Leung [3] and Borisov-Joyce [1]. More specifically, we consider the moduli scheme $M_{X,\beta}$ of one dimensional stable sheaves on X with Chern character $(0, 0, 0, \beta, 1)$. By the results of [3, 1], and the recent orientability result [2], there exists a DT_4 virtual class

$$[M_{X,\beta}]^{\text{vir}} \in H_2(M_{X,\beta}, \mathbb{Z}),$$

which depends on the choice of an orientation, i.e. on each connected component of $M_{X,\beta}$, there are two choices of orientations, which affect the corresponding contribution to the virtual class by a sign (for each connected component).

Since the virtual dimension is not zero, we require insertions to define invariants. Let

$$\tau: H^m(X) \rightarrow H^{m-2}(M_{X,\beta}), \quad \tau(\gamma) := \pi_{M*}(\pi_X^* \gamma \cup \text{ch}_3(\mathcal{E})),$$

where π_X, π_M are projections from $X \times M_{X,\beta}$ to corresponding factors, and $\text{ch}_3(\mathcal{E})$ is the Poincaré dual of the fundamental cycle of the universal sheaf \mathcal{E} .

For $\gamma_i \in H^{m_i}(X, \mathbb{Z})$ with $1 \leq i \leq n$, the DT_4 invariant is defined by

$$DT_4(X)(\beta \mid \gamma_1, \dots, \gamma_n) := \int_{[M_{X,\beta}]^{\text{vir}}} \prod_{i=1}^n \tau(\gamma_i).$$

Conjecture 0.1. ([5, Conjecture 1.3]) *For a suitable choice of orientation, we have the identity*

$$n_{0,\beta}(X)(\gamma_1, \dots, \gamma_n) = \text{DT}_4(X)(\beta \mid \gamma_1, \dots, \gamma_n),$$

i.e. we have a multiple cover formula

$$\text{GW}_{0,\beta}(X)(\gamma_1, \dots, \gamma_n) = \sum_{k \mid \beta} \frac{1}{k^{3-n}} \cdot \text{DT}_4(X)(\beta/k \mid \gamma_1, \dots, \gamma_n).$$

In [5], the authors did not specify how to choose the orientation in order to match invariants. They checked Conjecture 0.1 in examples, where the corresponding orientation is chosen through case by case study.

0.2. GV/DT₄ conjecture on Fano 3-folds. In this paper, we study Conjecture 0.1 when $X = K_Y$ is the total space of the canonical bundle of a Fano 3-fold Y . Although X is no longer compact, the moduli scheme $M_{X,\beta}$ of one dimensional stable sheaves is compact. In fact, by the negativity of K_Y , the zero section map $\iota : Y \hookrightarrow X$ induces an isomorphism

$$\iota_* : M_{Y,\beta} \cong M_{X,\beta}$$

between moduli schemes of one dimensional stable sheaves E 's with $[E] = \beta$ and $\chi(E) = 1$ on Y and X respectively.

Then the DT_4 virtual class of $M_{X,\beta}$ reduces to the DT_3 virtual class of $M_{Y,\beta}$ (ref. [3]). As for insertions, let

$$\tau : H^m(Y) \rightarrow H^{m-2}(M_{Y,\beta}), \quad \tau(\gamma) = \pi_{M*}(\pi_Y^* \gamma \cup \text{ch}_2(\mathcal{E})),$$

where π_Y, π_M are projections from $Y \times M_{Y,\beta}$ to corresponding factors, and $\text{ch}_2(\mathcal{E})$ is the Poincaré dual of the fundamental cycle of the universal sheaf \mathcal{E} over $M_{Y,\beta} \times Y$.

For $\gamma_i \in H^{m_i}(Y, \mathbb{Z})$, $1 \leq i \leq n$, the *twisted DT₃ invariant* is defined by

$$(0.1) \quad \text{DT}_3^{\text{twist}}(Y)(\beta \mid \gamma_1, \dots, \gamma_n) := (-1)^{c_1(Y) \cdot \beta - 1} \cdot \int_{[M_{Y,\beta}]^{\text{vir}}} \prod_{i=1}^n \tau(\gamma_i) \in \mathbb{Z},$$

where $[M_{Y,\beta}]^{\text{vir}} \in H_2(M_{Y,\beta}, \mathbb{Z})$ is the DT_3 virtual class [15].

As for GW invariants, one can identify GW invariants of X with the so-called *twisted GW invariants* of Y . For $\gamma_i \in H^{m_i}(Y, \mathbb{Z})$, $1 \leq i \leq n$, and the rank $(\int_{\beta} c_1(Y) - 1)$ vector bundle

$$(0.2) \quad -\mathbf{R}\pi_* f^* K_Y \in K(\overline{M}_{0,n}(Y, \beta)),$$

where $\pi : \mathcal{C} \rightarrow \overline{M}_{0,n}(X, \beta)$ is the universal curve and $f : \mathcal{C} \rightarrow Y$ is the universal map, one defines

$$(0.3) \quad \text{GW}_{0,\beta}^{\text{twist}}(Y)(\gamma_1, \dots, \gamma_n) := \int_{[\overline{M}_{0,n}(Y, \beta)]^{\text{vir}}} e(-\mathbf{R}\pi_* f^* K_Y) \cup \prod_{i=1}^n \text{ev}_i^*(\gamma_i) \in \mathbb{Q},$$

where $\text{ev}_i : \overline{M}_{0,n}(Y, \beta) \rightarrow Y$ is the i -th evaluation map.

Then Conjecture 0.1 can be rephrased completely on Y .

Conjecture 0.2. (Conjecture 1.2) *Let Y be a smooth Fano 3-fold. Then we have*

$$\text{GW}_{0,\beta}^{\text{twist}}(Y)(\gamma_1, \dots, \gamma_n) = \sum_{k \mid \beta} \frac{1}{k^{3-n}} \cdot \text{DT}_3^{\text{twist}}(Y)(\beta/k \mid \gamma_1, \dots, \gamma_n).$$

Remarkably, when restricting to local CY 4-fold K_Y , we conjecture the ambiguity in choosing the right orientation in Conjecture 0.1 can be fixed by introducing twisted DT_3 invariants.

In Section 1.3, we give a heuristic explanation of why the sign $(-1)^{c_1(Y) \cdot \beta - 1}$ in twisted DT_3 invariants should give the correct orientation for matching with twisted GW invariants. We also compute examples and check Conjecture 0.2 in those cases.

0.3. Verifications of Conjecture 0.2: compact examples. For compact Fano 3-folds, we check Conjecture 0.2 in the following examples.

For the line class on Fano hypersurfaces in \mathbb{P}^4 , we have

Proposition 0.3. (Proposition 2.1) *Let $Y_d \subseteq \mathbb{P}^4$ be a smooth hypersurface of degree $d \leq 4$. Then Conjecture 0.2 is true for the line class $\beta = [l] \in H_2(Y_d, \mathbb{Z})$.*

For multiple fiber classes for \mathbb{P}^1 -bundles, we have

Proposition 0.4. (Proposition 2.3) *Let S be a del-Pezzo surface, $Y = S \times \mathbb{P}^1$ be the product. Then Conjecture 0.2 is true for $\beta = n \cdot [\mathbb{P}^1]$ with $n \geq 1$.*

When $Y = S \times \mathbb{P}^1$ is the product of a del-Pezzo surface S with \mathbb{P}^1 and $\beta \in H_2(S, \mathbb{Z}) \subseteq H_2(Y, \mathbb{Z})$, we can identify twisted DT_3 (resp. GW) invariants with DT_3 (resp. GW) invariants of a non-compact Calabi-Yau 3-fold K_S up to multiplying some constant.

Proposition 0.5. (Proposition 2.5) *In the above setting, we have:*

(1) *If $\gamma = (\gamma_1, d) \in H^2(S) \otimes H^2(\mathbb{P}^1) \subseteq H^4(Y)$, then*

$$\begin{aligned} \text{DT}_3^{\text{twist}}(Y)(\beta \mid \gamma) &= d(\beta \cdot \gamma_1) \cdot \text{DT}_3(K_S)(\beta), \\ \text{GW}_{0,\beta}^{\text{twist}}(Y)(\gamma) &= d(\beta \cdot \gamma_1) \cdot \text{GW}_{0,\beta}(K_S). \end{aligned}$$

(2) *If $\gamma \in H^4(S) \subseteq H^4(Y)$, then*

$$\text{DT}_3^{\text{twist}}(Y)(\beta \mid \gamma) = \text{GW}_{0,\beta}^{\text{twist}}(Y)(\gamma) = 0.$$

Then Conjecture 0.2 is equivalent to Katz's conjecture [11] (see Conjecture 2.6). In particular, by [5, Corollary A.7.], we obtain

Theorem 0.6. (Theorem 2.8) *Let S be a toric del-Pezzo surface and $Y = S \times \mathbb{P}^1$ be the product. Then Conjecture 0.2 is true for $\beta \in H_2(S, \mathbb{Z}) \subseteq H_2(Y, \mathbb{Z})$.*

0.4. Verifications of Conjecture 0.2: non-compact examples. Let $Y = \text{Tot}_S(L) \rightarrow S$ be the total space of a negative line bundle L over a del-Pezzo surface S , and $i : S \rightarrow Y$ be the zero section. Although Y is non-compact, $M_{Y,\beta}$ is compact and has a well-defined virtual class as the push-forward morphism

$$i_* : M_{S,\beta} \rightarrow M_{Y,\beta}$$

is an isomorphism between moduli schemes of one dimensional stable sheaves on S and Y respectively. The virtual class of $M_{Y,\beta}$ is the Euler class of a vector bundle over $M_{S,\beta}$ and we can conclude the following symmetric property of twisted DT_3 invariants.

Proposition 0.7. (Proposition 3.2, Corollary 3.3) *Let S be a del-Pezzo surface and $Y_1 = \text{Tot}_S(L)$, $Y_2 = \text{Tot}_S(L^{-1} \otimes K_S)$ be the total space of ample line bundles L^{-1} , $L \otimes K_S^{-1}$ respectively on S . Then we have*

$$\text{DT}_3^{\text{twist}}(Y_1)(\beta \mid [\text{pt}]) = \text{DT}_3^{\text{twist}}(Y_2)(\beta \mid [\text{pt}]).$$

In particular, Conjecture 1.2 is true for Y_1 if and only if it is true for Y_2 .

Combining with previous computations [5, Sect. 3.2], we obtain

Proposition 0.8. (Proposition 3.4) *Let $Y = \mathcal{O}_{\mathbb{P}^2}(-1)$ or $\mathcal{O}_{\mathbb{P}^2}(-2)$ and $\beta = d[l] \in H_2(\mathbb{P}^2, \mathbb{Z})$ be the degree d class. Then Conjecture 1.2 is true when $d \leq 3$.*

Let $Y = \text{Tot}_{\mathbb{P}^1}(\mathcal{O}_{\mathbb{P}^1}(l_1) \oplus \mathcal{O}_{\mathbb{P}^1}(l_2))$ be the total space of a rank two bundle over \mathbb{P}^1 with Fano condition $l_1 + l_2 \geq -1$ and $T \cong (\mathbb{C}^*)^2$ be a torus acting on the fibers. The moduli scheme $M_{Y,d}$ of one dimensional stable sheaves F 's with $[F] = d[\mathbb{P}^1] \in \mathbb{Z}[\mathbb{P}^1]$ and $\chi(F) = 1$ is non-compact with compact T -fixed locus. We may define its twisted equivariant DT_3 invariants by virtual localization formula [9].

Definition 0.9. (Definition 3.7) *The equivariant twisted DT_3 invariant of $M_{Y,d}$ is*

$$\text{DT}_3^{\text{twist}}(Y)(d) := (-1)^{d(l_1+l_2)-1} \int_{[M_{Y,d}^T]^{\text{vir}}} e_T(N^{\text{vir}}) \in \mathbb{Q}(\lambda_1, \lambda_2),$$

where N^{vir} is the virtual normal bundle of $M_{Y,d}^T \hookrightarrow M_{Y,d}$.

An equivariant version of Conjecture 1.2 can be proposed as follows:

Conjecture 0.10. (Conjecture 3.8) *Let $Y = \text{Tot}_{\mathbb{P}^1}(\mathcal{O}_{\mathbb{P}^1}(l_1) \oplus \mathcal{O}_{\mathbb{P}^1}(l_2))$ with $l_1 + l_2 \geq -1$. Then*

$$\text{GW}_{0,d}^{\text{twist}}(Y) = \sum_{k \mid d} \frac{1}{k^3} \text{DT}_3^{\text{twist}}(Y)(d/k).$$

Here $\text{GW}_{0,d}^{\text{twist}}(Y)$ is the equivariant twisted GW invariant of Y and the multiple cover coefficient is $1/k^3$ instead of $1/k^2$ due to the absence of insertions.

Combining with [5, Thm. 4.12], we have:

Theorem 0.11. (Theorem 3.9) *Definition 0.9 for $Y = \text{Tot}_{\mathbb{P}^1}(\mathcal{O}_{\mathbb{P}^1}(l_1) \oplus \mathcal{O}_{\mathbb{P}^1}(l_2))$ recovers the definition of equivariant DT_4 invariants for $X = K_Y$ in (4.9), (4.17) of [5], i.e. for $d = 1, 2$,*

$$\text{DT}_3^{\text{twist}}(Y)(d) = \text{DT}_4(X)(d[\mathbb{P}^1]).$$

In particular, Conjecture 0.10 is true when (i) $d = 1$ and (ii) $d = 2$, $|l_1|, |l_2| \leq 10$.

1. DEFINITIONS AND CONJECTURES

1.1. Review of GV/DT₄ conjecture. Let X be a smooth projective Calabi-Yau 4-fold, i.e. $K_X \cong \mathcal{O}_X$. Fixing $\beta \in H_2(X, \mathbb{Z})$ and $n \geq 0$, the genus 0 Gromov-Witten invariants of X are defined using insertions: for integral classes $\gamma_i \in H^{m_i}(X, \mathbb{Z})$, $1 \leq i \leq n$, one defines

$$\mathrm{GW}_{0,\beta}(X)(\gamma_1, \dots, \gamma_n) := \int_{[\overline{M}_{0,n}(X,\beta)]^{\mathrm{vir}}} \prod_{i=1}^n \mathrm{ev}_i^*(\gamma_i),$$

where $\mathrm{ev}_i: \overline{M}_{0,n}(X, \beta) \rightarrow X$ is the i -th evaluation map.

The Gopakumar-Vafa type invariants

$$(1.1) \quad n_{0,\beta}(X)(\gamma_1, \dots, \gamma_n) \in \mathbb{Q}$$

are defined by Klemm-Pandharipande [13] in terms of the identity

$$\sum_{\beta > 0} \mathrm{GW}_{0,\beta}(X)(\gamma_1, \dots, \gamma_n) q^\beta = \sum_{\beta > 0} n_{0,\beta}(X)(\gamma_1, \dots, \gamma_n) \sum_{d=1}^{\infty} d^{n-3} q^{d\beta},$$

and conjectured to be integers.

In [5], Cao-Maulik-Toda gave a sheaf-theoretic interpretation of the above invariants (1.1) in terms of Donaldson-Thomas invariants for CY 4-folds (called DT₄ invariants). More specifically, we consider the moduli scheme $M_{X,\beta}$ of 1-dimensional stable sheaves on X with Chern character $(0, 0, 0, \beta, 1)$. Let $\mathcal{E} \rightarrow X \times M_{X,\beta}$ be its universal sheaf and

$$(1.2) \quad \mathcal{L} := \det(\mathbf{R}\mathcal{H}om_{\pi_M}(\mathcal{E}, \mathcal{E})) \in \mathrm{Pic}(M_{X,\beta}), \quad \pi_M: X \times M_{X,\beta} \rightarrow M_{X,\beta},$$

be the determinant line bundle, equipped with a symmetric pairing Q induced by Serre duality. By the results of [3, 1], if the structure group of the line bundle (\mathcal{L}, Q) can be reduced from $O(1, \mathbb{C})$ to $SO(1, \mathbb{C}) = \{1\}$ (this is always true by [2]), there exists a DT₄ virtual class

$$(1.3) \quad [M_{X,\beta}]^{\mathrm{vir}} \in H_2(M_{X,\beta}, \mathbb{Z}),$$

which depends on the choice of an orientation, i.e. the reduction of the structure group of (\mathcal{L}, Q) . The choices form a torsor for $H^0(M_{X,\beta}, \mathbb{Z}_2)$.

In order to define counting invariants, we require insertions. Define the map τ by

$$\tau: H^m(X) \rightarrow H^{m-2}(M_{X,\beta}), \quad \tau(\gamma) = \pi_{M*}(\pi_X^* \gamma \cup \mathrm{ch}_3(\mathcal{E})),$$

where π_X, π_M are projections from $X \times M_{X,\beta}$ to corresponding factors, and $\mathrm{ch}_3(\mathcal{E})$ is the Poincaré dual of the fundamental cycle of the universal sheaf \mathcal{E} .

For $\gamma_i \in H^{m_i}(X, \mathbb{Z})$, $1 \leq i \leq n$, the DT₄ invariant is defined by

$$\mathrm{DT}_4(X)(\beta \mid \gamma_1, \dots, \gamma_n) := \int_{[M_{X,\beta}]^{\mathrm{vir}}} \prod_{i=1}^n \tau(\gamma_i).$$

Conjecture 1.1. ([5, Conjecture 1.3]) *The identity*

$$n_{0,\beta}(X)(\gamma_1, \dots, \gamma_n) = \mathrm{DT}_4(X)(\beta \mid \gamma_1, \dots, \gamma_n)$$

holds for a suitable choice of orientation in defining the RHS.

In particular, we have the multiple cover formula

$$\mathrm{GW}_{0,\beta}(X)(\gamma) = \sum_{k \mid \beta} \frac{1}{k^2} \cdot \mathrm{DT}_4(X)(\beta/k \mid \gamma).$$

1.2. The conjecture on Fano 3-folds. When $X = K_Y$ is the total space of the canonical bundle of a smooth Fano 3-fold Y . Conjecture 1.1 can be completely rephrased on Y as follows.

Twisted GW invariants. Fixing $\beta \in H_2(Y, \mathbb{Z})$ and $n \geq 0$, the *twisted* genus 0 Gromov-Witten invariants of Y are defined using insertions: for integral classes $\gamma_i \in H^{m_i}(Y, \mathbb{Z})$, $1 \leq i \leq n$, and the rank $(\int_\beta c_1(Y) - 1)$ vector bundle

$$(1.4) \quad -\mathbf{R}\pi_* f^* K_Y \in K(\overline{M}_{0,n}(Y, \beta)),$$

where $\pi: \mathcal{C} \rightarrow \overline{M}_{0,n}(X, \beta)$ is the universal curve and $f: \mathcal{C} \rightarrow Y$ is the universal map, one defines

$$(1.5) \quad \mathrm{GW}_{0,\beta}^{\mathrm{twist}}(Y)(\gamma_1, \dots, \gamma_n) := \int_{[\overline{M}_{0,n}(Y,\beta)]^{\mathrm{vir}}} e(-\mathbf{R}\pi_* f^* K_Y) \cup \prod_{i=1}^n \mathrm{ev}_i^*(\gamma_i) \in \mathbb{Q},$$

where $\mathrm{ev}_i: \overline{M}_{0,n}(Y, \beta) \rightarrow Y$ is the i -th evaluation map.

Twisted DT₃ invariants. Let $M_{Y,\beta}$ be the moduli scheme of one dimensional stable sheaves E 's with $[E] = \beta$, $\chi(E) = 1$ (i.e. $\text{ch}(E) = (0, 0, \beta, 1 - \frac{c_1(Y) \cdot \beta}{2})$). We define the map τ by

$$\tau: H^m(Y) \rightarrow H^{m-2}(M_{Y,\beta}), \quad \tau(\gamma) = \pi_{M*}(\pi_Y^* \gamma \cup \text{ch}_2(\mathcal{E})),$$

where π_Y, π_M are projections from $Y \times M_{Y,\beta}$ to corresponding factors, and $\text{ch}_2(\mathcal{E})$ is the Poincaré dual of the fundamental cycle of the universal sheaf \mathcal{E} .

For $\gamma_i \in H^{m_i}(Y, \mathbb{Z})$, $1 \leq i \leq n$, the *twisted* DT₃ invariant is

$$(1.6) \quad \text{DT}_3^{\text{twist}}(Y)(\beta \mid \gamma_1, \dots, \gamma_n) := (-1)^{c_1(Y) \cdot \beta - 1} \cdot \int_{[M_{Y,\beta}]^{\text{vir}}} \prod_{i=1}^n \tau(\gamma_i) \in \mathbb{Z},$$

where $[M_{Y,\beta}]^{\text{vir}} \in H_2(M_{Y,\beta}, \mathbb{Z})$ is the DT₃ virtual class [15].

When restricting to local CY 4-fold K_Y , we conjecture that the ambiguity in choosing the right orientation in Conjecture 1.1 can be fixed by introducing twisted DT₃ invariants.

Conjecture 1.2. *Let Y be a smooth Fano 3-fold. Then we have*

$$\text{GW}_{0,\beta}^{\text{twist}}(Y)(\gamma_1, \dots, \gamma_n) = \sum_{k \mid \beta} \frac{1}{k^{3-n}} \cdot \text{DT}_3^{\text{twist}}(Y)(\beta/k \mid \gamma_1, \dots, \gamma_n).$$

Remark 1.3. *Note that (1.5) and (1.6) vanish unless*

$$\sum_{i=1}^n (m_i - 2) = 2.$$

If $m_n = 2$, we have

$$\begin{aligned} \text{GW}_{0,\beta}^{\text{twist}}(Y)(\gamma_1, \dots, \gamma_n) &= (\beta \cdot \gamma_n) \cdot \text{GW}_{0,\beta}^{\text{twist}}(Y)(\gamma_1, \dots, \gamma_{n-1}), \\ \text{DT}_3^{\text{twist}}(Y)(\beta \mid \gamma_1, \dots, \gamma_n) &= (\beta \cdot \gamma_n) \cdot \text{DT}_3^{\text{twist}}(Y)(\beta \mid \gamma_1, \dots, \gamma_{n-1}). \end{aligned}$$

Therefore we may assume that $m_i \geq 3$ for all i in Conjecture 1.2. There are two possibilities

- $n = 1$ and $m_1 = 4$,
- $n = 2$ and $m_1 = m_2 = 3$.

In particular, when $H^3(Y, \mathbb{Z}) = 0$, we only need to consider the first case.

1.3. Geometric explanation of the conjecture. Conjecture 1.2 on a Fano 3-fold Y can be formulated in terms of Conjecture 1.1 on a non-compact CY 4-fold $X = K_Y$.

More specifically, by a stability argument (e.g. [3]), one dimensional stable sheaves on X are scheme theoretically supported on the zero section $\iota: Y \rightarrow X$, so a moduli scheme $M_{X,\beta}$ of one dimensional stable sheaves on X is isomorphic to a moduli scheme $M_{Y,\beta}$ of one dimensional stable sheaves on Y via the push-forward by ι . Under this identification, the DT₄ virtual class of $M_{X,\beta}$ can be identified with the DT₃ virtual class of $M_{Y,\beta}$.

As for GW theory, the moduli stack of stable maps to X is the moduli stack of stable maps to the compact divisor $Y \subseteq X$, but the obstruction theories are different, and the difference is measured by the Euler class of the vector bundle (1.4). Hence Conjecture 1.2 can be regarded as a specialization of Conjecture 1.1 on the (non-compact) CY 4-fold $X = K_Y$, and a geometric explanation of the conjecture follows from the one given in [5, Sect. 1.4] modulo the issue of choosing an orientation.

When we restrict to $X = K_Y$, we expect to be able to fix the ambiguity of choosing an orientation by the following reason, where we only consider rational curves and ignore curves of higher genus.

Let Y be a ‘good’ Fano 3-fold, in the following sense:

- (1) any rational curve in Y comes with a compact smooth family of embedded rational curves, whose general member is smooth with normal bundle $\mathcal{O}_{\mathbb{P}^1}(a, b)$ specified as follows;
- (2) when $c_1(Y) \cdot \beta$ is even, then $a = b = \frac{1}{2}c_1(Y) \cdot \beta - 1$;
- (3) when $c_1(Y) \cdot \beta$ is odd, then $a = \frac{1}{2}(c_1(Y) \cdot \beta - 1)$, $b = \frac{1}{2}(c_1(Y) \cdot \beta - 3)$.

Let C be a rational curve in Y with $[C] = \beta \in H_2(Y, \mathbb{Z})$, and $\{C_t\}_{t \in T}$ be the smooth family C sits in. Any one-dimensional stable sheaf $[E] \in M_{Y,\beta}$ supported on $\{C_t\}_{t \in T}$ is \mathcal{O}_{C_t} for some

$t \in T$. For a general $t \in T$ such that $N_{C_t/Y} \cong \mathcal{O}_{\mathbb{P}^1}(a, b)$, there exist isomorphisms

$$\begin{aligned} \mathrm{Ext}_Y^1(E, E) &\cong H^0(C_t, N_{C_t/Y}) \cong \mathbb{C}^{c_1(Y) \cdot \beta}, \\ \mathrm{Ext}_Y^2(E, E) &\cong H^0(C_t, \wedge^2 N_{C_t/Y}) \cong \mathbb{C}^{c_1(Y) \cdot \beta - 1}. \end{aligned}$$

So the obstruction sheaf ($\mathrm{Ext}_Y^2(E, E)$'s) of $M_{Y,\beta}$ extends to a vector bundle ($H^0(C_t, \wedge^2 N_{C_t/Y})$'s) over the family T whose Euler class gives $[M_{Y,\beta}]^{\mathrm{vir}}$. The insertion $\tau(\gamma)$ imposes a codimension one constrain on the deformation space of curves in Y , whose integration against the virtual class gives the DT_3 invariant.

As for a stable map $f : \mathbb{P}^1 \rightarrow Y$, we may view it as a composition of a multiple cover $t : \mathbb{P}^1 \rightarrow \mathbb{P}^1$ and an embedding $i : \mathbb{P}^1 \hookrightarrow Y$ with some C_t as image. The usual obstruction space vanishes by our generic assumption. But to define twisted GW invariants (1.5), we need an extra ‘obstruction bundle’ whose fiber over such f is $H^1(\mathbb{P}^1, f^*K_Y)$. For an embedding f with image C_t , we have canonical isomorphisms

$$H^1(\mathbb{P}^1, f^*K_Y) \cong H^1(C_t, \wedge^2 N_{C_t/Y}^* \otimes K_{C_t}) \cong H^0(C_t, \wedge^2 N_{C_t/Y})^* \cong \mathrm{Ext}_Y^2(E, E)^*.$$

So modulo the multiple cover factor, we can see the obstruction bundle for $M_{Y,\beta}$ is dual to the extra obstruction bundle (with rank $(c_1(Y) \cdot \beta - 1)$) for defining twisted GW invariants. Thus,

$$\mathrm{GW}_{0,\beta}^{\mathrm{twist}}(Y)(\gamma) = \sum_{k|\beta} \frac{(-1)^{c_1(Y) \cdot \frac{\beta}{k} - 1}}{k^2} \cdot \int_{[M_{Y,\beta/k}]^{\mathrm{vir}}} \tau(\gamma),$$

where signs appear since the Euler class of the dual bundle of a bundle E differs with $e(E)$ by a sign $(-1)^{\mathrm{rk} E}$. Conjecture 1.2 in this case then follows from the definition of twisted DT_3 invariants (1.6). Of course, this only gives a geometric explanation (in the ‘good’ Fano 3-fold case) rather than a rigorous proof. We will check the conjecture in examples in the coming sections.

2. COMPACT EXAMPLES

2.1. Line classes on Fano hypersurfaces. We verify Conjecture 1.2 for line classes on Fano hypersurfaces in \mathbb{P}^4 .

Proposition 2.1. *Let $Y_d \subseteq \mathbb{P}^4$ be a smooth hypersurface of degree $d \leq 4$. Then Conjecture 1.2 is true for the line class $\beta = [l] \in H_2(Y_d, \mathbb{Z})$.*

Proof. From the deformation invariance of DT_3 and GW invariants, we may assume Y_d to be a general hypersurface. By [14, Thm. 4.3, pp. 266], the Hilbert scheme $\mathrm{Hilb}^{t+1}(Y_d)$ of lines on Y_d is connected and smooth of dimension $(5 - d)$. For any line $l \subseteq Y_d$, we have

$$N_{l/Y_d} \cong \mathcal{O}_{\mathbb{P}^1}(a_1) \oplus \mathcal{O}_{\mathbb{P}^1}(a_2).$$

By [14, Ex. 4.4, 4.5, pp. 269], we know $(2 - d) \leq a_i \leq 1$ and $a_i = 0$ or -1 if $d = 4$. Hence

$$\begin{aligned} N_{l/Y_d} &\cong \mathcal{O}_{\mathbb{P}^1}^{\oplus(d-1)} \oplus \mathcal{O}_{\mathbb{P}^1}(1)^{\oplus(3-d)}, \text{ if } d = 1, 2, \\ N_{l/Y_d} &\cong \mathcal{O}_{\mathbb{P}^1}(-1) \oplus \mathcal{O}_{\mathbb{P}^1}(1) \text{ or } N_{l/Y_d} \cong \mathcal{O}_{\mathbb{P}^1} \oplus \mathcal{O}_{\mathbb{P}^1}, \text{ if } d = 3, \\ N_{l/Y_d} &\cong \mathcal{O}_{\mathbb{P}^1} \oplus \mathcal{O}_{\mathbb{P}^1}(-1), \text{ if } d = 4. \end{aligned}$$

Any one dimensional stable sheaf supported on a line is its structure sheaf and the moduli space M_l of such one dimensional sheaves is isomorphic to the Hilbert scheme $\mathrm{Hilb}^{t+1}(Y_d)$.

By the possibility of the normal bundle of $l \subseteq Y_d$, the deformation and obstruction spaces of M_l satisfies

$$\begin{aligned} \mathrm{Ext}^1(\mathcal{O}_l, \mathcal{O}_l) &\cong H^0(l, N_{l/Y_d}) \cong \mathbb{C}^{5-d}, \\ (2.1) \quad \mathrm{Ext}^2(\mathcal{O}_l, \mathcal{O}_l) &\cong H^0(l, \wedge^2 N_{l/Y_d}) \cong H^1(l, K_{Y_d}|_l)^* \cong \mathbb{C}^{4-d}, \end{aligned}$$

so its virtual class is the Euler class of the obstruction bundle.

The deformation and obstruction spaces of $\overline{\mathcal{M}}_{0,0}(Y_d, [l])$ satisfies

$$H^0(l, N_{l/Y_d}) \cong \mathbb{C}^{5-d}, \quad H^1(l, N_{l/Y_d}) = 0.$$

So the virtual class is its usual fundamental class.

To relate them, note that we have an isomorphism

$$\begin{aligned} \overline{\mathcal{M}}_{0,0}(Y_d, [l]) &\rightarrow M_l, \\ (f : C \rightarrow Y_d) &\mapsto \mathcal{O}_{f(C)}, \end{aligned}$$

a forgetful map

$$\phi : \overline{\mathcal{M}}_{0,1}(Y_d, [l]) \rightarrow \overline{\mathcal{M}}_{0,0}(Y_d, [l]) \cong M_l,$$

$$\phi : (f : C \rightarrow Y_d, p \in C) \mapsto (f : C \rightarrow Y_d) \mapsto \mathcal{O}_{f(C)},$$

and an embedding

$$\begin{aligned} i &= (\phi, \text{ev}) : \overline{\mathcal{M}}_{0,1}(Y_d, [l]) \hookrightarrow M_l \times Y_d, \\ i(f : C \rightarrow Y_d, p \in C) &= (\mathcal{O}_{f(C)}, f(P)), \end{aligned}$$

whose image is the universal curve $\mathcal{C} \subseteq M_l \times Y_d$, where we use identification $M_l \cong \text{Hilb}^{t+1}(Y_d)$.

Since \mathcal{C} and $M_l \times Y_d$ are smooth,

$$(\mathbf{R}\pi_M)_* \mathbf{R}\mathcal{H}om(\mathcal{O}_{\mathcal{C}}, \mathcal{O}_{\mathcal{C}}) \cong (\mathbf{R}\pi_M)_* i_* \left(\bigwedge^* \mathcal{N}_{\mathcal{C}/(M_l \times Y_d)} \right),$$

where $\pi_M : M_l \times Y_d \rightarrow M_l$ is the natural projection.

By (2.1), the obstruction bundle Ob_{M_l} of M_l is

$$(\pi_M)_* i_* \left(\bigwedge^2 \mathcal{N}_{\mathcal{C}/(M_l \times Y_d)} \right) \cong (\pi_M)_* (i_*(K_{\mathcal{C}}) \otimes K_{M_l \times Y_d}^{-1}),$$

by the tangent-normal bundle exact sequence and the projection formula.

We claim there is an isomorphism of vector bundles:

$$(\mathbf{R}^1 \phi_* \text{ev}^* K_{Y_d})^* \cong (\pi_M)_* (i_*(K_{\overline{\mathcal{M}}_{0,1}(Y_d, [l])}) \otimes K_{M_l \times Y_d}^{-1}).$$

In fact, by the Grothendieck-Verdier duality (ref. [10, Thm. 3.34, pp. 86]), we have

$$\begin{aligned} \mathbf{R}\mathcal{H}om(\mathbf{R}\phi_* \text{ev}^* K_{Y_d}, \mathcal{O}_{M_l}) &\cong \mathbf{R}\mathcal{H}om(\mathbf{R}(\pi_M)_* i_* i^* \pi_Y^* K_{Y_d}, \mathcal{O}_{M_l}) \\ &\cong \mathbf{R}\mathcal{H}om(\mathbf{R}(\pi_M)_* (\pi_Y^* K_{Y_d} \otimes \mathcal{O}_{\mathcal{C}}), \mathcal{O}_{M_l}) \\ &\cong \mathbf{R}(\pi_M)_* \mathbf{R}\mathcal{H}om(\pi_Y^* K_{Y_d} \otimes \mathcal{O}_{\mathcal{C}}, \pi_Y^* K_{Y_d}[3]) \\ &\cong \mathbf{R}(\pi_M)_* \mathbf{R}\mathcal{H}om(\mathcal{O}_{\mathcal{C}}, \mathcal{O}_{M_l \times Y_d}[3]) \\ &\cong \mathbf{R}(\pi_M)_* (i_* K_{\overline{\mathcal{M}}_{0,1}(Y_d, [l])} \otimes K_{M_l \times Y_d}^{-1}[1]), \end{aligned}$$

where the last isomorphism is by (ref. [10, Cor. 3.40, pp. 89]) and $\pi_Y : M_l \times Y_d \rightarrow Y_d$ denotes the natural projection. Thus, we have an isomorphism

$$(2.2) \quad \text{Ob}_{M_l} \cong (\mathbf{R}^1 \phi_* \text{ev}^* K_{Y_d})^*$$

of rank $(4-d)$ vector bundles. As in [6, pp. 182], we have a commutative diagram

$$\begin{array}{ccccc} & & \text{ev}_2 & & \\ & & \curvearrowright & & \\ \overline{\mathcal{M}}_{0,2}(Y_d, [l]) & \xrightarrow{\phi_2} & \overline{\mathcal{M}}_{0,1}(Y_d, [l]) & \xrightarrow{\text{ev}_1} & Y_d \\ \downarrow \pi_2 & & \downarrow \pi_1 & & \\ \overline{\mathcal{M}}_{0,1}(Y_d, [l]) & \xrightarrow{\phi_1} & \overline{\mathcal{M}}_{0,0}(Y_d, [l]), & & \end{array}$$

where ϕ_i forgets the first marked point and π_i forgets the last marked point. Then base change gives an isomorphism

$$(2.3) \quad (\pi_2)_* \text{ev}_2^* K_{Y_d} \cong \phi_1^* (\pi_1)_* \text{ev}_1^* K_{Y_d}.$$

By the definition, for $\gamma \in H^4(Y_d)$, we have

$$\begin{aligned} \text{DT}_3^{\text{twist}}(Y_d)(l \mid \gamma) &= (-1)^{4-d} \cdot \int_{[M_l]_{\text{vir}}} (\pi_M)_* (\text{PD}[\mathcal{C}] \cup \pi_Y^* \gamma) \\ &= (-1)^{4-d} \cdot \int_{[M_l \times Y_d]} \text{PD}[\mathcal{C}] \cup \pi_Y^* \gamma \cup \pi_M^* e(\text{Ob}_{M_l}) \\ &= (-1)^{4-d} \cdot \int_{[\overline{\mathcal{M}}_{0,1}(Y_d, [l])]} i^* (\pi_Y^* \gamma \cup \pi_M^* e(\text{Ob}_{M_l})) \\ &= (-1)^{4-d} \cdot \int_{[\overline{\mathcal{M}}_{0,1}(Y_d, [l])]} \text{ev}^* \gamma \cup e(\phi^* \text{Ob}_{M_l}) \\ &= \int_{[\overline{\mathcal{M}}_{0,1}(Y_d, [l])]} \text{ev}^* \gamma \cup e(\phi^* \mathbf{R}^1 \phi_* \text{ev}^* K_{Y_d}) \\ &= \int_{[\overline{\mathcal{M}}_{0,1}(Y_d, [l])]} \text{ev}^* \gamma \cup e(\mathbf{R}^1(\pi_2)_* \text{ev}_2^* K_{Y_d}) \\ &= \text{GW}_{0,l}^{\text{twist}}(Y_d)(\gamma), \end{aligned}$$

where the second and third to last equality is by (2.2), (2.3).

As for insertions $\gamma_1, \gamma_2 \in H^3(Y, \mathbb{Z})$, we consider the case when $d = 4$ for simplicity (in this case, the obstruction bundle of M_l is zero), other cases can be proved in a similar way.

Let $Y = Y_4$, by considering the commutative diagram of embeddings

$$\begin{array}{ccc} \overline{\mathcal{M}}_{0,2}(Y, [l]) & \xrightarrow{i_1=(\phi_1, \text{ev}_1)} & \overline{\mathcal{M}}_{0,1}(Y, [l]) \times Y \\ \downarrow i_2=(\phi_2, \text{ev}_2) & \searrow i & \downarrow i_4:(f, y) \mapsto (\phi(f), y, \text{ev}(f)) \\ \overline{\mathcal{M}}_{0,1}(Y, [l]) \times Y & \xrightarrow{i_3=(\phi, \text{ev}) \times \text{Id}_Y} & \overline{\mathcal{M}}_{0,0}(Y, [l]) \times Y \times Y, \end{array}$$

where ϕ_i forgets the i -th marked point and ev_i is the i -th evaluation map, we get

$$\text{ev}_1^* \gamma_1 \cup \text{ev}_2^* \gamma_2 = i^*(1 \cup \gamma_1 \cup \gamma_2).$$

$$\begin{aligned} \text{GW}_{0,l}^{\text{twist}}(Y)(\gamma_1, \gamma_2) &= \int_{[\overline{\mathcal{M}}_{0,2}(Y, [l])]} \text{ev}_1^* \gamma_1 \cup \text{ev}_2^* \gamma_2 \\ &= \int_{[\overline{\mathcal{M}}_{0,2}(Y, [l])]} i^*(1 \cup \gamma_1 \cup \gamma_2) \\ &= \int_{[\overline{\mathcal{M}}_{0,0}(Y, [l]) \times Y \times Y]} (1 \cup \gamma_1 \cup \gamma_2) \cup \text{PD}\left([i(\overline{\mathcal{M}}_{0,2}(Y, [l]))]\right) \\ &= \int_{[\overline{\mathcal{M}}_{0,0}(Y, [l]) \times Y \times Y]} (1 \cup \gamma_1 \cup \gamma_2) \cup \text{PD}\left([i_3(\overline{\mathcal{M}}_{0,1}(Y, [l]) \times Y)]\right) \\ &\quad \cup \text{PD}\left([i_4(\overline{\mathcal{M}}_{0,1}(Y, [l]) \times Y)]\right) \\ &= \int_{[\overline{\mathcal{M}}_{0,0}(Y, [l])]} (\pi_M)_*(\text{PD}[\mathcal{C}] \cup \pi_Y^* \gamma_1) \cup (\pi_M)_*(\text{PD}[\mathcal{C}] \cup \pi_Y^* \gamma_2) \\ &= \text{DT}_3^{\text{twist}}(Y)(\beta \mid \gamma_1, \gamma_2), \end{aligned}$$

where we use $i(\overline{\mathcal{M}}_{0,2}(Y, [l])) = i_3(\overline{\mathcal{M}}_{0,1}(Y, [l]) \times Y) \cap i_4(\overline{\mathcal{M}}_{0,1}(Y, [l]) \times Y)$. \square

Remark 2.2. *If Y is a smooth intersection of $Gr(2, 5)$, a quadric and a \mathbb{P}^7 in \mathbb{P}^9 , then Y is a Fano 3-fold. For general choice of Y , the Hilbert scheme of lines on it is a connected smooth projective curve (ref. [7, Prop. 5.1]). One can easily check Proposition 2.1 also holds for Y .*

2.2. Multiple fiber classes for \mathbb{P}^1 -bundles.

Proposition 2.3. *Let S be a del-Pezzo surface and $Y = S \times \mathbb{P}^1$ be the product. Then Conjecture 1.2 is true for $\beta = n \cdot [\mathbb{P}^1]$ with $n \geq 1$.*

Proof. By a stability argument [5, Lemma 2.2], any one dimensional stable sheaf $E \in M_{Y,\beta}$ is scheme theoretically supported on $\iota : \mathbb{P}^1 \times \{p\} \hookrightarrow Y$ for some $p \in S$, hence $E = \iota_* \mathcal{O}_{\mathbb{P}^1}$ by the classification of stable vector bundle on \mathbb{P}^1 . So we have

$$M_{Y,\beta} \cong S, \text{ if } n = 1; \quad M_{Y,\beta} = \emptyset, \text{ if } n > 1.$$

We fix $n = 1$. Direct calculations give isomorphisms

$$\text{Ext}^1(\iota_* \mathcal{O}_{\mathbb{P}^1}, \iota_* \mathcal{O}_{\mathbb{P}^1}) \cong H^0(\mathbb{P}^1, \mathcal{O}_{\mathbb{P}^1}^{\oplus 2}) \cong \mathbb{C}^2,$$

$$\text{Ext}^2(\iota_* \mathcal{O}_{\mathbb{P}^1}, \iota_* \mathcal{O}_{\mathbb{P}^1}) \cong H^0(\mathbb{P}^1, \mathcal{O}_{\mathbb{P}^1}) \cong \mathbb{C}.$$

So local Kuranishi maps of $M_{Y,\beta}$ are zero and its virtual class is the Euler class of the corresponding obstruction bundle, which can be determined as follows.

Let $\Delta_{S \times S} \hookrightarrow S \times S$ be the diagonal and denote the closed subscheme

$$i : \mathcal{Z} := \Delta_{S \times S} \times \mathbb{P}^1 \hookrightarrow S \times S \times \mathbb{P}^1 \cong M_{Y,\beta} \times Y,$$

in the product $M_{Y,\beta} \times Y$. Then the universal sheaf of $M_{Y,\beta}$ is the structure sheaf $\mathcal{O}_{\mathcal{Z}}$ which satisfies

$$\mathbf{R}\mathcal{H}om(\mathcal{O}_{\mathcal{Z}}, \mathcal{O}_{\mathcal{Z}}) \cong i_* \left(\bigwedge^* \pi^* TS \right),$$

where $\pi : S \times \mathbb{P}^1 \rightarrow S$ is the projection and we identify $\Delta_{S \times S} \cong S$. Under the isomorphism $M_{Y,\beta} \cong S$, the obstruction bundle Ob of $M_{Y,\beta}$ satisfies

$$\text{Ob} \cong \bigwedge^2 TS.$$

As for insertions, by Remark 1.3, we only need to consider

$$\gamma \in H^4(Y, \mathbb{Z}) \cong H^4(S, \mathbb{Z}) \oplus H^2(\mathbb{P}^1, \mathbb{Z}) \otimes H^2(S, \mathbb{Z}),$$

since $H^3(Y, \mathbb{Z}) = 0$. Then

$$\tau(\gamma) = \pi_{M*}(\pi_Y^* \gamma \cup [\mathcal{Z}]) = 0, \text{ if } \gamma \in H^4(S, \mathbb{Z}),$$

$$\tau(\gamma) = d_1 \cdot \gamma_2, \text{ if } \gamma = d_1 \otimes \gamma_2 \in H^2(\mathbb{P}^1, \mathbb{Z}) \otimes H^2(S, \mathbb{Z}),$$

where π_M, π_Y are projections from $M_{Y, \beta} \times Y$ to corresponding factors. Hence

$$\text{DT}_3^{\text{twist}}(Y)(\beta \mid \gamma) = 0, \text{ if } \gamma \in H^4(S, \mathbb{Z}),$$

$$\text{DT}_3^{\text{twist}}(Y)(\beta \mid d_1 \otimes \gamma_2) = d_1 \int_S e\left(\bigwedge^2 TS\right) \cup \gamma_2, \text{ if } \gamma_2 \in H^2(S, \mathbb{Z}).$$

As for GW invariants, let $\beta = n \cdot [\mathbb{P}^1]$, we have

$$\overline{M}_{0,1}(Y, \beta) \cong \overline{M}_{0,1}(\mathbb{P}^1, n) \times S.$$

Given a map $t : \mathbb{P}^1 \rightarrow Y = \mathbb{P}^1 \times S$ with $t_*[\mathbb{P}^1] = n[\mathbb{P}^1]$, we have

$$H^*(\mathbb{P}^1, t^*TY) \cong H^*(\mathbb{P}^1, t^*T\mathbb{P}^1) \oplus H^*(\mathbb{P}^1, c^*TS),$$

$$H^*(\mathbb{P}^1, t^*K_Y) \cong H^*(\mathbb{P}^1, t^*K_{\mathbb{P}^1} \otimes c^*K_S),$$

where c is a constant map to some point of S . Hence

$$[\overline{M}_{0,1}(Y, \beta)]^{\text{vir}} = [\overline{M}_{0,1}(\mathbb{P}^1, n)]^{\text{vir}} \otimes [S] \in A_{2n+1}(\overline{M}_{0,1}(Y, \beta)),$$

and for the universal curve $\pi : \mathcal{C} \rightarrow \overline{M}_{0,1}(Y, \beta)$ and map $f : \mathcal{C} \rightarrow Y$,

$$\begin{aligned} e(-\mathbf{R}\pi_* f^* K_Y) &= c_{2n-1}(-\mathbf{R}\pi_* f^* K_Y) \\ &= c_{2n-2}(-\mathbf{R}\pi_* f^* K_{\mathbb{P}^1}) \cup e(K_S) + c_{2n-1}(-\mathbf{R}\pi_* f^* K_{\mathbb{P}^1}) \\ &= c_{2n-2}(-\mathbf{R}\pi_* f^* K_{\mathbb{P}^1}) \cup e(K_S), \end{aligned}$$

where $c_{2n-1}(-\mathbf{R}\pi_* f^* K_{\mathbb{P}^1})$ is the Euler class of the obstruction bundle for stable maps to $K_{\mathbb{P}^1}$, which is zero as $K_{\mathbb{P}^1}$ is holomorphic symplectic. In fact, one can easily construct a surjective cosection

$$\sigma : -\mathbf{R}\pi_* f^* K_{\mathbb{P}^1} \rightarrow \mathcal{O}_{\overline{M}_{0,1}(Y, \beta)}.$$

And $c_{2n-2}(-\mathbf{R}\pi_* f^* K_{\mathbb{P}^1}) = e(\text{Ker}(\sigma))$ is the Euler class of the reduced obstruction bundle [12].

So for $d_1 \otimes \gamma_2 \in H^2(\mathbb{P}^1, \mathbb{Z}) \otimes H^2(S, \mathbb{Z})$, we have

$$\begin{aligned} \text{GW}_{0, \beta}^{\text{twist}}(Y)(d_1 \otimes \gamma_2) &= d_1 \int_{[\overline{M}_{0,1}(\mathbb{P}^1, n)]^{\text{vir}}} c_{2n-2}(-\mathbf{R}\pi_* f^* K_{\mathbb{P}^1}) \cup \text{ev}^*([\text{pt}]) \cdot \int_S c_1(K_S) \cup \gamma_2 \\ &= d_1 \int_{[\overline{M}_{0,1}(K_{\mathbb{P}^1}, n)]_{\text{red}}^{\text{vir}}} \text{ev}^*([\text{pt}]) \cdot \int_S c_1(K_S) \cup \gamma_2 \\ &= d_1 \cdot n \cdot \deg([\overline{M}_{0,0}(K_{\mathbb{P}^1}, n)]_{\text{red}}^{\text{vir}}) \cdot \int_S c_1(K_S) \cup \gamma_2 \\ &= d_1 \cdot \frac{1}{n^2} \cdot \int_S c_1(K_S) \cup \gamma_2 \\ &= -d_1 \cdot \frac{1}{n^2} \cdot \int_S e\left(\bigwedge^2 TS\right) \cup \gamma_2. \end{aligned}$$

As for $\gamma \in H^4(S, \mathbb{Z})$, we obviously have vanishing $\text{GW}_{0, \beta}^{\text{twist}}(Y)(\gamma) = 0$. Thus

$$\text{GW}_{0, n[\mathbb{P}^1]}^{\text{twist}}(Y)(\gamma) = (1/n^2) \cdot \text{DT}_3^{\text{twist}}(Y)([\mathbb{P}^1] \mid \gamma)$$

holds for any $n \geq 1$ and $\gamma \in H^4(Y, \mathbb{Z})$ as $(c_1(Y) \cdot \beta - 1) = 2n + 1$ is always odd. \square

Remark 2.4. We can also consider a \mathbb{P}^1 -bundle $\pi : Y \rightarrow S$ over a del-Pezzo surface S with a section. For multiple fiber classes, similar result as Proposition 2.3 holds in this setting.

2.3. Product of del-Pezzo surface with \mathbb{P}^1 . We consider $Y = S \times \mathbb{P}^1$ for a del-Pezzo surface S with $\beta \in H_2(S, \mathbb{Z}) \subseteq H_2(Y, \mathbb{Z})$.

Proposition 2.5. *Let S be a del-Pezzo surface and $Y = S \times \mathbb{P}^1$ be the product. For $\beta \in H_2(S, \mathbb{Z}) \subseteq H_2(Y, \mathbb{Z})$, we have:*

(1) *If $\gamma = (\gamma_1, d) \in H^2(S) \otimes H^2(\mathbb{P}^1) \subseteq H^4(Y)$, then*

$$\begin{aligned} \mathrm{DT}_3^{\mathrm{twist}}(Y)(\beta \mid \gamma) &= d(\beta \cdot \gamma_1) \cdot \mathrm{DT}_3(K_S)(\beta), \\ \mathrm{GW}_{0,\beta}^{\mathrm{twist}}(Y)(\gamma) &= d(\beta \cdot \gamma_1) \cdot \mathrm{GW}_{0,\beta}(K_S). \end{aligned}$$

(2) *If $\gamma \in H^4(S) \subseteq H^4(Y)$, then*

$$\mathrm{DT}_3^{\mathrm{twist}}(Y)(\beta \mid \gamma) = \mathrm{GW}_{0,\beta}^{\mathrm{twist}}(Y)(\gamma) = 0.$$

Proof. Similar to Proposition 2.3, any $E \in M_{Y,\beta}$ is of type $E = (\iota_p)_*F$ for some one dimensional stable sheaf F on S , where $\iota_p : S = S \times \{p\} \rightarrow Y$ is the natural inclusion. Then we have an isomorphism

$$M_{Y,\beta} \cong M_{S,\beta} \times \mathbb{P}^1,$$

where $M_{S,\beta}$ is the moduli space of one dimensional stable sheaves F 's on S with $[F] = \beta$ and $\chi(F) = 1$. For $E = (\iota_p)_*F$, we have isomorphisms

$$\begin{aligned} \mathrm{Ext}_Y^1(E, E) &\cong \mathrm{Ext}_S^1(F, F) \oplus \mathbb{C}, \\ \mathrm{Ext}_Y^2(E, E) &\cong \mathrm{Ext}_S^1(F, F), \end{aligned}$$

as $\mathrm{Ext}_S^2(F, F) \cong \mathrm{Hom}_S(F, F \otimes K_S)^* = 0$ by the stability of F . Then

$$[M_{Y,\beta}]^{\mathrm{vir}} = \chi(M_{S,\beta}) \cdot [\mathbb{P}^1] \in H_2(M_{Y,\beta}, \mathbb{Z}).$$

The universal sheaf \mathcal{E}_{M_Y} over $M_{Y,\beta} \times Y$ can be identified with

$$\mathcal{E}_{M_Y} \cong \mathcal{E}_{M_S} \boxtimes \mathcal{O}_{\Delta_{\mathbb{P}^1}} \in \mathrm{Coh}(M_S \times S \times \mathbb{P}^1 \times \mathbb{P}^1),$$

where \mathcal{E}_{M_S} is the universal sheaf over $M_{S,\beta} \times S$.

Then for $\gamma \in H^4(S) \subseteq H^4(Y)$, we have

$$\mathrm{DT}_3^{\mathrm{twist}}(Y)(\beta \mid \gamma) = 0,$$

and for $\gamma = (\gamma_1, d) \in H^2(S) \otimes H^2(\mathbb{P}^1) \subseteq H^4(Y)$, we have

$$\begin{aligned} \mathrm{DT}_3^{\mathrm{twist}}(Y)(\beta \mid \gamma) &= (-1)^{c_1(S) \cdot \beta - 1} \cdot \int_{[M_{Y,\beta}]^{\mathrm{vir}}} \tau(\gamma) \\ &= (-1)^{c_1(S) \cdot \beta - 1} d \cdot \chi(M_{S,\beta}) \cdot (\pi_{M_S})_*([\mathcal{E}_{M_S}] \cup \pi_S^* \gamma_1) \\ &= (-1)^{c_1(S) \cdot \beta - 1} (\beta \cdot \gamma_1) \cdot d \cdot \chi(M_{S,\beta}), \end{aligned}$$

where π_S, π_{M_S} are projections from $M_{S,\beta} \times S$ to corresponding factors.

Meanwhile, the DT_3 invariant for the moduli space $M_{K_S,\beta}$ of one dimensional stable sheaves E 's on K_S with $[E] = \beta$ and $\chi(E) = 1$ satisfies

$$\mathrm{DT}_3(K_S)(\beta) = (-1)^{(1+\beta^2)} \cdot \chi(M_{S,\beta}),$$

where $(1 + \beta^2)$ is the dimension of the smooth moduli space $M_{S,\beta}$ (see e.g. [5, Lemma 3.6]).

We claim the parity of β^2 and $c_1(S) \cdot \beta$ are the same, i.e.

$$(2.4) \quad (-1)^{(1+\beta^2)} = (-1)^{(c_1(S) \cdot \beta - 1)}.$$

In fact, we may assume S to be a general del-Pezzo surface and given by the blow-up of r general points in \mathbb{P}^2 (the case of $\mathbb{P}^1 \times \mathbb{P}^1$ is trivial). Let H be the pull-back of the hyperplane class and $\{E_i\}_{i=1}^r$ are exceptional divisors. Then

$$\begin{aligned} c_1(S) &= 3H - \sum_{i=1}^r E_i, \quad \beta := dH + \sum_{i=1}^r d_i E_i, \text{ for some } d, d_i \in \mathbb{Z}, \\ c_1(S) \cdot \beta &= 3d + \sum_{i=1}^r d_i, \quad \beta^2 = d^2 - \sum_{i=1}^r d_i. \end{aligned}$$

Hence for $\gamma = (\gamma_1, d)$,

$$\mathrm{DT}_3^{\mathrm{twist}}(Y)(\beta \mid \gamma) = (\beta \cdot \gamma_1) \cdot d \cdot \mathrm{DT}_3(K_S)(\beta).$$

As for the corresponding GW theory, there is an isomorphism

$$\overline{M}_{0,n}(Y, \beta) \cong \overline{M}_{0,n}(S, \beta) \times \mathbb{P}^1, \quad n \geq 0.$$

And for $t : \mathbb{P}^1 \rightarrow Y = S \times \mathbb{P}^1$, there are isomorphisms

$$\begin{aligned} H^*(\mathbb{P}^1, t^*TY) &\cong H^*(\mathbb{P}^1, t^*TS) \oplus H^*(\mathbb{P}^1, c^*T\mathbb{P}^1), \\ H^*(\mathbb{P}^1, t^*K_Y) &\cong H^*(\mathbb{P}^1, t^*K_S \otimes c^*K_{\mathbb{P}^1}), \end{aligned}$$

where c is a constant map to some point of \mathbb{P}^1 .

The virtual class and the extra obstruction bundle satisfies

$$[\overline{M}_{0,1}(Y, \beta)]^{\text{vir}} = [\overline{M}_{0,1}(S, \beta)]^{\text{vir}} \otimes [\mathbb{P}^1].$$

For the universal curve $\pi : \mathcal{C} \rightarrow \overline{M}_{0,1}(Y, \beta)$ and map $f : \mathcal{C} \rightarrow Y$, by (2.3), we have

$$\mathbf{R}\pi_* f^* K_Y \cong \phi^* \mathbf{R}\phi_* \text{ev}^* K_Y,$$

where $\phi : \overline{M}_{0,1}(Y, \beta) \rightarrow \overline{M}_{0,0}(Y, \beta)$ is the forgetful map and $\text{ev} : \overline{M}_{0,1}(Y, \beta) \rightarrow Y$ is the evaluation map. Notice that

$$\begin{aligned} \phi &= (\phi_S, \text{id}) : \overline{M}_{0,1}(S, \beta) \times \mathbb{P}^1 \rightarrow \overline{M}_{0,0}(S, \beta) \times \mathbb{P}^1, \\ \text{ev} &= (\text{ev}_S, \text{id}) : \overline{M}_{0,1}(S, \beta) \times \mathbb{P}^1 \rightarrow S \times \mathbb{P}^1. \end{aligned}$$

Hence

$$\begin{aligned} \text{GW}_{0,\beta}^{\text{twist}}(Y)(\gamma) &= \int_{[\overline{M}_{0,1}(Y,\beta)]^{\text{vir}}} e(-\mathbf{R}\pi_* f^* K_Y) \cup \text{ev}^*(\gamma) \\ &= \int_{[\overline{M}_{0,1}(Y,\beta)]^{\text{vir}}} \phi^* e(\mathbf{R}^1 \phi_* \text{ev}^* K_Y) \cup \text{ev}^*(\gamma) \\ &= \int_{[\overline{M}_{0,1}(S,\beta)]^{\text{vir}} \otimes [\mathbb{P}^1]} e(\phi_S^*(\mathbf{R}^1(\phi_S)_* \text{ev}_S^* K_S) \boxtimes K_{\mathbb{P}^1}) \cup \text{ev}^*(\gamma). \end{aligned}$$

If $\gamma \in H^4(S) \subseteq H^4(Y)$, the above integration is zero by degree reason.

If $\gamma = (\gamma_1, d) \in H^2(S) \otimes H^2(\mathbb{P}^1) \subseteq H^4(Y)$, we have

$$\begin{aligned} \text{GW}_{0,\beta}^{\text{twist}}(Y)(\gamma) &= \int_{[\overline{M}_{0,1}(S,\beta)]^{\text{vir}} \otimes [\mathbb{P}^1]} e(\phi_S^*(\mathbf{R}^1(\phi_S)_* \text{ev}_S^* K_S) \boxtimes K_{\mathbb{P}^1}) \cup \text{ev}^*(\gamma) \\ &= \int_{[\overline{M}_{0,1}(S,\beta)]^{\text{vir}}} e(\phi_S^*(\mathbf{R}^1(\phi_S)_* \text{ev}_S^* K_S)) \cup \text{ev}_S^*(\gamma_1) \cdot \int_{\mathbb{P}^1} d [\text{pt}] \\ &= d(\beta \cdot \gamma_1) \int_{[\overline{M}_{0,0}(S,\beta)]^{\text{vir}}} e(\mathbf{R}^1(\phi_S)_* \text{ev}_S^* K_S) \\ &= d(\beta \cdot \gamma_1) \cdot \text{GW}_{0,\beta}(K_S), \end{aligned}$$

where we use the divisor equation of GW invariants in the second to last equality. \square

Thus twisted GW/DT₃ invariants of $S \times \mathbb{P}^1$ reduce to GW/DT₃ invariants of (non-compact) Calabi-Yau 3-fold K_S . We recall the following Katz's conjecture on Calabi-Yau 3-folds.

Conjecture 2.6. (Katz [11]) *Let Y be a (compact) Calabi-Yau 3-fold and $\beta \in H_2(Y, \mathbb{Z})$. Then*

$$\text{GW}_{0,\beta}(Y) = \sum_{k|\beta} \frac{1}{k^3} \text{DT}_3(Y)(\beta/k).$$

Although K_S is non-compact, the moduli spaces used to define GW/DT₃ invariants are compact, so Katz's conjecture still makes sense on K_S . By Proposition 2.5, it is easy to show:

Corollary 2.7. *In the setting of Proposition 2.5, Conjecture 1.2 is true for $Y = S \times \mathbb{P}^1$ with $\beta \in H_2(S, \mathbb{Z}) \subseteq H_2(Y, \mathbb{Z})$ if and only if Katz's conjecture holds for K_S with $\beta \in H_2(S, \mathbb{Z})$.*

Combining with the previous checks of Katz's conjecture, we obtain:

Theorem 2.8. *Let S be a toric del-Pezzo surface and $Y = S \times \mathbb{P}^1$ be the product. Then Conjecture 1.2 is true for $\beta \in H_2(S, \mathbb{Z}) \subseteq H_2(Y, \mathbb{Z})$.*

Proof. By Corollary 2.7, we are reduced to prove the Katz's conjecture for K_S . This is done by combining DT/PT/GW correspondence and geometric vanishing in [5, Corollary A.7]. \square

Remark 2.9. *When $Y = \mathcal{O}_{\mathbb{P}^1}(-1) \times \mathbb{P}^1$, Conjecture 1.2 is true and can be reduced to the Aspinwall-Morrison multiple cover formula*

$$\text{GW}_{0,d}(X) = \sum_{d|n} \frac{1}{d^3},$$

$$\text{DT}_3(X)(d) = 1, \text{ if } d = 1; \quad \text{DT}_3(X)(d) = 0, \text{ if } d > 1,$$

for $X = \mathcal{O}_{\mathbb{P}^1}(-1, -1)$.

3. NON-COMPACT EXAMPLES

3.1. Local surfaces. Let $(S, \mathcal{O}_S(1))$ be a smooth projective surface and

$$(3.1) \quad \pi: Y = \text{Tot}_S(L) \rightarrow S$$

be the total space of a line bundle L on S . When $L \otimes K_S^{-1}$ is ample, Y is a non-compact Fano 3-fold. We take a curve class

$$\beta \in H_2(Y, \mathbb{Z}) \cong H_2(S, \mathbb{Z}),$$

and consider the moduli scheme $M_{Y, \beta}$ (resp. $M_{S, \beta}$) of one dimensional stable sheaves F on Y (resp. on S) with $[F] = \beta$ and $\chi(F) = 1$. Note that $M_{S, \beta}$ is compact while $M_{Y, \beta}$ may not be compact. On the other hand, for the zero section $\iota: S \hookrightarrow Y$ of the projection (3.1), we have the push-forward embedding

$$(3.2) \quad \iota_*: M_{S, \beta} \hookrightarrow M_{Y, \beta}.$$

In the following setting, the morphism (3.2) is an isomorphism and $M_{Y, \beta}$ has a well-defined virtual class.

Lemma 3.1. ([5, Proposition 3.1]) *If S is a del-Pezzo surface and L^{-1} is ample, then (3.2) is an isomorphism, under which we have*

$$[M_{Y, \beta}]^{\text{vir}} = [M_{S, \beta}] \cdot e\left(\mathcal{E}xt_{\pi_{M_S}}^1(\mathbb{F}, \mathbb{F} \boxtimes L)\right),$$

where $\mathbb{F} \in \text{Coh}(S \times M_{S, \beta})$ is the universal sheaf and $\pi_{M_S}: S \times M_{S, \beta} \rightarrow M_{S, \beta}$ is the projection.

Proof. From the proof of [5, Proposition 3.1], for any $F \in M_{Y, \beta}$, there exists $\mathcal{E} \in M_{S, \beta}$ such that $F = \iota_*(\mathcal{E})$. To compare the deformation-obstruction theory, we have canonical isomorphisms

$$\begin{aligned} \text{Ext}_Y^1(\iota_*\mathcal{E}, \iota_*\mathcal{E}) &\cong \text{Ext}_S^1(\mathcal{E}, \mathcal{E}), \\ \text{Ext}_Y^2(\iota_*\mathcal{E}, \iota_*\mathcal{E}) &\cong \text{Ext}_S^1(\mathcal{E}, \mathcal{E} \otimes L), \end{aligned}$$

where $\text{Ext}_S^2(\mathcal{E}, \mathcal{E}) \cong \text{Ext}_S^0(\mathcal{E}, \mathcal{E} \otimes K_S)^* = 0$ and $\text{Ext}_S^0(\mathcal{E}, \mathcal{E} \otimes L) = 0$ by the stability of \mathcal{E} . \square

Note that twisted DT₃ invariants of $Y = \text{Tot}_S(L)$ are invariant under the transformation $L \mapsto L^{-1} \otimes K_S$.

Proposition 3.2. *Let S be a del-Pezzo surface and $Y_1 = \text{Tot}_S(L)$, $Y_2 = \text{Tot}_S(L^{-1} \otimes K_S)$ be the total space of ample line bundles L^{-1} , $L \otimes K_S^{-1}$ respectively on S . Then*

$$[M_{Y_1, \beta}]^{\text{vir}} = (-1)^{\beta^2} \cdot [M_{Y_2, \beta}]^{\text{vir}} \in H_2(M_{S, \beta}, \mathbb{Z}).$$

In particular, we have

$$\text{DT}_3^{\text{twist}}(Y_1)(\beta \mid [\text{pt}]) = \text{DT}_3^{\text{twist}}(Y_2)(\beta \mid [\text{pt}]).$$

Proof. By Lemma 3.1, we are left to determine the relation between $e(\mathcal{E}xt_{\pi_{M_S}}^1(\mathbb{F}, \mathbb{F} \boxtimes L))$ and $e(\mathcal{E}xt_{\pi_{M_S}}^1(\mathbb{F}, \mathbb{F} \boxtimes L^{-1} \otimes K_S))$. By the Grothendieck-Verdier duality,

$$\mathcal{E}xt_{\pi_{M_S}}^1(\mathbb{F}, \mathbb{F} \boxtimes L) \cong \mathcal{E}xt_{\pi_{M_S}}^1(\mathbb{F}, \mathbb{F} \boxtimes L^{-1} \otimes K_S)^*.$$

Hence,

$$e(\mathcal{E}xt_{\pi_{M_S}}^1(\mathbb{F}, \mathbb{F} \boxtimes L)) = (-1)^{\text{ext}^1(F, F \otimes L)} e(\mathcal{E}xt_{\pi_{M_S}}^1(\mathbb{F}, \mathbb{F} \boxtimes L^{-1} \otimes K_S)),$$

where $\text{ext}^1(F, F \otimes L) = \beta^2$ can be computed by the Riemann-Roch formula. Then

$$\begin{aligned} (-1)^{c_1(Y_1) \cdot \beta - 1} &= (-1)^{(c_1(S) + c_1(L)) \cdot \beta - 1} \\ &= (-1)^{c_1(L) \cdot \beta + \beta^2 - 1} \\ &= (-1)^{c_1(Y_2) \cdot \beta - 1} \cdot (-1)^{\beta^2}, \end{aligned}$$

where we use (2.4) in the second equality. \square

One can easily check that twisted GW invariants of Y_1 and Y_2 are also the same. Hence

Corollary 3.3. *Let S be a del-Pezzo surface and $Y_1 = \text{Tot}_S(L)$, $Y_2 = \text{Tot}_S(L^{-1} \otimes K_S)$ be the total space of ample line bundles L^{-1} , $L \otimes K_S^{-1}$ respectively on S . Then Conjecture 1.2 is true for Y_1 if and only if it is true for Y_2 .*

Combining with [5, Sect. 3.2], we then verify our conjecture for the following:

Proposition 3.4. *Let $Y = \mathcal{O}_{\mathbb{P}^2}(-1)$ or $\mathcal{O}_{\mathbb{P}^2}(-2)$ and $\beta = d[l] \in H_2(\mathbb{P}^2, \mathbb{Z})$ be the degree d class. Then Conjecture 1.2 is true when $d \leq 3$.*

3.2. Local curves. In [5, Section 4], we study local curves of form

$$X = \text{Tot}_{\mathbb{P}^1}(\mathcal{O}_{\mathbb{P}^1}(l_1) \oplus \mathcal{O}_{\mathbb{P}^1}(l_2) \oplus \mathcal{O}_{\mathbb{P}^1}(l_3)) \rightarrow \mathbb{P}^1,$$

where $l_1 + l_2 + l_3 = -2$ so that it is a non-compact CY 4-fold. Without loss of generality, we may assume $l_1 \geq l_2 \geq l_3$, so $l_3 < 0$. We can regard X as the total space of canonical bundle of

$$(3.3) \quad Y = \text{Tot}_{\mathbb{P}^1}(\mathcal{O}_{\mathbb{P}^1}(l_1) \oplus \mathcal{O}_{\mathbb{P}^1}(l_2)) \quad \text{with } l_1 + l_2 \geq -1.$$

In this section, we study an equivariant version of Conjecture 1.2 for non-compact Fano 3-fold Y as [5, Conjecture 4.10] for non-compact CY 4-fold X .

3.2.1. Equivariant GW invariants. As a toric variety, Y contains a torus $(\mathbb{C}^*)^3$ and has two invariant affine open subsets with transition map

$$(z_0, z_1, z_2) \mapsto (z_0^{-1}, z_0^{-l_1} z_1, z_0^{-l_2} z_2).$$

The action of $(\mathbb{C}^*)^3$ is given by

$$t \cdot (z_0, z_1, z_2) \mapsto (t_0 z_0, t_1 z_1, t_2 z_2).$$

Let \bullet denote $\text{Spec } \mathbb{C}$ with trivial $(\mathbb{C}^*)^3$ -action. Let $\mathbb{C} \otimes t_i$ be the one dimensional vector space with $(\mathbb{C}^*)^3$ -action with weight t_i , and $\lambda_i \in H_{(\mathbb{C}^*)^3}^*(\bullet)$ its 1st Chern class.

Let $T := \{(t_1, t_2) \mid (t_0, t_1, t_2) \in (\mathbb{C}^*)^3\}$ be the two dimensional subtorus which acts trivially on the base of Y . Note that

$$H_{(\mathbb{C}^*)^3}^*(\bullet) = \mathbb{C}[\lambda_0, \lambda_1, \lambda_2], \quad H_T^*(\bullet) = \mathbb{C}[\lambda_1, \lambda_2].$$

Let $j: \mathbb{P}^1 \hookrightarrow Y$ be the zero section. Note that we have $H_2(Y, \mathbb{Z}) = \mathbb{Z}[\mathbb{P}^1]$, where $[\mathbb{P}^1]$ is the fundamental class of $j(\mathbb{P}^1)$. For $d \in \mathbb{Z}_{>0}$, we consider the diagram

$$\begin{array}{ccc} \mathcal{C} & \xrightarrow{f} & \mathbb{P}^1 \\ \downarrow p & & \\ \overline{M}_{0,0}(\mathbb{P}^1, d), & & \end{array}$$

where \mathcal{C} is the universal curve and f is the universal stable map.

The T -equivariant twisted GW invariant of Y is defined by

$$\text{GW}_{0,d}^{\text{twist}}(Y) := \int_{[\overline{M}_{0,0}(\mathbb{P}^1, d)]} e_T(-\mathbf{R}p_* f^* N) \in \mathbb{Q}(\lambda_1, \lambda_2),$$

where

$$(3.4) \quad N := (\mathcal{O}_{\mathbb{P}^1}(l_1) \otimes t_1) \oplus (\mathcal{O}_{\mathbb{P}^1}(l_2) \otimes t_2) \oplus (\mathcal{O}_{\mathbb{P}^1}(-2 - l_1 - l_2) \otimes t_1^{-1} t_2^{-1})$$

is the T -equivariant normal bundle of $j(\mathbb{P}^1) \subset X$.

Remark 3.5. One can furthermore use the \mathbb{C}^* -action on the base \mathbb{P}^1 to do calculations.

For example in the $d = 1$ case, $\overline{M}_{0,0}(\mathbb{P}^1, 1)$ is one point and

$$(3.5) \quad \text{GW}_{0,1}^{\text{twist}}(Y) = \lambda_1^{-l_1-1} \lambda_2^{-l_2-1} (-\lambda_1 - \lambda_2)^{l_1+l_2+1}.$$

The $d = 2$ case is more complicated and can be computed by Kontsevich's localization method

$$(3.6) \quad \begin{aligned} \text{GW}_{0,2}^{\text{twist}}(Y) = & \frac{1}{8} \lambda_1^{-2l_1-1} \lambda_2^{-2l_2-1} (-\lambda_1 - \lambda_2)^{-2l_3-1} \left\{ (\bar{l}_1^2 - (\bar{l}_1 - 1)^2 + \dots) \lambda_1^{-2} + \right. \\ & (\bar{l}_2^2 - (\bar{l}_2 - 1)^2 + \dots) \lambda_2^{-2} + (\bar{l}_3^2 - (\bar{l}_3 - 1)^2 + \dots) (\lambda_1 + \lambda_2)^{-2} \\ & \left. + l_1 l_2 \lambda_1^{-1} \lambda_2^{-1} + l_2 l_3 \lambda_2^{-1} (-\lambda_1 - \lambda_2)^{-1} + l_1 l_3 \lambda_1^{-1} (-\lambda_1 - \lambda_2)^{-1} \right\}, \end{aligned}$$

where $l_3 := -l_1 - l_2 - 2$ and we write $\bar{l} = l$ for $l \geq 0$ and $\bar{l} = -l - 1$ for $l < 0$ (see e.g. [5, (4.6)]).

3.2.2. Equivariant DT₃ invariants. Let $M_{Y,d}$ (resp. $M_{X,d}$) be the moduli scheme of one dimensional stable sheaves F 's on $Y = \text{Tot}_{\mathbb{P}^1}(\mathcal{O}_{\mathbb{P}^1}(l_1) \oplus \mathcal{O}_{\mathbb{P}^1}(l_2))$ with $l_1 + l_2 \geq -1$ (resp. on $X = K_Y$) such that $[F] = d[\mathbb{P}^1] \in \mathbb{Z}[\mathbb{P}^1]$ and $\chi(F) = 1$. Note that $M_{Y,d}$ is non-compact and the virtual class is not well-defined, but the induced T -action on $M_{Y,d}$ has compact fixed locus.

Lemma 3.6.

(1) The zero section $\iota: Y \hookrightarrow X$ induces an isomorphism

$$\iota_*: M_{Y,d} \cong M_{X,d}.$$

(2) The T -fixed locus $M_{Y,d}^T$ is compact.

Proof. (1) We claim that any $[F] \in M_{X,d}$ is scheme theoretically supported on Y . In fact, we have an exact sequence

$$F \otimes \pi^* \mathcal{O}_{\mathbb{P}^1}(2 + l_1 + l_2) \rightarrow F \rightarrow F \otimes \mathcal{O}_Y \rightarrow 0,$$

where $\pi : X \rightarrow \mathbb{P}^1$ is the projection. By the stability of F and $2 + l_1 + l_2 \geq 1$, we know the first map is zero. Hence $F \cong F \otimes \mathcal{O}_Y$ and $l_* : M_{Y,d} \rightarrow M_{X,d}$ is an isomorphism.

(2) Let $\bar{Y} := \mathbb{P}(\mathcal{O}_{\mathbb{P}^1}(l_1) \oplus \mathcal{O}_{\mathbb{P}^1}(l_2) \oplus \mathcal{O}_{\mathbb{P}^1})$ be the projectification of Y with an extended T -action on the fiber. The inclusion $i : Y \rightarrow \bar{Y}$ induces a map

$$i_* : M_{Y,d} \rightarrow M_{\bar{Y},d},$$

to the moduli scheme $M_{\bar{Y},d}$ of one dimensional stable sheaves on \bar{Y} such that

$$M_{Y,d}^T \subseteq M_{\bar{Y},d}^T$$

is a closed and open subset. Note that $M_{\bar{Y},d}$ is compact since \bar{Y} is compact, then the T -fixed locus $M_{Y,d}^T$ is also compact. \square

Then we define equivariant twisted DT₃ invariant of $M_{Y,d}$ by virtual localization formula [9].

Definition 3.7. *In the above setting, the equivariant twisted DT₃ invariant of $M_{Y,d}$ is*

$$\text{DT}_3^{\text{twist}}(Y)(d) := (-1)^{d(l_1+l_2)-1} \int_{[M_{Y,d}^T]^{\text{vir}}} e_T(N^{\text{vir}}) \in \mathbb{Q}(\lambda_1, \lambda_2),$$

where N^{vir} is the virtual normal bundle of $M_{Y,d}^T \hookrightarrow M_{Y,d}$.

For example in the $d = 1$ case, $M_{Y,1}^T$ is one point and

$$\text{DT}_3^{\text{twist}}(Y)(1) = (-1)^{l_1+l_2-1} \lambda_1^{-l_1-1} \lambda_2^{-l_2-1} (\lambda_1 + \lambda_2)^{l_1+l_2+1},$$

which matches with $\text{GW}_{0,1}^{\text{twist}}(Y)$.

3.2.3. The equivariant conjecture. An equivariant version of Conjecture 1.2 for non-compact Fano 3-fold $Y = \text{Tot}_{\mathbb{P}^1}(\mathcal{O}_{\mathbb{P}^1}(l_1) \oplus \mathcal{O}_{\mathbb{P}^1}(l_2))$ can be stated in the following form, where the multiple cover coefficient is $1/k^3$ instead of $1/k^2$ due to the absence of insertions.

Conjecture 3.8. *Let $Y = \text{Tot}_{\mathbb{P}^1}(\mathcal{O}_{\mathbb{P}^1}(l_1) \oplus \mathcal{O}_{\mathbb{P}^1}(l_2))$ with $l_1 + l_2 \geq -1$. Then we have*

$$\text{GW}_{0,d}^{\text{twist}}(Y) = \sum_{k|d} \frac{1}{k^3} \text{DT}_3^{\text{twist}}(Y)(d/k).$$

In [5, (4.9), (4.17)], equivariant DT₄ invariants for $X = K_Y$ are defined for degree one and two classes by choosing suitable square root of the virtual normal bundle. Below we show that Definition 3.7 recovers them and can be viewed as a general definition of equivariant DT₄ invariants for all degree curves classes on such X .

Theorem 3.9. *Definition 3.7 for $Y = \text{Tot}_{\mathbb{P}^1}(\mathcal{O}_{\mathbb{P}^1}(l_1) \oplus \mathcal{O}_{\mathbb{P}^1}(l_2))$ recovers the definition of equivariant DT₄ invariants for $X = K_Y$ in (4.9), (4.17) of [5], i.e. for $d = 1, 2$,*

$$\text{DT}_3^{\text{twist}}(Y)(d) = \text{DT}_4(X)(d[\mathbb{P}^1]).$$

In particular, Conjecture 3.8 is true when (i) $d = 1$ and (ii) $d = 2$, $|l_1|, |l_2| \leq 10$.

Proof. When $d = 1$, it is already computed explicitly as above. When $d = 2$, say we take a T -fixed $\mathcal{O}_{\mathbb{P}^1}(l_1)$ -thickened sheaf $[F] \in M_{Y,2}^T$, then $F = F_0 + F_1 \cdot t_1^{-1}$ in the T -equivariant K -theory of Y , with F_0, F_1 are line bundles on \mathbb{P}^1 with weight 0, t_1^{-1} . Therefore,

$$\begin{aligned} \chi_Y(F, F) &= \chi_Y(j_* F_0, j_* F_0) + \chi_Y(j_* F_0, j_* F_1) \otimes t_i^{-1} + \chi_Y(j_* F_1, j_* F_0) \otimes t_i + \chi_Y(j_* F_1, j_* F_1) \\ &= 2(\chi(\mathcal{O}_{\mathbb{P}^1}) - \chi(N_{\mathbb{P}^1/Y}) + \chi(\wedge^2 N_{\mathbb{P}^1/Y})) + (\chi(A) - \chi(A \otimes N_{\mathbb{P}^1/Y}) + \chi(A \otimes \wedge^2 N_{\mathbb{P}^1/Y})) \otimes t_i^{-1} \\ &\quad + (\chi(A^{-1}) - \chi(A^{-1} \otimes N_{\mathbb{P}^1/Y}) + \chi(A^{-1} \otimes \wedge^2 N_{\mathbb{P}^1/Y})) \otimes t_i, \end{aligned}$$

where $N_{\mathbb{P}^1/Y} = \mathcal{O}(l_1) \otimes t_1 \oplus \mathcal{O}(l_2) \otimes t_2$ and $A := F_i \otimes F_0^\vee$. The movable part is

$$\begin{aligned} \chi_Y(F, F)^{\text{mov}} &= -2\chi(\mathcal{O}(l_1)) \otimes t_1 - 2\chi(\mathcal{O}(l_2)) \otimes t_2 + 2\chi(\mathcal{O}(l_1 + l_2)) \otimes t_1 t_2 \\ &\quad + \chi(A) \otimes t_1^{-1} - \chi(A \otimes \mathcal{O}(l_2)) \otimes t_1^{-1} t_2 + \chi(A \otimes \mathcal{O}(l_1 + l_2)) \otimes t_2 \\ &\quad + \chi(A^{-1}) \otimes t_1 - \chi(A^{-1} \otimes \mathcal{O}(l_1)) \otimes t_1^2 - \chi(A^{-1} \otimes \mathcal{O}(l_2)) \otimes t_1 t_2 \\ &\quad + \chi(A^{-1} \otimes \mathcal{O}(l_1 + l_2)) \otimes t_1^2 t_2. \end{aligned}$$

In [5, Sect. 4.4], we view F as a sheaf in $X = K_Y$ by push-forward via zero section and a square root of $\chi_X(F, F)$ is chosen as follows

$$\begin{aligned}\chi_X(F, F)^{1/2} &= 2\chi(\mathcal{O}) - \chi(\mathcal{O}(l_1)) \otimes t_1 - \chi(\mathcal{O}(l_2)) \otimes t_2 - \chi(\mathcal{O}(l_3)) \otimes t_3 \\ &\quad + \chi(\mathcal{O}(l_1 + l_2)) \otimes t_1 t_2 + \chi(\mathcal{O}(l_1 + l_3)) \otimes t_1 t_3 + \chi(\mathcal{O}(l_2 + l_3)) \otimes t_2 t_3 \\ &\quad + \chi(A) \otimes t_1^{-1} - \chi(A \otimes \mathcal{O}(l_1)) - \chi(A \otimes \mathcal{O}(l_2)) \otimes t_1^{-1} t_2 - \chi(A \otimes \mathcal{O}(l_3)) \otimes t_1^{-1} t_3 \\ &\quad + \chi(A \otimes \mathcal{O}(l_1 + l_2)) \otimes t_2 + \chi(A \otimes \mathcal{O}(l_1 + l_3)) \otimes t_3 + \chi(A \otimes \mathcal{O}(l_2 + l_3)) \otimes t_1^{-1} t_2 t_3 \\ &\quad - \chi(A \otimes \mathcal{O}(-2)) \otimes t_1^{-1},\end{aligned}$$

where $t_3 := t_1^{-1} t_2^{-1}$ and $l_3 := -l_1 - l_2 - 2$.

By Serre duality, terms in $\chi_X(F, F)^{1/2, \text{mov}}$ and $\chi_Y(F, F)^{\text{mov}}$ are in 1-1 correspondence and their equivariant Euler classes (after putting $\lambda_3 = -\lambda_1 - \lambda_2$ in $\chi_X(F, F)^{1/2, \text{mov}}$) differ by a sign. In fact, the difference appears whenever we use Serre duality, for instance, Serre duality gives

$$\chi(\mathcal{O}(l_3)) = -\chi(\mathcal{O}(l_1 + l_2)),$$

whose equivariant Euler class satisfies

$$e_T(\chi(\mathcal{O}(l_3)) \otimes t_3) = \lambda_3^{l_3+1}, \quad e_T(-\chi(\mathcal{O}(l_1 + l_2)) \otimes t_1 t_2) = (\lambda_1 + \lambda_2)^{l_3+1},$$

which differs by a sign $(-1)^{l_3+1}$ after taking $\lambda_3 = -\lambda_1 - \lambda_2$.

The product of all sign differences is $(-1)^\Delta$ with

$$\begin{aligned}\Delta &= \chi(\mathcal{O}(l_3)) + \chi(\mathcal{O}(l_1 + l_3)) + \chi(\mathcal{O}(l_2 + l_3)) \\ &\quad - \chi(A \otimes \mathcal{O}(l_3)) + \chi(A \otimes \mathcal{O}(l_1 + l_3)) + \chi(A \otimes \mathcal{O}(l_2 + l_3)) - \chi(A \otimes \mathcal{O}(-2)) \\ &= 2l_3 + 2(l_1 + l_2 + l_3) + 5.\end{aligned}$$

So $(-1)^\Delta = (-1) = (-1)^{2(l_1+l_2)+1}$, i.e. the sign in Definition 3.7.

Finally by [5, Thm. 4.12], we know Conjecture 3.8 is true for $d = 1$ and $d = 2$ when $|l_1|, |l_2| \leq 10$. \square

Acknowledgement. The author is grateful to the referee for carefully reading the manuscript and giving helpful comments which improve the exposition of the paper. This work is done while the author was in Oxford supported by The Royal Society Newton International Fellowship. The author is currently supported by the World Premier International Research Center Initiative (WPI), MEXT, Japan.

REFERENCES

- [1] D. Borisov and D. Joyce, *Virtual fundamental classes for moduli spaces of sheaves on Calabi-Yau four-folds*, Geom. Topol. (21), (2017) 3231-3311.
- [2] Y. Cao, J. Gross and D. Joyce, *Orientability of moduli spaces of Spin(7)-instantons and coherent sheaves on Calabi-Yau 4-folds*, arXiv:1811.09658, v2.
- [3] Y. Cao and N. C. Leung, *Donaldson-Thomas theory for Calabi-Yau 4-folds*, arXiv:1407.7659.
- [4] Y. Cao and N. C. Leung, *Orientability for gauge theories on Calabi-Yau manifolds*, Adv. in Math. (314), 2017, 48-70.
- [5] Y. Cao, D. Maulik and Y. Toda, *Genus zero Gopakumar-Vafa type invariants for Calabi-Yau 4-folds*, Adv. in Math. 338, (2018), 41-92.
- [6] D. Cox and S. Katz, *Mirror symmetry and algebraic geometry*, Mathematical Surveys and Monographs, 68. American Mathematical Society, Providence, RI, 1999.
- [7] O. Debarre, *Curves of low degrees in Fano varieties*, Lecture notes given at Università Roma 2 on May 2012.
- [8] R. Gopakumar and C. Vafa, *M-Theory and Topological Strings-II*, hep-th/9812127.
- [9] T. Graber and R. Pandharipande, *Localization of virtual classes*, Invent. Math. 135 (1999), no. 2, 487-518.
- [10] D. Huybrechts, *Fourier-Mukai transforms in algebraic geometry*, Oxford Mathematical Monographs. The Clarendon Press, Oxford University Press, Oxford, 2006.
- [11] S. Katz, *Genus zero Gopakumar-Vafa invariants of contractible curves*, J. Diff. Geom. 79 (2008), no. 2, 185-195.
- [12] Y. H. Kiem and J. Li, *Gromov-Witten invariants of varieties with holomorphic 2-forms*, arXiv:0707.2986.
- [13] A. Klemm and R. Pandharipande, *Enumerative geometry of Calabi-Yau 4-folds*, Comm. Math. Phys. 281, 621-653 (2008).
- [14] J. Kollár, *Rational curves on algebraic varieties*, Ergebnisse der Mathematik und ihrer Grenzgebiete 32, Springer Verlag, Berlin, 1996.
- [15] R. Thomas, *A holomorphic Casson invariant for Calabi-Yau 3-folds, and bundles on K3 fibrations*, J. Diff. Geom. 54 (2000) 367-438.