

ADMISSIBLE MODULES AND NORMALITY OF CLASSICAL NILPOTENT ORBITS

DAN BARBASCH AND KAYUE DANIEL WONG

ABSTRACT. In the case of complex classical groups, we find (\mathfrak{g}, K) -modules with the property that their K -structure matches the structure of regular functions on the closures of nilpotent orbits.

1. INTRODUCTION

In this manuscript, we study Zariski closures of complex classical nilpotent orbits.

In [Br], Rane Brylinski constructs a Dixmier algebra $\mathcal{B}(\overline{\mathcal{O}})$ associated to each Zariski closure of classical nilpotent orbit $\overline{\mathcal{O}}$. In particular, $\mathcal{B}(\overline{\mathcal{O}})$ has a (\mathfrak{g}_c, K_c) -module structure, with its $K_c \cong G$ spectrum isomorphic to $R(\overline{\mathcal{O}})$.

Normality of nilpotent orbit closures is of interest in both algebraic geometry and representation theory. We state a couple of representation theoretic perspectives on this topic. First of all, it affects the ‘quantizations’ of certain complex nilpotent orbits. Roughly speaking, quantization of a complex nilpotent orbit amounts to attaching a unitarizable (\mathfrak{g}_c, K_c) -module to the orbit. Work of Vogan, for example, [V1, Section 5 and 6], suggests that the lack of normality of certain nilpotent varieties complicates the quantization scheme of these orbits.

From another perspective, it is a standard fact in algebraic geometry that $\overline{\mathcal{O}}$ is not normal if and only if the ring of regular functions of \mathcal{O} is not isomorphic to that of $\overline{\mathcal{O}}$ as G -modules, i.e. $\overline{\mathcal{O}}$ is not normal iff $R(\mathcal{O}) \neq R(\overline{\mathcal{O}})$. It would be natural to ask which nilpotent varieties $\overline{\mathcal{O}}$ are not normal, and to make the discrepancy between $R(\mathcal{O})$ and $R(\overline{\mathcal{O}})$ explicit.

The classification of nilpotent varieties with non-normal closures is given by Kraft-Procesi [KP]. In fact, the construction of $\mathcal{B}(\overline{\mathcal{O}})$ given by R. Brylinski is based on that of Kraft and Procesi.

In this manuscript, we make the (\mathfrak{g}_c, K_c) -structure of $\mathcal{B}(\overline{\mathcal{O}})$ explicit in the classical cases $G = Sp(2n, \mathbb{C})$ and $O(n, \mathbb{C})$. In particular, since the K_c -structure of $\mathcal{B}(\overline{\mathcal{O}})$ and $R(\overline{\mathcal{O}})$ coincide, we get formulas for multiplicities of $K_c \cong G$ -types in the space of regular functions on the closure of a nilpotent orbit.

Date: January 25, 2022.

The techniques apply to all nilpotent orbits, but for clearness of exposition we focus on a special class of nilpotent orbits in Type C . This class of nilpotent orbits, given in section 1.3, exhibits all the difficulties.

The main result is Theorem 6.1.1, which provides the composition series of $\mathcal{B}(\overline{\mathcal{O}})$. As a corollary, one can compute the multiplicities of **small K -types** [B1, Definition 3.1] appearing in $\mathcal{B}(\overline{\mathcal{O}})$ and also $R(\overline{\mathcal{O}})$. There is an older definition of **small** in [V]. To avoid possible confusion, we will call the K -types we use **diminutive**.

Since orbits are normal if and only if $R(\mathcal{O}) = R(\overline{\mathcal{O}})$, our result, along with the results of [B5] and [Wo2] on the K -spectrum of $R(\mathcal{O})$, recover the normality criterion of [KP]; we do make use of some of the techniques introduced there. This also verifies a conjecture made by the second author [Wo2] on the first occurrence of the discrepancy between $R(\mathcal{O})$ and $R(\overline{\mathcal{O}})$ when $\overline{\mathcal{O}}$ is not normal.

The composition factors of $\mathcal{B}(\overline{\mathcal{Q}})$ for an orthogonal nilpotent orbit $\overline{\mathcal{Q}}$ corresponds to that of $\mathcal{B}(\overline{\mathcal{O}})$ under the dual pair correspondence, where $\overline{\mathcal{O}}$ is the symplectic nilpotent orbit with one extra column added to $\overline{\mathcal{Q}}$ (see Section 7 for details). It would be of interest to give a theoretical account for the observation.

The results provide algebraic geometric insight into how far sections on $R(\mathcal{O})$ extend to the closure $\overline{\mathcal{O}}$; we do not pursue this topic in this paper.

1.1. Langlands Parameters. We recall the Langlands parametrization of irreducible (\mathfrak{g}_c, K_c) -modules for a complex Lie group G viewed as a real Lie group. Fix a maximal compact subgroup K , and a pair $(B, H = TA)$ where B is a real Borel subgroup and H is a θ -stable Cartan subgroup such that $T = B \cap H$, and A the complement stabilized by θ .

The *Langlands parameter* of any irreducible module is a pair (λ_L, λ_R) such that $\mu := \lambda_L - \lambda_R$ is the parameter of a character of T in the decomposition of the θ -stable Cartan subalgebra $H = T \cdot A$, and $\nu := \lambda_L + \lambda_R$ the A -character. The *principal series representation* associated to (λ_L, λ_R) is the (\mathfrak{g}_c, K_c) -module

$$X(\lambda_L, \lambda_R) = \text{Ind}_B^G(e^\mu \otimes e^\nu \otimes 1)_{K\text{-finite}}.$$

The symbol Ind refers to Harish-Chandra induction. The infinitesimal character, when \mathfrak{g}_c is identified with $\mathfrak{g} \times \mathfrak{g}$, is (λ_L, λ_R) . Since $e^{(\lambda_L, \lambda_R)}|_T = e^\mu$,

$$X(\lambda_L, \lambda_R)|_K = \text{Ind}_T^K(e^\mu).$$

Let $\overline{X}(\lambda_L, \lambda_R)$ be the unique irreducible subquotient of $X(\lambda_L, \lambda_R)$ containing the K -type with extremal weight $\mu = \lambda_L - \lambda_R$. This is called the Langlands subquotient.

Proposition 1.1.1 (Parthasarathy-Rao-Varadarajan, Zhelobenko). *Let (λ_L, λ_R) and (λ'_L, λ'_R) be parameters. The following are equivalent:*

- $(\lambda_L, \lambda_R) = (w\lambda'_L, w\lambda'_R)$ for some $w \in W$.

- $X(\lambda_L, \lambda_R)$ and $X(\lambda'_L, \lambda'_R)$ have the same composition factors with same multiplicities.
- The Langlands subquotient of $X(\lambda_L, \lambda_R)$, written as $\overline{X}(\lambda_L, \lambda_R)$, is the same as that of $X(\lambda'_L, \lambda'_R)$.

Furthermore:

- Every irreducible (\mathfrak{g}_c, K_c) -module is equivalent to some $\overline{X}(\lambda_L, \lambda_R)$.
- When $\operatorname{Re} \nu$ is dominant with respect to the roots in B , $X(\lambda_L, \lambda_R)$ has a unique irreducible quotient identical to $\overline{X}(\lambda_L, \lambda_R)$; it is the image of the long intertwining operator given by integration see [Kn] for details.

If we need to specify the group, the standard module and Langlands quotient will acquire a subscript, e.g. $X_G(\lambda_L, \lambda_R)$ or $\overline{X}_G(\lambda_L, \lambda_R)$. If G is omitted in the notation, G is assumed Type C .

We will often write (λ_L, λ_R) in matrix notation $(\lambda_L, \lambda_R) \cong \begin{pmatrix} \lambda_L \\ \lambda_R \end{pmatrix}$. Also, we introduce some shorthand notations for the Langlands parameters:

- Let $x \leq y$ be two real numbers with $y - x \in \mathbb{N}$. We denote by $\Delta(x, y)$ the spherical Langlands parameter

$$\Delta(x, y) := \begin{pmatrix} x, & x+1, & \dots & y-1, & y \\ x, & x+1, & \dots & y-1, & y \end{pmatrix}.$$

- Let $2x', 2y' \in \mathbb{N}$ be such that $y' - x' \in \mathbb{N}$, and $x' < y'$. We write $\Delta(\underline{-x'}, y')$ for the non-spherical parameter

$$\Delta(\underline{-x'}, y') := \begin{pmatrix} -x', & \dots & x', & x'+1, & x'+2, & \dots & y' \\ -x'-1, & \dots & x'-1, & x', & x'+2, & \dots & y' \end{pmatrix}.$$

1.2. Strings and Intertwining Operators. We introduce more shorthand notation for induced modules.

Definition 1.2.1. Let Ψ_k be a representation of G_k with the same type as G of lower rank. The combination of strings

$$I^G(\Delta(a, A) \dots \Delta(b, B)\Psi_k)$$

denotes the induced module

$$\operatorname{Ind}_{GL^x \dots \times GL^y G_k}^G(\overline{X}_{GL}(\Delta(a, A)) \boxtimes \dots \boxtimes \overline{X}_{GL}(\Delta(b, B)) \boxtimes \Psi_k)$$

and similarly for $\Delta(\underline{-a'}, A')$, \dots , $\Delta(\underline{-b'}, B')$.

In particular, the Langlands quotient $\overline{X}_{GL}(\Delta(x, y))$ and $\overline{X}_{GL}(\Delta(\underline{-x'+1}, x'))$ are the one-dimensional representations $|\det|^{\frac{y-x}{2}}$ and \det respectively.

A careful study of various intertwining operators was done in Section 6 of [B4]. There are two types of intertwining operators that play a role:

$$\begin{aligned}\iota &: I^{Sp}(\dots \Delta(a, A)\Delta(b, B)\dots) \rightarrow I^{Sp}(\dots \Delta(b, B)\Delta(a, A)\dots) \\ \sigma &: I^{Sp}(\dots \Delta(b, B)T(L)) \rightarrow I^{Sp}(\dots \Delta(-B, -b)T(L)) \quad (\text{or without } T(L))\end{aligned}$$

where $T(L) := \overline{X}((1, \dots, L))$ denotes the trivial representation of $Sp(2L)$. We begin the study of intertwining operators for induced modules with only two strings:

Lemma 1.2.2. *Assume that $b \leq a \leq B \leq A$.*

(a) *Consider the intertwining operator*

$$I^{GL}(\Delta(a, A)\Delta(b, B)) \xrightarrow{\iota} I^{GL}(\Delta(b, B)\Delta(a, A))$$

(i) *If $b - a \notin \mathbb{Z}$, then ι is an isomorphism, in particular it is onto, and the kernel is equal to 0.*

(ii) *Whenever a, A, b, B satisfy*

$$\{a, a + 1, \dots, A\} \subseteq \{b, b + 1, \dots, B\} \quad \text{or} \quad \{b, b + 1, \dots, B\} \subseteq \{a, a + 1, \dots, A\},$$

ι is an isomorphism, i.e. it is onto and the kernel is 0.

(iii) *If $b - a \in \mathbb{Z}$ with $b < a \leq 1 \leq B < A$, then the kernel of ι is the irreducible module with parameter*

$$\begin{pmatrix} a - 1 & \dots & B & a & \dots & B + 1 & B + 2 & \dots & A & b & \dots & a - 2 \\ a & \dots & B + 1 & a - 1 & \dots & B & B + 2 & \dots & A & b & \dots & a - 2 \end{pmatrix}.$$

(b) *The image of ι is equal to*

$$I^{GL}(\Delta(b, A)\Delta(a, B))$$

Furthermore, the spherical vector in the induced module in the domain of ι is cyclic.

Proof. Omitted. □

Definition 1.2.3. *Case (a)(ii),*

$$\{a, a + 1, \dots, A\} \subseteq \{b, b + 1, \dots, B\} \quad \text{or} \quad \{b, b + 1, \dots, B\} \subseteq \{a, a + 1, \dots, A\}$$

*is called **strongly nested** [B4, Section 2].*

Proposition 1.2.4. *Assume that $b \leq a \leq B \leq A$ with $b - a \in \mathbb{Z}$.*

(a) *Consider the intertwining operator*

$$I^{Sp}(\Delta(a, A)\Delta(b, B)) \xrightarrow{\iota} I^{Sp}(\Delta(b, B)\Delta(a, A))$$

(i) *If $b = a$ or $B = A$, then $\ker \iota$ has no composition factors with diminutive lowest K -type.*

(ii) If $b < a \leq 1 \leq B < A$, the only composition factor in $\ker \iota$ with diminutive lowest K -type is

$$\begin{cases} \overline{X}(\Delta(\underline{-B, A})\Delta(\underline{a, -b})) & \text{for } a \leq 0 \\ \overline{X}(\Delta(\underline{-B, A})\Delta(1, -b)) & \text{for } a = 1 \end{cases}$$

(If $b = 0$, then the $\Delta(1, -b)$ term is not present).

(b) The image of ι is equal to

$$I^{Sp}(\Delta(b, A)\Delta(a, B))$$

Proof. The proposition on the image of ι follows from Lemma 1.2.2 and induction on intertwining operators. The same Lemma also implies that $\ker \iota$ is equal to

$$(1) \quad I^{Sp} \left(\overline{X}_{GL} \left(\left(\begin{array}{cccccccccccc} a-1 & \dots & B & a & \dots & B+1 & B+2 & \dots & A & b & \dots & a-2 \\ a & \dots & B+1 & a-1 & \dots & B & B+2 & \dots & A & b & \dots & a-2 \end{array} \right) \right) \right).$$

The above induced module contains the irreducible module

$$(2) \quad \overline{X} \left(\begin{array}{cccccccccccc} a-1 & \dots & B & a & \dots & B+1 & B+2 & \dots & A & b & \dots & a-2 \\ a & \dots & B+1 & a-1 & \dots & B & B+2 & \dots & A & b & \dots & a-2 \end{array} \right)$$

as a composition factor. By Proposition 1.1.1, this is equal to

$$\begin{aligned} & \overline{X} \left(\begin{array}{cccccccccccc} -B & \dots & -a+1 & a & \dots & B+1 & B+2 & \dots & A & -a+2 & \dots & -b \\ -B-1 & \dots & -a & a-1 & \dots & B & B+2 & \dots & A & -a+2 & \dots & -b \end{array} \right) \\ = & \overline{X} \left(\begin{array}{cccccccccccc} -B & \dots & B+1 & a & \dots & -a+1 & B+2 & \dots & A & -a+2 & \dots & -b \\ -B-1 & \dots & B & a-1 & \dots & -a & B+2 & \dots & A & -a+2 & \dots & -b \end{array} \right) \\ = & \overline{X} \left(\begin{array}{cccccccccccc} -B & \dots & B+1 & B+2 & \dots & A & a & \dots & -a+1 & -a+2 & \dots & -b \\ -B-1 & \dots & B & B+2 & \dots & A & a-1 & \dots & -a & -a+2 & \dots & -b \end{array} \right) \\ = & \begin{cases} \overline{X}(\Delta(\underline{-B, A})\Delta(\underline{a, -b})) & \text{for } a \leq 0 \\ \overline{X}(\Delta(\underline{-B, A})\Delta(1, -b)) & \text{for } a = 1 \end{cases}. \end{aligned}$$

Therefore we need to show the two modules (1) and (2) have the same diminutive K -type multiplicities.

Indeed, the diminutive K -type multiplicities of the module (1) can be computed by the difference between $I^{Sp}(\Delta(a, A)\Delta(b, B))$ and $I^{Sp}(\Delta(b, A)\Delta(a, B))$. On the other hand, the diminutive K -type multiplicities of (2) can be computed using **bottom layer K -type** arguments in Section 2.7 of [B1] (an example of such calculations can be found in Section 8.1). One can check that their multiplicities coincide with each other, hence the result follows. \square

Proposition 1.2.5. *The kernel of σ does not contain any **petite** K -types in the sense of [B2].*

Proof. This follows from a special case in Section 6 of [B4]. \square

Since all diminutive K -types are petite, the same result holds for diminutive K -types.

The intertwining operators of induced modules with more than two strings are induced from the operators in the above two propositions. If we only keep track on diminutive K -types, the composition factors of the kernel and image of the intertwining operators with diminutive, or more generally, petite lowest K -types can be deduced from the above propositions.

1.3. The Class of Nilpotent Orbits. As already mentioned in the introduction, we give details for $\mathcal{B}(\overline{\mathcal{O}})$ in the case of orbits \mathcal{O} of Type C only.

Recall that a nilpotent orbit in type C can be parametrized by the *columns* of its Jordan form $(c_0 \geq \cdots \geq c_{2p+1})$ (set $c_{2p+1} = 0$ if necessary), so that $c_{2i} + c_{2i+1}$ is even. A special nilpotent orbit in the sense of [L] satisfies the additional condition that if c_{2i} is odd, then $c_{2i} = c_{2i+1}$.

Definition 1.3.1. A chain \mathcal{C} is a sequence of integers of the form

$$[b_0 \geq (b_1 = b_2) \geq (b_3 = b_4) \geq \cdots \geq (b_{2k-1} = b_{2k}) \geq b_{2k+1}] \quad b_i \in 2\mathbb{N},$$

or

$$[b_0, b_1] \quad b_0 = b_1 \in 2\mathbb{N} + 1.$$

Definition 1.3.2 (Chains and Pairs). For each special nilpotent orbit \mathcal{O} of Type C with Jordan form $(c_0, c_1, \dots, c_{2p+1})$, we pair up the columns, and form a union of chains as follows:

- (1) An odd sized column only occurs as a pair $c_{2j} = c_{2j+1}$. Pair up all odd columns ($c_{2i} = c_{2i+1}$), and set them aside.
- (2) Pair up all even columns ($c_{2i-1} = c_{2i}$).
- (3) Suppose the two longest unpaired columns in \mathcal{O} are $c_{2i-2} \geq c_{2j+1}$ respectively. We pair up $(c_{2i-2} \geq c_{2j+1})$, and form a chain

$$\mathcal{C} = [c_{2i-2} \geq c_{2i-1} = c_{2i} \geq \cdots \geq c_{2j-1} = c_{2j} \geq c_{2j+1}].$$

If there are no paired columns between c_{2i-2} and c_{2j+1} , then $[c_{2i-2} \geq c_{2j+1}]$ form a chain of length 2.

- (4) Continue Step 3 until there are no unpaired columns left.

Suppose \mathcal{O} has l chains $\mathcal{C}_1, \dots, \mathcal{C}_l$, where the columns of \mathcal{C}_i are strictly larger than those of \mathcal{C}_j whenever $i < j$, then we denote the **chains** of \mathcal{O} as

$$\mathcal{C}(\mathcal{O}) := \mathcal{C}_1 \sqcup \mathcal{C}_2 \sqcup \cdots \sqcup \mathcal{C}_l.$$

Remark 1.3.3. Let $G = Sp(2n, \mathbb{C})$ and $\mathcal{C}(\mathcal{O}) = \mathcal{C}_1 \sqcup \mathcal{C}_2 \sqcup \cdots \sqcup \mathcal{C}_l$ be the chains of a nilpotent orbit \mathcal{O} in \mathfrak{g} . The fundamental group of the nilpotent orbit \mathcal{O} is isomorphic

to $A(\mathcal{O}) \cong (\mathbb{Z}/2\mathbb{Z})^l$. Suppose $l' \leq l$ of such chains end with a positive even integer. Then the Lusztig quotient (see [L] for details) is isomorphic to $\overline{A}(\mathcal{O}) \cong (\mathbb{Z}/2\mathbb{Z})^{l'}$.

Example 1.3.4. The orbit $\mathcal{O} = (10, 8, 8, 4, 4, 4, 3, 3, 2, 2)$ has three chains

$$\mathcal{C}_1 : [10 \geq 8 = 8 \geq 4 = 4 \geq 4], \quad \mathcal{C}_2 : [3 = 3], \quad \mathcal{C}_3 : [2 \geq 2],$$

and the column pairs are $(10, 4)$, $(8, 8)$, $(4, 4)$, $(3, 3)$, $(2, 2)$. Also, $A(\mathcal{O}) \cong (\mathbb{Z}/2\mathbb{Z})^3$ and $\overline{A}(\mathcal{O}) \cong (\mathbb{Z}/2\mathbb{Z})^2$.

We recall the necessary and sufficient conditions for a special orbit $\overline{\mathcal{O}}$ of Type C to be non-normal:

Theorem 1.3.5 (Kraft-Procesi, [KP]). *A special nilpotent orbit \mathcal{O} of Type C has non-normal closure if and only if one of the chains, $[b_0(b_1b_2) \dots (b_{2k-1}b_{2k})b_{2k+1}]$, satisfies*

$$b_{2i-2} > (b_{2i-1} = b_{2i}) = \dots = (b_{2j-1} = b_{2j}) > b_{2j+1}$$

for some $1 \leq i \leq j \leq k$.

Example 1.3.6. The orbit $\mathcal{O} = (4, 2, 2, 0)$ in $\mathfrak{sp}(8, \mathbb{C})$ is not normal:

$$b_0 = 4 > b_1 = 2 = b_2 = 2 > b_3 = 0.$$

The orbits $(4, 4, 4, 2)$ in $\mathfrak{sp}(14, \mathbb{C})$ and $(4, 2, 2, 2)$ in $\mathfrak{sp}(10, \mathbb{C})$ are normal.

We will study the Brylinski model $\mathcal{B}(\overline{\mathcal{O}})$ for the orbits \mathcal{O} satisfying the following assumptions.

Assumption 1.3.7. *We assume all columns of \mathcal{O} are even, and each chain appearing in \mathcal{O} satisfies*

$$[b_0 \geq (b_1 = b_2) > (b_3 = b_4) > \dots > (b_{2k-3} = b_{2k-2}) > (b_{2k-1} = b_{2k}) \geq b_{2k+1}].$$

Remark 1.3.8. *Assumption 1.3.7 guarantees that the infinitesimal character of $\mathcal{B}(\overline{\mathcal{O}})$ is integral (see Equation (9)). Standard modifications of the results hold in the absence of it.*

1.4. Outline of the Manuscript. For simplicity, we often express Langlands parameters by another set of notations. More precisely, let $m, n \in \mathbb{N}$ with $m \geq n$ be paired columns as in Definition 1.3.2. Denote by $(m, n)^+$ the spherical parameter corresponding to

$$(m, n)^+ := \begin{cases} \Delta(-\frac{n}{2} + 1, \frac{m}{2}) = \begin{pmatrix} -\frac{n}{2} + 1 & \dots & \frac{m}{2} \\ -\frac{n}{2} + 1 & \dots & \frac{m}{2} \end{pmatrix} & \text{if } a, b \in 2\mathbb{N}. \\ \Delta(-\frac{n-1}{2}, \frac{m-1}{2}) = \begin{pmatrix} -\frac{n-1}{2} & \dots & \frac{m-1}{2} \\ -\frac{n-1}{2} & \dots & \frac{m-1}{2} \end{pmatrix} & \text{if } a = b \in 2\mathbb{N} + 1 \end{cases}$$

If $m \geq n > 0$ are even, denote by $(m, n)^-$ the parameter corresponding to

$$(m, n)^- := \Delta\left(\underbrace{-\frac{n}{2} + 1, \frac{m}{2}}\right) = \begin{pmatrix} -\frac{n}{2} + 1 & \cdots & \frac{n}{2} & \frac{n}{2} + 1 & \cdots & \frac{m}{2} \\ -\frac{n}{2} & \cdots & \frac{n}{2} - 1 & \frac{n}{2} + 1 & \cdots & \frac{m}{2} \end{pmatrix}.$$

We outline the main results of this manuscript for orbits \mathcal{O} satisfying Assumption 1.3.7 with $\mathcal{C}(\mathcal{O}) = [c_0(c_1c_2) \cdots (c_{2p-1}c_{2p})c_{2p+1}]$.

In Section 2, we recall the construction of $\mathcal{B}(\overline{\mathcal{O}})$ in [Br]. The construction involves the dual pair correspondence, so we use the results in [AB] to prove the following in Section 3:

Theorem 1.4.1 (Corollary 3.1.6). *Let \mathcal{O} be an orbit satisfying Assumption 1.3.7 with $\mathcal{C}(\mathcal{O}) = [c_0(c_1c_2) \cdots (c_{2p-1}c_{2p})c_{2p+1}]$. There is a non-trivial (\mathfrak{g}_c, K_c) -equivariant map*

$$\mathcal{B}(\overline{\mathcal{O}}) \longrightarrow \mathcal{B}(\mathcal{O}) := I^G((c_1c_2)^+ \cdots (c_{2p-1}c_{2p})^+ \overline{X}((c_0c_{2p+1})^+)).$$

with the spherical vector in its image. In particular, $\mathcal{B}(\overline{\mathcal{O}})$ contains all the factors of the spherical cyclic submodule of $\mathcal{B}(\mathcal{O})$.

In Section 4 we compute all possible composition factors of $\mathcal{B}(\mathcal{O})$ and $\mathcal{B}(\overline{\mathcal{O}})$. The precise result is Proposition 4.3.1. A list of auxiliary orbits $Norm(\mathcal{O})$ is provided such that every composition factor of $\mathcal{B}(\mathcal{O})$ and $\mathcal{B}(\overline{\mathcal{O}})$ must have associated variety equal to $\overline{\mathcal{O}'}$ for some $\mathcal{O}' \in Norm(\mathcal{O})$.

In Section 5, we determine which factors stated in Proposition 4.3.1 appear in $\mathcal{B}(\mathcal{O})$ (Theorem 5.2.4). The main tool in the proof of the theorem is the study the (\mathfrak{g}, K) -module maps

$$(3) \quad \begin{array}{l} I^G((c_0c_{2p+1})^+(c_1c_2)^+ \cdots (c_{2p-1}c_{2p})^+) \xrightarrow{\iota_1} I^G((c_1c_2)^+(c_0c_{2p+1})^+ \cdots (c_{2p-1}c_{2p})^+) \\ \xrightarrow{\iota_2} \cdots \\ \xrightarrow{\iota_p} I^G((c_1c_2)^+ \cdots (c_{2p-1}c_{2p})^+(c_0c_{2p+1})^+) \\ \xrightarrow{q} \mathcal{B}(\mathcal{O}), \end{array}$$

where all ι_i 's are intertwining operators ι studied in Proposition 1.2.4, and q comes from the surjective map of an induced module to the Langlands quotient $I^G((c_0c_{2p+1})^+) \rightarrow \overline{X}((c_0c_{2p+1})^+)$.

Since q is surjective, the second to last module in Equation (3) contains all composition factors of $\mathcal{B}(\mathcal{O})$. They all have diminutive lowest K -types by Proposition 4.3.1. By Remark 4.3.2, every module in Equation (3) has the same diminutive K -type multiplicities, so all modules in Equation (3) have the same composition factors

with diminutive lowest K -types. In other words,

$$\begin{aligned}
 I^G((c_0 c_{2p+1})^+ (c_1 c_2)^+ \dots (c_{2p-1} c_{2p})^+) &\approx I^G((c_1 c_2)^+ (c_0 c_{2p+1})^+ \dots (c_{2p-1} c_{2p})^+) \\
 &\vdots \\
 (4) \quad &\approx I^G((c_1 c_2)^+ \dots (c_{2p-1} c_{2p})^+ (c_0 c_{2p+1})^+) \\
 &\approx \mathcal{B}(\mathcal{O}),
 \end{aligned}$$

where $X \approx Y$ means X and Y have the same composition factors of diminutive lowest K -types with multiplicities. In Section 5.3.2, we study the kernel and image of the ι 's in Equation (3) to determine the composition factors of $\mathcal{B}(\mathcal{O})$.

Section 6 begins with the following result.

Theorem 1.4.2 (Proposition 6.2.1). *The first module $I^G((c_0 c_{2p+1})^+ (c_1 c_2)^+ \dots (c_{2p-1} c_{2p})^+)$ appearing in Equation (3) is cyclic, i.e. it is generated by the trivial K -type.*

This enables us to compute the composition series of the cyclic submodule of $\mathcal{B}(\mathcal{O})$ by determining which factors in Proposition 4.3.1 appear in the image of $\iota_p \circ \dots \circ \iota_2 \circ \iota_1$ in Equation (3). They are given in Theorem 6.1.1. By Theorem 1.4.1 these composition factors must appear in $\mathcal{B}(\overline{\mathcal{O}})$ as well. Section 6.3 provides an upper bound for the diminutive K -type multiplicities of $\mathcal{B}(\overline{\mathcal{O}})$ by comparing $R(\overline{\mathcal{O}})$ with $R(\overline{\mathcal{O}}^\#)$ for a bigger orbit $\mathcal{O}^\#$. This enables us to conclude that all factors of $\mathcal{B}(\overline{\mathcal{O}})$ are accounted for by the cyclic submodule of $\mathcal{B}(\mathcal{O})$. This completes the proof of Theorem 6.1.1, the main theorem of the manuscript.

2. THE KRAFT-PROCESI MODEL

The Kraft Procesi model, introduced in Definition 6.1 in [Br], is an admissible (\mathfrak{g}_c, K_c) -module whose K -structure matches $R(\overline{\mathcal{O}})$. Under Assumption 1.3.7, it is well adapted to the Θ -correspondence. We recall the details.

Given a nilpotent orbit in type C with columns (c_0, \dots, c_{2p+1}) , let G_{2i} be $Sp(c_{2i} + \dots + c_{2p+1}, \mathbb{C})$ and $G_{2i+1} = O(c_{2i+1} + \dots + c_{2p+1}, \mathbb{C})$. We denote $G = G_0$. Let \mathfrak{g}_j be the corresponding Lie algebras, and K_j the maximal compact subgroups. Let $G_j \times G_{j+1}$ be a reductive dual pair, and Ω_{j+1} be the Oscillator representation as in [AB].

Let $K^1 := K_1 \times \dots \times K_{2p+1}$, and $(K^1)^0$ be the connected component of the identity.

Definition 2.0.1 (Kraft-Procesi Model). *Let*

$$\Omega(\overline{\mathcal{O}}) := \Omega_1 \otimes \Omega_2 \otimes \dots \otimes \Omega_{2p+1} / (\mathfrak{g}_1 \times \dots \times \mathfrak{g}_{2p+1})(\Omega_1 \otimes \dots \otimes \Omega_{2p+1}).$$

This is a (\mathfrak{g}, K) -module. Then $(K^1)^0$ acts trivially on $\Omega(\overline{\mathcal{O}})$. So $K^1 / (K^1)^0$ acts on $\Omega(\overline{\mathcal{O}})$.

Theorem 2.0.2 (Theorem 6.3 of [Br]). *Let $\mathcal{B}(\overline{\mathcal{O}}) := \Omega(\overline{\mathcal{O}})^{K^1/(K^1)^0}$, then*

$$\mathcal{B}(\overline{\mathcal{O}}) \cong R(\overline{\mathcal{O}})$$

as K_0 -modules. We have identified the complex group G_0 with the complexification of K_0 .

3. DUAL PAIRS

Since the definition of $\mathcal{B}(\overline{\mathcal{O}})$ involves oscillator representations, we first recall some results in [AB] on the dual pair correspondence:

Theorem 3.0.1 ([AB], Corollary 3.21). *Let $H_1 \times H_2$ be a reductive dual pair, and $P = MN = P_1 \times P_2 = M_1N_1 \times M_2N_2 \subset H_1 \times H_2$ a real parabolic subgroup:*

$$(5) \quad \begin{aligned} H_1 &= O(2m + \tau), & M_1 &= GL(k) \times O(2m + \tau - 2k); \\ H_2 &= Sp(2n), & M_2 &= GL(\ell) \times Sp(2n - 2\ell); \end{aligned}$$

with $\tau = 0, 1$. Suppose there is a nontrivial M -equivariant map $\Omega_M \rightarrow \sigma_1 \boxtimes \sigma_2$. Then there is a nontrivial map

$$(6) \quad \begin{aligned} \Omega &\rightarrow \text{Ind}_{P_1}^{H_1}[\alpha_1\sigma_1] \boxtimes \text{Ind}_{P_2}^{H_2}[\alpha_2\sigma_2], \text{ where} \\ \alpha_1(g_1) &= |\det g_1|^{2n-2m-\tau+k-\ell+1}, & \alpha_2(g_2) &= |\det g_2|^{-2n+2m+\tau-k+\ell-1}. \end{aligned}$$

Remark 3.0.2. *The exponents in α_1, α_2 are reversed because the parabolic subgroups in [AB] are opposite from the ones used here.*

The following corollary is an almost direct consequence of Theorem 3.0.1:

Corollary 3.0.3.

(a) *Let Ω be the oscillator representation corresponding to the dual pair $O(2n+2M) \times Sp(2n)$. Then there is a non-trivial map*

$$\Omega \rightarrow$$

$$I^{O(2n+2M)}(\Delta(a, A) \dots \Delta(b, B)\Delta(-M+1, L)) \boxtimes I^{Sp(2n)}(\Delta(-A, -a) \dots \Delta(-B, -b)T(L))$$

with the spherical vector in its image.

(b) *Let Ω be the oscillator representation corresponding to the dual pair $O(2m) \times Sp(2m+2L)$. Then there is a non-trivial map*

$$\Omega \rightarrow I^{O(2m)}(\Delta(-A, -a) \dots \Delta(-B, b)) \boxtimes I^{Sp(2m+2L)}(\Delta(a, A) \dots \Delta(b, B)T(L))$$

with the spherical vector in its image.

Proof. We prove (a) by induction on the number of GL -factors. When there are no GL -factors, i.e. $n = L$, consider the dual pairs

$$\begin{aligned} H_1 &= O(2L + 2M), & M_1 &= GL(L + M) \times O(0); \\ H_2 &= Sp(2L), & M_2 &= GL(0) \times Sp(2L); \end{aligned}$$

and $\Omega_M \rightarrow \text{triv} \boxtimes \text{triv}$. Then

$$2n - 2m - \tau + k - \ell + 1 = 2L - (2L + 2M) - 0 + (L + M) - 0 + 1 = L - M + 1$$

and Theorem 3.0.1 implies there is a non-trivial map

$$\begin{aligned} \Omega &\longrightarrow I^{O(2L+2M)}\left(\Delta\left(-\frac{L+M-1}{2} + \frac{L-M+1}{2}, \frac{L+M-1}{2} + \frac{L-M+1}{2}\right)\right) \boxtimes T(L) \\ &= I^{O(2L+2M)}(\Delta(-M+1, L)) \boxtimes T(L). \end{aligned}$$

Therefore the corollary holds for $k = 0$.

Now suppose (a) holds when there are k GL -factors, i.e. there is a non-trivial map
(7)
 $\Omega' \longrightarrow$

$$I^{O(2m')}\left(\Delta(a', A') \dots \Delta(b, B)\Delta(-M+1, L)\right) \boxtimes I^{Sp(2n')}\left(\Delta(-A', -a') \dots \Delta(-B, -b)T(L)\right)$$

with image containing the spherical vector. We denote the above correspondence by

$$\Omega' \longrightarrow \theta(\Pi) \otimes \Pi,$$

and consider the dual pairs

$$\begin{aligned} H_1 &= O(2m), & M_1 &= GL(A - a + 1) \times O(2m'); \\ H_2 &= Sp(2n), & M_2 &= GL(A - a + 1) \times Sp(2n'). \end{aligned}$$

The notation is

$$\begin{aligned} m - m' &= n - n' = (A - a + 1) \\ \Omega_M &\rightarrow (\chi \otimes \theta(\Pi)) \boxtimes (\chi^* \otimes \Pi), \\ \chi &= \Delta\left(a - \frac{2n' - 2m' + 1}{2}, A - \frac{2n' - 2m' + 1}{2}\right) \\ \chi^* &= \Delta\left(-A + \frac{2n' - 2m' + 1}{2}, -a + \frac{2n' - 2m' + 1}{2}\right). \end{aligned}$$

Theorem 3.0.1 and induction in stages imply the existence of a map

$$\Omega \longrightarrow I^{O(2m)}(\Delta(a, A) \theta(\Pi)) \boxtimes I^{Sp(2n)}(\Delta(-A, -a) \Pi)$$

with image containing the spherical vector.

For part (b), we give details when there is only one GL -factor. Consider the dual pairs

$$\begin{aligned} H_1 &= O(2m), & M_1 &= GL(m) \times O(0); \\ H_2 &= Sp(2m + 2L), & M_2 &= GL(m) \times Sp(2L) \end{aligned}$$

where $m = A - a + 1$, and $\Omega_M \rightarrow \chi^* \boxtimes (\chi \otimes \text{triv})$, with $\chi = |\det g|^x$ for some suitable x . We obtain a non-trivial map

$$\Omega \longrightarrow I^{O(2m)}(\Delta(-A, -a)) \boxtimes I^{Sp(2m+2L)}(\Delta(a, A)T(L)).$$

The rest of the proof of part (b) is omitted. \square

3.1. A non-trivial map to an induced module. We apply the results in the previous section to construct a non-trivial (\mathfrak{g}_c, K_c) -equivariant map from $\mathcal{B}(\overline{\mathcal{O}})$ to an induced module. As before, the spherical vector is in the image.

Lemma 3.1.1. *Let \mathcal{O} be a nilpotent orbit of Type C satisfying Assumption 1.3.7 with only one chain $\mathcal{C}(\mathcal{O}) = [c_0(c_1c_2) \dots (c_{2p-1}c_{2p})c_{2p+1}]$. There is a non-trivial (\mathfrak{g}_c, K_c) -equivariant map*

$$\Omega(\overline{\mathcal{O}}) \longrightarrow I^G(\Delta(-\frac{c_1}{2} + 1, \frac{c_2}{2}) \dots \Delta(-\frac{c_{2p-1}}{2} + 1, \frac{c_{2p}}{2})\Delta(0, \frac{c_{2p+1}}{2} - 1)T(\frac{c_0}{2}))$$

with the spherical vector in its image.

Proof. We do an induction on the number of columns of \mathcal{O} . First of all, consider the case when $p = 0$, i.e. $\mathcal{C}(\mathcal{O}) = [c_0, c_1]$.

If $c_1 = 0$, the result is obvious. Suppose $c_1 \neq 0$. There are non-trivial maps

$$\begin{aligned} \Omega_1 &\longrightarrow I^{O(c_1)}(\Delta(-\frac{c_1}{2} + 1, 0)) \boxtimes I^{Sp(c_0+c_1)}(\Delta(0, \frac{c_1}{2} - 1)T(\frac{c_0}{2})) \\ &\longrightarrow \overline{X}_{O(c_1)}(\Delta(-\frac{c_1}{2} + 1, 0)) \boxtimes I^{Sp(c_0+c_1)}(\Delta(0, \frac{c_1}{2} - 1)T(\frac{c_0}{2})), \end{aligned}$$

where the first map comes from Corollary 3.0.3(b) with the dual pair $O(c_1) \times Sp(c_0 + c_1)$, and the second map is the quotient map to the Langlands quotient.

Note that $\overline{X}_{O(c_1)}(\Delta(-\frac{c_1}{2} + 1, 0)) = \overline{X}_{O(c_1)}(\Delta(0, \frac{c_1}{2} - 1))$ is the trivial G_1 -module, so the above map factors through $\mathfrak{g}_1\Omega_1$, i.e.

$$\Omega_1/\mathfrak{g}_1\Omega_1 \longrightarrow I^G(\Delta(0, \frac{c_1}{2} - 1)T(\frac{c_0}{2})).$$

Hence the Proposition holds for $p = 1$.

Suppose the Proposition holds for $\mathcal{O}' = (c_2, c_3, \dots, c_{2p+1})$, i.e. there is a non-trivial map

$$\begin{aligned} \Omega_3 \otimes \dots \otimes \Omega_{2p+1}/(\mathfrak{g}_3 \times \dots \times \mathfrak{g}_{2p+1})(\Omega_3 \otimes \dots \otimes \Omega_{2p+1}) &\longrightarrow \\ \Pi_2 := I^{G_2}(\Delta(-\frac{c_3}{2} + 1, \frac{c_4}{2}) \dots \Delta(-\frac{c_{2p-1}}{2} + 1, \frac{c_{2p}}{2})\Delta(0, \frac{c_{2p+1}}{2} - 1)T(\frac{c_2}{2})) & \end{aligned}$$

Consider the oscillator representation Ω_2 with dual pair $G_1 \times G_2$. By Corollary 3.0.3(a) with $2M = c_1$, there is a non-trivial map

$$\Omega_2 \longrightarrow \Pi_1 \boxtimes \Pi_2^*$$

with Π_1 equal to

$$\Pi_1 := I^{G_1}(\Delta(-\frac{c_3}{2} + 1, \frac{c_4}{2}) \dots \Delta(-\frac{c_{2p-1}}{2} + 1, \frac{c_{2p}}{2}) \Delta(0, \frac{c_{2p+1}}{2} - 1) \Delta(-\frac{c_1}{2} + 1, \frac{c_2}{2}))$$

Therefore, we have a non-trivial map

$$\begin{aligned} \Omega_2 \otimes (\Omega_3 \otimes \dots \otimes \Omega_{2p+1}) / (\mathfrak{g}_3 \times \dots \times \mathfrak{g}_{2p+1}) (\Omega_3 \otimes \dots \otimes \Omega_{2p+1}) &\longrightarrow (\Pi_1 \boxtimes \Pi_2^*) \otimes \Pi_2 \\ &\longrightarrow \Pi_1 \boxtimes \text{triv}_{G_2}, \end{aligned}$$

where the last map is given by projecting $\Pi_2^* \otimes \Pi_2$ onto the trivial representation. Note that the final image does not depend on the G_2 -module structure, so the map factors through $\mathfrak{g}_2 \Omega_2$, and we obtain a non-trivial map

$$\Omega_2 \otimes \dots \otimes \Omega_{2p+1} / (\mathfrak{g}_2 \times \dots \times \mathfrak{g}_{2p+1}) (\Omega_2 \otimes \dots \otimes \Omega_{2p+1}) \longrightarrow \Pi_1.$$

Finally, Corollary 3.0.3(b) with $2L = c_0$ implies that there is a non-trivial map

$$\Omega_1 \longrightarrow \Pi_0 \boxtimes \Pi_1^*,$$

where Π_0 is the induced module

$$\Pi_0 := I^G(\Delta(-\frac{c_1}{2} + 1, \frac{c_2}{2}) \dots \Delta(-\frac{c_{2p-1}}{2} + 1, \frac{c_{2p}}{2}) \Delta(0, \frac{c_{2p+1}}{2} - 1) T(\frac{c_0}{2}))$$

as in the proposition. Then the above contraction argument applies again to give the result. We omit the details about the spherical vector. \square

Remark 3.1.2. *The proof of the above lemma also yields the analogous result to 3.1.1 for the closure of the nilpotent orbit $\overline{\mathcal{Q}}$, corresponding to $\mathcal{Q} = (c_1, \dots, c_{2p+1})$ in the orthogonal case.*

Before stating an analogous result of 3.1.1 for multiple chains, we make the following observation:

Corollary 3.1.3. *Let \mathcal{O} be a nilpotent orbit of Type C satisfying Assumption 1.3.7 with only one chain $\mathcal{C}(\mathcal{O}) = [c_0(c_1c_2) \dots (c_{2p-1}c_{2p})c_{2p+1}]$. There is a non-trivial (\mathfrak{g}_c, K_c) -equivariant map*

$$\Omega(\overline{\mathcal{O}}) \longrightarrow I^G((c_1c_2)^+ \dots (c_{2p-1}c_{2p})^+ \overline{X}((c_0c_{2p+1})^+)).$$

with the spherical vector in its image.

Proof. By Proposition 3.1.1, there is a non-trivial G -equivariant map

$$\Omega(\overline{\mathcal{O}}) \longrightarrow I^G(\Delta(-\frac{c_1}{2} + 1, \frac{c_2}{2}) \dots \Delta(-\frac{c_{2p-1}}{2} + 1, \frac{c_{2p}}{2}) \Delta(0, \frac{c_{2p+1}}{2} - 1) T(\frac{c_0}{2})).$$

with the same properties. The result follows from the fact that the induced module $I^{Sp}(\Delta(0, \frac{c_{2p+1}}{2}-1)T(\frac{c_0}{2}))$ maps surjectively onto the Langlands quotient $\overline{X}(\Delta(-\frac{c_{2p+1}}{2}+1, \frac{c_0}{2})) = \overline{X}((c_0 c_{2p+1})^+)$. \square

We now state a result on a non-trivial (\mathfrak{g}_c, K_c) -equivariant map from $\mathcal{B}(\overline{\mathcal{O}})$ to an induced module for \mathcal{O} with multiple chains:

Proposition 3.1.4. *Let \mathcal{O} be a nilpotent orbit of Type C satisfying Assumption 1.3.7, and $\mathcal{C}(\mathcal{O}) = \mathcal{C}_c \sqcup \cdots \sqcup \mathcal{C}_d$, where*

$$\mathcal{C}_c = [c_0(c_1 c_2) \cdots (c_{2p-1} c_{2p}) c_{2p+1}], \cdots, \mathcal{C}_d = [d_0(d_1 d_2) \cdots (d_{2q-1} d_{2q}) d_{2q+1}].$$

There is a (\mathfrak{g}_c, K_c) -equivariant map

$$\Omega(\overline{\mathcal{O}}) \longrightarrow I^G((c_1 c_2)^+ \cdots (c_{2p-1} c_{2p})^+ \cdots (d_1 d_2)^+ \cdots (d_{2q-1} d_{2q})^+ \overline{X}((c_0 c_{2p+1})^+ \cdots (d_0 d_{2q+1})^+))$$

with the spherical vector in its image.

Proof. We give details of the proof when \mathcal{O} has two chains, i.e.

$$\mathcal{C}(\mathcal{O}) = [c_0(c_1 c_2) \cdots (c_{2p-1} c_{2p}) c_{2p+1}] \sqcup [d_0(d_1 d_2) \cdots (d_{2q-1} d_{2q}) d_{2q+1}].$$

The results in Proposition 3.1.1 can be generalized such that there exists a non-trivial map

$$\begin{aligned} \Omega(\overline{\mathcal{O}}) \longrightarrow I^G(\Delta(-\frac{c_1}{2} + 1, \frac{c_2}{2}) \cdots \Delta(-\frac{c_{2p-1}}{2} + 1, \frac{c_{2p}}{2}) \Delta(-\frac{c_{2p+1}}{2} + 1, \frac{d_0}{2}) \\ \Delta(-\frac{d_1}{2} + 1, \frac{d_2}{2}) \cdots \Delta(-\frac{d_{2q-1}}{2} + 1, \frac{d_{2q}}{2}) \Delta(0, \frac{d_{2q+1}}{2} - 1) T(\frac{c_0}{2})) \end{aligned}$$

The intertwining operator

$$\begin{aligned} (8) \quad & I^G(\Delta(-\frac{c_1}{2} + 1, \frac{c_2}{2}) \cdots \Delta(-\frac{c_{2p-1}}{2} + 1, \frac{c_{2p}}{2}) \Delta(-\frac{c_{2p+1}}{2} + 1, \frac{d_0}{2}) \\ & \Delta(-\frac{d_1}{2} + 1, \frac{d_2}{2}) \cdots \Delta(-\frac{d_{2q-1}}{2} + 1, \frac{d_{2q}}{2}) \Delta(0, \frac{d_{2q+1}}{2} - 1) T(\frac{c_0}{2})) \longrightarrow \\ & I^G(\Delta(-\frac{c_1}{2} + 1, \frac{c_2}{2}) \cdots \Delta(-\frac{c_{2p-1}}{2} + 1, \frac{c_{2p}}{2}) \Delta(-\frac{d_1}{2} + 1, \frac{d_2}{2}) \cdots \\ & \Delta(-\frac{d_{2q-1}}{2} + 1, \frac{d_{2q}}{2}) \Delta(-\frac{c_{2p+1}}{2} + 1, \frac{d_0}{2}) \Delta(0, \frac{d_{2q+1}}{2} - 1) T(\frac{c_0}{2})) \end{aligned}$$

is an isomorphism on the diminutive K -types since the strings are strongly nested. The quotient map

$$\begin{aligned} & I^{Sp}(\Delta(-\frac{c_{2p+1}}{2} + 1, \frac{d_0}{2}) \Delta(0, \frac{d_{2q+1}}{2} - 1) T(\frac{c_0}{2})) \\ & \xrightarrow{q} \overline{X}(\Delta(-\frac{c_{2p+1}}{2} + 1, \frac{c_0}{2}) \Delta(-\frac{d_{2q+1}}{2} + 1, \frac{d_0}{2})) = \overline{X}((c_0 c_{2p+1})^+ (d_0 d_{2q+1})^+) \end{aligned}$$

and all the other intertwining operators are nonzero on the trivial K -type. Hence the last module in Equation (8) maps to $I^G((c_1c_2)^+ \dots (c_{2p-1}c_{2p})^+ (d_1d_2)^+ \dots (d_{2q-1}d_{2q})^+ \overline{X}((c_0c_{2p+1})^+(d_0d_{2q+1})^+))$ with the trivial K -type in its image, and the result follows. \square

Remark 3.1.5. *The irreducible module $\overline{X}((c_0c_{2p+1})^+ \dots (d_0d_{2q+1})^+)$ of Type C appearing in the proposition above is the spherical unipotent representation attached to the orbit $(c_0, c_{2p+1}, \dots, d_0, d_{2q+1})$.*

Corollary 3.1.6. *Under the setting of Proposition 3.1.4, there is a (\mathfrak{g}_c, K_c) -equivariant map*

$$\mathcal{B}(\overline{\mathcal{O}}) \longrightarrow I^G((c_1c_2)^+ \dots (c_{2p-1}c_{2p})^+ \dots (d_1d_2)^+ \dots (d_{2q-1}d_{2q})^+ \overline{X}((c_0c_{2p+1})^+ \dots (d_0d_{2q+1})^+))$$

with the spherical vector in its image.

Proof. The inclusion map

$$\mathcal{B}(\overline{\mathcal{O}}) := \Omega(\overline{\mathcal{O}})^{K^1/(K^1)^0} \hookrightarrow \Omega(\overline{\mathcal{O}})$$

has the trivial K -type in its image. Then the result follows directly from the above inclusion and Proposition 3.1.4. \square

4. PARAMETERS ATTACHED TO A NILPOTENT ORBIT

In this section, we list the possible Langlands parameters that *can* appear in the composition series of $\mathcal{B}(\overline{\mathcal{O}})$.

For $\mathcal{O} = (c_0, c_1, \dots, c_{2p+1})$ satisfying Assumption 1.3.7, the infinitesimal character attached to $\mathcal{B}(\overline{\mathcal{O}})$ is

$$(9) \quad c_{2i} \longrightarrow \left(\frac{c_{2i}}{2}, \dots, 1\right); \quad c_{2i+1} \longrightarrow \left(\frac{c_{2i+1}}{2} - 1, \dots, 1, 0\right).$$

This is implicit from the constructions in Proposition 3.1.1. It also matches the Θ -correspondence in, for example, [Pz].

Fix an orbit \mathcal{O} satisfying Assumption 1.3.7, and let λ be as in Equation (9). There is a unique maximal primitive ideal $I(\lambda) \subset U(\mathfrak{g})$ with a given infinitesimal character λ . This determines a nilpotent orbit \mathcal{O}_λ such that any admissible irreducible (\mathfrak{g}_c, K_c) -module will have at least $\overline{\mathcal{O}_\lambda}$ as its associated variety. In particular, the spherical irreducible module with infinitesimal character λ has associated variety precisely the closure of \mathcal{O}_λ (see [BV2] for details). The composition factors of $\mathcal{B}(\overline{\mathcal{O}})$ are irreducible (\mathfrak{g}_c, K_c) -modules with infinitesimal character λ and associated varieties contained in $\overline{\mathcal{O}}$ and containing $\overline{\mathcal{O}_\lambda}$.

4.1. **Definition of $Norm(\mathcal{O})$.** We list all the orbits between \mathcal{O} and \mathcal{O}_λ . The collection of such orbits is denoted by $Norm(\mathcal{O})$.

Definition 4.1.1. Let $(c_{2i}, c_{2i+1} = c_{2i+2}, c_{2i+3})$ be four even columns of a nilpotent orbit \mathcal{O} . A **fundamental collapse** of \mathcal{O} is

- (1) $(c_{2i} > c_{2i+1} = c_{2i+2} > c_{2i+3}) \rightarrow (c_{2i}, c_{2i+1} + 2, c_{2i+2} - 2, c_{2i+3})$.
- (2) $(c_{2i} = c_{2i+1} = c_{2i+2} > c_{2i+3}) \rightarrow (c_{2i} + 1, c_{2i+1} + 1, c_{2i+2} - 2, c_{2i+3})$.
- (3) $(c_{2i} > c_{2i+1} = c_{2i+2} = c_{2i+3}) \rightarrow (c_{2i}, c_{2i+1} + 2, c_{2i+2} - 1, c_{2i+3} - 1)$.
- (4) $(c_{2i} = c_{2i+1} = c_{2i+2} = c_{2i+3}) \rightarrow (c_{2i} + 1, c_{2i+1} + 1, c_{2i+2} - 1, c_{2i+3} - 1)$.

In the first two cases, we omit the columns $c_{2i+2} - 2, c_{2i+3}$ if both terms are equal to zero. Note that when $c_{2i} \neq c_{2i+1}$, the size of c_{2i} remains unchanged after collapse. Similarly, if $c_{2i+2} \neq c_{2i+3}$, the size of c_{2i+3} is the same after collapse.

Definition 4.1.2. Let \mathcal{O} be a nilpotent orbit satisfying Assumption 1.3.7. The collection of auxiliary orbits obtained as follows will be called $Norm(\mathcal{O})$.

- (1) Select an even column pair of the form $(c_{2i-1} = c_{2i})$ appearing in \mathcal{O} . Consider the four columns $c_{2i-2} \geq c_{2i-1} = c_{2i} \geq c_{2i+1}$. The columns c_{2i-2}, c_{2i+1} may or may not be paired with each other. Replace these columns with their corresponding fundamental collapse $c'_{2i-2}, c'_{2i-1}, c'_{2i}, c'_{2i+1}$ and obtain a new orbit.
- (2) Step (1) generates a set of new orbits, one for each pair $c_{2i-1} = c_{2i}$. Repeat Step (1) on the new orbits until there are no more $(c_{2j-1} = c_{2j})$'s.

Remark 4.1.3. The smallest orbit in $Norm(\mathcal{O})$ obtained in the above definition is precisely \mathcal{O}_λ mentioned at the beginning of this section. The orbits in $Norm(\mathcal{O})$ do not necessarily satisfy Assumption 1.3.7.

Example 4.1.4. Let $\mathcal{O} = (8, 6, 6, 4, 4, 2, 2, 0)$. The set $Norm(\mathcal{O})$ is

$$(10) \quad \begin{array}{c} (8, 6, 6, 4, 4, 2, 2, 0) \\ (8, 8, 4, 4, 4, 2, 2, 0) \quad (8, 6, 6, 6, 2, 2, 2, 0) \quad (8, 6, 6, 4, 4, 4) \\ (8, 8, 5, 5, 2, 2, 2, 0) \quad (8, 8, 4, 4, 4, 4) \quad (8, 6, 6, 6, 3, 3) \\ (8, 8, 5, 5, 3, 3) \end{array}$$

The orbits $(8, 8, 5, 5, 2, 2, 2, 0)$, $(8, 6, 6, 6, 3, 3)$ and $\mathcal{O}_\lambda = (8, 8, 5, 5, 3, 3)$ in $Norm(\mathcal{O})$ do not satisfy Assumption 1.3.7.

The next definition singles out the orbits \mathcal{O} for which only (1) of Definition 4.1.1 applies. They exhibit all the difficulties encountered for proving the main results.

Definition 4.1.5 (Generic orbit). *A nilpotent orbit \mathcal{O} of Type C satisfying Assumption 1.3.7 is called **generic**, if for every chain $\mathcal{C}_b = [b_0(b_1b_2) \dots (b_{2k-1}b_{2k})b_{2k+1}]$ of \mathcal{O} ,*

$$b_{2i} > b_{2i+1} \text{ for } i = 0, k \quad \text{and} \quad b_{2i} > b_{2i+1} + 2 \text{ for } i = 1, \dots, k-1.$$

It is clear that if \mathcal{O} is generic, then every $\mathcal{O}' \in \text{Norm}(\mathcal{O})$ is generic and satisfies Assumption 1.3.7. Also, (2)-(4) of Definition 4.1.1 do not apply.

4.2. Parameters attached to $\mathcal{O}' \in \text{Norm}(\mathcal{O})$. We list the parameters of the irreducible representations with associated variety contained in $\overline{\mathcal{O}}$, and infinitesimal character given in the beginning of this section.

Definition 4.2.1. *Let $\mathcal{C}(\mathcal{O}') = \mathcal{C}_{c'} \sqcup \dots \sqcup \mathcal{C}_{d'}$ be the chains of \mathcal{O}' . For each chain $\mathcal{C}_{b'} = [b'_0(b'_1b'_2) \dots (b'_{2k-1}b'_{2k})b'_{2k+1}]$ in \mathcal{O}' ,*

(a) *If $b'_{2k+1} \in 2\mathbb{N}^+$, we assign **two** parameters to $\mathcal{C}_{b'}$:*

$$(\mathcal{C}_{b'}, +) := [b_0(b'_1b'_2)^- \dots (b'_{2k-1}b'_{2k})^- b'_{2k+1}]^+ = (b'_1, b'_2)^- \dots (b'_{2k-1}, b'_{2k})^- (b'_0, b'_{2k+1})^+;$$

$$(\mathcal{C}_{b'}, -) := [b_0(b'_1b'_2)^- \dots (b'_{2k-1}b'_{2k})^- b'_{2k+1}]^- = (b'_1, b'_2)^- \dots (b'_{2k-1}, b'_{2k})^- (b'_0, b'_{2k+1})^-.$$

(b) *If $b'_{2k+1} = 0$ (which can only happen in the last chain $\mathcal{C}_{d'}$), assign **one** parameter*

$$(\mathcal{C}_{b'}, +) := [b_0(b'_1b'_2)^- \dots (b'_{2k-1}b'_{2k})^- 0]^+ = (b'_1, b'_2)^- \dots (b'_{2k-1}, b'_{2k})^- (b'_0, 0)^+.$$

(c) *If $\mathcal{C}_{b'} = [b'_0 = b'_1]$ with $b'_0, b'_1 \in 2\mathbb{N} + 1$, assign **one** parameter*

$$(\mathcal{C}_{b'}, +) := [b'_0 = b'_1]^+ = (b'_0, b'_1)^+.$$

Let $(\mathcal{C}_{c'}, \epsilon_{c'}), \dots, (\mathcal{C}_{d'}, \epsilon_{d'})$ be the parameters obtained above for chains $\mathcal{C}_{c'}, \dots, \mathcal{C}_{d'}$ respectively, then

$$(\mathcal{C}_{c'}, \epsilon_{c'}) \sqcup \dots \sqcup (\mathcal{C}_{d'}, \epsilon_{d'})$$

is a parameter attached to \mathcal{O}' .

By Remark 1.3.3, the number of parameters attached to \mathcal{O}' is equal to $|\overline{A}(\mathcal{O}')|$, the cardinality of the Lusztig quotient of \mathcal{O}' .

The fact that these parameters exhaust all possible factors of $\mathcal{B}(\overline{\mathcal{O}})$ is detailed in Proposition 4.3.1. As a result, the number of possible distinct irreducible modules appearing in $B(\overline{\mathcal{O}})$ is equal to

$$\sum_{\mathcal{O}' \in \text{Norm}(\mathcal{O})} |\overline{A}(\mathcal{O}')|.$$

Example 4.2.2. Let $\mathcal{O} = (8, 6, 6, 4, 4, 2, 2, 0)$. Then \mathcal{O} is not generic. The parameters attached to each $\mathcal{O}' \in \text{Norm}(\mathcal{O})$ are given by

$$\begin{aligned}
& [8(66)^-(44)^-(22)^-0]^+ \\
& \begin{array}{ccc}
[88]^+[4(44)^-(22)^-0]^+ & [8(66)^-6]^+[2(22)^-0]^+ & [8(66)^-(44)^-4]^+ \\
[88]^-[4(44)^-(22)^-0]^+ & [8(66)^-6]^-[2(22)^-0]^+ & [8(66)^-(44)^-4]^+
\end{array} \\
(11) \quad & \begin{array}{ccc}
& [88]^+[4(44)^-4]^+ & \\
[88]^+[55]^+[2(22)^-0]^+ & [88]^-[4(44)^-4]^+ & [8(66)^-6]^+[33]^+ \\
[88]^-[55]^+[2(22)^-0]^+ & [88]^+[4(44)^-4]^+ & [8(66)^-6]^-[33]^+ \\
& [88]^-[4(44)^-4]^+ & \\
& [88]^+[55]^+[33]^+ & \\
& [88]^-[55]^+[33]^+ &
\end{array}
\end{aligned}$$

Note that for each $\mathcal{O}' \in \text{Norm}(\mathcal{O})$, the signs of $\epsilon_{d'}$, \dots , $\epsilon_{d'}$ are given by the sign appearing at the end of each chain.

Example 4.2.3. Let $\mathcal{O} = (10, 8, 8, 4, 4, 2)$. Then \mathcal{O} is generic with

$$\text{Norm}(\mathcal{O}) = \{\mathcal{O}, (10, 8, 8, 6, 2, 2), (10, 10, 6, 4, 4, 2), (10, 10, 6, 6, 2, 2)\}.$$

The parameters attached to each $\mathcal{O}' \in \text{Norm}(\mathcal{O})$ are given by

$$\begin{aligned}
(12) \quad & \begin{array}{ccc}
& [10(88)^-(44)^-2]^+ & \\
& [10(88)^-(44)^-2]^+ & \\
& & \\
[10(88)^-6]^+[22]^+ & & [10, 10]^+[6(44)^-2]^+ \\
[10(88)^-6]^-[22]^+ & & [10, 10]^+[6(44)^-2]^+ \\
[10(88)^-6]^+[22]^+ & & [10, 10]^-[6(44)^-2]^+ \\
[10(88)^-6]^-[22]^+ & & [10, 10]^-[6(44)^-2]^+
\end{array} \\
& \begin{array}{c}
[10, 10]^+[66]^+[22]^+, [10, 10]^-[66]^+[22]^+ \\
[10, 10]^+[66]^-[22]^+, [10, 10]^-[66]^-[22]^+ \\
[10, 10]^+[66]^+[22]^-, [10, 10]^-[66]^+[22]^+ \\
[10, 10]^+[66]^-[22]^-, [10, 10]^-[66]^-[22]^+
\end{array}
\end{aligned}$$

Remark 4.2.4. When \mathcal{O} is not generic and $\mathcal{O}' \in \text{Norm}(\mathcal{O})$ has odd columns, the parameters attached to \mathcal{O}' in Definition 4.2.1 have a different infinitesimal character than $\mathcal{B}(\overline{\mathcal{O}'})$.

4.3. Construction of Irreducible Modules. The following proposition gives an explicit construction of the irreducible modules corresponding to each parameter

attached to $\mathcal{O}' \in \text{Norm}(\mathcal{O})$ in Definition 4.2.1. In particular, the lists exhaust all possibilities of the composition factors of $\mathcal{B}(\overline{\mathcal{O}})$. We will determine which of them appear in $\mathcal{B}(\overline{\mathcal{O}})$ in Section 6.

Proposition 4.3.1. *Let \mathcal{O} be a nilpotent orbit satisfying Assumption 1.3.7, and $\mathcal{O}' \in \text{Norm}(\mathcal{O})$ have chains $\mathcal{C}(\mathcal{O}') = \mathcal{C}_{c'} \sqcup \cdots \sqcup \mathcal{C}_{d'}$, where*

$$\mathcal{C}_{c'} = [c'_0(c'_1c'_2) \cdots (c'_{2p'-1}c'_{2p'})c'_{2p'+1}], \cdots, \mathcal{C}_{d'} = [d'_0(d'_1d'_2) \cdots (d'_{2q'-1}d'_{2q'})d'_{2q'+1}].$$

For each parameter $(\mathcal{C}_{c'}, \epsilon_{c'}) \sqcup \cdots \sqcup (\mathcal{C}_{d'}, \epsilon_{d'})$ attached to \mathcal{O}' (Definition 4.2.1), the irreducible module with this parameter equals the induced module from $M := GL(c'_2) \times \cdots \times GL(c'_{2p}) \times \cdots \times GL(d'_2) \times \cdots \times GL(d'_{2q}) \times Sp$,

$$(13) \quad I^G((c'_1c'_2)^- \cdots (c'_{2p'-1}c'_{2p'})^- \cdots (d'_1d'_2)^- \cdots (d'_{2q'-1}d'_{2q'})^- \overline{X}((c'_0c'_{2p'+1})^{\epsilon_{c'}} \cdots (d'_0d'_{2q'+1})^{\epsilon_{d'}})).$$

In this formula, $\overline{X}((c'_0c'_{2p'+1})^{\epsilon_{c'}} \cdots (d'_0d'_{2q'+1})^{\epsilon_{d'}})$ is the irreducible (unipotent) representation attached to the special orbit $(c'_0, c'_{2p'+1}, \cdots, d'_0, d'_{2q'+1})$, with the listed parameter.

Moreover, the irreducible modules constructed above exhaust all irreducible modules having associated variety contained in $\overline{\mathcal{O}'}$, and infinitesimal character given by Equation (9).

Proof. The proof is in Section 8.1. □

Remark 4.3.2. *The module $\overline{X}((c'_0c'_{2p'+1})^{\epsilon_{c'}} \cdots (d'_0d'_{2q'+1})^{\epsilon_{d'}})$ is a unipotent representation attached to the orbit $(c'_0, c'_{2p'+1}, \cdots, d'_0, d'_{2q'+1})$. The multiplicity of a diminutive K -type equals its multiplicity in the induced module*

$$I^G((c'_0c'_{2p'+1})^{\epsilon_{c'}} \cdots (d'_0d'_{2q'+1})^{\epsilon_{d'}}).$$

In particular, every module in Proposition 4.3.1 has diminutive lowest K -types. More explicitly, consider the Langlands quotient map

$$I^G((c'_0c'_{2p'+1})^{\epsilon_{c'}} \cdots (d'_0d'_{2q'+1})^{\epsilon_{d'}}) \twoheadrightarrow \overline{X}((c'_0c'_{2p'+1})^{\epsilon_{c'}} \cdots (d'_0d'_{2q'+1})^{\epsilon_{d'}})$$

which is isomorphic on diminutive K -types. From Equation (13), we have

$$\begin{aligned} & \overline{X}((\mathcal{C}_{c'}, \epsilon_{c'}) \sqcup \cdots \sqcup (\mathcal{C}_{d'}, \epsilon_{d'})) \approx \\ & I^G((c'_1c'_2)^- \cdots (c'_{2p'-1}c'_{2p'})^- \cdots (d'_1d'_2)^- \cdots (d'_{2q'-1}d'_{2q'})^- (c'_0c'_{2p'+1})^{\epsilon_{c'}} \cdots (d'_0d'_{2q'+1})^{\epsilon_{d'}}). \end{aligned}$$

5. COMPOSITION FACTORS OF $\mathcal{B}(\mathcal{O})$

5.1. Definition of $\mathcal{B}(\mathcal{O})$. In order to establish the composition series of $\mathcal{B}(\overline{\mathcal{O}})$, we introduce the module $\mathcal{B}(\mathcal{O})$ with K -spectrum isomorphic to $R(\mathcal{O})$ and infinitesimal character given by Equation (9).

Definition 5.1.1. *Let \mathcal{O} be a nilpotent orbit satisfying Assumption 1.3.7 with chains $\mathcal{C}(\mathcal{O}) = \mathcal{C}_c \sqcup \cdots \sqcup \mathcal{C}_d$, where*

$$\mathcal{C}_c = [c_0(c_1c_2) \cdots (c_{2p-1}c_{2p})c_{2p+1}], \cdots, \mathcal{C}_d = [d_0(d_1d_2) \cdots (d_{2q-1}d_{2q})d_{2q+1}].$$

Associated to it, is the Levi component $M = GL(c_2) \times \cdots \times GL(c_{2p}) \times \cdots \times GL(d_2) \times \cdots \times GL(d_{2q}) \times Sp$. The module $\mathcal{B}(\mathcal{O})^{\epsilon_c, \dots, \epsilon_d}$ is defined as

$$\mathcal{B}(\mathcal{O})^{\epsilon_c, \dots, \epsilon_d} := I^G((c_1c_2)^+ \cdots (c_{2p-1}c_{2p})^+ \cdots (d_1d_2)^+ \cdots (d_{2q-1}d_{2q})^+ \overline{X}((c_0c_{2p+1})^{\epsilon_c} \cdots (d_0d_{2q+1})^{\epsilon_d})).$$

Recall that $c_1 = c_2$ etc., so that the parameter makes sense.

By [B5] and [Wo2], each $\mathcal{B}(\mathcal{O})^{\epsilon_c, \dots, \epsilon_d}$ has K -spectrum equal to $R(\mathcal{O}, \varphi)$ for some local system φ . In particular, $\mathcal{B}(\mathcal{O}) := \mathcal{B}(\mathcal{O})^{+, \dots, +} \cong R(\mathcal{O})$. Also, Corollary 3.1.6 implies that there exists a nontrivial (\mathfrak{g}_c, K_c) -module morphism $\mathcal{B}(\overline{\mathcal{O}}) \rightarrow \mathcal{B}(\mathcal{O})$.

$\mathcal{B}(\mathcal{O})^{\epsilon_c, \dots, \epsilon_d}$ are generally not irreducible. Since $\mathcal{B}(\mathcal{O})^{\epsilon_c, \dots, \epsilon_d}$ has associated variety equal to $\overline{\mathcal{O}}$ and infinitesimal character (9), the composition factors of $\mathcal{B}(\mathcal{O})^{\epsilon_c, \dots, \epsilon_d}$ must come from the irreducible modules studied in Proposition 4.3.1. The main theorem in this section is to find its composition factors.

5.2. Decomposition of $\mathcal{B}(\mathcal{O})$. We first consider the case when \mathcal{O} is generic (Definition 4.1.5) and has only one chain

$$\mathcal{C}(\mathcal{O}) = [c_0(c_1c_2) \cdots (c_{2p-1}c_{2p})c_{2p+1}].$$

There are two cases, $c_{2p+1} > 0$ and $c_{2p+1} = 0$:

Case $c_{2p+1} > 0$ corresponds to the seesaw sequence in the Θ -correspondence starting with a symplectic group; there are two modules $\mathcal{B}(\mathcal{O})^\pm$.

Case $c_{2p+1} = 0$ corresponds to the seesaw sequence starting with an orthogonal group; there is one module $\mathcal{B}(\mathcal{O})$.

Proposition 5.2.1. *Let \mathcal{O} be generic and has only one chain. Suppose $(\mathcal{C}_{c'}, \epsilon_{c'}) \sqcup \cdots \sqcup (\mathcal{C}_{d'}, \epsilon_{d'})$ are the parameters attached to $\mathcal{O}' \in \text{Norm}(\mathcal{O})$ constructed in Proposition 4.3.1.*

(i) *Assume $c_{2p+1} > 0$.*

- *The composition factors of $\mathcal{B}(\mathcal{O})^+$ have parameters $(\mathcal{C}_{c'}, \epsilon_{c'}) \sqcup \cdots \sqcup (\mathcal{C}_{d'}, \epsilon_{d'})$ such that there is an even number of $-$'s in the expression of (13) with multiplicity 1.*
- *The composition factors of $\mathcal{B}(\mathcal{O})^-$ have parameters such that there is an odd number of $-$'s in the expression of (13) with multiplicity 1.*

(ii) *Assume $c_{2p+1} = 0$.*

- *All possible composition factors listed in 4.3.1 occur in $\mathcal{B}(\mathcal{O})^+$ with multiplicity 1.*

Example 5.2.2. Consider $\mathcal{O} = (10, 8, 8, 4, 4, 2)$. Then $\mathcal{B}(\mathcal{O})^+ =$

$$(14) \quad \begin{array}{l} [10(88)^-(44)^-2]^+ \\ [10(88)^-6]^- [22]^+ \qquad [10, 10]^+ [6(44)^-2]^- \\ [10(88)^-6]^+ [22]^- \qquad [10, 10]^- [6(44)^-2]^+ \\ [10, 10]^+ [66]^+ [22]^+ \\ [10, 10]^- [66]^- [22]^+ \\ [10, 10]^- [66]^+ [22]^- \\ [10, 10]^+ [66]^- [22]^- \end{array}$$

and $\mathcal{B}(\mathcal{O})^- =$

$$(15) \quad \begin{array}{l} [10(88)^-(44)^-2]^- \\ [10(88)^-6]^+ [22]^+ \qquad [10, 10]^+ [6(44)^-2]^+ \\ [10(88)^-6]^- [22]^- \qquad [10, 10]^- [6(44)^-2]^- \\ [10, 10]^- [66]^+ [22]^+ \\ [10, 10]^+ [66]^- [22]^+ \\ [10, 10]^+ [66]^+ [22]^- \\ [10, 10]^- [66]^- [22]^- \end{array}$$

Definition 5.2.3. Let \mathcal{O} be a generic orbit with a single chain $\mathcal{C}(\mathcal{O}) = \mathcal{C}$. We call all parameters $(\mathcal{C}_{c'}, \epsilon_{c'}) \sqcup \cdots \sqcup (\mathcal{C}_{d'}, \epsilon_{d'})$ attached to $\mathcal{O}' \in \text{Norm}(\mathcal{O})$ appearing in $\mathcal{B}(\mathcal{O})^\epsilon$, \mathcal{C}^ϵ -admissible.

In general, the composition factors of $\mathcal{B}(\mathcal{O})^{\epsilon_c, \dots, \epsilon_d}$ for all generic orbits \mathcal{O} are given by the following:

Theorem 5.2.4. Let \mathcal{O} be a generic orbit with chains

$$\mathcal{C}(\mathcal{O}) = \mathcal{C}_c \sqcup \cdots \sqcup \mathcal{C}_d.$$

The composition factors of $\mathcal{B}(\mathcal{O})^{\epsilon_c, \dots, \epsilon_d}$ are the irreducible modules with parameters

$$\mathcal{P}_c \sqcup \cdots \sqcup \mathcal{P}_d,$$

where each \mathcal{P}_j is a $\mathcal{C}_j^{\epsilon_j}$ -admissible parameter (Definition 5.2.3). Each factor occurs with multiplicity 1.

Remark 5.2.5. The statements of Proposition 5.2.1 and Theorem 5.2.4 hold for all orbits satisfying Assumption 1.3.7. The proofs are essentially the same; there are fewer parameters to consider.

In particular, the details of the proof of Proposition 5.2.1 for **all** chains \mathcal{C} of length 4 are in Section 5.3.2. We will only give an outline on their proofs in the more general setting in Remark 5.4.1.

Example 5.2.6. Let $\mathcal{O} = (10, 8, 8, 6, 4, 2, 2, 0)$ with $\mathcal{C}(\mathcal{O}) = [10(88)6] \sqcup [4(22)0]$. By Proposition 5.2.1, the $[10(88)6]^+$ -admissible and $[4(22)0]^+$ -admissible parameters are given by:

$$(16) \quad \begin{array}{cc} [10(88)^-6]^- & [4(22)^-0]^+ \\ [10, 10]^+[66]^+ & [44]^+ \\ [10, 10]^-[66]^- & [44]^- \end{array}$$

By Theorem 5.2.4, the composition series of $\mathcal{B}(\mathcal{O}) = \mathcal{B}(\mathcal{O})^{+,+}$ consists of the following parameters:

$$(17) \quad \begin{array}{cc} [10(88)^-6]^-[4(22)^-0]^+ & \\ [10(88)^-6]^-[44]^+ & [10, 10]^+[66]^+[4(22)^-0]^+ \\ [10(88)^-6]^-[44]^- & [10, 10]^-[66]^-[4(22)^-0]^+ \\ [10, 10]^+[66]^+[44]^+ & \\ [10, 10]^-[66]^-[44]^+ & \\ [10, 10]^+[66]^+[44]^- & \\ [10, 10]^-[66]^-[44]^- & \end{array}$$

The proof of Theorem 5.2.4 is by induction on the number of chains (see Section 5.4). In the special case when there is only one chain, i.e. Proposition 5.2.1, we will do an induction on the length of the chain.

5.3. One Chain. We now prove Proposition 5.2.1 by induction on the length of the chain.

5.3.1. *Initial Step.* When $\mathcal{O} = (c_0, c_1)$ has only two columns, then $Norm(\mathcal{O}) = \{\mathcal{O}\}$ and the result follows directly from [B5].

The next case is when there is only one bracket, i.e. orbits of the form $\mathcal{O} = (c_0, c_1, c_2, c_3)$ given in Definition 4.1.1 with $\mathcal{C}(\mathcal{O}) = [c_0(c_1c_2)c_3]$. Recall that $Norm(\mathcal{O}) = \{\mathcal{O}, \mathcal{O}'\}$, where $\mathcal{O}' := (c'_0, c'_1, c'_2, c'_3)$ is obtained from the fundamental collapse of \mathcal{O} .

Consider the $GL \times GL$ induced module

$$(18) \quad I^{Sp}((c_0c_3)^+(c_1c_2)^+) \approx I^{Sp}((c_1c_2)^+\overline{X}((c_0c_3)^+)) = \mathcal{B}(\mathcal{O})$$

(Recall in Equation (4) that $X \approx Y$ means the composition factors of X and Y with diminutive lowest K -types coincide). The composition factors of (18) with diminutive lowest K -types are precisely the factors that show up in $\mathcal{B}(\mathcal{O})$.

To show this, we study the intertwining operator between these induced modules:

$$(19) \quad I^{Sp}((c_0c_3)^+(c_1c_2)^+) \xrightarrow{\iota} I^{Sp}((c_1c_2)^+(c_0c_3)^+).$$

Note that the codomain of the above map also have the same diminutive K -type multiplicities as $\mathcal{B}(\mathcal{O})$. We will find the diminutive lowest K -type composition factors appearing in both $\ker \iota$ and $\text{im } \iota$, which add up to be the composition factors of $\mathcal{B}(\mathcal{O})$.

Kernel of ι : The following lemma gives $\ker \iota$ explicitly.

Lemma 5.3.1. *$\ker \iota$ does not contain any composition factor with diminutive lowest K -types unless $c_0 > (c_1c_2) > c_3$. In this case, the kernel is equal to*

$$\overline{X}([c_0, c_1 + 2]^- [c_2 - 2, c_3]^\mu)$$

(notation of Section 4), where $\mu = -$ if $c_3 \neq 0$, and $\mu = +$ if $c_3 = 0$. If $c_2 - 2 = c_3 = 0$, the term $[c_2 - 2, c_3]^\mu$ is omitted.

Proof. This is a direct application of Proposition 1.2.4. □

Image of ι : Again, by Proposition 1.2.4,

$$\text{im } \iota = I^{Sp}(\Delta(-\frac{c_1}{2} + 1, \frac{c_0}{2})\Delta(-\frac{c_3}{2} + 1, \frac{c_2}{2})).$$

To make $\text{im } \iota$ explicit, we compute the image and kernel of the composition of intertwining operators

$$(20) \quad \begin{aligned} \text{im } \iota &= I^{Sp}(\Delta(-\frac{c_1}{2} + 1, \frac{c_0}{2})\Delta(-\frac{c_3}{2} + 1, \frac{c_2}{2})) \xrightarrow{\sigma} \\ &I^{Sp}(\Delta(-\frac{c_1}{2} + 1, \frac{c_0}{2})\Delta(-\frac{c_2}{2}, \frac{c_3}{2} - 1)) \xrightarrow{\iota} \\ &I^{Sp}(\Delta(-\frac{c_2}{2}, \frac{c_3}{2} - 1)\Delta(-\frac{c_1}{2} + 1, \frac{c_0}{2})). \end{aligned}$$

They are computed by a repeated application of Propositions 1.2.4 and 1.2.5. For the image, note that the first map of (20) is an isomorphism on the diminutive K -types. The second map has image equal to

$$I^{Sp}(\Delta(-\frac{c_2}{2}, \frac{c_0}{2})\Delta(-\frac{c_1}{2} + 1, \frac{c_3}{2} - 1)) = I^{Sp}((c'_0c'_1)^+(c'_2c'_3)^+)$$

(If $c'_2 = c'_3 = 0$, we omit the term $[c'_2, c'_3]^+$).

For the kernel we only need the second map in (20). By Proposition 1.2.4 again, it is equal to $\overline{X}([c_0(c_1c_2)^-c_3]^\mu)$.

In conclusion, the kernel and image of ι in Equation (19) with diminutive lowest K -types are given as follows:

	$\mathcal{C}(\mathcal{O})$	$\ker \iota$	$\text{im } \iota$
(21)	$[c_0 > (c_1 c_2) > c_3]$	$[c'_0 c'_1]^- [c'_2 c'_3]^\mu$	$[c_0 (c_1 c_2)^- c_3]^\mu$ $[c'_0 c'_1]^+ [c'_2 c'_3]^+$
	$[c_0 = (c_1 c_2) > c_3]$	0	
	$[c_0 > (c_1 c_2) = c_3]$		
	$[c_0 = (c_1 c_2) = c_3]$		

When $c'_2 = c'_3 = 0$, the term $[c'_2 c'_3]^\mu$ or $[c'_2 c'_3]^+$ is not present.

Proposition 5.3.2. *Proposition 5.2.1 holds for all orbits \mathcal{O} satisfying Assumption 1.3.7 with $\mathcal{C}(\mathcal{O}) = [c_0(c_1 c_2)c_3]$.*

Proof. For $\mathcal{B}(\mathcal{O})$, the parameters appearing in $\ker \iota$ and $\text{im } \iota$ in Table (21) are precisely the ones given by Proposition 5.2.1(a). Hence the proof is complete for $\mathcal{B}(\mathcal{O})$.

For $\mathcal{B}(\mathcal{O})^-$, consider the kernel and image of the intertwining operator

$$(22) \quad \begin{aligned} & I^{Sp}(\Delta(-\frac{c_3}{2}, \frac{c_3}{2} - 1)\Delta(\frac{c_3}{2} + 1, \frac{c_0}{2})\Delta(-\frac{c_1}{2} + 1, \frac{c_2}{2})) \longrightarrow \\ & I^{Sp}(\Delta(-\frac{c_3}{2}, \frac{c_3}{2} - 1)\Delta(-\frac{c_1}{2} + 1, \frac{c_2}{2})\Delta(\frac{c_3}{2} + 1, \frac{c_0}{2})) \end{aligned}$$

The techniques used above to prove the Proposition for $\mathcal{B}(\mathcal{O})$ can be carried over to proving the Proposition for $\mathcal{B}(\mathcal{O})^-$. \square

5.3.2. *Induction step on the length of one chain.* We now prove Proposition 5.2.1, using induction on the length of a chain and the results in Section 5.4.

We only give the proof for $\mathcal{B}(\mathcal{O})^+$. Suppose the proposition holds for $p \leq k - 1$, consider the orbit \mathcal{O} with

$$\mathcal{C}(\mathcal{O}) = [c_0(c_1 c_2) \dots (c_{2k-1} c_{2k})(c_{2k-1} c_{2k}) c_{2k+1}].$$

We study the intertwining operators ι_l for $l = 1, \dots, k$:

$$(23) \quad \begin{aligned} & I^{Sp}((c_1 c_2)^+ \dots (c_0 c_{2k+1})^+ (c_{2l-1} c_{2l})^+ \dots (c_{2k-1} c_{2k})^+) \\ & \xrightarrow{\iota_l} I^{Sp}((c_1 c_2)^+ \dots (c_{2l-1} c_{2l})^+ (c_0 c_{2k+1})^+ \dots (c_{2k-1} c_{2k})^+) \end{aligned}$$

As in the previous subsection, we compute the kernels and images appearing in the intertwining operators ι_l for $l = 1, \dots, k$. The composition factors of $\mathcal{B}(\mathcal{O})$ follow from this calculation.

Kernel of ι_l : By applying the results in Table (21) for $(c_0, c_{2l-1}, c_{2l}, c_{2k+1})$ and the fact that the intertwining operators are induced from corresponding ones on Levi components, the composition factors with diminutive lowest K -types in $\ker \iota_l$ are given by $\ker \iota_l =$

$$I^G(((c_1 c_2)^+ \dots (c_{2l-3} c_{2l-2})^+ (c_{2l+1} c_{2l+2})^+ \dots (c_{2k-1} c_{2k})^+ \overline{X}((c_0, c_{2l-1} + 2)^- (c_{2l} - 2, c_{2k+1})^-)).$$

This is equal to $\mathcal{B}(\mathcal{O}_l)^{-,-}$, where $\mathcal{C}(\mathcal{O}_l) = \mathcal{C}_1 \sqcup \mathcal{C}_2$ with

$$\mathcal{C}_1 = [c_0(c_1c_2) \dots (c_{2l-3}c_{2l-2})c_{2l-1} + 2], \quad \mathcal{C}_2 = [c_{2l} - 2(c_{2l+1}c_{2l+2}) \dots (c_{2k-1}c_{2k})c_{2k+1}].$$

Note that the lengths of \mathcal{C}_i are smaller than that of $\mathcal{C}(\mathcal{O})$. By induction hypothesis and the result in Section 5.4, the composition factors of $\ker \iota_l = \mathcal{B}(\mathcal{O}_l)^{-,-}$ are of the form $\overline{X}(\mathcal{P}_1 \sqcup \mathcal{P}_2)$, where \mathcal{P}_i are \mathcal{C}_i^- -admissible parameters.

Image of ι_l : The composition factors can be read off from Table (21). The factors with diminutive lowest K -types in $\text{im } \iota_l$ contain the composition factors in

$$I^G((c_1c_2)^+ \dots (c_{2l-3}c_{2l-2})^+(c_{2l+1}c_{2l+2})^+ \dots (c_{2k-1}c_{2k})^+ \overline{X}((c_0, c_{2l-1} + 2)^+(c_{2l} - 2, c_{2k+1})^+)).$$

They equal $\mathcal{B}(\mathcal{O}_l)^{+,+}$. So the composition factors of $\text{im } \iota_l$ include the irreducible modules $\overline{X}(\mathcal{Q}_1 \sqcup \mathcal{Q}_2)$, where \mathcal{Q}_i are \mathcal{C}_i^+ -admissible parameters.

Consequently, $\ker \iota_l$ and $\text{im } \iota_l$ contain the factors $\overline{X}(\mathcal{P}_1 \sqcup \mathcal{P}_2)$ and $\overline{X}(\mathcal{Q}_1 \sqcup \mathcal{Q}_2)$. They include all parameters (with **even** number of minuses if $c_{2k+1} \neq 0$) attached to orbits $\mathcal{O}'_l \in \text{Norm}(\mathcal{O})$ smaller than or equal to $\mathcal{O}_l = (c_0, \dots, c_{2l-2}, c_{2l-1} + 2, c_{2l} - 2, c_{2l+1}, \dots, c_{2k+1})$. In other words, every irreducible module given in Proposition 5.2.1(a), whose parameters are attached to $\mathcal{O}' \neq \mathcal{O} \in \text{Norm}(\mathcal{O})$, shows up in the kernel or image of ι_l for some l , and hence in $\mathcal{B}(\mathcal{O})$ as well.

We are left to show that the (unique) parameter attached to $\mathcal{O} \in \text{Norm}(\mathcal{O})$ described in Proposition 5.2.1(a) constitutes a composition factor of $\mathcal{B}(\mathcal{O})$, and the multiplicities are 1.

Both statements can be verified by comparing the K -spectrum of $\mathcal{B}(\mathcal{O}) \cong R(\mathcal{O})$ (which is known by the work of [B5]) and the K -spectrum of the irreducible modules attached to $\mathcal{O}' \in \text{Norm}(\mathcal{O})$ (which is known by Remark 4.3.2). \square

Example 5.3.3. *We illustrate the above argument in the case of $\mathcal{O} = (10, 8, 8, 4, 4, 2)$, i.e. we will account for the factors of $\mathcal{B}(\mathcal{O})$ in Example 5.2.2.*

Consider the intertwining operators

$$I^{Sp}((10, 2)^+(88)^+(44)^+) \xrightarrow{\iota_1} I^{Sp}((88)^+(10, 2)^+(44)^+) \xrightarrow{\iota_2} I^{Sp}((88)^+(44)^+(10, 2)^+).$$

The kernel of ι_1 with diminutive K -type is given by $\mathcal{B}(10, 10, 6, 4, 4, 2)^{-,-}$. By Proposition 5.3.2 and the results in Section 5.4, its composition series consists of irreducible modules with parameters

$$\ker \iota_1 : [10, 10]^- [6(44)^- 2]^+, [10, 10]^- [66]^- [22]^+, [10, 10]^- [66]^+ [22]^-.$$

The image of ι_1 contains the induced module $\mathcal{B}(10, 10, 6, 4, 4, 2)^{+,+}$. Its composition series consists of irreducible modules with parameters

$$\text{im } \iota_1 : [10, 10]^+ [6(44)^- 2]^- , [10, 10]^+ [66]^+ [22]^+, [10, 10]^+ [66]^- [22]^-.$$

The kernel of ι_2 with diminutive K -type is given by $\mathcal{B}(10, 8, 8, 6, 2, 2)^{-, -}$. Its composition series consists of irreducible modules with parameters

$$\ker \iota_2 : [10(88)^-6]^+[22]^-, [10, 10]^+[66]^-[22]^-, [10, 10]^-[66]^+[22]^-$$

The image of ι_2 contains the induced module $\mathcal{B}(10, 8, 8, 6, 2, 2)^{+, +}$. Its composition series consists of irreducible modules with parameters

$$\text{im } \iota_2 : [10(88)^-6]^-[22]^+, [10, 10]^+[66]^+[22]^+, [10, 10]^-[66]^-[22]^+$$

By direct comparison, the kernel and image of ι_1 and ι_2 consists of the 8 factors of $\mathcal{B}(\mathcal{O})$ in Example 5.2.2 attached to the orbits $\mathcal{O}' < \mathcal{O}$ in $\text{Norm}(\mathcal{O})$.

If we consider the GK-dimension of $\mathcal{B}(\mathcal{O})$, it must contain a factor attached to $\mathcal{O} \in \text{Norm}(\mathcal{O})$. We pick the factor $\overline{X}([10(88)^-(44)^-2]^+)$ so that, using Remark 4.3.2, the diminutive K -type multiplicities of the sum of the 8 factors in the previous paragraph (with multiplicity one) with the factor $\overline{X}([10(88)^-(44)^-2]^+)$ is exactly equal to that of $R(\mathcal{O})$.

In conclusion, the 9 factors in Example 5.2.2 exhaust all composition factors of $\mathcal{B}(\mathcal{O})$.

5.4. If Theorem 5.2.4 holds for l chains, it holds for $l + 1$ chains. We show how the validity for one chain implies the validity for two chains, and the general case is analogous.

Assume $\mathcal{C}(\mathcal{O}) = \mathcal{C}_1 \sqcup \mathcal{C}_2$. Let

$$\mathcal{C}_1 = [c_0(c_1c_2) \dots (c_{2p-1}c_{2p})c_{2p+1}], \quad \mathcal{C}_2 = [d_0(d_1d_2) \dots (d_{2q-1}d_{2q})d_{2q+1}]$$

be the two chains. Then

$$\mathcal{B}(\mathcal{O})^{\epsilon_1, \epsilon_2} = I^{Sp}((c_1c_2)^+ \dots (c_{2p-1}c_{2p})^+(d_1d_2)^+ \dots (d_{2q-1}d_{2q})^+\overline{X}((c_0c_{2p+1})^{\epsilon_1}(d_0d_{2q+1})^{\epsilon_2}))$$

By Remark 4.3.2,

$$\begin{aligned} (24) \quad \mathcal{B}(\mathcal{O})^{\epsilon_1, \epsilon_2} &\approx I^{Sp}((c_1c_2)^+ \dots (c_{2p-1}c_{2p})^+(d_1d_2)^+ \dots (d_{2q-1}d_{2q})^+(c_0c_{2p+1})^{\epsilon_1}(d_0d_{2q+1})^{\epsilon_2}) \\ &\approx I^{Sp}((d_1d_2)^+ \dots (d_{2q-1}d_{2q})^+(d_0d_{2q+1})^{\epsilon_2}(c_1c_2)^+ \dots (c_{2p-1}c_{2p})^+(c_0c_{2p+1})^{\epsilon_1}) \\ &\approx I^{Sp}((d_1d_2)^+ \dots (d_{2q-1}d_{2q})^+(d_0d_{2q+1})^{\epsilon_2}I^{Sp}((c_1c_2)^+ \dots (c_{2p-1}c_{2p})^+(c_0c_{2p+1})^{\epsilon_1})). \end{aligned}$$

Let \mathcal{A}_1 be the collection of all $\mathcal{C}_1^{\epsilon_1}$ -admissible parameters (Definition 5.2.3). The induction hypothesis of Proposition 5.2.1 implies that

$$(25) \quad I^{Sp}((c_1c_2)^+ \dots (c_{2p-1}c_{2p})^+(c_0c_{2p+1})^{\epsilon_1}) \approx \mathcal{B}(c_0, \dots, c_{2p+1})^{\epsilon_1} \approx \bigoplus_{\mathcal{R}_1 \in \mathcal{A}_1} \overline{X}(\mathcal{R}_1).$$

Recall from Proposition 4.3.1 and Remark 4.3.2 that

$$\overline{X}(\mathcal{R}_1) \approx I^{Sp}((r_1r_2)^- \dots (r_{2i-1}r_{2i})^- \dots (s_1s_2)^- \dots (s_{2j-1}s_{2j})^-(r_0r_{2i+1})^{\epsilon_r} \dots (s_0s_{2j+1})^{\epsilon_s})$$

for some $r_0 \geq \dots \geq r_{2i+1} > \dots > s_0 \geq \dots \geq s_{2j+1}$ and $\epsilon_r, \dots, \epsilon_s \in \{+, -\}$. Let $I^{GL}(\mathcal{R}_1)$ be the module

$$I^{GL}(\mathcal{R}_1) := I^{GL}((r_1 r_2)^- \dots (r_{2i-1} r_{2i})^- \dots (s_1 s_2)^- \dots (s_{2j-1} s_{2j})^- (r_0 r_{2i+1})^{\epsilon_r} \dots (s_0 s_{2j+1})^{\epsilon_s})$$

such that $I^{Sp}(I^{GL}(\mathcal{R}_1)) \approx \overline{X}(\mathcal{R}_1)$. Using Equations (24) and (25), we have

$$\begin{aligned} \mathcal{B}(\mathcal{O})^{\epsilon_1, \epsilon_2} &\approx \bigoplus_{\mathcal{R}_1 \in \mathcal{A}_1} I^{Sp}((d_1 d_2)^+ \dots (d_{2q-1} d_{2q})^+ (d_0 d_{2q+1})^{\epsilon_2} \overline{X}(\mathcal{R}_1)) \\ (26) \quad &\approx \bigoplus_{\mathcal{R}_1 \in \mathcal{A}_1} I^{Sp}((d_1 d_2)^+ \dots (d_{2q-1} d_{2q})^+ (d_0 d_{2q+1})^{\epsilon_2} I^{GL}(\mathcal{R}_1)) \\ &\approx \bigoplus_{\mathcal{R}_1 \in \mathcal{A}_1} I^{Sp}(I^{GL}(\mathcal{R}_1) I^{Sp}((d_1 d_2)^+ \dots (d_{2q-1} d_{2q})^+ (d_0 d_{2q+1})^{\epsilon_2})) \end{aligned}$$

Let \mathcal{A}_2 be the collection of $\mathcal{C}_2^{\epsilon_2}$ -admissible parameters. Then the induction hypothesis for Proposition 5.2.1 implies

$$(27) \quad I^{Sp}((d_1 d_2)^+ \dots (d_{2q-1} d_{2q})^+ (d_0 d_{2q+1})^{\epsilon_2}) \approx \mathcal{B}(d_0, \dots, d_{2q+1})^{\epsilon_2} = \bigoplus_{\mathcal{R}_2 \in \mathcal{A}_2} \overline{X}(\mathcal{R}_2).$$

Plugging Equation (27) into (26), we obtain

$$\mathcal{B}(\mathcal{O})^{\epsilon_1, \epsilon_2} \approx \bigoplus_{\mathcal{R}_1 \in \mathcal{A}_1} \bigoplus_{\mathcal{R}_2 \in \mathcal{A}_2} I^{Sp}(I^{GL}(\mathcal{R}_1) \overline{X}(\mathcal{R}_2)).$$

By Proposition 4.3.1 (on the structure of $\overline{X}(\mathcal{R}_1 \sqcup \mathcal{R}_2)$) and Remark 4.3.2 (on the diminutive K -type multiplicities of $I^{Sp}(I^{GL}(\mathcal{R}_1) \overline{X}(\mathcal{R}_2))$ and $\overline{X}(\mathcal{R}_1 \sqcup \mathcal{R}_2)$), the irreducible module with parameter $\mathcal{R}_1 \sqcup \mathcal{R}_2$ satisfies

$$I^{Sp}(I^{GL}(\mathcal{R}_1) \overline{X}(\mathcal{R}_2)) \approx \overline{X}(\mathcal{R}_1 \sqcup \mathcal{R}_2).$$

Hence $\mathcal{B}(\mathcal{O})^{\epsilon_1, \epsilon_2} \approx \bigoplus_{\mathcal{R}_1 \in \mathcal{A}_1} \bigoplus_{\mathcal{R}_2 \in \mathcal{A}_2} \overline{X}(\mathcal{R}_1 \sqcup \mathcal{R}_2)$, and the result follows. \square

Remark 5.4.1. *We describe the modifications needed in order to generalize Proposition 5.2.1 and Theorem 5.2.4 to all orbits satisfying Assumption 1.3.7. In this case, $\ker \iota_1$ or $\ker \iota_k$ may be equal to 0 (see Table (21)), and the fundamental collapse of $(c_0 = c_1 = c_2 > c_{2k+1})$ or $(c_0 > c_{2k-1} = c_{2k} = c_{2k+1})$ may yield odd columns $(c_0 + 1, c_1 + 1, c_2 - 2, c_{2k+1})$ or $(c_0, c_{2k-1} + 2, c_{2k} - 1, c_{2k+1} - 1)$ respectively.*

In these cases, $\text{im } \iota_1$ or $\text{im } \iota_k$ contains

$$\mathcal{B}(c_0 + 1, c_1 + 1, c_2 - 2, c_3, \dots, c_{2k+1})^{+,+} \approx I^G((c_0 + 1, c_0 + 1)^+ \mathcal{B}(c_2 - 2, c_3, \dots, c_{2k+1})^+)$$

or

$$\mathcal{B}(c_0, c_1, \dots, c_{2k-1} + 2, c_{2k} - 1, c_{2k+1} - 1)^{+,+} \approx I^G((c_{2k} - 1, c_{2k+1} - 1)^+ \mathcal{B}(c_0, c_1, \dots, c_{2k-1} + 2)^+)$$

respectively. Then one can proceed by induction on studying the composition factors of $\mathcal{B}(c_2 - 2, c_3, \dots, c_{2k+1})^+$ or $\mathcal{B}(c_0, c_1, \dots, c_{2k-1} + 2)^+$ as above.

6. DECOMPOSITION OF $\mathcal{B}(\overline{\mathcal{O}})$

6.1. **The main Theorem.** We are in the position to state the main theorem of the manuscript:

Theorem 6.1.1. *Let \mathcal{O} be a nilpotent orbit satisfying Assumption 1.3.7, and*

$$(\mathcal{C}_{c'}, \epsilon_{c'}) \sqcup \cdots \sqcup (\mathcal{C}_{b'}, \epsilon_{b'}) \sqcup \cdots \sqcup (\mathcal{C}_{d'}, \epsilon_{d'})$$

be a parameter attached to $\mathcal{O}' \in \text{Norm}(\mathcal{O})$ with

$$(28) \quad (\mathcal{C}_{b'}, \epsilon_{b'}) := [b'_0(b'_1 b'_2)^- \cdots (b'_{2l-1} b'_{2l})^- b'_{2l+1}]^{\epsilon_{b'}}.$$

Then $\mathcal{B}(\overline{\mathcal{O}})$ consists of irreducible modules whose parameters satisfy the following condition:

$$\textbf{Suppose } b'_{2l+1} \in 2\mathbb{N}^+, \textbf{ then } \epsilon_{b'} = \text{sgn}(-1)^l.$$

In other words, when $b'_{2l+1} \in 2\mathbb{N}^+$, the sign of $\epsilon_{b'}$ is chosen such that the number of $-$'s in Equation (28) is even.

Remark 6.1.2.

- (1) Each $\mathcal{O}' \in \text{Norm}(\mathcal{O})$ contributes exactly one parameter to the composition series of $\mathcal{B}(\overline{\mathcal{O}})$. We denote $\Pi_{\mathcal{O}'}$ be the irreducible module corresponding to this parameter.
- (2) The composition factors of $\mathcal{B}(\overline{\mathcal{O}})$ are also composition factors of $\mathcal{B}(\mathcal{O})$ (given in Theorem 5.2.4). Furthermore, $\mathcal{B}(\mathcal{O}) = \mathcal{B}(\overline{\mathcal{O}})$ precisely when $\overline{\mathcal{O}}$ is normal, consistent with $R(\overline{\mathcal{O}}) = R(\mathcal{O})$ in this case. Also, when $\overline{\mathcal{O}}$ is not normal, the discrepancy between $R(\mathcal{O})$ and $R(\overline{\mathcal{O}})$ occurs at the level of diminutive K -types already. As a consequence, the multiplicities of the diminutive K -types in $R(\overline{\mathcal{O}})$ and $R(\mathcal{O})$ are explicitly computable (see Corollary 6.3.3 below). An example is computed in [Wo2, Example 5.2].

Example 6.1.3. Let $\mathcal{O} = (10, 8, 8, 4, 4, 2)$. The composition factors of $\mathcal{B}(\mathcal{O})$ are listed in Example 5.2.2. Among the 9 factors of $\mathcal{B}(\mathcal{O})$, only 4 of them show up in $\mathcal{B}(\overline{\mathcal{O}})$:

$$(29) \quad \begin{array}{l} [10(88)^-(44)^-2]^+ \\ [10(88)^-6]^- [22]^+ \qquad [10, 10]^+ [6(44)^-2]^- \\ [10, 10]^+ [66]^+ [22]^+ \end{array}$$

□

6.2. The Cyclic Submodule of $\mathcal{B}(\mathcal{O})$. Recall the notation from Definition 2.0.1 and Theorem 2.0.2. Corollary 3.1.6 establishes a (\mathfrak{g}, K) -module map with the spherical vector in its image,

$$\mathcal{B}(\overline{\mathcal{O}}) \longrightarrow \mathcal{B}(\mathcal{O}).$$

We will study the cyclic submodule (the submodule generated by the spherical vector) of $\mathcal{B}(\mathcal{O})$, which provides information on which composition factors show up in $\mathcal{B}(\overline{\mathcal{O}})$.

Proposition 6.2.1. *Let \mathcal{O} be a nilpotent orbit with chains*

$$\mathcal{C}(\mathcal{O}) = [c_0(c_1c_2) \dots (c_{2p-1}c_{2p})c_{2p+1}] \sqcup \dots \sqcup [d_0(d_1d_2) \dots (d_{2q-1}d_{2q})d_{2q+1}].$$

The submodule generated by the spherical vector in the induced module

$$(30) \quad I^G[(c_0c_{2p+1})^+(c_1c_2)^+ \dots (c_{2p-1}c_{2p})^+ \dots (d_0d_{2q+1})^+(d_1d_2)^+ \dots (d_{2q-1}d_{2q})^+]$$

contains all the factors in $\mathcal{B}(\mathcal{O})$.

The proof is in Section 8. Assuming its validity, consider the image of the composition of the following intertwining operators

$$(31) \quad \begin{aligned} & I^G[(c_0c_{2p+1})^+(c_1c_2)^+ \dots (c_{2p-1}c_{2p})^+, \dots (d_0d_{2q+1})^+(d_1d_2)^+ \dots (d_{2q-1}d_{2q})^+] \\ & \xrightarrow{\iota^c} I^G[(c_1c_2)^+ \dots (c_{2p-1}c_{2p})^+(c_0c_{2p+1})^+ \dots (d_0d_{2q+1})^+(d_1d_2)^+ \dots (d_{2q-1}d_{2q})^+] \\ & \rightarrow \dots \\ & \xrightarrow{\iota^d} I^G[(c_1c_2)^+ \dots, (c_{2p-1}c_{2p})^+(c_0c_{2p+1})^+ \dots (d_1d_2)^+ \dots (d_{2q-1}d_{2q})^+(d_0d_{2q+1})^+] \\ & \xrightarrow{\iota^{sn}} I^G[(c_1c_2)^+ \dots, (c_{2p-1}c_{2p})^+ \dots (d_1d_2)^+ \dots (d_{2q-1}d_{2q})^+(c_0c_{2p+1})^+ \dots (d_0d_{2q+1})^+] \\ & \xrightarrow{q} I^G[(c_1c_2)^+ \dots, (c_{2p-1}c_{2p})^+ \dots (d_1d_2)^+ \dots (d_{2q-1}d_{2q})^+ \overline{X}((c_0c_{2p+1})^+ \dots (d_0d_{2q+1})^+)] \\ & = \mathcal{B}(\mathcal{O}). \end{aligned}$$

where the intertwining operators

$$\iota^c := \iota_p^c \circ \dots \circ \iota_2^c \circ \iota_1^c, \quad \dots, \quad \iota^d := \iota_q^d \circ \dots \circ \iota_2^d \circ \iota_1^d$$

are studied in Section 5.3.2, and ι^{sn} is a composition of intertwining operators interchanging strongly nested Δ 's.

The initial module in Equation (31) is cyclic by Proposition 6.2.1. All maps in (31) are isomorphisms on the trivial K -type. So the cyclic submodule of $\mathcal{B}(\mathcal{O})$ is precisely the image of (31).

Proposition 6.2.2. *The composition factors of the cyclic submodule of $\mathcal{B}(\mathcal{O})$ are precisely the ones specified in Theorem 6.1.1.*

Proof. The images and kernels of the intertwining operators in Equation (31) were studied in detail in Section 5.3.2. Since ι^{sn} consists of operators involving strongly nested Δ 's, they are isomorphisms on the diminutive K -types. So we only need to study each ι^c, \dots, ι^d .

We will only focus on the image of ι^c when \mathcal{O}^c has a single chain $\mathcal{C}(\mathcal{O}^c) = [c_0(c_1c_2) \dots (c_{2p-1}c_{2p})c_{2p+1}]$. The general case of multiple chains follows directly from the description of composition factors of $\mathcal{B}(\mathcal{O})$ in Theorem 5.2.4.

Rather than looking at $\text{im } \iota^c = \text{im } (\iota_p^c \circ \dots \circ \iota_2^c \circ \iota_1^c)$ directly, we keep track of $\ker \iota_l^c$ defined in (23) separately. Namely, from the proof of Proposition 5.2.1 in Section 5.3.2, the factors of each $\ker \iota_l^c$ are of the form $\overline{X}(\mathcal{P}_1 \sqcup \mathcal{P}_2)$, where

$$\begin{aligned} \mathcal{P}_1 &\text{ is } [c_0(c_1c_2) \dots (c_{2l-3}c_{2l-2})c_{2l-1} + 2]^- && - \text{ admissible,} \\ \mathcal{P}_2 &\text{ is } [c_{2l} + 2(c_{2l+1}c_{2l+2}) \dots (c_{2p-1}c_{2p})c_{2p+1}]^- && - \text{ admissible.} \end{aligned}$$

It follows directly from the definition of \mathcal{C}^ϵ -admissible parameters (Definition 5.2.3) that the factors specified in Theorem 6.1.1 do *not* lie in any of the $\ker \iota$'s. So they must occur in $\text{im } \iota^c$.

The same argument shows that a parameter which contains a $(\mathcal{C}_{b'}, \epsilon_{b'})$ that ends with a coordinate in $2\mathbb{N}^+$ and has an odd number of minuses, must appear in some $\ker \iota_l^c$ for some l . \square

The next example illustrates $\ker \iota^c = \iota_2^c \circ \iota_1^c$ in the above proposition in a special case.

Example 6.2.3. *Let $\mathcal{O} = (10, 8, 8, 4, 4, 2)$ as before. The kernel of $\iota_2 \circ \iota_1$ is given in Example 5.3.3 by:*

$$\begin{aligned} \ker \iota_1 &: [10, 10]^- [6(44)^- 2]^+, [10, 10]^- [66]^- [22]^+, [10, 10]^- [66]^+ [22]^- . \\ \ker \iota_2 &: [10(88)^- 6]^+ [22]^- , [10, 10]^+ [66]^- [22]^- , [10, 10]^- [66]^+ [22]^- . \end{aligned}$$

These are precisely the parameters that appear in $\mathcal{B}(\mathcal{O})$ (Example 5.2.2) but do not show up in $\mathcal{B}(\overline{\mathcal{O}})$ (Example 6.1.3).

Corollary 6.2.4. *The factors listed in Theorem 6.1.1 all occur in the spherical cyclic submodule of $\mathcal{B}(\overline{\mathcal{O}})$.*

Proof. This follows directly from Proposition 6.2.2, and the non-trivial map $\mathcal{B}(\overline{\mathcal{O}}) \rightarrow \mathcal{B}(\mathcal{O})$ with the trivial K -type in its image. \square

6.3. Proof of Theorem 6.1.1. By Corollary 6.2.4, we only need to show there are no composition factors of $\mathcal{B}(\overline{\mathcal{O}})$ other than the ones given by Theorem 6.1.1.

The following proposition gives the multiplicities of the diminutive K -types in the sum of the factors in Theorem 6.1.1:

Proposition 6.3.1. *Let \mathcal{O} be a nilpotent orbit of Type C satisfying Assumption 1.3.7 with chains*

$$\mathcal{C}(\mathcal{O}) = [c_0(c_1c_2) \dots (c_{2p-1}c_{2p})c_{2p+1}] \sqcup \dots \sqcup [d_0(d_1d_2) \dots (d_{2q-1}d_{2q})d_{2q+1}].$$

For each $\mathcal{O}' \in \text{Norm}(\mathcal{O})$, let $\Pi_{\mathcal{O}'}$ be the irreducible modules specified in Theorem 6.1.1 (Remark 6.1.2). Then $\bigoplus_{\mathcal{O}' \in \text{Norm}(\mathcal{O})} \Pi_{\mathcal{O}'}$ has diminutive K -type multiplicities equal to

that of the induced module

$$\bigoplus_{\mathcal{O}' \in \text{Norm}(\mathcal{O})} \Pi_{\mathcal{O}'} \approx I^G((c_0c_1)^+ \dots (c_{2p}c_{2p+1})^+ \dots (d_0d_1)^+ \dots (d_{2q}d_{2q+1})^+).$$

First of all, we study the proposition for $\mathcal{C}(\mathcal{O}) = [c_0(c_1c_2)c_3]$:

Lemma 6.3.2. *Let $c_0 \geq c_1 = c_2 \geq c_3$ be even integers, and $c'_0 \geq c'_1 > c'_2 \geq c'_3$ be the fundamental collapse of $c_0 \geq c_1 = c_2 \geq c_3$ (Definition 4.1.1).*

(a) *The factors with diminutive lowest K -type of $I^G((c_0c_1)^+(c_2c_3)^+)$ are*

$$\overline{X}([c_0(c_1c_2)^-c_3]^\mu) \quad \text{and} \quad \overline{X}([c'_0c'_1]^+[c'_2c'_3]^+).$$

(b) *The factors with diminutive lowest K -type of $I^G((c_0c_1)^+(c_2c_3)^-)$ are*

$$\overline{X}([c_0(c_1c_2)^-c_3]^+) \quad \text{and} \quad \begin{cases} \overline{X}([c'_0c'_1]^+[c'_2c'_3]^\mu) & \text{if } c_2 \neq c_3 \\ 0 & \text{if } c_2 = c_3 \end{cases}.$$

Proof. The result is obvious when $c_1 = c_2 = c_3$.

When $c_0 = c_1 = c_2$ or $c_1 = c_2 = c_3$, the induced module in (a) is $\approx \mathcal{B}(c_0, c_1, c_2, c_3)^+$, so the results follow from Table (21).

When $c_0 = c_1 = c_2$, the induced module in (b) is $\approx \mathcal{B}(c_0, c_1, c_2, c_3)^-$, so Theorem 5.2.4(b)(ii) applies.

when $c_0 > c_1 = c_2 > c_3$, The arguments are exactly the same as in Section 5.3.2. we omit the details. \square

By Lemma 6.3.2(a), Proposition 6.3.1 holds for $\mathcal{C}(\mathcal{O}) = [c_0(c_1c_2)c_3]$. Indeed, the parameters of the two irreducible modules appearing in Lemma 6.3.2(a) are precisely $\Pi_{(c_0, c_1, c_2, c_3)}$ and $\Pi_{(c'_0, c'_1, c'_2, c'_3)}$ described in Theorem 6.1.1.

6.3.1. Proof of Proposition 6.3.1. We prove the proposition when \mathcal{O} has only one chain. The general result follows from the arguments in Section 5.4.

We do induction on the length of a chain. The initial step is given in Lemma 6.3.2.

Suppose Proposition 6.3.1 holds for

$$\mathcal{C}(\mathcal{O}_k) = [c_2(c_3c_4) \dots (c_{2k-1}c_{2k})c_{2k+1}].$$

We want to show it also holds for

$$\mathcal{C}(\mathcal{O}_{k+1}) = [c_0(c_1c_2)(c_3c_4) \dots (c_{2k-1}c_{2k})c_{2k+1}].$$

We make the relation between $Norm(\mathcal{O}_k)$ and $Norm(\mathcal{O}_{k+1})$ explicit. Let $\mathcal{O}'_k = (b_2, b_3, b_4, \dots, b_{2k+1}) \in Norm(\mathcal{O}_k)$. By Assumption 1.3.7, $c_2 \neq c_3$, so b_2 must be equal to c_2 . Consider the two orbits

$$(\mathcal{O}'_k)_u := (c_0, c_1, c_2, b_3, b_4, \dots, b_{2k+1}), \quad (\mathcal{O}'_k)_d := (c'_0, c'_1, c'_2, b_3, b_4, \dots, b_{2k+1}),$$

where $(\mathcal{O}'_k)_u$ is obtained by concatenating \mathcal{O}'_k with (c_0, c_1) on the left, and $(\mathcal{O}'_k)_d$ is obtained by performing the fundamental collapse on c_0, c_1, c_2, b_3 of $(\mathcal{O}'_k)_u$. Then

$$Norm(\mathcal{O}_{k+1}) = \bigcup_{\mathcal{O}'_k \in Norm(\mathcal{O}_k)} \{(\mathcal{O}'_k)_u, (\mathcal{O}'_k)_d\}.$$

In particular, the cardinality of $Norm(\mathcal{O}_{k+1})$ is twice of that of $Norm(\mathcal{O}_k)$. Let

$$\Pi_{\mathcal{O}'_k} = \overline{X}([b_2(b_3b_4)^- \dots (b_{2l-1}b_{2l})^- b_{2l+1}]^\epsilon \sqcup \dots)$$

be the irreducible module attached to $\mathcal{O}'_k \in Norm(\mathcal{O}_k)$ given in Theorem 6.1.1, where $\epsilon = +$ if l is even, and $\epsilon = -$ if l is odd. Consider the induced module

$$\begin{aligned} I^G((c_0c_1)^+ \Pi_{\mathcal{O}'_k}) &= I^G((c_0c_1)^+ \overline{X}([b_2(b_3b_4)^- \dots (b_{2l-1}b_{2l})^- b_{2l+1}]^\epsilon \sqcup \dots)) \\ (32) \quad &\approx I^G((b_3b_4)^- \dots (b_{2l-1}b_{2l})^- (c_0c_1)^+ (b_2b_{2l+1})^\epsilon \dots) \\ &\approx I^G((b_3b_4)^- \dots (b_{2l-1}b_{2l})^- \dots I^G((c_0c_1)^+ (b_2b_{2l+1})^\epsilon)). \end{aligned}$$

By applying Lemma 6.3.2 with $(c_0, c_1, b_2, b_{2l+1})$, the induced module in Equation (32) contains two diminutive lowest K -type modules

$$(33) \quad I^G((c_0c_1)^+ \Pi_{\mathcal{O}'_k}) \approx \Pi_{(\mathcal{O}'_k)_u} \oplus \Pi_{(\mathcal{O}'_k)_d},$$

where

$$\begin{aligned} \Pi_{(\mathcal{O}'_k)_u} &= \overline{X}([c_0(c_1c_2)^- (b_3b_4)^- \dots (b_{2l-1}b_{2l})^- b_{2l+1}]^{-\epsilon} \sqcup \dots), \\ \Pi_{(\mathcal{O}'_k)_d} &= \overline{X}([c'_0c'_1]^+ \sqcup [c'_2(b_3b_4)^- \dots (b_{2l-1}b_{2l})^- b_{2l+1}]^\epsilon \sqcup \dots). \end{aligned}$$

These are precisely the irreducible modules attached to the orbits $(\mathcal{O}'_k)_u, (\mathcal{O}'_k)_d \in \text{Norm}(\mathcal{O}_{k+1})$ as given in Theorem 6.1.1. Therefore, for diminutive K -type multiplicities,

$$\begin{aligned}
 \bigoplus_{\mathcal{O}'_{k+1} \in \text{Norm}(\mathcal{O}_{k+1})} \Pi_{\mathcal{O}'_{k+1}} &\approx \bigoplus_{\mathcal{O}'_k \in \text{Norm}(\mathcal{O}_k)} (\Pi_{(\mathcal{O}'_k)_u} \oplus \Pi_{(\mathcal{O}'_k)_d}) \\
 &\approx \bigoplus_{\mathcal{O}'_k \in \text{Norm}(\mathcal{O}_k)} I^G((c_0 c_1)^+ \Pi_{\mathcal{O}'_k}) \\
 &\approx I^G((c_0 c_1)^+ I^{Sp}((c_2 c_3)^+ \dots (c_{2k} c_{2k+1})^+)) \\
 &\approx I^G((c_0 c_1)^+ (c_2 c_3)^+ \dots (c_{2k} c_{2k+1})^+).
 \end{aligned}$$

The second equality comes from Equation (33), the third equality comes from induction hypothesis that $\bigoplus_{\mathcal{O}'_k \in \text{Norm}(\mathcal{O}_k)} \Pi_{\mathcal{O}'_k} \approx I^{Sp}((c_2 c_3)^+ \dots (c_{2k} c_{2k+1})^+)$, and the last equality comes from induction in stages. Hence the proposition follows. \square

Corollary 6.3.3. *Let \mathcal{O} be a nilpotent orbit give in 6.3.1. Then the sum of the factors appearing in Theorem 6.1.1 $\bigoplus_{\mathcal{O}' \in \text{Norm}(\mathcal{O})} \Pi_{\mathcal{O}'}$ has diminutive K -type multiplicities equal to that of $R(\overline{\mathcal{O}^\#})$, where*

$$\mathcal{O}^\# := \left(\frac{c_0 + c_1}{2}, \frac{c_0 + c_1}{2}, \dots, \frac{c_{2p} + c_{2p+1}}{2}, \frac{c_{2p} + c_{2p+1}}{2}, \dots, \right. \\
 \left. \frac{d_0 + d_1}{2}, \frac{d_0 + d_1}{2}, \dots, \frac{d_{2q} + d_{2q+1}}{2}, \frac{d_{2q} + d_{2q+1}}{2} \right).$$

Proof. By construction, $\mathcal{O}^\#$ is a Richardson orbit whose closure is normal (Theorem 1.3.5). Therefore the K -spectrum of $R(\overline{\mathcal{O}^\#}) = R(\mathcal{O}^\#)$ equals that of

$$\text{Ind}_{GL(\frac{c_0+c_1}{2}) \times \dots \times GL(\frac{c_{2p}+c_{2p+1}}{2}) \times \dots \times GL(\frac{d_0+d_1}{2}) \times \dots \times GL(\frac{d_{2q}+d_{2q+1}}{2})}^G (\text{triv}).$$

In turn this is isomorphic to the K -spectrum of

$$I^G((c_0 c_1)^+ \dots (c_{2p} c_{2p+1})^+ \dots (d_0 d_1)^+ \dots (d_{2q} d_{2q+1})^+).$$

\square

Proof of Theorem 6.1.1. We argue by contradiction. Suppose that $\mathcal{B}(\overline{\mathcal{O}})$ has a factor other than the $\Pi_{\mathcal{O}'}$ in Theorem 6.1.1. Let Ψ be one of such factor; its lowest K -type,

V_γ , must be diminutive by Theorem 4.3.1. Then

$$\begin{aligned}
(34) \quad [R(\overline{\mathcal{O}}) : V_\gamma] &= [\mathcal{B}(\overline{\mathcal{O}}) : V_\gamma] \\
&\geq [\Psi \oplus \bigoplus_{\mathcal{O}' \in \text{Norm}(\mathcal{O})} \Pi_{\mathcal{O}'} : V_\gamma] \\
&> [\bigoplus_{\mathcal{O}' \in \text{Norm}(\mathcal{O})} \Pi_{\mathcal{O}'} : V_\gamma] \\
&= [R(\overline{\mathcal{O}^\#}) : V_\gamma].
\end{aligned}$$

On the other hand, $\overline{\mathcal{O}} \hookrightarrow \overline{\mathcal{O}^\#}$ as G -varieties. This implies that there is a G -module surjection $R(\overline{\mathcal{O}^\#}) \twoheadrightarrow R(\overline{\mathcal{O}})$. In particular, $[R(\overline{\mathcal{O}}) : V] \leq [R(\overline{\mathcal{O}^\#}) : V]$ for **all** irreducible, finite-dimensional representations of G . This contradicts Equation (34), and consequently there are no composition factors of $\mathcal{B}(\overline{\mathcal{O}})$ other than the ones listed in Theorem 6.1.1. \square

7. FINAL REMARKS

7.1. Symplectic orbits with only even columns. All the results we have proved so far works only for nilpotent orbits satisfying Assumption 1.3.7 (which, roughly speaking, avoids a large number of columns with equal sizes). We would like to briefly mention how the results can be generalized to all symplectic orbits with even column sizes.

For any special symplectic orbit \mathcal{O} with even column sizes, we can separate as few column pairs (c_i, c_i) as possible such that:

$$(35) \quad \mathcal{O} = \mathcal{O}_1 \cup (c_1, c_1) \cup \cdots \cup (c_k, c_k),$$

where \mathcal{O}_1 satisfies Assumption 1.3.7. For example, if $\mathcal{O} = (8, 6, 6, 6, 6, 6, 4, 2, 2, 2, 2, 0)$, then $\mathcal{O} = (8, 6, 6, 6, 4, 2, 2, 0) \cup (6, 6) \cup (2, 2)$.

We claim that the composition factors of $\mathcal{B}(\overline{\mathcal{O}})$ are given by the irreducible factors of

$$I^G((c_1, c_1)^+ \cdots (c_k, c_k)^+ \Pi_{\mathcal{O}'_1}),$$

where $\Pi_{\mathcal{O}'_1}$ is the factor attached to the orbit $\mathcal{O}'_1 \in \text{Norm}(\mathcal{O}_1)$ given by Theorem 6.1.1. We will present an example below:

Example 7.1.1. Let $\mathcal{O} = (6, 4, 4, 4, 4, 2) = (6, 4, 4, 2) \cup (4, 4)$. As in Chapter 3 - Chapter 6, we study the intertwining operators

$$(36) \quad I^G((6, 2)^+(4, 4)^+(4, 4)^+) \rightarrow I^G((4, 4)^+(6, 2)^+(4, 4)^+) \rightarrow I^G((4, 4)^+(4, 4)^+(6, 2)^+).$$

These intertwining operators are induced from the Type A operators

$$I^{GL}((6, 2)^+(4, 4)^+(4, 4)^+) \rightarrow I^{GL}((4, 4)^+(6, 2)^+(4, 4)^+) \rightarrow I^{GL}((4, 4)^+(4, 4)^+(6, 2)^+).$$

Using Lemma 1.2.2, the image of the first map is $I^{GL}((6, 4)^+(4, 2)^+(4, 4)^+)$, which is irreducible since the strings are strongly nested. By Lemma 1.2.2 again, the irreducible module $I^{GL}((6, 4)^+(4, 2)^+(4, 4)^+)$ is not in the kernel of the second map, so the second map is an isomorphism in terms of diminutive K -types.

By induction of intertwining operators, the second map in Equation (36) has no kernel of diminutive K -types.

In general, for each (c_i, c_i) appearing in Equation (35), the intertwining operator involving

$$I^G(\cdots (*, *)^+(c_i, c_i)^+ \cdots) \rightarrow I^G(\cdots (c_i, c_i)^+ (*, *)^+ \cdots)$$

has no kernel of diminutive K -types. Therefore our claim on the composition factors of $\mathcal{B}(\overline{\mathcal{O}})$ follows from the same arguments in Chapter 4 - Chapter 6.

7.2. Symplectic orbits with only odd columns. We will mention the results for symplectic orbits \mathcal{O} with odd columns only.

In this case, the construction in Section 2 - Section 3 of $\mathcal{B}(\overline{\mathcal{O}})$ yield half-integral infinitesimal characters. However, the construction of $Norm(\mathcal{O})$ and irreducible parameters are exactly the same as in the even column cases, thanks to the Kazhdan-Lusztig conjecture (see Section 4.2 of [B1], for instance). The main results follow analogously as in the even column cases.

Example 7.2.1. Consider the orbit $(9, 7, 7, 3, 3, 1)$. Then

$$(37) \quad \mathcal{B}(\mathcal{O}) = \begin{array}{c} [99]_o^+ [5(33)_o^- 1]_o^- \\ [99]_o^- [5(33)_o^- 1]_o^+ \end{array} \quad \begin{array}{c} [9(77)_o^- 5]_o^+ [11]_o^- \\ [9(77)_o^- 5]_o^- [11]_o^+ \end{array}$$

$$\begin{array}{cc} [99]_o^+ [55]_o^+ [11]_o^+ & [99]_o^- [55]_o^+ [11]_o^- \\ [99]_o^- [55]_o^- [11]_o^+ & [99]_o^+ [55]_o^- [11]_o^- \end{array}$$

Note that the parameters expressed in the form $[c_0(c_1 c_2)_o^- \cdots (c_{2p-1} c_{2p})_o^- c_{2p+1}]_o^{\epsilon c}$ are different from the parameters $[c_0(c_1 c_2)^- \cdots (c_{2p-1} c_{2p})^- c_{2p+1}]^{\epsilon c}$ defined in Definition 4.2.1, in the sense that the former has half-integral infinitesimal character, while the latter has integral infinitesimal character.

On the other hand, the Brylinski model has composition factors

$$(38) \quad \mathcal{B}(\overline{\mathcal{O}}) = \begin{array}{c} [9(77)_o^- (33)_o^- 1]_o^+ \\ [99]_o^+ [5(33)_o^- 1]_o^- \\ [9(77)_o^- 5]_o^- [11]_o^+ \\ [99]_o^+ [55]_o^+ [11]_o^+ \end{array}$$

7.3. All symplectic nilpotent orbits. Let \mathcal{O} be any symplectic nilpotent orbit. In order to find the composition factors of $\mathcal{B}(\overline{\mathcal{O}})$, we divide \mathcal{O} into odd and even columns

$$\mathcal{O} = \mathcal{O}_{\text{even}} \cup \mathcal{O}_{\text{odd}}.$$

For example, the orbit $(12, 10, 10, 8, 7, 5, 5, 3, 2, 0)$ is written as $(12, 10, 10, 8, 2, 0) \cup (7, 5, 5, 3)$.

Since the parameters attached to $\mathcal{O}_{\text{even}}$ are integral, while those attached to \mathcal{O}_{odd} are half-integral, the intertwining operator

$$I^G(\cdots (a, b)^+(c, d)_o^+ \cdots) \rightarrow I^G(\cdots (c, d)_o^+(a, b)^+ \cdots)$$

is isomorphic on diminutive K -types (Lemma 1.2.2(i)). Consequently, in order to find the composition factors of $\mathcal{B}(\overline{\mathcal{O}})$, we only need to concatenate the results for $\mathcal{O}_{\text{even}}$ and \mathcal{O}_{odd} .

Example 7.3.1. *The orbit $\mathcal{O} = (12, 10, 10, 8, 7, 5, 5, 3, 2, 0)$ has*

$$\mathcal{O}_{\text{even}} = (12, 10, 10, 8, 2, 0), \quad \mathcal{O}_{\text{odd}} = (7, 5, 5, 3).$$

Using the notations in the previous sections,

$$(39) \quad \mathcal{B}(\mathcal{O}_{\text{even}}) = \begin{array}{l} [12(10, 10)^-8]^- [20]^+ \\ [12, 12]^+ [10, 10]^+ [20]^+ \\ [12, 12]^- [10, 10]^- [20]^+ \end{array}, \quad \mathcal{B}(\mathcal{O}_{\text{odd}}) = \begin{array}{l} [7(55)_o^-3]_o^- \\ [77]_o^+ [33]_o^+ \\ [77]_o^- [33]_o^- \end{array}.$$

Then $\mathcal{B}(\mathcal{O}) =$

$$(40) \quad \begin{array}{l} [12(10, 10)^-8]^- [7(55)_o^-3]_o^- [20]^+ \\ [12(10, 10)^-8]^- [77]_o^+ [33]_o^+ [20]^+ \quad [12, 12]^+ [10, 10]^+ [7(55)_o^-3]_o^- [20]^+ \\ [12(10, 10)^-8]^- [77]_o^- [33]_o^- [20]^+ \quad [12, 12]^- [10, 10]^- [7(55)_o^-3]_o^- [20]^+ \\ [12, 12]^+ [10, 10]^+ [77]_o^+ [33]_o^+ [20]^+ \\ [12, 12]^- [10, 10]^- [77]_o^+ [33]_o^+ [20]^+ \\ [12, 12]^+ [10, 10]^+ [77]_o^- [33]_o^- [20]^+ \\ [12, 12]^- [10, 10]^- [77]_o^- [33]_o^- [20]^+ \end{array}.$$

Also, Theorem 6.1.1 (and its generalization to orbits with odd columns) yields

$$(41) \quad \mathcal{B}(\overline{\mathcal{O}_{\text{even}}}) = \begin{array}{l} [12(10, 10)^-8]^- [20]^+ \\ [12, 12]^+ [10, 10]^+ [20]^+ \end{array}, \quad \mathcal{B}(\overline{\mathcal{O}_{\text{odd}}}) = \begin{array}{l} [7(55)_o^-3]_o^- \\ [77]_o^+ [33]_o^+ \end{array}.$$

Therefore, $\mathcal{B}(\overline{\mathcal{O}}) =$

$$(42) \quad \begin{aligned} & [12(10, 10)^- 8]^- [7(55)_o^- 3]^- [20]^+ \\ & [12(10, 10)^- 8]^- [77]_o^+ [33]_o^+ [20]^+ \quad [12, 12]^+ [10, 10]^+ [7(55)_o^- 3]^- [20]^+ \quad . \\ & [12, 12]^+ [10, 10]^+ [77]_o^+ [33]_o^+ [20]^+ \end{aligned}$$

7.4. Orthogonal nilpotent orbits. We will briefly mention how the results can be generalized to nilpotent orbits of Type B and Type D .

Let $\mathcal{Q} = (d_1, d_2, \dots, d_{2p+1})$ be a partition specified by its column sizes (we insist there is an *odd* number of columns by putting $d_{2p+1} = 0$ if necessary). Then \mathcal{Q} defines a nilpotent orbit of Type B or D iff $d_{2i} + d_{2i+1}$ is even. More explicitly, \mathcal{Q} defines an orbit of Type B (or Type D) if d_1 is odd (or even).

Since our underlying group is $O(k, \mathbb{C})$, we only have one nilpotent orbit corresponding to the very even partitions.

The special orbits are characterized by the following rule:

- Type B : If d_{2i} is even, then $d_{2i} = d_{2i+1}$.
- Type D : If d_{2i} is odd, then $d_{2i} = d_{2i+1}$.

Using the same techniques as in Chapter 4 - Chapter 6, the composition factors of $\mathcal{B}(\overline{\mathcal{Q}})$, in the orthogonal case, are given by the following rule:

- (1) Consider the *symplectic* orbit $\mathcal{O} := (d_1 + 2, d_1, \dots, d_{2p+1})$ by adding a column of size $d_1 + 2$ on the left.
- (2) Study the composition factors of $\mathcal{B}(\overline{\mathcal{O}})$, and denote them as $\Pi_{\mathcal{O}'}$ for all $\mathcal{O}' \in \text{Norm}(\mathcal{O})$.
- (3) The composition factors of $\mathcal{B}(\overline{\mathcal{Q}})$ are given by $\theta(\Pi_{\mathcal{O}'})$, where $\Pi_{\mathcal{O}'} \leftrightarrow \theta(\Pi_{\mathcal{O}'})$ are related under the theta correspondence $Sp(d_1 + 2 + \sum d_i) \times O(\sum d_i)$. The relationship between the Langlands parameters of $\Pi_{\mathcal{O}'}$ and $\theta(\Pi_{\mathcal{O}'})$ is given explicitly in Theorem 2.8 of [AB].

Example 7.4.1. Let $\mathcal{Q} = (6, 6, 2, 2, 0)$ be an orthogonal nilpotent orbit. We consider $\mathcal{O} = (8, 6, 6, 2, 2, 0)$ with

$$(43) \quad \begin{aligned} & [8(66)^- (22)^- 0]^+ \\ \mathcal{B}(\overline{\mathcal{O}}) = & [88]^+ [4(22)^- 0]^+ \quad [8(66)^- 4]^- . \\ & [88]^+ [44]^+ \end{aligned}$$

Then the Brylinski model $\mathcal{B}(\overline{\mathcal{Q}})$ is given by

$$(44) \quad \mathcal{B}(\overline{\mathcal{Q}}) = \begin{array}{c} [(66)^-(22)^-0]^+ \\ [8]^+[4(22)^-0]^+ \\ [8]^+[44]^+ \end{array} \quad [(66)^-4]^-.$$

For example, the composition factor of $\mathcal{B}(\overline{\mathcal{O}})$ with parameter

$$[8(66)^-4]^- = \begin{pmatrix} -2 & -1 & \dots & 2 & 3; & -1 & -0 & 1 & 2 & 3 & 4 \\ -3 & -2 & \dots & 1 & 2; & -2 & -1 & 0 & 1 & 3 & 4 \end{pmatrix}.$$

is in theta-correspondence with the composition factor of $\mathcal{B}(\overline{\mathcal{Q}})$ with parameter

$$[(66)^-4]^- = \begin{pmatrix} -2 & -1 & \dots & 2 & 3; & 0 & 1 & - \\ -3 & -2 & \dots & 1 & 2; & 0 & 1 & - \end{pmatrix}$$

8. PROOFS OF PROPOSITIONS 4.3.1 AND 6.2.1

8.1. Proof of Proposition 4.3.1. We will prove Proposition 4.3.1 for \mathcal{O} where the infinitesimal character λ in Equation (9) is integral. In other words, we require that \mathcal{O} contains columns of even sizes only. For the half-integral cases, one can obtain the same result using the Kazhdan-Lusztig conjecture (e.g. [B1, Section 4.2]). The results of this section can also be found in Chapter 7 of [Wo1].

Given the integral infinitesimal character (λ, λ) in Equation (9), we study the **left cone representation** corresponding to the orbit \mathcal{O}_λ as described in the beginning of Section 4:

$$\overline{V}^L(w_0w_\lambda) = \text{Ind}_{W(\lambda)}^W(\text{triv}),$$

where $W(\lambda)$ is the largest Weyl Levi subgroup of W fixing λ , and w_λ is the longest element in $W(\lambda)$.

For any special orbit \mathcal{O}' , the number of irreducible representations with infinitesimal character (λ, λ) and associated variety $\overline{\mathcal{O}'}$ is given by

$$[\overline{V}^L(w_0w_\lambda) : V_{\mathcal{O}'}] = [\text{Ind}_{W(\lambda)}^W(\text{triv}) : V_{\mathcal{O}'}],$$

where $V_{\mathcal{O}'}$ is the **left cell representation** corresponding to the orbit \mathcal{O}' . In particular, each $V_{\mathcal{O}'}$ contains $|\overline{A}(\mathcal{O}')|$ distinct irreducible representations of W (see [BV2], Proposition 5.28 for instance).

The decomposition of $\overline{V}^L(w_0w_\lambda) = \text{Ind}_{W(\lambda)}^W(\text{triv})$ into irreducible representations can be easily computed using the Robinson-Schensted algorithm. In particular, one can check that for any even \mathcal{O} and any $\mathcal{O}' \in \text{Norm}(\mathcal{O})$, each irreducible factor of $V_{\mathcal{O}'}$ appears in $\text{Ind}_{W(\lambda)}^W(\text{triv})$ exactly once.

Therefore, the number of irreducible modules with infinitesimal character (λ, λ) and associated variety $\overline{\mathcal{O}'}$ for each $\mathcal{O}' \in \text{Norm}(\mathcal{O})$ is equal to $|\overline{A}(\mathcal{O}')|$, which is precisely the number of parameters attached to \mathcal{O}' .

Example 8.1.1. Consider $\mathcal{O} = (6, 4, 4, 2, 2, 0)$ with $\lambda = (322111100)$. Then

$$\text{Norm}(\mathcal{O}) = \{\mathcal{O}, \mathcal{O}_1 := (6, 4, 4, 4), \mathcal{O}_2 := (6, 6, 2, 2, 2, 0), \mathcal{O}_\lambda := (6, 6, 3, 3)\}$$

and $W(\lambda) = W(A_0) \times W(A_1) \times W(A_3) \times W(C_2)$.

The left cell representation for each $\mathcal{O}' \in \text{Norm}(\mathcal{O})$ is given by the following:

- \mathcal{O} : (21×321) .
- \mathcal{O}_1 : $(22 \times 32), (43 \times 11)$.
- \mathcal{O}_2 : $(31 \times 311), (41 \times 211)$.
- \mathcal{O}_λ : $(32 \times 31), (42 \times 21)$.

Note that the Young diagrams are all described in terms of columns, i.e. $(43 \times 11) =$

$$\left(\begin{array}{|c|c|} \hline \square & \square \\ \hline \square & \square \\ \hline \square & \square \\ \hline \square & \\ \hline \end{array}, \begin{array}{|c|c|} \hline \square & \square \\ \hline \end{array} \right).$$

One can easily show that each irreducible representation shows up exactly once in $\text{Ind}_{W(\lambda)}^W(\text{triv})$. In other words, there is only one way to fill up the Young diagrams into semi-standard Young tableaux using the coordinates of λ with the first row of the left Young diagram filled with zeros. For example, the only way to fill up (43×11)

$$\text{is given by } \left(\begin{array}{|c|c|} \hline 0 & 0 \\ \hline 1 & 1 \\ \hline 2 & 2 \\ \hline 3 & \\ \hline \end{array}, \begin{array}{|c|c|} \hline 1 & 1 \\ \hline \end{array} \right).$$

For an arbitrary orbit \mathcal{O} with even columns, there is a systematic way of describing the left cell representations for all $\mathcal{O}' \in \text{Norm}(\mathcal{O})$ (c.f. [L, Chapter 4]) and the same results hold. We omit the details here.

We are left to show that all the modules $I_{\mathcal{O}', \epsilon'}$ corresponding to $\mathcal{O}' \in \text{Norm}(\mathcal{O})$ given in Proposition 4.3.1 are irreducible, have the correct associated variety and are of multiplicity one. The last two properties follow directly from the definition of $I_{\mathcal{O}', \epsilon'}$ in Equation (13), so we only need to check that they are irreducible.

We argue by induction on the size of $\mathcal{O}' \in \text{Norm}(\mathcal{O})$. For the smallest orbit $\mathcal{O}_\lambda \in \text{Norm}(\mathcal{O})$, its chains $\mathcal{C}(\mathcal{O}_\lambda)$ are all of length 2. Therefore, the modules $I_{\mathcal{O}_\lambda, \epsilon_\lambda}$ are all special unipotent representations, which are known to be irreducible.

Now consider any nilpotent orbit $\mathcal{O}' \in \text{Norm}(\mathcal{O})$. By induction hypothesis, we assume that all the modules $I_{\mathcal{O}'', \epsilon''}$ in Proposition 4.3.1 with parameters attached to $\mathcal{O}'' \subset \overline{\mathcal{O}'}$ are all irreducible.

Let $\overline{X_{\mathcal{O}', \epsilon'}}$ be the lowest K -type subquotient in the induced module $I_{\mathcal{O}', \epsilon'}$. By Remark 4.3.2, the lowest K -type of $\overline{X_{\mathcal{O}', \epsilon'}}$ is $V_{2z} := \underbrace{(1, \dots, 1, 0, \dots, 0)}_{2z}$ for some $z \in \mathbb{N}$.

For any $y \in \mathbb{N}$ with $y \geq z$, V_{2y} is a **m-bottom layer K-type** of $\overline{X_{\mathcal{O}', \epsilon'}}$ with

$$M = GL(c'_2) \times \cdots \times GL(c'_{2p}) \cdots \times GL(d'_2) \times \cdots \times GL(d'_{2q}) \times Sp$$

as given in Proposition 4.3.1.

Let $m := \sum_{i=1}^p c'_{2i} + \cdots + \sum_{j=1}^q d'_{2j}$. By Proposition 2.7 of [B1],

$$\begin{aligned} & [\overline{X_{\mathcal{O}', \epsilon'}} : V_{2y}]_G \\ &= [\det \otimes \cdots \otimes \det \otimes \overline{X}([c'_0 c'_{2p+1}]^{\epsilon_{c'}} \cdots [d'_0 d'_{2q+1}]^{\epsilon_{d'}}) : \det \otimes \cdots \otimes \det \otimes V_{2y-m}]_M \\ &= [\overline{X}([c'_0 c'_{2p+1}]^{\epsilon_{c'}} \cdots [d'_0 d'_{2q+1}]^{\epsilon_{d'}}) : V_{2y-m}]_{Sp} \end{aligned}$$

On the other hand, using Frobenius reciprocity, we have

$$[I_{\mathcal{O}', \epsilon'} : V_{2y}]_G = [\det \otimes \cdots \otimes \det \otimes \overline{X}([c'_0 c'_{2p+1}]^{\epsilon_{c'}} \cdots [d'_0 d'_{2q+1}]^{\epsilon_{d'}}) : V_{2y}|_M]_M.$$

Note that under the decomposition of $V_{2y}|_M = \det \otimes \cdots \otimes \det \otimes V_{2y-m} + \dots$ into irreducible modules, only the first factor gives non-zero multiplicity on the above equation. Hence

$$\begin{aligned} & [I_{\mathcal{O}', \epsilon'} : V_{2y}]_G \\ &= [\det \otimes \cdots \otimes \det \otimes \overline{X}([c'_0 c'_{2p+1}]^{\epsilon_{c'}} \cdots [d'_0 d'_{2q+1}]^{\epsilon_{d'}}) : \det \otimes \cdots \otimes \det \otimes V_{2y-m} + \dots]_M \\ &= [\overline{X}([c'_0 c'_{2p+1}]^{\epsilon_{c'}} \cdots [d'_0 d'_{2q+1}]^{\epsilon_{d'}}) : V_{2y-m}]_{Sp} + 0 \\ &= [\overline{X_{\mathcal{O}', \epsilon'}} : V_{2y}]_G. \end{aligned}$$

One can therefore check that the irreducible module $\overline{X_{\mathcal{O}', \epsilon'}}$ has different diminutive K -type multiplicities from $\overline{X_{\mathcal{O}'', \epsilon''}} = I_{\mathcal{O}'', \epsilon''}$ for any $\mathcal{O}'' \subsetneq \overline{\mathcal{O}'}$, hence $\overline{X_{\mathcal{O}', \epsilon'}}$ is different from all $\overline{X_{\mathcal{O}'', \epsilon''}}$, and has associated variety equal to $\overline{\mathcal{O}'}$.

Suppose on the contrary that W is a composition factor of $I_{\mathcal{O}', \epsilon}$ other than $\overline{X_{\mathcal{O}', \epsilon'}}$, then W must have associated variety $\overline{\mathcal{O}''} \subsetneq \overline{\mathcal{O}'}$ since $I_{\mathcal{O}', \epsilon}$ is of multiplicity one. Hence $W = \overline{X_{\mathcal{O}'', \epsilon''}} = I_{\mathcal{O}'', \epsilon''}$ for some ϵ'' . By Remark 4.3.2 again, W has diminutive lowest K -type V_{2w} with $w \geq z$. So the lowest K -type of W is a **m-bottom layer K-type**, and $[\overline{X_{\mathcal{O}', \epsilon'}} : V_{2w}] = [I_{\mathcal{O}', \epsilon'} : V_{2w}]$. Now we have

$$[I_{\mathcal{O}', \epsilon'} : V_{2w}] \geq [\overline{X_{\mathcal{O}', \epsilon'}} \oplus W : V_{2w}] = [I_{\mathcal{O}', \epsilon'} : V_{2w}] + [W : V_{2w}] = [I_{\mathcal{O}', \epsilon'} : V_{2w}] + 1,$$

which gives a contradiction. \square

8.2. Proof of Proposition 6.2.1. The proof of Proposition 6.2.1 is by induction on the number of chains. We give details for \mathcal{O} formed of two chains only, i.e.

$$\mathcal{C}(\mathcal{O}) = [c_0(c_1c_2) \dots (c_{2p-1}c_{2p})c_{2p+1}] \sqcup [d_0(d_1d_2) \dots (d_{2q-1}d_{2q})d_{2q+1}].$$

Rearrange the infinitesimal character given in Equation (9) into the form

$$\lambda = (c_0/2, c_0/2 - 1, \dots, \underbrace{r, \dots, r}_{n_r}, \dots, \underbrace{0, \dots, 0}_{n_0})$$

with $n_r \leq n_{r_1}$ and $n_1 \leq 2n_0 + 1$. This is the *dominant form* of the parameter. Properties of the standard intertwining operators (Propositions 1.2.4 - 1.2.5) imply that $I(\Delta(1, \frac{c_0}{2}), \lambda^1)$ is a homomorphic image of the cyclic spherical module $X(\lambda)$; so it is cyclic.

By induction, $I(\Delta(1, \frac{c_0}{2})\Delta(-\frac{c_2}{2} + 1, \frac{c_1}{2}), \lambda^2)$ is also a homomorphic image of $X(\lambda)$ (this can be seen from the arguments below too). Repeating the process with λ^2 , we find

$$I(\Delta(1, \frac{c_0}{2})(c_1c_2)^+ \dots (c_{2p-1}c_{2p})^+ \Delta(0, \frac{c_{2p+1}}{2} - 1), \lambda^3)$$

as a homomorphic image of $X(\lambda)$. Since $\Delta(0, \frac{c_{2p+1}}{2} - 1)$ include all coordinates λ_j^3 in λ^3 , i.e. $\Delta(0, \frac{c_{2p+1}}{2} - 1)$ and $\{\lambda_j^3\}$ are strongly nested, we can now pass $\Delta(0, \frac{c_{2p+1}}{2} - 1)$ past λ^3 to conclude that

$$(45) \quad I(\Delta(1, \frac{c_0}{2})(c_1c_2)^+ \dots (c_{2p-1}c_{2p})^+, \lambda^3, \overline{X}(\Delta(0, \frac{c_{2p+1}}{2} - 1)))$$

is a homomorphic image of $X(\lambda)$. Note that

$$\overline{X}(\Delta(0, \dots, \frac{c_{2p+1}}{2} - 1)) \approx I(\Delta(-\frac{c_{2p+1}}{2} + 1, 0)),$$

so we can replace \overline{X} by I in Equation (45). Next, we pass $\Delta(-\frac{c_{2p+1}}{2} + 1, 0)$ past λ^3 in (45). Note that d_0 is the largest coordinate in λ^3 .

Applying Proposition 1.2.4 again, the induced module

$$I(\Delta(1, \frac{c_0}{2})(c_1c_2)^+ \dots (c_{2p-1}c_{2p})^+ \Delta(-\frac{c_{2p+1}}{2} + 1, \frac{d_0}{2}), \lambda^4)$$

is a homomorphic image of $X(\lambda)$. The intertwining operator

$$I(\Delta(1, \frac{c_0}{2})(c_1c_2)^+ \dots (c_{2p-1}c_{2p})^+ \Delta(-\frac{c_{2p+1}}{2} + 1, \frac{d_0}{2}), \lambda^4) \rightarrow$$

$$I(\Delta(1, \frac{c_0}{2})\Delta(-\frac{c_{2p+1}}{2} + 1, \frac{d_0}{2})(c_1c_2)^+ \dots (c_{2p-1}c_{2p})^+, \lambda^4)$$

is an isomorphism, since the Δ 's are all strongly nested. Also, the intertwining operator

$$I(\Delta(1, \frac{c_0}{2})\Delta(-\frac{c_{2p+1}}{2} + 1, \frac{d_0}{2})(c_1c_2)^+ \dots (c_{2p-1}c_{2p})^+, \lambda^4) \rightarrow$$

$$I(\Delta(-\frac{c_{2p+1}}{2} + 1, \frac{d_0}{2})\Delta(1, \frac{c_0}{2})(c_1c_2)^+ \dots (c_{2p-1}c_{2p})^+, \lambda^4)$$

has image $I((c_0c_{2p+1})^+\Delta(1, \frac{d_0}{2})(c_1c_2)^+ \dots (c_{2p-1}c_{2p})^+, \lambda^4)$. By applying the intertwining operator on strongly nested Δ 's,

$$I((c_0c_{2p+1})^+\Delta(1, \frac{d_0}{2})(c_1c_2)^+ \dots (c_{2p-1}c_{2p})^+, \lambda^4) \rightarrow$$

$$I((c_0c_{2p+1})^+(c_1c_2)^+ \dots (c_{2p-1}c_{2p})^+\Delta(1, \frac{d_0}{2}), \lambda^4),$$

one can see the last induced module is a homomorphic image of $X(\lambda)$. Repeating the same arguments on $I(\Delta(1, \frac{d_0}{2}), \lambda^4)$, we conclude that the induced module (30) in the proposition is a homomorphic image of $X(\lambda)$. Since the trivial K -type generates $X(\lambda)$, so is the module in (30) and the result follows. \square

REFERENCES

- [AB] J. Adams, D. Barbasch *The Reductive Dual Pairs Correspondence for Complex Groups*, J. of Func. An. vol 132, 1995
- [B1] D. Barbasch *The unitary dual for complex classical Lie groups*, Invent. Math., vol 96, 1989, pp 103-176
- [B2] ——— *Unipotent representations for real reductive groups*, Proceedings of ICM 1990, Springer Verlag, Tokyo, 1991, pp. 769-777
- [B3] ——— *Orbital integrals of nilpotent orbits*, volume in honor of Harish-Chandra, Proc. Symp. Pure Math. 68 (2000) pp. 97-110.
- [B4] ——— *The unitary spherical spectrum for split classical groups*, Journal of Inst. of Math. Jussieu, 2010 9(2) pp 265-356
- [B5] ——— *Unipotent Representations and the Dual Pairs Correspondence*, to appear in volume in honor of Roger Howe
- [BV] N. Berline, M. Vergne *Fourier transforms of orbits of the coadjoint representation*, Representation theory of reductive groups, Birkhäuser-Boston, Progress in mathematics vol 40, 1983, pp. 53-69
- [BV1] D. Barbasch, D. Vogan *The local structure of characters*, J. Funct. Analysis, vol. 34 no. 1, 1980, pp. 27-55
- [BV2] D. Barbasch, D. Vogan *Unipotent representations of complex semisimple Lie groups*, Ann. of Math., 1985, vol 121, pp. 41-110
- [BV3] D. Barbasch, D. Vogan *Weyl group representations and nilpotent orbits*, Representation theory of reductive groups, Birkhäuser-Boston, Progress in mathematics vol 40, 1983, pp. 21-35
- [Br] R. Brylinski *Line bundles on the cotangent bundle of the flag variety*, Invent. Math., vol 113 Fasc. 1, 1993, pp. 1-20
- [BZ] I.N. Bernstein, A.V. Zelevensky *Induced representations of reductive p -adic groups. I*, Ann. Sci. cole Norm. Sup. (4) 10 (1977), no. 4, 441-472.
- [CM] D. Collingwood and W. M. McGovern, *Nilpotent orbits in semisimple Lie algebras*, Van Norstrand Reinhold Mathematics Series, 1993

- [G] W. Graham, *Functions on the universal cover of the principal nilpotent orbit*, Inv. Math. **108** (1992), 15-27.
- [HC] Harish-Chandra *Fourier transform on a semisimple Lie algebra II*, Amer. J. of Math., vol 79, 1957, pp. 193-257
- [HC1] Harish-Chandra *Harmonic analysis on semisimple Lie groups*, Bulletin AMS. vol 76, 1970, pp. 529-551
- [Kn] A. Knapp *Representation Theory of Semisimple Groups*, Princeton University Press, Princeton NJ, 1986
- [KnV] A. Knapp, D. Vogan *Cohomological induction and unitary representations*, Princeton University Press, Princeton NJ, 1995
- [KP] H-P. Kraft, C. Procesi *On the geometry of conjugacy classes in classical groups*, Comm. Math. Helv., 57, 1982, pp. 539-601
- [J] J. C. Jantzen *Nilpotent orbits in representation theory*, in volume Lie Theory, Lie algebras and Representations by J-P Anker and B. Orsted, Progress in Mathematics, Birkhäuser, 2004
- [L] G. Lusztig, *Characters of Reductive Groups over a Finite Field*, Ann. Math. Studies, vol. 107, Princeton University Press, 1984
- [LS] G. Lusztig, N. Spaltenstein *Induced unipotent classes*, J. London Math. Soc., vol. 19, 1979, pp. 41-52
- [McG] M. McGovern *Rings of regular functions on nilpotent orbits and their covers*, Invent. Math., 97, 1989, pp. 209-217
- [Pz] T. Przebinda *The dual correspondence of infinitesimal characters*, Colloq. Math. 70 (1996), pp. 93102
- [R] R. Rao *Orbital integrals in reductive groups*, Annals of Math., vol. 96 no. 3, 1972, pp. 505-510
- [Ro] W. Rossmann *Nilpotent orbital integrals in a real semisimple Lie algebra and representations of Weyl groups*, Operator algebras, unitary representations, enveloping algebras and invariant theory, Progress in Mathematics 92, Birkhäuser Boston, 1990, pp. 263-287
- [ScVi1] W. Schmid, K. Vilonen *Two geometric character formulas for reductive Lie groups*, J. Amer. math. Soc., vol. 11 no. 4, 1998, pp. 799-867
- [ScVi2] W. Schmid, K. Vilonen *Characteristic cycles and wave front cycles of representations of reductive groups*, Annals of Math. 151 (2000), pp. 1071 - 1118.
- [SV] B. Speh, D. Vogan *Reducibility of generalized principal series representations*, Acta Math. 145 (1980), pp. 227-299
- [V] D. Vogan *The Algebraic Structure of the Representations of Semisimple Lie Groups I*, Annals of Mathematics, Second Series, Vol. 109, No. 1 (Jan., 1979), pp. 1-60
- [V1] D. Vogan *The orbit method and primitive ideals for semisimple Lie algebras*, Canad. Math. Soc. Conf. Proc. Vol. 5 (1986), pp. 281-316
- [V2] D. Vogan *Associated varieties and unipotent representations*, Harmonic Analysis on reductive groups, Progress in Mathematics vol. 101, Birkhäuser, Boston-Basel-Berlin, 1991, pp. 315-388
- [V3] ——— *The method of coadjoint orbits for real reductive groups*, Representation Theory of Lie Groups, IAS/Park City Mathematics Series 8 (1999), pp. 179-238
- [W] N. Wallach *Invariant differential operators on a reductive Lie algebra and Weyl group representations*, J. Amer. Math. Soc., vol. 6 no. 4, 1993, pp. 779-816
- [Wo1] K. D. Wong, *Dixmier Algebras on Complex Classical Nilpotent Orbits and their Representation Theories*, Ph.D. Thesis, Cornell University, 2013
- [Wo2] K.D. Wong *Quantization of special symplectic nilpotent orbits and normality of their closures*, J. Algebra 462 (2016), pp. 37-53

DEPARTMENT OF MATHEMATICS, CORNELL UNIVERSITY
E-mail address: `barbasch@math.cornell.edu`

DEPARTMENT OF MATHEMATICS, CORNELL UNIVERSITY
E-mail address: `kayue.wong@cornell.edu`