

Level-zero van der Kallen modules and specialization of nonsymmetric Macdonald polynomials at $t = \infty^*$

Satoshi Naito

Department of Mathematics, Tokyo Institute of Technology,
2-12-1 Oh-okayama, Meguro-ku, Tokyo 152-8551, Japan
(e-mail: naito@math.titech.ac.jp)

and

Daisuke Sagaki

Institute of Mathematics, University of Tsukuba,
1-1-1 Tennodai, Tsukuba, Ibaraki 305-8571, Japan
(e-mail: sagaki@math.tsukuba.ac.jp)

Abstract

Let $\lambda \in P^+$ be a level-zero dominant integral weight, and w an arbitrary coset representative of minimal length for the cosets in W/W_λ , where W_λ is the stabilizer of λ in a finite Weyl group W . In this paper, we give a module $\mathbb{K}_w^-(\lambda)$ over the negative part of a quantum affine algebra whose graded character is identical to the specialization at $t = \infty$ of the nonsymmetric Macdonald polynomial $E_{w\lambda}(q, t)$ multiplied by a certain explicit finite product of rational functions of q of the form $(1 - q^{-r})^{-1}$ for a positive integer r . This module $\mathbb{K}_w^-(\lambda)$ (called a level-zero van der Kallen module) is defined to be the quotient module of the level-zero Demazure module $V_w^-(\lambda)$ by the sum of the submodules $V_z^-(\lambda)$ for all those coset representatives z of minimal length for the cosets in W/W_λ such that $z > w$ in the Bruhat order $<$ on W .

1 Introduction.

In our previous paper [NS], we computed the graded character $\text{gch } V_e^-(\lambda)$ of the Demazure submodule $V_e^-(\lambda)$ of a level-zero extremal weight module $V(\lambda)$ over the quantum affine algebra $U_q(\mathfrak{g}_{\text{af}})$ associated to a nontwisted affine Lie algebra \mathfrak{g}_{af} , where $\lambda \in P^+$ is a level-zero dominant integral weight and e is the identity element of the affine Weyl group W_{af} .

*Key words and phrases: semi-infinite Lakshmibai-Seshadri path, nonsymmetric Macdonald polynomial, extremal weight module

Mathematics Subject Classification 2010: Primary 17B37; Secondary 14N15, 14M15, 33D52, 81R10.

The main result of [NS] states that the graded character $\text{gch } V_e^-(\lambda)$ is identical to the specialization $E_{w_\circ\lambda}(q, 0)$ at $t = 0$ of the nonsymmetric Macdonald polynomial multiplied by the inverse of the finite product $\prod_{i \in I} \prod_{r=1}^{\langle \lambda, \alpha_i^\vee \rangle} (1 - q^{-r})$, where w_\circ is the longest element of the finite Weyl group $W \subset W_{\text{af}}$ and q denotes the formal exponential e^δ , with δ the null root of \mathfrak{g}_{af} . Also, in [NNS1], we computed the graded character $\text{gch } V_{w_\circ}^-(\lambda)$ of the Demazure submodule $V_{w_\circ}^-(\lambda)$ of $V(\lambda)$, and proved that it is identical to the specialization $E_{w_\circ\lambda}(q, \infty)$ at $t = \infty$ of the nonsymmetric Macdonald polynomial multiplied by the inverse of the same finite product as above. Moreover, in [NNS1], for an arbitrary element w of the finite Weyl group W , we obtained combinatorial formulas for the graded character $\text{gch } V_w^-(\lambda)$ of the Demazure submodule $V_w^-(\lambda)$ of $V(\lambda)$ and the specialization $E_{w\lambda}(q, \infty)$ at $t = \infty$ of the nonsymmetric Macdonald polynomial, described in terms of quantum Lakshmibai-Seshadri paths introduced in [LNS³3].

Independently, Feigin-Makedonskyi [FM] introduced a family of finite-dimensional modules (called generalized Weyl modules), indexed by the elements w of the finite Weyl group W , over the Iwahori subalgebra $\mathfrak{J} := \mathfrak{b} \oplus (z\mathbb{C}[z] \otimes \mathfrak{g})$ of the current algebra $\mathfrak{g}[z] := \mathbb{C}[z] \otimes \mathfrak{g}$ associated to the finite-dimensional simple Lie algebra $\mathfrak{g} \subset \mathfrak{g}_{\text{af}}$ with Borel subalgebra \mathfrak{b} , and proved that for the cases $w = e$ and $w = w_\circ$, their graded characters are identical to the specializations at $t = \infty$ and $t = 0$ of the nonsymmetric Macdonald polynomial $E_{w_\circ\lambda}(q, t)$, respectively. Here we mention that the graded character of a generalized Weyl module indexed by a general element $w \neq e, w_\circ$ of W is not identical to the specialization at $t = 0$ nor $t = \infty$ of a nonsymmetric Macdonald polynomial.

Afterward, in [Kat], for an arbitrary element w of the finite Weyl group W , Kato gave an algebro-geometric construction of a finite-dimensional \mathfrak{J} -module whose graded character is identical to the specialization $E_{w\lambda}(q, \infty)$ at $t = \infty$ of the nonsymmetric Macdonald polynomial. Based on this result, Feigin-Kato-Makedonskyi [FKM] gave an algebraic description of these \mathfrak{J} -modules (denoted by $\mathbb{U}_{\sigma(\lambda_-)}$, with λ_- antidominant and $\sigma \in W$) by generators and relations, which are similar to that of generalized Weyl modules given in [FM].

The purpose of this paper is to give a module (which we call a level-zero van der Kallen module) over the negative part $U_{\mathfrak{q}}^-(\mathfrak{g}_{\text{af}})$ of $U_{\mathfrak{q}}(\mathfrak{g}_{\text{af}})$ whose graded character is identical to the specialization $E_{w\lambda}(q, \infty)$ at $t = \infty$ of the nonsymmetric Macdonald polynomial multiplied by the inverse of the same product as above for an arbitrary $w \in W$; note that the Demazure submodule $V_w^-(\lambda)$ is a $U_{\mathfrak{q}}^-(\mathfrak{g}_{\text{af}})$ -submodule of the level-zero extremal weight module $V(\lambda)$. Also, we give an explicit description of the crystal basis of the level-zero van der Kallen module in terms of semi-infinite Lakshmibai-Seshadri paths (or, quantum Lakshmibai-Seshadri paths).

Let us explain our main result more precisely. Let $\lambda \in P^+$ be a level-zero dominant

integral weight. We set $S = S_\lambda := \{i \in I \mid \langle \lambda, \alpha_i^\vee \rangle = 0\}$, and let W^S denote the set of minimal-length coset representatives for the cosets in W/W_S , where $W_S := \langle s_i \mid i \in S \rangle$ is the subgroup of the finite Weyl group $W = \langle s_i \mid i \in I \rangle$; we denote by $w_\circ(S)$ the longest element of W_S . For $w \in W^S$, we set

$$\mathbb{K}_w^-(\lambda) := V_w^-(\lambda) \Big/ \sum_{z \in W^S, z > w} V_z^-(\lambda);$$

here we know from [NS] that $V_z^-(\lambda) \subset V_w^-(\lambda)$ for all $z \in W^S$ such that $z > w$ in the Bruhat order $<$ on the finite Weyl group W . We call the module $\mathbb{K}_w^-(\lambda)$ a level-zero van der Kallen module. Our main result is the following.

Theorem 1. *Let $\lambda \in P^+$ be a level-zero dominant integral weight, and let $w \in W^S$. Then the graded character $\text{gch } \mathbb{K}_w^-$ can be expressed as follows:*

$$\text{gch } \mathbb{K}_w^-(\lambda) = \left(\prod_{i \in I} \prod_{r=1}^{\langle \lambda, \alpha_i^\vee \rangle - \epsilon_i} (1 - q^{-r}) \right)^{-1} E_{w\lambda}(q, \infty),$$

where

$$\epsilon_i = \epsilon_i([w]) := \begin{cases} 1 & \text{if } [w]s_i > [w], \\ 0 & \text{if } [w]s_i < [w]. \end{cases}$$

Here, $[w] := ww_\circ(S) \in W$ denotes the maximal-length representative for the coset $wW_S \in W/W_S$.

Our proof of this theorem is a crystal-theoretic one, which is based on the formulas for the graded character $\text{gch } V_w^-(\lambda)$ and the specialization $E_{w\lambda}(q, \infty)$ at $t = \infty$ of the nonsymmetric Macdonald polynomial in terms of quantum Lakshmibai-Seshadri paths obtained in [NNS1]. In fact, the crystal basis of the level-zero van der Kallen module $\mathbb{K}_w^-(\lambda)$ can be realized as the set of those semi-infinite Lakshmibai-Seshadri paths π of shape λ whose final direction $\kappa(\pi)$ lies in the set

$$K_w^S := ((W^S)_{\text{af}})_{\succeq w} \setminus \bigcup_{z \in W^S, z > w} ((W^S)_{\text{af}})_{\succeq z},$$

where $((W^S)_{\text{af}})_{\succeq x} := \{y \in (W^S)_{\text{af}} \mid y \succeq x\}$ for $x \in (W^S)_{\text{af}}$, with $(W^S)_{\text{af}}$ the set of Peterson's coset representatives for the cosets in $W_{\text{af}}/(W_S)_{\text{af}}$ (for the notation, see Section 2). Also, in the course of the proof of Theorem 1, we obtain a non-recursive description of the (strange) subsets $\text{EQB}(w)$ for $w \in W$, introduced in [NNS1] and studied in [NNS2], in terms of the semi-infinite Bruhat graph (or, quantum Bruhat graph); see Section 4 for details. Here we should mention that in contrast to the arguments in [FKM], our proof works uniformly for all nontwisted affine Lie algebras and does not use the reduction to

the rank two cases, though the construction of the module $\mathbb{K}_w^-(\lambda)$ above is inspired by [FKM].

In Appendix A, we give another proof of Theorem 1, which is based on the recursive formula for the specialization at $t = \infty$ of the nonsymmetric Macdonald polynomials due to Cherednik-Orr ([CO]); we would like to thank a referee for suggesting this recursive proof. A key to this proof is Lemma A.2 (or rather equivalently, Proposition A.1). As a consequence of this lemma, we obtain a formula, which expresses the graded character of a Demazure submodule of a level-zero extremal weight module as an explicit sum of the graded characters of level-zero van der Kallen modules: for each $w \in W^S$,

$$\text{gch } V_w^-(\lambda) = \sum_{v \in W^S, v \geq w} \text{gch } \mathbb{K}_v^-(\lambda).$$

Finally, in Appendix B, we remark that the cyclic vector of (the classical limit, i.e., the limit at $q \rightarrow 1$ of) the level-zero van der Kallen module $\mathbb{K}_w^-(\lambda)$ for $w \in W^S$ satisfies the same relations (see Lemma B.1) as those for the negative-roots version of the \mathfrak{J} -modules $\mathbb{U}_{\sigma(\lambda_-)}$ for $\sigma \in W$ introduced in [FKM]. Because both of these modules have the same graded character (at least in simply-laced cases), they are in fact isomorphic if the notational convention is suitably adjusted. Hence level-zero van der Kallen modules $\mathbb{K}_w^-(\lambda)$ can be thought of as a quantum analog of the \mathfrak{J} -modules $\mathbb{U}_{\sigma(\lambda_-)}$ in [FKM].

This paper is organized as follows. In Section 2, we fix our notation for affine Lie algebras, and recall some basic facts about the (parabolic) semi-infinite Bruhat graph. Next, we briefly review fundamental results on level-zero extremal weight modules and their Demazure submodules. Also, we define level-zero van der Kallen modules and state our main result above. In Section 3, we review the realization of the crystal bases of level-zero extremal weight modules by semi-infinite Lakshmibai-Seshadri paths. In Section 4, we first recall some basic facts about the (parabolic) quantum Bruhat graph, and then review from [NNS1] a recursive description of the subsets $\text{EQB}(w) \subset W$, $w \in W$, which are needed in the formula for the specialization $E_{w\lambda}(q, \infty)$. Also, we obtain a condition for the final directions of semi-infinite Lakshmibai-Seshadri paths forming the crystal basis of a level-zero van der Kallen module. In Section 5, by using this condition, we give a proof of our main result above. In Appendix A, we give a recursive proof of our main result. In Appendix B, we mention some relations satisfied by the cyclic vector of the level-zero van der Kallen module.

Acknowledgments.

S.N. was partially supported by JSPS Grant-in-Aid for Scientific Research (B) 16H03920. D.S. was partially supported by JSPS Grant-in-Aid for Scientific Research (C) 15K04803.

2 Main result.

2.1 Affine Lie algebras.

Let \mathfrak{g} be a finite-dimensional simple Lie algebra over \mathbb{C} with Cartan subalgebra \mathfrak{h} . Denote by $\{\alpha_i^\vee\}_{i \in I}$ and $\{\alpha_i\}_{i \in I}$ the set of simple coroots and simple roots of \mathfrak{g} , respectively, and set $Q := \bigoplus_{i \in I} \mathbb{Z}\alpha_i$, $Q^+ := \sum_{i \in I} \mathbb{Z}_{\geq 0}\alpha_i$, and $Q^\vee := \bigoplus_{i \in I} \mathbb{Z}\alpha_i^\vee$, $Q^{\vee,+} := \sum_{i \in I} \mathbb{Z}_{\geq 0}\alpha_i^\vee$; for $\xi, \zeta \in Q^\vee$, we write $\xi \geq \zeta$ if $\xi - \zeta \in Q^{\vee,+}$. Let Δ , Δ^+ , and Δ^- be the set of roots, positive roots, and negative roots of \mathfrak{g} , respectively, with $\theta \in \Delta^+$ the highest root of \mathfrak{g} . For a root $\alpha \in \Delta$, we denote by α^\vee its dual root. We set $\rho := (1/2) \sum_{\alpha \in \Delta^+} \alpha$. Also, let ϖ_i , $i \in I$, denote the fundamental weights for \mathfrak{g} , and set

$$P := \bigoplus_{i \in I} \mathbb{Z}\varpi_i, \quad P^+ := \sum_{i \in I} \mathbb{Z}_{\geq 0}\varpi_i. \quad (2.1)$$

Let $\mathfrak{g}_{\text{af}} = (\mathbb{C}[z, z^{-1}] \otimes \mathfrak{g}) \oplus \mathbb{C}c \oplus \mathbb{C}d$ be the untwisted affine Lie algebra over \mathbb{C} associated to \mathfrak{g} , where c is the canonical central element, and d is the scaling element (or the degree operator), with Cartan subalgebra $\mathfrak{h}_{\text{af}} = \mathfrak{h} \oplus \mathbb{C}c \oplus \mathbb{C}d$. We regard an element $\mu \in \mathfrak{h}^* := \text{Hom}_{\mathbb{C}}(\mathfrak{h}, \mathbb{C})$ as an element of $\mathfrak{h}_{\text{af}}^*$ by setting $\langle \mu, c \rangle = \langle \mu, d \rangle := 0$, where $\langle \cdot, \cdot \rangle : \mathfrak{h}_{\text{af}}^* \times \mathfrak{h}_{\text{af}} \rightarrow \mathbb{C}$ denotes the canonical pairing of $\mathfrak{h}_{\text{af}}^* := \text{Hom}_{\mathbb{C}}(\mathfrak{h}_{\text{af}}, \mathbb{C})$ and \mathfrak{h}_{af} . Let $\{\alpha_i^\vee\}_{i \in I_{\text{af}}} \subset \mathfrak{h}_{\text{af}}$ and $\{\alpha_i\}_{i \in I_{\text{af}}} \subset \mathfrak{h}_{\text{af}}^*$ be the set of simple coroots and simple roots of \mathfrak{g}_{af} , respectively, where $I_{\text{af}} := I \sqcup \{0\}$; note that $\langle \alpha_i, c \rangle = 0$ and $\langle \alpha_i, d \rangle = \delta_{i0}$ for $i \in I_{\text{af}}$. Denote by $\delta \in \mathfrak{h}_{\text{af}}^*$ the null root of \mathfrak{g}_{af} ; recall that $\alpha_0 = \delta - \theta$. Also, let $\Lambda_i \in \mathfrak{h}_{\text{af}}^*$, $i \in I_{\text{af}}$, denote the fundamental weights for \mathfrak{g}_{af} such that $\langle \Lambda_i, d \rangle = 0$, and set

$$P_{\text{af}} := \left(\bigoplus_{i \in I_{\text{af}}} \mathbb{Z}\Lambda_i \right) \oplus \mathbb{Z}\delta \subset \mathfrak{h}_{\text{af}}^*, \quad P_{\text{af}}^0 := \{\mu \in P_{\text{af}} \mid \langle \mu, c \rangle = 0\}; \quad (2.2)$$

notice that $P_{\text{af}}^0 = P \oplus \mathbb{Z}\delta$, and that $\langle \mu, \alpha_0^\vee \rangle = -\langle \mu, \theta^\vee \rangle$ for $\mu \in P_{\text{af}}^0$. We remark that for each $i \in I$, ϖ_i is equal to $\Lambda_i - \langle \Lambda_i, c \rangle \Lambda_0$, which is called the level-zero fundamental weight in [Kas2].

Let $W := \langle s_i \mid i \in I \rangle$ and $W_{\text{af}} := \langle s_i \mid i \in I_{\text{af}} \rangle$ be the (finite) Weyl group of \mathfrak{g} and the (affine) Weyl group of \mathfrak{g}_{af} , respectively, where s_i is the simple reflection with respect to α_i for each $i \in I_{\text{af}}$. We denote by $\ell : W_{\text{af}} \rightarrow \mathbb{Z}_{\geq 0}$ the length function on W_{af} , whose restriction to W agrees with the one on W , by $e \in W \subset W_{\text{af}}$ the identity element, and by $w_0 \in W$ the longest element. Denote by \geq the (ordinary) Bruhat order on W . For each $\xi \in Q^\vee$, let $t_\xi \in W_{\text{af}}$ denote the translation in $\mathfrak{h}_{\text{af}}^*$ by ξ (see [Kac, Sect. 6.5]); for $\xi \in Q^\vee$, we have

$$t_\xi \mu = \mu - \langle \mu, \xi \rangle \delta \quad \text{if } \mu \in \mathfrak{h}_{\text{af}}^* \text{ satisfies } \langle \mu, c \rangle = 0. \quad (2.3)$$

Then, $\{t_\xi \mid \xi \in Q^\vee\}$ forms an abelian normal subgroup of W_{af} , in which $t_\xi t_\zeta = t_{\xi+\zeta}$ holds for $\xi, \zeta \in Q^\vee$. Moreover, we know from [Kac, Proposition 6.5] that

$$W_{\text{af}} \cong W \rtimes \{t_\xi \mid \xi \in Q^\vee\} \cong W \rtimes Q^\vee.$$

Denote by Δ_{af} the set of real roots of \mathfrak{g}_{af} , and by $\Delta_{\text{af}}^+ \subset \Delta_{\text{af}}$ the set of positive real roots; we know from [Kac, Proposition 6.3] that $\Delta_{\text{af}} = \{\alpha + n\delta \mid \alpha \in \Delta, n \in \mathbb{Z}\}$, and $\Delta_{\text{af}}^+ = \Delta^+ \sqcup \{\alpha + n\delta \mid \alpha \in \Delta, n \in \mathbb{Z}_{>0}\}$. For $\beta \in \Delta_{\text{af}}$, we denote by $\beta^\vee \in \mathfrak{h}_{\text{af}}$ its dual root, and $s_\beta \in W_{\text{af}}$ the corresponding reflection; if $\beta \in \Delta_{\text{af}}$ is of the form $\beta = \alpha + n\delta$ with $\alpha \in \Delta$ and $n \in \mathbb{Z}$, then $s_\beta = s_\alpha t_{n\alpha^\vee} \in W \rtimes Q^\vee$.

Finally, let $U_{\mathfrak{q}}(\mathfrak{g}_{\text{af}})$ (resp., $U'_{\mathfrak{q}}(\mathfrak{g}_{\text{af}})$) denote the quantized universal enveloping algebra over $\mathbb{C}(\mathfrak{q})$ associated to \mathfrak{g}_{af} (resp., $[\mathfrak{g}_{\text{af}}, \mathfrak{g}_{\text{af}}]$), with E_i and F_i , $i \in I_{\text{af}}$, the Chevalley generators corresponding to α_i and $-\alpha_i$, respectively. We denote by $U_{\mathfrak{q}}^-(\mathfrak{g}_{\text{af}})$ the negative part of $U_{\mathfrak{q}}(\mathfrak{g}_{\text{af}})$, that is, the $\mathbb{C}(\mathfrak{q})$ -subalgebra of $U_{\mathfrak{q}}(\mathfrak{g}_{\text{af}})$ generated by F_i , $i \in I_{\text{af}}$.

2.2 Parabolic semi-infinite Bruhat graph.

In this subsection, we take and fix an arbitrary subset $S \subset I$. We set $Q_S := \bigoplus_{i \in S} \mathbb{Z}\alpha_i$, $Q_S^\vee := \bigoplus_{i \in S} \mathbb{Z}\alpha_i^\vee$, $Q_S^{\vee,+} := \sum_{i \in S} \mathbb{Z}_{\geq 0}\alpha_i^\vee$, $\Delta_S := \Delta \cap Q_S$, $\Delta_S^\pm := \Delta^\pm \cap Q_S$, $W_S := \langle s_i \mid i \in S \rangle$, and $\rho_S := (1/2) \sum_{\alpha \in \Delta_S^+} \alpha$; we denote by $[\cdot]^S : Q^\vee \rightarrow Q_{I \setminus S}^\vee$ the projection from $Q^\vee = Q_{I \setminus S}^\vee \oplus Q_S^\vee$ onto $Q_{I \setminus S}^\vee$ with kernel Q_S^\vee . Let W^S denote the set of minimal-length coset representatives for the cosets in W/W_S ; we know from [BB, Sect. 2.4] that

$$W^S = \{w \in W \mid w\alpha \in \Delta^+ \text{ for all } \alpha \in \Delta_S^+\}. \quad (2.4)$$

For $w \in W$, we denote by $[w] = [w]^S \in W^S$ (resp., $[w] = [w]^S$) the minimal-length (resp., maximal-length) coset representative for the coset wW_S in W/W_S ; note that $[w] = [w]w_\circ(S)$, where $w_\circ(S)$ denotes the longest element of the subgroup W_S of W . Also, following [P] (see also [LS, Sect. 10]), we set

$$(\Delta_S)_{\text{af}} := \{\alpha + n\delta \mid \alpha \in \Delta_S, n \in \mathbb{Z}\} \subset \Delta_{\text{af}}, \quad (2.5)$$

$$(\Delta_S)_{\text{af}}^+ := (\Delta_S)_{\text{af}} \cap \Delta_{\text{af}}^+ = \Delta_S^+ \sqcup \{\alpha + n\delta \mid \alpha \in \Delta_S, n \in \mathbb{Z}_{>0}\}, \quad (2.6)$$

$$(W_S)_{\text{af}} := W_S \rtimes \{t_\xi \mid \xi \in Q_S^\vee\} = \langle s_\beta \mid \beta \in (\Delta_S)_{\text{af}}^+ \rangle, \quad (2.7)$$

$$(W^S)_{\text{af}} := \{x \in W_{\text{af}} \mid x\beta \in \Delta_{\text{af}}^+ \text{ for all } \beta \in (\Delta_S)_{\text{af}}^+\}; \quad (2.8)$$

if $S = \emptyset$, then $(W^\emptyset)_{\text{af}} = W_{\text{af}}$ and $(W_\emptyset)_{\text{af}} = \{e\}$. We know from [P] (see also [LS, Lemma 10.6]) that for each $x \in W_{\text{af}}$, there exist a unique $x_1 \in (W^S)_{\text{af}}$ and a unique $x_2 \in (W_S)_{\text{af}}$ such that $x = x_1 x_2$; let

$$\Pi^S : W_{\text{af}} \rightarrow (W^S)_{\text{af}}, \quad x \mapsto x_1, \quad (2.9)$$

denote the projection, where $x = x_1 x_2$ with $x_1 \in (W^S)_{\text{af}}$ and $x_2 \in (W_S)_{\text{af}}$.

Lemma 2.1.

(1) *It holds that*

$$\begin{cases} \Pi^S(w) = \lfloor w \rfloor & \text{for all } w \in W; \\ \Pi^S(xt_\xi) = \Pi^S(x)\Pi^S(t_\xi) & \text{for all } x \in W_{\text{af}} \text{ and } \xi \in Q^\vee; \end{cases} \quad (2.10)$$

in particular, $(W^S)_{\text{af}} = \{w\Pi^S(t_\xi) \mid w \in W^S, \xi \in Q^\vee\}$.

(2) *For each $\xi \in Q^\vee$, the element $\Pi^S(t_\xi) \in (W^S)_{\text{af}}$ is of the form: $\Pi^S(t_\xi) = ut_{\xi+\xi_1}$ for some $u \in W_S$ and $\xi_1 \in Q_S^\vee$.*

(3) *For $\xi, \zeta \in Q^\vee$, $\Pi^S(t_\xi) = \Pi^S(t_\zeta)$ if and only if $\xi - \zeta \in Q_S^\vee$.*

Proof. Part (1) follows from [LS, Proposition 10.10], and part (2) follows from [LNS³1, (3.7)]. The “if” part of part (3) is obvious by part (1) and the fact that $t_{\xi-\zeta} \in (W_S)_{\text{af}}$. The “only if” part of part (3) is obvious by part (2). \square

Definition 2.2. Let $x \in W_{\text{af}}$, and write it as $x = wt_\xi$ with $w \in W$ and $\xi \in Q^\vee$. We define the semi-infinite length $\ell^{\frac{\infty}{2}}(x)$ of x by: $\ell^{\frac{\infty}{2}}(x) = \ell(w) + 2\langle \rho, \xi \rangle$.

Definition 2.3 ([Lu1], [Lu2]; see also [P]).

- (1) The (parabolic) semi-infinite Bruhat graph $\text{BG}^{\frac{\infty}{2}}((W^S)_{\text{af}})$ is the Δ_{af}^+ -labeled directed graph whose vertices are the elements of $(W^S)_{\text{af}}$, and whose directed edges are of the form: $x \xrightarrow{\beta} y$ for $x, y \in (W^S)_{\text{af}}$ and $\beta \in \Delta_{\text{af}}^+$ such that $y = s_\beta x$ and $\ell^{\frac{\infty}{2}}(y) = \ell^{\frac{\infty}{2}}(x) + 1$. When $S = \emptyset$, we write $\text{BG}^{\frac{\infty}{2}}(W_{\text{af}})$ for $\text{BG}^{\frac{\infty}{2}}((W^\emptyset)_{\text{af}})$.
- (2) The (parabolic) semi-infinite Bruhat order is a partial order \preceq on $(W^S)_{\text{af}}$ defined as follows: for $x, y \in (W^S)_{\text{af}}$, we write $x \preceq y$ if there exists a directed path in $\text{BG}^{\frac{\infty}{2}}((W^S)_{\text{af}})$ from x to y ; we write $x \prec y$ if $x \preceq y$ and $x \neq y$.

Remark 2.4. In the case $S = \emptyset$, the semi-infinite Bruhat order on W_{af} is essentially the same as the generic Bruhat order introduced in [Lu1]; see [INS, Appendix A.3] for details. Also, for a general S , the parabolic semi-infinite Bruhat order on $(W^S)_{\text{af}}$ is essentially the same as the partial order on S -alcoves introduced in [Lu2] when we take a special point to be the origin.

Remark 2.5. It follows from the definition that the restriction of the semi-infinite Bruhat order \preceq on $(W^S)_{\text{af}}$ to $W^S \subset (W^S)_{\text{af}}$ agrees with the (ordinary) Bruhat order \geq on W^S .

In Section 4.2 below, we recall some of the basic properties of the semi-infinite Bruhat order.

2.3 Crystal bases of extremal weight modules.

In this subsection, we fix $\lambda \in P^+ \subset P_{\text{af}}^0$ (see (2.1) and (2.2)). Let $V(\lambda)$ denote the extremal weight module of extremal weight λ over $U_{\mathfrak{q}}(\mathfrak{g}_{\text{af}})$, which is an integrable $U_{\mathfrak{q}}(\mathfrak{g}_{\text{af}})$ -module generated by a single element v_λ with the defining relation that v_λ is an extremal weight vector of weight λ ; recall from [Kas2, Sect. 3.1] and [Kas3, Sect. 2.6] that v_λ is an extremal weight vector of weight λ if (v_λ is a weight vector of weight λ and) there exists a family $\{v_x\}_{x \in W_{\text{af}}}$ of weight vectors in $V(\lambda)$ such that $v_e = v_\lambda$, and such that for every $i \in I_{\text{af}}$ and $x \in W_{\text{af}}$ with $n := \langle x\lambda, \alpha_i^\vee \rangle \geq 0$ (resp., ≤ 0), the equalities $E_i v_x = 0$ and $F_i^{(n)} v_x = v_{s_i x}$ (resp., $F_i v_x = 0$ and $E_i^{(-n)} v_x = v_{s_i x}$) hold, where for $i \in I_{\text{af}}$ and $k \in \mathbb{Z}_{\geq 0}$, the $E_i^{(k)}$ and $F_i^{(k)}$ are the k -th divided powers of E_i and F_i , respectively; note that v_x is an extremal weight vector of weight $x\lambda$. We know from [Kas1, Proposition 8.2.2] that $V(\lambda)$ has a crystal basis $(\mathcal{L}(\lambda), \mathcal{B}(\lambda))$ and the corresponding global basis $\{G(b) \mid b \in \mathcal{B}(\lambda)\}$; we denote by u_λ the element of $\mathcal{B}(\lambda)$ such that $G(u_\lambda) = v_\lambda$. It follows from [Kas1, Sect. 7] that the affine Weyl group W_{af} acts on $\mathcal{B}(\lambda)$ by

$$s_i \cdot b := \begin{cases} f_i^n b & \text{if } n := \langle \text{wt } b, \alpha_i^\vee \rangle \geq 0, \\ e_i^{-n} b & \text{if } n := \langle \text{wt } b, \alpha_i^\vee \rangle \leq 0, \end{cases} \quad (2.11)$$

for $b \in \mathcal{B}(\lambda)$ and $i \in I_{\text{af}}$.

We know the following from [Kas2] (see also [NS, Sect. 5.2]). Let $i \in I$.

- (i) The crystal graph of $\mathcal{B}(\varpi_i)$ is connected, and $\mathcal{B}(\varpi_i)_{\varpi_i + k\delta} = \{u_{\varpi_i + k\delta}\}$ for all $k \in \mathbb{Z}$, where $u_{\varpi_i + k\delta} := t_{-k\alpha_i^\vee} \cdot u_{\varpi_i}$ for $k \in \mathbb{Z}$. Therefore, $\dim V(\varpi_i)_{\varpi_i + k\delta} = 1$ for all $k \in \mathbb{Z}$.
- (ii) There exists a $U'_{\mathfrak{q}}(\mathfrak{g}_{\text{af}})$ -module automorphism $z_i : V(\varpi_i) \rightarrow V(\varpi_i)$ that maps v_{ϖ_i} to $v_{\varpi_i + \delta} := G(u_{\varpi_i + \delta})$; thus, this map commutes with the Kashiwara operators e_j, f_j , $j \in I_{\text{af}}$, on $V(\varpi_i)$.
- (iii) We have $z_i(\mathcal{L}(\varpi_i)) \subset \mathcal{L}(\varpi_i)$. Hence the map $z_i : V(\varpi_i) \rightarrow V(\varpi_i)$ induces a \mathbb{C} -linear automorphism $z_i : \mathcal{L}(\varpi_i)/\mathfrak{q}\mathcal{L}(\varpi_i) \rightarrow \mathcal{L}(\varpi_i)/\mathfrak{q}\mathcal{L}(\varpi_i)$; this induced map commutes with the Kashiwara operators e_j, f_j , $j \in I_{\text{af}}$, on $\mathcal{L}(\varpi_i)/\mathfrak{q}\mathcal{L}(\varpi_i)$, and satisfies $z_i(u_{\varpi_i}) = u_{\varpi_i + \delta}$. Therefore, the map z_i preserves $\mathcal{B}(\varpi_i)$.

Let us write $\lambda \in P^+$ as $\lambda = \sum_{i \in I} m_i \varpi_i$, with $m_i \in \mathbb{Z}_{\geq 0}$ for $i \in I$. We fix an arbitrary total order on I , and then set $\tilde{V}(\lambda) := \bigotimes_{i \in I} V(\varpi_i)^{\otimes m_i}$. By [BN, Eq. (4.8) and Corollary 4.15], there exists an injective $U_{\mathfrak{q}}(\mathfrak{g}_{\text{af}})$ -module homomorphism $\Phi_\lambda : V(\lambda) \hookrightarrow \tilde{V}(\lambda)$ that maps v_λ to $\tilde{v}_\lambda := \bigotimes_{i \in I} v_{\varpi_i}^{\otimes m_i}$. For each $i \in I$ and $1 \leq k \leq m_i$, we define $z_{i,k}$ to be the $U'_{\mathfrak{q}}(\mathfrak{g}_{\text{af}})$ -module automorphism of $\tilde{V}(\lambda)$ which acts as z_i only on the k -th factor of $V(\varpi_i)^{\otimes m_i}$ in $\tilde{V}(\lambda)$, and as the identity map on the other factors of $\tilde{V}(\lambda)$; these $z_{i,k}$'s, $i \in I$, $1 \leq k \leq m_i$,

commute with each other. We define

$$\overline{\text{Par}(\lambda)} := \{ \boldsymbol{\rho} = (\rho^{(i)})_{i \in I} \mid \rho^{(i)} \text{ is a partition of length } \leq m_i \text{ for each } i \in I \}. \quad (2.12)$$

For $\boldsymbol{\rho} = (\rho^{(i)})_{i \in I} \in \overline{\text{Par}(\lambda)}$, we set

$$s_{\boldsymbol{\rho}}(z^{-1}) := \prod_{i \in I} s_{\rho^{(i)}}(z_{i,1}^{-1}, \dots, z_{i,m_i}^{-1}). \quad (2.13)$$

Here, for a partition $\chi = (\chi_1 \geq \dots \geq \chi_m)$ of length less than or equal to $m \in \mathbb{Z}_{\geq 0}$, $s_{\chi}(x) = s_{\chi}(x_1, \dots, x_m)$ denotes the Schur polynomial in the variables x_1, \dots, x_m corresponding to the partition χ . We can easily show (see [NS, Sect. 7.3]) that $s_{\boldsymbol{\rho}}(z^{-1})(\text{Image } \Phi_{\lambda}) \subset \text{Image } \Phi_{\lambda}$ for each $\boldsymbol{\rho} = (\rho^{(i)})_{i \in I} \in \overline{\text{Par}(\lambda)}$. Hence we can define a $U'_{\mathfrak{q}}(\mathfrak{g}_{\text{af}})$ -module homomorphism $z_{\boldsymbol{\rho}} : V(\lambda) \rightarrow V(\lambda)$ in such a way that the following diagram commutes:

$$\begin{array}{ccc} V(\lambda) & \xrightarrow{\Phi_{\lambda}} & \tilde{V}(\lambda) \\ z_{\boldsymbol{\rho}} \downarrow & & \downarrow s_{\boldsymbol{\rho}}(z^{-1}) \\ V(\lambda) & \xrightarrow{\Phi_{\lambda}} & \tilde{V}(\lambda); \end{array} \quad (2.14)$$

note that $z_{\boldsymbol{\rho}}v_{\lambda} = S_{\boldsymbol{\rho}}^{-}v_{\lambda}$ in the notation of [BN] (and [NS]). Here, recall that $\tilde{V}(\lambda)$ has the crystal basis $(\tilde{\mathcal{L}}(\lambda) := \bigotimes_{i \in I} \mathcal{L}(\varpi_i)^{\otimes m_i}, \tilde{\mathcal{B}}(\lambda) := \bigotimes_{i \in I} \mathcal{B}(\varpi_i)^{\otimes m_i})$. We see from part (iii) above that the map $s_{\boldsymbol{\rho}}(z^{-1})$ preserves $\tilde{\mathcal{L}}(\lambda)$. Also, we know from [BN, page 369, the 2nd line from below] that $\Phi_{\lambda}(\mathcal{L}(\lambda)) \subset \tilde{\mathcal{L}}(\lambda)$. Therefore, we deduce that the map $z_{\boldsymbol{\rho}} : V(\lambda) \rightarrow V(\lambda)$ preserves $\mathcal{L}(\lambda)$, and hence induces a \mathbb{C} -linear map $z_{\boldsymbol{\rho}} : \mathcal{L}(\lambda)/\mathfrak{q}\mathcal{L}(\lambda) \rightarrow \mathcal{L}(\lambda)/\mathfrak{q}\mathcal{L}(\lambda)$; this map commutes with the Kashiwara operators. It follows from [BN, p. 371] that

$$\mathcal{B}(\lambda) = \{ z_{\boldsymbol{\rho}}b \mid \boldsymbol{\rho} \in \overline{\text{Par}(\lambda)}, b \in \mathcal{B}_0(\lambda) \}, \quad (2.15)$$

where $\mathcal{B}_0(\lambda)$ denotes the connected component of $\mathcal{B}(\lambda)$ containing u_{λ} , and

$$\text{Par}(\lambda) := \{ \boldsymbol{\rho} = (\rho^{(i)})_{i \in I} \mid \rho^{(i)} \text{ is a partition of length } < m_i \text{ for each } i \in I \}; \quad (2.16)$$

we understand that a partition of length less than 0 is the empty partition \emptyset . For $\boldsymbol{\rho} \in \text{Par}(\lambda)$, we set

$$u^{\boldsymbol{\rho}} := z_{\boldsymbol{\rho}}u_{\lambda} \in \mathcal{B}(\lambda). \quad (2.17)$$

Remark 2.6. We see from [BN, Theorem 4.16(ii)] (see also the argument after [NS, (7.3.8)]) that $z_{\boldsymbol{\rho}}G(b) = G(z_{\boldsymbol{\rho}}b)$ for $b \in \mathcal{B}_0(\lambda)$ and $\boldsymbol{\rho} \in \overline{\text{Par}(\lambda)}$.

2.4 Level-zero van der Kallen modules and their graded characters.

Let $\lambda \in P^+ \subset P_{\text{af}}^0$, and set

$$S = S_{\lambda} := \{ i \in I \mid \langle \lambda, \alpha_i^{\vee} \rangle = 0 \} \subset I. \quad (2.18)$$

Let $\{v_x\}_{x \in W_{\text{af}}}$ be the family of extremal weight vectors in $V(\lambda)$ corresponding to v_λ (see Section 2.3). For each $x \in W_{\text{af}}$, we define the Demazure submodule $V_x^-(\lambda)$ of $V(\lambda)$ by

$$V_x^-(\lambda) := U_{\mathfrak{q}}^-(\mathfrak{g}_{\text{af}})v_x \subset V(\lambda). \quad (2.19)$$

We see that the Demazure submodule $V_x^-(\lambda)$ has the $(\mathfrak{h}_{\text{af}}^-)$ -weight space decomposition of the form:

$$V_x^-(\lambda) = \bigoplus_{k \in \mathbb{Z}} \left(\bigoplus_{\gamma \in Q} V_x^-(\lambda)_{\lambda + \gamma + k\delta} \right), \quad (2.20)$$

where each weight space $V_x^-(\lambda)_{\lambda + \gamma + k\delta}$ is finite-dimensional. Also, we know from [Kas3, Sect. 2.8] (see also [NS, Sect. 4.1]) that $V_x^-(\lambda)$ is compatible with the global basis of $V(\lambda)$, that is, there exists a subset $\mathcal{B}_x^-(\lambda)$ of the crystal basis $\mathcal{B}(\lambda)$ such that

$$V_x^-(\lambda) = \bigoplus_{b \in \mathcal{B}_x^-(\lambda)} \mathbb{C}(\mathfrak{q})G(b) \subset \bigoplus_{b \in \mathcal{B}(\lambda)} \mathbb{C}(\mathfrak{q})G(b) = V(\lambda); \quad (2.21)$$

by [NNS1, Lemma 5.4.1], we have

$$\mathcal{B}_x^-(\lambda) = \{z_\rho b \mid \rho \in \text{Par}(\lambda), b \in \mathcal{B}_x^-(\lambda) \cap \mathcal{B}_0(\lambda)\}. \quad (2.22)$$

Remark 2.7. By [NS, Lemma 4.1.2], we have $V_x^-(\lambda) = V_{\Pi^S(x)}^-(\lambda)$ for $x \in W_{\text{af}}$. Also, it follows from [NS, Lemma 5.2.3] that for $x, y \in (W^S)_{\text{af}}$,

$$V_y^-(\lambda) \subset V_x^-(\lambda) \iff \mathcal{B}_y^-(\lambda) \subset \mathcal{B}_x^-(\lambda) \iff y \succeq x. \quad (2.23)$$

Now, for $w \in W^S$, we define a quotient $U_{\mathfrak{q}}^-(\mathfrak{g}_{\text{af}})$ -module (level-zero van der Kallen module) of $V_w^-(\lambda)$ by

$$\mathbb{K}_w^-(\lambda) := V_w^-(\lambda) \Big/ \sum_{z \in W^S, z > w} V_z^-(\lambda); \quad (2.24)$$

note that $V_z^-(\lambda) \subset V_w^-(\lambda)$ for all $z \in W^S$ such that $z > w$ (see (2.23) and Remark 2.5). Also, for each $w \in W^S$, we define another quotient $U_{\mathfrak{q}}^-(\mathfrak{g})$ -module of $V_w^-(\lambda)$ by

$$\mathbf{K}_w^-(\lambda) := V_w^-(\lambda) \Big/ \left(\sum_{z \in W^S, z > w} V_z^-(\lambda) + \sum_{\rho \in \overline{\text{Par}(\lambda)}, \rho \neq (\emptyset)_{i \in I}} z_\rho V_w^-(\lambda) \right), \quad (2.25)$$

where $z_\rho : V(\lambda) \rightarrow V(\lambda)$ is as in (2.14). Here we know from [NNS1, (5.15) and (5.16)] that if we set

$$X_w^-(\lambda) := \sum_{\rho \in \overline{\text{Par}(\lambda)}, \rho \neq (\emptyset)_{i \in I}} z_\rho V_w^-(\lambda), \quad (2.26)$$

then $X_w^-(\lambda) = \bigoplus_{b \in \mathcal{B}(X_w^-(\lambda))} \mathbb{C}(\mathfrak{q})G(b)$, where

$$\mathcal{B}(X_w^-(\lambda)) := \{z_\rho b \mid \rho \in \text{Par}(\lambda), \rho \neq (\emptyset)_{i \in I}, b \in \mathcal{B}_w^-(\lambda) \cap \mathcal{B}_0(\lambda)\}. \quad (2.27)$$

By (2.22), we see that $\mathcal{B}(X_w^-(\lambda)) \subset \mathcal{B}_w^-(\lambda)$, and hence $X_w^-(\lambda) \subset V_w^-(\lambda)$. Observe that $\mathbb{K}_w^-(\lambda)$ and $\mathbf{K}_w^-(\lambda)$ have the $(\mathfrak{h}_{\text{af}}^-)$ -weight space decompositions induced by that of $V_w^-(\lambda)$ (see (2.20)):

$$\mathbb{K}_w^-(\lambda) = \bigoplus_{k \in \mathbb{Z}} \left(\bigoplus_{\gamma \in Q} \mathbb{K}_w^-(\lambda)_{\lambda + \gamma + k\delta} \right), \quad \mathbf{K}_w^-(\lambda) = \bigoplus_{k \in \mathbb{Z}} \left(\bigoplus_{\gamma \in Q} \mathbf{K}_w^-(\lambda)_{\lambda + \gamma + k\delta} \right).$$

By putting $q := e^\delta$, we define

$$\text{gch } \mathbb{K}_w^-(\lambda) := \sum_{\gamma \in Q, k \in \mathbb{Z}} (\dim \mathbb{K}_w^-(\lambda)_{\lambda + \gamma + k\delta}) e^{\lambda + \gamma} q^k,$$

$$\text{gch } \mathbf{K}_w^-(\lambda) := \sum_{\gamma \in Q, k \in \mathbb{Z}} (\dim \mathbf{K}_w^-(\lambda)_{\lambda + \gamma + k\delta}) e^{\lambda + \gamma} q^k.$$

The following is the main result of this paper.

Theorem 2.8 (cf. [FKM, Corollaries 3.19 and 3.20]). *Let $\lambda \in P^+$, and set $S = S_\lambda := \{i \in I \mid \langle \lambda, \alpha_i^\vee \rangle = 0\}$. For $w \in W^S$, the graded character $\text{gch } \mathbb{K}_w^-(\lambda)$ can be expressed as:*

$$\text{gch } \mathbb{K}_w^-(\lambda) = \left(\prod_{i \in I} \prod_{r=1}^{\langle \lambda, \alpha_i^\vee \rangle - \epsilon_i} (1 - q^{-r}) \right)^{-1} E_{w\lambda}(q, \infty), \quad (2.28)$$

where $E_{w\lambda}(q, \infty)$ is the specialization of the nonsymmetric Macdonald polynomial $E_{w\lambda}(q, t)$ at $t = \infty$, and for $i \in I$,

$$\epsilon_i = \epsilon_i([w]) := \begin{cases} 1 & \text{if } [w]s_i > [w], \\ 0 & \text{if } [w]s_i < [w]. \end{cases} \quad (2.29)$$

Moreover, it holds that

$$\text{gch } \mathbf{K}_w^-(\lambda) = E_{w\lambda}(q, \infty). \quad (2.30)$$

Remark 2.9. Keep the notation and setting of the theorem above. We see by (2.4) that $[w]\alpha_i \in \Delta^-$ for all $i \in S$ since $[w] = [w]w_o(S)$. Hence it follows that $[w]s_i < [w]$ for all $i \in S$.

We will give a proof of Theorem 2.8 in Section 5.

3 Semi-infinite Lakshmibai-Seshadri paths

3.1 Crystal structure on semi-infinite LS paths.

In this subsection, we fix $\lambda \in P^+ \subset P_{\text{af}}^0$ (see (2.1) and (2.2)), and take $S = S_\lambda$ as in (2.18).

Definition 3.1. For a rational number $0 < a < 1$, we define $\text{BG}_{a\lambda}^{\frac{\infty}{2}}((W^S)_{\text{af}})$ to be the subgraph of $\text{BG}_{\frac{\infty}{2}}((W^S)_{\text{af}})$ with the same vertex set but having only those directed edges of the form $x \xrightarrow{\beta} y$ for which $a\langle x\lambda, \beta^\vee \rangle \in \mathbb{Z}$ holds.

Definition 3.2. A semi-infinite Lakshmibai-Seshadri (LS for short) path of shape λ is a pair

$$\pi = (\mathbf{x}; \mathbf{a}) = (x_1, \dots, x_s; a_0, a_1, \dots, a_s), \quad s \geq 1, \quad (3.1)$$

of a strictly decreasing sequence $\mathbf{x} : x_1 \succ \dots \succ x_s$ of elements in $(W^S)_{\text{af}}$ and an increasing sequence $\mathbf{a} : 0 = a_0 < a_1 < \dots < a_s = 1$ of rational numbers satisfying the condition that there exists a directed path from x_{u+1} to x_u in $\text{BG}_{a_u\lambda}^{\frac{\infty}{2}}((W^S)_{\text{af}})$ for each $u = 1, 2, \dots, s-1$.

We denote by $\mathbb{B}^{\frac{\infty}{2}}(\lambda)$ the set of all semi-infinite LS paths of shape λ . Following [INS, Sect. 3.1] (see also [NS, Sect. 2.4]), we endow the set $\mathbb{B}^{\frac{\infty}{2}}(\lambda)$ with a crystal structure with weights in P_{af} as follows. Let $\pi \in \mathbb{B}^{\frac{\infty}{2}}(\lambda)$ be of the form (3.1). We define $\bar{\pi} : [0, 1] \rightarrow \mathbb{R} \otimes_{\mathbb{Z}} P_{\text{af}}$ to be the piecewise-linear, continuous map whose ‘‘direction vector’’ for the interval $[a_{u-1}, a_u]$ is $x_u\lambda \in P_{\text{af}}$ for each $1 \leq u \leq s$, that is,

$$\bar{\pi}(t) := \sum_{k=1}^{u-1} (a_k - a_{k-1})x_k\lambda + (t - a_{u-1})x_u\lambda \quad \text{for } t \in [a_{u-1}, a_u], 1 \leq u \leq s. \quad (3.2)$$

We know from [INS, Proposition 3.1.3] that $\bar{\pi}$ is an (ordinary) LS path of shape λ , introduced in [Li, Sect. 4]. We set

$$\text{wt}(\pi) := \bar{\pi}(1) = \sum_{u=1}^s (a_u - a_{u-1})x_u\lambda \in P_{\text{af}}. \quad (3.3)$$

We define root operators $e_i, f_i, i \in I_{\text{af}}$, in the same manner as in [Li, Sect. 2]. Set

$$\begin{cases} H_i^\pi(t) := \langle \bar{\pi}(t), \alpha_i^\vee \rangle & \text{for } t \in [0, 1], \\ m_i^\pi := \min\{H_i^\pi(t) \mid t \in [0, 1]\}. \end{cases} \quad (3.4)$$

As explained in [NS, Remark 2.4.3], all local minima of the function $H_i^\pi(t), t \in [0, 1]$, are integers; in particular, the minimum value m_i^π is a nonpositive integer (recall that $\bar{\pi}(0) = 0$, and hence $H_i^\pi(0) = 0$). We define $e_i\pi$ as follows. If $m_i^\pi = 0$, then we set $e_i\pi := \mathbf{0}$, where $\mathbf{0}$ is an additional element not contained in any crystal. If $m_i^\pi \leq -1$, then we set

$$\begin{cases} t_1 := \min\{t \in [0, 1] \mid H_i^\pi(t) = m_i^\pi\}, \\ t_0 := \max\{t \in [0, t_1] \mid H_i^\pi(t) = m_i^\pi + 1\}; \end{cases} \quad (3.5)$$

notice that $H_i^\pi(t)$ is strictly decreasing on the interval $[t_0, t_1]$. Let $1 \leq p \leq q \leq s$ be such that $a_{p-1} \leq t_0 < a_p$ and $t_1 = a_q$. Then we define $e_i\pi$ to be

$$\begin{aligned} e_i\pi := & (x_1, \dots, x_p, s_i x_p, s_i x_{p+1}, \dots, s_i x_q, x_{q+1}, \dots, x_s; \\ & a_0, \dots, a_{p-1}, t_0, a_p, \dots, a_q = t_1, \dots, a_s); \end{aligned} \quad (3.6)$$

if $t_0 = a_{p-1}$, then we drop x_p and a_{p-1} , and if $s_i x_q = x_{q+1}$, then we drop x_{q+1} and $a_q = t_1$. Similarly, we define $f_i \pi$ as follows. Note that $H_i^\pi(1) - m_i^\pi$ is a nonnegative integer. If $H_i^\pi(1) - m_i^\pi = 0$, then we set $f_i \pi := \mathbf{0}$. If $H_i^\pi(1) - m_i^\pi \geq 1$, then we set

$$\begin{cases} t_0 := \max\{t \in [0, 1] \mid H_i^\pi(t) = m_i^\pi\}, \\ t_1 := \min\{t \in [t_0, 1] \mid H_i^\pi(t) = m_i^\pi + 1\}; \end{cases} \quad (3.7)$$

notice that $H_i^\pi(t)$ is strictly increasing on the interval $[t_0, t_1]$. Let $0 \leq p \leq q \leq s-1$ be such that $t_0 = a_p$ and $a_q < t_1 \leq a_{q+1}$. Then we define $f_i \pi$ to be

$$\begin{aligned} f_i \pi := & (x_1, \dots, x_p, s_i x_{p+1}, \dots, s_i x_q, s_i x_{q+1}, x_{q+1}, \dots, x_s; \\ & a_0, \dots, a_p = t_0, \dots, a_q, t_1, a_{q+1}, \dots, a_s); \end{aligned} \quad (3.8)$$

if $t_1 = a_{q+1}$, then we drop x_{q+1} and a_{q+1} , and if $x_p = s_i x_{p+1}$, then we drop x_p and $a_p = t_0$. In addition, we set $e_i \mathbf{0} = f_i \mathbf{0} := \mathbf{0}$ for all $i \in I_{\text{af}}$.

Theorem 3.3 (see [INS, Theorem 3.1.5]).

- (1) The set $\mathbb{B}^{\frac{\infty}{2}}(\lambda) \sqcup \{\mathbf{0}\}$ is stable under the action of the root operators e_i and f_i , $i \in I_{\text{af}}$.
- (2) For each $\pi \in \mathbb{B}^{\frac{\infty}{2}}(\lambda)$ and $i \in I_{\text{af}}$, we set

$$\begin{cases} \varepsilon_i(\pi) := \max\{n \geq 0 \mid e_i^n \pi \neq \mathbf{0}\}, \\ \varphi_i(\pi) := \max\{n \geq 0 \mid f_i^n \pi \neq \mathbf{0}\}. \end{cases}$$

Then, the set $\mathbb{B}^{\frac{\infty}{2}}(\lambda)$, equipped with the maps wt , e_i , f_i , $i \in I_{\text{af}}$, and ε_i , φ_i , $i \in I_{\text{af}}$, defined above, is a crystal with weights in P_{af} .

Finally, if $\pi \in \mathbb{B}^{\frac{\infty}{2}}(\lambda)$ is of the form (3.1), then we set $\kappa(\pi) := x_s \in (W^S)_{\text{af}}$, and call it the final direction of π . For $x \in W_{\text{af}}$, we set

$$\mathbb{B}_{\succeq x}^{\frac{\infty}{2}}(\lambda) := \{\pi \in \mathbb{B}^{\frac{\infty}{2}}(\lambda) \mid \kappa(\pi) \succeq \Pi^S(x)\}. \quad (3.9)$$

3.2 Realization of the crystal bases of Demazure submodules by semi-infinite LS paths.

As in the previous subsection, we fix $\lambda \in P^+$, and take $S = S_\lambda$ as in (2.18). We write $\lambda \in P^+$ as $\lambda = \sum_{i \in I} m_i \varpi_i$ with $m_i \in \mathbb{Z}_{\geq 0}$ for $i \in I$; recall the definitions of $\overline{\text{Par}(\lambda)}$ and $\text{Par}(\lambda)$ from (2.12) and (2.16), respectively. For $\boldsymbol{\rho} = (\rho^{(i)})_{i \in I} \in \overline{\text{Par}(\lambda)}$, we set $|\boldsymbol{\rho}| := \sum_{i \in I} |\rho^{(i)}|$, where for a partition $\chi = (\chi_1 \geq \chi_2 \geq \dots \geq \chi_m)$, we set $|\chi| := \chi_1 + \dots + \chi_m$. We equip the set $\text{Par}(\lambda)$ with a crystal structure as follows: for each $\boldsymbol{\rho} = (\rho^{(i)})_{i \in I} \in \text{Par}(\lambda)$, we set

$$\begin{cases} e_j \boldsymbol{\rho} = f_j \boldsymbol{\rho} := \mathbf{0}, & \varepsilon_j(\boldsymbol{\rho}) = \varphi_j(\boldsymbol{\rho}) := -\infty \quad \text{for } j \in I_{\text{af}}, \\ \text{wt}(\boldsymbol{\rho}) := -|\boldsymbol{\rho}|\delta. \end{cases} \quad (3.10)$$

Let $\text{Conn}(\mathbb{B}^{\frac{\infty}{2}}(\lambda))$ denote the set of all connected components of $\mathbb{B}^{\frac{\infty}{2}}(\lambda)$, and let $\mathbb{B}_0^{\frac{\infty}{2}}(\lambda) \in \text{Conn}(\mathbb{B}^{\frac{\infty}{2}}(\lambda))$ denote the connected component of $\mathbb{B}^{\frac{\infty}{2}}(\lambda)$ containing $\pi_\lambda := (e; 0, 1) \in \mathbb{B}^{\frac{\infty}{2}}(\lambda)$, where e is the identity element of W_{af} .

Proposition 3.4. *Keep the notation and setting above.*

- (1) *Each connected component $C \in \text{Conn}(\mathbb{B}^{\frac{\infty}{2}}(\lambda))$ of $\mathbb{B}^{\frac{\infty}{2}}(\lambda)$ contains a unique element of the form:*

$$\pi^C = (\Pi^S(t_{\xi_1}), \Pi^S(t_{\xi_2}), \dots, \Pi^S(t_{\xi_{s-1}}), e; a_0, a_1, \dots, a_{s-1}, a_s) \quad (3.11)$$

for some $s \geq 1$ and $\xi_1, \xi_2, \dots, \xi_{s-1} \in Q^{\vee,+}$ (see [INS, Proposition 7.1.2]).

- (2) *There exists a bijection $\Theta : \text{Conn}(\mathbb{B}^{\frac{\infty}{2}}(\lambda)) \rightarrow \text{Par}(\lambda)$ such that $\text{wt}(\pi^C) = \lambda - |\Theta(C)|\delta = \lambda + \text{wt}(\Theta(C))$ (see [INS, Proposition 7.2.1 and its proof]).*
- (3) *Let $C \in \text{Conn}(\mathbb{B}^{\frac{\infty}{2}}(\lambda))$. Then, there exists an isomorphism $C \xrightarrow{\sim} \{\Theta(C)\} \otimes \mathbb{B}_0^{\frac{\infty}{2}}(\lambda)$ of crystals that maps π^C to $\Theta(C) \otimes \pi_\lambda$. Consequently, $\mathbb{B}^{\frac{\infty}{2}}(\lambda)$ is isomorphic as a crystal to $\text{Par}(\lambda) \otimes \mathbb{B}_0^{\frac{\infty}{2}}(\lambda)$ (see [INS, Proposition 3.2.4 and its proof]).*

We know the following from [INS, Theorem 3.2.1] and [NS, Theorem 4.2.1].

Theorem 3.5. *There exists an isomorphism $\Psi_\lambda : \mathcal{B}(\lambda) \xrightarrow{\sim} \mathbb{B}^{\frac{\infty}{2}}(\lambda)$ of crystals satisfying the following conditions (1) and (2).*

- (1) $\Psi_\lambda(u^\rho) = \pi^{\Theta^{-1}(\rho)}$ for every $\rho \in \text{Par}(\lambda)$, where $u^\rho = z_\rho u_\lambda \in \mathcal{B}(\lambda)$ is as defined in (2.17). In particular, $\Psi_\lambda(u_\lambda) = \pi_\lambda$.
- (2) $\Psi_\lambda(\mathcal{B}_x^-(\lambda)) = \mathbb{B}_{\sum x}^{\frac{\infty}{2}}(\lambda)$ for every $x \in (W^S)_{\text{af}}$.

Remark 3.6. Recall from (2.11) that the crystal basis $\mathcal{B}(\lambda)$ has an action of the affine Weyl group W_{af} . Hence, by Theorem 3.5, the crystal $\mathbb{B}^{\frac{\infty}{2}}(\lambda)$ also has the induced action of the affine Weyl group W_{af} , which we denote by $x \cdot \pi$ for $x \in W_{\text{af}}$ and $\pi \in \mathbb{B}^{\frac{\infty}{2}}(\lambda)$.

For $\gamma \in Q$ and $k \in \mathbb{Z}$, we set $\text{fin}(\lambda + \gamma + k\delta) := \lambda + \gamma \in P$ and $\text{nul}(\lambda + \gamma + k\delta) := k \in \mathbb{Z}$. Let $w \in W^S$. Then, from Theorem 3.5 and (2.21), we deduce that

$$\text{gch } \mathbb{K}_w^-(\lambda) = \sum_{\pi \in \mathbb{B}_{\sum w; \not> w}^{\frac{\infty}{2}}(\lambda)} \mathbf{e}^{\text{fin}(\text{wt}(\pi))} q^{\text{nul}(\text{wt}(\pi))}, \quad (3.12)$$

where

$$\mathbb{B}_{\sum w; \not> w}^{\frac{\infty}{2}}(\lambda) := \mathbb{B}_{\sum w}^{\frac{\infty}{2}}(\lambda) \setminus \bigcup_{z \in W^S, z > w} \mathbb{B}_{\sum z}^{\frac{\infty}{2}}(\lambda). \quad (3.13)$$

If we set $((W^S)_{\text{af}})_{\succeq x} := \{y \in (W^S)_{\text{af}} \mid y \succeq x\}$ for $x \in (W^S)_{\text{af}}$, and

$$K_w^S := ((W^S)_{\text{af}})_{\succeq w} \setminus \bigcup_{z \in W^S, z > w} ((W^S)_{\text{af}})_{\succeq z}, \quad (3.14)$$

then it is easily verified that

$$\mathbb{B}_{\succeq w; \not\succeq w}^{\infty}(\lambda) = \{\pi \in \mathbb{B}^{\infty}(\lambda) \mid \kappa(\pi) \in K_w^S\}. \quad (3.15)$$

4 Description of the crystal bases of level-zero van der Kallen modules.

4.1 Quantum Bruhat graph and the tilted Bruhat order.

In this subsection, we take and fix a subset S of I .

Definition 4.1. The (parabolic) quantum Bruhat graph $\text{QBG}(W^S)$ is the $(\Delta^+ \setminus \Delta_S^+)$ -labeled directed graph whose vertices are the elements of W^S , and whose directed edges are of the form: $u \xrightarrow{\beta} v$ for $u, v \in W^S$ and $\beta \in \Delta^+ \setminus \Delta_S^+$ such that $v = [us_\beta]$, and such that either of the following (i) or (ii) holds:

- (i) $\ell(v) = \ell(u) + 1$;
- (ii) $\ell(v) = \ell(u) + 1 - 2\langle \rho - \rho_S, \beta^\vee \rangle$.

An edge satisfying (i) (resp., (ii)) is called a Bruhat (resp., quantum) edge. When $S = \emptyset$, we write $\text{QBG}(W)$ for $\text{QBG}(W^\emptyset)$.

Remark 4.2. We know from [LNS³1, Remark 6.13] that for each $u, v \in W^S$, there exists a directed path in $\text{QBG}(W^S)$ from u to v .

Let $u, v \in W^S$, and let $\mathbf{p} : u = u_0 \xrightarrow{\beta_1} u_1 \xrightarrow{\beta_2} \cdots \xrightarrow{\beta_s} u_s = v$ be a directed path in $\text{QBG}(W^S)$ from u to v . Then we define the weight of \mathbf{p} by

$$\text{wt}^S(\mathbf{p}) := \sum_{\substack{1 \leq r \leq s; \\ u_{r-1} \xrightarrow{\beta_r} u_r \text{ is} \\ \text{a quantum edge}}} \beta_r^\vee \in Q^{\vee,+}; \quad (4.1)$$

when $S = \emptyset$, we write $\text{wt}(\mathbf{p})$ for $\text{wt}^\emptyset(\mathbf{p})$. We know the following proposition from [LNS³1, Proposition 8.1 and its proof].

Proposition 4.3. *Let $u, v \in W^S$. Let \mathbf{p} be a shortest directed path in $\text{QBG}(W^S)$ from u to v , and \mathbf{q} an arbitrary directed path in $\text{QBG}(W^S)$ from u to v . Then, $[\text{wt}^S(\mathbf{q}) - \text{wt}^S(\mathbf{p})]^S \in Q_{I \setminus S}^{\vee,+}$, where $[\cdot]^S : Q^\vee \rightarrow Q_{I \setminus S}^{\vee}$ is as defined in Section 2.2. Moreover, \mathbf{q} is also shortest if and only if $[\text{wt}^S(\mathbf{q})]^S = [\text{wt}^S(\mathbf{p})]^S$.*

For $u, v \in W^S$, we take a shortest directed path \mathbf{p} in $\text{QBG}(W^S)$ from u to v , and set $\text{wt}^S(u \Rightarrow v) := [\text{wt}^S(\mathbf{p})]^S \in Q_{I \setminus S}^{\vee,+}$. When $S = \emptyset$, we write $\text{wt}(u \Rightarrow v)$ for $\text{wt}^\emptyset(u \Rightarrow v)$.

Lemma 4.4 ([LNS³2, Lemma 7.2]). *Let $u, v \in W^S$, and let $u_1 \in uW_S, v_1 \in vW_S$. Then we have $\text{wt}^S(u \Rightarrow v) = [\text{wt}(u_1 \Rightarrow v_1)]^S$.*

For $u, v \in W$, we denote by $\ell(u \Rightarrow v)$ the length of a shortest directed path from u to v in $\text{QBG}(W)$.

Lemma 4.5 ([LNS³1, Lemma 7.7]). *Let $u, v \in W$, and $i \in I$.*

- (1) *If $u^{-1}\alpha_i \in \Delta^+$ and $v^{-1}\alpha_i \in \Delta^-$, then $\ell(u \Rightarrow v) = \ell(s_i u \Rightarrow v) + 1 = \ell(u \Rightarrow s_i v) + 1$, and $\text{wt}(u \Rightarrow v) = \text{wt}(u \Rightarrow s_i v) = \text{wt}(s_i u \Rightarrow v)$.*
- (2) *If $u^{-1}\alpha_i, v^{-1}\alpha_i \in \Delta^+$, or if $u^{-1}\alpha_i, v^{-1}\alpha_i \in \Delta^-$, then $\ell(u \Rightarrow v) = \ell(s_i u \Rightarrow s_i v)$, and $\text{wt}(u \Rightarrow v) = \text{wt}(s_i u \Rightarrow s_i v)$.*

Definition 4.6 ([BFP]). For each $w \in W$, we define the w -tilted Bruhat order \leq_w on W as follows: for $u, v \in W$,

$$u \leq_w v \iff \ell(w \Rightarrow v) = \ell(w \Rightarrow u) + \ell(u \Rightarrow v). \quad (4.2)$$

Namely, $u \leq_w v$ if and only if the concatenation of a shortest directed path from w to u and one from u to v is one from w to v .

4.2 Some lemmas on the semi-infinite Bruhat order.

In this subsection, we fix a subset S of I .

Lemma 4.7 ([INS, Lemma 6.1.1]). *If $x, y \in W_{\text{af}}$ satisfy $x \succeq y$ in W_{af} , then $\Pi^S(x) \succeq \Pi^S(y)$ in $(W^S)_{\text{af}}$.*

Lemma 4.8 ([NNS1, Lemmas 4.3.5, 4.3.6, and 4.3.7]). *Let $u, v \in W^S$, and $\xi, \zeta \in Q^\vee$. Then,*

$$u\Pi^S(t_\xi) \succeq v\Pi^S(t_\zeta) \iff [\xi]^S \geq \text{wt}^S(v \Rightarrow u) + [\zeta]^S. \quad (4.3)$$

Lemma 4.9 ([INS, Lemma 2.3.6]). *Let $x \in (W^S)_{\text{af}}$, and $i \in I_{\text{af}}$. Then, $x^{-1}\alpha_i \notin (\Delta_S)_{\text{af}}$ if and only if $s_i x \in (W^S)_{\text{af}}$. In particular, for $u \in W^S$ and $i \in I$, $u^{-1}\alpha_i \notin \Delta_S$ if and only if $s_i u \in W^S$ (see also [LNS³1, Proposition 5.10]).*

Remark 4.10. Keep the notation in Lemma 4.9. Let $\lambda \in P^+ \subset P_{\text{af}}^0$ be such that $\{i \in I \mid \langle \lambda, \alpha_i^\vee \rangle = 0\} = S$. Then, $x^{-1}\alpha_i \notin (\Delta_S)_{\text{af}}$ (resp., $u^{-1}\alpha_i \notin \Delta_S$) if and only if $\langle x\lambda, \alpha_i^\vee \rangle \neq 0$ (resp., $\langle u\lambda, \alpha_i^\vee \rangle \neq 0$).

Lemma 4.11 ([NS, Lemma 2.3.6]). *Let $x, y \in (W^S)_{\text{af}}$ be such that $x \preceq y$, and let $i \in I_{\text{af}}$.*

- (1) *If $x^{-1}\alpha_i \in (\Delta^+ \setminus \Delta_S^+) + \mathbb{Z}\delta$ and $y^{-1}\alpha_i \in (\Delta^- \cup \Delta_S) + \mathbb{Z}\delta$ (or equivalently, $\langle x\lambda, \alpha_i^\vee \rangle > 0$ and $\langle y\lambda, \alpha_i^\vee \rangle \leq 0$, where λ is as in Remark 4.10), then $s_i x \in (W^S)_{\text{af}}$ and $s_i x \preceq y$.*
- (2) *If $x^{-1}\alpha_i \in (\Delta^+ \cup \Delta_S) + \mathbb{Z}\delta$ and $y^{-1}\alpha_i \in (\Delta^- \setminus \Delta_S^-) + \mathbb{Z}\delta$ (or equivalently, $\langle x\lambda, \alpha_i^\vee \rangle \geq 0$ and $\langle y\lambda, \alpha_i^\vee \rangle < 0$), then $s_i y \in (W^S)_{\text{af}}$ and $x \preceq s_i y$.*
- (3) *If $x^{-1}\alpha_i, y^{-1}\alpha_i \in (\Delta^+ \setminus \Delta_S^+) + \mathbb{Z}\delta$ (or equivalently, $\langle x\lambda, \alpha_i^\vee \rangle > 0$ and $\langle y\lambda, \alpha_i^\vee \rangle > 0$), or if $x^{-1}\alpha_i, y^{-1}\alpha_i \in (\Delta^- \setminus \Delta_S^-) + \mathbb{Z}\delta$ (or equivalently, $\langle x\lambda, \alpha_i^\vee \rangle < 0$ and $\langle y\lambda, \alpha_i^\vee \rangle < 0$), then $s_i x \preceq s_i y$.*

By Remark 2.5, the next well-known lemma (see, e.g., [H, Lemma on page 151] for the case of $S = \emptyset$) also follows from Lemma 4.11 as a special case.

Lemma 4.12. *Let $u, v \in W^S$ be such that $u \preceq v$, and let $i \in I$.*

- (1) *If $u^{-1}\alpha_i \in \Delta^+ \setminus \Delta_S^+$ and $v^{-1}\alpha_i \in \Delta^- \cup \Delta_S$ (or equivalently, $\langle u\lambda, \alpha_i^\vee \rangle > 0$ and $\langle v\lambda, \alpha_i^\vee \rangle \leq 0$, where λ is as in Remark 4.10), then $s_i u \in W^S$ and $s_i u \leq v$.*
- (2) *If $u^{-1}\alpha_i \in \Delta^+ \cup \Delta_S$ and $v^{-1}\alpha_i \in \Delta^- \setminus \Delta_S^-$ (or equivalently, $\langle u\lambda, \alpha_i^\vee \rangle \geq 0$ and $\langle v\lambda, \alpha_i^\vee \rangle < 0$), then $s_i v \in W^S$ and $u \leq s_i v$.*
- (3) *If $u^{-1}\alpha_i, v^{-1}\alpha_i \in \Delta^+ \setminus \Delta_S^+$ (or equivalently, $\langle u\lambda, \alpha_i^\vee \rangle > 0$ and $\langle v\lambda, \alpha_i^\vee \rangle > 0$), or if $u^{-1}\alpha_i, v^{-1}\alpha_i \in \Delta^- \setminus \Delta_S^-$ (or equivalently, $\langle u\lambda, \alpha_i^\vee \rangle < 0$ and $\langle v\lambda, \alpha_i^\vee \rangle < 0$), then $s_i u \leq s_i v$.*

4.3 Definition and a recursive description of subsets $\text{EQB}(w)$.

For $w \in W$, we define the right descent set I_w for w by

$$I_w := \{j \in I \mid ws_j < w\} = \{j \in I \mid w\alpha_j \in \Delta^-\}. \quad (4.4)$$

Remark 4.13 (see Remark 2.9). Let S be a subset of I . Then we have $S \subset I_{[w]}$ for $w \in W^S$.

Lemma 4.14 ([NNS2, Lemma 3.1.1]). *Let $w \in W$ and $i \in I$ be such that $s_i w < w$.*

- (a) *$s_i w \notin wW_{I_w}$ if and only if $-w^{-1}\alpha_i$ is not a simple root. In this case, $I_{s_i w} = I_w$.*
- (b) *$s_i w \in wW_{I_w}$ if and only if $-w^{-1}\alpha_i$ is a simple root. In this case, $I_{s_i w} = I_w \setminus \{k\}$, where $\alpha_k = -w^{-1}\alpha_i$.*

Now, we recall from [NNS1, Sect. 3.2] and [NNS2, Sect. 2.2] the definition of the subsets $\text{EQB}(w) \subset W$ for $w \in W$. For each $w \in W$, take and fix a reduced expression

$$w = s_{i_1} s_{i_2} \cdots s_{i_p}. \quad (4.5)$$

Then we take $i_{-q}, i_{-q+1}, \dots, i_0 \in I$ in such a way that $s_{i_{-q}} \cdots s_{i_0} s_{i_1} \cdots s_{i_p}$ is a reduced expression for the longest element $w_\circ \in W$. We set

$$\beta_k := s_{i_p} \cdots s_{i_{k+1}} \alpha_{i_k} \in \Delta^+ \quad \text{for } -q \leq k \leq p. \quad (4.6)$$

Since $\beta_{-q} < \cdots < \beta_p$ is a reflection order on Δ^+ , we know (see, e.g., [LNS³1, Theorem 7.3]) that for each $u \in W$, there exists a unique shortest directed path

$$w = x_0 \xrightarrow{\beta_{j_1}} x_1 \xrightarrow{\beta_{j_2}} \cdots \xrightarrow{\beta_{j_s}} x_s = u \quad (4.7)$$

in $\text{QBG}(W)$ from w to u such that $-q \leq j_1 < j_2 < \cdots < j_s \leq p$, which we call the “label-increasing” directed path. The subset $\text{EQB}(w) \subset W$ is defined to be the set of all those elements $u \in W$ whose label-increasing directed path (4.7) from w to u satisfies $j_1 \geq 1$; we know from [NNS1, Proposition 3.2.5] that this definition of $\text{EQB}(w)$ does not depend on the choice of a reduced expression (4.5) for w . The subsets $\text{EQB}(w)$, $w \in W$, above are also determined through the following recursive formula by descending induction.

Proposition 4.15 ([NNS2, Proposition 3.2.3]).

- (1) For the longest element $w_\circ \in W$, it holds that $\text{EQB}(w_\circ) = W$.
- (2) Let $w \in W$ and $i \in I$ be such that $s_i w < w$. If $s_i w \notin wW_{I_w}$, then

$$\begin{cases} \text{EQB}(w) \cap \text{EQB}(s_i w) = \emptyset, \\ \text{EQB}(w) \cup s_i \text{EQB}(w) = \text{EQB}(w) \sqcup \text{EQB}(s_i w); \end{cases} \quad (4.8)$$

if $s_i w \in wW_{I_w}$, then

$$\begin{cases} \text{EQB}(s_i w) = \{v \in \text{EQB}(w) \mid s_i w \leq_w v\}, \\ s_i \text{EQB}(w) = \text{EQB}(w), \end{cases} \quad (4.9)$$

where \leq_w is the w -tilted Bruhat order on W (see Definition 4.6).

4.4 Description of K_w^S in terms of the quantum Bruhat graph.

In the case $S = \emptyset$, for $w \in W = W^\emptyset$, we write K_w for the subset K_w^\emptyset defined by (3.14), that is,

$$K_w = (W_{\text{af}})_{\succeq w} \setminus \bigcup_{z \in W, z > w} (W_{\text{af}})_{\succeq z}. \quad (4.10)$$

Proposition 4.16. For $w \in W$, the subset K_w is identical to the set $\{ut_\xi \in W_{\text{af}} \mid u \in \text{EQB}(w), \xi \in \text{wt}(w \Rightarrow u) + Q_{I_w}^{\vee,+}\}$, where $\text{EQB}(w)$, $\text{wt}(w \Rightarrow u)$, and I_w are as defined in Section 4.3, (4.1), and (4.4), respectively.

In order to prove Proposition 4.16, we need some lemmas.

Lemma 4.17. If $w = w_\circ$, then $K_{w_\circ} = \{ut_\xi \in W_{\text{af}} \mid u \in W, \xi \in Q^\vee, \xi \geq \text{wt}(w_\circ \Rightarrow u)\}$.

Proof. The assertion follows immediately from Lemma 4.8 (with $S = \emptyset$), since there is no $z \in W$ such that $z > w_\circ$. \square

Lemma 4.18. Let $w \in W$. We have

$$\text{EQB}(w) = \{u \in W \mid \text{there does not exist } z \in W \text{ such that } z > w \text{ and } z \leq_w u\}. \quad (4.11)$$

Proof. Let u be an element of the set on the right-hand side in (4.11). Define β_k , $-q \leq k \leq p$, as in (4.6), and take the label-increasing directed path

$$w = x_0 \xrightarrow{\beta_{j_1}} x_1 \xrightarrow{\beta_{j_2}} \cdots \xrightarrow{\beta_{j_s}} x_s = u \quad (4.12)$$

in $\text{QBG}(W)$ from w to u (see (4.7)). Suppose, for a contradiction, that $j_1 \leq 0$. Then we see that

$$w\beta_{j_1} = (s_{i_1} \cdots s_{i_p})(s_{i_p} \cdots s_{i_1} s_{i_0} \cdots s_{i_{j_1+1}} \alpha_{i_{j_1}}) = s_{i_0} \cdots s_{i_{j_1+1}} \alpha_{i_{j_1}} \in \Delta^+,$$

which implies that $w = x_0 \xrightarrow{\beta_{j_1}} x_1$ is a Bruhat edge, and hence $z := x_1 > w$. Since (4.12) is a shortest directed path from w to u passing through $z = x_1$, we see that $z \leq_w u$, which is a contradiction. Thus we obtain $j_1 \geq 1$, and hence $u \in \text{EQB}(w)$.

Next, we prove the opposite inclusion \subset by descending induction on $\ell(w)$. If $w = w_\circ$, then the assertion is obvious since there does not exist $z \in W$ such that $z > w_\circ$. Let $w \in W$ and $i \in I$ be such that $s_i w < w$; note that $w^{-1}\alpha_i \in \Delta^-$. Then we can take a reduced expression (4.5) for w such that $i_1 = i$; in this case, $\beta_1 = -w^{-1}\alpha_i$. Assume that (4.11) holds for w (the induction hypothesis), and suppose, for a contradiction, that for some $u \in \text{EQB}(s_i w)$, there exists $z \in W$ such that $z > s_i w$ and $z \leq_{s_i w} u$; note that

$$\ell(s_i w \Rightarrow u) = \ell(s_i w \Rightarrow z) + \ell(z \Rightarrow u). \quad (4.13)$$

Case 1. Assume that $s_i u > u$, or equivalently, $u^{-1}\alpha_i \in \Delta^+$. We see from Proposition 4.15(2) that $s_i u \in \text{EQB}(w)$. Indeed, if $s_i w \notin wW_{I_{[w]}}$, then it follows from the second equality of (4.8) that $u \in \text{EQB}(w)$ or $s_i u \in \text{EQB}(w)$. However, since $\text{EQB}(w) \cap \text{EQB}(s_i w) = \emptyset$ by the first equality of (4.8), and since $u \in \text{EQB}(s_i w)$, we obtain $u \notin \text{EQB}(w)$, and hence $s_i u \in \text{EQB}(w)$. Also, if $s_i w \in wW_{I_{[w]}}$, then we have $u \in \text{EQB}(s_i w) \subset \text{EQB}(w)$ by the first equality of (4.9). Hence we have $s_i u \in \text{EQB}(w)$ by the second equality of (4.9).

Subcase 1.1. Assume that $s_i z < z$, or equivalently, $z^{-1}\alpha_i \in \Delta^-$. By Lemmas 4.12 (1) and 4.5 (1), we deduce that $z \geq w$ and $\ell(w \Rightarrow z) = \ell(s_i w \Rightarrow z) - 1$. Since $u^{-1}\alpha_i \in \Delta^+$, we have a Bruhat edge $u \xrightarrow{u^{-1}\alpha_i} s_i u$, which implies that $\ell(z \Rightarrow s_i u) \leq \ell(z \Rightarrow u) + 1$. Also, since $u^{-1}\alpha_i \in \Delta^+$ and $(s_i w)^{-1}\alpha_i \in \Delta^+$, it follows from Lemma 4.5 (2) that $\ell(s_i w \Rightarrow u) = \ell(w \Rightarrow s_i u)$. Combining these, we see that

$$\begin{aligned} \ell(s_i w \Rightarrow u) &= \ell(w \Rightarrow s_i u) \leq \ell(w \Rightarrow z) + \ell(z \Rightarrow s_i u) \\ &\leq \ell(s_i w \Rightarrow z) - 1 + \ell(z \Rightarrow u) + 1 = \ell(s_i w \Rightarrow z) + \ell(z \Rightarrow u) \\ &= \ell(s_i w \Rightarrow u) \quad \text{by (4.13)}. \end{aligned}$$

In particular, we obtain $\ell(w \Rightarrow s_i u) = \ell(w \Rightarrow z) + \ell(z \Rightarrow s_i u)$, which implies that $z \leq_w s_i u$. Here we recall that $s_i u \in \text{EQB}(w)$ and $z \geq w$, as seen above. Therefore, by the induction hypothesis, we must have $z = w$. In particular, we obtain $w \leq_{s_i w} u$.

By concatenating the label-increasing (shortest) directed path (4.12) in $\text{QBG}(W)$ from w to u with the Bruhat edge $s_i w \xrightarrow{\beta_1} w$, we obtain

$$s_i w \xrightarrow{\beta_1} w = x_0 \xrightarrow{\beta_{j_1}} x_1 \xrightarrow{\beta_{j_2}} \cdots \xrightarrow{\beta_{j_s}} x_s = u, \quad (4.14)$$

which is a shortest directed path from $s_i w$ to u since $w \leq_{s_i w} u$. Let

$$s_i w = y_0 \xrightarrow{\beta_{k_1}} y_1 \xrightarrow{\beta_{k_2}} \cdots \xrightarrow{\beta_{k_{s+1}}} y_{s+1} = u \quad (4.15)$$

be the shortest directed path from $s_i w$ to u such that $-q \leq k_1 < \cdots < k_{s+1} \leq p$. Since $u \in \text{EQB}(s_i w)$ by our assumption, we see by [NNS2, Remark 23] that $k_1 \geq 2$, and hence $\beta_1 < \beta_{k_1}$ in our fixed reflection order. However, we know (see, e.g., [LNS³1, Theorem 7.3]) that the shortest directed path (4.15) is lexicographically minimal, that is, for every shortest directed path $s_i w = y'_0 \xrightarrow{\gamma_1} y'_1 \xrightarrow{\gamma_2} \cdots \xrightarrow{\gamma_{s+1}} y'_{s+1} = u$ in $\text{QBG}(W)$ from $s_i w$ to u , there exists $1 \leq a \leq s + 1$ such that $\gamma_b = \beta_{k_b}$ for $1 \leq b \leq a - 1$, and $\gamma_a > \beta_{k_a}$. Therefore, we obtain $\beta_{k_1} \leq \beta_1$, which is a contradiction.

Subcase 1.2. Assume that $s_i z > z$, or equivalently, $z^{-1}\alpha_i \in \Delta^+$. By Lemma 4.12 (3), we see that $s_i z > w$. Also, we deduce from Lemma 4.5 (2) that $\ell(s_i w \Rightarrow z) = \ell(w \Rightarrow s_i z)$, $\ell(s_i w \Rightarrow u) = \ell(w \Rightarrow s_i u)$, and $\ell(z \Rightarrow u) = \ell(s_i z \Rightarrow s_i u)$. Substituting these equalities into (4.13), we obtain $\ell(w \Rightarrow s_i u) = \ell(w \Rightarrow s_i z) + \ell(s_i z \Rightarrow s_i u)$, which implies that $s_i z \leq_w s_i u$. Since $s_i u \in \text{EQB}(w)$ and $s_i z > w$, as seen above, the inequality $s_i z \leq_w s_i u$ contradicts the induction hypothesis.

Case 2. Assume that $s_i u < u$, or equivalently, $u^{-1}\alpha_i \in \Delta^-$. Then we have $s_i w \in wW_{I_w}$. Indeed, since $s_i w < w$, $s_i(s_i u) = u > s_i u$, and $s_i u \in \text{EQB}(w)$, as seen above, it follows

from [NNS2, Lemma 24 (2)] that $s_i(s_i u) = u \in \text{EQB}(w)$. Thus we have $u \in \text{EQB}(w) \cap \text{EQB}(s_i w)$, which contradicts Proposition 4.15 (2). Hence we conclude that $s_i w \in wW_{I_w}$. In particular, we obtain $u \in \text{EQB}(w)$ by Proposition 4.15 (2).

Subcase 2.1. Assume that $s_i z < z$, or equivalently, $z^{-1}\alpha_i \in \Delta^-$. By Lemmas 4.12 (1) and 4.5 (1), we see that $z \geq w$ and $\ell(w \Rightarrow z) = \ell(s_i w \Rightarrow z) - 1$. Similarly, we deduce by Lemma 4.5 (1) that $\ell(w \Rightarrow u) = \ell(s_i w \Rightarrow u) - 1$. Substituting these equalities into (4.13), we obtain $\ell(w \Rightarrow u) = \ell(w \Rightarrow z) + \ell(z \Rightarrow u)$, which implies that $z \leq_w u$. Since $u \in \text{EQB}(w)$ and $z \geq w$, as seen above, it follows from the induction hypothesis that $z = w$. In particular, we have $w \leq_{s_i w} u$. In exactly the same way as in the second paragraph of Subcase 1.1, we obtain a contradiction from this inequality.

Subcase 2.2. Assume that $s_i z > z$, or equivalently, $z^{-1}\alpha_i \in \Delta^+$. By Lemmas 4.12 (3) and 4.5 (2), we see that $s_i z > w$ and $\ell(w \Rightarrow s_i z) = \ell(s_i w \Rightarrow z)$. Also, we deduce by Lemma 4.5 (1) that $\ell(w \Rightarrow u) = \ell(s_i w \Rightarrow u) - 1$, and $\ell(s_i z \Rightarrow u) = \ell(z \Rightarrow u) - 1$. Substituting these equalities into (4.13), we obtain $\ell(w \Rightarrow u) = \ell(w \Rightarrow s_i z) + \ell(s_i z \Rightarrow u)$, which implies that $s_i z \leq_w u$. Since $u \in \text{EQB}(w)$ and $s_i z > w$, as seen above, the inequality $s_i z \leq_w u$ contradicts the induction hypothesis.

This completes the proof of the lemma. \square

Lemma 4.19. *Let $w \in W$ and $i \in I$ be such that $s_i w < w$. For every $u \in \text{EQB}(s_i w)$, we have $s_i u > u$, or equivalently, $u^{-1}\alpha_i \in \Delta^+$.*

Proof. Suppose, for a contradiction, that there exists $u \in \text{EQB}(s_i w)$ such that $s_i u < u$, or equivalently, $u^{-1}\alpha_i \in \Delta^-$. Since $s_i w < w$, it follows from Lemma 4.5 (1) that $\ell(s_i w \Rightarrow u) = \ell(w \Rightarrow u) + 1$. Also, we have a Bruhat edge $s_i w \xrightarrow{-w^{-1}\alpha_i} w$, and hence $\ell(s_i w \Rightarrow w) = 1$. Therefore, we obtain $\ell(s_i w \Rightarrow w) + \ell(w \Rightarrow u) = \ell(s_i w \Rightarrow u)$, which implies that $w \leq_{s_i w} u$. Since $w > s_i w$ and $u \in \text{EQB}(s_i w)$ by the assumption, the equality $w \leq_{s_i w} u$ contradicts Lemma 4.18. This proves the lemma. \square

We set

$$\text{fin}(K_w) := \{u \in W \mid ut_\xi \in K_w \text{ for some } \xi \in Q^\vee\}. \quad (4.16)$$

Remark 4.20. We deduce by Lemma 4.8 (with $S = \emptyset$) that $u \in \text{fin}(K_w)$ if and only if $ut_{\text{wt}(w \Rightarrow u)} \in K_w$. Indeed, the “if” part is obvious from the definition. Let us show the “only if” part. Let $u \in \text{fin}(K_w)$. By the definition, there exists $\xi \in Q^\vee$ such that $ut_\xi \in K_w$. Since $ut_\xi \in K_w \subset (W_{\text{af}})_{\succeq w}$, we see from Lemma 4.8 that $\xi \geq \text{wt}(w \Rightarrow u)$, and hence $ut_\xi \succeq ut_{\text{wt}(w \Rightarrow u)}$. If $ut_{\text{wt}(w \Rightarrow u)} \in (W_{\text{af}})_{\succeq z}$ for some $z \in W$ such that $z > w$, then we have $ut_\xi \succeq ut_{\text{wt}(w \Rightarrow u)} \succeq z$, which contradicts the fact that $ut_\xi \in K_w$.

Lemma 4.21. *For every $w \in W$, we have $\text{fin}(K_w) = \text{EQB}(w)$.*

Proof. First we prove that $\text{fin}(K_w) \subset \text{EQB}(w)$. Let $u \in \text{fin}(K_w)$ (note that $ut_{\text{wt}(w \Rightarrow u)} \in K_w$ by Remark 4.20), and suppose, for a contradiction, that $u \notin \text{EQB}(w)$. By Lemma 4.18, there exists $z \in W$ such that $z > w$ and $z \leq_w u$. Since $z \leq_w u$, we see that $\text{wt}(z \Rightarrow u) \leq \text{wt}(w \Rightarrow u)$, which implies that $ut_{\text{wt}(w \Rightarrow u)} \succeq ut_{\text{wt}(z \Rightarrow u)} \succeq z$ by Lemma 4.8 (with $S = \emptyset$). This contradicts the fact that $ut_{\text{wt}(w \Rightarrow u)} \in K_w$.

Next we prove that $\text{fin}(K_w) \supset \text{EQB}(w)$. Let $u \in \text{EQB}(w)$. It suffices to show that $ut_{\text{wt}(w \Rightarrow u)} \in K_w$. It is obvious from Lemma 4.8 that $ut_{\text{wt}(w \Rightarrow u)} \in (W_{\text{af}})_{\succeq w}$. Suppose, for a contradiction, that $ut_{\text{wt}(w \Rightarrow u)} \succeq z$ for some $z \in W$ such that $z > w$; it follows from Lemma 4.8 that $\text{wt}(z \Rightarrow u) \leq \text{wt}(w \Rightarrow u)$. Take arbitrary shortest directed paths

$$w = x_0 \xrightarrow{\gamma_1} \cdots \xrightarrow{\gamma_a} x_a = z, \quad z = x_a \xrightarrow{\gamma_{a+1}} \cdots \xrightarrow{\gamma_b} x_b = u$$

in $\text{QBG}(W)$, and concatenate these as:

$$\mathbf{p} : w = x_0 \xrightarrow{\gamma_1} \cdots \xrightarrow{\gamma_a} x_a = z \xrightarrow{\gamma_{a+1}} \cdots \xrightarrow{\gamma_b} x_b = u.$$

Since $z > w$ in the (ordinary) Bruhat order on W , we deduce that all the edges in the shortest directed path above from w to z are Bruhat edges, which implies that $\text{wt}(\mathbf{p}) = \text{wt}(z \Rightarrow u)$. Also, it follows from Proposition 4.3 (with $S = \emptyset$) that $\text{wt}(\mathbf{p}) \geq \text{wt}(w \Rightarrow u)$. Therefore, we obtain

$$\text{wt}(z \Rightarrow u) \leq \text{wt}(w \Rightarrow u) \leq \text{wt}(\mathbf{p}) = \text{wt}(z \Rightarrow u),$$

and hence $\text{wt}(w \Rightarrow u) = \text{wt}(\mathbf{p})$. In particular, we deduce from Proposition 4.3 that \mathbf{p} is a shortest directed path from w to u . Hence we obtain $z \leq_w u$. Since $u \in \text{EQB}(w)$ and $z > w$ by our assumption, the inequality $z \leq_w u$ contradicts Lemma 4.18. Thus we have shown that $u \in \text{fin}(K_w)$. This proves the lemma. \square

Proof of Proposition 4.16. For $w \in W$, we set

$$K'_w := \{ut_\xi \in W_{\text{af}} \mid u \in \text{EQB}(w), \xi \in \text{wt}(w \Rightarrow u) + Q_{I_w}^{\vee,+}\}. \quad (4.17)$$

We prove the equality $K_w = K'_w$ by descending induction on $\ell(w)$. If $w = w_\circ$, then the equality follows from Lemma 4.17. Let $w \in W$ and $i \in I$ be such that $s_i w < w$. Assume that $K_w = K'_w$ (the induction hypothesis). We will prove that $K_{s_i w} = K'_{s_i w}$.

Step 1. We prove the inclusion $K_{s_i w} \subset K'_{s_i w}$. Let $x \in K_{s_i w}$, and write it as $x = ut_\xi \in K_{s_i w}$ for some $u \in W$ and $\xi \in Q^\vee$. We see from Lemma 4.21 that $u \in \text{EQB}(s_i w)$. Hence it remains to show that $\xi \in \text{wt}(s_i w \Rightarrow u) + Q_{I_{s_i w}}^{\vee,+}$. Since $u \in \text{EQB}(s_i w)$ and $s_i w < w$,

it follows from Lemma 4.19 that $u^{-1}\alpha_i \in \Delta^+$. Therefore, from the assumption that $x = ut_\xi \succeq s_iw$, we see by Lemma 4.11 (3) that $s_iut_\xi \succeq w$, and hence $s_iut_\xi \in (W_{\text{af}})_{\succeq w}$. Suppose, for a contradiction, that $s_iut_\xi \succeq z$ for some $z \in W$ such that $z > w$. If $z^{-1}\alpha_i \in \Delta^+$, then we see by Lemma 4.11 (1) that $ut_\xi \succeq z$. Hence we have $ut_\xi \succeq z > w > s_iw$, which contradicts the assumption that $ut_\xi \in K_{s_iw}$. Similarly, if $z^{-1}\alpha_i \in \Delta^-$, then we see by Lemma 4.11 (3) that $ut_\xi \succeq s_iz$. Hence we have $ut_\xi \succeq s_iz > s_iw$ (by Lemma 4.12 (3)), which contradicts the assumption that $ut_\xi \in K_{s_iw}$. Thus we conclude that $s_iut_\xi \in K_w$. From this, by the induction hypothesis, we obtain $\xi \in \text{wt}(w \Rightarrow s_iu) + Q_{I_w}^{\vee,+}$; note that $\text{wt}(w \Rightarrow s_iu) = \text{wt}(s_iw \Rightarrow u)$ by Lemma 4.5 (2).

Case 1.1 Assume that $s_iw \notin wW_{I_w}$. In this case, we have $I_{s_iw} = I_w$ by Lemma 4.14 (a), and hence $\xi \in \text{wt}(s_iw \Rightarrow u) + Q_{I_{s_iw}}^{\vee,+}$, as desired.

Case 1.2 Assume that $s_iw \in wW_{I_w}$; by Lemma 4.14 (b), $I_{s_iw} = I_w \setminus \{k\}$, where $\alpha_k = -w^{-1}\alpha_i$. Suppose that $\xi \in \text{wt}(s_iw \Rightarrow u) + (Q_{I_w}^{\vee,+} \setminus Q_{I_{s_iw}}^{\vee,+})$, namely, that the coefficient of α_k^\vee in $\xi - \text{wt}(s_iw \Rightarrow u)$ is greater than 0. Since $u \in \text{EQB}(s_iw)$, we deduce from Proposition 4.15 (2) that $\text{wt}(w \Rightarrow u) = \text{wt}(s_iw \Rightarrow u) + \alpha_k$. Hence we see that $\xi - \text{wt}(w \Rightarrow u) \in Q^{\vee,+}$, which implies that $ut_\xi \succeq w$ by Lemma 4.8. Since $w > s_iw$ by the assumption, the inequality $ut_\xi \succeq w$ contradicts the assumption that $x = ut_\xi \in K_{s_iw}$. Thus, we have shown that $\xi \in \text{wt}(s_iw \Rightarrow u) + Q_{I_{s_iw}}^{\vee,+}$, as desired.

Step 2. We prove the opposite inclusion $K_{s_iw} \supset K'_{s_iw}$. Let $x = ut_\xi \in K'_{s_iw}$, where $u \in \text{EQB}(s_iw)$ and $\xi \in \text{wt}(s_iw \Rightarrow u) + Q_{I_{s_iw}}^{\vee,+}$; note that $s_iu > u$, or equivalently, $u^{-1}\alpha_i \in \Delta^+$ by Lemma 4.19. Hence it follows from Lemma 4.5 (2) that $\text{wt}(s_iw \Rightarrow u) = \text{wt}(w \Rightarrow s_iu)$. Here, by Lemma 4.14, we have $I_{s_iw} \subset I_w$, and hence $Q_{I_{s_iw}}^{\vee,+} \subset Q_{I_w}^{\vee,+}$. Therefore, we have $\xi \in \text{wt}(w \Rightarrow s_iu) + Q_{I_w}^{\vee,+}$. Also, we see from Proposition 4.15 (2) that $s_iu \in \text{EQB}(w)$. Hence we conclude that $s_iut_\xi \in K_w$ by the induction hypothesis; namely, we have $s_iut_\xi \succeq w$ and $s_iut_\xi \not\succeq z$ for any $z \in W$ such that $z > w$. Since $s_iu > u$ and $s_iw < w$, it follows from Lemma 4.11 (3) that $ut_\xi \succeq s_iw$, i.e., $ut_\xi \in (W_{\text{af}})_{\succeq s_iw}$. Suppose, for a contradiction, that $ut_\xi \in (W_{\text{af}})_{\succeq z}$ for some $z \in W$ such that $z > s_iw$. Assume first that $s_iz > z$, or equivalently, $z^{-1}\alpha_i \in \Delta^+$. We see by Lemmas 4.11 (3) and 4.12 (3) that $s_iut_\xi \succeq s_iz > w$, which contradicts the fact that $s_iut_\xi \in K_w$. Assume next that $s_iz < z$, or equivalently, $z^{-1}\alpha_i \in \Delta^-$. We see by Lemma 4.12 (1) that $z > w$. Also, since $s_iu > u$, we have $s_iut_\xi \succeq ut_\xi$ (by the definition of \succeq). Combining these inequalities, we obtain $s_iut_\xi \succeq ut_\xi \succeq z > w$, which contradicts the fact that $s_iut_\xi \in K_w$. Thus we conclude that $ut_\xi \in K_{s_iw}$.

This completes the proof of Proposition 4.16. \square

Proposition 4.22. *Let S be a subset of I , and let $w \in W^S$; recall from Remark 4.13 that $S \subset I_{\lceil w \rceil}$. Then, the subset*

$$K_w^S = ((W^S)_{\text{af}})_{\succeq w} \setminus \bigcup_{z \in W^S, z > w} ((W^S)_{\text{af}})_{\succeq z},$$

defined in (3.14), is identical to the set

$$\Pi^S(K_{\lceil w \rceil}) = \{u\Pi^S(t_\xi) \in (W^S)_{\text{af}} \mid u \in [\text{EQB}(\lceil w \rceil)], \xi \in \text{wt}^S(w \Rightarrow u) + Q_{I_{\lceil w \rceil} \setminus S}^{\vee,+}\}. \quad (4.18)$$

Remark 4.23. The equality in (4.18) follows from Proposition 4.16, Lemma 4.4, and Lemma 2.1 (1), (3).

Remark 4.24. Keep the setting of Proposition 4.22. We see by Lemma 2.1 (3) that the map

$$[\text{EQB}(\lceil w \rceil)] \times Q_{I_{\lceil w \rceil} \setminus S}^{\vee,+} \rightarrow K_w^S, \quad (u, \gamma) \mapsto u\Pi^S(t_{\text{wt}^S(w \Rightarrow u) + \gamma}),$$

is bijective.

Proof of Proposition 4.22. First, we claim that for arbitrary $u \in W^S$ and $u_1 \in uW_S$,

$$\Pi^S((W_{\text{af}})_{\succeq u_1}) = ((W^S)_{\text{af}})_{\succeq u}. \quad (4.19)$$

The inclusion \subset follows from Lemma 4.7 and (2.10). Let us prove the opposite inclusion \supset . Let $x \in ((W^S)_{\text{af}})_{\succeq u}$, and write it as $x = v\Pi^S(t_\xi)$ for some $v \in W^S$ and $\xi \in Q^\vee$. Since $x \succeq u$ by the assumption, we see from Lemma 4.8 that $[\xi]^S \geq \text{wt}^S(u \Rightarrow v)$; note that $\text{wt}^S(u \Rightarrow v) = [\text{wt}(u_1 \Rightarrow v)]^S$ by Lemma 4.4. Hence we can take $\zeta \in Q^\vee$ such that $[\zeta]^S = [\xi]^S$ and $\zeta \geq \text{wt}(u_1 \Rightarrow v)$. We set $y := v t_\zeta \in W_{\text{af}}$. Then it follows from Lemma 4.8 (with $S = \emptyset$) that $y \succeq u_1$. Also, we have $\Pi^S(y) = \Pi^S(v)\Pi^S(t_\zeta) = v\Pi^S(t_\xi) = x$ by Lemma 2.1 (1) and (3). This proves the opposite inclusion \supset , and hence (4.19).

From (4.19) and the definitions of $K_{\lceil w \rceil}$ and K_w^S (by a standard set-theoretical argument), we see easily that $\Pi^S(K_{\lceil w \rceil}) \supset K_w^S$. Let us prove the opposite inclusion \subset . Let $x \in K_{\lceil w \rceil}$; by (4.19), we have $\Pi^S(x) \in ((W^S)_{\text{af}})_{\succeq w}$. Suppose, for a contradiction, that $\Pi^S(x) \in ((W^S)_{\text{af}})_{\succeq z}$ for some $z \in W^S$ such that $z > w$. By Proposition 4.16, we can write the x as $x = ut_\xi$ for some $u \in \text{EQB}(\lceil w \rceil)$ and $\xi \in \text{wt}(\lceil w \rceil \Rightarrow u) + Q_{I_{\lceil w \rceil}}^{\vee,+}$. Hence we have $\Pi^S(x) = [u]\Pi^S(t_\xi) = u_1 t_{\xi + \xi_1}$ for some $u_1 \in uW_S$ and $\xi_1 \in Q_S^\vee$ by Lemma 2.1 (1) and (2). Here, by [NNS2, Lemma 29], the subset $\text{EQB}(\lceil w \rceil)$ of W is a disjoint union of some cosets in $W/W_{I_{\lceil w \rceil}}$; since $S \subset I_{\lceil w \rceil}$, the subset $\text{EQB}(\lceil w \rceil)$ of W is also a disjoint union of some cosets in W/W_S . Hence it follows that u_1 is contained in $\text{EQB}(\lceil w \rceil)$. Also, since $\Pi^S(x) \succeq z$ by our assumption, it follows from Lemma 4.8 that $[\xi]^S \geq \text{wt}^S(z \Rightarrow [u])$; by Lemma 4.4, we have $\text{wt}^S(z \Rightarrow [u]) = [\text{wt}(\lceil z \rceil \Rightarrow u_1)]^S$, and hence $[\xi]^S \geq [\text{wt}(\lceil z \rceil \Rightarrow u_1)]^S$.

We set

$$\zeta := \text{wt}([w] \Rightarrow u_1) + \underbrace{\xi - \text{wt}([w] \Rightarrow u)}_{\in Q_{I_{[w]}}^{\vee,+}} = \xi + \underbrace{\text{wt}([w] \Rightarrow u_1) - \text{wt}([w] \Rightarrow u)}_{\in Q_S^{\vee} \text{ by Lemma 4.4}}.$$

Since $[\xi]^S \geq [\text{wt}([z] \Rightarrow u_1)]^S$, as seen above, we can take $\zeta_1 \in Q_S^{\vee,+}$ such that $\zeta + \zeta_1 \geq \text{wt}([z] \Rightarrow u_1)$; since $S \subset I_{[w]}$, we have $\zeta + \zeta_1 \in \text{wt}([w] \Rightarrow u_1) + Q_{I_{[w]}}^{\vee,+}$. Hence it follows from Proposition 4.16 that $y := u_1 t_{\zeta+\zeta_1} \in K_{[w]}$. However, since $\zeta + \zeta_1 \geq \text{wt}([z] \Rightarrow u_1)$, we deduce by Lemma 4.8 (with $S = \emptyset$) that $y = u_1 t_{\zeta+\zeta_1} \succeq [z] > [w]$, which contradicts the fact that $y \in K_{[w]}$. This proves the opposite inclusion $\Pi^S(K_{[w]}) \subset K_w^S$, and hence completes the proof of Proposition 4.22. \square

5 Proof of Theorem 2.8.

5.1 Quantum Lakshmibai-Seshadri paths and the degree function.

We fix $\lambda \in P^+$, and take $S = S_\lambda$ as in (2.18).

Definition 5.1. For a rational number $0 < a < 1$, we define $\text{QBG}_{a\lambda}(W^S)$ to be the subgraph of $\text{QBG}(W^S)$ with the same vertex set but having only those edges of the form $u \xrightarrow{\beta} v$ for which $a\langle \lambda, \beta^\vee \rangle \in \mathbb{Z}$ holds.

Definition 5.2. A quantum LS path of shape λ is a pair

$$\eta = (\mathbf{w}; \mathbf{a}) = (w_1, \dots, w_s; a_0, a_1, \dots, a_s), \quad s \geq 1, \quad (5.1)$$

of a sequence $\mathbf{w} : w_1, \dots, w_s$ of elements in W^S with $w_u \neq w_{u+1}$ for any $1 \leq u \leq s-1$ and an increasing sequence $\mathbf{a} : 0 = a_0 < a_1 < \dots < a_s = 1$ of rational numbers satisfying the condition that there exists a directed path from w_{u+1} to w_u in $\text{QBG}_{a_u\lambda}(W^S)$ for each $u = 1, 2, \dots, s-1$.

We denote by $\text{QLS}(\lambda)$ the set of all quantum LS paths of shape λ . If $\eta \in \text{QLS}(\lambda)$ is of the form (5.1), then we set $\kappa(\eta) := w_s \in W^S$, and call it the final direction of η . For $w \in W$, we set (see [NNS1, Sect. 3.2] and [NNS2, Sect. 2.3])

$$\text{QLS}^{w\lambda, \infty}(\lambda) := \{\eta \in \text{QLS}(\lambda) \mid \kappa(\eta) \in [\text{EQB}[w]]\}. \quad (5.2)$$

We define a projection $\text{cl} : (W^S)_{\text{af}} \rightarrow W^S$ by $\text{cl}(x) := w$ for $x \in (W^S)_{\text{af}}$ of the form $x = w\Pi^S(t_\xi)$ with $w \in W^S$ and $\xi \in Q^\vee$. For $\pi = (x_1, \dots, x_s; a_0, a_1, \dots, a_s) \in \mathbb{B}^{\frac{\infty}{2}}(\lambda)$, we define

$$\text{cl}(\pi) := (\text{cl}(x_1), \dots, \text{cl}(x_s); a_0, a_1, \dots, a_s);$$

here, for each $1 \leq p < q \leq s$ such that $\text{cl}(x_p) = \cdots = \text{cl}(x_q)$, we drop $\text{cl}(x_p), \dots, \text{cl}(x_{q-1})$ and a_p, \dots, a_{q-1} . We know from [NS, Sect. 6.2] that $\text{cl}(\pi) \in \text{QLS}(\lambda)$ for all $\pi \in \mathbb{B}_0^{\infty}(\lambda)$, and that the map $\text{cl} : \mathbb{B}_0^{\infty}(\lambda) \rightarrow \text{QLS}(\lambda)$, $\pi \mapsto \text{cl}(\pi)$, is surjective. We also know the following lemma from [NS, Lemma 6.2.3]; recall that $\mathbb{B}_0^{\infty}(\lambda)$ denotes the connected component of $\mathbb{B}_0^{\infty}(\lambda)$ containing $\pi_\lambda = (e; 0, 1)$.

Lemma 5.3. *For each $\eta \in \text{QLS}(\lambda)$, there exists a unique $\pi_\eta \in \mathbb{B}_0^{\infty}(\lambda)$ such that $\text{cl}(\pi_\eta) = \eta$ and $\kappa(\pi_\eta) = \kappa(\eta) \in W^S$.*

Now we define the (tail) degree function $\text{deg}_\lambda : \text{QLS}(\lambda) \rightarrow \mathbb{Z}_{\leq 0}$ as follows. Let $\eta \in \text{QLS}(\lambda)$, and take $\pi_\eta \in \mathbb{B}_0^{\infty}(\lambda)$ as in Lemma 5.3. We see from the argument in [NS, Sect. 6.2] that $\text{wt}(\pi_\eta) = \lambda - \gamma + k\delta$ for some $\gamma \in Q^+$ and $k \in \mathbb{Z}_{\leq 0}$. Then we set $\text{deg}_\lambda(\eta) := k$. We know the following description of deg_λ from [LNS³², Corollary 4.8]; for the definition of $\text{wt}^S(u \Rightarrow v)$, see Section 4.1.

Proposition 5.4. *For $\eta = (w_1, \dots, w_s; a_0, a_1, \dots, a_s) \in \text{QLS}(\lambda)$, it holds that*

$$\text{deg}_\lambda(\eta) = - \sum_{u=1}^{s-1} a_u \langle \lambda, \text{wt}^S(w_{u+1} \Rightarrow w_u) \rangle. \quad (5.3)$$

Also, for $\eta = (w_1, \dots, w_s; a_0, a_1, \dots, a_s) \in \text{QLS}(\lambda)$ and $w \in W^S$, we define the degree of η at $w\lambda$ (see [NNS1, Sect. 3.2] and [NNS2, Sect. 2.3]) by

$$\text{deg}_{w\lambda}(\eta) := - \sum_{u=1}^s a_u \langle \lambda, \text{wt}^S(w_{u+1} \Rightarrow w_u) \rangle, \quad \text{with } w_{s+1} := w. \quad (5.4)$$

Theorem 5.5 ([NNS1, Theorem 3.2.7]). *Let $\lambda \in P^+$, and take $S = S_\lambda$ as in (2.18). Then,*

$$\sum_{\eta \in \text{QLS}^{w\lambda, \infty}(\lambda)} e^{\text{wt}(\eta)} q^{\text{deg}_{w\lambda}(\eta)} = E_{w\lambda}(q, \infty) \quad \text{for } w \in W^S. \quad (5.5)$$

5.2 Proof of the graded character formula for $\mathbb{K}_w^-(\lambda)$.

Let $\lambda = \sum_{i \in I} m_i \varpi_i \in P^+$, and take $S = S_\lambda$ as in (2.18). Let $w \in W^S$. Recall from (3.12) that

$$\text{gch } \mathbb{K}_w^-(\lambda) = \sum_{\pi \in \mathbb{B}_{\geq w; \neq w}^{\infty}(\lambda)} e^{\text{fin}(\text{wt}(\pi))} q^{\text{nul}(\text{wt}(\pi))},$$

where $\mathbb{B}_{\geq w; \neq w}^{\infty}(\lambda) = \{\pi \in \mathbb{B}_0^{\infty}(\lambda) \mid \kappa(\pi) \in K_w^S\}$ by (3.15). Because $\mathbb{B}_{\geq w; \neq w}^{\infty}(\lambda) = \bigsqcup_{\eta \in \text{QLS}(\lambda)} (\text{cl}^{-1}(\eta) \cap \mathbb{B}_{\geq w; \neq w}^{\infty}(\lambda))$, we deduce that

$$\text{gch } \mathbb{K}_w^-(\lambda) = \sum_{\eta \in \text{QLS}(\lambda)} \underbrace{\left(\sum_{\pi \in \text{cl}^{-1}(\eta) \cap \mathbb{B}_{\geq w; \neq w}^{\infty}(\lambda)} e^{\text{fin}(\text{wt}(\pi))} q^{\text{nul}(\text{wt}(\pi))} \right)}_{=: F_\eta}. \quad (5.6)$$

Let us compute F_η for each $\eta \in \text{QLS}(\lambda)$. We fix an arbitrary $\eta \in \text{QLS}(\lambda)$, and take $\pi_\eta \in \mathbb{B}_0^{\frac{\infty}{2}}(\lambda)$ as in Lemma 5.3. We take and fix a monomial X_η in root operators e_i and f_i , $i \in I_{\text{af}}$, such that $\pi_\eta = X_\eta \pi_\lambda$; we see by [NS, Lemma 6.2.2] that

$$\text{cl}^{-1}(\eta) = \{X_\eta(t_\xi \cdot \pi^C) \in \mathbb{B}^{\frac{\infty}{2}}(\lambda) \mid C \in \text{Conn}(\mathbb{B}^{\frac{\infty}{2}}(\lambda)), \xi \in Q_{I \setminus S}^\vee\}; \quad (5.7)$$

for the definitions of $\pi^C \in \mathbb{B}^{\frac{\infty}{2}}(\lambda)$ and $x \cdot \pi$ with $x \in W_{\text{af}}$ and $\pi \in \mathbb{B}^{\frac{\infty}{2}}(\lambda)$, see (3.11) and Remark 3.6, respectively. By the argument after [NNS1, (5.4)], we see that

$$\kappa(X_\eta(t_\xi \cdot \pi^C)) = \kappa(\eta) \Pi^S(t_\xi) \quad \text{for } C \in \text{Conn}(\mathbb{B}^{\frac{\infty}{2}}(\lambda)) \text{ and } \xi \in Q_{I \setminus S}^\vee. \quad (5.8)$$

Because

$$K_w^S = \{u \Pi^S(t_\xi) \in (W^S)_{\text{af}} \mid u \in [\text{EQB}(\lceil w \rceil)], \xi \in \text{wt}^S(w \Rightarrow u) + Q_{I_{\lceil w \rceil} \setminus S}^{\vee,+}\}$$

by Proposition 4.22, we deduce the following:

- (i) $\text{cl}^{-1}(\eta) \cap \mathbb{B}_{\geq w; \not\geq w}^{\frac{\infty}{2}}(\lambda) \neq \emptyset \iff \kappa(\eta) \in [\text{EQB}(\lceil w \rceil)] \iff \eta \in \text{QLS}^{w\lambda, \infty}(\lambda)$; the implication \Leftarrow in the first equivalence follows from (5.8) and the fact that $\text{wt}^S(w \Rightarrow u) + Q_{I_{\lceil w \rceil} \setminus S}^{\vee,+} \subset Q_{I \setminus S}^\vee$.

- (ii) If $\eta \in \text{QLS}^{w\lambda, \infty}(\lambda)$, then

$$\begin{aligned} & \text{cl}^{-1}(\eta) \cap \mathbb{B}_{\geq w; \not\geq w}^{\frac{\infty}{2}}(\lambda) \\ &= \left\{ X_\eta(t_\xi \cdot \pi^C) \in \mathbb{B}^{\frac{\infty}{2}}(\lambda) \mid \begin{array}{l} C \in \text{Conn}(\mathbb{B}^{\frac{\infty}{2}}(\lambda)), \\ \xi \in \text{wt}^S(w \Rightarrow \kappa(\eta)) + Q_{I_{\lceil w \rceil} \setminus S}^{\vee,+} \end{array} \right\}; \end{aligned} \quad (5.9)$$

remark that for $C, C' \in \text{Conn}(\mathbb{B}^{\frac{\infty}{2}}(\lambda))$ and $\xi, \xi' \in \text{wt}^S(w \Rightarrow \kappa(\eta)) + Q_{I_{\lceil w \rceil} \setminus S}^{\vee,+}$,

$$X_\eta(t_\xi \cdot \pi^C) = X_\eta(t_{\xi'} \cdot \pi^{C'}) \iff C = C' \text{ and } \xi = \xi'. \quad (5.10)$$

Indeed, the implication \Leftarrow is obvious. Let us show the implication \Rightarrow . Since $X_\eta(t_\xi \cdot \pi^C) \in C$ and $X_\eta(t_{\xi'} \cdot \pi^{C'}) \in C'$, we have $C = C'$. Also, since $\kappa(\eta) \Pi^S(t_\xi) = \kappa(X_\eta(t_\xi \cdot \pi^C)) = \kappa(X_\eta(t_{\xi'} \cdot \pi^{C'})) = \kappa(\eta) \Pi^S(t_{\xi'})$, as seen above, we deduce that $\xi - \xi' \in Q_S^\vee$ by Lemma 2.1 (3). Therefore, by the assumption that $\xi, \xi' \in \text{wt}^S(w \Rightarrow \kappa(\eta)) + Q_{I_{\lceil w \rceil} \setminus S}^{\vee,+}$, we obtain $\xi = \xi'$.

Let $C \in \text{Conn}(\mathbb{B}^{\frac{\infty}{2}}(\lambda))$, and write $\Theta(C) \in \text{Par}(\lambda)$ as $\Theta(C) = (\rho^{(i)})_{i \in I}$, where $\rho^{(i)} = (\rho_1^{(i)} \geq \dots \geq \rho_{m_i-1}^{(i)} \geq \rho_{m_i}^{(i)} = 0)$ for each $i \in I \setminus S$, and $\rho^{(i)} = \emptyset$ for all $i \in S$. Also, let $\xi \in \text{wt}^S(w \Rightarrow \kappa(\eta)) + Q_{I_{\lceil w \rceil} \setminus S}^{\vee,+}$, and write it as:

$$\xi = \text{wt}^S(w \Rightarrow \kappa(\eta)) + \sum_{i \in I_{\lceil w \rceil} \setminus S} c_i \alpha_i^\vee,$$

where $c_i \in \mathbb{Z}_{\geq 0}$ for $i \in I_{[w]} \setminus S$; by convention, we set $c_i := 0$ for all $i \in S$. For each $i \in I_{[w]}$, we set $\tilde{\rho}^{(i)} := (c_i + \rho_1^{(i)} \geq \cdots \geq c_i + \rho_{m_i-1}^{(i)} \geq c_i + \rho_{m_i-1}^{(i)} = c_i)$, which is a partition of length less than or equal to m_i ; note that $\tilde{\rho}^{(i)} = \emptyset$ for all $i \in S$. Also, for each $i \in I \setminus I_{[w]}$, we set $\tilde{\rho}^{(i)} := \rho^{(i)}$, which is a partition of length less than m_i . Then, $(\tilde{\rho}^{(i)})_{i \in I}$ is an element of the set $\text{Par}_w(\lambda)$ of I -tuples $\boldsymbol{\chi} = (\chi^{(i)})_{i \in I}$ of partitions such that for each $i \in I_{[w]}$ (resp., $i \in I \setminus I_{[w]}$), $\chi^{(i)}$ is a partition of length less than or equal to (resp., less than) m_i . By the same calculations as those after [NS, (6.3.3)] and [NNS1, (5.5)], we deduce that

$$\text{wt}(X(t_\xi \cdot \pi^C)) = \text{wt}(\eta) + \underbrace{(\deg_\lambda(\eta) - \langle \lambda, \text{wt}^S(w \Rightarrow \kappa(\eta)) \rangle)}_{= \deg_{w\lambda}(\eta) \text{ by (5.3) and (5.4)}} \delta - |(\tilde{\rho}^{(i)})_{i \in I}| \delta. \quad (5.11)$$

Summarizing the above, for each $\eta \in \text{QLS}^{w\lambda, \infty}(\lambda)$,

$$\begin{aligned} F_\eta &= \sum_{\pi \in \text{cl}^{-1}(\eta) \cap \mathbb{B}_{\sum w; \neq w}^{\infty}(\lambda)} \mathbf{e}^{\text{fin}(\text{wt}(\pi))} q^{\text{nul}(\text{wt}(\pi))} \\ &= \sum_{\substack{C \in \text{Conn}(\mathbb{B}_{\sum w}^{\infty}(\lambda)) \\ \xi \in \text{wt}^S(w \Rightarrow \kappa(\eta)) + Q_{I_{[w]} \setminus S}^{\vee, +}}} \mathbf{e}^{\text{fin}(\text{wt}(X_\eta(t_\xi \cdot \pi^C)))} q^{\text{nul}(\text{wt}(X_\eta(t_\xi \cdot \pi^C)))} \quad \text{by (5.9) and (5.10)} \\ &= \mathbf{e}^{\text{wt}(\eta)} q^{\deg_{w\lambda}(\eta)} \sum_{\boldsymbol{\chi} \in \text{Par}_w(\lambda)} q^{-|\boldsymbol{\chi}|} \quad \text{by (5.11),} \end{aligned}$$

where $|\boldsymbol{\chi}| = \sum_{i \in I} |\chi^{(i)}|$ for $\boldsymbol{\chi} = (\chi^{(i)})_{i \in I} \in \text{Par}_w(\lambda)$ (see Section 3.2). Here we recall that if $\boldsymbol{\chi} = (\chi^{(i)})_{i \in I} \in \text{Par}_w(\lambda)$, then the partition $\chi^{(i)}$ is a partition of length less than or equal to $m_i - \epsilon_i$ for each $i \in I$, where ϵ_i is as in (2.29). Therefore, we deduce that

$$F_\eta = \mathbf{e}^{\text{wt}(\eta)} q^{\deg_{w\lambda}(\eta)} \sum_{\boldsymbol{\chi} \in \text{Par}_w(\lambda)} q^{-|\boldsymbol{\chi}|} = \mathbf{e}^{\text{wt}(\eta)} q^{\deg_{w\lambda}(\eta)} \left(\prod_{i \in I} \prod_{r=1}^{m_i - \epsilon_i} (1 - q^{-r}) \right)^{-1}.$$

Also, we have $F_\eta = 0$ for all $\eta \in \text{QLS}(\lambda) \setminus \text{QLS}^{w\lambda, \infty}(\lambda)$, since $\text{cl}^{-1} \cap \mathbb{B}_{\sum w; \neq w}^{\infty}(\lambda) = \emptyset$, as seen in (i) above. Substituting these F_η 's into (5.6), we obtain

$$\begin{aligned} \text{gch } \mathbb{K}_w^-(\lambda) &= \sum_{\eta \in \text{QLS}^{w\lambda, \infty}(\lambda)} \mathbf{e}^{\text{wt}(\eta)} q^{\deg_{w\lambda}(\eta)} \left(\prod_{i \in I} \prod_{r=1}^{m_i - \epsilon_i} (1 - q^{-r}) \right)^{-1} \\ &= \left(\prod_{i \in I} \prod_{r=1}^{m_i - \epsilon_i} (1 - q^{-r}) \right)^{-1} E_{w\lambda}(q, \infty) \quad \text{by Theorem 5.5.} \end{aligned}$$

This completes the proof of equation (2.28).

5.3 Proof of the graded character formula for $\mathbf{K}_w^-(\lambda)$.

Let $\lambda \in P^+$, and take $S = S_\lambda$ as in (2.18). Let $w \in W^S$. For each $\eta \in \text{QLS}(\lambda)$, we take and fix a monomial X_η in root operators e_i and f_i , $i \in I_{\text{af}}$, such that $\pi_\eta = X_\eta \pi_\lambda$, where we take π_η as in Lemma 5.3. We see from [NNS1, Theorem 5.12] that under the isomorphism $\Psi_\lambda : \mathcal{B}(\lambda) \xrightarrow{\sim} \mathbb{B}_{\geq w}^{\infty}(\lambda)$ of crystals (see Theorem 3.5), the subset $\mathcal{B}(X_w^-(\lambda))$ of $\mathcal{B}(\lambda)$ (see (2.27)) is mapped to

$$\mathbb{X}_w^-(\lambda) := \mathbb{B}_{\geq w}^{\infty}(\lambda) \setminus \{X_\eta(t_{\text{wt}^S(w \Rightarrow \kappa(\eta))} \cdot \pi_\lambda) \mid \eta \in \text{QLS}(\lambda)\}. \quad (5.12)$$

Therefore, we have

$$\text{gch } \mathbf{K}_w^-(\lambda) = \sum_{\pi \in \mathbb{B}_{\geq w; \not\geq w}^{\infty}(\lambda) \setminus \mathbb{X}_w^-(\lambda)} \mathbf{e}^{\text{fin}(\text{wt}(\pi))} q^{\text{nul}(\text{wt}(\pi))}. \quad (5.13)$$

Here, by (i) and (ii) in the previous subsection, we see that

$$\mathbb{B}_{\geq w; \not\geq w}^{\infty}(\lambda) = \left\{ X_\eta(t_\xi \cdot \pi^C) \in \mathbb{B}_{\geq w}^{\infty}(\lambda) \mid \begin{array}{l} C \in \text{Conn}(\mathbb{B}_{\geq w}^{\infty}(\lambda)), \eta \in \text{QLS}^{w\lambda, \infty}(\lambda), \\ \xi \in \text{wt}^S(w \Rightarrow \kappa(\eta)) + Q_{I_{[w]}^V, +} \end{array} \right\}.$$

From this, we obtain

$$\mathbb{B}_{\geq w; \not\geq w}^{\infty}(\lambda) \setminus \mathbb{X}_w^-(\lambda) = \{X_\eta(t_{\text{wt}^S(w \Rightarrow \kappa(\eta))} \cdot \pi_\lambda) \mid \eta \in \text{QLS}^{w\lambda, \infty}(\lambda)\}. \quad (5.14)$$

Combining (5.13) and (5.14), we conclude by the same computation as in the previous subsection (this time, only the term corresponding to $\chi = (\emptyset)_{i \in I} \in \text{Par}_w(\lambda)$ remains) that

$$\text{gch } \mathbf{K}_w^-(\lambda) = \sum_{\eta \in \text{QLS}^{w\lambda, \infty}(\lambda)} \mathbf{e}^{\text{wt}(\eta)} q^{\text{deg}_{w\lambda}(\eta)} = E_{w\lambda}(q, \infty).$$

This completes the proof of equation (2.30).

Appendix.

A Recursive proof of the formula (2.28) for $\mathbb{K}_w^-(\lambda)$.

Proposition A.1. *Let S be a subset of I . For each $x \in (W^S)_{\text{af}}$ of the form $x = w\Pi^S(t_\xi)$ with $w \in W^S$ and $\xi \in Q^{V, +} = \sum_{i \in I} \mathbb{Z}_{\geq 0} \alpha_i^\vee$, the subset $\{v \in W^S \mid v \leq x\}$ of W^S has a unique maximal element in the Bruhat order \leq on W , which we denote by v_x .*

Proof. By Lemma 4.7 and [BB, Proposition 2.5.1], it suffices to prove the assertion in the case that $S = \emptyset$.

We take and fix $\gamma \in Q^\vee$ such that $\langle \gamma, \alpha_i^\vee \rangle > 0$ for all $i \in I$. Since $W \cup \{x\}$ is a finite set, we deduce from [S, Claim 4.14] (see also [P, Lectures 12, 13] and [INS, Appendix A.3]) that there exists $N \in \mathbb{Z}_{\geq 0}$ such that for $y, z \in W \cup \{x\}$,

- (a) if $y \preceq z$, then $yt_{n\gamma} \leq zt_{n\gamma}$ for all $n \geq N$ in the (ordinary) Bruhat order \geq on W_{af} ;
- (b) if $yt_{n\gamma} \leq zt_{n\gamma}$ for some $n \geq N$, then $y \preceq z$.

Let $n \geq N$. Because W is a parabolic subgroup of W_{af} , it follows from [LLM, Lemma 11 (ii)] (note that $t_{n\gamma} \leq xt_{n\gamma}$ by (a) since $n \geq N$ and $e \preceq x$) that the subset $\{vt_{n\gamma} \mid v \in W, vt_{n\gamma} \leq xt_{n\gamma}\}$ of W_{af} has a unique maximal element in the Bruhat order on W_{af} , which we write as $v_n t_{n\gamma}$ with $v_n \in W$. Here we see by (a) and (b) that v_n does not depend on $n \geq N$, and is a unique maximal element of $\{v \in W \mid v \preceq x\}$ in the semi-infinite Bruhat order \preceq on W_{af} (and hence in the Bruhat order on W ; see Remark 2.5). This proves the proposition. \square

Let S be a subset of I . For $w \in W^S$, we set $W_{\geq w}^S := \{v \in W^S \mid v \geq w\}$. Also, recall from (3.14) the definition of K_v^S for $v \in W^S$.

Lemma A.2. *Keep the notation and setting above. It holds that*

$$((W^S)_{\text{af}})_{\succeq w} = \bigsqcup_{v \in W_{\geq w}^S} K_v^S. \quad (\text{A.1})$$

Proof. First we prove the equality:

$$((W^S)_{\text{af}})_{\succeq w} = \bigcup_{v \in W_{\geq w}^S} K_v^S. \quad (\text{A.2})$$

Let $x \in ((W^S)_{\text{af}})_{\succeq w}$; note that $w \in \{v \in W^S \mid v \preceq x\}$. By Proposition A.1, the subset $\{v \in W^S \mid v \preceq x\}$ of W^S has a unique maximal element v_x . Hence it follows from the definition of v_x that $w \leq v_x \preceq x$; in particular, $x \in ((W^S)_{\text{af}})_{\succeq v_x}$. If $x \notin K_{v_x}^S$, then $x \in ((W^S)_{\text{af}})_{\succeq u}$ for some $u \in W^S$ such that $u > v_x$. Since $u \in \{v \in W^S \mid v \preceq x\}$, this contradicts the maximality of v_x . Hence, $x \in K_{v_x}^S \subset \bigcup_{v \in W_{\geq w}^S} K_v^S$. Thus we have shown the inclusion \subset . Let us show the opposite inclusion \supset . Let $x \in \bigcup_{v \in W_{\geq w}^S} K_v^S$, and take $v \in W_{\geq w}^S$ such that $x \in K_v^S$. Then, by the definition of K_v^S , we have $x \in ((W^S)_{\text{af}})_{\succeq v}$. Since $v \geq w$, we have $((W^S)_{\text{af}})_{\succeq v} \subset ((W^S)_{\text{af}})_{\succeq w}$, and hence $x \in ((W^S)_{\text{af}})_{\succeq w}$. Thus we have shown the inclusion \supset , as desired.

The remaining task is to prove that $K_{v_1}^S \cap K_{v_2}^S = \emptyset$ for all $v_1, v_2 \in W^S$ with $v_1 \neq v_2$. Suppose, for a contradiction, that $K_{v_1}^S \cap K_{v_2}^S \neq \emptyset$ for some $v_1, v_2 \in W^S$ with $v_1 \neq v_2$, and let $x \in K_{v_1}^S \cap K_{v_2}^S$; note that x is of the form $x = w\Pi^S(t_\xi)$ for some $w \in W^S$ and $\xi \in Q^{\vee,+}$ since $x \succeq v_1 \in W^S$ (see Lemmas 4.8 and 2.1(3)). Therefore, by Proposition A.1, the subset $\{v \in W^S \mid v \preceq x\}$ of W^S has a unique maximal element v_x in the Bruhat order \leq on W . Since $x \in K_{v_1}^S \subset ((W^S)_{\text{af}})_{\succeq v_1}$, we have $v_1 \in \{v \in W^S \mid v \preceq x\}$, and hence $v_1 \leq v_x$ by the maximality of v_x . Since $x \in ((W^S)_{\text{af}})_{\succeq v_x}$ by the definition of v_x and $v_x \geq v_1$, and since $x \in K_{v_1}^S$, we see by the definition of $K_{v_1}^S$ that $v_1 = v_x$. Similarly, we see that $v_2 = v_x$, and hence $v_1 = v_2$, which is a contradiction. This proves the lemma. \square

We fix $\lambda \in P^+ \subset P_{\text{af}}^0$ (see (2.1) and (2.2)), and take $S = S_\lambda$ as in (2.18). By Lemma A.2 and (3.15), we have

$$\text{gch } V_w^-(\lambda) = \sum_{v \in W_{\geq w}^S} \text{gch } \mathbb{K}_v^-(\lambda) \quad \text{for each } w \in W^S. \quad (\text{A.3})$$

For each $i \in I$, we define a $\mathbb{C}(q)$ -linear operator D_i on $\mathbb{C}(q)[P]$ by

$$D_i \mathbf{e}^\nu := \frac{\mathbf{e}^{\nu-\rho} - \mathbf{e}^{s_i(\nu-\rho)}}{1 - \mathbf{e}^{\alpha_i}} \mathbf{e}^\rho = \frac{\mathbf{e}^\nu - \mathbf{e}^{\alpha_i} \mathbf{e}^{s_i \nu}}{1 - \mathbf{e}^{\alpha_i}} \quad \text{for } \nu \in P, \text{ where } q = \mathbf{e}^\delta;$$

note that $D_i^2 = D_i$, and that

$$D_i \mathbf{e}^\nu = \begin{cases} \mathbf{e}^\nu (1 + \mathbf{e}^{\alpha_i} + \mathbf{e}^{2\alpha_i} + \dots + \mathbf{e}^{-\langle \nu, \alpha_i^\vee \rangle \alpha_i}) & \text{if } \langle \nu, \alpha_i^\vee \rangle \leq 0, \\ 0 & \text{if } \langle \nu, \alpha_i^\vee \rangle = 1, \\ -\mathbf{e}^\nu (\mathbf{e}^{-\alpha_i} + \mathbf{e}^{-2\alpha_i} + \dots + \mathbf{e}^{-(\langle \nu, \alpha_i^\vee \rangle + 1)\alpha_i}) & \text{if } \langle \nu, \alpha_i^\vee \rangle \geq 2. \end{cases} \quad (\text{A.4})$$

Also, we define a $\mathbb{Q}(q)$ -linear operator T_i on $\mathbb{C}(q)[P]$ by

$$T_i := D_i - 1, \quad \text{that is,} \quad T_i \mathbf{e}^\nu = \frac{\mathbf{e}^{\alpha_i} (\mathbf{e}^\nu - \mathbf{e}^{s_i \nu})}{1 - \mathbf{e}^{\alpha_i}};$$

note that $T_i^2 = -T_i$ and $T_i D_i = D_i T_i = 0$.

Proposition A.3 ([NS]; see also [NOS, Proposition 6.6 and Remark 6.7]). *Let $w \in W^S$ and $i \in I$. Then,*

$$T_i \text{gch } V_w^-(\lambda) = \begin{cases} \text{gch } V_{s_i w}^-(\lambda) - \text{gch } V_w^-(\lambda) & \text{if } \langle w\lambda, \alpha_i^\vee \rangle < 0, \\ 0 & \text{if } \langle w\lambda, \alpha_i^\vee \rangle \geq 0. \end{cases} \quad (\text{A.5})$$

Proposition A.4. *Let $w \in W^S$ and $i \in I$. Then,*

$$T_i \text{gch } \mathbb{K}_w^-(\lambda) = \begin{cases} \text{gch } \mathbb{K}_{s_i w}^-(\lambda) & \text{if } \langle w\lambda, \alpha_i^\vee \rangle < 0, \\ -\text{gch } \mathbb{K}_w^-(\lambda) & \text{if } \langle w\lambda, \alpha_i^\vee \rangle > 0, \\ 0 & \text{if } \langle w\lambda, \alpha_i^\vee \rangle = 0. \end{cases} \quad (\text{A.6})$$

Proof. We show (A.6) by descending induction on w . Assume first that $w = [w_\circ]$; note that $\langle [w_\circ]\lambda, \alpha_i^\vee \rangle = \langle w_\circ\lambda, \alpha_i^\vee \rangle \leq 0$. Also, by (A.3), we have $\text{gch } V_{[w_\circ]}^-(\lambda) = \text{gch } \mathbb{K}_{[w_\circ]}^-(\lambda)$. If $\langle [w_\circ]\lambda, \alpha_i^\vee \rangle < 0$, then it follows from (A.5) that

$$\begin{aligned} T_i \text{gch } \mathbb{K}_{[w_\circ]}^-(\lambda) &= T_i \text{gch } V_{[w_\circ]}^-(\lambda) = \text{gch } V_{s_i [w_\circ]}^-(\lambda) - \text{gch } V_{[w_\circ]}^-(\lambda) \\ &= \text{gch } \mathbb{K}_{s_i [w_\circ]}^-(\lambda), \end{aligned}$$

where the last equality follows from (A.3) and the fact that $W_{\geq s_i [w_\circ]}^S = \{s_i [w_\circ], [w_\circ]\}$. If $\langle [w_\circ]\lambda, \alpha_i^\vee \rangle = 0$, then it follows from (A.5) that

$$T_i \text{gch } \mathbb{K}_{[w_\circ]}^-(\lambda) = T_i \text{gch } V_{[w_\circ]}^-(\lambda) = 0.$$

This proves (A.6) in the case that $w = \lfloor w_o \rfloor$.

Assume next that $w < \lfloor w_o \rfloor$. We set

$$\begin{aligned} (W_{\geq w}^S)_0 &:= \{v \in W_{\geq w}^S \mid \langle v\lambda, \alpha_i^\vee \rangle = 0\}, & (W_{\geq w}^S)_+ &:= \{v \in W_{\geq w}^S \mid \langle v\lambda, \alpha_i^\vee \rangle > 0\}, \\ (W_{\geq w}^S)'_- &:= \{v \in W_{\geq w}^S \mid \langle v\lambda, \alpha_i^\vee \rangle < 0 \text{ and } s_i v \in W_{\geq w}^S\}, \\ (W_{\geq w}^S)''_- &:= \{v \in W_{\geq w}^S \mid \langle v\lambda, \alpha_i^\vee \rangle < 0 \text{ and } s_i v \notin W_{\geq w}^S\}; \end{aligned}$$

observe that

$$W_{\geq w}^S = (W_{\geq w}^S)_0 \sqcup (W_{\geq w}^S)_+ \sqcup (W_{\geq w}^S)'_- \sqcup (W_{\geq w}^S)''_-, \quad (W_{\geq w}^S)'_- = s_i(W_{\geq w}^S)_+.$$

Case 1. Assume that $\langle w\lambda, \alpha_i^\vee \rangle < 0$; note that $w \in (W_{\geq w}^S)''_-$. By (A.3), we have

$$\mathbb{T}_i \text{gch } V_w^-(\lambda) = \sum_{v \in W_{\geq w}^S} \mathbb{T}_i \text{gch } \mathbb{K}_v^-(\lambda). \quad (\text{A.7})$$

The right-hand side of (A.7) is equal to:

$$\begin{aligned} \sum_{v \in (W_{\geq w}^S)_0} \mathbb{T}_i \text{gch } \mathbb{K}_v^-(\lambda) + \sum_{v \in (W_{\geq w}^S)_+} \mathbb{T}_i \text{gch } \mathbb{K}_v^-(\lambda) \\ + \sum_{v \in (W_{\geq w}^S)'_-} \mathbb{T}_i \text{gch } \mathbb{K}_v^-(\lambda) + \sum_{v \in (W_{\geq w}^S)''_-} \mathbb{T}_i \text{gch } \mathbb{K}_v^-(\lambda). \end{aligned}$$

Here, by the induction hypothesis, we see that

$$\begin{aligned} \sum_{v \in (W_{\geq w}^S)_0} \mathbb{T}_i \text{gch } \mathbb{K}_v^-(\lambda) = 0, \quad \sum_{v \in (W_{\geq w}^S)_+} \mathbb{T}_i \text{gch } \mathbb{K}_v^-(\lambda) = - \sum_{v \in (W_{\geq w}^S)_+} \text{gch } \mathbb{K}_v^-(\lambda), \\ \sum_{v \in (W_{\geq w}^S)'_-} \mathbb{T}_i \text{gch } \mathbb{K}_v^-(\lambda) = \sum_{v \in (W_{\geq w}^S)'_-} \text{gch } \mathbb{K}_{s_i v}^-(\lambda) = \sum_{v \in (W_{\geq w}^S)_+} \text{gch } \mathbb{K}_v^-(\lambda). \end{aligned}$$

Therefore, the right-hand side of (A.7) is equal to $\sum_{v \in (W_{\geq w}^S)''_-} \mathbb{T}_i \text{gch } \mathbb{K}_v^-(\lambda)$. Hence, again by the induction hypothesis, we deduce that

$$\mathbb{T}_i \text{gch } V_w^-(\lambda) = \mathbb{T}_i \text{gch } \mathbb{K}_w^-(\lambda) + \sum_{v \in (W_{\geq w}^S)''_-, v \neq w} \text{gch } \mathbb{K}_{s_i v}^-(\lambda). \quad (\text{A.8})$$

Also, by (A.5) and (A.3), we have

$$\mathbb{T}_i \text{gch } V_w^-(\lambda) = \text{gch } V_{s_i w}^-(\lambda) - \text{gch } V_w^-(\lambda) = \sum_{v \in W_{\geq s_i w}^S \setminus W_{\geq w}^S} \text{gch } \mathbb{K}_v^-(\lambda). \quad (\text{A.9})$$

We claim that $W_{\geq s_i w}^S \setminus W_{\geq w}^S = s_i(W_{\geq w}^S)''_-$. Let $v \in W_{\geq s_i w}^S \setminus W_{\geq w}^S$. If $\langle v\lambda, \alpha_i^\vee \rangle \leq 0$, then we deduce from Lemma 4.12(1), together with the assumption $\langle s_i w\lambda, \alpha_i^\vee \rangle > 0$ of Case 1,

that $v \in W_{\geq w}^S$, which is a contradiction. As a result, we have $\langle v\lambda, \alpha_i^\vee \rangle > 0$, and hence $\langle s_i v \lambda, \alpha_i^\vee \rangle < 0$. Since $\langle s_i w \lambda, \alpha_i^\vee \rangle > 0$, it follows from Lemma 4.12 (3) that $s_i v \in W_{\geq w}^S$. Since $s_i(s_i v) = v \notin W_{\geq w}^S$, we obtain $s_i v \in (W_{\geq w}^S)''$, and hence $v \in s_i(W_{\geq w}^S)''$. Thus we have shown $W_{\geq s_i w}^S \setminus W_{\geq w}^S \subset s_i(W_{\geq w}^S)''$. In order to show the opposite inclusion, let $v \in (W_{\geq w}^S)''$. Then we see from Lemma 4.12 (3) that $s_i v \in W_{\geq s_i w}^S$. In addition, by the definition of $(W_{\geq w}^S)''$, we have $s_i v \notin W_{\geq w}^S$. Thus we obtain $s_i v \in W_{\geq s_i w}^S \setminus W_{\geq w}^S$, as desired.

By the equality $W_{\geq s_i w}^S \setminus W_{\geq w}^S = s_i(W_{\geq w}^S)''$ and (A.9), we see that

$$\begin{aligned} \mathsf{T}_i \text{gch } V_w^-(\lambda) &= \sum_{v \in s_i(W_{\geq w}^S)''} \text{gch } \mathbb{K}_v^-(\lambda) = \sum_{v \in (W_{\geq w}^S)''} \text{gch } \mathbb{K}_{s_i v}^-(\lambda) \\ &= \text{gch } \mathbb{K}_{s_i w}^-(\lambda) + \sum_{v \in (W_{\geq w}^S)'', v \neq w} \text{gch } \mathbb{K}_{s_i v}^-(\lambda). \end{aligned}$$

Combining this equality and (A.8), we obtain $\mathsf{T}_i \text{gch } \mathbb{K}_w^-(\lambda) = \text{gch } \mathbb{K}_{s_i w}^-(\lambda)$, as desired.

Case 2. Assume that $\langle w\lambda, \alpha_i^\vee \rangle > 0$; note that $w \in (W_{\geq w}^S)_+$. We see by Lemma 4.12 (2) that if $v \in W_{\geq w}^S$ satisfies $\langle v\lambda, \alpha_i^\vee \rangle < 0$, then $s_i v \in W_{\geq w}^S$. Hence it follows that $(W_{\geq w}^S)'' = \emptyset$, $(W_{\geq w}^S)'_-$ is identical to $(W_{\geq w}^S)_- := \{v \in W_{\geq w}^S \mid \langle v\lambda, \alpha_i^\vee \rangle < 0\}$, and $s_i(W_{\geq w}^S)_- = (W_{\geq w}^S)_+$. By (A.3), we have

$$\mathsf{T}_i \text{gch } V_w^-(\lambda) = \sum_{v \in (W_{\geq w}^S)_0} \mathsf{T}_i \text{gch } \mathbb{K}_v^-(\lambda) + \sum_{v \in (W_{\geq w}^S)_+} \mathsf{T}_i \text{gch } \mathbb{K}_v^-(\lambda) + \sum_{v \in (W_{\geq w}^S)_-} \mathsf{T}_i \text{gch } \mathbb{K}_v^-(\lambda). \quad (\text{A.10})$$

Here we remark that $\mathsf{T}_i \text{gch } V_w^-(\lambda) = 0$ by (A.5). Also, by the induction hypothesis, we have

$$\sum_{v \in (W_{\geq w}^S)_0} \mathsf{T}_i \text{gch } \mathbb{K}_v^-(\lambda) = 0,$$

$$\begin{aligned} \sum_{v \in (W_{\geq w}^S)_+} \mathsf{T}_i \text{gch } \mathbb{K}_v^-(\lambda) &= \mathsf{T}_i \text{gch } \mathbb{K}_w^-(\lambda) + \sum_{v \in (W_{\geq w}^S)_+, v \neq w} \mathsf{T}_i \text{gch } \mathbb{K}_v^-(\lambda) \\ &= \mathsf{T}_i \text{gch } \mathbb{K}_w^-(\lambda) - \sum_{v \in (W_{\geq w}^S)_+, v \neq w} \text{gch } \mathbb{K}_v^-(\lambda), \end{aligned}$$

$$\sum_{v \in (W_{\geq w}^S)_-} \mathsf{T}_i \text{gch } \mathbb{K}_v^-(\lambda) = \sum_{v \in (W_{\geq w}^S)_-} \text{gch } \mathbb{K}_{s_i v}^-(\lambda) = \sum_{v \in (W_{\geq w}^S)_+} \text{gch } \mathbb{K}_v^-(\lambda).$$

Substituting these equalities into (A.10), we obtain $\mathsf{T}_i \text{gch } \mathbb{K}_w^-(\lambda) = -\text{gch } \mathbb{K}_w^-(\lambda)$, as desired.

Case 3. Assume that $\langle w\lambda, \alpha_i^\vee \rangle = 0$; note that $w \in (W_{\geq w}^S)_0$. By the same argument as in Case 2, we see that $(W_{\geq w}^S)''_- = \emptyset$, $(W_{\geq w}^S)'_- = (W_{\geq w}^S)_-$, and $s_i(W_{\geq w}^S)_- = (W_{\geq w}^S)_+$; hence the equality (A.10) holds also in this case. Here we remark that $\mathbb{T}_i \text{gch } V_w^-(\lambda) = 0$ by (A.5), and that

$$\begin{aligned} \sum_{v \in (W_{\geq w}^S)_0} \mathbb{T}_i \text{gch } \mathbb{K}_v^-(\lambda) &= \mathbb{T}_i \text{gch } \mathbb{K}_w^-(\lambda), \\ \sum_{v \in (W_{\geq w}^S)_+} \mathbb{T}_i \text{gch } \mathbb{K}_v^-(\lambda) &= - \sum_{v \in (W_{\geq w}^S)_+} \text{gch } \mathbb{K}_v^-(\lambda), \\ \sum_{v \in (W_{\geq w}^S)_-} \mathbb{T}_i \text{gch } \mathbb{K}_v^-(\lambda) &= \sum_{v \in (W_{\geq w}^S)_+} \text{gch } \mathbb{K}_v^-(\lambda) \end{aligned}$$

by the induction hypothesis. Substituting these equalities into (A.10), we conclude that $\mathbb{T}_i \text{gch } \mathbb{K}_w^-(\lambda) = 0$, as desired.

This completes the proof of Proposition A.4. \square

Recall the following theorem from [CO] and [NNS2, Theorem 35].

Theorem A.5. *Let $w \in W^S$ and $i \in I$ be such that $\langle w\lambda, \alpha_i^\vee \rangle < 0$.*

(1) *If $-[w]^{-1}\alpha_i$ is not a simple root, then*

$$\mathbb{T}_i E_{w\lambda}(q, \infty) = E_{s_i w\lambda}(q, \infty). \quad (\text{A.11})$$

(2) *If $-[w]^{-1}\alpha_i$ is a simple root, then*

$$\mathbb{T}_i E_{w\lambda}(q, \infty) = (1 - q^{\langle \lambda, [w]^{-1}\alpha_i^\vee \rangle}) E_{s_i w\lambda}(q, \infty). \quad (\text{A.12})$$

For $w \in W^S$, let $F_w(q)$ denote the right-hand side of (2.28), that is,

$$F_w(q) := \underbrace{\prod_{i \in I} \prod_{r=1}^{\langle \lambda, \alpha_i^\vee \rangle - \epsilon_i} \frac{1}{1 - q^{-r}}}_{=: c_w(q)} E_{w\lambda}(q, \infty), \quad (\text{A.13})$$

where $\epsilon_i = \epsilon_i([w])$ is as in (2.29).

Lemma A.6. *Let $w \in W^S$ and $i \in I$ be such that $\langle w\lambda, \alpha_i^\vee \rangle < 0$. Then,*

$$\mathbb{T}_i F_w(q) = F_{s_i w}(q). \quad (\text{A.14})$$

Proof. Since $s_i w \in W^S$ (see Lemma 4.9 and Remark 4.10), we have $[s_i w] = s_i w w_\circ(S) = s_i [w]$, where $w_\circ(S) \in W_S$ is the longest element of W_S (see Section 2.2).

Assume first that $-[w]^{-1}\alpha_i$ is not a simple root. Then, for each $k \in I$, we have

$$(\epsilon_k([w]) = 1 \iff) \quad [w]s_k > [w] \iff [s_iw]s_k > [s_iw] \quad (\iff \epsilon_k([s_iw]) = 1). \quad (\text{A.15})$$

Indeed, this is verified as follows. Assume that $[w]s_k > [w]$. Then, $[w]\alpha_k \in \Delta^+$. Also, since $\langle [w]\lambda, \alpha_i^\vee \rangle = \langle w\lambda, \alpha_i^\vee \rangle < 0$, we have $[w]^{-1}\alpha_i \in \Delta^-$. It follows that $[w]\alpha_k \neq \alpha_i$, and hence $s_i[w]\alpha_k \in \Delta^+$, which implies that $[s_iw]s_k = s_i[w]s_k > s_i[w]$. Assume that $[s_iw]s_k > [s_iw]$; we have $s_i[w]\alpha_k = [s_iw]\alpha_k \in \Delta^+$. Suppose, for a contradiction, that $[w]\alpha_k \in \Delta^-$. Then we have $[w]\alpha_k = -\alpha_i$, and hence $-[w]^{-1}\alpha_i = \alpha_k$. However, this contradicts the assumption that $-[w]^{-1}\alpha_i$ is not a simple root. Thus we have shown (A.15). By (A.15), we deduce that $c_w(q) = c_{s_iw}(q)$. Therefore, (A.14) follows from (A.11) in this case.

Assume next that $-[w]^{-1}\alpha_i = \alpha_m$ for some $m \in I$. Then, by exactly the same argument as for (A.15), we have

$$\begin{cases} [w]s_k > [w] \iff [s_iw]s_k > [s_iw] & \text{for all } k \in I \setminus \{m\}, \\ [w]s_m < [w] & \text{and } [s_iw]s_m > [s_iw], \end{cases} \quad (\text{A.16})$$

and hence

$$\frac{1}{1 - q^{-\langle \lambda, \alpha_m^\vee \rangle}} c_{s_iw}(q) = c_w(q); \quad (\text{A.17})$$

notice that $-\langle \lambda, \alpha_m^\vee \rangle = \langle \lambda, [w]^{-1}\alpha_i^\vee \rangle$. Therefore, (A.14) follows from (A.12) in this case. This proves the lemma. \square

Now, we prove the formula (2.28), that is,

$$\text{gch } \mathbb{K}_w^-(\lambda) = F_w(q) \quad \text{for } w \in W^S \quad (\text{A.18})$$

by descending induction on w . Assume first that $w = [w_\circ]$. Then we see by (A.3) that $\text{gch } \mathbb{K}_{[w_\circ]}^-(\lambda) = \text{gch } V_{[w_\circ]}^-(\lambda)$. Since $[w_\circ] = w_\circ$, it follows that $\epsilon_k([w_\circ]) = 0$ for all $k \in I$, and hence

$$F_{[w_\circ]}(q) = \prod_{i \in I} \prod_{r=1}^{\langle \lambda, \alpha_i^\vee \rangle} \frac{1}{1 - q^{-r}} E_{w_\circ\lambda}(q, \infty).$$

Therefore, we deduce that $\text{gch } \mathbb{K}_{[w_\circ]}^-(\lambda) = F_{[w_\circ]}(q)$ by [NNS1, Theorem 5.1.2]. Assume next that $w < [w_\circ]$, and take $i \in I$ such that $\langle w\lambda, \alpha_i^\vee \rangle > 0$; we see by Lemma 4.9 and Remark 4.10 that $s_iw \in W^S$ and $s_iw > w$. Hence, by the induction hypothesis, we obtain $\text{gch } \mathbb{K}_{s_iw}^-(\lambda) = F_{s_iw}(q)$. Here, by Proposition A.4, we have $\text{gch } \mathbb{K}_w^-(\lambda) = \mathbb{T}_i \text{gch } \mathbb{K}_{s_iw}^-(\lambda)$. Also, by Lemma A.6, we have $F_w(q) = \mathbb{T}_i F_{s_iw}(q)$. From these equalities, we conclude that

$$\text{gch } \mathbb{K}_w^-(\lambda) = \mathbb{T}_i \text{gch } \mathbb{K}_{s_iw}^-(\lambda) = \mathbb{T}_i F_{s_iw}(q) = F_w(q),$$

as desired. This completes the proof of the formula (2.28).

Remark A.7. Recall from (A.3) that $\text{gch } V_w^-(\lambda) = \sum_{v \in W_{\geq w}^S} \text{gch } \mathbb{K}_v^-(\lambda)$ for each $w \in W^S$. By the Möbius inversion formula, together with [BB, Corollary 2.7.10], we deduce that

$$\text{gch } \mathbb{K}_w^-(\lambda) = \sum_{v \in W_{\geq w}^S, [w, v] \subset W^S} (-1)^{\ell(v) - \ell(w)} \text{gch } V_v^-(\lambda) \quad \text{for each } w \in W^S,$$

where we set $[w, v] := \{u \in W \mid w \leq u \leq v\}$.

B Relations satisfied by the cyclic vector $u_w \in \mathbb{K}_w^-(\lambda)$.

Let $\lambda \in P^+$, and take $S = S_\lambda$ as in (2.18). For each $w \in W^S$, let $u_w \in \mathbb{K}_w^-(\lambda)$ denote the image of the cyclic vector $v_w \in V_w^-(\lambda)$ under the canonical projection $V_w^-(\lambda) \rightarrow \mathbb{K}_w^-(\lambda)$. Recall from Section 2.1 that for $i \in I_{\text{af}}$, $F_i \in U_{\mathfrak{q}}(\mathfrak{g}_{\text{af}})$ denotes the Chevalley generator corresponding to $-\alpha_i$.

Lemma B.1 (cf. [FKM, §3.1]).

(1) For each $i \in I$, it holds that

$$\begin{cases} F_i^{\langle w\lambda, \alpha_i^\vee \rangle} u_w = 0 & \text{if } \langle w\lambda, \alpha_i^\vee \rangle > 0 \ (\Leftrightarrow w^{-1}\alpha_i \in \Delta^+ \setminus \Delta_S^+), \\ F_i u_w = 0 & \text{if } \langle w\lambda, \alpha_i^\vee \rangle \leq 0 \ (\Leftrightarrow w^{-1}\alpha_i \in \Delta^- \cup \Delta_S). \end{cases}$$

(2) It holds that

$$\begin{cases} F_0^{-\langle w\lambda, \theta^\vee \rangle + 1} u_w = 0 & \text{if } \langle w\lambda, \theta^\vee \rangle < 0 \ (\Leftrightarrow w^{-1}\theta \in \Delta^- \setminus \Delta_S^-), \\ F_0 u_w = 0 & \text{if } \langle w\lambda, \theta^\vee \rangle \geq 0 \ (\Leftrightarrow w^{-1}\theta \in \Delta^+ \cup \Delta_S). \end{cases}$$

(3) It holds that $U_{\mathfrak{q}}^-(\mathfrak{g}_{\text{af}})_{w\alpha + m\delta} u_w = \{0\}$ for all $\alpha \in \Delta^+$ and $m \in \mathbb{Z}$.

Proof. The first equality of part (1) follows from the fact that $F_i^{\langle w\lambda, \alpha_i^\vee \rangle} v_w$ is identical to $v_{s_i w} \in V_{s_i w}^-(\lambda)$; notice that $s_i w \in W^S$ and $s_i w > w$ (see [LNS³1, Lemma 5.9]). The second equality of part (1) and part (2) follow from the fact that v_w is an extremal weight vector of weight $w\lambda$. Part (3) follows from [Kas2, Theorem 5.3]. \square

References

- [BB] A. Björner and F. Brenti, “Combinatorics of Coxeter Groups”, Graduate Texts in Mathematics Vol. 231, Springer, New York, 2005.
- [BFP] F. Brenti, S. Fomin, A. Postnikov, Mixed Bruhat operators and Yang-Baxter equations for Weyl groups, *Int. Math. Res. Not.* **1999** (1999), no. 8, 419–441.
- [BN] J. Beck and H. Nakajima, Crystal bases and two-sided cells of quantum affine algebras, *Duke Math. J.* **123** (2004), 335–402.

- [CO] I. Cherednik and D. Orr, Nonsymmetric difference Whittaker functions, *Math. Z.* **279** (2015), 879–938.
- [FKM] E. Feigin, S. Kato, and I. Makedonskyi, Representation theoretic realization of nonsymmetric Macdonald polynomials at infinity, arXiv:1703.04108.
- [FM] E. Feigin and I. Makedonskyi, Generalized Weyl modules, alcove paths and Macdonald polynomials, *Selecta Math.* (N.S.) **23** (2017), 2863–2897.
- [HK] J. Hong and S.-J. Kang, “Introduction to Quantum Groups and Crystal Bases”, Graduate Studies in Mathematics Vol. 42, Amer. Math. Soc., Providence, RI, 2002.
- [H] J. E. Humphreys, “Reflection Groups and Coxeter Groups”, Cambridge Studies in Advanced Mathematics Vol. 29, Cambridge University Press, Cambridge, 1990.
- [INS] M. Ishii, S. Naito, and D. Sagaki, Semi-infinite Lakshmibai-Seshadri path model for level-zero extremal weight modules over quantum affine algebras, *Adv. Math.* **290** (2016), 967–1009.
- [Kac] V. G. Kac, “Infinite Dimensional Lie Algebras”, 3rd Edition, Cambridge University Press, Cambridge, UK, 1990.
- [Kas1] M. Kashiwara, Crystal bases of modified quantized enveloping algebra, *Duke Math. J.* **73** (1994), 383–413.
- [Kas2] M. Kashiwara, On level-zero representations of quantized affine algebras, *Duke Math. J.* **112** (2002), 117–175.
- [Kas3] M. Kashiwara, Level zero fundamental representations over quantized affine algebras and Demazure modules, *Publ. Res. Inst. Math. Sci.* **41** (2005), 223–250.
- [Kat] S. Kato, Demazure character formula for semi-infinite flag varieties, *Math. Ann.* **371** (2018), 1769–1801.
- [LLM] V. Lakshmibai, P. Littelmann, and P. Magyar, Standard monomial theory for Bott-Samelson varieties, *Compositio Math.* **130** (2002), 293–318.
- [LS] T. Lam and M. Shimozono, Quantum cohomology of G/P and homology of affine Grassmannian, *Acta Math.* **204** (2010), 49–90.
- [LNS³1] C. Lenart, S. Naito, D. Sagaki, A. Schilling, and M. Shimozono, A uniform model for Kirillov-Reshetikhin crystals I: Lifting the parabolic quantum Bruhat graph, *Int. Math. Res. Not.* **2015** (2015), 1848–1901.
- [LNS³2] C. Lenart, S. Naito, D. Sagaki, A. Schilling, and M. Shimozono, A uniform model for Kirillov-Reshetikhin crystals II: Alcove model, path model, and $P = X$, *Int. Math. Res. Not.* **2017** (2017), 4259–4319.
- [LNS³3] C. Lenart, S. Naito, D. Sagaki, A. Schilling, and M. Shimozono, Quantum Lakshmibai-Seshadri paths and root operators, in “Schubert Calculus – Osaka 2012”, pp.267–294, Adv. Stud. Pure Math. Vol. 71, Math. Soc. Japan, Tokyo, 2016.
- [LNS³4] C. Lenart, S. Naito, D. Sagaki, A. Schilling, and M. Shimozono, A uniform model for Kirillov-Reshetikhin crystals III: Nonsymmetric Macdonald polynomials at $t = 0$ and Demazure characters, *Transform. Groups* **22** (2017), 1041–1079.
- [Li] P. Littelmann, Paths and root operators in representation theory, *Ann. of Math.* (2) **142** (1995), 499–525.
- [Lu1] G. Lusztig, Hecke algebras and Jantzen’s generic decomposition patterns, *Adv. Math.* **37** (1980), 121–164.

- [Lu2] G. Lusztig, Periodic W -graphs, *Represent. Theory* **1** (1997), 207–279.
- [NNS1] S. Naito, F. Nomoto, and D. Sagaki, Specialization of nonsymmetric Macdonald polynomials at $t = \infty$ and Demazure submodules of level-zero extremal weight modules, *Trans. Amer. Math. Soc.* **370** (2018), 2739–2783.
- [NNS2] S. Naito, F. Nomoto, and D. Sagaki, Representation-theoretic interpretation of Cherednik-Orr’s recursion formula for the specialization of nonsymmetric Macdonald polynomials at $t = \infty$, to appear in *Transform. Groups*, DOI:10.1007/s00031-017-9468-0.
- [NOS] S. Naito, D. Orr, and D. Sagaki, Pieri-Chevalley formula for anti-dominant weights in the equivariant K -theory of semi-infinite flag manifolds, preprint 2018, arXiv:1808.01468.
- [NS] S. Naito and D. Sagaki, Demazure submodules of level-zero extremal weight modules and specializations of Macdonald polynomials, *Math. Z.* **283** (2016), 937–978.
- [P] D. Peterson, Quantum Cohomology of G/P , Lecture notes, Massachusetts Institute of Technology, Spring 1997.
- [S] W. Soergel, Kazhdan-Lusztig polynomials and a combinatoric for tilting modules, *Represent. Theory* **1** (1997), 83–114.