

Left-orderability for surgeries on $(-2, 3, 2s + 1)$ -pretzel knots

Zipei Nie

December 10, 2021

Abstract

In this paper, we prove that the fundamental group of the manifold obtained by Dehn surgery along a $(-2, 3, 2s + 1)$ -pretzel knot ($s \geq 3$) with slope $\frac{p}{q}$ is not left orderable if $\frac{p}{q} \geq 2s + 3$, and that it is left orderable if $\frac{p}{q}$ is in a neighborhood of zero depending on s .

1 Introduction

For a nontrivial group G , we say it is left-orderable if it admits a total ordering that is invariant under left multiplication, that is, $g < h$ implies $fg < fh$ for all elements $f, g, h \in G$. By convention, the trivial group is not left-orderable.

For any rational homology 3-sphere Y , the Euler characteristic of the Heegaard Floer homology $\widehat{HF}(Y)$ is $|H_1(Y; \mathbb{Z})|$. We say a closed 3-manifold Y is an L-space if it is a rational homology 3-sphere and the rank of $\widehat{HF}(Y)$ is $|H_1(Y; \mathbb{Z})|$.

The following conjecture proposed by Boyer, Gordon, and Watson [2] suggests a close relation between the existence of a left order and the L-space.

Conjecture 1. *An irreducible rational homology 3-sphere is an L-space if and only if its fundamental group is not left-orderable.*

In [2], Conjecture 1 is verified for geometric, non-hyperbolic 3-manifolds and the 2-fold branched covers of non-split alternating links.

The surgery formula of the Heegaard Floer homology in [14] gives rise to the following relation between the L-space and Dehn surgery. It is a way to produce many hyperbolic L-spaces.

Theorem 1. *For any knot K , if a positive surgery of K yields an L-space, then the $\frac{p}{q}$ -surgery of K is an L-space if and only if $\frac{p}{q} \geq 2g(K) - 1$, where $g(K)$ is the genus of K .*

According to [1], the $(-2, 3, 2s + 1)$ -pretzel knots for integers $s \geq 3$ are the only hyperbolic knots with L-space surgeries up to mirroring among all Montesinos knots. It is no wonder these knots draw attention in the literature. We state Conjecture 1 restricted on these knots as follows.

Conjecture 2. *The fundamental group of the manifold obtained by Dehn surgery along a $(-2, 3, 2s + 1)$ -pretzel knot ($s \geq 3$) with slope $\frac{p}{q}$ is left orderable if and only if $\frac{p}{q} < 2s + 3$.*

In 2004, Jinha Jun [11] proved that, for $s = 3$ and the slope $\frac{p}{q} \geq 10$ and p is odd, then the manifold obtained by Dehn surgery does not contain an \mathbb{R} -covered foliation. In 2013, Yasuharu Nakae [12] extends Jun's result to the case where $\frac{p}{q} \geq 4s + 7$ and p is odd. In the viewpoint of left-orderable group, they essentially proved that the fundamental group is not left-orderable for $s = 3$ and $\frac{p}{q} \geq 10$, and for $\frac{p}{q} \geq 4s + 7$, respectively.

With a different strategy, Adam Clay and Liam Watson [5] in 2012 proved that the fundamental group is not left-orderable for $\frac{p}{q} \geq 2s + 11$.

In Section 2, following the strategy set up by Jun and Nakae, we improve their results by proving the following theorem.

Theorem 2. *If $\frac{p}{q} \geq 2s + 3$, then the fundamental group of the manifold obtained by Dehn surgery along a $(-2, 3, 2s + 1)$ -pretzel knot ($s \geq 3$) with slope $\frac{p}{q}$ is not left orderable.*

Marc Culler and Nathan Dunfield [6] defined the knot complement of K to be lean, if every closed essential surface in the manifold $S_0^3(K)$ obtained by longitudinal Dehn surgery is a fiber in a fibration over circle. In the general case where S^3 is replaced by another integral homology sphere, we need an extra condition that the resulting manifold is prime. We do not need to bother about it here, because Gabai [8] showed that $S_0^3(K)$ is irreducible for any nontrivial knot K . In particular, $S_0^3(K)$ is prime for any knot K .

Culler and Dunfield proved the following result by considering the representations $\pi_1(S^3 - K) \rightarrow \widetilde{\text{PSL}}_2\mathbf{R}$ that are elliptic on the fundamental group of the torus boundary.

Theorem 3. *For an knot K in S^3 with lean knot complement, if its Alexander polynomial $\Delta_K(t)$ has a simple root on the unit circle, then there exists $\varepsilon > 0$ such that the fundamental group of the manifold obtained by Dehn surgery along K with any slope $\frac{p}{q}$ in $(-\varepsilon, \varepsilon)$ is left orderable.*

As an example demonstrated in their work, they proved that the fundamental group of the manifold obtained by Dehn surgery along a $(-2, 3, 7)$ -pretzel knot with slope $\frac{p}{q}$ is left orderable if $\frac{p}{q} < 6$.

In Section 3, we prove the following result by verifying the conditions of Theorem 3 for all $(-2, 3, 2s + 1)$ -pretzel knots ($s \geq 3$).

Theorem 4. *For each integer $s \geq 3$, there exists $\varepsilon > 0$ such that the fundamental group of the manifold obtained by Dehn surgery along a $(-2, 3, 2s + 1)$ -pretzel knot with any slope $\frac{p}{q}$ in $(-\varepsilon, \varepsilon)$ is left orderable.*

Acknowledgement

I would like to thank my advisor, Professor Zoltán Szabó, for suggesting me this problem and helping me throughout my research project.

2 The Proof of Theorem 2

Let K_s be a $(-2, 3, 2s + 1)$ -pretzel knot with $s \geq 3$. It is shown [12, Proposition 2.1] that the knot group of K_s has following presentation.

Theorem 5. *The knot group of K_s has a presentation*

$$G_{K_s} = \langle c, l | clcl\bar{c}l^s\bar{c}lcl^{s-1} \rangle,$$

and an element which represents the meridian M is c and an element of the longitude L is $\bar{c}^{2s-2}lcl^scl^scl\bar{c}^{2s+9}$.

For coprime positive integers p and q , let $G_{K_s}(p, q)$ denote the fundamental group of the manifold obtained by Dehn surgery along K_s with slope $\frac{p}{q}$. Then we have the following presentation of $G_{K_s}(p, q)$:

$$G_{K_s}(p, q) = \langle c, l | clcl\bar{c}l^s\bar{c}lcl^{s-1}, M^pL^q \rangle.$$

We quote a lemma [12, Lemma 3.1] below. The element k in the statement is a generator of the cyclic group generated by M and L .

Lemma 1. *There exists an element k of $G_{K_s}(p, q)$ such that $M = k^q$ and $L = k^{-p}$.*

By the result from the standard exercise [4, Problem 2.25] in Adam Clay and Liam Watson's book, for a countable group G with a left ordering $<$, the dynamic realization of the left ordering has no global fixed points. In other words, we have the following theorem.

Theorem 6. *For a countable group G with a left ordering $<$, there exists a monomorphism $\rho : G \rightarrow \text{Homeo}_+(\mathbf{R})$. Additionally, there is no $x \in \mathbf{R}$ such that $\rho(g)(x) = x$ for all g in G .*

Now we assume $G_{K_s}(p, q)$ has a left ordering and fix such a monomorphism as in Theorem 6 for the group $G_{K_s}(p, q)$. For abbreviation, we write gx instead of $\rho(g)(x)$ for $g \in G_{K_s}(p, q)$ and $x \in \mathbf{R}$.

First, We prove that $\rho(k)$ has no fixed points.

According to [9], a Seifert surface can be found by taking a candidate surface of connected boundary associated to a certain type III admissible system of monochromatic edgepaths. For the knot K_s , the edgepaths in this system always move upwards before heading to the common ending point $\langle \infty \rangle$.

A non-constant edgepath in any admissible edgepath system for the knot K_s either always move upwards or always move downwards. Therefore, the slope of the boundary curves of a candidate surface associated to a type II or type III admissible edgepath system for the knot K_s has at most $2^3 = 8$ possible values, which can be computed similarly to [9, p.462] and [18, Table 2]. The possible slopes are listed in the table below. The only zero entry corresponds to the unique edgepath system that produces Seifert surfaces.

Slope List

Directions of edgepaths	Type II	Type III
+++	Not admissible	0
++-	$4s + 4$	$4s + 2$
+ - +	8	6
+ - -	$4s + 8$	$4s + 8$
- + +	6	4
- + -	$4s + 6$	$4s + 6$
- - +	10	10
- - -	$4s + 10$	$4s + 12$

By the proof of [17, Lemma 4.4], the slope of the boundary curves of a candidate surface associated to a type I admissible edgepath system for the knot K_s is always positive.

Therefore, any incompressible, ∂ -incompressible connected orientable surface in the knot complement of K_s with 0-slope boundary curves are isomorphic to a Seifert surface. Since K_s is fibered [7] and any Seifert surface of a fibered knot is isotopic to a given fiber [3], it follows that K_s is lean.

3.2 Roots of the Alexander polynomial of K_s

By [10, Theorem 1.2], for an odd number k and positive integers p_1, \dots, p_k , the Alexander polynomial of a $(p_1, \dots, p_k, -1)$ -pretzel link can be written as $Q_{p_1, \dots, p_k}(-x)$, where we define

$$Q_{p_1, \dots, p_k}(x) = [p_1][p_2] \cdots [p_k] \left(x - k + 1 + \sum_{i=1}^k \frac{1}{[p_i]} \right),$$

and $[n] = \frac{1-x^n}{1-x}$. In the case where $p_1 = 2$, the $(2, p_2, \dots, p_k, -1)$ -pretzel link is equivalent to the $(-2, p_2, \dots, p_k)$ -pretzel link by a single flipping operation. By [15, Theorem 2.6], the polynomial $Q_{p_1, \dots, p_k}(x)$ is also the denominator of the growth function of the (p_1, \dots, p_k) -Coxeter reflection group $\langle g_1, \dots, g_k | g_i^2, (g_i g_{i+1})^{p_i} \rangle$, and if the Coxeter system is hyperbolic, then the polynomial $Q_{p_1, \dots, p_k}(x)$ is a product of distinct irreducible cyclotomic polynomials and exactly one Salem polynomial. Hence, if

$$\sum_{i=1}^k \frac{1}{p_i} < k - 2,$$

then all roots of $Q_{p_1, \dots, p_k}(x)$ are simple, with exactly two of them real, and the rest on the unit circle. Therefore, the Alexander polynomial of K_s has $2s + 2$ simple roots on the unit circle.

In conclusion, by Theorem 3, we proved Theorem 4.

References

- [1] K. L. Baker and A. H. Moore. "Montesinos knots, Hopf plumbings, and L-space surgeries." J. Math. Soc. Japan. Vol. 70. No. 1. (2018): 95-110.
- [2] S. Boyer, C. Gordon and L. Watson. "On L-spaces and left-orderable fundamental groups." Mathematische Annalen 356.4 (2013): 1213-1245.

- [3] G. Burde and H. Zieschang. "Neuwirthsche knoten und flächenabbildungen." Abhandlungen aus dem Mathematischen Seminar der Universität Hamburg. Vol. 31. No. 3-4. Springer-Verlag, 1967.
- [4] A. Clay and D. Rolfsen. Ordered groups and topology. Vol. 176. American Mathematical Society, 2016.
- [5] A. Clay and L. Watson. "Left-orderable fundamental groups and Dehn surgery." International Mathematics Research Notices 2013.12 (2012): 2862-2890.
- [6] M. Culler and N. Dunfield. Orderability and Dehn filling, arXiv: 1602.03793.
- [7] D. Gabai. "Detecting fibred links in S^3 ." Commentarii Mathematici Helvetici 61.1 (1986): 519-555.
- [8] D. Gabai. "Foliations and the topology of 3-manifolds. III." Journal of Differential Geometry 26.3 (1987): 479-536.
- [9] A. Hatcher and U. Oertel. "Boundary slopes for Montesinos knots." Topology 28.4 (1989): 453-480.
- [10] E. Hironaka. "The Lehmer polynomial and pretzel links." Canadian Mathematical Bulletin 44.4 (2001): 440-451.
- [11] J. Jun. " $(-2, 3, 7)$ -pretzel knot and Reebless foliation." Topology and its Applications 145.1-3 (2004): 209-232.
- [12] Y. Nakae. "A Good Presentation of $(-2, 3, 2s + 1)$ -type Pretzel Knot Group and R-covered Foliation." Journal of Knot Theory and Its Ramifications 22.01 (2013): 1250143
- [13] U. Oertel. 'Closed incompressible surfaces in complements of star links,' Pac. J. Math. 111 (1984) 209-230.
- [14] P. S. Ozsváth and Z. Szabó. "Knot Floer homology and rational surgeries." Algebraic & Geometric Topology 11.1 (2011): 1-68.
- [15] W. Parry. "Growth series of Coxeter groups and Salem numbers." J. Algebra 154.2 (1993): 406-415.
- [16] W. P. Thurston. The geometry and topology of three-manifolds. Princeton, NJ: Princeton University, 1979.
- [17] J. Rodríguez-Viorato. "Alternating Montesinos knots and Conjecture \mathbb{Z} ." arXiv preprint arXiv:1606.07033 (2016).
- [18] J. Rodríguez-Viorato and F. J. González-Acuña. "On pretzel knots and Conjecture \mathbb{Z} ." Journal of Knot Theory and Its Ramifications 25.02 (2016): 1650012.