

Geometric Random Graphs and Rado Sets of Continuous functions

Anthony Bonato* Jeannette Janssen† Anthony Quas‡

Received 2 March 2018; Revised 25 May 2020; Published 7 April 2021

Abstract: We prove the existence of geometric Rado sets in the Banach space of continuous functions on $[0, 1]$. Starting from a countable dense set S , we form a random graph by independently joining pairs of elements of S that are less than 1 apart by an edge with probability $p \in (0, 1)$. If any two realizations of this random graph are almost surely isomorphic, then the set S is said to be *geometrically Rado*.

We show that for suitable measures which we construct, almost all countable dense sets of piecewise linear functions and of polynomials are geometrically Rado. The same holds for countably many independent samples of integrated Brownian motion. Moreover the almost sure random graph obtained in the three cases is the same.

We also study Brownian motion paths. Again, we show for a natural measure on Brownian paths, almost all countable subsets are geometrically Rado and the resulting graphs are almost surely of a unique isomorphism type. However, the almost sure Rado graph in the Brownian motion case is not isomorphic to the Rado graph in the previous cases.

Key words and phrases: geometric random graph, Rado graph

1 Overview

We consider *geometric graphs*, where the vertex set is a dense subset of a separable metric space, and the edge set has the property that there exists $a > 0$ such that if the vertices x and y are joined by an edge, then $0 < d(x, y) < a$. In this paper the metric space will be a separable Banach space, X , and a will be 1.

*Supported by NSERC

†Supported by NSERC

‡Supported by NSERC

If V is a countable dense subset of X , and $p \in (0, 1)$ is a parameter, then we define a collection of random geometric graphs with vertex set V by independently joining each pair x and y in V by an edge with probability p if $0 < \|x - y\| < 1$. We call this distribution the *local area random graph* on V with parameter p and denote it by $\text{LARG}(V, p)$.

Our main interest in this paper is in questions of isomorphism between random geometric graphs. First, we ask, for a fixed countable dense vertex set V , whether the graphs in $\text{LARG}(V, p)$ are almost surely isomorphic. If they are, then we say that the set V has the *geometric Rado property*, by analogy with the discovery that random subgraphs on a countable vertex set where all pairs of vertices are independently joined with probability p are almost surely isomorphic to a universal graph.

Second, if a Banach space has countable dense subsets with the Rado property, we ask whether there is a natural measure μ on X such that for $\mu^{\mathbb{N}}$ -almost every countable dense V (simply the set of elements appearing in the sequence in $X^{\mathbb{N}}$), the elements of $\text{LARG}(V, p)$ are almost all isomorphic. In this case, we also ask whether for two independently chosen countable dense subsets V and V' and for all $p, p' \in (0, 1)$, it is the case that almost every G in $\text{LARG}(V, p)$ and $G' \in \text{LARG}(V', p')$ are isomorphic. If it is, then we say that $\mu^{\mathbb{N}}$ is *universally geometrically Rado*. If X supports natural measures generating universally geometrically Rado vertex sets, then it is natural to ask if different measures lead to different universal graphs.

In [3] where these notions were introduced, Bonato and Janssen showed the surprising result that if a finite-dimensional ℓ^∞ space is equipped with an absolutely continuous probability measure μ with full support (that is, every open set has positive measure), then $\mu^{\mathbb{N}}$ is universally geometrically Rado, and also the isomorphism class does not depend on the measure μ . The fact that μ is absolutely continuous guarantees that, almost surely, all of the points sampled are *integer distance free*, that is that no two distinct points have a coordinate in which the points differ by an integer. In fact, their proof shows that if V and W are dense countable integer distance free subsets of ℓ^∞ , then elements of $\text{LARG}(V, p)$ and $\text{LARG}(W, p')$ are almost surely isomorphic. They also show that in \mathbb{R}^2 equipped with the Euclidean norm, countable dense sets never have the Rado property.

In the elegant paper [2], Balister, Bollobás, Gunderson, Leader, and Walters give a necessary and sufficient criterion for the existence of Rado sets in finite-dimensional Banach spaces. They uniquely express X as an “ ∞ -decomposition” ($U \oplus \ell_\infty^d$) into a maximal ℓ^∞ part and the rest. They then show that the space admits geometric Rado sets if and only if $d > 0$, and it has the universal geometric Rado property if and only if $U = \{0\}$ (that is, if and only if X is isometrically isomorphic to ℓ_∞^d for some d). They also show that there is a countable dense subset V of ℓ^∞ that is not integer distance free which does not have the geometric Rado property, thereby emphasizing the role of the integer distance free property.

In [4], we studied the existence of Rado sets in the separable infinite-dimensional subspaces of ℓ^∞ : c_0 (of sequences converging to 0) and c (of convergent sequences). We exhibited fully supported measures on c_0 and c with the universal geometric Rado property and showed that in both cases, the universal graph obtained was the same for all measures in a reasonably general class (but the universal graphs for c_0 and c are not isomorphic; see Corollary 2). In [1], the authors considered Rado sets in metric spaces on circles of a fixed circumference.

In this paper, we study the Banach space $C[0, 1]$ of continuous real-valued functions on $[0, 1]$, a separable subspace of $L^\infty[0, 1]$. We exhibit probability measures μ on $C[0, 1]$ supported on polynomials, on piecewise linear functions, and on integrated Brownian motions, such that $\mu^{\mathbb{N}}$ is universally geometrically

Rado. Further, the universal random graph is the same for these classes. Additionally, we exhibit a probability measure ν on $C[0, 1]$ supported on Brownian motion functions such that $\nu^{\mathbb{N}}$ is universally geometrically Rado. However, it turns out that the universal random graph for the Brownian motion functions is not isomorphic to the universal random graph for the polynomials and piecewise linear functions. Informally, the universal random graphs “see the intersections between pairs of functions”. The fact that pairs of Brownian motions cross infinitely often if they cross at all leads to non-isomorphism of the graphs.

The current paper revolves around two themes:

1. showing that the ℓ^∞ -like space $C[0, 1]$ supports natural measures on countable dense subsets that exhibit the universal geometric Rado property;
2. understanding the geometric information about countable dense subsets that may be obtained from the isomorphism type of the local area random graphs that they support (in particular exhibiting two natural classes of distributions on $C[0, 1]$ with the universal geometric Rado property, and showing that their universal Rado graphs fail to be isomorphic).

The *crossings* between elements functions f and g (that is, points $x \in [0, 1]$ such that $f(x) - g(x) \in \mathbb{Z}$) play a key role in all of this that is similar to the role played by the *integer distance free* property in [3].

2 The Banach space and random graph geometries

A geometric graph G is said to be *well-connected* if it has the property that for every finite subset S of $V(G)$ and for every non-empty open set $U \subseteq X$ such that $\|u - x\| < 1$ for all $u \in U$ and $x \in S$, there exists $y \in U \cap V(G)$ such that y is adjacent to each $x \in S$. By a Borel-Cantelli argument, it is straightforward to show that if V is a countable dense subset of X , then almost every G sampled from $\text{LARG}(V, p)$ is well-connected.

Lemma 1. *If G and G' are well-connected geometric graphs in X and X' and $\theta: G \rightarrow G'$ is a graph isomorphism, then we have the following:*

1. $\lfloor \|x - y\| \rfloor = \lfloor \|\theta(x) - \theta(y)\| \rfloor$ for all $x, y \in V(G)$ satisfying $\|x - y\| \geq 2$; and
2. $\lceil \|x - y\| \rceil = \lceil \|\theta(x) - \theta(y)\| \rceil$ for all $x, y \in V(G)$.

In particular, if $\|x - y\|$ is an integer and is at least 2, then $\|\theta(x) - \theta(y)\| = \|x - y\|$.

For the time being, in the infinite-dimensional case, we are unable to rule out the possibility that a pair of vertices separated by a distance of 1 are mapped to vertices separated by a distance less than 1 that are not connected by an edge. However, this can be ruled out in the finite-dimensional case; see Corollary 4 below.

The following corollary of Lemma 1 was established in [4]. The corollary was established using a theorem of Dilworth [5] showing that approximately surjective approximate isometries may be uniformly approximated by surjective isometries, and the theorem of Mazur and Ulam stating that surjective isometries of real Banach spaces may be expressed as the composition of a translation and a linear isometry.

Corollary 2. *If G and G' are isomorphic well-connected geometric graphs in X and X' , then the Banach spaces X and X' are isometrically isomorphic.*

The corollary implies that a well-connected geometric graph with vertices in a separable Banach space determines the Banach space structure.

A map between two sets V and V' is called a *step-isometry* if $\lfloor \|\theta(x) - \theta(y)\| \rfloor = \lfloor \|x - y\| \rfloor$ for all $x, y \in V$. As shown in Lemma 1, a graph isomorphism between two well-connected graphs is almost a step-isometry (modulo the possibility mentioned above). We use this in the context of $C[0, 1]$ to build graph isomorphisms: given graphs G and G' sampled from $\text{LARG}(V, p)$ and $\text{LARG}(V', p')$, we build a graph isomorphism using a back-and-forth technique, ensuring that at each stage of the construction (at which time only finitely many vertices are matched), the part of the isomorphism so far constructed is a step-isometry. As long as all of the vertices in both graphs are exhausted by this process, this yields an isomorphism between G and G' .

The above shows (as first observed in [BJ1]), there is an intimate connection between step-isometries and geometric graph isomorphisms. This is elaborated upon in [2]; in particular, it is shown that if a finite-dimensional space X has ∞ -decomposition $(U \oplus \ell_\infty^d)_\infty$, then a step-isometry may be factorized over the decomposition as a surjective linear isometry (on U) and a step-isometry on the ℓ_∞ factor. Establishing the Rado property depends on the availability of a plentiful supply of step-isometries and this gives rise to the failure of the Rado property on spaces that are not isomorphic to ℓ_∞^d .

Proof of Lemma 1. It was shown in [3] that if x, y are elements of $V(G)$ for any integer $k \geq 2$, $\|x - y\| < k$ implies $d_G(x, y) \leq k$. To see this, consider the $k - 1$ equally spaced points $x + \frac{i}{k}(y - x)$ for $i = 1, \dots, k - 1$ along the chord $[x, y]$. Since $\frac{\|y - x\|}{k} < 1$, there are open balls around these points with the property that each point in the i th ball is less than one unit distant from each point in the next ball and the first and last balls are within 1 unit of x and y , respectively. For $i = 1$, by well-connectedness there exists a vertex x_1 in the first ball adjacent to x . Continuing inductively one may select x_1, \dots, x_{k-2} such that each is adjacent to the next. Finally, x_{k-1} is chosen to be adjacent to both x_{k-2} and y . On the other hand, it is straightforward that for any $k \in \mathbb{N}$, $d_G(x, y) \leq k$ implies $\|x - y\| < k$. It follows that for any $k \geq 3$, $d_G(x, y) = k$ if and only if $\|x - y\| \in [k - 1, k)$. In particular, if $\|x - y\| \geq 2$, then $\lfloor \|x - y\| \rfloor = d_G(x, y) - 1 = d_{G'}(\theta(x), \theta(y)) - 1 = \lfloor \|\theta(x) - \theta(y)\| \rfloor$.

For the second part, suppose $\|x - y\| > k \in \mathbb{N}_0$. By well-connectedness, there exist w adjacent to x and z adjacent to y such that $\|w - z\| > k + 2$ (for example, let $u = (y - x)/\|y - x\|$ be the direction joining x to y and choose w in a small ball around $x - u$ and z in a small ball around $y + u$). Hence, we have that $d_G(w, z) \geq k + 3$, so that $d_{G'}(\theta(w), \theta(z)) \geq k + 3$ and by the previous part, $\|\theta(w) - \theta(z)\| \geq k + 2$. Now, it follows that $\|\theta(x) - \theta(y)\| \geq \|\theta(w) - \theta(z)\| - \|\theta(w) - \theta(x)\| - \|\theta(z) - \theta(y)\| > k$. \square

Recall that a sequence (y_n) in a Banach space X is said to *converge weakly* to y if $\phi(y_n) \rightarrow \phi(y)$ for all ϕ in X^* , the dual space of X .

Lemma 3. *Let G and G' be well-connected graphs on the Banach space X and suppose that they are isomorphic by the map θ . If (x_n) is a sequence of elements of $V(G)$ such that $x_n \rightarrow x \in V(G)$, then $\theta(x_n)$ converges weakly to $\theta(x)$ in $V(G')$.*

In particular, if X is finite-dimensional, θ is continuous.

Proof. We prove this by contradiction. Let $y = \theta(x)$ and $y_n = \theta(x_n)$. Suppose that ϕ is a linear functional of norm 1 so that $\phi(y_n) \not\rightarrow \phi(y)$. Since $x_n \rightarrow x$, all but finitely many x_n 's lie in the closed ball of radius 1 about x , so that by the second statement of Lemma 1, for all except these finitely many n , y_n lies in the closed ball of radius 1 about y . Hence, we may extract a subsequence such that $\phi(y_n) \rightarrow \phi(y) - a$, where we may assume, by negating ϕ if necessary, that $a \in (0, 1)$. Since $\|\phi\| = 1$ and $V(G')$ is dense, there exists $z = \theta(w) \in V(G')$ such that $\|z - y\| < 2$; $\phi(z) - \phi(y) > 2 - \frac{a}{2}$. For all sufficiently large n , we have that $\phi(z) - \phi(y_n) > 2$, so that $\|z - y_n\| > 2$. By the second statement of Lemma 1, $\|w - x_n\| > 2$, so that $\|w - x\| \geq 2$. We then have that $d_G(w, x) \geq 3$. However, $\|z - y\| < 2$ implies that $d_{G'}(z, y) = 2$ by the first paragraph of the proof of Lemma 1, which contradicts the assumption that G and G' are isomorphic.

If X is finite-dimensional, then weak convergence is equivalent to convergence in norm. □

Corollary 4. *If X is a finite-dimensional Banach space and G and G' are two well-connected geometric graphs on X , that are isomorphic by a map θ , then $\|\theta(x) - \theta(y)\| = 1$ if and only if $\|x - y\| = 1$.*

Hence, in the finite-dimensional case, for any $k \in \mathbb{N}_0$, $\|x - y\| = k$ if and only if $\|\theta(x) - \theta(y)\| = k$; $k < \|x - y\| < k + 1$ if and only if $k < \|\theta(x) - \theta(y)\| < k + 1$.

Proof. Let $x, y \in V(G)$ satisfy $\|x - y\| = 1$. By Lemma 1, we know that $\|\theta(x) - \theta(y)\| \in (0, 1]$. Let $x_n \rightarrow x$ and $y_n \rightarrow y$ with $x_n, y_n \in V(G)$ and $\|y_n - x_n\| > 1$ (the existence of such sequences is guaranteed by density of $V(G)$): let $u = y - x$ be the unit vector joining x and y ; let x_n be a vertex in a ball of radius $\frac{1}{2n}$ around $x - \frac{u}{n}$ and let y_n be a vertex in a ball of radius $\frac{1}{2n}$ around $y + \frac{u}{n}$. We then have that $\|\theta(y_n) - \theta(x_n)\| > 1$ by the second statement of Lemma 1 and $\theta(x_n) \rightarrow \theta(x)$ and $\theta(y_n) \rightarrow \theta(y)$ by Lemma 3, so that $\|\theta(y) - \theta(x)\| \geq 1$, completing the proof of the first part. The second part follows by combining this with Lemma 1. □

We know from Lemma 1 that the isometries of well-connected geometric graphs give rise to step-isometries of the vertex sets. It is therefore useful to have a large collection of step-isometries of $C[0, 1]$. The following lemma gives a class of step-isometries that underlie the inductive proof that we give of the Rado properties.

Informally, we can think of the step-isometries that we build as acting on graphs of functions on $[0, 1]$ as follows. We first apply a homeomorphism to the x coordinates. We then apply a step-isometry (depending continuously on x) to the graphs of the functions over the x coordinate. This should be compared with the Banach-Stone theorem; see Theorem 15.

Lemma 5. *Let $\Phi: [0, 1] \times \mathbb{R} \rightarrow \mathbb{R}$ be a continuous function such that $\Phi(y, t)$ is a strictly increasing function of t satisfying $\Phi(y, t + 1) = \Phi(y, t) + 1$ for all $y \in [0, 1]$, $t \in \mathbb{R}$. If $\psi: [0, 1] \rightarrow [0, 1]$ is an increasing continuous bijection, then the map $\Theta: C[0, 1] \rightarrow C[0, 1]$ defined by $\Theta[f](y) = \Phi(y, f(\psi(y)))$ is a bijective continuous step-isometry of $C[0, 1]$.*

Proof. We first establish that Θ is a homeomorphism of $C[0, 1]$, and that its inverse is of the same form. Note that the map $f \mapsto f \circ \psi$ is a bijective linear isometry of $C[0, 1]$ and that Θ may be expressed as a composition of $g \mapsto \Phi(y, g(y))$ with this bijection. Hence, it suffices to consider the case where $\psi(y) = y$ and $\Theta[f](y) = \Phi(y, f(y))$. Let ϕ_y be defined by $\phi_y(t) = \Phi(y, t)$. By hypothesis, ϕ_y is a bijection of \mathbb{R} .

The inverse of Θ is given by $\Theta^{-1}(g)(y) = \phi_y^{-1}(g(y))$. We claim that Θ^{-1} is continuous. To see this, let $\varepsilon > 0$ and let $\delta = \min_{y \in [0, 1]} \min_{t \in \mathbb{R}} [\phi_y(t + \varepsilon) - \phi_y(t)]$ (this quantity is positive by the fact that ϕ_y is

periodic-plus-identity and by compactness of $[0,1]$). Now for any $y \in [0, 1]$ and $a \in \mathbb{R}$, $\phi_y(\phi_y^{-1}a - \varepsilon) \leq \phi_y(\phi_y^{-1}a) - \delta = a - \delta$, so that $\phi_y^{-1}(a - \delta) \geq \phi_y^{-1}a - \varepsilon$ and similarly $\phi_y^{-1}(a + \delta) \leq \phi_y^{-1}a + \varepsilon$. In particular, if $|a - b| < \delta$, then $|\phi_y^{-1}b - \phi_y^{-1}a| < \varepsilon$. Now if $\|f - g\| < \delta$, we see that $\|\Theta^{-1}f - \Theta^{-1}g\| < \varepsilon$. (We have shown in this paragraph that the inversion map from the set of bijective homeomorphisms of the line such that $\phi(t + 1) = \phi(t) + 1$ to itself is continuous.)

If $\lfloor \|f - g\| \rfloor = k$, then we have that $|f(x) - g(x)| < k + 1$ for each x . We note that $\Theta[f](y) - \Theta[g](y) = \Phi(y, f(y)) - \Phi(y, g(y))$. Since $-(k + 1) < f(y) - g(y) < k + 1$, it follows that $-(k + 1) < \Phi(y, f(y)) - \Phi(y, g(y)) < k + 1$ for each x , so that $\|\Theta(f) - \Theta(g)\| \leq k = \lfloor \|f - g\| \rfloor$. Applying this argument to Θ^{-1} (which we have already shown to be of the same form), we deduce that $\|\Theta(f) - \Theta(g)\| = \lfloor \|f - g\| \rfloor$, as required. \square

3 Transversely dense functions

Lemma 1 suggests that pairs of functions separated by an integer (norm) distance are particularly problematic. In fact more is true: not only the maximum distance is important, but also places where pairs of functions differ by an integer (even if that integer is not the largest distance between the functions). A key issue underlying most of this paper is that any graph isomorphism has to preserve “crossing patterns”. Two continuous functions f and g *cross* at x if $f(x) - g(x) \in \mathbb{Z}$. We define $\text{cross}(f, g)$ to be $\{x: f(x) - g(x) \in \mathbb{Z}\}$. If \mathcal{F} is a set of functions, then $\text{cross}(\mathcal{F})$ denotes $\bigcup_{f, g \in \mathcal{F}; f \neq g} \text{cross}(f, g)$. If \mathcal{F} is a collection of continuous functions, then notice that for all $f, g \in \mathcal{F}$, $\lfloor f - g \rfloor$ is constant on each component of $[0, 1] \setminus \text{cross}(\mathcal{F})$.

For the first class of distributions on $X^{\mathbb{N}}$ that we show to be geometrically Rado, we impose the condition that crossing sets are discrete and disjoint. A subset V of $C[0, 1]$ is said to be *transverse* if the following two properties hold:

- (Tr1) For each pair of distinct functions, $f, g \in V$, $\text{cross}(f, g)$ is finite; does not contain 0 or 1; and does not contain any local maxima or minima of $f - g$.
- (Tr2) The sets $\text{cross}(f, g)$ are disjoint as (f, g) runs over the pairs of distinct functions in V .

Given a finite transverse collection of continuous functions, f_1, \dots, f_m , its *crossing data*, $\text{CD}(f_1, \dots, f_m)$ consists of a *crossing tuple* (x_1, \dots, x_{k-1}) and an array $(d_{ab}^{(i)})_{1 \leq a < b \leq m; 1 \leq i \leq k}$ of integers so that

$$\text{cross}(\{f_1, \dots, f_m\}) = \{x_1, \dots, x_{k-1}\};$$

$x_i < x_{i+1}$ for each i ; and $(f_b - f_a)$ takes values in the range $(d_{ab}^{(i)}, d_{ab}^{(i)} + 1)$ for $x \in (x_{i-1}, x_i)$ where x_0 is taken to be 0 and x_k is taken to be 1.

If $\text{CD}(f_1, \dots, f_m) = ((x_1, \dots, x_{k-1}), (d_{ab}^{(i)}))$ and $\text{CD}(g_1, \dots, g_m) = ((x'_1, \dots, x'_{k'-1}), (d'_{ab}{}^{(i)}))$, then we write $\text{CD}(f_1, \dots, f_m) \sim_{\varepsilon} \text{CD}(g_1, \dots, g_m)$ if $k = k'$; $|x_i - x'_i| < \varepsilon$ for each i ; and $d'_{ab}{}^{(i)} = d_{ab}^{(i)}$ for all $1 \leq a < b \leq m$, $1 \leq i \leq k$. If $k = k'$ and $d'_{ab}{}^{(i)} = d_{ab}^{(i)}$ for all i, a, b with no restriction on $|x_i - x'_i|$, then we write $\text{CD}(f_1, \dots, f_m) \sim_1 \text{CD}(g_1, \dots, g_m)$.

Finally, a countable subset, V , of $C[0, 1]$ is *transversely dense* if the following properties hold:

(TD1) V is transverse.

(TD2) For each finite collection $\mathcal{F} = \{f_1, \dots, f_m\}$ in V , for each $f \in C[0, 1] \setminus \mathcal{F}$ such that $\mathcal{F} \cup \{f\}$ is transverse, for each $\varepsilon > 0$, there exist infinitely many $g \in V \setminus \mathcal{F}$ such that $\|f - g\| < \varepsilon$ and $\text{CD}(f_1, \dots, f_m, g) \sim_\varepsilon \text{CD}(f_1, \dots, f_m, f)$.

In particular, a transversely dense subset of $C[0, 1]$ is dense. We say a sequence (f_1, f_2, \dots) of distinct functions is transversely dense if $\{f_1, f_2, \dots\}$ is transversely dense. A function g satisfying (TD2) is called an ε -approximation of f relative to \mathcal{F} . These ε -approximations are the key to the back-and-forth technique that we use.

Theorem 6. *Let V and W be countable transversely dense subsets of $C[0, 1]$ and let $0 < p, p' < 1$. If G and G' are sampled from $\text{LARG}(V, p)$ and $\text{LARG}(W, p')$, respectively, then G and G' are almost surely isomorphic.*

In particular, there exists a graph G_{TD} such that if μ is a measure on X where $\mu^{\mathbb{N}}$ -a.e. sequence is transversely dense, then μ is universally geometrically Rado and for $\mu^{\mathbb{N}}$ -a.e. vertex set V , almost every element of $\text{LARG}(V, p)$ is isomorphic to G_{TD} for any $0 < p < 1$.

Proof. Let $V = \{f_1, f_2, \dots\}$ and $W = \{g_1, g_2, \dots\}$ and let G and G' be sampled from $\text{LARG}(V, p)$ and $\text{LARG}(W, p')$, respectively. We describe a countable-step algorithm to build an isomorphism from G to G' that fails at each time-step with probability 0; and if it does not fail at any time-step, yields an isomorphism of G and G' .

We use a back-and-forth process to (almost surely) build two sequences (i_n) and (j_n) with the properties:

(T6-1) For each $n \in \mathbb{N}$, the graphs induced by G on $\{f_{i_1}, \dots, f_{i_n}\}$ and by G' on $\{g_{j_1}, \dots, g_{j_n}\}$ are isomorphic by the map sending f_{i_i} to g_{j_i} ;

(T6-2) For each $n \in \mathbb{N}$, $\{i_1, \dots, i_n\}$ and $\{j_1, \dots, j_n\}$ contain $\{m \in \mathbb{N} : m \leq \frac{n}{2}\}$;

(T6-3) $\text{CD}(f_{i_1}, \dots, f_{i_n}) \sim_1 \text{CD}(g_{j_1}, \dots, g_{j_n})$.

It is evident that if properties (T6-1) and (T6-2) are satisfied, then G and G' are isomorphic. Property (T6-3) ensures that the partial isomorphism can always be extended.

To describe the algorithm, we initially set up two queues: Q and Q' , initially consisting of (f_1, f_2, \dots) and (g_1, g_2, \dots) . As usual in the back-and-forth process, we take an unmatched vertex in one of the graphs and match it to a suitable unmatched vertex in the other graph. To find a suitable vertex, transverse density is shown to yield infinitely many suitable vertices to match to (based only on the crossing pattern). We then apply the Borel-Cantelli lemma to ensure that we pick a vertex with the correct adjacencies to the previously matched vertices to ensure that the enlarged matching remains a graph isomorphism. A critical hypothesis here is that all of the events being tested in a single application of the lemma are independent, and it is helpful if they all have equal probabilities. For this reason, when searching for a match for a particular vertex in one of the graphs, we *only test vertices in the other graph that have not previously been tested*. That is, we take elements from the appropriate queue and stop when we find a match. Elements that are tested are removed from the queue. We make a further induction hypothesis.

(T6-4) The queues Q and Q' consist of all but finitely many vertices of G and G' .

For the base case, we set $i_1 = j_1 = 1$ and remove f_1 and g_1 from Q and Q' . We then inductively assume that i_1, \dots, i_{n-1} and j_1, \dots, j_{n-1} have been chosen so that the induction hypotheses are satisfied. If n is even, then let i_n be the first unmatched vertex in G , while if n is odd, let j_n be the first unmatched vertex in G' . We then seek suitable j_n or i_n , respectively. Since the processes are essentially identical, we deal only with the case where n is even, so we are looking for a suitable j_n .

We first identify a target function g in $C[0, 1]$, and then find a suitable approximation in W . Let $\text{cross}(f_{i_1}, \dots, f_{i_{n-1}}) = \{x_1, \dots, x_{k-1}\}$. By the induction hypothesis (T6-3), we may write

$$\text{cross}(g_{j_1}, \dots, g_{j_{n-1}}) = \{y_1, \dots, y_{k-1}\},$$

and define $x_0 = y_0 = 0$ and $x_k = y_k = 1$. We build the target function g as follows. Let $\psi: [0, 1] \rightarrow [0, 1]$ be the continuous increasing piecewise linear map sending each $[y_{l-1}, y_l]$ to $[x_{l-1}, x_l]$. By (T6-3), we have $g_{j_a}(y) + p < g_{j_b}(y) + q$ if and only if $f_{i_a}(\psi(y)) + p < f_{i_b}(\psi(y)) + q$ for $a, b \in \{1, \dots, n-1\}$ and $p, q \in \mathbb{Z}$. Let ϕ_y be the piecewise linear map from \mathbb{R} to \mathbb{R} satisfying

$$\phi_y(f_{i_m}(\psi(y)) + p) = g_{j_m}(y) + p \tag{1}$$

for each $1 \leq m < n$ and $p \in \mathbb{Z}$ that is linear between consecutive values of $f_{i_m}(y) + p$'s. This satisfies $\phi_y(t+1) = \phi_y(t) + 1$ for each $t \in \mathbb{R}$ and $y \in [0, 1]$. We write $f_{i,p}(x)$ for $f_i(x) + p$ and similarly $g_{j,p}(y) = g_j(y) + p$ for any $x, y \in [0, 1]$ and $p \in \mathbb{Z}$.

The map obtained sending $f_{i_m,p}$ to $g_{j_m,p}$ is of the type described in Lemma 5 (also described informally prior to the statement of that lemma). The map ψ is the piecewise linear homeomorphism sending the original crossing points of the (finite collection of) f 's to the crossing points of the corresponding g 's. The step-isometries over each x coordinate are piecewise linear, and send integer translates of each f to the corresponding integer translate of a g .

The fact that $\{f_{i_1}, \dots, f_{i_{n-1}}\}$ and $\{g_{j_1}, \dots, g_{j_{n-1}}\}$ are transverse (specifically the fact that f_{i_a} and f_{i_b} cross at $\psi(y)$ if and only if g_{j_a} and g_{j_b} cross at y) ensures that the map $(y, t) \mapsto \phi_y(t)$ is continuous. To see this, notice that by (1), for $t \notin S(y) := \{f_{i_a}(\psi(y)) + p : 1 \leq a < n; p \in \mathbb{Z}\}$, the piecewise linear function ϕ_y satisfies

$$\phi_y(t) = \frac{[f_{i_M,P}(\psi(y)) - t]g_{j_m,P}(y)}{f_{i_M,P}(\psi(y)) - f_{i_m,P}(\psi(y))} + \frac{[t - f_{i_m,P}(\psi(y))]g_{j_M,P}(y)}{f_{i_M,P}(\psi(y)) - f_{i_m,P}(\psi(y))},$$

where m and p are such that $f_{i_m,p}(\psi(y)) = \max(S(y) \cap (-\infty, t))$; and M and P are such that $f_{i_M,P}(\psi(y)) = \min(S(y) \cap (t, \infty))$. If t is close to $f_{i_a,b}(\psi(y_0))$ and y is close to y_0 , then $\phi_y(t)$ is close to $g_{j_a,b}(y_0)$.

The target function is defined by

$$g(y) = \phi_y(f_{i_n}(\psi(y))).$$

By construction, the places where g crosses $g_{j_1}, \dots, g_{j_{n-1}}$ are the pre-images under ψ of the places where f_{i_n} crosses $f_{i_1}, \dots, f_{i_{n-1}}$. We see that $g(y) < g_{j_m,p}(y)$ if and only if $f_{i_n}(\psi(y)) < f_{i_m,p}(\psi(y))$. It follows that $\text{CD}(f_{i_1}, \dots, f_{i_n}) \sim_1 \text{CD}(g_{j_1}, \dots, g_{j_{n-1}}, g)$. Hence, by the transverse density hypothesis, there exist infinitely many ℓ 's such that $\text{CD}(f_{i_1}, \dots, f_{i_n}) \sim_1 \text{CD}(g_{j_1}, \dots, g_{j_{n-1}}, g_\ell)$. Notice that $\|f_{i_a} - f_{i_n}\| < 1$ if and only if $\|g_{j_a} - g_\ell\| < 1$ for each of these ℓ 's and each $1 \leq a \leq n-1$. Sequentially testing only those ℓ 's

(all but finitely many of these) that remain in Q' , each has probability at least $\min(p', 1 - p')^{n-1}$ (where p' is the parameter in $\text{LARG}(W, p')$) to have the same pattern of adjacencies to $g_{j_1}, \dots, g_{j_{n-1}}$ as f_{i_n} has to $f_{i_1}, \dots, f_{i_{n-1}}$, establishing the induction hypothesis for n . The (almost surely finitely many) tested g 's should then be removed from Q' and f_{i_n} should be removed from Q . \square

We now exhibit two examples of measures on $C[0, 1]$ satisfying the hypotheses of the theorem, supported on polynomials and piecewise linear functions, respectively. Let P be the map from $\mathbb{N} \times \mathbb{R}^{\mathbb{N}_0}$ to $C[0, 1]$ defined by $P(k, a_0, a_1, \dots) = f$, where $f(t) = \sum_{i=0}^k a_i t^i$. Let $\mathbb{N} \times \mathbb{R}^{\mathbb{N}_0}$ be equipped with the product measure, where the first coordinate is geometrically distributed with parameter $\frac{1}{2}$ (that is, $\mathbb{P}(N = k) = 1/2^k$ for $k = 1, 2, \dots$). The remaining coordinates are standard normal random variables. Let μ_{poly} be the measure on $C[0, 1]$ obtained as the push-forward of \mathbb{P} under P (that is, a randomly sampled element is a polynomial of degree k where k is a geometric random variable whose coefficients are sampled from independent standard normal random variables).

Theorem 7. *Let μ_{poly} be as above. For $\mu_{\text{poly}}^{\mathbb{N}}$ -a.e. subset V of $C[0, 1]$, and each $0 < p < 1$, almost every realization of $\text{LARG}(V, p)$ is isomorphic to G_{TD} .*

Proof. Let a randomly sampled V be $\{f_1, f_2, \dots\}$. We show that V is transversely dense, so that the desired result follows from Theorem 6

We first establish that V is almost surely transverse. Since μ_{poly} is a continuous measure, the (f_i) are almost surely pairwise distinct, so that $\text{cross}(f_i, f_j)$ is finite since the functions are polynomials. If f is sampled from μ_{poly} , we can consider $f(0)$ and $f(1)$ as random variables. These also have continuous distributions, so the probability that $\text{cross}(f_i, f_j)$ intersects $\{0, 1\}$ is 0 for each of the countably many (i, j) pairs.

For the local maxima and minima of $f_i - f_j$, the (finitely many) locations are determined by the non-constant terms of the pair of $f_i(t) - f_j(t)$. Conditioned on the non-constant terms, the value of the difference at the locations of the critical points is again continuously distributed, so that almost surely, $f_i - f_j$ does not differ by an integer at any critical point for each of the countably many (i, j) pairs. We have established (Tr1).

Given a pair of functions f_i and f_j , there are finitely many crossings. Now given the coefficients of the polynomial f_k except for the constant term, we see that conditioned on all of this information, the probability that f_i, f_j and f_k have a common crossing is 0. A similar argument shows that the probability that two other functions f_k and f_l cross at one of the crossings of f_i and f_j is 0, establishing (Tr2).

Finally, we establish (TD2). Let $\mathcal{F} = \{f_1, \dots, f_m\}$ (this may be done without loss of generality by expanding the finite collection if necessary). Let $f \in C[0, 1]$ be such that $\mathcal{F} \cup \{f\}$ is transverse and let $\varepsilon > 0$. We may assume without loss of generality that ε is less than $\frac{1}{2} \min\{|x - y| : x, y \in \text{cross}(f_1, \dots, f_m, f), x \neq y\}$. First, by the transverse assumption, we may replace f by a continuously differentiable function g such that $\text{CD}(f_1, \dots, f_m, g) = \text{CD}(f_1, \dots, f_m, f)$ and $(g - f_i)'$ is non-zero at each crossing of g and f_i . Applying the Weierstrass approximation theorem to g' , there is a polynomial q , of degree d say, and a $\delta > 0$ such that if $\|Dh - q\| < \delta$ and $|h(0) - g(0)| < \delta$, then $\|h - g\| < \varepsilon$ and $\text{CD}(f_1, \dots, f_m, h) \sim_{\varepsilon} \text{CD}(f_1, \dots, f_m, g)$. Now there is an open collection of polynomials of degree $d + 1$ satisfying these constraints, so that there are, as required, infinitely many f_j such that $\text{CD}(f_1, \dots, f_m, f_j) \sim_{\varepsilon} \text{CD}(f_1, \dots, f_m, f)$. \square

We now prove a similar result for piecewise linear functions. Given an $n \in \mathbb{N}$, a sequence of distinct $(x_i)_{i \geq 1}$ taking values in $(0, 1)$ and a sequence of $(y_i)_{i \geq 0}$ taking real values, let $\Phi(n, (x_i), (y_i))$ be the piecewise linear function with n pieces joining (\hat{x}_{i-1}, y_{i-1}) to (\hat{x}_i, y_i) for $i = 1, \dots, n$ by a straight line where $\hat{x}_0 = 0$, $\hat{x}_n = 1$ and \hat{x}_i is the i th smallest element of $\{x_1, \dots, x_{n-1}\}$. Equip $\mathbb{N} \times (0, 1)^{\mathbb{N}} \times \mathbb{R}^{\mathbb{N}_0}$ with the product of the geometric distribution with parameter $\frac{1}{2}$ on the first coordinate; independent identically distributed Lebesgue measure on the x coordinates; and independent standard normals on the y coordinates, and let μ_{PL} denote the push-forward measure under the map Φ on the space $C[0, 1]$.

Theorem 8. *Let μ_{PL} be as above. For $\mu_{PL}^{\mathbb{N}}$ -a.e. subset V of $C[0, 1]$, and each $0 < p < 1$, almost every realization of $\text{LARG}(V, p)$ is isomorphic to G_{TD} .*

Proof. Let a randomly sampled V be $\{f_1, f_2, \dots\}$. We show that V is transversely dense, so that the desired result follows from Theorem 6. First notice that the slope of each piece of a function sampled from μ_{PL} is continuously distributed, so that almost surely, no two pieces of any f_i and f_j have the same slope. This ensures that $\text{cross}(f_i, f_j)$ is finite for each i, j . As before, the distributions of $f(0)$ and $f(1)$ are continuous, so with probability 1, no crossings occur at 0 or 1. For any two piecewise linear functions f_i and f_j with different slopes, local maxima occur only at the endpoints of the intervals. Given the function f_j and the x values (but not the y values) defining f_i , the conditional probability that a crossing occurs at one of these values is 0, so that almost surely no crossings occur at local maxima or minima of $f_i - f_j$, establishing (Tr1).

To show (Tr2), condition on f_i and f_j . These functions almost surely have finitely many crossings. Since at each of these crossings, the values of f_k and f_l have independent continuous distributions, the probability that f_k crosses f_i and f_j at one of these points; or that f_k and f_l cross at one of these points is 0.

Finally, let $\mathcal{F} = \{f_1, \dots, f_m\}$ and let $\mathcal{F} \cup \{f\}$ be transverse. There is a piecewise linear function g with $\|f - g\|$ arbitrarily small such that $\text{CD}(f_1, \dots, f_m, g) = \text{CD}(f_1, \dots, f_m, f)$. Any suitably close approximation with the same number of pieces will satisfy $\text{CD}(f_1, \dots, f_m, h) \sim_{\varepsilon} \text{CD}(f_1, \dots, f_m, f)$, so that (TD2) follows. \square

We exhibit integrated Brownian motion as a third example. Let ν be the Wiener measure on standard Brownian motions on $[0, 1]$. We then define a map $\Phi: \mathbb{R} \times C[0, 1] \rightarrow C[0, 1]$ by

$$[\Phi(a, f)](x) = a + \int_0^x f(t) dt.$$

We then define a measure μ_{IBM} on $C[0, 1]$ that is the push-forward of the product of a standard normal distribution and ν under the map Φ . In other words, a function sampled from μ_{IBM} is obtained by adding a normally-distributed initial condition to the anti-derivative of a standard Brownian motion.

Theorem 9. *Let μ_{IBM} be as above. For $\mu_{IBM}^{\mathbb{N}}$ -a.e. subset V of $C[0, 1]$, and each $0 < p < 1$, almost every realization of $\text{LARG}(V, p)$ is isomorphic to G_{TD} .*

We give a brief proof, primarily emphasizing the parts that differ from the previous examples.

Proof. Recall that the difference of two independent standard normal random variables is a normal random variable with variance 2. Similarly, the difference of two independent normal standard Brownian motions

is a Brownian motion with parameters $\mu = 0$ and $\sigma^2 = 2$. Hence, the difference of two independent realizations of μ_{IBM} is, up to scaling by a factor of $\sqrt{2}$, of the same form. Fubini's theorem implies that the zero set of a Brownian motion almost surely has measure 0, so that the measure of the set of critical points of a realization of μ_{IBM} is almost surely 0. Since realizations of Brownian motion on $[0,1]$ are continuous, they are bounded, so that each realization of μ_{IBM} is almost surely Lipschitz. Conditioned on the Wiener process f , it follows that for all values of a except for a set of measure 0, $\Phi(a, f)$ does not take an integer value at any critical point. It follows that almost surely, if f_1 and f_2 are realizations of μ_{IBM} , their difference has non-zero derivative at each element of $\text{cross}(f_1, f_2)$. This ensures that $\text{cross}(f_1, f_2)$ is closed, and almost surely discrete, hence finite. The remainder of the proof that a countable set of independently chosen realizations of integrated Brownian motions are transverse is similar to the previous proofs.

Next, to show that a countable set of independently chosen realizations of integrated Brownian motions satisfies (TD2), let $\mathcal{F} = \{f_1, \dots, f_m\}$ and suppose $f \in C[0, 1] \setminus \mathcal{F}$ is such that $\mathcal{F} \cup \{f\}$ is transverse. Let $\varepsilon > 0$. There is a C^1 function g such that $g'(0) = 0$; $\|g - f\|_{C[0,1]} < \frac{\varepsilon}{2}$; $\text{cross}(\mathcal{F} \cup \{g\}) = \text{cross}(\mathcal{F} \cup \{f\})$; and so that $(f_i - g)'$ is non-zero at each crossing. Then there is a C^1 -neighbourhood, N , of g so that all elements h of N satisfy $\|f - h\|_{C_0[0,1]} < \varepsilon$ and $\text{CD}(f_1, \dots, f_m, h) \sim_\varepsilon \text{CD}(f_1, \dots, f_m, f)$. Since the measure on Brownian motions gives positive measure to every open subset of $\{k \in C_0[0, 1] : k(0) = 0\}$, the measure of the pairs (a, f) such that $\Phi(a, f)$ lies in N is positive, so that there are almost surely infinitely many f_k 's satisfying the conditions of (TD2). \square

4 IC-dense sets and Brownian motion

By definition, transversely dense sets must contain functions that cross finitely often and transversely with other functions in the set. In this section, we establish results for functions in $C[0, 1]$ that exhibit the opposite type of behaviour: pairs of functions cross infinitely often. We will call sets of such functions *infinite crossing dense* (IC-dense). An example of such functions is formed by Brownian motions. We will first show that any two LARG graphs with IC-dense vertex sets are almost surely isomorphic. We will then define a natural probability measure on the set of Brownian motions, and show that under this measure almost all countable sets of Brownian motions are IC-dense.

A countable subset V of $C[0, 1]$ will be called *IC-dense* if it has the following properties:

- (IC1) The set V is dense in $C[0, 1]$;
- (IC2) The sets $\text{cross}(f, g)$ are disjoint for distinct pairs $f, g \in V$;
- (IC3) For any $f, g \in V$, $\text{cross}(f, g)$ does not contain 0 or 1;
- (IC4) For each distinct pair $f, g \in V$, if $x \in \text{cross}(f, g)$ for $x \in (0, 1)$, then there exists a sequence of points x_n converging to x monotonically such that $f(x_{2n}) - g(x_{2n}) < f(x) - g(x)$ and $f(x_{2n+1}) - g(x_{2n+1}) > f(x) - g(x)$ for each n .

We comment on (IC4). The requirement that the sequence (x_n) converges to x monotonically is to ensure that there is an infinite sequence of crossings of f and g converging to x (whereas if f and g cross transversely, it is straightforward to build a non-monotonic sequence of x_n 's with the alternating property).

The reason that we only require that there is an infinite sequence of x_n 's on one side of x is that the set of crossings of two Brownian motions is perfect (each crossing is a limit of other crossings), but while some crossings may be approached by crossings on either side, there are crossings that may be only approached on one side. To see this, notice that for any pair of continuous functions that cross, the set of crossings is closed, and therefore has a maximum.

4.1 IC-dense sets are geometrically Rado

We will now show that IC-dense sets have the geometric Rado property. As in the case for transversely dense sets, the proof is based on the inductive construction of a graph isomorphism. While in the transversely dense case we required that the isomorphism exactly preserves the crossing behaviour of the functions, here we wish to preserve the decomposition of the interval $[0, 1]$ into subintervals where pairs of functions cross infinitely often. Before we state the main theorem, we give definitions and useful lemmas.

Crossing partitions. If \mathcal{F} is a finite subset of an IC-dense of $C[0, 1]$, then we build a finite partition of $[0, 1]$ into open and closed sub-intervals by the following procedure. Let

$$a_1 = \min\{x \in [0, 1] : g(x) - h(x) \in \mathbb{Z} \text{ for some distinct } g, h \in \mathcal{F}\}. \tag{2}$$

Suppose a_1, \dots, a_n have been found, and let $(g_i, h_i, m_i)_{i=1}^n$ be the tuples in $\mathcal{F} \times \mathcal{F} \times \mathbb{Z}$ with $g_i \neq h_i$ so that $g_i(a_i) - h_i(a_i) = m_i$ for each $1 \leq i \leq n$. We then define

$$a_{n+1} = \min\{x \in (a_n, 1] : h(x) - g(x) = m \in \mathbb{Z} \text{ for some distinct } g, h \in \mathcal{F} \\ \text{such that } \{g, h, m\} \neq \{g_n, h_n, m_n\}\}. \tag{3}$$

If a_{n+1} is defined (that is, the set is not empty), then set $g_{n+1} = g$, $h_{n+1} = h$ and $m_{n+1} = m$. If not, then the procedure terminates with a_n .

Note that it is possible that $\{g_{n+1}, h_{n+1}\} = \{g_n, h_n\}$, but only if they differ by one more or less at a_{n+1} than a_n . This, in turn, is only possible if there are exactly two functions. If there are any more functions, then g_n or h_n would have to cross another element of \mathcal{F} before crossing each other again.

Lemma 10. *The procedure to generate the sequence (a_n) as defined in (2) and (3) terminates in a finite number of steps.*

Proof. Suppose, for a contradiction, that there are infinitely many a_n 's. Since (a_n) is an increasing sequence lying in $[0, 1]$, let its limit be a . By the Pigeonhole Principle, there are functions f^1, f^2, f^3 and f^4 in \mathcal{F} (not necessarily distinct, but such that $f^1 \neq f^2, f^3 \neq f^4$ and $\{f^1, f^2\} \neq \{f^3, f^4\}$) and a subsequence (n_j) such that $\{g_{n_j}, h_{n_j}\} = \{f^1, f^2\}$ and $\{g_{n_{j+1}}, h_{n_{j+1}}\} = \{f^3, f^4\}$ for all j . We have $f^1(a_{n_j}) - f^2(a_{n_j}) \in \mathbb{Z}$ and $f^3(a_{n_{j+1}}) - f^4(a_{n_{j+1}}) \in \mathbb{Z}$ for each j . Taking a limit, and using continuity of the functions involved, we see that $f^1(a) - f^2(a) \in \mathbb{Z}$ and $f^3(a) - f^4(a) \in \mathbb{Z}$. That is, $a \in \text{cross}(f^1, f^2) \cap \text{cross}(f^3, f^4)$. This contradicts (IC2), establishing the above procedure terminates as required. \square

Now for each i such that a_i is defined, let $b_i = \max\{x \in [a_i, a_{i+1}] : g_i(x) - h_i(x) \in \mathbb{Z}\}$. Note that, since $\text{cross}(g_i, h_i)$ is closed, this maximum is defined, and since h_i and g_i cross at b_i and a different pair h_{i+1}, g_{i+1} crosses at a_{i+1} , we have that $b_i < a_{i+1}$. The *crossing partition* $\mathcal{P}(\mathcal{F})$ is then given by

$$\mathcal{P}(\mathcal{F}) = \{[0, a_1], [a_1, b_1], (b_1, a_2), [a_2, b_2], \dots, [a_n, b_n], (b_n, 1]\},$$

so that on the *stable intervals*, that is the (relatively) open sets $[0, a_1), (b_1, a_2), \dots, (b_{n-1}, a_n)$ and $(b_n, 1]$, there are no crossings, while on the *crossing intervals*, $[a_i, b_i]$, g_i and h_i cross infinitely often, but there are no other crossings.

The *crossing data* for \mathcal{F} consists of the crossing partition together with a collection of integers $d_{f,g}^I$, one for each pair of distinct elements of \mathcal{F} and interval $I \in \mathcal{P}(\mathcal{F})$ except that $d_{g_i, h_i}^{[a_i, b_i]}$ is not defined. Except in this case, for all x in I and all $f, g \in \mathcal{F}$, then $g(x) - f(x) \in (d_{f,g}^I, d_{f,g}^I + 1)$. We denote this data by $\text{CD-inf}(\mathcal{F})$. As a consequence, setting $\bar{\mathcal{F}} = \{f + n : f \in \mathcal{F}, n \in \mathbb{Z}\}$, the elements of $\bar{\mathcal{F}}$ maintain a constant ordering on the stable sets. The elements of $\bar{\mathcal{F}} \setminus \{g_i\}$ and $\bar{\mathcal{F}} \setminus \{h_i\}$ maintain a constant ordering on the crossing intervals.

We write $\text{CD-inf}(f_1, \dots, f_k) \sim_\varepsilon \text{CD-inf}(g_1, \dots, g_k)$ if $\mathcal{P}(\{f_1, \dots, f_k\})$ contains the same number of intervals as $\mathcal{P}(\{g_1, \dots, g_k\})$; the functions crossing in each crossing interval of $\mathcal{P}(\{f_1, \dots, f_k\})$ have the same indices as the functions crossing in the corresponding crossing interval of $\mathcal{P}(\{g_1, \dots, g_k\})$; $|a'_i - a_i| < \varepsilon$ and $|b'_i - b_i| < \varepsilon$ for each i ; and $d_{f_i, f_j}^I = d_{g_i, g_j}^{I'}$ whenever these are defined (where I and I' are corresponding intervals of $\mathcal{P}(\{f_1, \dots, f_k\})$ and $\mathcal{P}(\{g_1, \dots, g_k\})$). As before $\text{CD-inf}(f_1, \dots, f_k) \sim_1 \text{CD-inf}(g_1, \dots, g_k)$ requires that the partitions have the same number of elements; the same pairs cross in the corresponding crossing intervals and the gap between a pair of functions has the same floor in each interval of the crossing partition as the corresponding pair has in the corresponding interval (with the exception of the pair that is crossing in a crossing interval).

Theorem 11. *Let V and W be countable IC-dense subsets of $C[0, 1]$ and let $0 < p, p' < 1$. If G and G' are sampled from $\text{LARG}(V, p)$ and $\text{LARG}(W, p')$, respectively, then G and G' are almost surely isomorphic.*

In particular, there exists a graph G_{ICD} such that if μ is a measure on $C[0, 1]$ such that $\mu^{\mathbb{N}}$ -a.e. sequence is IC-dense, then μ is universally geometrically Rado, and for $\mu^{\mathbb{N}}$ -a.e. vertex set V , almost every element of $\text{LARG}(V, p)$ is isomorphic to G_{ICD} for any $0 < p < 1$.

Proof. Let V and W be infinite crossing dense subsets of $C([0, 1])$ and let $p, p' \in (0, 1)$. We show that a realization of $\text{LARG}(V, p)$ is almost surely isomorphic to a realization of $\text{LARG}(W, p')$. As in the proof of Theorem 6 for transversely dense sets, we construct an isomorphism via a back-and-forth argument.

The key step of the proof is to show that if \mathcal{F} and \mathcal{G} are finite subsets of V and W such that $\text{CD-inf}(\mathcal{F}) \sim_1 \text{CD-inf}(\mathcal{G})$ then given any additional $f \in V \setminus \mathcal{F}$, there is almost surely a $g \in W \setminus \mathcal{G}$ such that $\text{CD-inf}(\mathcal{F} \cup \{f\}) \sim_1 \text{CD-inf}(\mathcal{G} \cup \{g\})$ and g has the same adjacencies to \mathcal{F} as f has to \mathcal{G} .

Let G and G' be sampled from $\text{LARG}(V, p)$ and $\text{LARG}(W, p')$, respectively. As before, for the purposes of maintaining independence when applying the Borel-Cantelli lemma, we maintain queues of functions to test, ensuring that we never test whether a pair of vertices is adjacent more than once. Let $Q = (f_1, f_2, \dots)$ be an enumeration of \mathcal{F} and $Q' = (g_1, g_2, \dots)$ be an enumeration of \mathcal{G} . As before, we build sequences (i_n) and (j_n) satisfying the following:

- (T11-1) For each n , the graphs induced by G on $\{f_{i_1}, \dots, f_{i_n}\}$ and by G' on $\{g_{j_1}, \dots, g_{j_n}\}$ are isomorphic by mapping f_{i_i} to g_{j_i} ;
- (T11-2) For each $n \in \mathbb{N}$, $\{i_1, \dots, i_n\}$ and $\{j_1, \dots, j_n\}$ contain the set $\{m \in \mathbb{N} : m \leq \frac{n}{2}\}$;
- (T11-3) $\text{CD-inf}(f_{i_1}, \dots, f_{i_n}) \sim_1 \text{CD-inf}(g_{j_1}, \dots, g_{j_n})$;

(T11-4) the remaining elements of the queues Q and Q' at the n th stage consist of all but finitely many elements of V and W respectively.

When $n = 1$, we let $i_1 = j_1 = 1$ and remove f_1 and g_1 from the respective queues. We focus on a single step in one direction. Suppose that the induction hypotheses are satisfied after the $(n - 1)$ st step. If n is even, let i_n be $\min(\mathbb{N} \setminus \{i_1, \dots, i_{n-1}\})$, while if n is odd, let $j_n = \min(\mathbb{N} \setminus \{j_1, \dots, j_{n-1}\})$. For even n , we then seek a suitable j_n to identify with i_n , while if n is odd, we seek a suitable i_n . We focus on the case of even n , the case of odd n being essentially identical.

Let $\mathcal{F} = \{f_{i_1}, \dots, f_{i_{n-1}}\}$ and $\mathcal{G} = \{g_{j_1}, \dots, g_{j_{n-1}}\}$. Let the crossing partitions of $[0, 1]$ of \mathcal{F} and \mathcal{G} be

$$\begin{aligned} \mathcal{P}(\mathcal{F}) &= \{[0, a_1], [a_1, b_1], (b_1, a_2), [a_2, b_2], \dots, [a_N, b_N], (b_N, 1]\} \text{ and} \\ \mathcal{P}(\mathcal{G}) &= \{[0, a'_1], [a'_1, b'_1], (b'_1, a'_2), [a'_2, b'_2], \dots, [a'_N, b'_N], (b'_N, 1]\}. \end{aligned}$$

Notice that $\mathcal{P}(\mathcal{F} \cup \{f_{i_n}\})$ is a refinement of $\mathcal{P}(\mathcal{F})$: the elements of $\mathcal{P}(\mathcal{F})$ may be sub-divided in $\mathcal{P}(\mathcal{F} \cup \{f_{i_n}\})$ due to intersections of f_{i_n} with elements of \mathcal{F} . For an element I of $\mathcal{P}(\mathcal{F})$ the sub-partition \mathcal{P}_I of I is obtained by taking those elements of $\mathcal{P}(\mathcal{F} \cup \{f_{i_n}\})$ that lie in I . (Note that \mathcal{P}_I may be the trivial partition of I if f_{i_n} does not cross any element of \mathcal{F} in I).

If $I = [a_k, b_k]$ is a crossing interval in $\mathcal{P}(\mathcal{F})$, then two functions f_{i_l} and f_{i_m} cross infinitely often in I . By construction, no other functions of \mathcal{F} cross in I . The sub-partition, \mathcal{P}_I consists of crossing intervals in which f_{i_l} and f_{i_m} cross infinitely often; stable intervals (with no crossings); and crossing intervals in which f_{i_n} crosses exactly one element of \mathcal{F} (possibly f_{i_l} or f_{i_m}), again infinitely often. The situation is illustrated schematically in Figure 1. If I is a stable interval in $\mathcal{P}(\mathcal{F})$, then \mathcal{P}_I consists of stable intervals, and intervals in which f_{i_n} crosses exactly one element of \mathcal{F} infinitely often.

In each crossing interval of \mathcal{P}_I , there are crossings between exactly one pair of functions in $\mathcal{F} \cup \{f_{i_n}\}$. As in the proof of Theorem 6, for $a \in \mathbb{Z}$, we write $f_{i,a}(x) = f_i(x) + a$.

We build a target function g such that all $h \in W \setminus \mathcal{G}$ sufficiently close to g satisfy $\text{CD-inf}(f_{i_1}, \dots, f_{i_n}) \sim_1 \text{CD-inf}(g_{j_1}, \dots, g_{j_{n-1}}, h)$, which will allow us to maintain (T11-3). As in the proof of Theorem 6, this function is built by interpolating various convex combinations of the elements of \mathcal{G} .

Let I' be the element of $\mathcal{P}(\mathcal{G})$ corresponding to an interval I of $\mathcal{P}(\mathcal{F})$. We consider separately the cases where I is a stable interval of $\mathcal{P}(\mathcal{F})$ and where I is a crossing interval of $\mathcal{P}(\mathcal{F})$. In the case where I is a stable interval of \mathcal{F} , let ψ be the increasing affine map sending I' to I and for each y , as in Theorem 6, let ϕ_y be the piecewise linear function sending $f_{i_k}(\psi(y)) + a$ to $g_{j_k}(y) + a$. The restriction of g to I' is then given by

$$g|_{I'}(y) = \phi_y(f_{i_n}(\psi(y))). \tag{4}$$

In the case where I is a crossing interval, suppose that the functions that cross in I are f_{i_l} and f_{i_m} . We build a target function $g|_{I'}$ in a similar manner to the above, but additional care is needed in the elements of \mathcal{P}_I on which f_{i_l} and f_{i_m} cross.

We first build a suitable partition $\mathcal{Q}_{I'}$ of I' corresponding to \mathcal{P}_I . Let the intervals of \mathcal{P}_I on which f_{i_l} and f_{i_m} cross be $[\alpha_1, \beta_1], \dots, [\alpha_s, \beta_s]$ (with $\alpha_1 = \min I$ and $\beta_s = \max I$). On each of the open intervals separating these crossing intervals, (β_r, α_{r+1}) , $f_{i_m} - f_{i_l}$ takes values in $(d_r, d_r + 1)$ for some $d_r \in \mathbb{Z}$. We then build a partition of I' consisting of sub-intervals $[\alpha'_1, \beta'_1], \dots, [\alpha'_s, \beta'_s]$ (with $\alpha'_1 = \min I'$ and $\beta'_s = \max I'$) of I' containing all crossings of g_{j_l} and g_{j_m} , so that crossings occur at the endpoints and such that the restriction of $g_{j_m} - g_{j_l}$ to $(\beta'_r, \alpha'_{r+1})$ takes values in $(d_r, d_r + 1)$. This is possible since $\text{CD-inf}(f_{i_1}, \dots, f_{i_{n-1}}) \sim_1$




Figure 1: Schematic indicating the refinement of a crossing interval $[a_j, b_j]$ in which the two central functions, f_{i_l} and f_{i_m} are crossing when a new function f (in bold) is added to the collection. The graph illustrates $f_{i_l, p}$ and $f_{i_m, q}$ for suitable p, q so that the functions agree at the crossing, rather than differing by an integer. The closed sub-intervals in the refinement are indicated below.

CD-inf($g_{j_1}, \dots, g_{j_{n-1}}$) and since W satisfies (IC4). Now let $Q_{J'} = \{[\alpha'_1, \beta'_1], (\beta'_1, \alpha'_2), \dots, [\alpha'_s, \beta'_s]\}$ and let ψ be the increasing piecewise affine map that sends each α' and β' to the corresponding α or β . Let J be an element of \mathcal{P}_I and J' be the corresponding element of $Q_{J'}$. If J is a stable interval or an interval on which f_{i_n} crosses an f_{i_a} other than f_{i_l} or f_{i_m} , then define ϕ_y and $g|_{J'}$ as in (4).

If J is an interval on which f_{i_l} and f_{i_m} cross (so that f_{i_n} does not cross either f_{i_l} or f_{i_m}), for $y \in J'$, then let $f_{i_n} - f_{i_l}$ take values in $(a, a + 1)$ and $f_{i_n} - f_{i_m}$ take values in $(b, b + 1)$ on J . We then set $u(x) = \max(f_{i_l}(x) + a, f_{i_m}(x) + b)$ and $v(x) = \min(f_{i_l}(x) + a + 1, f_{i_m}(x) + b + 1)$ on J ; and $u'(y) = \max(g_{j_l}(y) + a, f_{j_m}(y) + b)$ and $v'(y) = \min(g_{j_l}(y) + a + 1, f_{j_m}(y) + b + 1)$ on J' . Notice that since the only functions crossing in J are f_{i_l} and f_{i_m} , there is a single function, either u or a function of the form $f_{i,c}$ with $i \in \{1, \dots, n - 1\} \setminus \{l, m\}$ and $c \in \mathbb{Z}$ that is the closest approximation from below of f_{i_n} throughout the interval J . Similarly, there is a single function $f_{i,d}$ or v that is the closest approximation from above of f_{i_n} . We call these functions $m(x)$ and $M(x)$. Let the corresponding functions on J' be $m'(y)$ and $M'(y)$. Let ϕ_y be the affine function sending $m(\psi(y))$ to $m'(y)$ and $M(\psi(y))$ to $M'(y)$. We then define $g|_{J'}$ as in (4). One may check that the function g defined in this way is continuous since the maps ϕ_y being applied at the endpoints of the intervals agree. For example in Figure 1, ϕ_y is defined at β_1 using $M(x)$ and the function f lying immediately above f_{i_n} , while at β_1^+ , assuming $f_{i,a} > f_{i_m,b}$ on (β_1, α_2) , ϕ_y is defined at β_1^+ using $f_{i,a}$ and f . These agree since $M(\beta_1) = f_{i,a}(\beta_1)$.

By construction, in any of the intervals of $\mathcal{P}(\mathcal{G})$, if f_{i_n} crosses f_{i_a} at $\psi(y)$, then g crosses g_{j_a} at y . Now for any sufficiently small $\varepsilon > 0$, any h satisfying $\|g - h\| < \varepsilon$ has nearby crossings. For sufficiently small $\varepsilon > 0$, if $h \in W$ satisfies $\|g - h\| < \varepsilon$, then $\text{CD-inf}(f_{i_1}, \dots, f_{i_{n-1}}, f_{i_n}) \sim_1 \text{CD-inf}(g_{j_1}, \dots, g_{j_{n-1}}, h)$. In particular, $\|h - g_{j_a}\| = \|f_{i_n} - f_{i_a}\|$ for each a . Each h selected from Q' therefore has probability at least $\min(p', 1 - p')^{n-1}$ of being adjacent to exactly those g_{j_a} 's corresponding to the f_{i_a} 's that are adjacent to f_{i_n} . Take the first suitable g_b from the queue and discard all elements of Q' up to and including g_b . Let $j_n = b$ and delete i_n from Q .

The induction hypotheses almost surely remain satisfied. □

4.2 Brownian motion

We consider the following procedure to select random functions: a starting point (the value at 0) is chosen from a standard normal distribution and from there the function follows a standard Brownian path. Formally, let $(N_n)_{n \in \mathbb{N}}$ be a family of countably many independent standard normal random variables and let $(X_n(t))_{n \in \mathbb{N}}$ be a family of countably many independent standard Brownian motions (with $X_n(0) = 0$, $\mathbb{E}X_n(t) = 0$ and $\text{Var}X_n(t) = t$ for each $n \in \mathbb{N}$ and $t \in [0, 1]$). Let $f_n(t) = N_n + X_n(t)$ be a random function obtained in this manner, which we refer to as a *shifted Brownian motion*.

Theorem 12. *If $(f_n)_{n \in \mathbb{N}}$ is a family of countably many independent shifted Brownian motions sampled from the above distribution, then the collection is almost surely IC-dense.*

Proof. Let $\varepsilon > 0$ be given. Let f be a continuous function. There exists a function g where $g(x_i) = f(x_i)$ for $i = 0, \dots, n$ with $0 = x_0 < x_1 < \dots < x_n = 1$ and g is affine in between such that $\|g - f\| < \frac{\varepsilon}{2}$. By standard properties of Brownian motion, it is known that $\mathbb{P}(|N - y_0| < \frac{\varepsilon}{2n}) > 0$ and that $\mathbb{P}(|X(x_j) - X(x_{j-1}) - (y_j - y_{j-1})| < \frac{\varepsilon}{2n} \mid X(x_1), \dots, X(x_{j-1})) > 0$. Finally, it is known that conditioned on the values of $X(x_1), \dots, X(x_n)$, the probability that X never strays by more than $\frac{\varepsilon}{2}$ from the affine function joining $X(x_i)$ and $X(x_{i+1})$ is positive (the distribution of $(X(t))_{x_i \leq t \leq x_{i+1}}$ conditioned on $X(x_i)$ and $X(x_{i+1})$ is a

so-called *Brownian bridge*). Multiplying these probabilities together, we note that the probability that $\|f_j - f\| < \varepsilon$ is positive, and is the same for each j . Hence, by the second Borel-Cantelli lemma, there almost surely exists a j such that $\|f_j - f\| < \varepsilon$. Since $C([0, 1])$ is separable, the above argument shows that (f_n) is almost surely dense.

Condition (IC2) follows from a corollary of Lévy’s theorem (Corollary 2.23 in [7]) which states that for any non-zero point p in \mathbb{R}^2 , the standard 2-dimensional Brownian motion almost surely does not pass through p . To see (IC2) from this, we deal with two cases: we first show that there are no three functions all crossing at a single x value; and then show that there are not two distinct pairs of functions with both pairs crossing at a single x value. For the first of these, it suffices (since there are countably many triples) to show that almost surely, f_1, f_2 and f_3 do not all cross at any x . That is, it suffices to show that for any integers m and n , with probability 1, there does not exist an x such that $f_2(x) - f_1(x) = m$ and $f_3(x) - f_1(x) = n$. Conditional on $N_1 = y_1, N_2 = y_2$ and $N_3 = y_3$, the values at $x = 0$, we therefore have to show that

$$\mathbb{P}(\exists x: X_2(x) - X_1(x) = m + y_2 - y_1 \text{ and } X_3(x) - X_1(x) = n + y_3 - y_1) = 0.$$

This does not immediately follow from the theorem mentioned above about 2-dimensional Brownian motion since $((X_2 - X_1)(x), (X_3 - X_1)(x))$ is not a standard Brownian motion: the coordinates are correlated. It is, however, expressible as a fixed linear transformation of a standard Brownian motion (see, for example, [7] page 14) and hence, the previous theorem applies.

Similarly, to show that f_1, f_2, f_3 and f_4 do not have the property that f_1 and f_2 cross at some x and f_3 and f_4 cross at the same x , we need to show that

$$\mathbb{P}(\exists x: X_2(x) - X_1(x) = m + y_1 - y_2 \text{ and } X_4(x) - X_3(x) = n + y_3 - y_4) = 0.$$

This follows directly from the quoted theorem since $((X_2 - X_1)(x), (X_4 - X_3)(x))$ is a scaled copy of the standard Brownian motion (by a scale factor of $\sqrt{2}$).

To see that f_1 and f_2 do not cross at 0, we observe that $\mathbb{P}(N_2 = N_1 + n | N_1 = x) = 0$ and so integrating over x , the probability of a crossing at 0 is 0. A similar argument involving $N_1, N_2, X_1(1)$ and $X_2(1)$ shows that the probability of a crossing at 1 is 0, and this proves (IC3).

To verify the last condition, notice that $f_2(x) - f_1(x) = X_2(x) - X_1(x) + N_2 - N_1$. By Theorem 2.11 of [7], a Brownian motion has countably many local extrema and by Theorem 5.14 of [7], no local points of increase or decrease (a point of local increase for a Brownian motion $Z(x)$ is an x_0 such that for some $\delta > 0$, on $(x_0, x_0 + \delta)$, $Z(x) > Z(x_0)$ and on $(x_0 - \delta, x_0)$, $Z(x) < Z(x_0)$, with a point of local decrease being defined similarly). Given X_1 and X_2 , the probability that $N_2 - N_1$ is such that value taken at one of the countably many local extrema of $f_2 - f_1$ is an integer is 0 (since this only occurs for countably many values of $N_2 - N_1$ and N_1 and N_2 are independent and have continuous distributions). Hence, we have that almost surely, each crossing point of $f_2 - f_1$ is not a local extremum or a point of local increase or decrease. Since these possibilities are ruled out, the remaining possibility is the IC-density condition (IC4). \square

Corollary 13. *If V is a countable collection of independent samples of shifted Brownian motion, then for all $p \in (0, 1)$, $\text{LARG}(V, p)$ is almost surely isomorphic to G_{ICD} .*

Proof. By Theorem 12, a countable collection of realizations of Brownian motions with random starting points is almost surely IC-dense. By Theorem 11, if one builds a LARG graph from such a realization, it is almost surely isomorphic to G_{ICD} . \square

5 Non-isomorphism

In the previous sections we showed that LARG graphs on transversely dense sets are almost surely isomorphic to G_{TD} , while LARG graphs on IC-dense sets are almost surely isomorphic to G_{ICD} . In this section, we will show that these two graphs are not isomorphic. By Corollary 2, we already know they are not isomorphic to the infinite graphs obtained from LARG graphs on other Banach spaces.

Before we give this theorem, we recall some classical results which will be used in the proof. A theorem of Mazur and Ulam [6] states that a surjective isometry of real normed spaces is necessarily affine. Dilworth [5] extended this by showing that an approximate surjective isometry of Banach spaces may be uniformly approximated by a linear isometry. Recall from Lemma 1 that any isomorphism between well-connected LARG graphs satisfies $\lceil \|x - y\rceil \rceil = \lceil \theta(x) - \theta(y)\rceil \rceil$. The following corollary of Dilworth’s result states that any step-isometry gives rise to a surjective linear isometry that bounds the distance between pre-image and image.

Theorem 14. (*[4]*) *Suppose (f_n) and (g_n) are dense sequences in separable Banach spaces X and Y and $\lceil \|f_i - f_j\rceil \rceil = \lceil \|g_i - g_j\rceil \rceil$ for each i, j . We then have that there exists a surjective linear isometry, L , from X to Y and a constant $\kappa > 0$ such that $\|g_i - L(f_i)\| < \kappa$ for all i .*

The following classical result classifies the surjective isometries between Banach spaces of real continuous functions.

Theorem 15 (Banach–Stone). *If A and B are compact Hausdorff spaces and L is a surjective linear isometry from $C(A, \mathbb{R})$ to $C(B, \mathbb{R})$, then there exists a homeomorphism $\varphi: B \rightarrow A$ and a function $g \in C(B, \mathbb{R})$ with $|g(y)| = 1$ for all y such that*

$$L[f](y) = g(y)f(\varphi(y)) \text{ for all } f \in C(A, \mathbb{R}) \text{ and } y \in B.$$

We apply this in the special case that X and Y are both the closed unit interval, so that g is a constant function, either $g \equiv 1$ or $g \equiv -1$. This will lead to necessary conditions for two sets to give rise to isomorphic graphs, as stated in the following theorem.

Theorem 16. *If $\{f_n: n \in \mathbb{N}\}$ and $\{g_n: n \in \mathbb{N}\}$ are countable dense sets in $C([0, 1])$; and G_1 and G_2 are well-connected geometric graphs with these vertex sets, then there exists a homeomorphism ϕ from $[0, 1]$ to itself and a permutation of \mathbb{N} , $i \mapsto n_i$, such that one of the following holds:*

1. (*Order-preserving case*) *For each $x_0 \in [0, 1]$, if $f_i(x_0) > f_j(x_0)$ then $g_{n_i}(\phi(x_0)) \geq g_{n_j}(\phi(x_0))$;*
2. (*Order-reversing case*) *For each $x_0 \in [0, 1]$, if $f_i(x_0) > f_j(x_0)$ then $g_{n_i}(\phi(x_0)) \leq g_{n_j}(\phi(x_0))$.*

The idea of the proof (in the order-preserving case) is that we find a function f_k where $f_i - f_k$ and $f_j - f_k$ are non-negative, $\lfloor \|f_i - f_k\| \rfloor > \lfloor \|f_j - f_k\| \rfloor$ and $f_i - f_k$ and $f_j - f_k$ take values close to their maximum only on a small neighbourhood of x_0 . We use the facts that $g_a \approx f_a \circ \psi$ and $\lfloor \|g_{n_a} - g_{n_b}\| \rfloor = \lfloor \|f_a - f_b\| \rfloor$ for each a, b to deduce that g_{n_i} dominates g_{n_j} somewhere arbitrarily close to $\psi^{-1}(x_0)$.

Proof. Let $\{f_n\}$ and $\{g_n\}$ be as described and let G_1 and G_2 be isomorphic well-connected geometric graphs on these sets. We may re-label (g_n) so that for each n , f_n is identified with g_n in the isomorphism. By Lemma 1, such an isomorphism must almost surely satisfy: $\lceil \|f_i - f_j\| \rceil = \lceil \|g_i - g_j\| \rceil$ for each i, j . By Theorems 14 and 15, there exist a self-homeomorphism ψ of $[0, 1]$, a constant $\rho = \pm 1$ and a $\kappa > 0$ such that

$$\|\rho f_i \circ \psi - g_i\| < \kappa \text{ for all } i. \tag{5}$$

We deal with the order-preserving case, $\rho = 1$, although the case $\rho = -1$ is analogous. Let $\phi = \psi^{-1}$.

Let $x_0 \in [0, 1]$ and suppose that $f_i(x_0) > f_j(x_0)$. We let $\Delta = f_i(x_0) - f_j(x_0)$. For any $l, N \in \mathbb{N}$, define

$$h_{l,N}(x) = \max(f_i(x), f_j(x) + \Delta) - N - \frac{\Delta}{2} + l|x - x_0|.$$

Given $l \in \mathbb{N}$, let $N_l \in \mathbb{N}_{\geq 2}$ be chosen sufficiently large that

$$h_{l,N_l}(x) < \min(f_i(x), f_j(x)) - 2\kappa - \Delta \text{ for all } x \in [0, 1]. \tag{6}$$

We write h for h_{l,N_l} . We now have that, for each $x \in [0, 1]$,

$$\begin{aligned} 2\kappa + \Delta &\leq f_i(x) - h(x) \leq N + \frac{\Delta}{2} - l|x - x_0| \\ 2\kappa + \Delta &\leq f_j(x) - h(x) \leq N - \frac{\Delta}{2} - l|x - x_0|, \end{aligned}$$

where the upper bounds for $f_i - h$ and $f_j - h$ are equalities at $x = x_0$. In particular, we see $\|f_i - h\| = N + \frac{\Delta}{2}$ and $\|f_j - h\| = N - \frac{\Delta}{2}$, so $\lfloor \|f_i - h\| \rfloor \geq N$ while $\lfloor \|f_j - h\| \rfloor < N$.

Let $k \in \mathbb{N}$ be such that $\|f_k - h\| < \frac{\Delta}{2}$. This implies that $\lfloor \|f_i - f_k\| \rfloor \geq N$ and $\lfloor \|f_j - f_k\| \rfloor < N$. By Lemma 1, we have $\lfloor \|g_i - g_k\| \rfloor \geq N$ and $\lfloor \|g_j - g_k\| \rfloor < N$ also.

By (5), we have that

$$\begin{aligned} -\kappa &< f_n \circ \psi(y) - g_n(y) < \kappa \text{ for } n \in \mathbb{N} \text{ and } y \in [0, 1]; \text{ so} \\ -\kappa - \frac{\Delta}{2} &< h \circ \psi(y) - g_k(y) < \kappa + \frac{\Delta}{2} \text{ for } y \in [0, 1]. \end{aligned}$$

Combining this with (6), it follows that $g_i - g_k$ and $g_j - g_k$ are non-negative.

We also see that

$$\begin{aligned} g_i(y) - g_k(y) &\leq f_i(\psi(y)) - h(\psi(y)) + 2\kappa + \frac{\Delta}{2} \\ &\leq N + \Delta + 2\kappa - l|\psi(y) - x_0| \end{aligned}$$

Notice that $0 \leq g_i(y) - g_k(y) < N$ whenever $|\psi(y) - x_0| > (\Delta + 2\kappa)/l$. Since $\|g_i - g_k\| \geq N$, it follows that there exists y_l such that $|\psi(y_l) - x_0| \leq (\Delta + 2\kappa)/l$ with $g_i(y_l) - g_k(y_l) \geq N$. On the other hand, since $\|g_j - g_k\| < N$, we see $g_j(y_l) - g_k(y_l) < N$. In particular, we deduce that $g_i(y_l) > g_j(y_l)$. We now have a sequence of points y_l where $g_i(y_l) > g_j(y_l)$. We also have $|\psi(y_l) - x_0| \rightarrow 0$ as $l \rightarrow \infty$, which implies $y_l \rightarrow y_0 = \phi(x_0)$. By continuity of g_i and g_j , we deduce that $g_i(\phi(x_0)) \geq g_j(\phi(x_0))$. \square

Corollary 17. *The graphs G_{ICD} and G_{TD} are non-isomorphic.*

Proof. Let (f_n) be IC-dense and (g_n) be a transversely dense sequence of polynomials (such a sequence exists by Theorem 7). Let G_1 and G_2 be LARG graphs with vertices (f_n) and (g_n) , respectively, so that G_1 is almost surely isomorphic to G_{ICD} and G_2 is almost surely isomorphic to G_{TD} . Suppose for a contradiction that G_1 and G_2 are isomorphic.

We now apply Theorem 16, and assume without loss of generality that we are in the orientation-preserving case. Thus, there exists a permutation $i \mapsto n_i$ of \mathbb{N} and a homeomorphism ψ of $[0, 1]$ such that $f_i(x) > f_j(x)$ implies $g_{n_i}(\psi(x)) \geq g_{n_j}(\psi(x))$ for all x . By the infinite crossing dense condition, there must be two functions that intersect. That is, there exist $i \neq j$ so that $f_i(x) = f_j(x)$ for some $x \in (0, 1)$. By (IC4), there exists a strictly monotonic sequence $x_m \rightarrow x$ such that $f_i(x_m) = f_j(x_m)$ and between any pair x_m and x_{m+1} , there exist y_m and z_m such that $f_i(y_m) > f_j(y_m)$ and $f_i(z_m) < f_j(z_m)$. It follows that $g_{n_i}(\psi(y_m)) \geq g_{n_j}(\psi(y_m))$ and $g_{n_i}(\psi(z_m)) \geq g_{n_j}(\psi(z_m))$. In particular, by the Intermediate Value Theorem, there exists w_m between y_m and z_m such that $g_{n_i}(\psi(w_m)) = g_{n_j}(\psi(w_m))$. Hence, the two polynomials g_{n_i} and g_{n_j} agree at infinitely many points, and so are equal, which is a contradiction. \square

Acknowledgments

We thank the referee for a careful reading of the paper, and for making helpful suggestions regarding the presentation.

References

- [1] O. Angel, Y. Spinka, Geometric random graphs on circles, Preprint 2021. [2](#)
- [2] P. Balister, B. Bollobás, K. Gunderson, I. Leader, M. Walters, Random geometric graphs and isometries of normed spaces, *Trans. Amer. Math. Soc.* **370** (2018) 7361–7389. [2](#), [4](#)
- [3] A. Bonato, J. Janssen, Infinite random geometric graphs, *Annals of Combinatorics* **15** (2011) 597–617. [2](#), [3](#), [4](#)
- [4] A. Bonato, J. Janssen, A. Quas, Geometric random graphs and Rado sets in sequence spaces, *Eur. J. Combin.* **79** (2019) 1–14. [2](#), [3](#), [18](#)
- [5] S. Dilworth, Approximate isometries on finite-dimensional normed spaces, *Bull. London Math. Soc.* **31** (1999) 471–476. [3](#), [18](#)
- [6] S. Mazur, S. Ulam, Sur les transformations isométriques d’espaces vectoriels normés, *C. R. Acad. Sci. Paris* **194** (1932) 946–948. [18](#)
- [7] P. Mörters, Y. Peres, *Brownian Motion*, Cambridge University Press, Cambridge, 2010. [17](#)

AUTHORS

Anthony Bonato
Department of Mathematics
Ryerson University
Toronto, ON, CANADA
abonato@ryerson.ca

Jeannette Janssen
Department of Mathematics and Statistics
Dalhousie University
Halifax, NS, CANADA
jeannette.janssen@dal.ca

Anthony Quas
Department of Mathematics and Statistics
University of Victoria
Victoria, BC, CANADA
aquas@uvic.ca