

AN EXTENSION OF POLYNOMIAL INTEGRABILITY TO DUAL QUERMASINTEGRALS

VLAD YASKIN

ABSTRACT. A body K is called polynomially integrable if its parallel section function $V_{n-1}(K \cap \{\xi^\perp + t\xi\})$ is a polynomial of t (on its support) for every ξ . A complete characterization of such bodies was given recently. Here we obtain a generalization of these results in the setting of dual quermassintegrals. We also address the associated smoothness issues.

1. INTRODUCTION

Let K be a convex body in \mathbb{R}^n . The *parallel section function* of K in the direction $\xi \in S^{n-1}$ is defined by

$$A_{K,\xi}(t) = V_{n-1}(K \cap \{\xi^\perp + t\xi\}),$$

where $\xi^\perp = \{x \in \mathbb{R}^n : \langle x, \xi \rangle = 0\}$ and V_{n-1} is the $(n-1)$ -dimensional volume (Lebesgue measure).

Following Agranovsky [1], we say that a body K is *polynomially integrable* if

$$A_{K,\xi}(t) = \sum_{k=0}^N a_k(\xi) t^k,$$

for some integer N , all $\xi \in S^{n-1}$ and all t for which the set $K \cap \{\xi^\perp + t\xi\}$ is non-empty. Here, a_k are functions on the sphere.

It is not difficult to see that ellipsoids are polynomially integrable in odd dimensions. Agranovsky asked whether there are other polynomially integrable bodies. This question was answered in [1] and [10]. In even dimensions there are no polynomially integrable bodies, while in odd dimensions the only polynomially integrable bodies are ellipsoids. Note that these results were obtained under the C^∞ smoothness condition. Below we will address this issue.

Agranovsky's question is motivated by a problem about algebraically integrable domains that goes back to Newton [14]. The problem is to describe bounded domains with smooth boundaries for which the volume cut off by a plane from such a domain depends algebraically on the plane. The case of convex bodies in \mathbb{R}^2 was settled by Newton, who showed that such bodies do not exist. Arnold (see problems 1987-14, 1988-13, and 1990-27 in [2]) asked for extensions of Newton's result

2010 *Mathematics Subject Classification*. Primary 52A20, 33C55.

Key words and phrases. Convex bodies, dual quermassintegrals, spherical harmonics.

The author is supported in part by NSERC.

to other dimensions and non-convex domains. In even dimensions Arnold's question was answered by Vassiliev [16].

In this paper we deal with a generalization of the result in [10] to dual quermass-integrals. Before we introduce the latter, let us recall some facts from the classical Brunn-Minkowski theory. In the heart of this theory is the behavior of volume under the Minkowski addition. Intrinsic volumes are important concepts in this theory and arise as coefficients in the Steiner formula. If K is a convex body in \mathbb{R}^n and B_2^n is the unit Euclidean ball, then

$$V_n(K + \epsilon B_2^n) = \sum_{i=0}^n \binom{n}{i} W_i(K) \epsilon^i = \sum_{i=0}^n \kappa_{n-i} V_i(K) \epsilon^{n-i},$$

where the addition is the Minkowski addition and κ_i is the volume of the unit Euclidean ball in \mathbb{R}^i . The coefficients $W_i(K)$ and $V_i(K)$ are known as quermassintegrals and intrinsic volumes, correspondingly. Clearly, $V_i(K)$ is the same as $W_{n-i}(K)$, up to a constant factor. The standard reference for the Brunn-Minkowski theory is the book of Schneider [15].

The dual Brunn-Minkowski theory was initiated by Lutwak [11]. In this theory the Minkowski addition of convex bodies is replaced by the radial addition of star bodies. It served as a foundation for many developments in modern convex geometry, especially in problems related to sections of convex or star bodies. One of the famous examples is the Busemann-Petty problem, where one of the main ingredients were Lutwak's intersection bodies introduced within the dual Brunn-Minkowski theory; see [12]. For the history of this problem the reader is referred to [9].

It turns out that there are many analogies between the Brunn-Minkowski theory and its dual counterpart. For example, there is a version of the Steiner formula in the dual setting. Let K be a star body in \mathbb{R}^n and let B_2^n be the unit Euclidean ball. Consider the star body $K \tilde{+} \epsilon B_2^n$, where $\tilde{+}$ is the radial addition. Then

$$V_n(K \tilde{+} \epsilon B_2^n) = \sum_{i=0}^n \binom{n}{i} \tilde{W}_i(K) \epsilon^i = \sum_{i=0}^n \binom{n}{i} \tilde{V}_i(K) \epsilon^{n-i},$$

where the coefficients $\tilde{W}_i(K)$ and $\tilde{V}_i(K)$ are called the dual quermassintegrals and dual volumes, correspondingly. Note that they depend on the choice of the origin. Explicitly, they can be written as follows:

$$\tilde{V}_i(K) = \tilde{W}_{n-i}(K) = \frac{1}{n} \int_{S^{n-1}} \rho_K^i(\theta) d\theta,$$

where the integration is with respect to the spherical Lebesgue measure and ρ_K is the radial function of K .

Analogous to the Kubota integral recursion formula, which gives the i -th intrinsic volume as the average volume of projections of the body onto i -dimensional subspaces, is its dual version, which some authors call the dual Kubota integral recursion formula,

$$\tilde{V}_i(K) = \frac{\kappa_n}{\kappa_i} \int_{G(n,i)} V_i(K \cap H) dH, \quad (1)$$

where the integration is with respect to the Haar probability measure on the Grassmanian $G(n, i)$. This formula, as well as a brief introduction into the dual Brunn-Minkowski theory, can be found in [4].

Dual quermassintegrals are particular cases of dual mixed volumes. For some results about applications and characterizations of dual mixed volumes, the reader is referred to [3], [7], [8], [13].

Now we are ready to state our results. Let K be a convex body in \mathbb{R}^n that contains the origin in its interior and let m be an integer, $1 \leq m \leq n-1$. Consider the function $A_{K,m,\xi}(t)$ that gives the m th dual volume of the section $K \cap \{\xi^\perp + t\xi\}$ with respect to the point $t\xi$, i.e.,

$$\begin{aligned} A_{K,m,\xi}(t) &= \tilde{V}_m(K \cap \{\xi^\perp + t\xi\}) = \frac{1}{n-1} \int_{S^{n-1} \cap \xi^\perp} \rho_{K-t\xi}^m(\theta) d\theta \\ &= \frac{\kappa_{n-1}}{\kappa_m} \int_{G(\xi^\perp, m)} V_m((K - t\xi) \cap H) dH, \end{aligned} \quad (2)$$

where $G(\xi^\perp, m)$ denotes the Grassmanian of m -dimensional subspaces of ξ^\perp . Note that $t\xi$ is not an interior point of the section $K \cap \{\xi^\perp + t\xi\}$ when $t \geq \rho_K(\xi)$. The latter is the radial function in the direction ξ ; see the next section for its definition. One can also use representation (2) as a definition of $A_{K,m,\xi}(t)$ for all t , but we do not really need this.

We say that K has a polynomial m th dual section function if there is a positive integer N such that

$$A_{K,m,\xi}(t) = \sum_{k=0}^N a_k(\xi) t^k$$

for all $\rho_K(-\xi) \leq t \leq \rho_K(\xi)$, where a_k are continuous functions on the sphere.

In this paper we show that for even m the only convex bodies with this property are ellipsoids, while for odd m such bodies do not exist. Note that the results of [10] can be obtained as a particular case of our results when $m = n - 1$. Since we do not impose any smoothness assumptions on the bodies, it means that the C^∞ assumption used in [10] can be dropped. The proof here is based on spherical harmonics, instead of the Fourier transform as in [10].

2. DEFINITIONS AND AUXILIARY FACTS

Let K be a compact set in \mathbb{R}^n that is star-shaped about the origin o ; the latter means that for every point $x \in K$ the closed segment $[o, x]$ is also contained in K . The *radial function* of K is defined by

$$\rho_K(\xi) = \max\{a \geq 0 : a\xi \in K\}, \quad \xi \in S^{n-1}.$$

We say that K is a *star body* if it is compact, star-shaped about the origin and its radial function is a positive continuous function on S^{n-1} .

The *Minkowski functional* of K is defined by

$$\|x\|_K = \min\{a \geq 0 : x \in aK\}, \quad x \in \mathbb{R}^n.$$

Clearly, $\|\cdot\|_K$ is a 1-homogeneous function on \mathbb{R}^n and $\|\xi\|_K = \rho_K(\xi)^{-1}$ for $\xi \in S^{n-1}$.

If K and L are star bodies, and α and β are positive numbers, then the radial sum $\alpha K \tilde{+} \beta L$, which we saw in the introduction, is defined as the star body with the radial function

$$\rho_{\alpha K \tilde{+} \beta L} = \alpha \rho_K + \beta \rho_L.$$

Let h be an integrable function on \mathbb{R} that is C^∞ -smooth in a neighborhood of the origin. If $q \in \mathbb{C}$, $-1 < \Re q < 0$, then the *fractional derivative* of the function h of order q at zero is defined by

$$h^{(q)}(0) = \frac{1}{\Gamma(-q)} \int_0^\infty t^{-1-q} h(t) dt.$$

For other values of q it is defined using analytic continuation (with respect to q). In particular, if q is not an integer and $-1 < \Re q < m$ for some integer m , then

$$\begin{aligned} h^{(q)}(0) &= \frac{1}{\Gamma(-q)} \int_0^1 t^{-1-q} \left(h(t) - h(0) - \dots - h^{(m-1)}(0) \frac{t^{m-1}}{(m-1)!} \right) dt \\ &\quad + \frac{1}{\Gamma(-q)} \int_1^\infty t^{-1-q} h(t) dt + \frac{1}{\Gamma(-q)} \sum_{k=0}^{m-1} \frac{h^{(k)}(0)}{k!(k-q)}; \end{aligned}$$

see e.g., [9].

If $k \geq 0$ is an integer, then passing to the limit in the previous expression we get ordinary derivatives (up to a sign),

$$h^{(k)}(0) = (-1)^k \frac{d^k}{dt^k} h(t) \Big|_{t=0}.$$

In this paper we will extensively use spherical harmonics; see [6] for a comprehensive introduction into this subject. Recall that a *spherical harmonic* of degree m and dimension n is the restriction to S^{n-1} of a harmonic homogeneous polynomial of degree m on \mathbb{R}^n . Given $f \in L^2(S^{n-1})$, one can associate with it its spherical harmonic expansion $\sum_{m=0}^\infty H_m$, where H_m is a spherical harmonic of order m .

The following is the well-known Funk-Hecke theorem. If Φ is an integrable function on $[-1, 1]$ and H_m is a spherical harmonic of order m , then

$$\int_{S^{n-1}} \Phi(\langle \xi, \theta \rangle) H_m(\theta) d\theta = \lambda_m(\Phi) H_m(\xi),$$

where

$$\lambda_m(\Phi) = \int_{-1}^1 \Phi(t) P_m^n(t) (1-t^2)^{(n-3)/2} dt$$

and P_m^n is the Legendre polynomial of degree m and dimension n .

3. MAIN RESULTS

Lemma 1. *Let E be an ellipsoid that contains the origin in its interior. Then E has a polynomial m th dual section function for every even integer m , $2 \leq m \leq n-1$.*

Proof. By formula (2), we have

$$\begin{aligned} A_{E,m,\xi}(t) &= \frac{\kappa_{n-1}}{\kappa_m} \int_{G(\xi^\perp, m)} V_m((E - t\xi) \cap H) dH \\ &= \frac{\kappa_{n-1}}{\kappa_m} \int_{G(\xi^\perp, m)} V_m(E \cap (H \vee \xi) \cap \{\xi^\perp + t\xi\}) dH, \end{aligned}$$

where $H \vee \xi = \text{span}(H, \xi)$.

We now use the fact that $E \cap (H \vee \xi)$ is an ellipsoid, and therefore polynomially integrable in $H \vee \xi$ if m is even (since $H \vee \xi$ is $(m+1)$ -dimensional). Thus

$$V_m(E \cap (H \vee \xi) \cap \{\xi^\perp + t\xi\}) = \sum_{k=0}^m a_k(H, \xi) t^k,$$

for all $\rho_K(-\xi) \leq t \leq \rho_K(\xi)$.

Note that $a_k(H, \xi)$ are continuous functions of H and ξ , and thus denoting

$$a_k(\xi) = \frac{\kappa_{n-1}}{\kappa_m} \int_{G(\xi^\perp, m)} a_k(H, \xi) dH,$$

we get

$$A_{E,m,\xi}(t) = \sum_{k=0}^m a_k(\xi) t^k,$$

for all $\rho_K(-\xi) \leq t \leq \rho_K(\xi)$. □

Theorem 1. *There are no convex bodies with polynomial m th dual section functions if m is odd.*

Proof. Let K be a convex body in \mathbb{R}^n containing the origin as an interior point. For a fixed t , $|t| < \min_{\xi \in S^{n-1}} \rho_K(\xi)$, consider the integral

$$\int_{S^{n-1}} A_{K,m,\xi}(t) d\xi = \frac{\kappa_{n-1}}{\kappa_m} \int_{S^{n-1}} \int_{G(\xi^\perp, m)} V_m(K \cap (H \vee \xi) \cap \{\xi^\perp + t\xi\}) dH d\xi,$$

where $H \vee \xi = \text{span}(H, \xi)$, as before.

Observe that

$$\begin{aligned} V_m(K \cap (H \vee \xi) \cap \{\xi^\perp + t\xi\}) &= \lim_{\epsilon \rightarrow 0^+} \frac{\epsilon}{2} \int_{K \cap (H \vee \xi)} |\langle x, \xi \rangle - t|^{-1+\epsilon} dx \\ &= \lim_{\epsilon \rightarrow 0^+} \frac{\epsilon}{2} \int_{S^{n-1} \cap (H \vee \xi)} \int_0^{\rho_K(\theta)} r^m |r \langle \theta, \xi \rangle - t|^{-1+\epsilon} dr d\theta \\ &= \lim_{\epsilon \rightarrow 0^+} \frac{\epsilon}{2} \int_{S^{n-1} \cap H} \int_{-1}^1 (1-z^2)^{(m-2)/2} \int_0^{\rho_K(z\xi + \sqrt{1-z^2}\eta)} r^m |rz - t|^{-1+\epsilon} dr dz d\eta. \end{aligned}$$

Note that for a function f , integrable on the corresponding sub-spheres, one has

$$\int_{G(\xi^\perp, m)} \int_{S^{n-1} \cap H} f(\eta) d\eta dH = \frac{m\kappa_m}{(n-1)\kappa_{n-1}} \int_{S^{n-1} \cap \xi^\perp} f(\eta) d\eta.$$

Therefore,

$$\begin{aligned}
& \int_{S^{n-1}} A_{K,m,\xi}(t) d\xi \\
&= \lim_{\epsilon \rightarrow 0^+} \frac{\epsilon m}{2(n-1)} \int_{S^{n-1}} \int_{S^{n-1} \cap \xi^\perp} \int_{-1}^1 (1-z^2)^{(m-2)/2} \int_0^{\rho_K(z\xi + \sqrt{1-z^2}\eta)} r^m |rz - t|^{-1+\epsilon} dr dz d\eta d\xi \\
&= \lim_{\epsilon \rightarrow 0^+} \frac{\epsilon m}{2(n-1)} \int_{S^{n-1}} \int_{S^{n-1}} (1 - \langle \theta, \xi \rangle^2)^{(m-n+1)/2} \int_0^{\rho_K(\theta)} r^m |r\langle \theta, \xi \rangle - t|^{-1+\epsilon} dr d\theta d\xi \\
&= \lim_{\epsilon \rightarrow 0^+} \frac{\epsilon m}{2(n-1)} \int_{S^{n-1}} \int_0^{\rho_K(\theta)} r^m \int_{S^{n-1}} (1 - \langle \theta, \xi \rangle^2)^{(m-n+1)/2} |r\langle \theta, \xi \rangle - t|^{-1+\epsilon} d\xi dr d\theta.
\end{aligned}$$

Let us now compute

$$\begin{aligned}
& \lim_{\epsilon \rightarrow 0^+} \frac{\epsilon}{2} \int_{S^{n-1}} (1 - \langle \theta, \xi \rangle^2)^{(m-n+1)/2} |r\langle \theta, \xi \rangle - t|^{-1+\epsilon} d\xi \\
&= \lim_{\epsilon \rightarrow 0^+} \frac{\epsilon}{2} r^{-1+\epsilon} \int_{S^{n-1} \cap \theta^\perp} \int_{-1}^1 (1-z^2)^{(m-2)/2} \left| z - \frac{t}{r} \right|^{-1+\epsilon} dz d\eta \\
&= \begin{cases} r^{-1} |S^{n-2}| \left(1 - \frac{t^2}{r^2}\right)^{(m-2)/2}, & \text{if } t \leq r, \\ 0, & \text{if } t > r. \end{cases}
\end{aligned}$$

Thus,

$$\begin{aligned}
\int_{S^{n-1}} A_{K,m,\xi}(t) d\xi &= m\kappa_{n-1} \int_{S^{n-1}} \int_t^{\rho_K(\theta)} r^{m-1} \left(1 - \frac{t^2}{r^2}\right)^{(m-2)/2} dr d\theta \\
&= m\kappa_{n-1} \int_{S^{n-1}} \int_t^{\rho_K(\theta)} r (r^2 - t^2)^{(m-2)/2} dr d\theta \\
&= \kappa_{n-1} \int_{S^{n-1}} (\rho_K^2(\theta) - t^2)^{m/2} d\theta.
\end{aligned}$$

If K had a polynomial m th dual section function, then $\int_{S^{n-1}} A_{K,m,\xi}(t) d\xi$ would be a polynomial of t , for $|t| < \min_{\xi \in S^{n-1}} \rho_K(\xi)$. But, clearly, $\int_{S^{n-1}} (\rho_K^2(\theta) - t^2)^{m/2} d\theta$ is not a polynomial, since m is odd. \square

Theorem 2. *For even m , the only convex bodies with a polynomial m th dual section function are ellipsoids.*

Proof. Let K be convex body in \mathbb{R}^n that contains the origin in its interior. Assume that K has a polynomial m th dual section function for some even m .

For a fixed ξ , let us compute fractional derivatives of $A_{K,m,\xi}(t)$ at zero. Assume that $-1 < \Re q < 0$. Then

$$\begin{aligned}
A_{K,m,\xi}^{(q)}(0) &= \frac{1}{\Gamma(-q)} \int_0^\infty t^{-1-q} A_{K,m,\xi}(t) dt \\
&= \frac{1}{2\Gamma(-q)} \int_{-\infty}^\infty |t|^{-1-q} (1 + \operatorname{sgn} t) A_{K,\xi,m}(t) dt
\end{aligned}$$

$$\begin{aligned}
 &= \frac{\kappa_{n-1}}{\kappa_m} \frac{1}{2\Gamma(-q)} \int_{G(\xi^\perp, m)} \int_{-\infty}^{\infty} |t|^{-1-q} (1 + \operatorname{sgn} t) V_m(K \cap (H \vee \xi) \cap \{\xi^\perp + t\xi\}) dt dH \\
 &= \frac{\kappa_{n-1}}{\kappa_m} \frac{1}{2\Gamma(-q)} \int_{G(\xi^\perp, m)} \int_{K \cap (H \vee \xi)} |\langle x, \xi \rangle|^{-1-q} (1 + \operatorname{sgn} \langle x, \xi \rangle) dx dH \\
 &= \frac{\kappa_{n-1}}{\kappa_m} \frac{1}{2\Gamma(-q)} \int_{G(\xi^\perp, m)} \int_{S^{n-1} \cap (H \vee \xi)} \int_0^{\rho_K(\theta)} r^{m-1-q} dr |\langle \theta, \xi \rangle|^{-1-q} (1 + \operatorname{sgn} \langle \theta, \xi \rangle) d\theta dH \\
 &= \frac{\kappa_{n-1}}{\kappa_m} \frac{1}{2\Gamma(-q)(m-q)} \int_{G(\xi^\perp, m)} \int_{S^{n-1} \cap (H \vee \xi)} \rho_K^{m-q}(\theta) |\langle \theta, \xi \rangle|^{-1-q} (1 + \operatorname{sgn} \langle \theta, \xi \rangle) d\theta dH \\
 &= \frac{\kappa_{n-1}}{\kappa_m} \frac{1}{2\Gamma(-q)(m-q)} \int_{G(\xi^\perp, m)} \int_{S^{n-1} \cap H} \int_{-1}^1 \rho_K^{m-q}(z\xi + \sqrt{1-z^2}\eta) |z|^{-1-q} (1 + \operatorname{sgn} z) \\
 &\quad \times (1-z^2)^{(m-2)/2} dz d\eta dH \\
 &= \frac{m}{2\Gamma(-q)(m-q)(n-1)} \int_{S^{n-1} \cap \xi^\perp} \int_{-1}^1 \rho_K^{m-q}(z\xi + \sqrt{1-z^2}\eta) |z|^{-1-q} (1 + \operatorname{sgn} z) \\
 &\quad \times (1-z^2)^{(m-2)/2} dz d\eta \\
 &= \frac{m}{2\Gamma(-q)(m-q)(n-1)} \int_{S^{n-1}} \rho_K^{m-q}(\theta) |\langle \theta, \xi \rangle|^{-1-q} (1 + \operatorname{sgn} \langle \theta, \xi \rangle) \\
 &\quad \times (1 - \langle \theta, \xi \rangle^2)^{(m-n+1)/2} d\theta.
 \end{aligned}$$

For $-1 < \Re q < 0$ and a function $f \in C(S^{n-1})$ we will be interested in the operator I_q defined by

$$I_q f(\xi) = \frac{1}{2\Gamma(-q)} \int_{S^{n-1}} |\langle \theta, \xi \rangle|^{-1-q} (1 + \operatorname{sgn} \langle \theta, \xi \rangle) (1 - \langle \theta, \xi \rangle^2)^{(m-n+1)/2} f(\theta) d\theta.$$

If H_l is a spherical harmonic of degree l , then by Funk-Hecke's theorem,

$$I_q H_l = \lambda_l(q) H_l,$$

where

$$\lambda_l(q) = \frac{1}{\Gamma(-q)} \int_0^1 P_l^n(z) z^{-1-q} (1-z^2)^{(m-2)/2} dz,$$

where P_l^n is the Legendre polynomial of degree l and dimension n .

Such multipliers were studied before, e.g., by Goodey [5]. But he was interested in the case when $-1-q$ is a non-negative integer, while our main interest is when $-1-q$ is a negative integer.

Note that $\lambda_l(q)$ is the fractional derivative of order q (with $-1 < \Re q < 0$) at zero of the function $P_l^n(z)(1-z^2)^{(m-2)/2}$. Thus $\lambda_l(q)$ can be extended to all q using analytic continuation. So, if q is equal to an integer $k \geq 0$, then

$$\lambda_l(k) = (-1)^k \frac{d^k}{dz^k} (P_l^n(z)(1-z^2)^{(m-2)/2}) \Big|_{z=0}.$$

Note that $P_l^n(z)(1-z^2)^{(m-2)/2}$ is a polynomial of degree $l+m-2$ and thus $\lambda_l(k) = 0$ if $m < k - l + 2$. In addition, $\lambda_l(k) = 0$ if l and k are of different parity. Let us now show that $\lambda_l(k) \neq 0$ for all $m \geq k - l + 2$ if $k - l$ is even.

We have

$$\begin{aligned}\lambda_l(k) &= (-1)^k \frac{d^k}{dz^k} (P_l^n(z)(1-z^2)^{(m-2)/2}) \Big|_{z=0} \\ &= (-1)^k \sum_{j=0}^k \frac{d^{k-j}}{dz^{k-j}} P_l^n(z) \Big|_{z=0} \frac{d^j}{dz^j} (1-z^2)^{(m-2)/2} \Big|_{z=0}.\end{aligned}$$

Since

$$(1-z^2)^{(m-2)/2} = \sum_{i=0}^{(m-2)/2} (-1)^i \binom{(m-2)/2}{i} z^{2i},$$

it follows that

$$\frac{d^j}{dz^j} (1-z^2)^{(m-2)/2} \Big|_{z=0} = (-1)^{j/2} \binom{(m-2)/2}{j/2} j!,$$

if j is even and $j \leq m-2$, and this derivative is zero if j is odd.

Also, by [6, Lemma 3.3.9], we have

$$\frac{d^{k-j}}{dz^{k-j}} P_l^n(z) = c_{n,l,j} P_{l-k+j}^{n+2k-2j}(z),$$

if $k-j \leq l$ and it is zero otherwise. Here $c_{n,l,j}$ is a positive constant.

Now use [6, Lemma 3.3.8],

$$P_{l-k+j}^{n+2k-2j}(0) = (-1)^{(l-k+j)/2} b_{n,l,j},$$

if $l-k+j$ is a non-negative even number, and it is zero otherwise. Here $b_{n,l,j}$ is a positive constant.

Summarizing, we have

$$\lambda_l(k) = (-1)^l \sum_{\substack{j=0 \\ j \text{ even}}}^k \binom{(m-2)/2}{j/2} j! c_{n,l,j} b_{n,l,j}.$$

Note that inside the summation all the terms are non-negative, and there is at least one term that is strictly positive, since there is j that satisfies $l-k \leq j \leq m-2$. Thus, we have proved that $\lambda_l(k) \neq 0$ if and only if $k-l$ is even and $m \geq k-l+2$.

Note that we extended $\lambda_l(q)$ to all non-negative values of q , but we did not do this for $I_q f$. For $I_q f$ this is also possible, but we would need f to be infinitely smooth (which we want to avoid). Instead of extending $I_q f$, we will proceed as follows.

Let H_l be a spherical harmonic of order l . First consider the case $-1 < \Re q < 0$. Then

$$\begin{aligned}\int_{S^{n-1}} A_{K,m,\xi}^{(q)}(0) H_l(\xi) d\xi &= \frac{m}{(m-q)(n-1)} \int_{S^{n-1}} I_q(\rho_K^{m-q})(\xi) H_l(\xi) d\xi \\ &= \frac{m}{(m-q)(n-1)} \int_{S^{n-1}} \rho_K^{m-q}(\xi) I_q(H_l)(\xi) d\xi \\ &= \frac{m}{(m-q)(n-1)} \lambda_l(q) \int_{S^{n-1}} \rho_K^{m-q}(\xi) H_l(\xi) d\xi.\end{aligned}$$

The left and right-hand sides of the equality are analytic functions of q in the domain $\{q \in \mathbb{C} : \Re(q) > -1 \text{ and } q \neq m\}$. Thus they are equal when q is an integer $k > \max\{m, N\}$. Since $A_{K,m,\xi}$ is a polynomial of t for every ξ , we have

$$A_{K,m,\xi}^{(k)}(0) = 0,$$

for all $\xi \in S^{n-1}$, and so

$$\lambda_l(k) \int_{S^{n-1}} \rho_K^{m-k}(\xi) H_l(\xi) d\xi = 0.$$

If $l \geq k - m + 2$ and $k - l$ is even, then $\lambda_l(k) \neq 0$, and thus ρ_K^{m-k} does not have spherical harmonics of order l , where $l = k \pmod{2}$, in its expansion. Thus if k is even, then $\rho_K^{m-k} + \rho_{-K}^{m-k}$ is a combination of spherical harmonics of even degrees less than $k - m + 2$. If k is odd, then $\rho_K^{m-k} - \rho_{-K}^{m-k}$ is a combination of spherical harmonics of odd degrees less than $k - m + 2$.

Note that spherical harmonics are restrictions of homogeneous harmonic polynomials to the sphere. Thus, as a function on \mathbb{R}^n , we have, when k is even,

$$\|x\|_K^{k-m} + \|-x\|_K^{k-m} = \sum_{\substack{l=0 \\ l \text{ even}}}^{k-m} H_l\left(\frac{x}{|x|}\right) |x|^{k-m} = \sum_{l=0}^{k-m} H_l(x) |x|^{k-m-l}.$$

Since $k - m - l$ is even, the right-hand side is a polynomial on \mathbb{R}^n .

Similarly, if k is odd,

$$\|x\|_K^{k-m} - \|-x\|_K^{k-m} = \sum_{\substack{l=0 \\ l \text{ odd}}}^{k-m} H_l\left(\frac{x}{|x|}\right) |x|^{k-m} = \sum_{l=0}^{k-m} H_l(x) |x|^{k-m-l}$$

is also a polynomial on \mathbb{R}^n .

Now we proceed as in the proof of Theorem 3.7 in [10]. It was shown there that if K is a convex body such that $\|x\|_K^l + \|-x\|_K^l$ is a polynomial for even l and $\|x\|_K^l - \|-x\|_K^l$ is a polynomial for odd l , then K is an ellipsoid. □

4. FINAL REMARKS

One can also consider the function $P_{K,m,\xi}(t)$ that gives the m th intrinsic volume of the section $K \cap \{\xi^\perp + t\xi\}$,

$$P_{K,m,\xi}(t) = V_m(K \cap \{\xi^\perp + t\xi\}).$$

If m is even and K is an ellipsoid, it is easy to see that $P_{K,m,\xi}(t)$ is a polynomial of t of degree m (on its support), for every ξ . The argument goes as in Lemma 1. Using the Kubota integral recursion formula, we have

$$P_{K,m,\xi}(t) = \frac{\kappa_{n-1}}{\kappa_m} \int_{G(\xi^\perp, m)} V_m((K|(H \vee \xi)) \cap \{\xi^\perp + t\xi\}) dH.$$

We now use the fact that $K|(H \vee \xi)$ is an ellipsoid, and therefore polynomially integrable in $H \vee \xi$ if m is even (since $H \vee \xi$ is $(m+1)$ -dimensional).

Question. Are ellipsoids the only convex bodies in \mathbb{R}^n for which $P_{K,m,\xi}(t)$ is a polynomial of t for every ξ ?

REFERENCES

- [1] M. AGRANOVSKY, *On polynomially integrable domains in Euclidean spaces*, In: Agranovsky M., Golberg A., Jacobzon F., Shoikhet D., Zalcman L. (eds) *Complex Analysis and Dynamical Systems*. Trends in Mathematics. Birkhäuser, Cham (2018).
- [2] V. I. ARNOLD, *Arnold's problems*, 2nd edition, Springer-Verlag, Berlin (2004).
- [3] P. DULIO, R. J. GARDNER, C. PERI, *Characterizing the dual mixed volume via additive functionals*, *Indiana Univ. Math. J.* **65** (2016), no. 1, 69–91.
- [4] R. J. GARDNER, *Geometric Tomography*. Second edition. Encyclopedia of Mathematics and its Applications, **58**. Cambridge University Press, Cambridge, 2006.
- [5] P. GOODEY, *Minkowski sums of projections of convex bodies*, *Mathematika* **45** (1998), no. 2, 253–268.
- [6] H. GROEMER, *Geometric Applications of Fourier Series and Spherical Harmonics*, Cambridge University Press, New York, 1996.
- [7] C. JIMÉNEZ, I. VILLANUEVA, *Characterization of dual mixed volumes via polymeasures*, *J. Math. Anal. Appl.* **426** (2015), no. 2, 688–699.
- [8] D. KLAIN, *Star valuations and dual mixed volumes*, *Adv. Math.* **121** (1996), 80–108.
- [9] A. KOLDOBSKY, *Fourier Analysis in Convex Geometry*, American Mathematical Society, Providence RI, 2005.
- [10] A. KOLDOBSKY, A. MERKURJEV, AND V. YASKIN, *On polynomially integrable convex bodies*, *Adv. Math.* **320**, (2017), 876–886.
- [11] E. LUTWAK, *Dual mixed volumes*, *Pacific J. Math.* **58** (1975), 531–538.
- [12] E. LUTWAK, *Intersection bodies and dual mixed volumes*, *Advances in Math.* **71** (1988), 232–261.
- [13] E. MILMAN, *Dual mixed volumes and the slicing problem*, *Advances in Math.* **207** (2006), 566–598.
- [14] I. NEWTON, *Philosophiae Naturalis Principia Mathematica*, London, 1687.
- [15] R. SCHNEIDER, *Convex bodies: the Brunn-Minkowski theory*, Second expanded edition. Encyclopedia of Mathematics and its Applications, **151**. Cambridge University Press, Cambridge, 2014.
- [16] V. A. VASSILIEV, *Newton's lemma XXVIII on integrable ovals in higher dimensions and reflection groups*, *Bull. Lond. Math. Soc.* **47** (2015), 290–300.

DEPARTMENT OF MATHEMATICAL AND STATISTICAL SCIENCES, UNIVERSITY OF ALBERTA, EDMONTON, ALBERTA T6G 2G1, CANADA

E-mail address: yaskin@ualberta.ca