

The inverses of tails of the Riemann zeta function

Donggyun Kim and Kyunghwan Song*

Department of Mathematics, Korea University, Seoul 02841, Republic of Korea

Abstract

We present some bounds of the inverses of tails of the Riemann zeta function on $0 < s < 1$ and compute the integer parts of the inverses of tails of the Riemann zeta function for $s = \frac{1}{2}, \frac{1}{3}$ and $\frac{1}{4}$.

Key words— Riemann zeta function; tails of Riemann zeta function; inverses of tails of the Riemann zeta function

Mathematics Subject Classification 2010: 11M06, 11B83

1 Introduction

The Riemann zeta function $\zeta(s)$ in the real variable s was introduced by L. Euler [2] in connection with questions about the distribution of prime numbers. Later B. Riemann [6] derived deeper results about a dual correspondence between the distribution of prime numbers and the complex zeros of $\zeta(s)$ in the complex variable s . In these developments, he asserted that all the non-trivial zeros of $\zeta(s)$ are on the line $\operatorname{Re}(s) = \frac{1}{2}$, and this has been one of the most important unsolved problems in mathematics, called the Riemann hypothesis. A vast amount of research on calculation of $\zeta(s)$ for the line $\operatorname{Re}(s) = \frac{1}{2}$ which is called the critical line, and on the strip $0 < \operatorname{Re}(s) < 1$ which is called the critical strip, has been conducted using various methods [1].

The *Riemann zeta function* and a *tail of the Riemann zeta function from n* for an integer $n \geq 1$ are defined, respectively, by for $\operatorname{Re}(s) > 1$,

$$\zeta(s) = \sum_{k=1}^{\infty} \frac{1}{k^s} \quad \text{and} \quad \zeta_n(s) = \sum_{k=n}^{\infty} \frac{1}{k^s}$$

and for $0 < \operatorname{Re}(s) < 1$,

$$\zeta(s) = \frac{1}{1-2^{1-s}} \sum_{k=1}^{\infty} \frac{(-1)^{k+1}}{k^s} \quad \text{and} \quad \zeta_n(s) = \frac{1}{1-2^{1-s}} \sum_{k=n}^{\infty} \frac{(-1)^{k+1}}{k^s}.$$

To understand the values of $\zeta(s)$, it would be helpful to understand the values of tails of $\zeta(s)$, for example, the integer parts of their inverses, $[\zeta_n(s)^{-1}]$, where $[x]$ denotes the greatest integer that is less than or equal to x .

Some values of $[\zeta_n(s)^{-1}]$ for small positive integers s are known recently. L. Xin [7] showed that for $s = 2$ and 3,

$$[\zeta_n(2)^{-1}] = n - 1 \quad \text{and} \quad [\zeta_n(3)^{-1}] = 2n(n - 1).$$

For $s = 4$, L. Xin and L. Xiaoxue [8] showed that

$$[\zeta_n(4)^{-1}] = 3n^3 - 5n^2 + 4n - 1 + \left\lceil \frac{(2n+1)(n-1)}{4} \right\rceil$$

for any integer $n \geq 2$ and H. Xu [9] showed that for $s = 5$,

$$[\zeta_n(5)^{-1}] = 4n^4 - 8n^3 + 9n^2 - 5n + \left\lceil \frac{(n+1)(n-2)}{3} \right\rceil$$

for any integer $n \geq 4$. W. Hwang and K. Song [3] provided an alternative proof of the case when $s = 5$ and a formula when $s = 6$ as follows. For an integer n , write n_{48} for the remainder when n is divided by 48, then

$$[\zeta_n(6)^{-1}] = \begin{cases} 5n^5 - \frac{25}{2}n^4 + \frac{75}{4}n^3 - \frac{125}{8}n^2 + \frac{185}{48}n - \frac{5n_{48}}{48} - \left\lceil \frac{35-5n_{48}}{48} \right\rceil, & \text{if } n \text{ is even,} \\ 5n^5 - \frac{25}{2}n^4 + \frac{75}{4}n^3 - \frac{125}{8}n^2 + \frac{185}{48}n - \frac{5n_{48}+18}{48} - \left\lceil \frac{17-5n_{48}}{48} \right\rceil, & \text{if } n \text{ is odd} \end{cases}$$

for any integer $n \geq 829$. For the integer s greater than 6, no such a formula is known.

There are other interesting results related to this theme such as bounds of $\zeta(3)$ in greater precision in [4] and [5].

We study the inverses of tails of the Riemann zeta function, $\zeta_n(s)^{-1}$, for s on the critical strip $0 < s < 1$. The following notation is needed to explain our results.

Definition 1.1. For any positive integer n and real number s with $0 < s < 1$, we define

$$A_{n,s} = \left(\frac{1}{n^s} - \frac{1}{(n+1)^s} \right) + \left(\frac{1}{(n+2)^s} - \frac{1}{(n+3)^s} \right) + \cdots$$

and

$$B_{n,s} = \left(-\frac{1}{n^s} + \frac{1}{(n+1)^s} \right) + \left(-\frac{1}{(n+2)^s} + \frac{1}{(n+3)^s} \right) + \cdots.$$

Now the tail of the Riemann zeta function can be written as, for $0 < s < 1$,

$$\zeta_n(s) = \begin{cases} -\frac{1}{1-2^{1-s}} A_{n,s}, & \text{if } n \text{ is even,} \\ -\frac{1}{1-2^{1-s}} B_{n,s}, & \text{if } n \text{ is odd.} \end{cases} \quad (1.1)$$

In this paper, we present the bounds of $A_{n,s}^{-1}$ and $B_{n,s}^{-1}$ hence the bounds of the inverses of tails of the Riemann zeta function, $\zeta_n(s)^{-1}$, for $0 < s < 1$ in Section 2.1, and compute the values $[A_{n,s}^{-1}]$ and $[B_{n,s}^{-1}]$ hence the values of the inverses of tails of the Riemann zeta function, $[\frac{1}{1-2^{1-s}} \zeta_n(s)^{-1}]$, for $s = \frac{1}{2}$, $\frac{1}{3}$ and $\frac{1}{4}$ in Section 2.2.

2 Main Results

2.1 The bounds of the inverses of $\zeta_n(s)$ for $0 < s < 1$

In this section, we present the bounds of $A_{n,s}^{-1}$ and $B_{n,s}^{-1}$ in Definition 1.1, hence the bounds of the inverses of tails of the Riemann zeta function, $\zeta_n(s)^{-1}$, for $0 < s < 1$.

Proposition 2.1. *Let s be a real number with $0 < s < 1$. Then for any positive even number n ,*

$$2(n-1)^s < A_{n,s}^{-1} < 2n^s$$

and for any positive odd number n ,

$$-2n^s < B_{n,s}^{-1} < -2(n-1)^s.$$

Proof. Let n be a positive even number. For every positive integer k , it is easy to see that

$$\begin{aligned} & \left(\frac{1}{(n+1+2k)^s} - \frac{1}{(n+2+2k)^s} \right) \\ & < \left(\frac{1}{(n+2k)^s} - \frac{1}{(n+1+2k)^s} \right) \\ & < \left(\frac{1}{(n-1+2k)^s} - \frac{1}{(n+2k)^s} \right). \end{aligned}$$

The summations of each term over k give

$$A_{n+1,s} < A_{n,s} < A_{n-1,s}$$

and

$$\frac{1}{2}(A_{n+1,s} + A_{n,s}) < A_{n,s} < \frac{1}{2}(A_{n-1,s} + A_{n,s}).$$

Therefore, we have

$$\frac{1}{2n^s} < A_{n,s} < \frac{1}{2(n-1)^s},$$

which gives the first statement.

The second statement can be shown similarly. □

Since every proof of the case when n is an odd number is analogous to that of the case when n is an even number, we omit all the proofs of the odd number cases in this paper.

Now we find tighter bounds for $A_{n,s}^{-1}$ and $B_{n,s}^{-1}$.

Proposition 2.2. *Let s be a real number with $0 < s < 1$. Then for any positive even number n ,*

$$2 \left(n - \frac{1}{2} \right)^s < A_{n,s}^{-1}$$

and for any positive odd number n ,

$$B_{n,s}^{-1} < -2 \left(n - \frac{1}{2} \right)^s.$$

Proof. Let n be a positive even number. We will show that

$$A_{n,s} < \frac{1}{2 \left(n - \frac{1}{2} \right)^s}.$$

Rewriting each of the both sides as a series,

$$A_{n,s} = \sum_{k=\frac{n}{2}}^{\infty} \left(\frac{1}{(2k)^s} - \frac{1}{(2k+1)^s} \right) \quad \text{and}$$

$$\frac{1}{2(n-\frac{1}{2})^s} = \sum_{k=\frac{n}{2}}^{\infty} \left(\frac{1}{2(2k-\frac{1}{2})^s} - \frac{1}{2(2k+\frac{3}{2})^s} \right),$$

we will show that for any positive integer k ,

$$\frac{1}{(2k)^s} - \frac{1}{(2k+1)^s} < \frac{1}{2(2k-\frac{1}{2})^s} - \frac{1}{2(2k+\frac{3}{2})^s}.$$

For this, we let

$$f(x) = \left(\frac{1}{2(2x-\frac{1}{2})^s} - \frac{1}{2(2x+\frac{3}{2})^s} \right) - \left(\frac{1}{(2x)^s} - \frac{1}{(2x+1)^s} \right)$$

and will show that $f(x)$ is positive for $x \geq 1$ and $0 < s < 1$. With

$$g(x) = \frac{1}{2(2x-\frac{1}{2})^s} + \frac{1}{2(2x+\frac{1}{2})^s} - \frac{1}{(2x)^s},$$

we have $f(x) = g(x) - g(x+\frac{1}{2})$. Consider the derivative of $g(x)$,

$$g'(x) = -2s \left(\frac{1}{2(2x-\frac{1}{2})^{s+1}} + \frac{1}{2(2x+\frac{1}{2})^{s+1}} - \frac{1}{(2x)^{s+1}} \right).$$

Since the function $\frac{1}{x^{s+1}}$ is convex, we obtain that

$$\frac{1}{2(2x-\frac{1}{2})^{s+1}} + \frac{1}{2(2x+\frac{1}{2})^{s+1}} - \frac{1}{(2x)^{s+1}} \geq 0$$

and therefore $g'(x)$ is negative, that is, $g(x)$ is decreasing. We conclude that $f(x)$ is positive which gives the statement. \square

Proposition 2.3. *Let s be a real number with $0 < s < 1$. Then for any positive even number n ,*

$$A_{n,s}^{-1} < 2 \left(n - \frac{1}{4} \right)^s$$

and for any positive odd number n ,

$$-2 \left(n - \frac{1}{4} \right)^s < B_{n,s}^{-1}.$$

Proof. Let n be a positive even number. We will show that

$$\frac{1}{2 \left(n - \frac{1}{4} \right)^s} < A_{n,s}.$$

Rewriting each of the both sides as a series,

$$A_{n,s} = \sum_{k=\frac{n}{2}}^{\infty} \left(\frac{1}{(2k)^s} - \frac{1}{(2k+1)^s} \right) \quad \text{and}$$

$$\frac{1}{2(n-\frac{1}{4})^s} = \sum_{k=\frac{n}{2}}^{\infty} \left(\frac{1}{2(2k-\frac{1}{4})^s} - \frac{1}{2(2k+\frac{7}{4})^s} \right),$$

we need to show that for any positive integer k ,

$$\frac{1}{2(2k-\frac{1}{4})^s} - \frac{1}{2(2k+\frac{7}{4})^s} < \frac{1}{(2k)^s} - \frac{1}{(2k+1)^s}.$$

For this, we let

$$f(x) = \left(\frac{1}{(2x)^s} - \frac{1}{(2x+1)^s} \right) - \left(\frac{1}{2(2x-\frac{1}{4})^s} - \frac{1}{2(2x+\frac{7}{4})^s} \right).$$

We check that $f(1) > 0$ and now we will show that $f(x)$ is positive for $x \geq 2$ and $0 < s < 1$.

With

$$g(x) = \frac{1}{(2x)^s} - \left(\frac{1}{2(2x-\frac{1}{4})^s} + \frac{1}{2(2x+\frac{3}{4})^s} \right),$$

we have $f(x) = g(x) - g(x+\frac{1}{2})$, so we only need to show that $g(x)$ is decreasing. Consider the derivative of $g(x)$,

$$g'(x) = s \left(-\frac{2}{(2x)^{s+1}} + \left(\frac{1}{(2x-\frac{1}{4})^{s+1}} + \frac{1}{(2x+\frac{3}{4})^{s+1}} \right) \right)$$

$$= s \left(\left(\frac{1}{(2x-\frac{1}{4})^{s+1}} - \frac{1}{(2x)^{s+1}} \right) - \left(\frac{1}{(2x)^{s+1}} - \frac{1}{(2x+\frac{3}{4})^{s+1}} \right) \right).$$

Since the function $\frac{1}{x^{s+1}}$ is decreasing and convex, by comparing slopes at $(2x-\frac{1}{4})$ and $(2x+\frac{3}{4})$, we obtain

$$\frac{1}{(2x-\frac{1}{4})^{s+1}} - \frac{1}{(2x)^{s+1}} < \frac{1}{4}(s+1) \frac{1}{(2x-\frac{1}{4})^{s+2}}$$

and

$$\frac{1}{(2x)^{s+1}} - \frac{1}{(2x+\frac{3}{4})^{s+1}} > \frac{1}{4}(s+1) \frac{3}{(2x+\frac{3}{4})^{s+2}}.$$

Therefore,

$$g'(x) < \frac{1}{4}s(s+1) \left(\frac{1}{(2x-\frac{1}{4})^{s+2}} - \frac{3}{(2x+\frac{3}{4})^{s+2}} \right).$$

Consider $h(x, s) := \frac{1}{3} \left(\frac{2x+3/4}{2x-1/4} \right)^{s+2}$ which is the ratio of two terms on the right-hand side of the above expression. We check that $h(x, s) < 1$ for $x \geq 2$ and $0 < s < 1$. Since $h(2, 1) = 6859/10125$ and $\lim_{x \rightarrow \infty} h(x, s) = \frac{1}{3}$ for $0 < s < 1$, we obtain that $g'(x)$ is negative and therefore $g(x)$ is decreasing which gives the statement. \square

We combine the results of Proposition 2.2 and Proposition 2.3.

Theorem 2.4. *Let s be a real number with $0 < s < 1$. Then for any positive even number n ,*

$$2 \left(n - \frac{1}{2} \right)^s < A_{n,s}^{-1} < 2 \left(n - \frac{1}{4} \right)^s$$

and for any positive odd number n ,

$$-2 \left(n - \frac{1}{4} \right)^s < B_{n,s}^{-1} < -2 \left(n - \frac{1}{2} \right)^s.$$

We express these bounds in terms of $\zeta_n(s)$ using expression (1.1).

Corollary 2.5. *Let s be a real number with $0 < s < 1$. Then for any positive even number n ,*

$$2(1 - 2^{1-s}) \left(n - \frac{1}{4} \right)^s < \zeta_n(s)^{-1} < 2(1 - 2^{1-s}) \left(n - \frac{1}{2} \right)^s$$

and for any positive odd number n ,

$$-2(1 - 2^{1-s}) \left(n - \frac{1}{2} \right)^s < \zeta_n(s)^{-1} < -2(1 - 2^{1-s}) \left(n - \frac{1}{4} \right)^s.$$

Furthermore, we have tighter bounds of $A_{n,s}^{-1}$ and $B_{n,s}^{-1}$ for a sufficiently large number n .

Theorem 2.6. *For any positive number ϵ and any real number s with $0 < s < 1$,*

$$2 \left(n - \frac{1}{2} \right)^s < A_{n,s}^{-1} < 2 \left(n - \frac{1}{2} + \epsilon \right)^s$$

for a sufficiently large even number n and

$$-2 \left(n - \frac{1}{2} + \epsilon \right)^s < B_{n,s}^{-1} \leq -2 \left(n - \frac{1}{2} \right)^s$$

for a sufficiently large odd number n .

Proof. From Theorem 2.4, it suffices to show that for a sufficiently large even number n ,

$$\frac{1}{2 \left(n - \frac{1}{2} + \epsilon \right)^s} < A_{n,s}.$$

Rewriting each of the both sides as a series,

$$A_{n,s} = \sum_{k=\frac{n}{2}}^{\infty} \left(\frac{1}{(2k)^s} - \frac{1}{(2k+1)^s} \right) \text{ and}$$

$$\frac{1}{2 \left(n - \frac{1}{2} + \epsilon \right)^s} = \sum_{k=\frac{n}{2}}^{\infty} \left(\frac{1}{2 \left(2k - \frac{1}{2} + \epsilon \right)^s} - \frac{1}{2 \left(2k + \frac{3}{2} + \epsilon \right)^s} \right),$$

we need to show that for a sufficiently large even number n and every integer $k \geq \frac{n}{2}$,

$$\frac{1}{2(2k - \frac{1}{2} + \epsilon)^s} - \frac{1}{2(2k + \frac{3}{2} + \epsilon)^s} < \frac{1}{(2k)^s} - \frac{1}{(2k+1)^s}.$$

For this, let

$$f(x) = \left(\frac{1}{(2x)^s} - \frac{1}{(2x+1)^s} \right) - \left(\frac{1}{2(2x - \frac{1}{2} + \epsilon)^s} - \frac{1}{2(2x + \frac{3}{2} + \epsilon)^s} \right)$$

and will show that $f(x)$ is positive for $x \geq x_0$ where x_0 is a sufficiently large number. With

$$g(x) = \frac{1}{(2x)^s} - \left(\frac{1}{2(2x - \frac{1}{2} + \epsilon)^s} + \frac{1}{2(2x + \frac{1}{2} + \epsilon)^s} \right),$$

we have that $f(x) = g(x) - g(x + \frac{1}{2})$, so we only need to show that $g(x)$ is decreasing. Consider the derivative of $g(x)$,

$$\begin{aligned} g'(x) &= s \left(-\frac{2}{(2x)^{s+1}} + \frac{1}{(2x - \frac{1}{2} + \epsilon)^{s+1}} + \frac{1}{(2x + \frac{1}{2} + \epsilon)^{s+1}} \right) \\ &= s \left(\left(\frac{1}{(2x - \frac{1}{2} + \epsilon)^{s+1}} - \frac{1}{(2x)^{s+1}} \right) - \left(\frac{1}{(2x)^{s+1}} - \frac{1}{(2x + \frac{1}{2} + \epsilon)^{s+1}} \right) \right). \end{aligned}$$

Since $\frac{1}{x^{s+1}}$ is decreasing and convex, by comparing slopes at $(2x - \frac{1}{2} + \epsilon)$ and $(2x + \frac{1}{2} + \epsilon)$, we obtain

$$\frac{1}{(2x - \frac{1}{2} + \epsilon)^{s+1}} - \frac{1}{(2x)^{s+1}} < (s+1) \frac{\frac{1}{2} - \epsilon}{(2x - \frac{1}{2} + \epsilon)^{s+2}}$$

and

$$\frac{1}{(2x)^{s+1}} - \frac{1}{(2x + \frac{1}{2} + \epsilon)^{s+1}} > (s+1) \frac{\frac{1}{2} + \epsilon}{(2x + \frac{1}{2} + \epsilon)^{s+2}}.$$

Therefore

$$g'(x) < s(s+1) \left(\frac{\frac{1}{2} - \epsilon}{(2x - \frac{1}{2} + \epsilon)^{s+2}} - \frac{\frac{1}{2} + \epsilon}{(2x + \frac{1}{2} + \epsilon)^{s+2}} \right).$$

Consider $h(x) := \frac{\frac{1}{2} - \epsilon}{\frac{1}{2} + \epsilon} \left(\frac{2x + \frac{1}{2} + \epsilon}{2x - \frac{1}{2} + \epsilon} \right)^{s+2}$ which is the ratio of two terms on the right-hand side of the above expression. We need to show that $h(x) < 1$ for every $x > x_0$ where x_0 is a sufficiently large number. We check that

$$h(x) < 1 \iff \frac{2x + \frac{1}{2} + \epsilon}{2x - \frac{1}{2} + \epsilon} < \left(\frac{\frac{1}{2} + \epsilon}{\frac{1}{2} - \epsilon} \right)^{\frac{1}{s+2}}.$$

For any $\epsilon > 0$ and $0 < s < 1$, we have that $1 < \left(\frac{\frac{1}{2} + \epsilon}{\frac{1}{2} - \epsilon} \right)^{1/(s+2)}$ and $\frac{2x + \frac{1}{2} + \epsilon}{2x - \frac{1}{2} + \epsilon}$ is larger than 1, decreasing and converges to 1 as x goes to infinite, so there is x_0 so that for every $x > x_0$, $h(x) < 1$. Therefore the proof is complete. \square

We express these bounds in terms of $\zeta_n(s)$ using expression (1.1).

Corollary 2.7. *For any positive number ϵ and any real number s with $0 < s < 1$,*

$$2(1 - 2^{1-s}) \left(n - \frac{1}{2} + \epsilon \right)^s < \zeta_n(s)^{-1} < 2(1 - 2^{1-s}) \left(n - \frac{1}{2} \right)^s$$

for a sufficiently large even number n and

$$-2(1 - 2^{1-s}) \left(n - \frac{1}{2} \right)^s < \zeta_n(s)^{-1} < -2(1 - 2^{1-s}) \left(n - \frac{1}{2} + \epsilon \right)^s$$

for a sufficiently large odd number n .

2.2 The value of the inverse of $\zeta_n(s)$ for $s = \frac{1}{2}, \frac{1}{3}$ and $\frac{1}{4}$

We study firstly the value of the inverse of $\zeta_n(\frac{1}{2})$, where $\zeta_n(\frac{1}{2})$ is the tail of the Riemann zeta function from n at $s = \frac{1}{2}$.

Theorem 2.8. *For any positive even number n ,*

$$[A_{n,1/2}^{-1}] = \left[2 \left(n - \frac{1}{2} \right)^{1/2} \right]$$

and for any positive odd number n ,

$$[B_{n,1/2}^{-1}] = \left[-2 \left(n - \frac{1}{2} \right)^{1/2} \right].$$

Proof. Let n be a positive even number. By Theorem 2.4, we have that

$$2 \left(n - \frac{1}{2} \right)^{1/2} < A_{n,1/2}^{-1} < 2 \left(n - \frac{1}{4} \right)^{1/2}.$$

Note that $2(n - \frac{1}{4})^{1/2} - 2(n - \frac{1}{2})^{1/2} < 1$ for $n \geq 2$, and it implies that there is at most one integer in the open interval from $2(n - \frac{1}{2})^{1/2}$ to $2(n - \frac{1}{4})^{1/2}$. Suppose that there is an integer h in the open interval, i.e.,

$$2 \left(n - \frac{1}{2} \right)^{1/2} < h < 2 \left(n - \frac{1}{4} \right)^{1/2} \quad \text{or} \quad 4n - 2 < h^2 < 4n - 1.$$

There is, however, no integer in the open interval from $4n - 2$ to $4n - 1$, therefore such an integer h does not exist. This gives the statement. \square

We express this result in terms of $\zeta_n(s)$ using expression (1.1).

Corollary 2.9. *For any positive integer n ,*

$$\left[\frac{1}{1 - 2^{1/2}} \zeta_n \left(\frac{1}{2} \right)^{-1} \right] = \left[(-1)^{n+1} 2 \left(n - \frac{1}{2} \right)^{1/2} \right].$$

We study secondly the value of the inverse of $\zeta_n(\frac{1}{3})$, where $\zeta_n(\frac{1}{3})$ is the tail of the Riemann zeta function from n at $s = \frac{1}{3}$.

Theorem 2.10. For any positive even number n ,

$$[A_{n,1/3}^{-1}] = \left[2 \left(n - \frac{1}{2} \right)^{1/3} \right]$$

and for any positive odd number n ,

$$[B_{n,1/3}^{-1}] = \left[-2 \left(n - \frac{1}{2} \right)^{1/3} \right].$$

Proof. Let n be a positive even number. By Theorem 2.4, we have that

$$2 \left(n - \frac{1}{2} \right)^{1/3} < A_{n,1/3}^{-1} < 2 \left(n - \frac{1}{4} \right)^{1/3}.$$

Note that $2(n - \frac{1}{4})^{1/3} - 2(n - \frac{1}{2})^{1/3} < 1$ for $n \geq 2$, and it implies that there is at most one integer in the open interval from $2(n - \frac{1}{2})^{1/3}$ to $2(n - \frac{1}{4})^{1/3}$. Suppose that there is an integer h in the open interval, i.e.,

$$2(n - \frac{1}{2})^{1/3} < h < 2(n - \frac{1}{4})^{1/3} \quad \text{or} \quad 8n - 4 < h^3 < 8n - 2.$$

This shows that the integer h is of the form $h = 2(n - \frac{3}{8})^{1/3}$. If we show $A_{n,1/3}^{-1} < 2(n - \frac{3}{8})^{1/3}$ or equivalently, $\frac{1}{2(n - \frac{3}{8})^{1/3}} < A_{n,1/3}$, then our proof will be done. Let us rewrite

$$A_{n,1/3} = \sum_{k=\frac{n}{2}}^{\infty} \left(\frac{1}{(2k)^{1/3}} - \frac{1}{(2k+1)^{1/3}} \right) \quad \text{and}$$

$$\frac{1}{2(n - \frac{3}{8})^{1/3}} = \sum_{k=\frac{n}{2}}^{\infty} \left(\frac{1}{2(2k - \frac{3}{8})^{1/3}} - \frac{1}{2(2k + \frac{13}{8})^{1/3}} \right).$$

Now it suffices to show that for any positive integer k ,

$$\frac{1}{2(2k - \frac{3}{8})^{1/3}} - \frac{1}{2(2k + \frac{13}{8})^{1/3}} < \frac{1}{(2k)^{1/3}} - \frac{1}{2(2k+1)^{1/3}}.$$

For this, we let

$$f(x) = \left(\frac{1}{(2x)^{1/3}} - \frac{1}{(2x+1)^{1/3}} \right) - \left(\frac{1}{2(2x - \frac{3}{8})^{1/3}} - \frac{1}{2(2x + \frac{13}{8})^{1/3}} \right)$$

and we will show that $f(x)$ is positive for any positive integer x .

We check that $f(1) = 0.00053\dots$ and $f(2) = 0.00081\dots$, so it suffices to show $f(x) > 0$ for $x \geq 3$. With

$$g(x) = \frac{1}{(2x)^{1/3}} - \left(\frac{1}{2(2x - \frac{3}{8})^{1/3}} + \frac{1}{2(2x + \frac{5}{8})^{1/3}} \right),$$

we have that $f(x) = g(x) - g(x + \frac{1}{2})$, so we only need to show that $g(x)$ is decreasing for $x \geq 3$. Consider the derivative of $g(x)$,

$$\begin{aligned} g'(x) &= \frac{1}{3} \left(-\frac{2}{(2x)^{4/3}} + \frac{1}{(2x - \frac{3}{8})^{4/3}} + \frac{1}{(2x + \frac{5}{8})^{4/3}} \right) \\ &= \frac{1}{3} \left(\left(\frac{1}{(2x - \frac{3}{8})^{4/3}} - \frac{1}{(2x)^{4/3}} \right) - \left(\frac{1}{(2x)^{4/3}} - \frac{1}{(2x + \frac{5}{8})^{4/3}} \right) \right). \end{aligned}$$

Since $\frac{1}{x^{4/3}}$ is decreasing and convex, by comparing slopes at $(2x - \frac{3}{8})$ and $(2x + \frac{5}{8})$, we obtain

$$\frac{1}{(2x - \frac{3}{8})^{4/3}} - \frac{1}{(2x)^{4/3}} < 2 \cdot \frac{3}{16} \cdot \frac{4}{3} \cdot \frac{1}{(2x - \frac{3}{8})^{7/3}}$$

and

$$\frac{1}{(2x)^{4/3}} - \frac{1}{(2x + \frac{5}{8})^{4/3}} > 2 \cdot \frac{5}{16} \cdot \frac{4}{3} \cdot \frac{1}{(2x + \frac{5}{8})^{7/3}}.$$

Therefore

$$g'(x) < \frac{1}{18} \left(\frac{3}{(2x - \frac{3}{8})^{7/3}} - \frac{5}{(2x + \frac{5}{8})^{7/3}} \right).$$

Consider $h(x) := \frac{3}{5} \left(\frac{2x+5/8}{2x-3/8} \right)^{7/3}$ which is the ratio of two terms of the right-hand side of the above expression. We check that $h(x) < 1$ for $x \geq 3$ because $h(3) = 0.87 \dots$ and $\lim_{x \rightarrow \infty} h(x) = \frac{3}{5}$ and $h'(x) < 0$ for $x \geq 3$. Hence we obtain that $g'(x)$ is negative and so $g(x)$ is decreasing for $x \geq 3$ which proves the statement. \square

We express this result in terms of $\zeta_n(s)$ using expression (1.1).

Corollary 2.11. *For any positive integer n ,*

$$\left[\frac{1}{1 - 2^{2/3}} \zeta_n \left(\frac{1}{3} \right)^{-1} \right] = \left[(-1)^{n+1} 2 \left(n - \frac{1}{2} \right)^{1/3} \right].$$

We study lastly the value of the inverse of $\zeta_n(\frac{1}{4})$, which is the tail of the Riemann zeta function from n at $s = \frac{1}{4}$.

Theorem 2.12. *For any positive even number n ,*

$$[A_{n,1/4}^{-1}] = \left[2 \left(n - \frac{1}{2} \right)^{1/4} \right]$$

and for any positive odd number n ,

$$[B_{n,1/4}^{-1}] = \left[-2 \left(n - \frac{1}{2} \right)^{1/4} \right].$$

Proof. Let n be a positive even number. By Theorem 2.4, we have that

$$2 \left(n - \frac{1}{2} \right)^{1/4} < A_{n,1/4}^{-1} < 2 \left(n - \frac{1}{4} \right)^{1/4}.$$

Note that $2(n - \frac{1}{4})^{1/4} - 2(n - \frac{1}{2})^{1/4} < 1$ for $n \geq 2$ and it implies that there is at most one integer in the open interval from $2(n - \frac{1}{2})^{1/4}$ to $2(n - \frac{1}{4})^{1/4}$. Suppose that there is an integer h in the open interval, i.e.,

$$2(n - \frac{1}{2})^{1/4} < h < 2(n - \frac{1}{4})^{1/4} \quad \text{or} \quad 16n - 8 < h^4 < 16n - 4.$$

This shows that the integer h^4 is one of the form $16n - 7$, $16n - 6$ or $16n - 5$. For any integer h , however $h^4 \equiv 0$ or $1 \pmod{16}$, hence such an integer h does not exist. Therefore this gives the statement. \square

We express this result in terms of $\zeta_n(s)$ using expression (1.1).

Corollary 2.13. *For any positive integer n ,*

$$\left[\frac{1}{1 - 2^{3/4}} \zeta_n \left(\frac{1}{4} \right)^{-1} \right] = \left[(-1)^{n+1} 2 \left(n - \frac{1}{2} \right)^{1/4} \right].$$

We express the results of Theorems 2.8, 2.10 and 2.12 in a single statement.

Theorem 2.14. *For $s = \frac{1}{2}$, $\frac{1}{3}$, or $\frac{1}{4}$, and for any positive even number n ,*

$$[A_{n,s}^{-1}] = \left[2 \left(n - \frac{1}{2} \right)^s \right]$$

and for any positive odd number n ,

$$[B_{n,s}^{-1}] = \left[-2 \left(n - \frac{1}{2} \right)^s \right].$$

We express the results of Corollary 2.9, 2.11 and 2.13 in a single statement.

Corollary 2.15. *For any positive integer n and $s = \frac{1}{2}$, $\frac{1}{3}$, or $\frac{1}{4}$,*

$$\left[\frac{1}{1 - 2^{1-s}} \zeta_n(s)^{-1} \right] = \left[(-1)^{n+1} 2 \left(n - \frac{1}{2} \right)^s \right].$$

3 Conclusion

In this paper, we present the bounds of $A_{n,s}^{-1}$ and $B_{n,s}^{-1}$ hence the bounds of the inverses of tails of the Riemann zeta function, $\zeta_n(s)^{-1}$, for $0 < s < 1$, and compute the values $[A_{n,s}^{-1}]$ and $[B_{n,s}^{-1}]$ hence the values of the inverses of tails of the Riemann zeta function, $\left[\frac{1}{1 - 2^{1-s}} \zeta_n(s)^{-1} \right]$, for $s = \frac{1}{2}$, $\frac{1}{3}$ and $\frac{1}{4}$. For other values of s , for example $s = \frac{1}{5}$ or $\frac{2}{3}$, etc, the values of $A_{n,s}$ and $B_{n,s}$ don't seem to have simple expressions.

Authors' Contributions

All authors equally contributed to the manuscript.

Acknowledgements

The authors would like to express their thanks to the referees and the editors for their helpful comments and advice.

References

- [1] Borwein, P, Choi, S, Rooney, B, Weirathmuellerer, A: The Riemann hypothesis. CMS Books in Mathematics/Ouvrages de Mathématiques de la SMC, Springer, New York (2008).
- [2] Euler, L: *Variae observationes circa series infinitas*. *Commentarii academiae scientiarum Petropolitanae* 9 160–188 (1744), (presented to the St. Petersburg Academy in 1737) = *Opera Omnia Ser.* **114**, 217–244. URL: <https://www.biodiversitylibrary.org>
- [3] Hwang, W, Song, K: A reciprocal sum related to the Riemann zeta function at $s = 6$. Preprint. URL: <https://arxiv.org/abs/1709.07994>
- [4] Luo, Qiu-Ming, Guo, Bai-Ni, Qi, Feng: On evaluation of Riemann zeta function $\zeta(s)$. *Advanced Studies in Contemporary Mathematics (Kyungshang)*. **7**(2), 135–144 (2003).
- [5] Luo, Qiu-Ming, Wei, Zong-Li, Qi, Feng: Lower and upper bounds of $\zeta(3)$. *Advanced Studies in Contemporary Mathematics (Kyungshang)*. **6**(1), 47–51 (2003).
- [6] Riemann, B: Über die Anzahl der Primzahlen unter einer gegebenen Grösse. *Monatsberichte der Berliner Akademie*. 671–680 (1859).
- [7] Xin, L: Some identities related to Riemann zeta-function. *Journal of Inequalities and Applications*. **2016**(1), 32 (2016). DOI: 10.1186/s13660-016-0980-9
- [8] Xin L, Xiaoxue, L: A reciprocal sum related to the Riemann ζ - function. *Journal of mathematical inequalities*. **11**(1), 209–215 (2017). DOI: 10.7153/jmi-11-20
- [9] Xu, H: Some computational formulas related to the Riemann zeta-function tails. *Journal of Inequalities and Applications*. **2016**(1), 132 (2016). DOI: 10.1186/s13660-016-1068-2