

# VERTEX TYPES IN THRESHOLD AND CHAIN GRAPHS

M. ANDELIĆ, E. GHORBANI, AND S.K. SIMIĆ

*In honour of Domingos M. Cardoso on the occasion of his 65th birthday*

ABSTRACT. A graph is called a chain graph if it is bipartite and the neighborhoods of the vertices in each color class form a chain with respect to inclusion. A threshold graph can be obtained from a chain graph by making adjacent all pairs of vertices in one color class. Given a graph  $G$ , let  $\lambda$  be an eigenvalue (of the adjacency matrix) of  $G$  with multiplicity  $k \geq 1$ . A vertex  $v$  of  $G$  is a downer, or neutral, or Parter depending whether the multiplicity of  $\lambda$  in  $G - v$  is  $k - 1$ , or  $k$ , or  $k + 1$ , respectively. We consider vertex types in the above sense in threshold and chain graphs. In particular, we show that chain graphs can have neutral vertices, disproving a conjecture by Alazemi *et al.*

## 1. INTRODUCTION

This paper is a successor of [4] in which vertex types (see the Abstract) in the lexicographic products of an arbitrary graph over cliques and/or co-cliques were investigated. Such class of graphs includes threshold graphs and chain graphs as particular instances. Both of these types (or classes) of graphs were discovered, and also rediscovered by various researchers in different contexts (see, for example, [5, 6, 12], and references therein). Needless to say, they were named by different names mostly depending on applications in which they arise. It is also noteworthy that threshold graphs are subclass of cographs, i.e. of  $P_4$ -free graphs. Recall that threshold graphs are  $\{P_4, 2K_2, C_4\}$ -free graphs, while chain graphs are  $\{2K_2, C_3, C_5\}$ -free graphs - see [1, 3] for more details. Note, if these graphs are not connected then (since  $2K_2$  is forbidden) at most one of its components is non-trivial (others are trivial, i.e. isolated vertices). Moreover, stars are the only connected graphs which belong to both of two classes of graphs under consideration.

Recall, these graphs play a very important role in Spectral Graph Theory, since the maximizers for the largest eigenvalue of the adjacency matrix (for graphs of fixed order and size, either connected or disconnected) belong to these classes (threshold graphs in general case,

---

2000 *Mathematics Subject Classification.* 05C50.

*Key words and phrases.* adjacency spectrum, chain graphs, threshold graphs, vertex types.

and chain graphs in bipartite case). Such graphs (in both classes) have a very specific structure (embodied in nesting property), and this fact enables us to tell more on the type of certain vertices. Here, we also disprove Conjecture 3.1 from [3].

Throughout, we will consider simple graphs, i.e. finite undirected graphs without loops or multiple edges. In addition, without loss of generality, we will assume that any such graph is connected. For a graph  $G$  we denote its vertex set by  $V(G)$ , and by  $n = |V(G)|$  its order. An  $n \times n$  matrix  $A(G) = [a_{ij}]$  is its *adjacency matrix* if  $a_{ij} = 1$  whenever vertices  $i$  and  $j$  are adjacent, or  $a_{ij} = 0$  otherwise. For a vertex  $v$  of  $G$ , let  $N(v)$  denote the *neighborhood* of  $v$ , i.e. the set of all vertices of  $G$  adjacent to  $v$ .

The eigenvalues of  $G$  are the eigenvalues of its adjacency matrix. In non-increasing order they are denoted by

$$\lambda_1(G) \geq \lambda_2(G) \geq \cdots \geq \lambda_n(G),$$

or by

$$\mu_1(G) > \mu_2(G) > \cdots > \mu_r(G)$$

if only distinct eigenvalues are considered. If understandable from the context we will drop out graph names from the notation of eigenvalues (or other related objects). The eigenvalues comprise (together with multiplicities, say  $k_1, k_2, \dots, k_r$ , respectively) the spectrum of  $G$ , denoted by  $\text{Spec}(G)$ . The *characteristic polynomial* of  $G$ , denoted by  $\phi(x; G)$ , is the characteristic polynomial of its adjacency matrix. Both, the spectrum and characteristic polynomial of a graph  $G$  are its invariants. Further on, all spectral invariants (and other relevant quantities) associated to the adjacency matrix will be prescribed to the corresponding graph. For a given eigenvalue  $\lambda \in \text{Spec}(G)$ ,  $\text{mult}(\lambda, G)$  denotes its multiplicity, while  $\mathcal{E}(\lambda; G)$  its eigenspace (provided  $G$  is a labeled graph). The equation  $A\mathbf{x} = \lambda\mathbf{x}$ , is called the eigenvalue equation for  $\lambda$ . Here  $A$  is the adjacency matrix, while  $\mathbf{x}$  a  $\lambda$ -eigenvector also of the labeled graph  $G$ . If  $G$  is of order  $n$ , then  $\mathbf{x}$  can be seen as an element of  $\mathbb{R}^n$ , or a mapping  $\mathbf{x} : V(G) \rightarrow \mathbb{R}^n$  (so its  $i$ -th entry can be denoted by  $x_i$  or  $\mathbf{x}(i)$ ). Eigenspaces (as the eigenvector sets) are not graph invariants, since the eigenvector entries become permuted if the vertices of  $G$  are relabeled.

An eigenvalue  $\lambda \in \text{Spec}(G)$  is *main* if the corresponding eigenspace  $\mathcal{E}(\lambda; G)$  is not orthogonal to all-1 vector  $\mathbf{j}$ ; otherwise, it is *non-main*.

Given a graph  $G$ , let  $\lambda$  be its eigenvalue of multiplicity  $k \geq 1$  and  $v \in V(G)$ . Then  $v$  is a *downer*, or *neutral*, or *Parter* vertex of  $G$ , depending whether the multiplicity of  $\lambda$  in  $G - v$  is  $k - 1$ , or  $k$ , or  $k + 1$ , respectively. Recall, neutral and Parter vertices of  $G$  are also called *Fiedler* vertices. For more details, about the above vertex types see, for example, [19].

**Remark 1.1.** *Sum rule: Let  $\mathbf{x}$  be a  $\lambda$ -eigenvector of a graph  $G$ . Then the entries of  $\mathbf{x}$  satisfy the following equalities:*

$$(1) \quad \lambda \mathbf{x}(v) = \sum_{u \sim v} \mathbf{x}(u), \text{ for all } v \in V(G).$$

From (1) it follows that if  $\lambda \neq 0$ , then  $N(u) = N(v)$  implies that  $\mathbf{x}(u) = \mathbf{x}(v)$  and if  $\lambda \neq -1$ ,  $N(u) \cup \{u\} = N(v) \cup \{v\}$  implies that  $\mathbf{x}(u) = \mathbf{x}(v)$ .

In sequel, we will need the following interlacing property for graph eigenvalues (or, eigenvalues of Hermitian matrices, see [8, Theorem 2.5.1]).

**Theorem 1.1.** *Let  $G$  be a graph of order  $n$  and  $G'$  be an induced subgraph of  $G$  of order  $n'$ . If  $\lambda_1 \geq \lambda_2 \geq \dots \geq \lambda_n$  and  $\lambda'_1 \geq \lambda'_2 \geq \dots \geq \lambda'_{n'}$  are their eigenvalues respectively, then*

$$(2) \quad \lambda_i \geq \lambda'_i \geq \lambda_{n-n'+i} \text{ for } i = 1, 2, \dots, n'.$$

In particular, if  $n' = n - 1$ , then

$$\lambda_1 \geq \lambda'_1 \geq \lambda_2 \geq \lambda'_2 \geq \dots \geq \lambda_{n-1} \geq \lambda'_{n-1} \geq \lambda_n.$$

In the case of equality in (2) (see [8, Theorem 2.5.1]) the following holds.

**Lemma 1.1.** *If  $\lambda'_i = \lambda_i$  or  $\lambda'_i = \lambda_{n-n'+i}$  for some  $i \in \{1, 2, \dots, n'\}$ , then  $G'$  has an eigenvector  $\mathbf{x}'$  for  $\lambda'_i$  such that  $\begin{pmatrix} \mathbf{0} \\ \mathbf{x}' \end{pmatrix}$  is an eigenvector of  $G$  for  $\lambda'_i$ , where  $\mathbf{0}$  is a zero vector whose entries correspond to the vertices from  $V(G) \setminus V(G')$ .*

**Remark 1.2.** *A vertex  $v$  is a downer for a fixed eigenvalue  $\lambda$ , if there exists in the corresponding eigenspace an eigenvector whose  $v$ -th component is non-zero. Otherwise, it is a Fiedler vertex. Let  $W$  be the eigenspace corresponding to  $\lambda$ . If for each  $\mathbf{x} \in W$ , we have  $\mathbf{x}(v) = 0$ , then  $v$  cannot be a downer vertex as for any  $\mathbf{x} \in W$ , the vector  $\mathbf{x}'$  obtained by deleting the  $v$ -th component, is a  $\lambda$ -eigenvector of  $G - v$ , and therefore we have*

$$\text{mult}(\lambda, G - v) \geq \dim \{\mathbf{x}' : \mathbf{x} \in W\} = \dim W = \text{mult}(\lambda, G).$$

From this and Lemma 1.1 it follows, if  $\text{mult}(\lambda, G) = 1$  that there exists a  $\lambda$ -eigenvector  $\mathbf{x}$  with  $\mathbf{x}(v) = 0$  if and only if  $v$  is not a downer vertex for  $\lambda$ .

The rest of the paper is organized as follows: in Section 2 we give some particular results about vertex types in threshold graphs, while in Section 3 we put focus on chain graphs, and among others we disprove Conjecture 3.1 from [1], which states that in any chain graph, every vertex is a downer with respect to every non-zero eigenvalue. Besides we point out that some weak versions of the same conjecture are true.

## 2. VERTEX TYPES IN THRESHOLD GRAPHS

Any (connected) threshold graph  $G$  is a split graph i.e., it admits a partition of its vertex set into two subsets, say  $U$  and  $V$ , such that the vertices of  $U$  induce a co-clique, while the vertices of  $V$  induce a clique. All other edges join a vertex in  $U$  with a vertex in  $V$ . Moreover, if  $G$  is connected, then both  $U$  and  $V$  are partitioned into  $h \geq 1$  non-empty cells such that  $U = \bigcup_{i=1}^h U_i$  and  $V = \bigcup_{i=1}^h V_i$  and the following holds for (cross) edges: each vertex in  $U_i$  is adjacent to all vertices in  $V_1 \cup \dots \cup V_i$  (a nesting property). Accordingly, connected threshold graphs are also called *nested split graphs* (or NSG for short). If  $m_i = |U_i|$  and  $n_i = |V_i|$ , then we write

$$(3) \quad G = \text{NSG}(m_1, \dots, m_h; n_1, \dots, n_h),$$

(see Fig. 1). We denote by  $M_h (= \sum_{i=1}^h m_i)$  the size of  $U$ , and by  $N_h (= \sum_{i=1}^h n_i)$  the size of  $V$ .




FIGURE 1. The threshold graph  $G = \text{NSG}(m_1, \dots, m_h; n_1, \dots, n_h)$ .

The following Theorem states the essential spectral properties of threshold graphs (see [1, 15, 18]).

**Theorem 2.1.** *Let  $G = \text{NSG}(m_1, \dots, m_h; n_1, \dots, n_h)$ . Then the spectrum of  $G$  contains:*

- $h$  positive simple eigenvalues;
- $h - 1$  simple eigenvalues less than  $-1$  if  $m_h = 1$ , or otherwise if  $m_h \geq 2$ ,  $h$  simple eigenvalues less than  $-1$ ;
- eigenvalue  $0$  of multiplicity  $M_h - h$ , and  $-1$  of multiplicity  $N_h - h + 1$  if  $m_h = 1$ , or of multiplicity  $N_h - h$  if  $m_h > 1$ .

*In addition, if  $\lambda \neq 0, -1$  then  $\lambda$  is a main eigenvalue.*

**Remark 2.1.** *If  $\lambda \neq 0, -1$ , then any vertex of a threshold graph is either downer or neutral. Parter vertices may arise only for  $\lambda = 0$  or*

−1. Any vertex deleted subgraph  $G - v$  of a threshold graph  $G$  is a threshold graph as well. By Theorem 2.1 one can easily determine the multiplicities of 0 and −1 in both  $G$  and  $G - v$  and, consequently, the vertex type for  $v$  of  $\lambda = 0$  or −1.

Recall that any vertex of a connected graph is downer for the largest eigenvalue, see [10, Proposition 1.3.9.]. In addition, if  $\lambda \neq 0, -1$ , then the corresponding eigenvector  $\mathbf{x}$  is unique (up to scalar multiple) and constant on each of the sets  $U_i$  and  $V_i$  ( $i = 1, \dots, h$ ); in particular, if  $m_h = 1$  then it is constant on the set  $U_h \cup V_h$ . These facts will be used repeatedly further on without any recall.

**Theorem 2.2.** *Let  $G = \text{NSG}(m_1, \dots, m_h; n_1, \dots, n_h)$  and  $\lambda \neq 0, -1$  its eigenvalue other than the largest one. Then all vertices in  $U_1 \cup V_1$  are downers for  $\lambda$ . The same holds for vertices in  $U_h$ , and also in  $V_h$  unless  $\lambda = -m_h$  and  $m_h \geq 2$ .*

*Proof.* Let  $u_1 \in U_1$  and  $v_1 \in V_1$ . Then, by the sum rule,  $\lambda \mathbf{x}(u_1) = n_1 \mathbf{x}(v_1)$ . Since  $\lambda \neq 0, -1$ ,  $u_1$  and  $v_1$  are both downer or Fiedler vertices (see Remark 1.2). Let  $X = \sum_{w \in V(G)} \mathbf{x}(w)$ , and by the way of contradiction assume that  $u_1$  and  $v_1$  are both Fiedler vertices, i.e.  $\mathbf{x}(u_1) = \mathbf{x}(v_1) = 0$ . Again, by the sum rule, we have  $\lambda \mathbf{x}(v_1) = X - \mathbf{x}(v_1)$ , and therefore  $X = 0$ , a contradiction since  $\lambda \neq 0, -1$  is a simple and non-main eigenvalue (see Theorem 2.1).

Let  $u_h \in U_h$ ,  $v_h \in V_h$  and  $Y = \sum_{w \in V_1 \cup \dots \cup V_h} \mathbf{x}(w)$ . Then,  $\lambda \mathbf{x}(u_h) = Y$  and  $\lambda \mathbf{x}(v_h) = Y - \mathbf{x}(v_h) + m_h \mathbf{x}(u_h)$ . For a contradiction, let  $\mathbf{x}(u_h) = 0$ . Then it easily follows that  $\mathbf{x}(v_h) = 0$ . We next claim that for  $2 \leq i \leq h$ ,  $\mathbf{x}(u_i) = \mathbf{x}(v_i) = 0$  implies  $\mathbf{x}(u_{i-1}) = \mathbf{x}(v_{i-1}) = 0$ . To see this, since  $\mathbf{x}(v_i) = 0$  by the sum rule we obtain

$$\lambda \mathbf{x}(u_i) = Y - \sum_{j=i+1}^h n_j \mathbf{x}(v_j) = Y - \sum_{j=i}^h n_j \mathbf{x}(v_j) = \lambda \mathbf{x}(u_{i-1}),$$

and therefore  $\lambda \mathbf{x}(u_{i-1}) = 0$ . Similarly, since  $\mathbf{x}(u_i) = 0$  and

$$\begin{aligned} \lambda \mathbf{x}(v_i) &= Y - \mathbf{x}(v_i) + \sum_{j=i}^h m_j \mathbf{x}(u_j) \\ &= Y - \mathbf{x}(v_i) + \sum_{j=i-1}^h m_j \mathbf{x}(u_j) = (\lambda + 1) \mathbf{x}(v_{i-1}), \end{aligned}$$

it follows  $\mathbf{x}(v_{i-1}) = 0$ . Consequently, we obtain  $\mathbf{x}(u_h) = \dots = \mathbf{x}(u_1) = 0$  and  $\mathbf{x}(v_h) = \dots = \mathbf{x}(v_1) = 0$ , i.e.  $\mathbf{x} = \mathbf{0}$ , a contradiction. This proves that all vertices in  $U_h$  are downers for  $\lambda$ .

For the last part of the theorem, let  $\lambda \neq -m_h$ . Then we have

$$\lambda \mathbf{x}(u_h) = Y, \quad \lambda \mathbf{x}(v_h) = Y - \mathbf{x}(v_h) + m_h \mathbf{x}(u_h),$$

and so  $(\lambda + 1)\mathbf{x}(v_h) = (\lambda + m_h)\mathbf{x}(u_h)$ . Hence, if  $\mathbf{x}(v_h) = 0$ , then  $\mathbf{x}(u_h) = 0$  and we reach a contradiction as above. Consequently, all vertices in  $V_h$  are downers.  $\square$

**Remark 2.2.** *The following example shows that in unresolved case when  $\lambda = -m_h$  and  $m_h \geq 2$  vertices in  $V_h$  may be neutral.*

*Let  $G = \text{NSG}(2, 2, 2; 2, 3, 2)$ . Then all vertices in  $U_3$  are downers, while all vertices in  $V_3$  are neutral for  $\lambda = -2$ . So, an unresolved case from Theorem 2.2 can be an exceptional one.*

*So, the following question arises: Can we find an example when  $\lambda = -m_h$ ,  $m_h \geq 2$  and that each vertex in  $V_h$  is a downer?*

**Theorem 2.3.** *Let  $G = \text{NSG}(m_1, \dots, m_h; n_1, \dots, n_h)$  and let  $\lambda \neq 0, -1$  be its eigenvalue. Then, for any  $i = 1, \dots, h-1$ , at least one of  $U_i, U_{i+1}$  (resp.  $V_i, V_{i+1}$ ) contains only downer vertices for  $\lambda$ .*

*Proof.* Recall first that all vertices within  $U_k$  or  $V_k$  ( $k = 1, \dots, h$ ) are of the same type for  $\lambda$ , and that  $\lambda$  is a simple eigenvalue. Assume on the contrary that all vertices in  $U_i$  and  $U_{i+1}$  are neutral and let  $\mathbf{x}$  be a  $\lambda$ -eigenvector. Then, for  $u_i \in U_i$  and  $u_{i+1} \in U_{i+1}$ ,  $\mathbf{x}(u_i) = \mathbf{x}(u_{i+1}) = 0$ . By the sum rule it easily follows that for any  $v_{i+1} \in V_{i+1}$ ,  $\mathbf{x}(v_{i+1}) = 0$ . Next, we have

$$(4) \quad \lambda \mathbf{x}(v_i) = \sum_{j=1}^h n_j \mathbf{x}(v_j) - \mathbf{x}(v_i) + \sum_{j=i}^h m_j \mathbf{x}(u_j),$$

$$(5) \quad \lambda \mathbf{x}(v_{i+1}) = \sum_{j=1}^h n_j \mathbf{x}(v_j) - \mathbf{x}(v_{i+1}) + \sum_{j=i+1}^h m_j \mathbf{x}(u_j).$$

By subtracting (5) from (4) we obtain  $\lambda \mathbf{x}(v_i) = -\mathbf{x}(v_i)$ . Since  $\lambda \neq -1$ ,  $\mathbf{x}(v_i) = 0$  and consequently  $\mathbf{x}(u_{i-1}) = 0$ . Proceeding in the similar way, we conclude that  $\mathbf{x}(u_1) = 0$ , which contradicts Theorem 2.2.

The proof for vertices in  $V_i, V_{i+1}$  is similar, and therefore omitted.  $\square$

Next examples show that in an nested split graph  $G$  neutral vertices for the same eigenvalue may be distributed in different  $U_i$ 's,  $V_i$ 's and at the same time in both  $U$  and  $V$ .

**Example 2.1.** *If  $G = \text{NSG}(4, 1, 3, 1, 1; 1, 1, 1, 2, 1)$ , then all vertices in  $U_2$  and  $U_4$  are neutral vertices for  $\lambda_3 = 1$ .*

*If  $G = \text{NSG}(2, 4, 4, 2; 1, 1, 1, 2)$ , then all vertices in  $V_2$  and  $V_4$  are neutral for  $\lambda_{16} = -2$ .*

**Example 2.2.** *In  $G = \text{NSG}(2, 2, 5, 1; 1, 1, 1, 1)$  all vertices in  $U_3$  and in  $V_2$  are neutral vertices for  $\lambda_2 = 1$ .*

In what follows we assume that all vertices in  $U_s$  (resp.  $V_s$ ) of a nested split graph  $G$  are neutral for some  $s$  with respect to some  $\lambda_i \neq 0, -1$ . If so, we will show that this assumption imply some restrictions on position of  $\lambda_i$  in the spectrum of  $G$ .

**Theorem 2.4.** *Let  $G = \text{NSG}(m_1, \dots, m_h; n_1, \dots, n_h)$  such that all vertices in  $U_s$  for some  $2 \leq s \leq h-1$  are neutral for  $\lambda_i \neq 0, -1$ . If  $G' = \text{NSG}(m_{s+1}, \dots, m_h; n_{s+1}, \dots, n_h)$ ,  $n' = |V(G')|$  and  $\text{Spec}(G') = \{\lambda'_1, \dots, \lambda'_{n'}\}$  then  $\lambda_i = \lambda'_j$  for some  $j \in \{1, \dots, n'\}$ . Moreover,  $j < i < n - n' + j$ , and if  $i \leq n'$  then  $\lambda_i \neq \lambda'_{n'}$ .*

*Proof.* Let  $G'$  be the induced subgraph of  $G$  obtained by deleting all vertices in  $U_1, \dots, U_s, V_1, \dots, V_s$  i.e.

$$G' = \text{NSG}(m_{s+1}, \dots, m_h; n_{s+1}, \dots, n_h).$$

Let  $n' = |V(G')| = \sum_{j=s+1}^h (m_j + n_j)$  and let  $\mathbf{x}$  be a  $\lambda$ -eigenvector of  $G$ . Denote by  $\mathbf{x}'$  the vector obtained from  $\mathbf{x}$  by deleting all entries corresponding to deleted vertices from  $G$ . Since

$$0 = \lambda_i \mathbf{x}(u_s) = \sum_{j=1}^s n_j \mathbf{x}(v_j),$$

for any  $k \geq s+1$ , we obtain

$$\begin{aligned} \lambda_i \mathbf{x}'(u_k) &= \lambda_i \mathbf{x}(u_k) = \sum_{j=1}^k n_j \mathbf{x}(v_j) = \sum_{j=s+1}^k n_j \mathbf{x}(v_j) = \sum_{j=s+1}^k n_j \mathbf{x}'(v_j) \\ \lambda_i \mathbf{x}'(v_k) &= \lambda_i \mathbf{x}(v_k) = \sum_{j=1}^h n_j \mathbf{x}(v_j) - \mathbf{x}(v_k) + \sum_{j=k}^h m_j \mathbf{x}(u_j) \\ &= \sum_{j=s+1}^h n_j \mathbf{x}'(v_j) - m_j \mathbf{x}'(v_k) + \sum_{j=k}^h \mathbf{x}'(u_j) \end{aligned}$$

and therefore  $\mathbf{x}'$  is an eigenvector of  $G'$  for  $\lambda_i$ , i.e.  $\lambda_i \in \text{Spec}(G')$ . Suppose  $\lambda_i = \lambda'_j$  for some  $j \in \{1, \dots, n'\}$ . From interlacing it follows that

$$(6) \quad \lambda_{n-n'+i} \leq \lambda'_i \leq \lambda_i = \lambda'_j, \quad \text{if } i \leq n'.$$

as well as

$$(7) \quad \lambda_{n-n'+j} \leq \lambda_i = \lambda'_j \leq \lambda_j.$$

If in (6), at least one of inequalities holds as an equality then, by Lemma 1.1,  $G'$  has an eigenvector  $\mathbf{y}'$  for  $\lambda'_i$  such that  $\begin{pmatrix} \mathbf{0} \\ \mathbf{y}' \end{pmatrix}$  is an eigenvector of  $G$  for  $\lambda'_i$ . By the sum rule for any vertex in  $V_s$  we obtain that the sum of all entries of  $\mathbf{y}'$  is 0 and accordingly that  $\lambda'_i$  is non-main eigenvalue of  $G'$ . Hence,  $\lambda'_i = 0$  or  $\lambda'_i = -1$  which implies  $\lambda'_i < \lambda_i$ . Similarly, in (7) we conclude that  $\lambda'_j = \lambda_i$  is a non-main eigenvalue of  $G'$ , a contradiction, by Theorem 2.1. Therefore, the interlacing in these cases reads

$$(8) \quad \lambda_{n-n'+i} \leq \lambda'_i < \lambda_i, \quad i \leq n',$$

$$(9) \quad \lambda_{n-n'+j} < \lambda'_j = \lambda_i < \lambda_j.$$

Moreover, (9) implies  $j < i < n - n' + j$ . Also, if  $i \leq n'$ ,  $\lambda_i \neq \lambda'_{n'}$  holds. Otherwise,  $\lambda_{n-n'+i} \leq \lambda'_i < \lambda_i = \lambda'_{n'}$ , a contradiction.  $\square$

If all vertices in  $V_s$  for some  $s$  are neutral for  $\lambda_i \neq 0, -1$ , then bearing in mind that

$$G - V_s = \text{NSG}(m_1, \dots, m_{s-1} + m_s, \dots, m_h; n_1, \dots, n_{s-1}, n_{s+1}, \dots, n_h)$$

we can similarly conclude the following.

**Theorem 2.5.** *Let  $G = \text{NSG}(m_1, \dots, m_h; n_1, \dots, n_h)$  such that all vertices in  $V_s$  for some  $2 \leq s \leq h$  are neutral for  $\lambda_i \neq 0, -1$ . If*

$$H_s = \text{NSG}(m_1, \dots, m_{s-1} + m_s, \dots, m_h; n_1, \dots, n_{s-1}, n_{s+1}, \dots, n_h),$$

and  $\text{Spec}(H_s) = \{\lambda'_1, \dots, \lambda'_{n-n_s}\}$ , then  $\lambda_i = \lambda'_j$  for some  $j \in \{1, \dots, n - n_s\}$ . Moreover,  $j < i < n_s + j$  and if  $i \leq n - n_s$  then  $\lambda_i \neq \lambda'_{n-n_s}$ .

**Corollary 2.1.** *Let  $G = \text{NSG}(m_1, \dots, m_h; n_1, \dots, n_h)$  of order  $n$ . Then all vertices in  $V(G)$  are downer vertices for  $\lambda_n$ .*

*Proof.* If  $\lambda_n = -1$ , then  $G$  is a complete graph and all vertices are downers for it. So, we assume that  $\lambda_n \neq 0, -1$ . Suppose on the contrary that there exists at least one neutral vertex  $u$  for  $\lambda_n$ . If  $u \in U_s$ , then  $\mathbf{x}(u) = 0$ , where  $\mathbf{x}$  is a  $\lambda$ -eigenvector of  $G$ . As shown in the proof of Theorem 2.4,  $\lambda_n = \lambda'_j \in \text{Spec}(G')$ , for some  $j \in \{1, \dots, n'\}$  and  $\lambda_{n-n'+j} < \lambda'_j < \lambda_j$ , i.e.  $\lambda_{n-n'+j} < \lambda_n < \lambda_j$ , a contradiction.

The proof is similar if  $v \in V_s$  for some  $s$  and hence omitted here.  $\square$

**Theorem 2.6.** *Let  $G = \text{NSG}(m_1, \dots, m_h; n_1, \dots, n_h)$  such that all vertices in  $U_s$  are neutral for  $\lambda \neq 0, -1$  and  $G'' = \text{NSG}(m_1, \dots, m_s; n_1, \dots, n_s)$ . Then*

$$\lambda_{n''}(G'') < \lambda < \lambda_1(G''),$$

where  $n'' = |V(G'')| = \sum_{i=1}^s (m_i + n_i)$ .

*Proof.* The graph  $G''$  is an induced subgraph of  $G$  with vertex set  $V(G'') = \bigcup_{j=1}^s (U_j \cup V_j)$ . The adjacency matrix  $A$  of the whole graph is equal to:

$$\begin{bmatrix} A'' & B \\ B^T & A' \end{bmatrix},$$

where  $A', A''$  are adjacency matrices of

$$G' = \text{NSG}(m_{s+1}, \dots, m_h; n_{s+1}, \dots, n_h)$$

and  $G''$ , respectively, and

$$B = \begin{bmatrix} O_{M_s, n'} \\ J_{N_s, n'} \end{bmatrix},$$



where  $M_s = \sum_{j=1}^s m_j$ ,  $N_s = \sum_{j=1}^s n_j$  and  $n' = |V(G')|$ . The corresponding eigenvector  $\mathbf{x}$  can be represented as  $\mathbf{x} = \begin{pmatrix} \mathbf{x}_1 \\ \mathbf{x}_2 \end{pmatrix}$  and the eigenvalue system reads:

$$(10) \quad A''\mathbf{x}_1 + B\mathbf{x}_2 = \lambda\mathbf{x}_1$$

$$(11) \quad B^T\mathbf{x}_1 + A'\mathbf{x}_2 = \lambda\mathbf{x}_2.$$

As we have seen in the proof of Theorem 2.4,  $\lambda$  is an eigenvalue of  $A'$ , the corresponding eigenvector is  $\mathbf{x}_2$  and further  $\mathbf{x}_1 \neq 0$ . Therefore, it follows that  $B^T\mathbf{x}_1 = 0$ , i.e. the sum of some entries of  $\mathbf{x}_1$  is 0. From (10), we obtain

$$(\lambda I - A'')\mathbf{x}_1 = B\mathbf{x}_2$$

and then by multiplying by  $\mathbf{x}_1^T$  from the left we obtain

$$\mathbf{x}_1^T(\lambda I - A'')\mathbf{x}_1 = 0$$

and consequently

$$\min_{\mathbf{y} \neq 0} \frac{\mathbf{y}^T(\lambda I - A'')\mathbf{y}}{\mathbf{y}^T\mathbf{y}} \leq \frac{\mathbf{x}_1^T(\lambda I - A'')\mathbf{x}_1}{\mathbf{x}_1^T\mathbf{x}_1} \leq \max_{\mathbf{y} \neq 0} \frac{\mathbf{y}^T(\lambda I - A'')\mathbf{y}}{\mathbf{y}^T\mathbf{y}}.$$

Hence,

$$\lambda_{n''}(\lambda I - A'') \leq 0 \leq \lambda_1(\lambda I - A''),$$

where  $n'' = |V(G'')| = M_s + N_s$ . Since,  $\lambda_{n''}(\lambda I - A'') = \lambda - \lambda_1(G'')$  and  $\lambda_1(\lambda I - A'') = \lambda - \lambda_{n''}(G'')$  it follows

$$(12) \quad \lambda_{n''}(G'') \leq \lambda \leq \lambda_1(G'').$$

Moreover,  $\lambda \neq \lambda_1(G'')$ . Equality holds if and only if  $\mathbf{x}_1$  is an eigenvector of  $G''$  for  $\lambda_1(G'')$ , that is not possible due to the condition (11) and positivity of  $\mathbf{x}_1$  as an eigenvector corresponding to the largest eigenvalue of a connected graph. Similarly, if  $\lambda = \lambda_{n''}(G'')$ , then  $\mathbf{x}_1$  is the corresponding eigenvector and from (10) it follows  $B\mathbf{x}_2 = 0$ . This implies that  $\lambda$  is a non-main eigenvalue of a nested split graph  $G'$ , a contradiction by Theorem 2.1.  $\square$

**Corollary 2.2.** *Let*

$$\begin{aligned} G &= \text{NSG}(m_1, \dots, m_h; n_1, \dots, n_h), \\ G_s'' &= \text{NSG}(m_1, \dots, m_s; n_1, \dots, n_s), \end{aligned}$$

$I_s = (\lambda_{n_s''}(G_s''), \lambda_1(G_s''))$ , where  $n_s'' = |V(G_s'')|$  and  $\lambda \in \text{Spec}(G)$ . If  $\lambda \notin \bigcup_{s=2}^{h-1} I_s$ , then all vertices in  $U$  are downer vertices for  $\lambda$ .

**Example 2.3.** *Let  $G = \text{NSG}(1, 1, 5; 1, 1, 8)$ . Then  $I_2 = (-1.48, 2.17)$  and besides  $\lambda_1$  and  $\lambda_n$  all vertices in  $U$  are downer for  $\lambda_{n-2}$  and  $\lambda_{n-1}$ , as well.*

## 3. VERTEX TYPES IN CHAIN GRAPHS

Chain graph can be defined as follows: a graph is a chain graph if and only if it is bipartite and the neighborhoods of the vertices in each color class form a chain with respect to inclusion. For this reason, if connected (as was the case with threshold graphs), it is also called *double nested graph* [5].

Non-zero eigenvalues of chain graphs are simple (see Theorem 3.1 below). As the subgraphs of any chain graph are also chain graphs, it follows that there is no Parter vertex in any chain graphs with respect to non-zero eigenvalues. A question raises whether they can have neutral vertices. In [1] it is conjectured that this cannot be the case.

**Conjecture 3.1.** ([1]) *In any chain graph, every vertex is downer with respect to every non-zero eigenvalue.*

We disprove Conjecture 3.1 in this section. Indeed, Theorems 3.4 and 3.5 will show that there are infinitely many counterexamples for this conjecture. In spite of that, a couple of weak versions of the conjecture are true.

**Remark 3.1.** (Structure of chain graphs) *As it was observed in [5], the color classes of any chain graph  $G$  can be partitioned into  $h$  non-empty cells  $U_1, \dots, U_h$  and  $V_1, \dots, V_h$  such that  $N(u) = V_1 \cup \dots \cup V_{h-i+1}$  for any  $u \in U_i$ ,  $1 \leq i \leq h$ . If  $m_i = |U_i|$  and  $n_i = |V_i|$ , then we write  $\text{DNG}(m_1, \dots, m_h; n_1, \dots, n_h)$  (see Fig. 2).*




FIGURE 2. The chain graph  $G = \text{DNG}(m_1, \dots, m_h; n_1, \dots, n_h)$ .

The spectrum of any chain graph has the following properties (see [1]):

**Theorem 3.1.** *Let  $G = \text{DNG}(m_1, \dots, m_h; n_1, \dots, n_h)$ . Then the spectrum of  $G$  is symmetric about the origin and it contains:*

- $h$  positive simple eigenvalues greater than  $\frac{1}{2}$ ;
- $h$  negative simple eigenvalues less than  $-\frac{1}{2}$ ;
- eigenvalue 0 of multiplicity  $M_h + N_h - 2h$ .

**Remark 3.2.** *On the contrary from threshold graphs nonzero eigenvalues of chain graphs need not be main. For more information see [3].*

**Theorem 3.2.** *Let  $G = \text{DNG}(m_1, \dots, m_h; n_1, \dots, n_h)$  be a chain graph. Then the vertices in  $U_1 \cup U_h \cup V_1 \cup V_h$  are downer for any non-zero eigenvalue.*

*Proof.* Let  $\mathbf{x}$  be any  $\lambda$ -eigenvector of  $G$ . Assume that  $u_1 \in U_1$  and  $v_h \in V_h$ . By the sum rule  $\lambda \mathbf{x}(v_h) = m_1 \mathbf{x}(u_1)$ . Since,  $\lambda \neq 0$ ,  $u_1$  and  $v_h$  are both downer or neutral. Let  $X = \sum_{w \in V} \mathbf{x}(w)$  and assume on the contrary that  $\mathbf{x}(u_1) = \mathbf{x}(v_h) = 0$ . Again, by the sum rule  $\lambda \mathbf{x}(u_2) = X - n_h \mathbf{x}(v_h) = 0$  and consequently  $\mathbf{x}(u_2) = 0$ , for any  $u_2 \in U_2$  as well as  $\mathbf{x}(v_{h-1}) = 0$  for any  $v_{h-1} \in V_{h-1}$ . Next, for any  $u_3 \in U_3$ ,

$$\lambda \mathbf{x}(u_3) = X - n_{h-1} \mathbf{x}(v_{h-1}) - n_h \mathbf{x}(v_h) = 0$$

It follows that  $\mathbf{x}$  is zero on  $U_3$ , too. Continuing this argument, it follows that  $\mathbf{x} = \mathbf{0}$ , a contradiction.  $\square$

The following proposition states some facts related to vertex types in chain graphs. The proofs are similar to those in Section 2 and therefore omitted here.

**Theorem 3.3.** *Let*

$$\begin{aligned} G &= \text{DNG}(m_1, \dots, m_h; n_1, \dots, n_h), \\ G'_s &= \text{DNG}(m_1, \dots, m_{s-1}; n_{h-s+2}, \dots, n_h), \\ G''_s &= \text{DNG}(m_s, \dots, m_h; n_1, \dots, n_{h-s+1}), \end{aligned}$$

$1 < s < h$ ,  $\lambda_i \in \text{Spec}(G) \setminus \{0\}$ ,  $n'_s = \sum_{j=1}^{s-1} (m_j + n_{h-j+1})$ ,  $n''_s = n - n'_s$ ,  $\text{Spec}(G'_s) = \{\lambda'_1, \dots, \lambda'_{n'_s}\}$ . Then

- For any  $j = 1, \dots, h-1$  at least one of  $U_j, U_{j+1}$  contains only downer vertices for  $\lambda_i$ .
- If all vertices in  $U_s$  for some  $2 < s < h-1$  are neutral for  $\lambda_i$ , then
  - $\lambda_i$  is an eigenvalue of  $G'_s$  and  $\lambda_i = \lambda'_j$ , for some  $j \in \{1, \dots, n'_s\}$ . If  $\lambda_i$  is main, then  $j < i < n - n'_s + j$ . If  $i \leq n'_s$  then  $\lambda_i \neq \lambda'_{n'_s}$ .
  - $\lambda_i \in [\lambda_{n'_s}''(G''_s), \lambda_1(G''_s)]$ .
  - If  $\lambda_i$  is a main eigenvalue then  $\lambda_i \in (\lambda_{n'_s}''(G''_s), \lambda_1(G''_s))$ .
  - If  $\lambda_i \notin \bigcup_{s=2}^{h-1} [\lambda_{n'_s}''(G''_s), \lambda_1(G''_s)]$  then all vertices in  $V(G)$  are downer vertices for  $\lambda_i$ .

A chain graph for which  $|U_1| = \dots = |U_h| = |V_1| = \dots = |V_h| = 1$  is called a *half graph*. Here we denote it by  $H(h)$ . As we will see in what follows, specific half graphs provide counterexamples to Conjecture 3.1. Let

$$(a_1, \dots, a_6) := (1, 0, -1, -1, 0, 1).$$

In what follows, for convenience, we will instead of column vectors use row vectors, especially for eigenvectors.

Let

$$\mathbf{x} := (x_1, \dots, x_h)$$

where  $x_i = a_s$  if  $i \equiv s \pmod{6}$ . In the next theorem, we show that the vector  $(\mathbf{x}, \mathbf{x})$  (each  $\mathbf{x}$  corresponds to a color class) is an eigenvector of a non-zero eigenvalue of  $H(h)$  for some  $h$ . In view of Remark 1.2, this disproves Conjecture 3.1.

**Theorem 3.4.** *In any half graph  $H(h)$ , the vector  $(\mathbf{x}, \mathbf{x})$  is an eigenvector for  $\lambda = 1$  if  $h \equiv 1 \pmod{6}$  and it is an eigenvector for  $\lambda = -1$  if  $h \equiv 4 \pmod{6}$ .*

*Proof.* From Table 1, we observe that for  $1 \leq s \leq 6$ ,

$$\sum_{i=1}^{5-s} a_i = -a_s \text{ and } \sum_{i=1}^{2-s} a_i = a_s,$$

where we consider  $5-s$  and  $2-s$  modulo 6 as elements of  $\{1, \dots, 6\}$ .

$s$	$a_s$	$5-s$	$\sum_{i=1}^{5-s} a_i$	$2-s$	$\sum_{i=1}^{2-s} a_i$
1	1	4	-1	1	1
2	0	3	0	6	0
3	-1	2	1	5	-1
4	-1	1	1	4	-1
5	0	6	0	3	0
6	1	5	-1	2	1

TABLE 1. The values of  $\sum_{i=1}^{5-s} a_i$  and  $\sum_{i=1}^{2-s} a_i$

Note that, since  $\sum_{i=1}^6 a_i = 0$ , if  $1 \leq \ell \leq h$ ,  $1 \leq s \leq 6$  and  $\ell \equiv s \pmod{6}$ , then

$$\sum_{i=1}^{\ell} x_i = \sum_{i=1}^s a_i.$$

Let  $\{u_1, \dots, u_h\}$  and  $\{v_1, \dots, v_h\}$  be the color classes of  $H(h)$ . Let  $h = 6t + 4$ . We show that  $(\mathbf{x}, \mathbf{x})$  satisfies the sum rule for  $\lambda = -1$ . By the symmetry, we only need to show this for  $u_i$ 's. Let  $i = 6t' + s$  for

some  $1 \leq s \leq 6$ . Then  $n - i + 1 = 6(t - t') + 5 - s$ .

$$\sum_{j: v_j \sim u_i} x_j = \sum_{j=1}^{n-i+1} x_j = \sum_{j=1}^{5-s} a_j = -a_s = -x_i.$$

Now, let  $h = 6t + 1$ . We show that in this case  $(\mathbf{x}, \mathbf{x})$  satisfies the sum rule for  $\lambda = 1$ . Let  $i = 6t' + s$  for some  $1 \leq s \leq 6$ . Then  $n - i + 1 = 6(t - t') + 2 - s$ .

$$\sum_{j: v_j \sim u_i} x_j = \sum_{j=1}^{n-i+1} x_j = \sum_{j=1}^{2-s} a_j = a_s = x_i.$$

□

Now we give another class of counterexamples to Conjecture 3.1. For this, let

$$\omega^2 + \omega - 1 = 0,$$

and

$$(b_1, \dots, b_{10}) := (\omega, -1, 0, 1, -\omega, -\omega, 1, 0, -1, \omega).$$

Let

$$\mathbf{x} := (x_1, \dots, x_h)$$

where  $x_i = b_s$  if  $i \equiv s \pmod{10}$ .

**Theorem 3.5.** *In any half graph  $H(h)$ , the vector  $(\mathbf{x}, \mathbf{x})$  is an eigenvector for  $\lambda = \omega$  if  $h \equiv 7 \pmod{10}$  and it is an eigenvector for  $\lambda = -\omega$  if  $h \equiv 2 \pmod{10}$ .*

*Proof.* From Table 2, we observe that for  $1 \leq s \leq 10$ ,

$$\sum_{i=1}^{8-s} b_i = \omega b_s \text{ and } \sum_{i=1}^{3-s} b_i = -\omega b_s,$$

where we consider  $8 - s$  and  $3 - s$  modulo 10 as elements of  $\{1, \dots, 10\}$ .

$s$	$b_s$	$8 - s$	$\sum_{i=1}^{8-s} b_i$	$3 - s$	$\sum_{i=1}^{3-s} b_i$
1	$\omega$	7	$1 - \omega$	2	$\omega - 1$
2	-1	6	$-\omega$	1	$\omega$
3	0	5	0	10	0
4	1	4	$\omega$	9	$-\omega$
5	$-\omega$	3	$\omega - 1$	8	$1 - \omega$
6	$-\omega$	2	$\omega - 1$	7	$1 - \omega$
7	1	1	$\omega$	6	$-\omega$
8	0	10	0	5	0
9	-1	9	$-\omega$	4	$\omega$
10	$\omega$	8	$1 - \omega$	3	$\omega - 1$

TABLE 2. The values of  $\sum_{i=1}^{8-s} b_i$  and  $\sum_{i=1}^{3-s} b_i$

Note that, since  $\sum_{i=1}^{10} b_i = 0$ , if  $1 \leq \ell \leq k$ ,  $1 \leq s \leq 10$  and  $\ell \equiv s \pmod{10}$ , then

$$\sum_{i=1}^{\ell} x_i = \sum_{i=1}^s b_i.$$

Let  $k = 10t + 7$ . Then  $(\mathbf{x}, \mathbf{x})$  satisfies the sum rule for  $\lambda = \omega$ . Let  $i = 10t' + s$  for some  $1 \leq s \leq 10$ . Then  $n - i + 1 = 10(t - t') + 8 - s$ .

$$\sum_{j: v_j \sim u_i} x_j = \sum_{j=0}^{n-i+1} x_j = \sum_{j=1}^{8-s} b_j = \omega b_s = \omega x_i.$$

Now, let  $h = 10t + 2$ . Assume that  $i = 10t' + s$  for some  $1 \leq s \leq 10$ . Then  $n - i + 1 = 6(t - t') + 3 - s$ .

$$\sum_{j: v_j \sim u_i} x_j = \sum_{j=1}^{n-i+1} x_j = \sum_{j=1}^{3-s} b_j = -\omega b_s = -\omega x_i.$$

It follows that in this case  $(\mathbf{x}, \mathbf{x})$  satisfies the sum rule for  $\lambda = -\omega$ .  $\square$

**Remark 3.3.** *The following two facts deserve to be mentioned:*

(i) *Given  $(\mathbf{x}, \mathbf{x})$  as eigenvector of  $H(h)$  for  $\lambda \in \{\pm 1, \pm \omega\}$ , then  $(\mathbf{x}, -\mathbf{x})$  is an eigenvector of  $H(h)$  for  $-\lambda$ . This gives more eigenvalues of  $H(h)$  with eigenvectors containing zero components.*

(ii) *Let  $\mathbf{x}$  be an eigenvector for eigenvalue  $\lambda \neq 0$  of a graph  $G$  with  $\mathbf{x}_v = 0$ , for some vertex  $v$ . If we add a new vertex  $u$  with  $N(u) = N(v)$  and add a zero component to  $\mathbf{x}$  corresponding to  $u$ , then the new vector is an eigenvector of  $H$  for  $\lambda$ . So, we can extend any graph presented in Theorems 3.4 or 3.5 to construct infinitely many more counterexamples for Conjecture 3.1.*

#### ACKNOWLEDGMENTS

The research of the second author was in part supported by a grant from IPM.

#### REFERENCES

- [1] A. Alazemi, M. Anđelić, S.K. Simić, Eigenvalue location for chain graphs, *Linear Algebra Appl.* **505** (2016), 194–210.
- [2] M. Anđelić, S.K. Simić, Some notes on the threshold graphs, *Discrete Math.* **310** (2010), 2241–2248.
- [3] M. Anđelić, E. Andrade, D.M. Cardoso, C.M. da Fonseca, S.K. Simić, D.V. Tošić, Some new considerations about double nested graphs, *Linear Algebra Appl.* **483** (2015), 323–341.
- [4] M. Anđelić, F. Ashraf, C.M. da Fonseca, S.K. Simić, Vertex types in some lexicographic products of graphs, submitted.
- [5] F.K. Bell, D. Cvetković, P. Rowlinson, S.K. Simić, Graphs for which the least eigenvalue is minimal, II, *Linear Algebra Appl.* **429** (2008), 2168–2179.

- [6] A. Bhattacharya, S. Friedland, U.N. Peled, On the first eigenvalue of bipartite graphs, *Electron. J. Combin.* **15** (2008), #R144.
- [7] A. Brandstädt, V.B. Le, J.P. Spinrad, *Graph Classes: A Survey*, Society for Industrial and Applied Mathematics (SIAM), Philadelphia, PA, 1999.
- [8] A.E. Brouwer, W.H. Haemers, *Spectra of Graphs*, Springer, New York, 2012.
- [9] D.M. Cardoso, M.A.A. Freitas, E.A. Martins, M. Robbiano, Spectra of graphs obtained by a generalization of the join graph operation, *Discrete Math.* **313** (2013), 733–741.
- [10] D. Cvetković, P. Rowlinson, S. Simić, *An Introduction to the Theory of Graph Spectra*, London Mathematical Society Student Texts, 75, Cambridge University Press, Cambridge, 2010.
- [11] C.D. Godsil, B.D. McKay, A new graph product and its spectrum, *Bull. Austral. Math. Soc.* **18(1)** (1978), 21–28.
- [12] P.L. Hammer, U.N. Peled, X. Sun, Difference graphs, *Discrete Appl. Math.* **28** (1990), 35–44.
- [13] F. Harary, The structure of threshold graphs, *Riv. Mat. Sci. Econom. Social.* **2** (1979), 169–172.
- [14] R.A. Horn, C.R. Johnson, *Matrix Analysis*, Cambridge University Press, 2013.
- [15] D. Jacobs, V. Trevisan, F. Tura, Eigenvalues and energy in threshold graphs, *Linear Algebra Appl.* **465** (2015), 412–425.
- [16] P. Manca, On a simple characterisation of threshold graphs, *Riv. Mat. Sci. Econom. Social.* **2** (1979), 3–8.
- [17] N.V.R. Mahadev, U.N. Peled, *Threshold Graphs and Related Topics*, Annals of Discrete Mathematics, NorthHolland Publishing Co., Amsterdam, 1995.
- [18] I. Sciriha, S. Farrugia, On the Spectrum of Threshold Graphs, *ISRN Discrete Mathematics*, **2011** (2011), Article ID 108509, 21 pages.
- [19] S.K.Simić, M. Anđelić, C.M. da Fonseca, D. Živković, On the multiplicities of eigenvalues of graphs and their vertex deleted subgraphs: old and new results, *Electron. J. Linear Algebra* **30** (2015), 85–105.

DEPARTMENT OF MATHEMATICS, KUWAIT UNIVERSITY, SAFAT 13060, KUWAIT  
*E-mail address:* milica@sci.kuniv.edu.kw

DEPARTMENT OF MATHEMATICS, K. N. TOOSI UNIVERSITY OF TECHNOLOGY, P. O. BOX 16315-1618, TEHRAN, IRAN, AND SCHOOL OF MATHEMATICS, INSTITUTE FOR RESEARCH IN FUNDAMENTAL SCIENCES (IPM), P. O. BOX 19395-5746, TEHRAN, IRAN  
*E-mail address:* e\_ghorbani@ipm.ir

STATE UNIVERSITY OF NOVI PAZAR, VUKA KARADŽIĆA BB, 36 300 NOVI PAZAR, SERBIA, AND MATHEMATICAL INSTITUTE SANU, KNEZA MIHAILA 36, 11 000 BELGRADE, SERBIA  
*E-mail address:* sksimic@mi.sanu.ac.rs