

# ROUNDING THE ARITHMETIC MEAN VALUE OF THE SQUARE ROOTS OF THE FIRST $n$ INTEGERS

THOMAS P. WIHLER

ABSTRACT. In this article we study the arithmetic mean value  $\Sigma(n)$  of the square roots of the first  $n$  integers. For this quantity, we develop an asymptotic expression, and derive a formula for its integer part which has been conjectured recently in the work of M. Merca. Furthermore, we address the numerical evaluation of  $\Sigma(n)$  for large  $n \gg 1$ .

The aim of this article is to derive an explicit formula for the integer part of the arithmetic mean value of the square roots of the first  $n$  integers. More precisely, we consider the sequence

$$\Sigma(n) = \frac{1}{n} \sum_{k=1}^n \sqrt{k}, \quad n \in \mathbb{N},$$

and show the following identity.

**Theorem 1.** *For any  $n \in \mathbb{N}$ , there holds that*

$$(1) \quad \lfloor \Sigma(n) \rfloor = \lfloor A(n) \rfloor,$$

where we define the function

$$(2) \quad A(x) = \frac{2}{3} \sqrt{x+1} \left( 1 + \frac{1}{4x} \right),$$

for  $x \geq 1$ . Here,  $\lfloor \cdot \rfloor$  signifies the integer part of a positive real number.

This result is motivated by the recent work [Mer17, see Conjecture 2], where Theorem 1 has been conjectured.

## 1. AN ASYMPTOTIC RESULT

In order to prove Theorem 1, we begin by deriving an asymptotic result for the sum of the square roots of the first  $n$  integers. Here, we employ an idea presented in [Mer17], which is based on using the trapezium rule for the numerical approximation of integrals. In this context, we also point to the related work [She13], where upper and lower Riemann sums have been applied. In comparison to the analysis pursued in [Mer17], in the current paper, we use a different approach to control the error in the trapezium rule. Thereby, we arrive at a slightly sharper asymptotic representation for large  $n$ . Incidentally, an asymptotic representation has been derived already in the early work [Ram00].

**Theorem 2.** *For any  $\nu, n \in \mathbb{N}$ , with  $\nu < n$ , there holds*

$$\sum_{k=\nu}^n \sqrt{k} = nA(n) - \frac{2}{3} \sqrt{\nu} \left( \nu - \frac{3}{4} \right) - \frac{\delta_{\nu,n}}{24},$$

with

$$(3) \quad \sigma(\nu+2, n+2) < \delta_{\nu,n} < \sigma(\nu, n),$$

where

$$\sigma(\nu, n) = \begin{cases} 3/2 - n^{-1/2} & \text{if } \nu = 1, \\ (\nu - 1)^{-1/2} - n^{-1/2} & \text{if } \nu \geq 2. \end{cases}$$

*Proof.* Let us consider the function  $f(x) = \sqrt{x}$ , for  $x \geq 1$ . We interpolate it by a piecewise linear function  $\ell$  in the points  $x = 1, 2, \dots$ , i.e., for any  $k \in \mathbb{N}$  there holds  $\ell(k) = \sqrt{k}$ , and  $\ell$  is a linear polynomial on the interval  $[k, k+1]$ . We define the remainder term

$$\widehat{\delta}_{\nu, n} := \int_{\nu}^{n+1} (\sqrt{x} - \ell(x)) \, dx = \frac{2}{3} \left( (n+1)^{3/2} - \nu^{3/2} \right) - \int_{\nu}^{n+1} \ell(x) \, dx.$$

Then, we note that

$$\int_{\nu}^{n+1} \ell(x) \, dx = \frac{1}{2} \sqrt{\nu} + \sum_{k=\nu+1}^n \sqrt{k} + \frac{1}{2} \sqrt{n+1};$$

this is the trapezium rule for the numerical integration of  $f$ . Therefore,

$$\begin{aligned} \sum_{k=\nu+1}^n \sqrt{k} &= \frac{2}{3} \left( (n+1)^{3/2} - \nu^{3/2} \right) - \frac{1}{2} \sqrt{\nu} - \frac{1}{2} \sqrt{n+1} - \widehat{\delta}_{\nu, n} \\ &= \frac{2}{3} \sqrt{n+1} \left( n + \frac{1}{4} \right) - \frac{2}{3} \sqrt{\nu} \left( \nu + \frac{3}{4} \right) - \widehat{\delta}_{\nu, n}. \end{aligned}$$

Hence,

$$\begin{aligned} \sum_{k=\nu}^n \sqrt{k} &= \frac{2}{3} \sqrt{n+1} \left( n + \frac{1}{4} \right) - \frac{2}{3} \sqrt{\nu} \left( \nu - \frac{3}{4} \right) - \widehat{\delta}_{\nu, n} \\ &= nA(n) - \frac{2}{3} \sqrt{\nu} \left( \nu - \frac{3}{4} \right) - \widehat{\delta}_{\nu, n}. \end{aligned}$$

It remains to study the error term  $\widehat{\delta}_{\nu, n}$ . For this purpose, applying twofold integration by parts, for  $k \in \mathbb{N}$ , we note that

$$\begin{aligned} \int_k^{k+1} (\sqrt{x} - \ell(x)) \, dx &= -\frac{1}{8} \int_k^{k+1} (\sqrt{x} - \ell(x)) \frac{d^2}{dx^2} (1 - 4(x - k - 1/2)^2) \, dx \\ &= \frac{1}{32} \int_k^{k+1} x^{-3/2} (1 - 4(x - k - 1/2)^2) \, dx \\ &< \frac{1}{32} k^{-3/2} \int_k^{k+1} (1 - 4(x - k - 1/2)^2) \, dx \\ &= \frac{1}{48} k^{-3/2}. \end{aligned}$$

Thus, if  $\nu \geq 2$ , we obtain

$$\begin{aligned} \widehat{\delta}_{\nu, n} &= \sum_{k=\nu}^n \int_k^{k+1} (\sqrt{x} - \ell(x)) \, dx \\ &< \frac{1}{48} \sum_{k=\nu}^n k^{-3/2} \\ &< \frac{1}{48} \int_{\nu-1}^n x^{-3/2} \, dx \\ &= \frac{1}{24} \left( (\nu - 1)^{-1/2} - n^{-1/2} \right). \end{aligned}$$

Otherwise, if  $\nu = 1$ , the above bound implies

$$\widehat{\delta}_{1,n} < \frac{1}{48} + \widehat{\delta}_{2,n} < \frac{1}{24} \left( \frac{3}{2} - n^{-1/2} \right).$$

Similarly, we have

$$\begin{aligned} \widehat{\delta}_{\nu,n} &> \frac{1}{48} \sum_{k=\nu}^n (k+1)^{-3/2} \\ &> \frac{1}{48} \int_{\nu+1}^{n+2} x^{-3/2} dx \\ &= \frac{1}{24} \left( (\nu+1)^{-1/2} - (n+2)^{-1/2} \right). \end{aligned}$$

This completes the proof.  $\square$

For  $\nu = 1$  the above result implies the identity

$$(4) \quad \Sigma(n) = A(n) - \frac{1}{6n} - \frac{\delta_{1,n}}{24n},$$

which will be crucial in the analysis below.

**Remark 1.** Proceeding in the same way as in the proof of Theorem 2, a formula for the more general case of the arithmetic mean value of the  $r$ -th roots of the first  $n$  integers, with  $r \geq 1$ , can be derived: More precisely, for any  $\nu, n \in \mathbb{N}$ , with  $\nu < n$ , there holds

$$\sum_{k=\nu}^n k^{1/r} = \frac{r}{r+1} (n+1)^{1/r} \left( n + \frac{1-1/r}{2} \right) - \frac{r}{r+1} \nu^{1/r} \left( \nu - \frac{1+1/r}{2} \right) - \frac{\delta_{\nu,n,r}}{12r},$$

with  $\delta_{\nu,n,1} = 0$  (i.e., for  $r = 1$ ), and  $\sigma_r(\nu+2, n+2) < \delta_{\nu,n,r} < \sigma_r(\nu, n)$  for  $r > 1$ , where

$$\sigma_r(\nu, n) = \begin{cases} 2 - 1/r - n^{-1+1/r} & \text{if } \nu = 1, \\ (\nu-1)^{-1+1/r} - n^{-1+1/r} & \text{if } \nu \geq 2. \end{cases}$$

## 2. PROOF OF THEOREM 1

The proof of Theorem 1 is based on the ensuing two auxiliary results. The first lemma provides tight upper and lower bounds on  $A$  from (2). The purpose of the second lemma is to identify any points where the integer part of  $A$  changes.

**Lemma 1.** *The function  $A$  from (2) is strictly monotone increasing for  $x \geq 1$ . Furthermore, there hold the bounds*

$$(5) \quad A(x) < \frac{2}{3} \sqrt{x+2} \quad \text{for } x \geq 2,$$

and

$$(6) \quad A(x) > \frac{2}{3} \sqrt{x + \frac{5}{4}} + \frac{1}{4x} \quad \text{for } x \geq 6.$$

*Proof.* The strict monotonicity of  $A$  follows directly from the fact that

$$A'(x) = \frac{4x^2 - x - 2}{12x^2 \sqrt{x+1}} > 0,$$

for any  $x \geq 1$ . Furthermore, we notice that the graph of  $A$  and of the function

$$h(x) = \frac{2}{3} \sqrt{x+2},$$

which is the upper bound in (5), have exactly one positive intersection point at  $x^* = (9 + \sqrt{113})/16 < 2$ . Moreover, choosing  $x = 2 > x^*$ , for instance, we have

$$A(2) = \frac{3\sqrt{3}}{4} < \frac{4}{3} = h(2).$$

Thus, we conclude that  $A(x) < h(x)$  for any  $x > x^*$ ; this yields (5). The lower bound (6) follows from an analogous argument.  $\square$

**Lemma 2.** *For any  $m \in \mathbb{N}_0$ , let  $\alpha(m) = 9/4(m+1)^2 - 2$ . Then, for  $n, m \in \mathbb{N}$ , with  $n \leq \alpha(m)$ , there holds that  $A(n) < m+1$ . Conversely, we have  $A(n) - 1/4n > m+1$ , for any integer  $n > \alpha(m)$ .*

*Proof.* Consider  $m, n \in \mathbb{N}$  such that  $1 \leq n \leq \alpha(m)$ . Applying the monotonicity of  $A$ , cf. Lemma 1, together with (5), we infer

$$A(n) \leq A(\alpha(m)) < \frac{2}{3}\sqrt{\alpha(m)+2} = m+1.$$

Furthermore, if  $m = 2s$ , with  $s \in \mathbb{N}$ , is even, then we have  $\alpha(2s) = 9s^2 + 9s + 1/4$ ; moreover, if  $m = 2s-1$ , with  $s \in \mathbb{N}$ , is odd, then there holds  $\alpha(2s-1) = 9s^2 - 2$ . In particular, we conclude that  $n \geq \alpha(m) + 3/4$  for any  $n \in \mathbb{N}$  with  $n > \alpha(m)$ . Then, involving (6), it follows that

$$A(n) - \frac{1}{4n} > \frac{2}{3}\sqrt{n + \frac{5}{4}} \geq \frac{2}{3}\sqrt{\alpha(m)+2} \geq m+1,$$

which yields the lemma.  $\square$

**Proof of Theorem 1.** We are now ready to prove the identity (1). To this end, given  $n \in \mathbb{N}$ , we define

$$m := \min\{k \in \mathbb{N} : \alpha(k) \geq n\} \in \mathbb{N}.$$

Evidently, there holds  $n \leq \alpha(m)$  as well as  $n > \alpha(m-1)$ . By virtue of (4) and due to Lemma 2, there holds

$$1 \leq \Sigma(n) < A(n) < m+1.$$

If  $m = 1$ , the proof of the theorem is complete. Otherwise, if  $m \geq 2$ , then by means of Theorem 2, we notice that  $\delta_{1,n} < 3/2$ . Then, recalling (4), this leads to

$$\Sigma(n) > A(n) - \frac{1}{6n} - \frac{1}{16n} = A(n) - \frac{11}{48n}.$$

Hence, upon employing Lemma 2 (with  $n > \alpha(m-1)$ ), it follows that

$$\Sigma(n) > A(n) - \frac{1}{4n} > m.$$

Combining the above estimates, we deduce the bounds

$$m < \Sigma(n) < A(n) < m+1.$$

This shows (1).

### 3. NUMERICAL EVALUATION OF $\Sigma(n)$

For large values of  $n$  the straightforward computation of  $\Sigma(n)$ , i.e., simply adding the numbers  $\sqrt{1}, \sqrt{2}, \sqrt{3}, \dots, \sqrt{n}$ , and dividing by  $n$ , is computationally slow and prone to roundoff errors. For this reason, we propose an alternative approach: if  $n \gg 1$ , we choose  $\nu \in \mathbb{N}$ , with  $\nu+1 < n$ , of moderate size (so that the numerical evaluation of  $\Sigma(\nu)$  is well-conditioned and accurate). Then, we write

$$\Sigma(n) = \frac{1}{n} \left( \nu \Sigma(\nu) + \sum_{k=\nu+1}^n \sqrt{k} \right).$$

Here, employing Theorem 2, we notice that

$$\sum_{k=\nu+1}^n \sqrt{k} = nA(n) - \nu A(\nu) - \frac{\delta_{\nu+1,n}}{24}.$$

Hence, upon defining the approximation

$$\tilde{\Sigma}(\nu, n) := \frac{1}{n} (nA(n) + \nu\Sigma(\nu) - \nu A(\nu)),$$

it follows that

$$\left| \Sigma(n) - \tilde{\Sigma}(\nu, n) \right| \leq \frac{\delta_{\nu+1,n}}{24n},$$

with

$$\delta_{\nu+1,n} < \sigma(\nu+1, n) = \nu^{-1/2} - n^{-1/2} = n^{-1/2} \left( \left( \frac{\nu}{n} \right)^{-1/2} - 1 \right);$$

cf. (3). In this way, we infer the error estimate

$$(7) \quad \left| \Sigma(n) - \tilde{\Sigma}(\nu, n) \right| < \frac{n^{-3/2}}{24} \left( \left( \frac{\nu}{n} \right)^{-1/2} - 1 \right).$$

In particular, given a prescribed tolerance  $\epsilon > 0$ , we require

$$\nu = \left\lceil n \left( 24\epsilon n^{3/2} + 1 \right)^{-2} \right\rceil,$$

with  $\nu \leq n - 2$ , in order to deduce the guaranteed bound

$$\left| \Sigma(n) - \tilde{\Sigma}(\nu, n) \right| < \epsilon.$$

To give an example, we consider  $n = 10^7$ . Then, choosing  $\nu = 100$  leads to the numerical value

$$\tilde{\Sigma}(n) = 2108.1852648724285 \dots$$

In this particular case, the error bound (7) gives

$$\left| \Sigma(n) - \tilde{\Sigma}(\nu, n) \right| \leq 4.1535 \times 10^{-10}.$$

This estimate is fairly sharp; indeed, the true error is approximately  $4.1328 \times 10^{-10}$ .

#### REFERENCES

- [Mer17] M. Merca, *On the arithmetic mean of the square roots of the first  $n$  positive integers*, *College Math. J.* **48** (2017), no. 2, 129–133.
- [Ram00] S. Ramanujan, *On the sum of the square roots of the first  $n$  natural numbers* [*J. Indian Math. Soc.* **7** (1915), 173–175], *Collected papers of Srinivasa Ramanujan*, AMS Chelsea Publ., Providence, RI, 2000, pp. 47–49.
- [She13] S. Shekatkar, *The sum of the  $r$ th root of first  $n$  natural numbers and new formula for factorial*, Tech. report, <https://arxiv.org/abs/1204.0877>, 2013.