

BOUNDING χ BY A FRACTION OF Δ FOR GRAPHS WITHOUT LARGE CLIQUES

MARTHE BONAMY, TOM KELLY, PETER NELSON, AND LUKE POSTLE

ABSTRACT. The greedy coloring algorithm shows that a graph of maximum degree at most Δ has chromatic number at most $\Delta + 1$, and this is tight for cliques. Much attention has been devoted to improving this “greedy bound” for graphs without large cliques. Brooks famously proved that this bound can be improved by one if $\Delta \geq 3$ and the graph contains no clique of size $\Delta + 1$. Reed’s Conjecture states that the “greedy bound” can be improved by k if the graph contains no clique of size $\Delta + 1 - 2k$. Johansson proved that the “greedy bound” can be improved by a factor of $\Omega(\ln(\Delta)^{-1})$ or $\Omega\left(\frac{\ln(\ln(\Delta))}{\ln(\Delta)}\right)$ for graphs with no triangles or no cliques of any fixed size, respectively.

Notably missing is a *linear* improvement on the “greedy bound” for graphs without large cliques. In this paper, we prove that for sufficiently large Δ , if G is a graph with maximum degree at most Δ and no clique of size ω , then

$$\chi(G) \leq 72\Delta \sqrt{\frac{\ln(\omega)}{\ln(\Delta)}}.$$

This implies that for sufficiently large Δ , if $\omega^{(72c)^2} \leq \Delta$ then $\chi(G) \leq \Delta/c$.

This bound actually holds for the list-chromatic and even the correspondence chromatic number (also known as DP-chromatic number). In fact, we prove what we call a “local version” of it, a result implying the existence of a coloring when the number of available colors for each vertex depends on local parameters, like the degree and the clique number of its neighborhood. We prove that for sufficiently large Δ , if G is a graph of maximum degree at most Δ and minimum degree at least $\ln^2(\Delta)$ with list-assignment L , then G is L -colorable if for each $v \in V(G)$,

$$|L(v)| \geq 72 \deg(v) \cdot \min \left\{ \frac{\log_2(\chi(v) + 1)}{\ln(\deg(v))}, \frac{\omega(v) \ln(\ln(\deg(v)))}{\ln(\deg(v))}, \sqrt{\frac{\ln(\omega(v))}{\ln(\deg(v))}} \right\},$$

where $\chi(v)$ denotes the chromatic number of the neighborhood of v and $\omega(v)$ denotes the size of a largest clique containing v . This simultaneously implies the linear improvement over the “greedy bound” and the two aforementioned results of Johansson.

1. INTRODUCTION

Let G be a graph, and for each $v \in V(G)$, let $L(v)$ be a set which we call the *available colors* for v . If each set $L(v)$ is non-empty, then we say that L is a *list-assignment* for G . If k is a positive integer and $|L(v)| \geq k$ for every $v \in V(G)$, then we say that L is a *k -list-assignment* for G . An L -*coloring* of G is a mapping ϕ with domain $V(G)$ such that $\phi(v) \in L(v)$ for every $v \in V(G)$ and $\phi(u) \neq \phi(v)$ for every pair of adjacent vertices $u, v \in V(G)$. We say that a graph G is *k -list-colorable*, or *k -choosable*, if G has an L -coloring for every k -list-assignment L . If $L(v) = \{1, \dots, k\}$ for every $v \in V(G)$, then we call an L -coloring of G a *k -coloring*, and we say G is *k -colorable* if G has a k -coloring. The *chromatic number* of G , denoted $\chi(G)$, is

the smallest k such that G is k -colorable. The *list-chromatic number* of G , denoted $\chi_\ell(G)$, is the smallest k such that G is k -list-colorable.

In 1996, Johansson [10] famously proved that if G is a triangle-free graph of maximum degree at most Δ , then $\chi_\ell(G) = O\left(\frac{\Delta}{\ln(\Delta)}\right)$. Determining the best possible value of the leading constant in this bound is of general interest. The best known lower bound, using Δ -regular graphs, is $\frac{\Delta}{2\ln(\Delta)}$. In 1995, Kim [12] proved that the upper bound holds with a leading constant of $1 + o(1)$ for graphs of girth at least five. In 2015, Pettie and Su [15] improved the leading constant in the upper bound for triangle-free graphs to $4 + o(1)$, and in 2017, Molloy [14], in the following theorem, improved it to $1 + o(1)$, matching the bound of Kim.

Theorem 1.1 ([14]). *If G is a triangle-free graph of maximum degree at most Δ , then*

$$\chi_\ell(G) \leq (1 + o(1)) \frac{\Delta}{\ln(\Delta)}.$$

Johansson [11] also proved that for any fixed $\omega \geq 4$, if G is a graph of maximum degree at most Δ with no clique of size greater than ω , then $\chi_\ell(G) = O\left(\frac{\Delta \ln(\ln(\Delta))}{\ln(\Delta)}\right)$; however, the proof was never published. Molloy [14] proved the following stronger result, which holds even when ω is not fixed.

Theorem 1.2 ([14]). *If G is a graph of maximum degree at most Δ with no clique of size greater than ω , then*

$$\chi_\ell(G) \leq 200\omega \frac{\Delta \ln(\ln(\Delta))}{\ln(\Delta)}.$$

In 1998, Reed [16] conjectured the following, sometimes referred to as “Reed’s ω, Δ, χ Conjecture.”

Conjecture 1.3 ([16]). *If G is a graph of maximum degree at most Δ with no clique of size greater than ω , then*

$$\chi(G) \leq \lceil \frac{1}{2}(\Delta + 1 + \omega) \rceil.$$

It is possible that Conjecture 1.3 is also true for the list-chromatic number. As evidence for his conjecture, Reed [16] proved the following.

Theorem 1.4 ([16]). *There exists $\varepsilon > 0$ such that the following holds. If G is a graph of maximum degree at most Δ with no clique of size greater than ω , then*

$$\chi(G) \leq \lceil (1 - \varepsilon)(\Delta + 1) + \varepsilon\omega \rceil.$$

Note that Theorem 1.4 holds for $\varepsilon = \frac{1}{2}$ if and only if Conjecture 1.3 is true. In 2016, Bonamy, Perrett, and Postle [6] proved that Theorem 1.4 holds for $\varepsilon = \frac{1}{26}$ when Δ is sufficiently large. In 2017, Delcourt and Postle [7] proved that Theorem 1.4 holds for the list-chromatic number for $\varepsilon = \frac{1}{13}$ when Δ is sufficiently large. Results from Ramsey Theory imply that Theorem 1.4 is not true for any value of $\varepsilon > \frac{1}{2}$; for example, Spencer [18] showed the existence of a graph on n vertices with independence number 2 (and thus chromatic number at least $n/2$) such that every clique has size at most $n^{\frac{1}{2} + o(1)}$. The blowup of a 5-cycle, i.e. the Cartesian product of a clique and a 5-cycle, also demonstrates that Theorem 1.4 is not true when $\varepsilon > \frac{1}{2}$, and even that the rounding in Conjecture 1.3 is necessary.

Theorem 1.2 implies Conjecture 1.3 when $\omega = o\left(\frac{\ln(\Delta)}{\ln(\ln(\Delta))}\right)$. It is natural to ask if a bound stronger than that of Conjecture 1.3 can be proved if $\omega = o(\Delta)$ even if $\omega \geq \frac{\ln(\Delta)}{200 \ln(\ln(\Delta))}$. Spencer's result implies that the bound can not be improved if $\omega = \Omega(\Delta^{1/2})$. Considering this, we were motivated to answer the following question.

Question 1.5. *Does there exist a function $f : \mathbb{R} \rightarrow \mathbb{R}$ such that, for every $c > 1$ and every graph G of maximum degree at most Δ with no clique of size greater than $\Delta^{1/f(c)}$, we have $\chi_\ell(G) \leq \Delta/c$?*

Our first result in this paper is the following theorem.

Theorem 1.6. *If G is a graph of maximum degree at most Δ with no clique of size greater than ω , then*

$$\chi_\ell(G) = O\left(\Delta \sqrt{\frac{\ln(\omega)}{\ln(\Delta)}}\right).$$

Theorem 1.6 answers Question 1.5 in the affirmative with a function $f(c)$ that is quadratic in c ; for large enough Δ , the function $f(c) = (72c)^2$ suffices. Determining the best possible function f that confirms Question 1.5 would be very interesting. As mentioned, Spencer [18] showed that $f(2) \geq 2$. This result actually provides a lower bound on $f(c)$ that is linear in c . Spencer [18] proved that the Ramsey number $R(c, \omega)$ is at least $\Omega\left((\omega/\ln(\omega))^{\frac{c+1}{2}}\right)$ as $\omega \rightarrow \infty$ for fixed $c \geq 3$. Therefore there exists a graph G on n vertices with no independent set of size c (and thus chromatic number at least $n/(c-1)$) and no clique of size ω where n is at least $\omega^{\frac{c+1}{2}-o(1)}$. Since the maximum degree of a graph is at most its number of vertices, it follows that $f(c) \geq c/2 + 1$ if $c \in \mathbb{N}$.

The bound of Spencer [18] was improved by Kim in [13] for $c = 3$ by a factor of $\ln \omega$ (matching the upper bound of Ajtai, Komlós, and Szemerédi [1] up to a constant factor), by Bohman in [4] for $c = 4$ by a factor of $\sqrt{\ln \omega}$, and by Bohman and Keevash in [5] for $c \geq 5$ by a factor of $\ln^{\frac{1}{c-2}} \omega$, but these improvements do not change the resulting lower bound on $f(c)$.

1.1. Local Versions. We actually prove a result much stronger than Theorem 1.6. One might wonder if the bounds on $|L(v)|$ supplied by Theorems 1.1, 1.2, 1.4, and 1.6 can be relaxed to depend on local parameters, such as the degree of the vertex v , or the size of a largest clique containing v , rather than the global parameters Δ and ω . To that end, for a vertex v , we let $\deg(v)$ denote the degree of v , $\omega(v)$ denote the size of a largest clique containing v , and $\chi(v)$ denote the chromatic number of the neighborhood of v .

We are interested in proving that a graph G is L -colorable whenever every vertex v satisfies $|L(v)| \geq f(v)$ where $f(v)$ depends on parameters such as $\deg(v)$ and $\omega(v)$. The archetypal example is the classical theorem of Erdős, Rubin, and Taylor [9] that a graph is *degree-choosable* (meaning L -colorable for any list-assignment L satisfying $|L(v)| = \deg(v)$ for every vertex v) unless every block is a clique or an odd cycle. We call such a Theorem a “local version.” Our main result implies local versions of Theorems 1.1, 1.2, and 1.6 simultaneously, although we do not match the leading constant in Theorem 1.1.

In fact, we prove the theorem for *correspondence coloring*, a generalization of list-coloring introduced by Dvořák and Postle [8] in 2015, and also known as *DP-coloring*. We provide a definition in Section 2; the theorem as stated below can also be read as if L is a list assignment.

Theorem 1.7. *For all sufficiently large Δ the following holds. Let G be a graph of maximum degree at most Δ with correspondence assignment (L, M) . For each $v \in V(G)$, let*

$$f(v) = 72 \cdot \min \left\{ \sqrt{\frac{\ln(\omega(v))}{\ln(\deg(v))}}, \frac{\omega(v) \ln(\ln(\deg(v)))}{\ln(\deg(v))}, \frac{\log_2(\chi(v) + 1)}{\ln(\deg(v))} \right\}.$$

If for each $v \in V(G)$,

$$|L(v)| \geq \deg(v) \cdot f(v)$$

and $\deg(v) \geq \ln^2(\Delta)$, then G is (L, M) -colorable.

Recently, Bernshteyn [3] proved that Theorems 1.1 and 1.2 hold for the *correspondence chromatic number*, which is always at least as large as the list-chromatic number. Our Theorem 1.7 implies that “local versions” of these theorems are true for correspondence coloring, as follows.

Corollary 1.8. *For some constant C the following holds. If G is a triangle-free graph of maximum degree at most Δ with correspondence assignment (L, M) such that for each $v \in V(G)$,*

$$|L(v)| \geq C \frac{\deg(v)}{\ln(\deg(v))},$$

and $\deg(v) \geq \ln^2(\Delta)$, then G is (L, M) -colorable.

Corollary 1.9. *For some constant C the following holds. If G is a graph of maximum degree at most Δ with correspondence assignment (L, M) such that for each $v \in V(G)$,*

$$|L(v)| \geq C \deg(v) \frac{\omega(v) \ln(\ln(\deg(v)))}{\ln(\deg(v))},$$

and $\deg(v) \geq \ln^2(\Delta)$, then G is (L, M) -colorable.

We also derive the following “local version” of a result of Johansson [11] on graphs that are *locally r -colorable*, meaning the neighborhood of every vertex is r -colorable.

Corollary 1.10. *For some constant C the following holds. If G is a locally r -colorable graph of maximum degree at most Δ with correspondence assignment (L, M) such that for each $v \in V(G)$,*

$$|L(v)| \geq C \deg(v) \frac{\log_2(r + 1)}{\ln(\deg(v))},$$

and $\deg(v) \geq \ln^2(\Delta)$, then G is (L, M) -colorable.

Of course, Theorem 1.7 also implies a “local version” of Theorem 1.6, as follows.

Corollary 1.11. *For some constant C the following holds. If G is a graph of maximum degree at most Δ with correspondence assignment (L, M) such that for each $v \in V(G)$,*

$$|L(v)| \geq C \deg(v) \sqrt{\frac{\ln(\omega(v))}{\ln(\deg(v))}},$$

and $\deg(v) \geq \ln^2(\Delta)$, then G is (L, M) -colorable.

We now argue that Theorem 1.6 follows from Corollary 1.11.

Proof of Theorem 1.6 assuming Corollary 1.11. Let G be a graph of maximum degree at most Δ with no clique of size greater than ω . We may assume that G has minimum degree at least one. If G has minimum degree at least $\ln^2(\Delta)$, then Corollary 1.11 implies $\chi_\ell(G) \leq C\Delta \sqrt{\frac{\ln(\omega)}{\ln(\Delta)}}$, as desired. Otherwise, we use the following standard procedure to obtain a graph of larger minimum degree containing G as a subgraph. We duplicate the graph G , and we add an edge between each vertex of minimum degree and its duplicate. Note that the minimum degree is increased by one, and that for every vertex v , the size of a largest clique containing v in the new graph does not increase. We repeat this procedure until we obtain a graph G' , having G as a subgraph, and with minimum degree at least $\ln^2(\Delta)$. The result now follows by applying Corollary 1.11 to G' . \square

Although we can not match the leading constant in Theorem 1.1 in our “local version,” we can get the leading constant within a factor of $4 \ln(2)$, as follows.

Theorem 1.12. *For every $\xi > 0$, if Δ is sufficiently large and G is a graph of maximum degree at most Δ with correspondence assignment (L, M) such that for each $v \in V(G)$,*

$$|L(v)| \geq (4 + \xi) \frac{\deg(v)}{\log_2(\deg(v))}$$

and $\deg(v) \geq \ln^2(\Delta)$, then G is (L, M) -colorable.

1.2. A More General Theorem. As mentioned, Bernshteyn [3] proved that Theorems 1.1 and 1.2 hold for the correspondence chromatic number. Many aspects of Bernshteyn’s proofs are similar to those of Molloy’s [14]; however, Bernshteyn’s proof is much shorter and simpler. Molloy used a proof technique known as “entropy compression,” which proves that a random algorithm terminates. Bernshteyn cleverly realized that the use of entropy compression in Molloy’s proof can be replaced with the Lopsided Lovász Local Lemma, resulting in a substantial simplification of the proof.

Both proofs can be applied in the more general setting of graphs in which the average size of an independent set is somewhat large in comparison to the number of independent sets. We make this precise by extracting a more general theorem from their proofs, and we actually prove a “local version” of it, as follows.

For a graph H , let $\bar{\alpha}(H)$ and $i(H)$ denote the average size of an independent set and the number of independent sets in H respectively.

Theorem 1.13. *Let G be a graph of maximum degree at most Δ with correspondence-assignment (L, M) , and $\varepsilon \in (0, 1/2)$. Let $\ell, t : V(G) \rightarrow \mathbb{N}$, and for each $v \in V(G)$,*

let $\alpha_{\min}(v)$ be the minimum of $\bar{\alpha}(H)$ taken over all induced subgraphs $H \subseteq G[N(v)]$ such that $i(H) \geq t(v)$. If for each $v \in V(G)$,

$$|L(v)| \geq \max \left\{ (1 + 2\varepsilon) \frac{\deg(v)}{\alpha_{\min}(v)}, \frac{2t(v)\ell(v)}{\varepsilon(\varepsilon - 2\varepsilon^2)} \right\},$$

and

- (1) $\ell(v) \geq 18 \ln(3\Delta)$,
- (2) $\binom{d(v)}{\ell(v)} / \ell(v)! < \Delta^{-3}/8$,

then G is (L, M) -colorable.

We think that proving Theorem 1.13 separately makes the proof easier to understand, and we think that Theorem 1.13 may have applications not listed in this paper.

1.3. Outline of the Paper. In Section 4, we prove Theorems 1.7 and 1.12 using Theorem 1.13. In order to apply Theorem 1.13, one needs to find a lower bound on $\alpha_{\min}(v)$. We do this by proving a general bound on $\bar{\alpha}(H)$ for a graph H in terms of $i(H)$ and $\omega(H)$. For large values of $\omega(H)$, our bound is better than the bound used by Molloy [14], and this yields the improvement in Theorem 1.6. The condition that $|L(v)| \geq (1 + 2\varepsilon) \frac{\deg(v)}{\alpha_{\min}(v)}$ in Theorem 1.7 results in the bound $|L(v)| \geq \deg(v) \cdot f(v)$ in Theorem 1.7. The condition that $|L(v)| \geq \frac{2t(v)\ell(v)}{\varepsilon(\varepsilon - 2\varepsilon^2)}$ restricts the choice of functions ℓ and t .

In Section 3, we prove Theorem 1.13. The proof is similar to Bernshteyn's proof of Theorem 1.2 from [3]; however, we prove the more general theorem, and some changes are necessary in order to prove the "local version" of it.

In Section 2, we formally define correspondence coloring. We also discuss some notation about "partial colorings" and some probabilistic tools that are needed in the proof of Theorem 1.13.

2. PRELIMINARIES

2.1. Correspondence Coloring. In this subsection we define correspondence coloring.

Definition 2.1. Let G be a graph with list-assignment L .

- If M is a function defined on $E(G)$ where for each $e = uv \in E(G)$, M_e is a matching of $\{u\} \times L(u)$ and $\{v\} \times L(v)$, we say (L, M) is a *correspondence-assignment* for G .
- An (L, M) -*coloring* of G is a function $\varphi : V(G) \rightarrow \mathbb{N}$ such that $\varphi(u) \in L(u)$ for every $u \in V(G)$, and for every $e = uv \in E(G)$, $(u, \varphi(u))(v, \varphi(v)) \notin M_e$. If G has an (L, M) -coloring, then we say G is (L, M) -colorable.

A correspondence-assignment (L, M) is a k -*correspondence-assignment* if L is a k -list-assignment, and the *correspondence chromatic number* of G , denoted $\chi_c(G)$, is the minimum k such that for every k -correspondence-assignment (L, M) , G is (L, M) -colorable.

For convenience, if $uv \in E(G)$, $c_1 \in L(u)$, $c_2 \in L(v)$, and $(u, c_1)(v, c_2) \in M_{uv}$, we will just say $c_1 c_2 \in M_{uv}$. We will also say c_1 *corresponds* to c_2 . Note that if for each $e = uv \in E(G)$ and $c \in L(u) \cap L(v)$, $cc \in M_{uv}$, then an (L, M) -coloring is an L -coloring. Hence, for every graph G , $\chi_\ell(G) \leq \chi_c(G)$.

2.2. Partial Colorings. The proof of Theorem 1.13 relies on analyzing a “partial coloring” of the graph chosen uniformly at random. In this subsection, we define some notation that will be useful for this analysis.

Definition 2.2. Let G be a graph with correspondence-assignment (L, M) , and let BLANK be a color not in $L(v)$ for any $v \in V(G)$. A *partial (L, M) -coloring* of G is a mapping ϕ with domain $V(G)$ such that for each $v \in V(G)$, $\phi(v) \in L(v) \cup \{\text{BLANK}\}$. If $\phi(v) = \text{BLANK}$, we say v is *ϕ -uncolored*. For each ϕ -uncolored vertex v , we let $L_\phi(v)$ denote the set of colors $c \in L(v)$ such that for every neighbor u of v , $\phi(u)$ does not correspond to c , and we let M_ϕ denote the restriction of M to edges between ϕ -uncolored vertices. If ϕ' is a partial (L_ϕ, M_ϕ) -coloring of the ϕ -uncolored vertices of G , we let

$$(\phi + \phi')(v) = \begin{cases} \phi(v) & \text{if } \phi(v) \neq \text{BLANK}, \\ \phi'(v) & \text{otherwise.} \end{cases}$$

We will show that with nonzero probability the random partial coloring can be extended to a coloring of the whole graph. Using the following proposition, it will suffice to show that if ϕ is a partial coloring of G chosen uniformly at random, then the ϕ -uncolored vertices induce a subgraph that is (L_ϕ, M_ϕ) -colorable.

Proposition 2.3. *If G is a graph with correspondence-assignment (L, M) , ϕ is a partial (L, M) coloring of G , and ϕ' is a (L_ϕ, M_ϕ) -coloring of the graph induced by G on the ϕ -uncolored vertices, then $\phi + \phi'$ is an (L, M) -coloring of G .*

2.3. Probabilistic Tools. We will need the following version of the Chernoff bounds for negatively correlated random variables.

Definition 2.4. We say that boolean random variables X_1, \dots, X_m are *negatively correlated* if for every $I \subseteq \{1, \dots, m\}$,

$$\mathbb{P}[\wedge_{i \in I} X_i] \leq \prod_{i \in I} \mathbb{P}[X_i].$$

Lemma 2.5 (Chernoff Bounds). *Let X_1, \dots, X_m be negatively correlated boolean random variables, and let $X = \sum_{i=1}^m X_i$. Then for any $0 < t \leq \mathbb{E}[X]$,*

$$\mathbb{P}[|X - \mathbb{E}[X]| > t] \leq 2 \exp\left(-\frac{t^2}{3\mathbb{E}[X]}\right).$$

In our application of Lemma 2.5, we will have a random partial (L, M) -coloring for some correspondence assignment (L, M) and a boolean random variable for each color $c \in L(v)$ indicating if v has a neighbor whose color corresponds to c .

We will also need the Lovász Local Lemma.

Lemma 2.6 (Lovász Local Lemma). *Let I be a finite set, and for each $i \in I$, let B_i be a random event. Suppose that for every $i \in I$, there is a set $\Gamma(i) \subseteq I$ such that $|\Gamma(i)| \leq d$ and for all $Z \subseteq I \setminus \Gamma(i)$,*

$$\mathbb{P}\left[B_i \mid \bigcap_{j \in Z} \overline{B_j}\right] \leq p.$$

If $4pd \leq 1$, then with nonzero probability none of the events B_i occur.

3. PROOF OF THEOREM 1.13

In this section, we prove Theorem 1.13. We assume G , (L, M) , Δ , ℓ , and t satisfy the conditions of Theorem 1.13 throughout the section.

3.1. Completing a partial coloring. We prove Theorem 1.13 by finding a partial (L, M) -coloring of G that we can greedily extend to an (L, M) -coloring. The following lemma provides the existence of such a partial (L, M) -coloring.

Lemma 3.1. *There exists a partial (L, M) -coloring ϕ of G such that for every ϕ -uncolored vertex v ,*

- (1) $|L_\phi(v)| \geq \ell(v)$, and
- (2) v has fewer than $\ell(v)$ ϕ -uncolored neighbors u such that $\ell(u) \geq \ell(v)$.

Lemma 3.1 generalizes Lemma 4.1 in the proof of Bernshteyn [3], and the partial (L, M) -coloring in Lemma 3.1 generalizes the “flaw-free” coloring output by the random algorithm of Molloy [14]. When the function ℓ is not constant, our second condition is slightly weaker, so we are not necessarily able to complete the partial coloring greedily in any order as in their proofs.

Proof of Theorem 1.13 assuming Lemma 3.1. Let ϕ be the partial (L, M) -coloring of G satisfying (1) and (2) of Lemma 3.1. By Proposition 2.3, it suffices to show that the ϕ -uncolored vertices induce a graph that is (L_ϕ, M_ϕ) -colorable. This follows by ordering the ϕ -uncolored vertices v by $\ell(v)$ from greatest to least, breaking ties arbitrarily, and coloring greedily. \square

3.2. Analyzing a random partial coloring. We prove Lemma 3.1 by analyzing a partial (L, M) -coloring of G chosen uniformly at random and using the Local Lemma to show that with nonzero probability, the random partial coloring satisfies Lemma 3.1. Instead of using the Local Lemma, Molloy [14] used the entropy compression technique. The key insight of Bernshteyn [3] was that a clever application of the Local Lemma is sufficient, and this greatly simplified the proof. In order to apply the Local Lemma to prove Lemma 3.1, we will need the following lemma.

Lemma 3.2. *Let $v \in V(G)$ and fix a partial (L, M) -coloring ϕ_1 of $G - N[v]$. Let ϕ_2 be a partial (L_{ϕ_1}, M_{ϕ_1}) coloring of $G[N(v)]$ chosen uniformly at random, and let $\phi = \phi_1 + \phi_2$. Let*

- (1) A_{v, ϕ_1} be the event that $|L_\phi(v)| < \ell(v)$ and
- (2) B_{v, ϕ_1} be the event that v has at least $\ell(v)$ ϕ -uncolored neighbors u such that $|L_\phi(u)| \geq \ell(u) \geq \ell(v)$.

Then $\mathbb{P}[A_{v, \phi_1}], \mathbb{P}[B_{v, \phi_1}] \leq \Delta^{-3}/8$.

Lemma 3.2 generalizes Lemma 4.2 in the proof of Bernshteyn [3] and Lemma 12 in the proof of Molloy [14].

To bound $\mathbb{P}[A_{v, \phi_1}]$ in Lemma 3.2, we show that $|L_\phi(v)|$ is large in expectation and with high probability is concentrated around its expectation, as in the following lemma.

Lemma 3.3. *Under the assumptions of Lemma 3.2,*

$$(1) \quad \mathbb{E}[|L_\phi(v)|] \geq \varepsilon(\varepsilon - 2\varepsilon^2) \frac{|L(v)|}{t(v)},$$

and

$$(2) \quad \mathbb{P} \left[|L_\phi(v)| - \mathbb{E}[|L_\phi(v)|] > \frac{1}{2} \mathbb{E}[|L_\phi(v)|] \right] \leq 2 \exp \left(\frac{-\mathbb{E}[|L_\phi(v)|]}{12} \right).$$

Before proving Lemma 3.3, we need some definitions. For the remainder of this subsection, we assume v, ϕ_1, ϕ_2 , and ϕ are as in Lemma 3.2.

Definition 3.4. For each $c \in L(v)$, let the random variable

$$\mathbf{appear}_c = |\{u \in N(v) : c\phi_2(u) \in M_{vu}\}|,$$

i.e. the number of neighbors u of v such that $\phi_2(u)$ corresponds to c .

Note that

$$(3) \quad \mathbb{E}[|L_\phi(v)|] = \sum_{c \in L(v)} \mathbb{P}[\mathbf{appear}_c = 0].$$

By (3), in order to prove Lemma 3.3, we need some bounds on $\mathbb{P}[\mathbf{appear}_c = 0]$ for the colors $c \in L(v)$. These bounds will depend on the average size and number of independent sets of certain subgraphs induced by neighbors u of v such that $L(u)$ contains a color corresponding to c .

Definition 3.5. Fix $c \in L(v)$ and a partial (L_{ϕ_1}, M_{ϕ_1}) coloring ϕ'_2 of $G[N(v)]$ such that for no neighbor u of v , the color $\phi'_2(u)$ corresponds to c . Let $\text{col}(c, \phi'_2)$ denote the ϕ'_2 -uncolored neighbors u of v such that $L_{\phi_1+\phi'_2}(u)$ contains c , i.e. the ϕ'_2 -uncolored neighbors of v that can be colored c without creating conflicts.

Definition 3.6. For each $c \in L(v)$, let ϕ_2^ε be the partial coloring obtained from ϕ_2 by uncoloring any neighbor u of v such that $\phi_2(u)$ corresponds to c .

Proposition 3.7. Fix $c \in L(v)$ and a partial (L_{ϕ_1}, M_{ϕ_1}) -coloring ϕ'_2 such that for no neighbor u of v , the color $\phi'_2(u)$ corresponds to c . Then

- (i) $\mathbb{E}[\mathbf{appear}_c | \phi_2^\varepsilon = \phi'_2] = \bar{\alpha}(G[\text{col}(c, \phi'_2)])$, and
- (ii) $\mathbb{P}[\mathbf{appear}_c = 0 | \phi_2^\varepsilon = \phi'_2] = i(\text{col}(c, \phi'_2))^{-1}$.

Definition 3.8. Let $\mathbf{col}(c)$ denote the random set of neighbors u of v such that $L_{\phi_1}(u)$ contains c and $\phi_2(u) \in \{c, \text{BLANK}\}$.

We can now prove Lemma 3.3.

Proof of Lemma 3.3. First we prove that (1) holds. We divide $L(v)$ into two parts in the following way. For each $c \in L(v)$, we let $c \in L_1(v)$ if $\mathbb{P}[i(\mathbf{col}(c)) \leq t(v)] \geq \varepsilon$, and otherwise we let $c \in L_2(v)$.

First we claim that $|L_2(v)| \leq |L(v)|/(1+\varepsilon-2\varepsilon^2)$. If $c \in L_2(v)$, by Proposition 3.7 and the definition of $\alpha_{\min}(v)$,

$$(4) \quad \mathbb{E}[\mathbf{appear}_c] \geq (1-\varepsilon)\alpha_{\min}(v).$$

However,

$$(5) \quad \sum_{c \in L_2(v)} \mathbb{E}[\mathbf{appear}_c] \leq \deg(v).$$

By (4) and (5), $(1-\varepsilon)\alpha_{\min}(v)|L_2(v)| \leq \deg(v)$. Since $|L(v)| \geq (1+2\varepsilon)\frac{\deg(v)}{\alpha_{\min}(v)}$, by rearranging terms,

$$|L_2(v)| \leq \frac{\deg(v)}{(1-\varepsilon)\alpha_{\min}(v)} \leq \frac{|L(v)|}{(1-\varepsilon)(1+2\varepsilon)},$$

as claimed. Since $|L(v)| = |L_1(v)| + |L_2(v)|$, this implies $|L_1(v)| \geq (\varepsilon - 2\varepsilon^2)|L(v)|$.

If $c \in L_1(v)$, by Proposition 3.7 and the definition of $L_1(v)$,

$$(6) \quad \mathbb{P}[\mathbf{appear}_c = 0] \geq \frac{\varepsilon}{t(v)}.$$

By (3) and (6),

$$\mathbb{E}[|L_\phi(v)|] \geq \frac{\varepsilon}{t(v)}|L_1(v)| \geq \varepsilon(\varepsilon - 2\varepsilon^2) \frac{|L(v)|}{t(v)},$$

as desired.

Now we prove that (2) holds. By (3), $\mathbb{E}[|L_\phi(v)|]$ is a sum of Boolean random variables

$$X_c = \begin{cases} 1 & \text{if } \mathbf{appear}_c = 0, \\ 0 & \text{if } \mathbf{appear}_c > 0, \end{cases}$$

for each $c \in L(v)$. Since the random variables X_c are negatively correlated, it follows from Lemma 2.5 with $t = \mathbb{E}[L_\phi(v)]/2$. \square

We can now prove Lemma 3.2.

Proof of Lemma 3.2. First we prove that $\mathbb{P}[A_{v,\phi_1}] \leq \Delta^{-3}/8$. Since $|L(v)| \geq \frac{2\ell(v)t(v)}{\varepsilon(\varepsilon - 2\varepsilon^2)}$, Lemma 3.3 implies that

$$(7) \quad \mathbb{E}[|L_\phi(v)|] \geq 2\ell(v).$$

Therefore by the definition of A_{v,ϕ_1} ,

$$\mathbb{P}[A_{v,\phi_1}] \leq \mathbb{P}\left[|L_\phi(v)| - \mathbb{E}[|L_\phi(v)|] > \frac{1}{2}\mathbb{E}[|L_\phi(v)|]\right].$$

Now by Lemma 3.3, (7), and the hypothesis that $\ell(v) \geq 18 \ln(3\Delta)$,

$$\mathbb{P}[A_{v,\phi_1}] \leq 2 \exp(-\ell(v)/6) \leq \Delta^{-3}/8,$$

as desired.

It remains to bound $\mathbb{P}[B_{v,\phi_1}]$. Let X be any set of $\ell(v)$ neighbors u of v such that $|L_\phi(u)| \geq \ell(u) \geq \ell(v)$. For any partial (L_{ϕ_1}, M_{ϕ_1}) coloring ϕ'_2 of $N(v)$ such that the vertices in X are ϕ'_2 -uncolored, there are at least $\ell(v)!$ partial $(L_{\phi_1+\phi'_2}, M_{\phi_1+\phi'_2})$ colorings of X , and in only one of them all of X is uncolored. Therefore the probability that every vertex of X is ϕ -uncolored is at most $\frac{1}{\ell(v)!}$. By the Union Bound and the assumption that $\binom{\deg(v)}{\ell(v)}/\ell(v)! \leq \Delta^{-3}/8$, this implies $\mathbb{P}[B_{v,\phi_1}] \leq \Delta^{-3}/8$, as desired. \square

3.3. Finding the partial coloring. In this subsection, we prove Lemma 3.1. Recall that Lemma 3.1 implies Theorem 1.13.

Proof of Lemma 3.1. Let ϕ be a partial (L, M) -coloring of G chosen uniformly at random. For each $v \in V(G)$, let A_v be the event that $|L_\phi(v)| < \ell(v)$, let B_v be the event that v has at least $\ell(v)$ ϕ -uncolored neighbors u such that $|L_\phi(u)| \geq \ell(u) \geq \ell(v)$, and let $\Gamma(v)$ denote the set of vertices of distance at most three from v in G . Note that for all $v \in V(G)$, $|\Gamma(v)| \leq \Delta^3$.

First we claim that with nonzero probability, none of the events $(A_v \cup B_v)$ occur in the random partial coloring ϕ . By the Local Lemma (Lemma 2.6), it suffices to show that for each $v \in V(G)$ and $Z \subseteq V(G) \setminus \Gamma(v)$, $\mathbb{P}[A_v | \bigcap_{u \in Z} \overline{A_u \cup B_u}] \leq \Delta^{-3}/8$ and $\mathbb{P}[B_v | \bigcap_{u \in Z} \overline{A_u \cup B_u}] \leq \Delta^{-3}/8$. This follows from Lemma 3.2.

Therefore there is a partial (L, M) -coloring ϕ' for which none of the events $(A_v \cup B_v)$ occur. We claim that ϕ' satisfies Lemma 3.1. Suppose not. If for some $v \in V(G)$, condition (1) is not satisfied, then A_v holds, a contradiction. Therefore we may assume for some $v \in V(G)$, condition (2) is not satisfied, that is v has fewer than $\ell(v)$ ϕ' -uncolored neighbors u such that $\ell(u) \geq \ell(v)$. Since B_v holds, for some neighbor u , $|L_{\phi'}(u)| < \ell(u)$, contradicting that A_u does not hold. Therefore ϕ' satisfies Lemma 3.1, as claimed. \square

4. PROOFS OF THEOREMS 1.7 AND 1.12

In this section we prove Theorems 1.7 and 1.12. In this section, \log means the base 2 logarithm.

4.1. Bounding the Average Size of an Independent Set. We will use Theorem 1.13, so we will need a lower bound on $\alpha_{\min}(v)$. We do this by bounding the average size of an independent set in terms of the total number. In the proof of Theorem 1.2, Molloy [14] and Bernshteyn [3] use the following result of Shearer [17], which we will also need.

Lemma 4.1 ([17]). *If H is a graph with no clique of size greater than ω , then*

$$\bar{\alpha}(H) \geq \frac{\log(i(H))}{2\omega \log(\log(i(H)))}.$$

We will also need the following result of Alon [2].

Lemma 4.2 ([2]). *If H is a graph on n vertices, then*

$$\bar{\alpha}(H) \geq \frac{\log(i(H))}{10 \log(n/\log(i(H)) + 1)}.$$

Since $\log(i(H)) \geq \alpha(H)$, we can replace the $\log(i(H))$ in the denominator of the bound in Lemma 4.2 with $\alpha(H)$ to get a suitable bound if H contains a large independent set.

The following lemma provides an improvement over Lemma 4.1 for larger values of ω .

Lemma 4.3. *If H is a graph on n vertices with no clique of size greater than ω and n is sufficiently large, then*

$$\bar{\alpha}(H) \geq \frac{1}{24} \sqrt{\frac{\log(i(H))}{\log(\omega)}}.$$

We will actually use Lemma 4.2 to prove Lemma 4.3. To apply Lemma 4.2, we need to upper bound $\log(n)$ in terms of $\log(i(H))$ and $\omega(H)$, as in the following lemma.

Lemma 4.4. *If H is a graph on n vertices with no clique of size greater than ω and n is sufficiently large, then*

$$\log(i(H)) \geq \frac{\log^2(n)}{2e \log(\omega)}.$$

Proof. We may assume $\omega \geq 3$, or else H has an independent set of size at least \sqrt{n} , and the result follows.

Let α be some positive integer to be determined later, and let $s = R(\alpha, \omega + 1)$, the Ramsey number. We will actually prove there are at least $2^{\frac{\log^2(n)}{2e \log(\omega)}}$ independent sets of size α .

By the definition of s , every subset of $V(H)$ of size s has an independent set of size α . Since every independent set of size α is contained in at most $\binom{n-\alpha}{s-\alpha}$ subsets of $V(H)$ of size s , there are at least

$$(8) \quad \binom{n}{s} / \binom{n-\alpha}{s-\alpha} \geq \left(\frac{n-\alpha}{s} \right)^\alpha = 2^{\alpha(\log(n-\alpha) - \log(s))}$$

independent sets of size α .

We let $\alpha = \frac{\log(n)}{e \log(\omega)} + 1$. By (8), it suffices to show that $\log(n-\alpha) - \log(s) \geq \log(n)/2$. It is well-known that $R(\alpha, \omega+1) \leq \binom{\alpha+\omega-1}{\alpha-1} \leq \left(\frac{\alpha+\omega-1}{\alpha-1} \cdot e \right)^{\alpha-1}$. Therefore

$$\log(s) \leq (\alpha-1) \log \left(\frac{\alpha+\omega-1}{\alpha-1} \cdot e \right) \leq \frac{\log(n)}{e \log(\omega)} \log \left(e + \omega \frac{\log(\omega)}{\log(n)} \right).$$

Since $\alpha = o(n)$ and $\omega \geq 3$, for n sufficiently large, $\log(n-\alpha) - \log(s) \geq \log(n)/2$, as desired. \square

Now we can prove Lemma 4.3.

Proof of Lemma 4.3. By Lemma 4.2,

$$\bar{\alpha}(H) \geq \frac{\log(i(H))}{10 \log(n)}.$$

By Lemma 4.4,

$$\log(n) \leq \sqrt{2e \log(i(H)) \log(\omega)},$$

and the result follows. \square

4.2. The Proofs. Lemmas 4.1, 4.2, and 4.3 provide good enough bounds for $\alpha_{\min}(v)$. Now we are able to prove Theorems 1.7 and 1.12.

Proof of Theorem 1.7. Let $\varepsilon = 1/4$, and let $v \in V(G)$. We show that v satisfies the conditions of Theorem 1.13. Let $\ell(v) = \deg(v)^{5/8}$, and let $t(v) = \deg(v)^{1/4}$. By Lemma 4.3,

$$\alpha_{\min}(v) \geq \frac{1}{48} \sqrt{\frac{\log(\deg(v))}{\log(\omega(v))}}.$$

By Lemma 4.1,

$$\alpha_{\min}(v) \geq \frac{\log(\deg(v))}{8\omega(v) \log(\log(\deg(v)))}.$$

Note that $\log(i(H)) \geq \alpha(H)$ for any graph H . Hence if $H \subseteq G[N(v)]$, then $|V(H)| / \log(i(H)) \leq \chi(H) \leq \chi(v)$. Therefore by Lemma 4.2,

$$\alpha_{\min}(v) \geq \frac{\log(\deg(v))}{40 \log(\chi(v) + 1)}.$$

Since $|L(v)| \geq 72 \deg(v) f(v)$, it follows that $|L(v)| \geq (1 + 2\varepsilon) \frac{\deg(v)}{\alpha_{\min}(v)}$, as desired. Note that $f(v)^8 = o(\deg(v))$. Since Δ is sufficiently large and $\deg(v) \geq \log^2(\Delta)$, we may assume $f(v) \geq 168 \deg(v)^{-1/8}$. Since $t(v)\ell(v) = \deg(v)^{7/8}$, $|L(v)| \geq \frac{2t(v)\ell(v)}{\varepsilon(\varepsilon - 2\varepsilon^2)}$, as desired.

Since $\ell(v) \geq \ln^{5/4}(\Delta)$ and Δ is sufficiently large, $\ell(v) \geq 18 \ln(3\Delta)$, as desired. It remains to show that $\binom{\deg(v)}{\ell(v)}/\ell(v)! < \Delta^{-3}/8$. We will use the following form of Stirling's approximation:

$$n! \geq \sqrt{2\pi n} n^{n+1/2} e^{-n}.$$

Therefore

$$(9) \quad \binom{\deg(v)}{\ell(v)}/\ell(v)! < \frac{\deg(v)^{\ell(v)}}{(\ell(v)!)^2} \leq \left(\frac{e^2 \deg(v)}{2\pi \ell(v)^{2+1/\ell(v)}} \right)^{\ell(v)}.$$

Since $\deg(v)/\ell(v)^{2+1/\ell(v)} \leq \ell(v)^{-2/5}$, by taking the logarithm of the bound in (9), it suffices to show that

$$\ell(v) \ln \left(\frac{2\pi \ell(v)^{2/5}}{e^2} \right) \geq 3 \ln(8\Delta).$$

Since $\ell(v) \geq \ln^{5/4}(\Delta)$ and Δ is sufficiently large, this follows. \square

Proof of Theorem 1.12. Let $\varepsilon = \xi/10$ and $\varepsilon' > 0$ be some constant to be chosen later, and let $v \in V(G)$. Let $\ell(v) = \deg(v)^{(1+\varepsilon')/2}$, and let $t(v) = \deg(v)^{(1-2\varepsilon')/2}$. Since G is triangle-free,

$$\alpha_{\min}(v) \geq \frac{\log(t(v))}{2} = (1 - 2\varepsilon') \log(\deg(v))/4.$$

Since $|L(v)| \geq (4 + \xi) \deg(v)/\log(\deg(v))$, it follows that $|L(v)| \geq (1 + \xi/4)(1 - 2\varepsilon') \frac{\deg(v)}{\alpha_{\min}(v)}$. We choose ε' sufficiently small so that $(1 + \xi/4)(1 - 2\varepsilon') \geq 1 + 2\varepsilon$.

Note that $t(v)\ell(v) = \deg(v)^{1-\varepsilon'/2}$. Hence we may assume $\deg(v)$ is sufficiently large so that $|L(v)| \geq \frac{2t(v)\ell(v)}{\varepsilon(2-2\varepsilon^2)}$, as desired.

Note also that $\ell(v) \geq \ln^{1+\varepsilon'}(\Delta)$. The rest of the proof is similar to the proof of Theorem 1.7, so we omit it. \square

REFERENCES

- [1] M. Ajtai, J. Komlós, and E. Szemerédi. A note on Ramsey numbers. *J. Combin. Theory Ser. A*, 29(3):354–360, 1980.
- [2] N. Alon. Independence numbers of locally sparse graphs and a Ramsey type problem. *Random Structures Algorithms*, 9(3):271–278, 1996.
- [3] A. Bernshteyn. The Johansson–Molloy Theorem for DP-Coloring. *ArXiv e-prints*, August 2017.
- [4] T. Bohman. The triangle-free process. *Adv. Math.*, 221(5):1653–1677, 2009.
- [5] T. Bohman and P. Keevash. The early evolution of the H -free process. *Invent. Math.*, 181(2):291–336, 2010.
- [6] M. Bonamy, T. Perrett, and L. Postle. Colouring graphs with sparse neighbourhoods: Bounds and applications. submitted.
- [7] M. Delcourt and L. Postle. On the list coloring version of reed's conjecture. manuscript.
- [8] Z. Dvořák and L. Postle. Correspondence coloring and its application to list-coloring planar graphs without cycles of lengths 4 to 8. *J. Combin. Theory Ser. B*, 129:38–54, 2018.
- [9] P. Erdős, A. L. Rubin, and H. Taylor. Choosability in graphs. In *Proceedings of the West Coast Conference on Combinatorics, Graph Theory and Computing (Humboldt State Univ., Arcata, Calif., 1979)*, Congress. Numer., XXVI, pages 125–157. Utilitas Math., Winnipeg, Man., 1980.
- [10] A. Johansson. Asymptotic choice number for triangle free graphs. Unpublished Manuscript, 1996.
- [11] A. Johansson. The choice number of sparse graphs. Unpublished Manuscript, 1996.
- [12] J. H. Kim. On Brooks' theorem for sparse graphs. *Combin. Probab. Comput.*, 4(2):97–132, 1995.

- [13] J. H. Kim. The Ramsey number $R(3, t)$ has order of magnitude $t^2 / \log t$. *Random Structures Algorithms*, 7(3):173–207, 1995.
- [14] M. Molloy. The list chromatic number of graphs with small clique number. *ArXiv e-prints*, January 2017.
- [15] S. Pettie and H-H. Su. Distributed coloring algorithms for triangle-free graphs. *Inform. and Comput.*, 243:263–280, 2015.
- [16] B. Reed. ω , Δ , and χ . *J. Graph Theory*, 27(4):177–212, 1998.
- [17] J. B. Shearer. On the independence number of sparse graphs. *Random Structures Algorithms*, 7(3):269–271, 1995.
- [18] J. Spencer. Asymptotic lower bounds for Ramsey functions. *Discrete Math.*, 20(1):69–76, 1977/78.

CNRS, LABRI, UNIVERSITÉ DE BORDEAUX
E-mail address: marthe.bonamy@u-bordeaux.fr

UNIVERSITY OF WATERLOO
E-mail address: t9kelly@uwaterloo.ca
E-mail address: apnelson@uwaterloo.ca
E-mail address: lpostle@uwaterloo.ca