

The groups G_{k+1}^k and fundamental groups of configuration spaces

V.O.Manturov *

To Leonid Arkadievich Bokut' on the occasion of his 80th birthday

AMS MSC 57M25,57M27

Keywords: group, configuration space, braid, word problem, conjugacy problem

Abstract

In [10], the author has constructed natural maps from fundamental groups of topological spaces (restricted configuration spaces) to the groups G_n^k . In the present paper, we show that in the case of $n = k + 1$, the group G_{k+1}^k is isomorphic to the fundamental group of some (quotient space of) some configuration space. In particular, this leads to the solution of word and conjugacy problems in G_4^3 and sheds light on G_{k+1}^k for higher k .

1 Introduction

In [8], the author defined a family of groups G_n^k depending on two natural numbers $n > k$, and formulated the following principle: *if a dynamical system describing a motion of n particles, admits some “good” codimension one property governed by exactly k particles, then this dynamical system has a topological invariant valued in G_n^k .* These groups are related to many problems in topology and combinatorial group theory, see, e.g., [7],[11], [12].

For $n \in \mathbb{N}$, let $[n] = \{1, \dots, n\}$.

The groups G_n^k are defined as follows.

$$G_n^k = \langle a_m \mid (1), (2), (3) \rangle,$$

where the generators a_m are indexed by all k -element subsets of $[n]$, the relation (1) means

$$(a_m)^2 = 1 \text{ for any unordered sets } m \subset \{1, \dots, n\}, \text{Card}(m) = k; \quad (1)$$

(2) means

$$a_m a_{m'} = a_{m'} a_m, \text{ if } \text{Card}(m \cap m') < k - 1; \quad (2)$$

*Research is carried out with the support of Russian Science Foundation (project no. 16-11-10291).

and, finally, the relations (3) look as follows. For every set $U \subset [n]$ of cardinality $(k + 1)$, let us order all its k -element subsets arbitrarily and denote them by m^1, \dots, m^{k+1} . Then (3) looks as

$$(a_{m^1} \cdots a_{m^{k+1}})^2 = 1. \quad (3)$$

In view of (1), we can rewrite (3) in the form

$$a_{m^1} \cdots a_{m^{k+1}} = a_{m^{k+1}} \cdots a_{m^1}. \quad (3')$$

Note that in the case $n = k + 1$, the relations (2) are void, since any two distinct subsets m, m' of $[n]$ of cardinality k have intersection of cardinality $k - 1$.

In [10], groups G_n^k are related to fundamental groups of the following (reduced) configuration spaces. We take n -point sets in the $(k - 1)$ -dimensional real projective plane $\mathbb{R}P^{k-1}$, such that no $(k - 1)$ of these points belong to the same (projective) $(k - 3)$ -plane. In particular, for $k = 3$ we deal with n -strand $\mathbb{R}P^2$ -braids. There is a map from the fundamental group of the (ordered) configuration space to G_n^k .

These maps correspond to the good property “some k points are not in general position” (i.e. belong the same $(k - 2)$ -projective plane). Note that the we have imposed the restriction that any $(k - 1)$ points should be in general position.

The maps constructed in [10] have some obvious kernel corresponding to rotations of $\mathbb{R}P^{k-1}$. It turns out that in the lowest level (for $n = k + 1$), after factorisation by this kernel, we get an isomorphism of groups.

The kernel looks as follows: identifying $\mathbb{R}P^{k-1}$ with the quotient space of S^{k-1} by the involution, we can just rotate S^{k-1} about some axis by π ; the corresponding motion of points is homotopically non-trivial; on the other hand, if points are in general position from the very beginning, they remain in the general position during the rotation. Hence, no singular moment occurs and the corresponding word is empty.

To remedy this, we define the configuration space $\tilde{C}'_n(\mathbb{R}P^{k-1})$ of all unordered n -tuples of points where the first $(k - 1)$ points are fixed and any $k - 1$ points are in general position. For example, for $\mathbb{R}P^2$, we can consider the path when one point x_1 is fixed and three other points lie in the small neighbourhood of x_1 ; when rotating them around x_1 , we get a non-trivial element of the fundamental group of the corresponding configuration space, and during the whole path, there are no moments when any three points are collinear. Then every closed path in the configuration space giving rise to the empty word is homotopic to the trivial path. See Lemma 3.

More precisely, we shall prove the following

Theorem 1. *There is a subgroup \tilde{G}_{k+1}^k of the group G_{k+1}^k of index 2^{k-1} is isomorphic to $\pi_1(\tilde{C}'_{k+1}(\mathbb{R}P^{k-1}))$.*

The space \tilde{C}'_n and the group \tilde{G}_{k+1}^k will be defined later.

The simplest case of the above Theorem is

Theorem 2. *The group \tilde{G}_4^3 (subgroup of index four of G_4^3) is isomorphic to $\pi_1(FBr_4(\mathbb{R}P^2))$, the 4-strand pure braid group on $\mathbb{R}P^2$ with two point fixed.*

The paper is organized as follows. In the next section, we give all necessary definitions and construct maps from configuration spaces of points in $\mathbb{R}P^{k-1}$ to G_{k+1}^k . In Section 3, we prove Theorem

1. In Section 4, we shall discuss an algebraic lemma about reduction of the word problem in G_{k+1}^k to the word problem in some subgroup of it.

We conclude the paper by Section 4 by discussing some open problems for further research.

2 Basic definitions

Let us now pass to the definition of spaces $C'_n(\mathbb{R}P^{k-1})$ and maps from the corresponding fundamental groups to the groups G_n^k .

Let us fix a pair of natural number $n > k$. A point in $C'_n(\mathbb{R}P^{k-1})$ is an ordered set of n pairwise distinct points in $\mathbb{R}P^{k-1}$, such that any $(k-1)$ of them are in general position. Thus, for instance, if $k=3$, then the only condition is that these points are pairwise distinct. For $k=4$ for points $x_1, \dots, x_n \in \mathbb{R}P^3$ we impose the condition that no three of them belong to the same line (though some four are allowed to belong to the same plane), and for $k=5$ a point in $C'_n(\mathbb{R}P^4)$ is a set of ordered n points in $\mathbb{R}P^4$, with no four of them belonging to the same 2-plane.

Let us use the projective coordinates $(a_1 : a_2 : \dots : a_k)$ in $\mathbb{R}P^{k-1}$ and let us fix the following $k-1$ points in general position, $y_1, y_2, \dots, y_{k-1} \in \mathbb{R}P^{k-1}$, where $a_i(y_j) = \delta_i^j$. Let us define the subspace $\tilde{C}'_n(\mathbb{R}P^{k-1})$ taking those n -tuples of points $x_1, \dots, x_n \in \mathbb{R}P^{k-1}$ for which $x_i = y_i$ for $i=1, \dots, k-1$.

We say that a point $x \in C'_n(\mathbb{R}P^{k-1})$ is *singular*, if the set of points $x = (x_1, \dots, x_n)$, corresponding x , contains some subset of k points lying on the same $(k-2)$ -plane. Let us fix two non-singular points $x, x' \in C'_n(\mathbb{R}P^{k-1})$.

We shall consider smooth paths $\gamma_{x,x'} : [0, 1] \rightarrow C'_n(\mathbb{R}P^{k-1})$. For each such path there are values t for which $\gamma_{x,x'}(t)$ is not in the general position (some k of them belong to the same $(k-2)$ -plane). We call these values $t \in [0, 1]$ *singular*.

On the path γ , we say that the moment t of passing through the singular point x , corresponding to the set x_{i_1}, \dots, x_{i_k} , is *transverse* (or *stable*) if for any sufficiently small perturbation $\tilde{\gamma}$ of the path γ , in the neighbourhood of the moment t there exists exactly one time moment t' corresponding to some set of points x_{i_1}, \dots, x_{i_k} not in general position.

Definition 1. We say that a path is good and stable if the following holds:

1. The set of singular values t is finite;
2. For every singular value $t = t_l$ corresponding to n points representing $\gamma_{x,x'}(t_l)$, there exists only one subset of k points belonging to a $(k-2)$ -plane;
3. Each singular value is stable.

Definition 2. We say that the path without singular values is void.

Definition 3. By a braid we mean a smooth path in the above configuration space whose initial and terminal configurations of points coincide as sets. We say that a braid whose ends x, x' coincide with respect to the order, is pure. We say that two braids γ, γ' with endpoints x, x' are isotopic if there exists a continuous family $\gamma_{x,x'}^s, s \in [0, 1]$ of smooth paths with fixed ends such that $\gamma_{x,x'}^0 = \gamma, \gamma_{x,x'}^1 = \gamma'$.

Remark 1. *By a small perturbation, any path can be made good and stable (if endpoints are generic, we may also require that the endpoints remain fixed).*

Definition 4. *A path from x to x' is called a braid if the points representing x are the same as those representing x' (possibly, in different orders); if x coincides with x' with respect to order, then such a braid is called pure.*

There is an obvious concatenation structure on the set of braids: for paths $\gamma_{x,x'}$ and $\gamma'_{x',x''}$, the concatenation is defined as a path $\gamma''_{x,x''}$ such that $\gamma''(t) = \gamma(2t)$ for $t \in [0, \frac{1}{2}]$ and $\gamma''(t) = \gamma'(2t - 1)$ for $t \in [\frac{1}{2}, 1]$; this path can be smoothed in the neighbourhood of $t = \frac{1}{2}$; the result of such smoothing is well defined up to isotopy.

Thus, the sets of braids and pure braids (for fixed x) admit a group structure. This latter group is certainly isomorphic to the fundamental group $\pi_1(C'_n(\mathbb{R}^{k-1}))$. The former group is isomorphic to the fundamental group of the quotient space by the action of the permutation group.

3 The realisability of G_{k+1}^k

The main idea of the proof of Theorem 1 is to associate with every word in G_{k+1}^k a braid in $\tilde{C}'_{k+1}(\mathbb{R}P^{k-1})$.

Let us start with the main construction from [10].

With each good and stable path from $PB_n(\mathbb{R}P^2)$ we associate an element of the group G_n^k as follows. We enumerate all singular values of our path $0 < t_1 < \dots < t_l < 1$ (we assume that 0 and 1 are not singular). For each singular value t_p we have a set m_p of k indices corresponding to the numbers of points which are not in general position. With this value we associate the letter a_{m_p} . With the whole path γ (braid) we associate the product $f(\gamma) = a_{m_1} \dots a_{m_l}$.

Theorem 3. [10] *The map f takes isotopic braids to equal elements of the group G_n^k . For pure braids, the map f is a homomorphism $f : \pi_1(C'_n(\mathbb{R}P^2)) \rightarrow G_n^k$.*

We are interested in the case $k = 3$.

Now we claim that

Every word from G_{k+1}^k can be realised by a path of the above form.

Note that if we replace $\mathbb{R}P^{k-1}$ with \mathbb{R}^{k-1} , the corresponding statement will fail. Namely, starting with the configuration of four points, $x_i, i = 1, \dots, 4$, where x_1, x_2, x_3 form a triangle and x_4 lies inside this triangle, we see that any path starting from this configuration will lead to a word starting from $a_{124}, a_{134}, a_{234}$ but not from a_{123} . In some sense the point 4 is “locked” and the points are not in the same position.

The following well known theorem (see, e.g., [2]) plays a crucial role in the construction

Theorem 4. *For any two sets of $k + 1$ points in general position in $\mathbb{R}P^{k-1}$, (x_1, \dots, x_{k+1}) and (y_1, \dots, y_{k+1}) there is an action of $PGL(k, \mathbb{R})$ taking all x_i to y_i .*

For us, this will mean that there is no difference between all possible “non-degenerate starting positions” for $k + 1$ points in $\mathbb{R}P^k$.




Figure 1: The “locked” position for the move a_{123}

We shall deal with paths in $\tilde{C}'_{k+1}(\mathbb{R}P^{k-1})$ similar to braids. Namely, we shall fix a set of 2^{k-1} points such that all paths will start and end at these points.

We shall denote homogeneous coordinates in $\mathbb{R}P^{k-1}$ by $(a_1 : \dots : a_k)$ in contrast to points (which we denote by (x_1, \dots, x_{k+1})).

3.1 Constructing a braid from a word in G_{k+1}^k

Our main goal is to construct a braid by a word. To this end, we need a base point for the braid. For the sake of convenience, we shall use not one, but rather 2^{k-1} reference points. For the first k points $y_1 = (1 : 0 : \dots : 0), \dots, y_k = (0 : \dots : 0 : 1)$ fixed, we will have 2^{k-1} possibilities for the choice of the last point. Namely, let us consider all possible strings of length k of ± 1 with the last coordinate $+1$: $(1, 1, \dots, 1, 1), (1, \dots, 1, -1, 1), \dots, (-1, -1, \dots, -1, 1)$ with $a_k \sim 1$. We shall denote these points by y_s where s records the first $(k-1)$ coordinates of the point.

Now, for each string s of length k of ± 1 , we set $z_s = (y_1, y_2, \dots, y_k, y_s)$.

The following lemma is evident.

Lemma 1. *For every point $z \in \mathbb{R}P^{k-1}$ with projective coordinates $(a_1(z) : \dots : a_{k-1}(z) : 1)$, let $\tilde{z} = (\text{sign}(a_1(z)) : \text{sign}(a_2(z)) : \dots : \text{sign}(a_{k-1}(z)) : 1)$. Then there is a path between (y_1, \dots, y_k, z) and $(y_1, \dots, y_k, \tilde{z})$ in $\tilde{C}'_{k+1}(\mathbb{R}P^k)$ with the first points y_1, \dots, y_k fixed, and the corresponding path in \tilde{C}'_{k+1} is void.*

Proof. Indeed, it suffices just to connect z to \tilde{z} by a geodesic. □

From this we easily deduce the following

Lemma 2. *Every point $y \in \tilde{C}'_{k+1}(\mathbb{R}P^{k-1})$ can be connected by a void path to some (y_1, \dots, y_k, y_s) for some s .*

Proof. Indeed, the void path can be constructed in two steps. At the first step, we construct a path which moves both y_k and y_{k+1} , so that y_k becomes $(0 : \dots : 0 : 1)$, and at the second step, we use Lemma 1. To realise the first step, we just use linear maps which keep the hyperplane $a_k = 0$ fixed. □

The lemma below shows that the path mentioned in Lemma 2 is unique up to homotopy.

Lemma 3. *Let γ be a closed path in $\tilde{C}'_{k+1}(\mathbb{R}P^{k-1})$ such that the word $f(\gamma)$ is empty. Then γ is homotopic to the trivial braid.*

Proof. In \tilde{C}'_{k+1} , we deal with the motion of points, where all but x_k, x_{k+1} are fixed.

Consider the projective hyperplane \mathcal{P}_1 passing through x_1, \dots, x_{k-1} given by the equation $a_k = 0$.

We know that none of the points x_k, x_{k+1} is allowed to belong to \mathcal{P}_1 . Hence, we may fix the last coordinate $a_k(x_k) = a_k(x_{k+1}) = 1$.

Now, we may pass to the affine coordinates of these two points (still to be denoted by a_1, \dots, a_k).

Now, the condition $\forall i = 1, \dots, k-1 : a_j(x_k) \neq a_j(x_{k+1})$ follows from the fact that the points $x_1, \dots, \hat{x}_j, x_{k+1}$ are generic.

This means that $\forall i = 1, \dots, k-1$ the sign of $a_i(x_k) - a_i(x_{k+1})$ remains fixed.

Now, the motion of points x_k, x_{k+1} is determined by their coordinates a_1, \dots, a_j , and since their signs are fixed, the configuration space for this motions is simply connected.

This means that the configuration space for γ is the direct product of the two configuration spaces: the one for the point x_k and the one for the point x_{k+1} . Having x_1, \dots, x_{k-1} fixed on $\mathbb{R}P^{k-1}$, the configuration space of admissible positions for x_k is nothing but the hemisphere; for fixed x_k , the admissible positions for x_{k+1} are regulated by inequalities in coordinates; this space is also simply connected. □

3.2 The surjectivity

In order to prove that some map is a bijection, it is crucial to understand the surjectivity. For the case of braid groups and words in G_n^k , this surjectivity is attained in the case of $n = k + 1$.

Namely, our next strategy is as follows. Having a word in G_{k+1}^k , we shall associate with this word a path in $\tilde{C}'_{k+1}(\mathbb{R}P^{k-1})$. After each letter, we shall get to (y_1, \dots, y_k, y_s) for some s .

Let us start from $(y_1, \dots, y_k, y_{1, \dots, 1})$.

After making the final step, one can calculate the coordinate of the $(k + 1)$ -th points. They will be governed by Lemma 6 (see ahead). As we shall see later, those words we have to check for the solution of the word problem in G_{k+1}^k , will lead us to closed paths, i.e., pure braids.

Let us be more detailed.

Lemma 4. *Given a non-singular set of points y in $\mathbb{R}P^{k-1}$. Then for every set of k numbers $i_1, i_2, \dots, i_k \in [n]$, there exists a path $y_{i_1 \dots i_k}(t) = y(t)$ in $C'_n(\mathbb{R}P^{k-1})$, having $y(0) = y(1) = y$ as the starting point and the final point and with only one singular moment corresponding to the numbers i_1, \dots, i_k which encode the points not in general position; moreover, we may assume that at this moment all points except i_1 , are fixed during the path.*

Moreover, the set of paths possessing this property is connected: any other path $\tilde{y}(t)$, possessing the above properties, is homotopic to $y(t)$ in this class.

Proof. Indeed, for the first statement of the Lemma, it suffices to construct a path for some initial position of points and then apply Theorem 4.

For the second statement, let us take two different paths γ_1 and γ_2 satisfying the conditions of the Lemma. By a small perturbation, we may assume that for both of them, $t = \frac{1}{2}$ is a singular moment with the same position of y_{i_1} .

Now, we can contract the loop formed by $\gamma_1|_{t \in [\frac{1}{2}, 1]}$ and the inverse of $\gamma_2|_{t \in [\frac{1}{2}, 1]}$ by using Lemma 3 as this is a small perturbation of a void braid. We are left with $\gamma_1|_{t \in [0, \frac{1}{2}]}$ and the inverse of $\gamma_2|_{t \in [0, \frac{1}{2}]}$ which is contractible by Lemma 3 again. \square

Remark 2. Note that in the above lemma, we deal with the space $C'_n(\mathbb{R}P^{k-1})$, not with $\tilde{C}'_n(\mathbb{R}P^{k-1})$. On the other hand, we may always choose $i_1 \in \{k, k+1\}$; hence, the path in question can be chosen in $\tilde{C}'(\mathbb{R}P^{k-1})$.

Now, for every subset $m \subset [n]$, $Card(m) = k+1$ we can create a path p_m starting from any of the base points listed above and ending at the corresponding basepoints.

Now, we construct our path step-by step by applying Lemma 4 and returning to some of base points by using Lemma 2.

From [10], we can easily get the following

Lemma 5. Let i_1, \dots, i_{k+1} be some permutation of $1, \dots, k+1$. Then the concatenation of paths $p_{i_1 i_2 \dots i_k} p_{i_1 i_3 i_4 \dots i_{k+1}} \dots p_{i_2 i_3 \dots i_k}$ is homotopic to the concatenation of paths in the inverse order $p_{i_2 i_3 \dots i_k} \dots p_{i_1 i_3 i_4 \dots i_{k+1}} p_{i_1 i_2 \dots i_k}$.

Proof. Indeed, in [10], some homotopy corresponding to the above mentioned relation corresponding to some permutation is discussed. However, since all basepoints are similar to each other as discussed above, we can transform the homotopy from [10] to the homotopy for any permutation. \square

Lemma 6. For the path starting from the point (y_1, \dots, y_k, y_s) constructed as in Lemma 4 for the set of indices j , we get to the point $(y_1, \dots, y_k, y_{s'})$ such that:

1. if $j = 1, \dots, k$, then s' differs from s only in coordinate a_j ;
2. if $j = k+1$, all coordinates of s' differ from those coordinates of s by sign.

Remark 3. Note that in general the path constructed in the above lemma is generally not a braid, but almost a braid: for the fixed position of the initial point, there are only infinitely many possibilities for the position of the final point.

Denote the map from words in G_{k+1}^k to paths between basepoints by g .

By construction, we see that for every word w we have $f(g(w)) = w \in G_{k+1}^k$.

Now, we define the group \tilde{G}_{k+1}^k as the subgroup of G_{k+1} which is taken by g to braids, i.e., to those paths with coinciding initial and final points. From lemma 6, we see that this is a subgroup of index $(k-1)$: there are exactly $(k-1)$ coordinates.

3.3 Equal words lead to homotopic paths

Let us pass to the proof of Theorem 2. Our next goal is to see that equal words can originate only from homotopic paths.

To this end, we shall first show that the map f from Theorem 3 is an isomorphism for $n = k + 1$. To perform this goal, we should construct the inverse map $g : \tilde{G}_{k+1}^k \rightarrow \pi_1(\tilde{C}'_{k+1}(\mathbb{R}P(k-1)))$.

Note that for $k = 3$ we deal with the pure braids $PB_4(\mathbb{R}P^2)$.

Let us fix a point $x \in C'_4(\mathbb{R}P^2)$. With each generator $a_m, m \subset [n], Card(m) = k$ we associate a path $g(m) = y_m(t)$, described in Lemma 4. This path is not a braid: we can return to any of the 2^{k-1} base points. However, once we take the concatenation of paths corresponding to \tilde{G}_{k+1}^k , we get a braid.

By definition of the map f , we have $f(g(a_m)) = a_m$. Thus, we have chosen that the map f is a surjection.

Now, let us prove that the kernel of the map f is trivial. Indeed, assume there is a pure braid γ such that $f(\gamma) = 1 \in G_{k+1}^k$. We assume that γ is good and stable. If this path has l critical points, then we have the word corresponding to it $a_{m_1} \cdots a_{m_l} \in G_{k+1}^k$.

Let us perform the transformation $f(\gamma) \rightarrow 1$ by applying the relations of G_{k+1}^k to it and transforming the path γ respectively. For each relation of the sort $a_m a_m = 1$ for a generator a_m of the group G_{k+1}^k , we see that the path γ contains two segments whose concatenation is homotopic to the trivial loop (as follows from the second statement of Lemma 4).

Whenever we have a relation of length $2k + 2$ in the group G_{k+1} , we use the Lemma 5 to perform the homotopy of the loops.

Thus, we have proved that if the word $f(\gamma)$ corresponding to a braid $\gamma \in G_{k+1}^k$ is equal to 1 in G_{k+1}^k then the braid γ is isotopic to a braid γ' such that the word corresponding to it is empty. Now, by Lemma 3, this braid is homotopic to the trivial braid.

4 The group H_k and the algebraic lemma

The aim of the present section is to reduce the word problem in G_{k+1}^k to the word problem in a certain subgroup of it, denoted by H_k .

Let us rename all generators of G_{k+1}^k lexicographically: $b_1 = a_{1,2,\dots,k}, \dots, b_{k+1} = a_{2,3,\dots,k+1}$.

Let H_k be the subgroup of G_{k+1}^k consisting of all elements $x \in G_{k+1}^k$ that can be represented by words with no occurrences of the last letter b_{k+1} .

Our task is to understand whether a word in G_{k+1}^k represents an element in H . To this end, we recall the map from [11]. Consider the group $F_k = \mathbb{Z}_2^{*2^k} = \langle c_m | c_m^2 = 1 \rangle$, where all generators c_m are indexed by k -element strings of 0 and 1 with only relations being that the square of each generator is equal to 1. We shall construct a map¹ from G_{k+1}^k to F_k as follows.

Take a word w in generators of G_{k+1}^k and list all occurrences of the last letter $b_{k+1} = a_{2,\dots,k+1}$ in this word. With any such occurrence we first associate the string of indices 0, 1 of length k . The j -th index is the number of letters b_j preceding this occurrence of b_{k+1} modulo 2. Thus, we get a string of length k .

¹This map becomes a homomorphism when restricted to a finite index subgroup

Now, we identify “opposite pairs” of strings: we set $(x_1, \dots, x_k) \sim (x_1 + 1, \dots, x_k + 1)$. Now, we may think that the last (k -th) element of our string is always 0, so, we can restrict ourselves with $(x_1, \dots, x_{k-1}, 0)$. Such a string of length $k - 1$ is called the *index* of the occurency of b_{k+1} .

Having this done, we associate with each occurency of b_{k+1} having index m the generator c_m of F_k . With the word w , we associate the word $f(w)$ equal to the product of all generators c_m in order.

In [11], the following Lemma is proved

Lemma 7. *The map $f : G_{k+1}^k \rightarrow F_k$ is well defined.*

Now, let us prove the following

Lemma 8. *If $f(w) = 1$ then $w \in H_k$.*

In other words, the free group F_k yields the only obstruction for an element from G_{k+1}^k to have a presentation with no occurency of the last letter.

Proof. Let w be a word such that $f(w) = 1$. If $f(w)$ is empty, then there is nothing to prove. Since f is trivial in F_k , it contains some two adjacent occurencies of the same generator b_x, \dots, b_x . This means that in the initial word w is of the type $w = Ab_{k+1}Bb_{k+1}C$, where A and C are some words, and B contains no occurencies of b_{k+1} and the number of occurencies of b_1, b_2, \dots, b_k in B are of the same parity.

Our aim is to show that w is equal to a word with smaller number of b_{k+1} in G_{k+1}^k . Then we will be able to induct on the number of b_{k+1} until we get a word without b_{k+1} .

Thus, it suffices for us to show that $b_{k+1}Bb_{k+1}$ is equal to a word from H_k . We shall induct on the length of B . Without loss of generality, we may assume that B is reduced, i.e., it does not contain adjacent $b_j b_j$.

Let us read the word B from the very beginning $B = b_{i_1} b_{i_2} \dots$. If all elements i_1, i_2, \dots are distinct, then, having in mind that the number of occurencies of all generators in B should be of the same parity, we conclude that $b_{k+1}B = B^{-1}b_{k+1}$, hence $b_{k+1}Bb_{k+1} = B^{-1}b_{k+1}b_{k+1} = B^{-1}$ is a word without occurencies of b_{k+1} .

Now assume $i_1 = i_p$. Then we have $b_{k+1}B = b_{k+1}b_{i_1} \dots b_{i_{p-1}} b_{i_1} B'$. Now we collect all indices distinct from $i_1, \dots, i_{p-1}, k + 1$ and write the word P containing exactly one generator for each index (the order does not matter). Then the word $W = Pb_{k+1}b_{i_1} \dots b_{i_{p-1}}$ contains any letter exactly once and we can invert the word W as follows: $W^{-1} = b_{i_{p-1}} \dots b_{i_1} b_{k+1} P$. Thus, $b_{k+1}B = P^{-1}(Pb_{k+1}b_{i_1} \dots b_{i_{p-1}})b_{i_1} B' = P^{-1}b_{i_{p-1}} \dots b_{i_1} b_{k+1} P^{-1}b_{i_1} B'$.

We know that the letters in P (hence, those in P^{-1}) do not contain b_{i_1} . Thus, the word $P^{-1}b_{i_1}$ consists of different letters. Now we perform the same trick: create the word $Q = b_{i_2} b_{i_3} \dots b_{i_{p-1}}$ consisting of remaining letters from $\{1, \dots, k\}$, we get:

$$\begin{aligned} & P^{-1}b_{i_{p-1}} \dots b_{i_1} b_{k+1} P^{-1}b_{i_1} B' \\ &= P^{-1}b_{i_{p-1}} \dots b_{i_1} Q^{-1} Q b_{k+1} P^{-1}b_{i_1} B' \\ &= P^{-1}b_{i_{p-1}} \dots b_{i_1} Q^{-1} b_{i_1} P b_{k+1} Q B'. \end{aligned}$$

Thus, we have moved b_{k+1} to the right and between the two occurrences of the letter b_{k+1} , we replaced $b_{i_1} \cdots b_{i_{p-1}} b_{i_1}$ with just $b_{i_2} \cdots b_{i_{p-1}}$, thus, we have shortened the distance between the two adjacent occurrences of b_{k+1} .

Arguing as above, we will finally cancel these two letters and perform the induction step. □

Theorem 5. *If the word problem in H_k is solvable, then the word problem in G_{k+1}^k is solvable. Moreover, the solution of the latter is constructive once the solution of the former is constructive.*

Proof. Indeed, having a word w in generators of G_{k+1}^k , we can look at the image of this word by the map f in the group H_k . If it is not equal to 1, then, from [11], it follows that w is non-trivial, otherwise we can construct a word \tilde{w} in H_k equal to w in G_{k+1}^k . □

Certainly, to be able to solve the word problem in H_k , one needs to know a presentation for H_k . It is natural to take b_1, \dots, b_k for generators of H_k . Obviously, they satisfy the relations $b_j^2 = 1$ for every j .

To understand the remaining relations for different k , we shall need geometrical arguments.

5 Concluding remarks

We have completely constructed the isomorphism between the (finite index subgroup) of the group G_{k+1}^k and a fundamental group of some configuration space.

This completely solves the word problem in G_4^3 for braid groups in projective spaces are very well studied, see, e.g., [1, 3, 4, 5]. The same can be said about the conjugacy problem in \tilde{G}_4^3 .

Besides, we have seen that the word problem for the case of general G_{k+1}^k can be reduced to the case of H_k .

In a subsequent paper, we shall completely describe the relations in H_k by geometric reasons and apply it to the word problem in G_5^4 . Here we just mention that it was proved by A.B.Karpov (unpublished) that the only relations in H_3 are $a^2 = b^2 = c^2$ which also follows from the geometrical techniques of the present paper.

The main open question which remains unsolved is how to construct a configuration space which can realise G_n^k for $n > k + 1$. Even in the case of G_4^2 this problem seems very attractive though the word and conjugacy problems for G_n^2 can be solved by algebraic methods, see [7].

The word problem and the partial case of conjugacy problem in G_4^3 was first solved by A.B.Karpov, but the author has not yet seen any complete text of it.

It would be very interesting to compare the approach of G_n^k with various generalizations of braid groups, e.g., Manin-Schechtmann groups [13, 6].

The author is very grateful to I.M.Nikonov, L.A.Bokut' and Jie Wu for extremely useful discussions and comments.

References

- [1] J. van Buskirk, Braid groups of compact 2-manifolds with elements of finite order. *Trans. Amer. Math. Soc.*, **122**, P.8197 (1966).
- [2] A.J.Berrick, F.R.Cohen, Y.L.Wong, J.Wu, Configurations, braids, and homotopy groups, *Journal AMS*, **19**, 2, Pp. 265-326 (2005).
- [3] D. L. Gonçalves and J. Guaschi, The braid groups of the projective plane, *Algebr. Geom. Topol.*, **4**, P. 757780 (2004)
- [4] D. L. Gonçalves and J. Guaschi, The braid groups of the projective plane and the FadellNeuwirth short exact sequence, *Geom. Dedicata*, **130**, P. 93107, (2007)
- [5] D.L. Gonçalves, J. Guaschi, The lower central and derived series of the braid groups of the projective plane. *J. Algebra*, **331**, P. 96129 (2011).
- [6] M.Kapranov, V.Voevodsky, Braided Monoidal 2-Categories and Manin-Schechtmann Higher Braid Groups, *Journal of Pure and Applied Mathematics*, N. 92, 1994, P. 241-167.
- [7] V.O.Manturov, On the Groups G_n^2 and Coxeter Groups, *Russ. Math.Surv.*, 2017, **72:2**, P.378-380, <http://arxiv.org/abs/1512.09273>.
- [8] V.O.Manturov, Non-Reidemeister Knot Theory and Its Applications in Dynamical Systems, Geometry, and Topology, <http://arxiv.org/abs/1501.05208>
- [9] V.O.Manturov, Invariants of Classical Braids Valued in G_n^2 , <http://arxiv.org/abs/1611.07434v2>.
- [10] V.O.Manturov, The Groups G_n^k and fundamental groups of configuration spaces, *J. Knot Theory & Ramifications*, 2015, Vol.26
- [11] V.O.Manturov, I.M.Nikonov, On braids and groups G_n^k *J. Knot Theory & Ramifications*, 2015, Vol. 24, No.13.
- [12] S.Kim, V.O.Manturov, The Braid Groups and Imaginary Generators, <http://arxiv.org/abs/1612.03486v1>.
- [13] Y.I.Manin, V.V.Schechtmann, Arrangements of Hyperplanes, higher braid groups and higher Bruhat orders, *Adv. Stud. Pure Math.*, 17 (1990), P. 289-308