

# 0-1 matrices with zero trace whose squares are 0-1 matrices

Zejun Huang,<sup>\*</sup> Zhenhua Lyu,<sup>†</sup>

## Abstract

In this paper, we determine the maximum number of nonzero entries in 0-1 matrices of order  $n$  with zero trace whose squares are 0-1 matrices when  $n \geq 8$ . The extremal matrices attaining this maximum number are also characterized.

**Key words:** 0-1 matrix, digraph, Turán problem, walk

**AMS subject classifications:** 15A36, 05C35

## 1 Introduction

Denote by  $M_n\{0, 1\}$  the set of 0-1 matrices of order  $n$ . In 2007, Zhan [11] proposed the following problem.

**Problem 1.** *Given two integers  $n$  and  $k$ , what is the maximum number of nonzero entries in a matrix  $A \in M_n\{0, 1\}$  such that  $A^k \in M_n\{0, 1\}$ , and what are the extremal matrices attaining this maximum number?*

Wu [10] solved the case  $k = 2$ . Huang and Zhan [6] solved the case  $k \geq n - 1$  and they attained the maximum number for the case  $k = n - 2$  and  $n - 3$ . The authors of [5] solved the case  $k \geq 5$  and they attained the maximum number for the case  $k = 4$ . In this paper, we consider the following related problem.

**Problem 2.** *Given integers  $n$  and  $k$ , determine the maximum number of nonzero entries in a matrix  $A \in M_n\{0, 1\}$  such that*

$$\text{tr}(A) = 0 \text{ and } A^k \in M_n\{0, 1\}.$$

*Characterize the extremal matrices that attain this maximum number.*

<sup>\*</sup>Institute of Mathematics, Hunan University, Changsha 410082, P.R. China. (mathzejun@gmail.com)

<sup>†</sup>College of Mathematics and Econometrics, Hunan University, Changsha 410082, P.R. China. (lyuzhh@outlook.com)

We solve the case  $k = 2$  for Problem 2 in this paper. Our approach is transferring the problem to an equivalent problem on digraphs and applying detail analysis on the structures of certain digraphs. We need the following definitions and notations.

We abbreviate directed walks, directed paths and directed cycles as walks, paths and cycles, respectively. The *length* of a walk, path or cycle is its number of arcs. The number of vertices in a digraph is called its *order* and the number of arcs its *size*. Let  $u, w$  be two vertices. The notation  $(u, w)$  or  $u \rightarrow w$  means there exists an arc from  $u$  to  $w$ ;  $u \nrightarrow w$  means there exists no arc from  $u$  to  $w$ ;  $u \leftrightarrow w$  means both  $u \rightarrow w$  and  $w \rightarrow u$ . If there exists an arc from  $u$  to  $w$ , then  $u$  is a *predecessor* of  $w$ , and  $w$  is a *successor* of  $u$ .

Let  $D = (\mathcal{V}, \mathcal{A})$  be a digraph with vertex set  $\mathcal{V} = \{v_1, v_2, \dots, v_n\}$  and arc set  $\mathcal{A}$ . Its *adjacency matrix*  $A_D = (a_{ij})$  is defined by

$$a_{ij} = \begin{cases} 1, & (v_i, v_j) \in \mathcal{A}; \\ 0, & \text{otherwise.} \end{cases} \quad (1.1)$$

Conversely, given an  $n \times n$  0-1 matrix  $A = (a_{ij})$ , we can define its digraph  $D(A) = (\mathcal{V}, \mathcal{A})$  on vertices  $v_1, v_2, \dots, v_n$  by (1.1), whose adjacency matrix is  $A$ .

For a subset  $X \subset \mathcal{V}$ ,  $D(X)$  denotes the subdigraph of  $D$  induced by  $X$ . The outdegree  $d^+(u)$  is the number of arcs with tail  $u$  and the indegree  $d^-(u)$  is the number of arcs with head  $u$ . For convenience, if  $X = \{x\}$  is a singleton, it will be abbreviated as  $x$ .

For  $S, T \subset \mathcal{V}$ , denote by  $\mathcal{A}(S, T)$  the set of arcs from  $S$  to  $T$ . If  $S = T$  we use  $\mathcal{A}(S)$  instead of  $\mathcal{A}(S, S)$ . Let  $e(S, T) = |\mathcal{A}(S, T)|$  and  $e(D) = |\mathcal{A}(\mathcal{V})|$ .  $S \rightarrow T$  means for every vertex  $i \in T$  there exists exactly one vertex  $j \in S$  such that  $j \rightarrow i$ ;  $S \nrightarrow T$  means there is no arc from  $S$  to  $T$ . If every vertex in  $S$  has exactly one successor in  $T$  and each vertex in  $T$  has exactly one predecessor in  $S$ , we say  $S$  *matches*  $T$ . Note that  $S$  matching  $T$  indicates  $|S| = |T|$ .

For  $W \subset \mathcal{V}$ , denote by

$$N_W^+(u) = \{x \in W \mid (u, x) \in \mathcal{A}\},$$

$$N_W^-(u) = \{x \in W \mid (x, u) \in \mathcal{A}\},$$

and

$$N_W^+(S) = \bigcup_{u \in S} N_W^+(u),$$

where  $S \subset \mathcal{V}$ . When  $W = \mathcal{V}$ , we simply write  $N^+(u)$ ,  $N^-(u)$  and  $N^+(S)$  respectively.


The following problem is equivalent with Problem 1 and Problem 2.

**Problem 3.** *Given two integers  $n$  and  $k$ , determine the maximum size of digraphs of order  $n$  avoiding two distinct directed walks of a given length  $k$  with the same initial and terminal vertices. Characterize the extremal digraphs attaining this maximum size.*

Using the digraphs of 0-1 matrices, we see that for a matrix  $A \in M_n\{0, 1\}$ ,  $A^k \in M_n\{0, 1\}$  if and only if  $D(A)$  avoids distinct directed walks of length  $k$  with the same initial vertex and the same terminal vertex. Hence, for digraphs allow loops but donot allow parallel arcs, Problem 3 is equivalent with Problem 1; for strict digraphs, i.e., digraphs do not allow loops or parallel arcs, Problem 3 is equivalent with Problem 2.

For strict digraphs, the solution to Problem 3 for the case  $k \geq 5$  follows straightforward from [5, 6], since the extremal digraphs in [5, 6] are loopless. In this paper we consider the case  $k = 2$  for Problem 3 on strict digraphs.

In what follows digraphs are strict. Given a family of digraphs  $\mathcal{H}$  and a digraph  $D$ ,  $D$  is  $\mathcal{H}$ -free if  $D$  contains no member of  $\mathcal{H}$  as its subgraph. Let  $\mathcal{F} = \{P_{2,2}, C_{2,2}\}$  where  $P_{2,2}$  and  $C_{2,2}$  are defined as follows.




It is obvious that a digraph avoids two distinct 2-walks with the same initial and terminal vertices if and only if it is  $\mathcal{F}$ -free. Denote by  $ex(n)$  and  $EX(n)$  the maximum size of  $\mathcal{F}$ -free digraphs of order  $n$  and the set of  $\mathcal{F}$ -free digraphs of order  $n$  attaining the maximum size, respectively.

Note that Problem 3 is a Turán type problem, which concerns the study of extremal graphs that avoid given subgraphs. Turán problem is a hot topic in graph theory with a long history; see [1, 2, 3, 7, 8, 9]. In [4] the authors studied a closely related Turán problem. They determined the maximum size of  $P_{2,2}$ -free digraphs as well as the extremal  $P_{2,2}$ -free digraphs.

## 2 Main results


If a digraph  $D$  is acyclic and there is a vertex  $u$  such that there is a unique directed path from  $u$  to any other vertex, then we say  $D$  is an *arborescence* with root  $u$ . If the maximum length of these paths is at most  $r$ , then we say  $D$  is an  *$r$ -arborescence*. Moreover, if  $D$  is a 1-arborescence, we also say  $D$  is an *out-star*. Throughout this article, we assume each arborescence has more than one vertices.

We will use  $S(x)$ ,  $S_x(y)$ ,  $T(x)$  and  $T(x, y)$  to denote the following digraphs, whose orders will be clear from the context. Note that  $S(x)$  is an out-star with root  $x$ ;  $S_x(y)$  is the union of a 2-cycle  $x \leftrightarrow y$  and an out-star with root  $y$ ;  $T(x)$  is a 2-arborescence with root  $x$ ;  $T(x, y)$  is the union of a 2-cycle  $x \leftrightarrow y$  and two 2-arborescences with roots  $x$  and  $y$ .



Here we denote a single vertex by  $C_1$ . The digraph obtained by taking the union of digraphs  $D$  and  $H$  with disjoint vertex sets is the disjoint union, written  $D + H$ . In general,  $mD$  is the digraph consisting of  $m$  pairwise disjoint copies of  $D$ . Denote by  $D'$  the reverse of  $D$ , which is obtained by reversing the directions of all arcs of  $D$ . The reverse of an out-star is called an *in-star*.

Now we present the following six classes of digraphs on  $n$  vertices, where  $n$  is even for  $D_1$  and odd for the others. Each of these diagrams has vertex partition  $\mathcal{V}_1 \cup \mathcal{V}_2$  with  $|\mathcal{V}_1| = \lfloor \frac{n}{2} \rfloor + 1$  and  $|\mathcal{V}_2| = \lceil \frac{n}{2} \rceil - 1$ .



In  $D_1$ ,  $D(\mathcal{V}_1) = T(y_1, y_2)$ ;  $D(\mathcal{V}_2)$  is empty;  $f$  means

$$\mathcal{V}_1 \setminus \{y_1, y_2\} \text{ matches } \mathcal{V}_2;$$

$g$  means

$$u \rightarrow \mathcal{V}_1 \text{ for all } u \in \mathcal{V}_2.$$

In  $D_2$ ,  $D(\mathcal{V}_1) = S_y(x)$ ;  $D(\mathcal{V}_2)$  is the disjoint union of  $T'(w)$  and isolated vertices, where  $T'(w)$  may vanish.  $\mathcal{V}_2$  is partitioned as  $\mathcal{V}_2 = \mathcal{V}_3 \cup \mathcal{V}_4$ , where  $\mathcal{V}_3$  consists of all the isolated vertices of  $\mathcal{V}_2$  and  $|\mathcal{V}_3| \geq 1$ ;  $f$  means

$$\mathcal{V}_1 \setminus \{y, x\} \text{ matches } \mathcal{V}_2 \setminus \{w\} \text{ and } x \rightarrow w;$$

$g$  means

$$u \rightarrow \mathcal{V}_1 \text{ for all } u \in \mathcal{V}_3 \text{ and } u \rightarrow \mathcal{V}_1 \setminus \{x\} \text{ for all } u \in \mathcal{V}_4.$$

In  $D_3$ ,  $D(\mathcal{V}_1) = T(y)$ ;  $D(\mathcal{V}_2)$  is empty;  $f$  means

$$\mathcal{V}_1 \setminus \{y\} \text{ matches } \mathcal{V}_2;$$

$g$  means

$$u \rightarrow \mathcal{V}_1 \text{ for all } u \in \mathcal{V}_2.$$

In  $D_4$ ,  $D(\mathcal{V}_1) = S(x)$ ;  $D(\mathcal{V}_2)$  is the disjoint union of  $T'(w)$ , in-stars and isolated vertices, and each of them may vanish.  $\mathcal{V}_2$  is partitioned as  $\mathcal{V}_2 = \mathcal{V}_3 \cup \mathcal{V}_4$ , where  $\mathcal{V}_3$  consists of all isolated vertices of  $\mathcal{V}_2$  and the roots of  $T'(w)$  and the in-stars;  $f$  means

$$\mathcal{V}_1 \setminus \{x, y\} \text{ matches } \mathcal{V}_2 \setminus \{w\} \text{ and } x \rightarrow w,$$

where  $y$  is an arbitrary vertex in  $\mathcal{V}_1 \setminus \{x\}$ ;  $g$  means

$$u \rightarrow \mathcal{V}_1 \text{ for all } u \in \mathcal{V}_3 \text{ and } u \rightarrow \mathcal{V}_1 \setminus \{x\} \text{ for all } u \in \mathcal{V}_4.$$

In  $D_5$ ,  $D(\mathcal{V}_1) = T(y_1, y_2)$ ;  $D(\mathcal{V}_2)$  is empty;  $f$  means

$$\mathcal{V}_1 \setminus \{y_1, y_2\} \text{ matches } \mathcal{V}_2 \setminus \{x\},$$

where  $x$  is an arbitrary vertex in  $\mathcal{V}_2$ ;  $g$  means

$$u \rightarrow \mathcal{V}_1 \text{ for all } u \in \mathcal{V}_2.$$

In  $D_6$ ,  $D(\mathcal{V}_1) = S_y(x)$ ;  $D(\mathcal{V}_2)$  is the disjoint union of  $T'(w, z)$  and isolated vertices.  $\mathcal{V}_2$  is partitioned as  $\mathcal{V}_2 = \mathcal{V}_3 \cup \mathcal{V}_4$ , where  $\mathcal{V}_3$  consists of all the isolated vertices of  $\mathcal{V}_2$  and  $|\mathcal{V}_3| \geq 1$ ;  $f$  means

$$\mathcal{V}_1 \setminus \{y, y', x\} \text{ matches } \mathcal{V}_2 \setminus \{z, w\} \text{ and } x \rightarrow w,$$

where  $y'$  is an arbitrary vertex in  $\mathcal{V}_1 \setminus \{y, x\}$ ;  $g$  means

$$u \rightarrow \mathcal{V}_1 \text{ for all } u \in \mathcal{V}_3 \text{ and } u \rightarrow \mathcal{V}_1 \setminus \{x\} \text{ for all } u \in \mathcal{V}_4.$$

We say that digraph  $D$  is an isomorphism of  $H$  if there exists a bijection  $f: \mathcal{V}(D) \rightarrow \mathcal{V}(H)$  such that  $(u, v) \in \mathcal{A}(D)$  if and only if  $(f(u), f(v)) \in \mathcal{A}(H)$ . Now we post our main result as follows.

**Theorem 4.** *Let  $D$  be a digraph on  $n$  vertices with  $n \geq 8$ . Then*

$$ex(n) = \lfloor \frac{n^2 + 4n}{4} \rfloor - 1.$$

*Moreover,  $D \in EX(n)$  if and only if*

- (1)  $n$  is even, and  $D$  or  $D'$  is an isomorphism of  $D_1$ ;  
(2)  $n$  is odd, and  $D$  or  $D'$  is an isomorphism of  $D_i$  with  $i \in \{2, \dots, 6\}$ .

Equivalently, we have the following solution to Problem 2 when  $k = 2$ .

**Theorem 4\***. Let  $n \geq 8$  be an integer. Suppose  $A \in M_n\{0, 1\}$  such that

$$\text{tr}(A) = 0 \text{ and } A^2 \in M_n\{0, 1\}.$$

Then  $A$  has at most  $\text{ex}(n)$  nonzero entries, and  $A$  has  $\text{ex}(n)$  nonzero entries if and only if  $D(A)$  or its reverse is an isomorphism of  $D_i$  with  $i \in \{1, \dots, 6\}$ .

### 3 Lemmas

In this section we give preparatory lemmas for the proof of the main theorem.

**Lemma 5.** Let  $n \geq 3$  be a positive integer. Then  $D_i$  is  $\mathcal{F}$ -free for  $i = 1, 2, \dots, 6$ . Moreover, we obtain

$$\text{ex}(n) \geq \begin{cases} \frac{n^2+4n-5}{4}, & n \text{ is odd}; \\ \frac{n^2+4n-4}{4}, & n \text{ is even}. \end{cases} \quad (3.1)$$

*Proof.* Suppose  $D \in \{D_1, D_2, \dots, D_6\}$  contains  $u_1 \rightarrow u_2 \rightarrow u_4$  and  $u_1 \rightarrow u_3 \rightarrow u_4$  with  $u_2 \neq u_3$ .

For  $i \in \{1, \dots, 6\}$ , every vertex of  $D_i$  has at most one predecessor in  $\mathcal{V}_1$ . Hence  $u_2$  and  $u_3$  can not both belong to  $\mathcal{V}_1$ . On the other hand, since each vertex has at most one successor in  $\mathcal{V}_2$ ,  $u_2$  and  $u_3$  can not both belong to  $\mathcal{V}_2$ . Without loss of generality, we suppose  $u_2 \in \mathcal{V}_1$  and  $u_3 \in \mathcal{V}_2$ . We have the following 4 cases.

*Case 1.*  $u_1, u_4 \in \mathcal{V}_1$ . In  $D_1, D_3, D_5$ , each 2-walk in  $D(\mathcal{V}_1)$  originates at a vertex who has no successors in  $\mathcal{V}_2$ . In  $D_4$ , there exists no 2-walks in  $D(\mathcal{V}_1)$ . In  $D_2, D_6$ , among the vertices in  $\mathcal{V}_1$ , only  $x$  has successors in both  $\mathcal{V}_1$  and  $\mathcal{V}_2$ . The only 2-walk in  $D(\mathcal{V}_1)$  with initial vertex  $x$  is  $x \rightarrow y \rightarrow x$ . But the unique successor of  $x$  in  $\mathcal{V}_2$ , namely  $w$ , is not a predecessor of  $x$ .

*Case 2.*  $u_1 \in \mathcal{V}_1$  and  $u_4 \in \mathcal{V}_2$ . In  $D_1, D_3$  and  $D_5$ ,  $D(\mathcal{V}_2)$  contains no arcs, and hence there exists no  $u_3 \rightarrow u_4$ . In  $D_2$  and  $D_4$ , among the vertices of  $\mathcal{V}_1$  only  $x$  has successors in both  $\mathcal{V}_1$  and  $\mathcal{V}_2$ . We know  $x$  has a unique successor in  $\mathcal{V}_2$ , namely  $w$ , but  $w$  has no successor in  $\mathcal{V}_2$ . Then there exists no 2-walks from  $x$  via  $\mathcal{V}_2$  to  $\mathcal{V}_2$ . In  $D_6$ , among the vertices of  $\mathcal{V}_1$  only  $x$  has successors in both  $\mathcal{V}_1$  and  $\mathcal{V}_2$ . We know  $x$  has a unique successor in  $\mathcal{V}_2$ , say  $w$ , and  $w$  has a unique successor  $z \in \mathcal{V}_2$ . But there exists no 2-walk from  $x$  to  $z$  via  $\mathcal{V}_1$ .

*Case 3.*  $u_1 \in \mathcal{V}_2$  and  $u_4 \in \mathcal{V}_1$ . In  $D_1, D_3$  and  $D_5$ ,  $D(\mathcal{V}_2)$  contains no arcs. Then there exists no  $u_1 \rightarrow u_3$ . In  $D_2, D_4$  and  $D_6$ , if there exists  $u_1$  has a successor  $u_3 \in \mathcal{V}_2$ , then we have  $N_{\mathcal{V}_1}^+(u_1) \rightarrow N_{\mathcal{V}_1}^+(u_3)$ .

*Case 4.*  $u_1, u_4 \in \mathcal{V}_2$ . In  $D_1, D_3, D_5, D(\mathcal{V}_2)$  contains no 2-walks, then  $D$  contains no  $u_1 \rightarrow u_3 \rightarrow u_4$ . In  $D_2$  and  $D_4$ , all 2-walks in  $D(\mathcal{V}_2)$  end at  $w$ . For all  $u \in \mathcal{V}_2$  with a successor in  $\mathcal{V}_2$ , we have  $N_{\mathcal{V}_1}^+(u) = \mathcal{V}_1 \setminus \{x\}$ . Combining with  $\mathcal{V}_1 \setminus \{x\} \rightarrow w$ , there exists no  $u_1 \rightarrow u_2 \rightarrow u_4$ . In  $D_6$ , all 2-walks in  $D(\mathcal{V}_2)$  end at  $w$  or  $z$ . For all  $u \in \mathcal{V}_2$  with successors in  $\mathcal{V}_2$ , we have  $N_{\mathcal{V}_1}^+(u) = \mathcal{V}_1 \setminus \{x\}$ . Combining with  $\mathcal{V}_1 \setminus \{x\} \rightarrow \{w, z\}$ , there exists no  $u_1 \rightarrow u_2 \rightarrow u_4$ .

In the above cases, none of  $D_1, \dots, D_6$  contains the required 2-walks  $u_1 \rightarrow u_2 \rightarrow u_4$  and  $u_1 \rightarrow u_3 \rightarrow u_4$ , a contradiction. Therefore,  $D_1, \dots, D_6$  are  $\mathcal{F}$ -free. By direct computation, we obtain  $e(D_1) = \frac{n^2+4n-4}{4}$  and  $e(D_i) = \frac{n^2+4n-5}{4}$  for  $i = 2, \dots, 6$ . Hence, we obtain (3.1).  $\square$

Let  $D = (\mathcal{V}, \mathcal{A})$  be a digraph. For a fixed vertex  $v \in \mathcal{V}$ , denote by

$$\mathcal{V}_1(v) = N^+(v) \text{ and } \mathcal{V}_2(v) = \mathcal{V} \setminus \mathcal{V}_1(v),$$

$$\mathcal{V}_3(v) = \{u \in \mathcal{V}_2(v) \mid N^+(u) = \mathcal{V}_1(v)\} \text{ and } \mathcal{V}_4(v) = \mathcal{V}_2(v) \setminus \mathcal{V}_3(v).$$

The index  $v$  will be omitted if no confusion from the context.

The following lemma is obvious.

**Lemma 6.** *Let  $D = (\mathcal{V}, \mathcal{A})$  be an  $\mathcal{F}$ -free digraph. Then*

- (i) *two distinct successors of a vertex share no common successor;*
- (ii) *given any  $v \in \mathcal{V}$ ,  $e(\mathcal{V}_1(v), u) \leq 1$  for all  $u \in \mathcal{V}$ .*

**Lemma 7.** *Let  $D = (\mathcal{V}, \mathcal{A})$  be an  $\mathcal{F}$ -free digraph and  $v \in \mathcal{V}$ . If there exists a 2-walk*

$$t_1 \rightarrow t_2 \rightarrow t_3 \text{ in } D(\mathcal{V}_1),$$

and

$$N^+(u) = \mathcal{V}_1 \text{ for all } u \in \mathcal{V}_2,$$

then  $t_1$  has no successor in  $\mathcal{V}_2$ .

*Proof.* Suppose  $t_1$  has a successor  $t_4 \in \mathcal{V}_2$ . Since  $N^+(t_4) = \mathcal{V}_1$ , we have  $t_4 \rightarrow t_3$ . Hence we obtain two distinct 2-walks from  $t_1$  to  $t_3$ , a contradiction with  $D$  being  $\mathcal{F}$ -free.  $\square$

Let  $\Delta^+(D)$  and  $\Delta^-(D)$  denote the maximum outdegree and indegree of  $D$ . If no confusion arises, we write  $\Delta^+$  and  $\Delta^-$ , respectively.

**Lemma 8.** *Let  $D = (\mathcal{V}, \mathcal{A}) \in EX(n)$ . If  $n$  is odd, then  $\Delta^+ \in \{\frac{n-1}{2}, \frac{n+1}{2}\}$ ; if  $n$  is even, then  $\Delta^+ \in \{\frac{n}{2} - 1, \frac{n}{2}, \frac{n}{2} + 1\}$ .*

*Proof.* Let  $v \in \mathcal{V}$  such that  $d^+(v) = \Delta^+$ . We count the size of  $D$  in the following way

$$e(D) = e(\mathcal{V}_1, \mathcal{V}) + e(\mathcal{V}_2, \mathcal{V}) \leq n + (n - \Delta^+)\Delta^+ = -(\Delta^+ - \frac{n}{2})^2 + \frac{n^2 + 4n}{4}.$$

The inequality is derived from Lemma 6 and the definition of  $\Delta^+$ . From (3.1), we get the lemma.  $\square$

Let

$$\alpha = \max_{\substack{v \in \mathcal{V}, \\ d^+(v) = \Delta^+}} \max_{u \in \mathcal{V}} e(u, \mathcal{V}_2(v)).$$

We give an upper bound on  $\alpha$  as follows.

**Lemma 9.** *Let  $D = (\mathcal{V}, \mathcal{A}) \in EX(n)$  and  $n \geq 8$ . Then  $\alpha \leq 1$ .*

*Proof.* Suppose  $\alpha \geq 3$ . Then there exist  $v$  and  $u$  such that  $d^+(v) = \Delta^+$  and  $u$  has 3 successors  $u_1, u_2, u_3 \in \mathcal{V}_2(v)$ . By Lemma 6, we have  $\sum_{i=1}^3 d^+(u_i) \leq n$  and

$$\begin{aligned} e(D) &= \sum_{w \in \mathcal{V}_2 \setminus \{u_1, u_2, u_3\}} d^+(w) + \sum_{i=1}^3 d^+(u_i) + \sum_{w \in \mathcal{V}} e(\mathcal{V}_1, w) \\ &\leq (n - \Delta^+ - 3)\Delta^+ + n + n \\ &= -(\Delta^+ - \frac{n-3}{2})^2 + \frac{n^2 + 2n + 9}{4}. \end{aligned}$$

From (3.1), we obtain  $e(D) < ex(n)$ , a contradiction. Hence, we have  $\alpha \leq 2$ .

Now suppose  $\alpha = 2$ . Then there exist  $v$  and  $u$  such that  $d^+(v) = \Delta^+$  and  $u$  has two successors  $u_1, u_2 \in \mathcal{V}_2(v)$ . We claim that one of  $u_1, u_2$  has  $\Delta^+$  successors. Otherwise, we have  $d^+(u_1) + d^+(u_2) \leq 2\Delta^+ - 2$ . Then

$$\begin{aligned} e(D) &= \sum_{w \in \mathcal{V}_2 \setminus \{u_1, u_2\}} d^+(w) + \sum_{i=1}^2 d^+(u_i) + \sum_{w \in \mathcal{V}} e(\mathcal{V}_1, w) \\ &\leq (n - \Delta^+ - 2)\Delta^+ + 2\Delta^+ - 2 + n \\ &= (n - \Delta^+)\Delta^+ - 2 + n \\ &\leq \frac{n^2 + 4n}{4} - 2. \end{aligned}$$

From (3.1) we obtain  $e(D) < ex(n)$ , a contradiction. Hence we could assume  $d^+(u_1) = \Delta^+$ .

By Lemma 6,  $u_2$  shares no common successor with  $u_1$ , i.e.,

$$e(u_2, \mathcal{V}_1(u_1)) = 0.$$



By the definition of  $\alpha$ , we have

$$e(u_2, \mathcal{V}_2(u_1)) \leq \alpha.$$

It follows that  $d^+(u_2) \leq \alpha = 2$ . Hence, by (3.1) we obtain

$$\begin{aligned} e(D) &= \sum_{w \in \mathcal{V}_2 \setminus \{u_1, u_2\}} d^+(w) + \sum_{i=1}^2 d^+(u_i) + \sum_{w \in \mathcal{V}} e(\mathcal{V}_1, w) \\ &\leq (n - \Delta^+ - 2)\Delta^+ + \Delta^+ + 2 + n \\ &= -(\Delta^+ - \frac{n-1}{2})^2 + \frac{n^2 + 2n + 9}{4} \\ &< ex(n). \end{aligned}$$

Therefore, we have  $\alpha \leq 1$ . □

Let  $D = (\mathcal{V}, \mathcal{A})$  be a digraph. Let  $v \in \mathcal{V}$  with  $d^+(v) = \Delta^+$  and  $u \in \mathcal{V}_2$ . If  $e(u, \mathcal{V}_1) = \Delta^+ - 1$ , we denote the unique vertex of  $\mathcal{V}_1 \setminus N_{\mathcal{V}_1}^+(u)$  by  $u'$ .

**Lemma 10.** *Let  $D = (\mathcal{V}, \mathcal{A}) \in EX(n)$  with  $n \geq 8$  and let  $v \in \mathcal{V}$  with  $d^+(v) = \Delta^+$ . If  $u_1 \rightarrow u_2$  is in  $D(\mathcal{V}_2(v))$ , then*

$$N_{\mathcal{V}_1(v)}^+(u_1) \not\rightarrow N_{\mathcal{V}_1(v)}^+(u_2).$$

Moreover, if  $\mathcal{V}_1(v) \rightarrow \mathcal{V}_1(v)$  and  $d^+(u_1) = d^+(u_2) = \Delta^+$ , then

$$u'_1 \rightarrow N_{\mathcal{V}_1(v)}^+(u_2), \quad u_2 \in \mathcal{V}_4(v) \text{ and } N_{\mathcal{V}_1(v)}^+(u_2) = \mathcal{V}_1(v) \setminus \{u'_1\}.$$

*Proof.* Suppose  $N_{\mathcal{V}_1(v)}^+(u_1) \rightarrow N_{\mathcal{V}_1(v)}^+(u_2)$ . Then there exists  $u_3 \in N_{\mathcal{V}_1(v)}^+(u_1)$  and  $u_4 \in N_{\mathcal{V}_1(v)}^+(u_2)$  such that  $u_3 \rightarrow u_4$ . We have  $u_1 \rightarrow u_3 \rightarrow u_4$  and  $u_1 \rightarrow u_2 \rightarrow u_4$ , a contradiction.

For the second part, since  $u_1 \rightarrow u_2$  is in  $D(\mathcal{V}_2(v))$ , by Lemma 9 we have  $e(u_1, \mathcal{V}_2(v)) = 1$ . It follows that

$$e(u_1, \mathcal{V}_1(v)) = \Delta^+ - 1 \text{ and } N_{\mathcal{V}_1(v)}^+(u_1) = \mathcal{V}_1 \setminus \{u'_1\}.$$

Since  $N_{\mathcal{V}_1(v)}^+(u_1) \not\rightarrow N_{\mathcal{V}_1(v)}^+(u_2)$  and  $\mathcal{V}_1(v) \rightarrow \mathcal{V}_1(v)$ , we have  $u'_1 \rightarrow N_{\mathcal{V}_1(v)}^+(u_2)$ . It follows that  $u'_1 \notin N^+(u_2)$  and  $u_2 \in \mathcal{V}_4(v)$ . By Lemma 9, we have  $e(u_2, \mathcal{V}_2(v)) \leq 1$  and  $e(u_2, \mathcal{V}_1(v)) \geq \Delta^+ - 1$ . It follows that  $N_{\mathcal{V}_1(v)}^+(u_2) = \mathcal{V}_1 \setminus \{u'_1\}$ . □

**Lemma 11.** *Let  $D = (\mathcal{V}, \mathcal{A}) \in EX(n)$  with  $n \geq 8$  and let  $v \in \mathcal{V}$  with  $d^+(v) = \Delta^+$ . If  $\mathcal{V}_1 \rightarrow \mathcal{V}$  and*

$$d^+(u) = \Delta^+ \text{ for all } u \in \mathcal{V}_2,$$

then

$$N^+(u) = \mathcal{V}_1 \text{ for all } u \in \mathcal{V}_2.$$

*Proof.* Suppose there exists an arc  $(u_1, u_2)$  in  $D(\mathcal{V}_2)$ . By Lemma 10, we have  $u'_1 \rightarrow \mathcal{V}_1 \setminus \{u'_1\}$  and  $u_2 \in \mathcal{V}_4$ . Hence  $u_2$  has a successor  $u_3 \in \mathcal{V}_2$ . Since  $u_1 \rightarrow u_2 \rightarrow u_3$  and  $u_1 \rightarrow \mathcal{V}_1 \setminus \{u'_1\}$ , we get  $\mathcal{V}_1 \setminus \{u'_1\} \rightarrow u_3$ . Since  $\mathcal{V}_1 \rightarrow \mathcal{V}$ , we obtain  $u'_1 \rightarrow u_3$ . By Lemma 10, we have  $u_3 \in \mathcal{V}_4$ . Then it has a successor  $u_4 \in \mathcal{V}_2$ , which implies  $u'_1 \rightarrow u_3 \rightarrow u_4$ . By Lemma 9,  $u'_1 \rightarrow u_4$ . From  $\mathcal{V}_1 \rightarrow \mathcal{V}$  we have  $u'_1 \rightarrow \mathcal{V}_1 \setminus \{u'_1\} \rightarrow u_4$ , which contradicts  $D \in EX(n)$ .  $\square$

**Lemma 12.** *Let  $D = (\mathcal{V}, \mathcal{A}) \in EX(n)$  with  $n \geq 8$ . If  $\Delta^+ \geq \Delta^-$ , then  $\Delta^+ = \lceil \frac{n+1}{2} \rceil$ .*

*Proof.* Let  $v \in \mathcal{V}$  such that  $d^+(v) = \Delta^+$ . Since  $\Delta^+ \geq \Delta^-$ , then

$$e(\mathcal{V}, \mathcal{V}_1) = \sum_{u \in \mathcal{V}_1} d^-(u) \leq \Delta^+ \Delta^- \leq (\Delta^+)^2. \quad (3.2)$$

Applying Lemma 9 we obtain

$$e(\mathcal{V}, \mathcal{V}_2) = \sum_{u \in \mathcal{V} \setminus \{v\}} e(u, \mathcal{V}_2) \leq n - 1. \quad (3.3)$$

It follows that

$$e(D) = e(\mathcal{V}, \mathcal{V}_1) + e(\mathcal{V}, \mathcal{V}_2) \leq (\Delta^+)^2 + n - 1. \quad (3.4)$$

From (3.1) we obtain  $e(D) < ex(n)$  when  $\Delta^+ < \frac{n}{2}$ . Hence,  $\Delta^+ \geq \frac{n}{2}$ . Combining with Lemma 8, we see that  $\Delta^+ = \frac{n+1}{2}$  when  $n$  is odd and  $\Delta^+ \in \{\frac{n}{2}, \frac{n}{2} + 1\}$  when  $n$  is even.

Now we consider the case  $n$  is even and  $\Delta^+ = \frac{n}{2}$ . By (3.1), (3.4) and  $D \in EX(n)$ , we have  $e(D) = \frac{n^2+4n-4}{4}$ . Combining with (3.2), (3.3) and (3.4), we obtain

$$d^-(u) = \Delta^+ \text{ for all } u \in \mathcal{V}_1. \quad (3.5)$$

and

$$e(\mathcal{V}, \mathcal{V}_1) = \frac{n^2}{4}, \quad e(\mathcal{V}, \mathcal{V}_2) = n - 1.$$

By Lemma 9, each vertex in  $\mathcal{V} \setminus \{v\}$  has a unique successor in  $\mathcal{V}_2$ .

By Lemma 6, we have  $e(\mathcal{V}_1, \mathcal{V}) \leq n$ . It follows from (3.1) that

$$e(\mathcal{V}_2, \mathcal{V}) = e(D) - e(\mathcal{V}_1, \mathcal{V}) \geq (n - \Delta^+) \Delta^+ - 1,$$

which implies there exists at least  $\Delta^+ - 1$  vertices in  $\mathcal{V}_2$  with outdegree  $\Delta^+$ .

Let  $u_1 \in \mathcal{V}_2 \setminus \{v\}$  such that  $d^+(u_1) = \Delta^+$  and  $u_1$  has a successor  $u_2 \in \mathcal{V}_2$ . We assert that either  $u_2 \rightarrow u'_1$  or  $u'_1$  has no predecessor in  $\mathcal{V}_1$ . Otherwise we have  $u_2 \rightarrow u'_1$  and there exists  $u_1^* \in \mathcal{V}_1 \setminus \{u'_1\}$  such that  $u_1^* \rightarrow u'_1$ . Then there exist two distinct 2-walks  $u_1 \rightarrow u_2 \rightarrow u'_1$  and  $u_1 \rightarrow u_1^* \rightarrow u'_1$ , a contradiction. By Lemma 6, we obtain  $e(\mathcal{V}_1, u'_1) \leq 1$ . It follows that

$$d^-(u'_1) = e(\mathcal{V}_2 \setminus \{u_1, u_2\}, u'_1) + e(\{u_1, u_2\}, u'_1) + e(\mathcal{V}_1, u'_1) \leq \Delta^+ - 1,$$

which contradicts (3.5). Hence, we have  $\Delta^+ = \frac{n}{2} + 1$ .

$\square$

## 4 Proof of Theorem 4

Now we are ready to present the proof of Theorem 4.

*Proof.* Let  $D = (\mathcal{V}, \mathcal{A}) \in EX(n)$ . Note that  $D \in EX(n)$  if and only if  $D' \in EX(n)$ . Without loss of generality, we assume  $\Delta^+ \geq \Delta^-$ . Let  $v \in \mathcal{V}$  such that  $d^+(v) = \Delta^+$ . Keep in mind that given  $u \in \mathcal{V}_2$  with outdegree  $\Delta^+$ ,  $u \in \mathcal{V}_4$  if and only if  $u$  has exactly one successor in  $\mathcal{V}_2$ .

By Lemma 6, we have

$$e(\mathcal{V}_1) \leq |\mathcal{V}_1| \text{ and } e(\mathcal{V}_1, \mathcal{V}_2) \leq |\mathcal{V}_2|.$$

It follows that

$$e(D) = e(\mathcal{V}_1) + e(\mathcal{V}_1, \mathcal{V}_2) + e(\mathcal{V}_2, \mathcal{V}) \leq |\mathcal{V}_1| + |\mathcal{V}_2| + |\mathcal{V}_2|\Delta^+ = n + (n - \Delta^+)\Delta^+. \quad (4.1)$$

We distinguish two cases according to the parity of  $n$ .

(1)  $n$  is even. By Lemma 12, we have  $\Delta^+ = \frac{n}{2} + 1$ . Then from (4.1) we obtain

$$e(D) \leq \frac{n^2 + 4n - 4}{4}.$$

It follows from (3.1) that

$$e(D) = ex(n) = \frac{n^2 + 4n - 4}{4}. \quad (4.2)$$

Now (4.1) and (4.2) lead to

$$d^+(u) = \Delta^+ \text{ for all } u \in \mathcal{V}_2$$

and

$$e(\mathcal{V}_1) = |\mathcal{V}_1| \text{ and } e(\mathcal{V}_1, \mathcal{V}_2) = |\mathcal{V}_2|,$$

which implies

$$\mathcal{V}_1 \rightarrow \mathcal{V}_1 \text{ and } \mathcal{V}_1 \rightarrow \mathcal{V}_2. \quad (4.3)$$

By Lemma 11, we have

$$N^+(u) = \mathcal{V}_1 \text{ for all } u \in \mathcal{V}_2. \quad (4.4)$$

Since  $\alpha \leq 1$ , there exist  $\frac{n}{2} - 1$  vertices in  $\mathcal{V}_1$  with exactly one successor in  $\mathcal{V}_2$ , leaving two vertices  $y_1$  and  $y_2$  in  $\mathcal{V}_1$  with no successors in  $\mathcal{V}_2$ . Moreover,

$$\mathcal{V}_1 \setminus \{y_1, y_2\} \text{ matches } \mathcal{V}_2. \quad (4.5)$$

If there exists  $z \in \mathcal{V}_1 \setminus \{y_1, y_2\}$  such that  $z \rightarrow y_2$ , we know  $z$  has a predecessor in  $\mathcal{V}_1$ . By Lemma 7, we have  $y_1 \rightarrow z$  or  $y_2 \rightarrow z$ . If the latter one holds, we obtain  $z \rightarrow y_2 \rightarrow z$ , a contradiction with Lemma 7. If the former one holds, we consider the predecessor of  $y_1$  in

$\mathcal{V}_1$ . By Lemma 7, we have  $y_2 \rightarrow y_1$ . Thus, we obtain  $z \rightarrow y_2 \rightarrow y_1$ , a contradiction with Lemma 7. Hence, we have  $y_1 \rightarrow y_2$ . Similarly,  $y_2 \rightarrow y_1$ .

By (4.3) every vertex  $u \in \mathcal{V}_1 \setminus \{y_1, y_2\}$  has a unique predecessor  $u_1 \in \mathcal{V}_1$ . By Lemma 7, either  $u \in N^+(y_1) \cup N^+(y_2)$  or  $u_1 \in N^+(y_1) \cup N^+(y_2)$ . By the first part of (4.3), we have  $D(\mathcal{V}_1) = T(y_1, y_2)$ . Combining with (4.4) and (4.5),  $D$  is an isomorphism of  $D_1$ .

(2) For  $n$  is odd. By Lemma 12, we have  $\Delta^+ = \frac{n+1}{2}$ . It follows from (4.1) that

$$ex(n) \leq \frac{n^2 + 4n - 1}{4}. \quad (4.6)$$

Suppose equality in (4.6) holds. From (4.1), we obtain  $d^+(u) = \Delta^+$  for all  $u \in \mathcal{V}_2$  and  $\mathcal{V}_1 \rightarrow \mathcal{V}$ . By Lemma 11, (4.4) holds. Since  $|\mathcal{V}_1| = |\mathcal{V}_2| + 1$  and  $\alpha \leq 1$ , there exists  $y \in \mathcal{V}_1$  with no successor in  $\mathcal{V}_2$  and

$$\mathcal{V}_1 \setminus \{y\} \text{ matches } \mathcal{V}_2. \quad (4.7)$$

Since  $\mathcal{V}_1 \rightarrow \mathcal{V}_1$ , there must exist a cycle whose length is larger than or equal to 2. It follows that there exists a walk  $t_1 \rightarrow t_2 \rightarrow t_3$  with  $t_1 \neq y$ . By (4.7)  $t_1$  has a successor  $t_4 \in \mathcal{V}_2$ , which contradicts Lemma 7.

Therefore,  $ex(n) \leq \frac{n^2+4n-5}{4}$ . From (3.1), we obtain

$$e(D) = ex(n) = \frac{n^2 + 4n - 5}{4}.$$

By Lemma 6 we have  $e(\mathcal{V}_1, \mathcal{V}) \leq n$ , which implies that  $e(\mathcal{V}_2, \mathcal{V}) \geq \frac{n^2-5}{4}$ . Hence, at least  $\frac{n-3}{2}$  vertices of  $\mathcal{V}_2$  have outdegrees  $\Delta^+$  and  $d^+(u) \geq \frac{n-1}{2}$  for all  $u \in \mathcal{V}_2$ . Since  $\alpha \leq 1$ , we have

$$e(u, \mathcal{V}_1) \geq \frac{n-3}{2} \text{ for all } u \in \mathcal{V}_2.$$

We will use the following claim repeatedly.

**Claim 1.** *Let  $u \in \mathcal{V}_1$  with  $e(u, \mathcal{V}_1) \geq 3$ . If it has a predecessor  $u^*$ , then  $u^*$  has no successors in  $\mathcal{V}_2$ .*

Otherwise,  $u^*$  has a successor  $u_1 \in \mathcal{V}_2$ . Combining with  $e(u_1, \mathcal{V}_1) \geq \frac{n-3}{2}$  and  $e(u, \mathcal{V}_1) \geq 3$ ,  $N_{\mathcal{V}_1}^+(u) \cap N_{\mathcal{V}_1}^+(u_1)$  is not empty, which contradicts Lemma 6.

As shown in (4.1), the size of  $D$  is the sum of three parts:  $e(\mathcal{V}_1)$ ,  $e(\mathcal{V}_1, \mathcal{V}_2)$ ,  $e(\mathcal{V}_2, \mathcal{V})$ . We have proven that the maximum size of  $D$  is less by 1 than the sum of the maximum numbers of these three parts. Hence, we can deduce that two of three parts achieve the maximum numbers and one of them misses one from the maximum number. We present these three maximum numbers as follows.

$$e(\mathcal{V}_1, \mathcal{V}_1) = \Delta^+, \quad (4.8)$$

$$e(\mathcal{V}_1, \mathcal{V}_2) = n - \Delta^+, \quad (4.9)$$

and

$$d^+(u) = \Delta^+ \text{ for all } u \in \mathcal{V}_2. \quad (4.10)$$

Moreover, (4.8) is equivalent to

$$\mathcal{V}_1 \rightarrow \mathcal{V}_1; \quad (4.11)$$

and (4.9) is equivalent to

$$\mathcal{V}_1 \rightarrow \mathcal{V}_2. \quad (4.12)$$

According to the above analysis, we distinguish three cases:

*Case 1.* (4.8) and (4.9) holds. Then  $\frac{n-3}{2}$  vertices of  $\mathcal{V}_2$  have outdegrees  $\frac{n+1}{2}$  and a vertex  $w \in \mathcal{V}_2$  has outdegree  $\frac{n-1}{2}$ . By Lemma 9, we have  $\alpha \leq 1$ . Then there exist  $|\mathcal{V}_2|$  vertices of  $\mathcal{V}_1$  having exactly one successor in  $\mathcal{V}_2$ , leaving only one vertex  $y \in \mathcal{V}_1$  having no successor in  $\mathcal{V}_2$ . It follows that

$$\mathcal{V}_1 \setminus \{y\} \text{ matches } \mathcal{V}_2. \quad (4.13)$$

Now we distinguish three cases.

*Subcase 1.1.*  $D(\mathcal{V}_2)$  contains no arc. Then

$$N^+(u) = \mathcal{V}_1 \text{ for all } u \in \mathcal{V}_2 \setminus \{w\} \text{ and } N^+(w) \subset \mathcal{V}_1. \quad (4.14)$$

We assert that if there exists  $t_1 \rightarrow t_2 \rightarrow t_3$  in  $D(\mathcal{V}_1)$ , then either  $t_1 = y$  or  $t_1 \rightarrow w$ . Otherwise, there exists  $t \in \mathcal{V}_2 \setminus \{w\}$  such that  $t_1 \rightarrow t$ . From (4.14), we obtain a 2-walk  $t_1 \rightarrow t \rightarrow t_3$ , a contradiction.

Since  $\mathcal{V}_1 \rightarrow \mathcal{V}_1$ ,  $D(\mathcal{V}_1)$  must contain a cycle. From the above assertion, we obtain  $D(\mathcal{V}_1)$  contains only one cycle  $z \leftrightarrow y$ , where  $z \in \mathcal{V}_1$  is the predecessor of  $w$ . Combining with  $z \rightarrow w \rightarrow \mathcal{V}_1 \setminus \{w'\}$ , we obtain  $z \notin \mathcal{V}_1 \setminus \{w'\}$ . Otherwise we have  $z \rightarrow w \rightarrow z$  and  $z \rightarrow y \rightarrow z$ , a contradiction. Thus we have  $z = w'$ , which implies  $w' \rightarrow w$ . Moreover,  $y$  has only one successor in  $\mathcal{V}_1$ , i.e.,

$$N_{\mathcal{V}_1}^+(y) = w' \quad (4.15)$$

We assert

$$N_{\mathcal{V}_1}^+(w') = \mathcal{V}_1 \setminus \{w'\}. \quad (4.16)$$

Otherwise, there exists  $t \in \mathcal{V}_1 \setminus \{w'\}$  with  $w' \rightarrow t$ . We know  $t$  has a predecessor  $u \in \mathcal{V}_1$ . Since 2-walks in  $D(\mathcal{V}_1)$  have to emanate from  $y$  or  $w'$ , from (4.15) we have  $w' \rightarrow u \rightarrow t$ . At the same time, we have  $w' \rightarrow w \rightarrow t$  since  $N^+(w) = \mathcal{V}_1 \setminus \{w'\}$ , a contradiction. Hence, we obtain  $w' \rightarrow \mathcal{V}_1 \setminus \{w'\}$ . It follows that

$$D(\mathcal{V}_1) = S_y(w'). \quad (4.17)$$

Since (4.13) and  $w' \rightarrow w$ , we have  $\mathcal{V}_1 \setminus \{y, w'\}$  matches  $\mathcal{V}_2 \setminus \{w\}$ . Combining with (4.14), (4.16) and (4.17),  $D$  is an isomorphism of  $D_2$  with  $T'(w)$  vanishing.

*Subcase 1.2.* There exists an arc in  $D(\mathcal{V}_2 \setminus \{w\})$ . Suppose  $u_1 \rightarrow u_2$  with  $u_1, u_2 \in \mathcal{V}_2 \setminus \{w\}$ . By Lemma 10, we obtain  $u'_1 = u'_2$  and  $u'_1 \rightarrow \mathcal{V}_1 \setminus \{u'_1\}$ . Since  $\mathcal{V}_1 \rightarrow \mathcal{V}_1$ ,  $u'_1$  has a predecessor  $t \in \mathcal{V}_1$ . By Claim 1 we have  $y \rightarrow u'_1$ . It follows that

$$D(\mathcal{V}_1) = S_y(u'_1).$$

Moreover,  $u'_1$  has a unique successor  $s \in \mathcal{V}_2$ .

If  $s \in \mathcal{V}_3$ , we have  $u'_1 \rightarrow s \rightarrow u'_1$ , a contradiction with  $u'_1 \rightarrow y \rightarrow u'_1$ . If  $s$  has a successor  $s_1 \in \mathcal{V}_2$ , we have  $u'_1 \rightarrow s \rightarrow s_1$ . On the other hand, from (4.13) we obtain  $u'_1 \rightarrow \mathcal{V}_1 \setminus \{u'_1\} \rightarrow s_1$ , a contradiction. Hence, we have

$$s = w \text{ and } N^+(w) \subset \mathcal{V}_1. \quad (4.18)$$

We assert that

$$u'_1 = w'.$$

Otherwise, we have  $w \rightarrow u'_1 \rightarrow \mathcal{V}_1 \setminus \{u'_1\}$ . On the other hand, since  $w \rightarrow \mathcal{V}_1 \setminus \{w'\}$ , then  $\mathcal{V}_1 \setminus \{w'\} \rightarrow \mathcal{V}_1 \setminus \{u'_1\}$ . Combining with (4.11), we obtain  $w' \rightarrow \mathcal{V}_1 \setminus \{u'_1\}$ . Recalling  $u'_1 \rightarrow \mathcal{V}_1 \setminus \{u'_1\}$ , we obtain a contradiction with Lemma 6. Then we have

$$D(\mathcal{V}_1) = S_y(w'). \quad (4.19)$$

We assert that

$$N_{\mathcal{V}_1}^+(u) = \mathcal{V}_1 \setminus \{w'\} \text{ for all } u \in \mathcal{V}_4. \quad (4.20)$$

Otherwise, there exists a vertex  $u \in \mathcal{V}_4 \setminus \{w\}$  such that  $u \rightarrow w'$ . By Claim 1, we have  $u$  has no successor in  $\mathcal{V}_2$ , which contradicts  $u \in \mathcal{V}_4 \setminus \{w\}$ .

We assert that if there exists a 2-walk  $u_3 \rightarrow u_4 \rightarrow u_5$  in  $D(\mathcal{V}_2)$ , then  $u_5 = w$ . Otherwise,  $u_5 \neq w$ . We have  $u_3 \neq w$  from (4.18). By (4.13) and  $u'_1 \rightarrow w$ , we obtain  $u_3 \rightarrow \mathcal{V}_1 \setminus \{u'_1\} \rightarrow u_5$ , a contradiction. Therefore, for any  $(t_1, t_2)$  in  $D(\mathcal{V}_2)$ , we have either  $t_2 = w$  or  $t_2 \in N^-(w)$ . It follows that  $D(\mathcal{V}_2) = T'(w) + aC_1$  with  $a \geq 1$ . Combining with (4.13), (4.18), (4.19) and (4.20),  $D$  is an isomorphism of  $D_2$ .

Notice that there exists  $(u_1, u_2)$  in  $D(\mathcal{V}_2 \setminus \{w\})$ . There must exist a 2-walk in  $D(\mathcal{V}_2)$ . Thus, in this case,  $T'(w)$  must contain a 2-walk.

*Subcase 1.3.*  $D(\mathcal{V}_2 \setminus \{w\})$  contains no arc but  $D(\mathcal{V}_2)$  contains arcs. We distinguish two cases.

*Subcase 1.3.1.*  $N^+(w) \subset \mathcal{V}_1$ . Then  $\mathcal{V}_4 \setminus \{w\}$  is not empty and

$$u \rightarrow w \text{ for all } u \in \mathcal{V}_4 \setminus \{w\},$$

which leads to

$$D(\mathcal{V}_2) = S'(w) + aC_1, \quad (4.21)$$

where  $a \geq 1$ . By Lemma 10, for any  $u \in \mathcal{V}_4 \setminus \{w\}$  we have  $\mathcal{V}_1 \setminus \{u'\} \rightarrow \mathcal{V}_1 \setminus \{w'\}$ . Since  $\mathcal{V}_1 \rightarrow \mathcal{V}_1$ , we have  $u' \rightarrow \mathcal{V}_1 \setminus \{w'\}$ , which implies that  $u' \notin \mathcal{V}_1 \setminus \{w'\}$ . It follows that

$$u' = w' \text{ for all } u \in \mathcal{V}_4. \quad (4.22)$$

Moreover,  $w' \rightarrow \mathcal{V}_1 \setminus \{w'\}$ . From (4.11)  $w'$  has a predecessor in  $\mathcal{V}_1$ . By Claim 1, we have  $y \rightarrow w'$ . Hence, we obtain

$$D(\mathcal{V}_1) = S_y(w'). \quad (4.23)$$

Let  $w_1$  be the successor of  $w'$  in  $\mathcal{V}_2$ . If  $w_1$  has a successor  $w_2 \in \mathcal{V}_2$ . Then by (4.13) we have  $w' \rightarrow w_1 \rightarrow w_2$  and  $w' \rightarrow \mathcal{V}_1 \setminus \{w'\} \rightarrow w_2$ , a contradiction. If  $w_1 \in \mathcal{V}_3$ , then  $w_1 \rightarrow w'$ . Hence, we obtain  $w' \rightarrow w_1 \rightarrow w'$  and  $w' \rightarrow y \rightarrow w'$ , a contradiction. Thus, we get  $w_1 = w$ , i.e.,  $w$  is the successor of  $w'$  in  $\mathcal{V}_2$ . It follows from (4.13) that

$$\mathcal{V}_1 \setminus \{y, w'\} \text{ matches } \mathcal{V}_2 \setminus \{w\} \text{ and } w' \rightarrow w. \quad (4.24)$$

Combining with (4.21), (4.22), (4.23) and (4.24),  $D$  is an isomorphism of  $D_2$ , where  $T'(w)$  is  $S'(w)$ . Note that we could consider  $S'(w)$  as a special case of  $T'(w)$ .

*Subcase 1.3.2.*  $w$  has a successor  $w_0 \in \mathcal{V}_2$ . Let  $\{w_1, w_2\} = \mathcal{V}_1 \setminus N_{\mathcal{V}_1}^+(w)$ . Suppose  $N^+(w_0) = \mathcal{V}_1$ . By Lemma 10, we have  $N_{\mathcal{V}_1}^+(w) \rightarrow \mathcal{V}_1$ . From (4.11), we have

$$\{w_1, w_2\} \rightarrow \mathcal{V}_1. \quad (4.25)$$

Since  $D$  is loopless, we get

$$w_1 \leftrightarrow w_2. \quad (4.26)$$

By (4.13) at least one of  $\{w_1, w_2\}$  has a successor in  $\mathcal{V}_2$ . Without loss of generality, we assume  $w_1$  has a successor  $w_3 \in \mathcal{V}_2$ . If  $N^+(w_3) = \mathcal{V}_1$ , we have  $w_1 \rightarrow w_2 \rightarrow w_1$  and  $w_1 \rightarrow w_3 \rightarrow w_1$ , a contradiction. Thus,  $w_3 \in \mathcal{V}_4$ .

If  $w_3 \neq w$ , since  $w_1 \rightarrow w_2 \rightarrow N_{\mathcal{V}_1}^+(w_2)$  and  $w_1 \rightarrow w_3 \rightarrow \mathcal{V}_1 \setminus \{w_3'\}$ , we have  $N_{\mathcal{V}_1}^+(w_2) = w_3'$ . From (4.25) and (4.26), we obtain

$$w_1 = w_3' \text{ and } D(\mathcal{V}_1) = S_{w_2}(w_1). \quad (4.27)$$

Since  $w_3 \in \mathcal{V}_4 \setminus \{w\}$ ,  $w_3$  has a successor  $w_4 \in \mathcal{V}_2$ . Hence, from (4.13) and (4.27) we have  $w_1 \rightarrow w_3 \rightarrow w_4$  and  $w_1 \rightarrow \mathcal{V}_1 \setminus \{w_1\} \rightarrow w_4$ , a contradiction. Therefore, we obtain  $w_3 = w$ , i.e.,

$$w_1 \rightarrow w. \quad (4.28)$$

Since  $w_1 \rightarrow w \rightarrow \mathcal{V}_1 \setminus \{w_1, w_2\}$  and  $w_1 \rightarrow w_2$ , we obtain  $w_2 \rightarrow \mathcal{V}_1 \setminus \{w_1, w_2\}$ . Combining with  $\{w_1, w_2\} \rightarrow \mathcal{V}_1$  and  $w_2 \rightarrow w_2$ , we have  $w_1 \rightarrow \mathcal{V}_1 \setminus \{w_1\}$ . It follows that

$$D(\mathcal{V}_1) = S_{w_2}(w_1). \quad (4.29)$$

By (4.13) and (4.28), we have  $\mathcal{V}_1 \setminus \{w_1\} \rightarrow w_0$ , which implies  $w_1 \rightarrow \mathcal{V}_1 \setminus \{w_1\} \rightarrow w_0$ . On the other hand, we have  $w_1 \rightarrow w \rightarrow w_0$ , a contradiction.

Hence,  $N^+(w_0) \neq \mathcal{V}_1$  and  $w_0$  has a successor in  $\mathcal{V}_2$ . If this vertex is not  $w$ , then  $D(\mathcal{V}_2 \setminus \{w\})$  contains an arc, which contradicts the given condition of Subcase 3.3. Hence, we have

$$w_0 \rightarrow w. \quad (4.30)$$

By Lemma 10, we have  $\mathcal{V}_1 \setminus \{w'_0\} \nrightarrow \mathcal{V}_1 \setminus \{w_1, w_2\}$ . Since  $\mathcal{V}_1 \rightarrow \mathcal{V}_1$ , we have  $w'_0 \rightarrow \mathcal{V}_1 \setminus \{w_1, w_2\}$ . Since  $D$  is loopless, we have  $w'_0 \in \{w_1, w_2\}$ . Without loss of generality, we suppose  $w'_0 = w_1$ . It follows that

$$w_0 \rightarrow \mathcal{V}_1 \setminus \{w_1\}. \quad (4.31)$$

and

$$w_1 \rightarrow \mathcal{V}_1 \setminus \{w_1, w_2\}. \quad (4.32)$$

By  $w_0 \rightarrow w \rightarrow w_0$  and (4.31), we have  $\mathcal{V}_1 \setminus \{w_1\} \nrightarrow w_0$ . Combining with (4.13), we obtain

$$w_1 \rightarrow w_0.$$

Similarly, since  $w \rightarrow w_0 \rightarrow w$  and  $w \rightarrow \mathcal{V}_1 \setminus \{w_1, w_2\}$ , we have  $\{w_1, w_2\} \rightarrow w$ . Hence,

$$w_2 \rightarrow w.$$

Applying Lemma 10 on  $w \rightarrow w_0$ , we have

$$\mathcal{V}_1 \setminus \{w_1, w_2\} \nrightarrow \mathcal{V}_1 \setminus \{w_1\}.$$

Hence, by (4.32) and (4.13), we have  $w_1 \rightarrow \mathcal{V}_1 \setminus \{w_1\}$ . Now we have  $w_1 \rightarrow w_2 \rightarrow w$  and  $w_1 \rightarrow w_0 \rightarrow w$ , a contradiction.

*Case 2.* (4.9) and (4.10) hold. Then

$$e(\mathcal{V}_1) = \Delta^+ - 1. \quad (4.33)$$

(4.33) implies that there exists a vertex  $x \in \mathcal{V}_1$ , which has no predecessor in  $\mathcal{V}_1$ , such that

$$\mathcal{V}_1 \rightarrow \mathcal{V}_1 \setminus \{x\} \quad (4.34)$$

Note that  $|\mathcal{V}_1| = |\mathcal{V}_2| + 1$ . By Lemma 9, there exists a unique vertex  $y \in \mathcal{V}_1$  such that  $e(y, \mathcal{V}_2) = 0$  and

$$\mathcal{V}_1 \setminus \{y\} \text{ matches } \mathcal{V}_2. \quad (4.35)$$

Now we distinguish three cases.

*Subcase 2.1.*  $\mathcal{V}_4$  is empty, which implies that

$$N^+(u) = \mathcal{V}_1 \text{ for all } u \in \mathcal{V}_2. \quad (4.36)$$

Suppose  $y \neq x$ . Then  $y$  has a predecessor in  $\mathcal{V}_1$ . By Lemma 7,  $y$  has no successor in  $\mathcal{V}_1$ . It follows that there exists no 2-walks in  $D(\mathcal{V}_1)$ . Recalling (4.34), among the vertices of  $\mathcal{V}_1$



only  $x$  has successors in  $\mathcal{V}_1$ , which implies  $D(\mathcal{V}_1) = S(x)$  where  $x \in \mathcal{V}_1 \setminus \{y\}$ . Moreover, by (4.35)  $x$  has a successor in  $\mathcal{V}_2$ . Combining with (4.35) and (4.36),  $D$  is an isomorphism of a special case of  $D_4$ , where  $\mathcal{V}_3 = \mathcal{V}_2$ .

Suppose  $y = x$ . By Lemma 7, all 2-walks originate at  $y$ . For any  $(u_1, u_2)$  in  $D(\mathcal{V}_1)$ , either  $u_1 = y$  or  $u_1 \in N^+(y)$ . From (4.34) we have  $D(\mathcal{V}_1) = T(y)$ . Combing with (4.35) and (4.36),  $D$  is an isomorphism of  $D_3$ .

*Subcase 2.2.* There exists an arc  $(u_1, u_2)$  in  $D(\mathcal{V}_2)$  with  $N^+(u_2) = \mathcal{V}_1$ . By Lemma 10, we have  $N_{\mathcal{V}_1}^+(u_1) \not\rightarrow \mathcal{V}_1$ . By Lemma 9, we have  $e(u_1, \mathcal{V}_2) = 1$  and  $e(u_1, \mathcal{V}_1) = \frac{n-1}{2}$ . Hence, we have  $N_{\mathcal{V}_1}^+(u_1) = \mathcal{V}_1 \setminus \{u'_1\}$ . By (4.34), we obtain  $u'_1 \rightarrow \mathcal{V}_1 \setminus \{x\}$ . Since  $D$  is loopless, we have  $u'_1 = x$  and

$$D(\mathcal{V}_1) = S(x). \quad (4.37)$$

By Lemma 9 and (4.10), we obtain

$$e(u, \mathcal{V}_1) \geq \Delta^+ - 1 \text{ for all } u \in \mathcal{V}_2.$$

We assert that

$$N_{\mathcal{V}_1}^+(u) = \mathcal{V}_1 \setminus \{x\} \text{ for all } u \in \mathcal{V}_4. \quad (4.38)$$

Otherwise, there exists an arc  $(t_1, t_2)$  in  $D(\mathcal{V}_2)$  with  $t_1 \rightarrow x$ . Then  $e(x, \mathcal{V}_1) = \frac{n-1}{2}$  and  $e(t_2, \mathcal{V}_1) \geq \frac{n-1}{2}$  contradict Lemma 6.

If  $x \neq y$ , then  $x$  has a successor  $x_1 \in \mathcal{V}_2$ . We assert that

$$N^+(x_1) = \mathcal{V}_1. \quad (4.39)$$

Otherwise,  $x_1$  has a successor  $x_2 \in \mathcal{V}_2$  and  $x \rightarrow x_1 \rightarrow x_2$ . On the other hand, by (4.35) and  $x \rightarrow x_1$  we have  $x \rightarrow \mathcal{V}_1 \setminus \{x\} \rightarrow x_2$ . We get two 2-walks from  $x$  to  $x_2$ , a contradiction.

We also assert that if there exists  $u_3 \rightarrow u_4 \rightarrow u_5$  in  $D(\mathcal{V}_2)$ , then  $u_5 = x_1$ . Otherwise, from (4.38) we have  $u_3 \rightarrow \mathcal{V}_1 \setminus \{x\}$ . By (4.35) and  $x \rightarrow x_1$ , we have  $u_3 \rightarrow \mathcal{V}_1 \setminus \{x\} \rightarrow u_5$ , a contradiction.

It follows that all 2-walks in  $D(\mathcal{V}_2)$  share the same terminal vertex  $x_1$ . Hence for any arc  $t_1 \rightarrow t_2$  in  $D(\mathcal{V}_2)$ , if  $N^+(t_2) \neq \mathcal{V}_1$ , then  $x_1$  is the successor of  $t_2$  in  $\mathcal{V}_2$ . Therefore,

$$D(\mathcal{V}_2) = T'(x_1) + \sum_{i=1}^k S'(u_i) + aC_1,$$

with  $\mathcal{V}_3 \neq \mathcal{V}_2$ . Combining with (4.35), (4.37) and (4.38),  $D$  is an isomorphism of  $D_4$  with  $\mathcal{V}_3 \neq \mathcal{V}_2$ .

Now suppose  $x = y$ . Then  $x$  has no successor in  $\mathcal{V}_2$ . Let us turn back to consider  $u_1 \rightarrow u_2$  with  $u_1, u_2 \in \mathcal{V}_2$ . From (4.38) we obtain  $N_{\mathcal{V}_1}^+(u_1) = \mathcal{V}_1 \setminus \{x\}$ . Then  $\mathcal{V}_2(u_1) = \{x\} \cup \mathcal{V}_2 \setminus \{u_2\}$ . Moreover, in  $\mathcal{V}_2(u_1)$  there exist  $\frac{n-3}{2}$  vertices with outdegree  $\Delta^+$  and one vertex with outdegree

$\Delta^+ - 1$ . Replace the role of  $v$  by  $u_1$  and apply the same arguments as in Case 1,  $D$  is an isomorphism of  $D_2$ .

*Subcase 2.3.*  $\mathcal{V}_4$  is a proper subset of  $\mathcal{V}_2$  and we have  $u_2 \in \mathcal{V}_4$  for every arc  $(u_1, u_2)$  in  $D(\mathcal{V}_2)$ . Suppose  $t_1 \rightarrow t_2$  with  $t_1, t_2 \in \mathcal{V}_2$  and  $e(t_2, \mathcal{V}_1) = \Delta^+ - 1$ . Then  $t_2$  has a successor  $t_3 \in \mathcal{V}_2$ . Similarly,  $t_3$  has a successor  $t_4 \in \mathcal{V}_2$ . Since  $t_1 \rightarrow t_2 \rightarrow t_3$  and  $t_1 \rightarrow \mathcal{V}_1 \setminus \{t'_1\}$ , then  $\mathcal{V}_1 \setminus \{t'_1\} \rightarrow t_3$ . Combining with (4.35), we have  $t'_1 \rightarrow t_3$ . Similarly,  $t'_2 \rightarrow t_4$ . From (4.35) we obtain  $t'_1 \neq t'_2$ . Since  $t_1 \rightarrow t_2 \rightarrow \mathcal{V}_1 \setminus \{t'_2\}$  and  $t_1 \rightarrow \mathcal{V}_1 \setminus \{t'_1\}$ , we have  $\mathcal{V}_1 \setminus \{t'_1\} \rightarrow \mathcal{V}_1 \setminus \{t'_2\}$ . From (4.34), we obtain  $t'_1 \rightarrow \mathcal{V}_1 \setminus \{x, t'_2\}$ , which implies  $e(t'_1, \mathcal{V}_1) \geq \Delta^+ - 2$ . Similarly,  $e(t'_2, \mathcal{V}_1) \geq \Delta^+ - 2$ . Hence there exists  $t \in \mathcal{V}_1$  such that  $t'_i \rightarrow t$  for  $i = 1, 2$ , which contradicts Lemma 6.

*Case 3.* (4.8) and (4.10) holds, and

$$e(\mathcal{V}_1, \mathcal{V}_2) = n - \Delta^+ - 1. \quad (4.40)$$

By Lemma 9, there exist  $|\mathcal{V}_2| - 1$  vertices of  $\mathcal{V}_1$  having exactly one successor in  $\mathcal{V}_2$ . Applying Lemma 6, there exists one vertex  $x \in \mathcal{V}_2$  with no predecessor in  $\mathcal{V}_1$ . Note that  $|\mathcal{V}_1| = |\mathcal{V}_2| + 1$ . There exist  $y_1, y_2 \in \mathcal{V}_1$  such that  $e(y_i, \mathcal{V}_2) = 0$  for  $i = 1, 2$  and

$$\mathcal{V}_1 \setminus \{y_1, y_2\} \text{ matches } \mathcal{V}_2 \setminus \{x\}. \quad (4.41)$$

Now we distinguish two cases.

*Subcase 3.1.*  $\mathcal{V}_4$  is empty. Then (4.36) holds. From (4.11), there exist cycles in  $D(\mathcal{V}_1)$ . Assume  $t_1 \in \mathcal{V}_1$  is in a cycle of  $D(\mathcal{V}_1)$ . Then there exists  $t_1 \rightarrow t_2 \rightarrow t_3$  in  $D(\mathcal{V}_1)$ . Lemma 7 guarantees that only  $y_1, y_2$  could be in a cycle, which implies there is only one cycle  $y_1 \leftrightarrow y_2$  in  $D(\mathcal{V}_1)$ . (4.11) implies that each vertex in  $\mathcal{V}_1$  has a unique predecessor in  $\mathcal{V}_1$ . By Lemma 7,  $\mathcal{V}_1 \subset N^+(y_1) \cup N^+(y_2) \cup N^+(N^+(y_1) \cup N^+(y_2))$ . Hence, we have  $D(\mathcal{V}_1) = T(y_1, y_2)$ . Combining with (4.36) and (4.41),  $D$  is an isomorphism of  $D_5$ .

*Subcase 3.2.*  $\mathcal{V}_4$  is a proper subset of  $\mathcal{V}_2$ . Then there exists an arc  $u_1 \rightarrow u_2$  in  $D(\mathcal{V}_2)$ . By Lemma 10, we have  $u'_1 \rightarrow \mathcal{V}_1 \setminus \{u'_1\}$ .

We assert that

$$u \rightarrow \mathcal{V}_1 \setminus \{u'_1\} \text{ for all } u \in \mathcal{V}_4. \quad (4.42)$$

Otherwise, there exists  $u \in \mathcal{V}_4$  such that  $u' \neq u'_1$ . Then  $u$  has a successor  $u^* \in \mathcal{V}_2$ . By Lemma 10, we obtain  $u' \rightarrow \mathcal{V}_1 \setminus \{u'\}$ , a contradiction with Lemma 6.

We assert if there exists  $t_1 \rightarrow t_2 \rightarrow t_3$  in  $D(\mathcal{V}_2)$ , then either  $u'_1 \rightarrow t_3$  or  $t_3 = x$ . Otherwise,  $t_3$  has a predecessor  $t_4 \in \mathcal{V}_1 \setminus \{u'_1\}$ . Since  $t_1 \in \mathcal{V}_4$  and (4.42), we have  $t_1 \rightarrow t_4 \rightarrow t_3$ , a contradiction.

Let us turn back to consider  $u_1 \rightarrow u_2$  with  $u_1, u_2 \in \mathcal{V}_2$ . Now by Lemma 10, we have  $u_2 \in \mathcal{V}_4$ , which means  $u_2$  has a successor in  $\mathcal{V}_2$ . Applying Lemma 10 repeatedly, we see that  $D(\mathcal{V}_2)$  must contain a cycle  $C_s$ . Since  $u'_1$  has a unique successor  $z \in \mathcal{V}_2$ , according to the

above assertion  $D(\mathcal{V}_2)$  contains only one 2-cycle  $z \leftrightarrow x$ . Moreover, for any arc  $(t_1, t_2)$  in  $D(\mathcal{V}_2)$ , either  $t_1 \in N^-(x) \cup N^-(z)$  or  $t_2 \in N^-(x) \cup N^-(z)$ . By Lemma 9, we have

$$D(\mathcal{V}_2) = T'(x, z) + aC_1, \quad (4.43)$$

where  $a \geq 1$ .

From (4.11),  $u'_1$  has a unique predecessor  $u_1^* \in \mathcal{V}_1$ . By Claim 1 we have  $u_1^* \in \{y_1, y_2\}$ . Without loss of generality, we let  $y_1 \rightarrow u_1^*$ . Note that  $x \in \mathcal{V}_4$  as  $x \rightarrow z$ . From (4.42) we get  $x' = u'_1$ . Hence,  $D(\mathcal{V}_1) = S_{y_1}(x')$ . Combining with (4.41), (4.42) and (4.43),  $D$  is an isomorphism of  $D_6$ .

So far we have proved that if  $e(D) = ex(n)$ , then  $D$  is an isomorphism of  $D_i$  with  $i \in \{1, \dots, 6\}$ . On the other hand, by Lemma 5,  $D_i$  is  $\mathcal{F}$ -free for  $i \in \{1, \dots, 6\}$ . Hence, we get the second part of Theorem 4.

This completes the proof. □

## Acknowledgement

Partial of this work was done when Huang was visiting Georgia Institute of Technology and Lyu was visiting Auburn University with the financial support of China Scholarship Council. They thank China Scholarship Council, Georgia Tech and Auburn University for their support. The second author also thanks Professor Tin-Yau Tam for helpful discussions on matrix theory during his visit.

## References

- [1] B. Bollobás, Extremal graph theory, Handbook of combinatorics, Vol. 1, 2, 1231-1292, Elsevier Sci. B. V., Amsterdam, 1995.
- [2] W.G. Brown and F. Harary, Extremal digraphs, Combinatorial theory and its applications, I (Proc. Colloq., Balatonfüred, 1969), pp. 135-198. North-Holland, Amsterdam, 1970.
- [3] W. G. Brown, M. Simonovits, Extremal multigraph and digraph problems, Paul Erdős and his mathematics, II (Budapest, 1999), 157-203, Bolyai Soc. Math. Stud., 11, János Bolyai Math. Soc., Budapest, 2002.
- [4] Z. Huang, Z. Lyu, Extremal digraphs avoiding an orientation of  $C_4$ , arXiv:1802.03431.

- [5] Z. Huang, Z. Lyu, P. Qiao, Turán problems for digraphs avoiding different walks of a given length with the same endpoints, arXiv:1608.06170.
- [6] Z. Huang, X. Zhan, Digraphs that have at most one walk of a given length with the same endpoints, *Discrete Math.* 311 (2011) 70-79.
- [7] M. Simonovits, A method for solving extremal problems in graph theory, stability problems, *Theory of Graphs (Proc. Colloq., Tihany, 1966)*, pp. 279-319, Academic Press, New York, 1968.
- [8] P. Turán, Eine Extremalaufgabe aus der Graphentheorie. (Hungarian) *Mat. Fiz. Lapok* 48 (1941) 436-452.
- [9] P. Turán, On the theory of graphs, *Colloq. Math.* 3 (1954) 19-30.
- [10] H. Wu, On the 0-1 matrices whose squares are 0-1 matrices, *Linear Algebra Appl.* 432 (2010) 2909-2924.
- [11] X. Zhan, *Matrix theory*, Graduate Studies in Mathematics 147, American Mathematical Society, Providence, RI, 2013.