

Vertex operator algebras generated by Ising vectors of σ -type

Cuipo Jiang*

School of Mathematical Science, Shanghai Jiao Tong University
Shanghai, 200240, China
email: cpjiang@sjtu.edu.cn

Ching Hung Lam[†]

Institute of Mathematics, Academia Sinica, Taipei, Taiwan 10617
e-mail: chlam@math.sinica.edu.tw

Hiroshi Yamauchi[‡]

Department of Mathematics, Tokyo Woman's Christian University
2-6-1 Zempukuji, Suginami-ku, Tokyo 167-8585, Japan
e-mail: yamauchi@lab.twcu.ac.jp

2010 Mathematics Subject Classification: Primary 17B69; Secondary 20B25, 20D08.

Abstract

We prove the uniqueness of the simple vertex operator algebra of OZ-type generated by Ising vectors of σ -type. We also prove that the simplicity can be omitted if the Griess algebra is isomorphic to the Matsuo algebra associated with the root system of type A_n .

1 Introduction

An *Ising vector* is a Virasoro vector of a vertex operator algebra (VOA) generating a simple Virasoro sub VOA of central charge $1/2$. Vertex operator algebras generated by Ising vectors are interesting from the view point of (finite) group theory since each Ising

*Partially supported by China NSF grants 11771281, 11531004.

[†]Partially supported by MoST grant 104-2115-M-001-004-MY3 of Taiwan.

[‡]Partially supported by JSPS KAKENHI Grant Number JP16K05073.

vector defines an involution called Miyamoto involution acting on them (cf. [Mi96]). The group generated by Miyamoto involutions associated with Ising vectors of σ -type (see Section 3 for definition) forms a 3-transposition group. When the VOAs have compact real forms, the 3-transposition groups obtained in this manner are of symplectic type (cf. [CH95]) and are completely classified in [Ma05]. In [Ma05], a complete list of 3-transposition groups generated by Miyamoto involutions associated with Ising vectors of σ -type as well as examples of VOAs realizing those groups are presented. The purpose of the paper [Ma05] is mainly on the classification of 3-transposition groups but not of VOAs. The classification of VOAs realizing 3-transposition groups has not been established so far. In this paper, we will prove the uniqueness of simple VOA structure of a VOA generated by Ising vectors of σ -type when it is of OZ-type, i.e., it is of CFT-type and the weight one subspace is trivial. Our result complements Matsuo's classification in the sense that the examples of VOAs in [Ma05] are indeed unique under the conditions considered there.

Let us explain our results precisely. Let V be a VOA of OZ-type and E_V the set of Ising vectors of V of σ -type. For $e \in E_V$, one can define the Miyamoto involution σ_e acting on V . It is shown in [Mi96] that the group G_V generated by $\{\sigma_e \mid e \in E_V\}$ is a 3-transposition group. The linear span of E_V forms a subalgebra of the Griess algebra of V which has a description in terms of the 3-transposition group G_V . This algebra structure is called the Matsuo algebra $B_{1/2,1/2}(G_V)$ associated with G_V (cf. [HRS15, Ma05]). Suppose V is generated by E_V as a VOA. The Griess algebra of V coincides with a homomorphic image of the Matsuo algebra $B_{1/2,1/2}(G_V)$ and G_V is a center-free 3-transposition group in $\text{Aut}(V)$. We will prove that the bilinear form on V is uniquely determined by its Griess algebra structure in Proposition 3.7. Since V is assumed to be of OZ-type, it has a unique simple quotient given by the non-degenerate quotient with respect to the bilinear form on V . Since the radical of the bilinear form on V is uniquely determined by the Griess algebra, we will prove in Theorem 3.6 the uniqueness of the VOA structure of V if it is simple. When V is simple, the Griess algebra of V is isomorphic to the quotient of Matsuo algebra by the radical of the bilinear form and uniquely determined by G_V . So our result shows that V is uniquely determined by the 3-transposition group G_V if V is simple and G_V is realized by Miyamoto involutions associated with Ising vectors of σ -type. In the case G_V is isomorphic to the symmetric group \mathfrak{S}_n of degree n , the Matsuo algebra $B_{1/2,1/2}(\mathfrak{S}_n)$ is already non-degenerate and we will prove in Theorem 4.6 that V is uniquely determined without assuming its simplicity. The simple VOA with $G_V = \mathfrak{S}_n$ is well studied in [LS08, JL16] and our result gives a characterization of this VOA in terms of the Griess algebra.

As a by-product, we can weaken the assumption on the positivity in [Ma05]. By the uniqueness for the VOA corresponding to \mathfrak{S}_3 , we can prove in Proposition 3.13 that the

3-transposition group G_V generated by Ising vectors of σ -type is always of symplectic type without assuming that V has a compact real form. However, since the existence of Ising vectors already requires the positivity on local subalgebras, it seems that all VOAs satisfying our conditions have compact real forms and we conjecture that Matsuo's list in [Ma05] is complete without assuming the positivity (cf. Conjecture 3.16).

The organization of this article is as follows. In Section 2, we review Matsuo algebras associated with 3-transposition groups in a general setting. In Section 3, we will prove the uniqueness of simple VOA structure of a VOA under the assumptions that it is of OZ-type and generated by Ising vectors of σ -type. In Section 4, we consider a VOA of OZ-type which is generated by Ising vectors of σ -type and whose Griess algebra is isomorphic to the Matsuo algebra associated with the symmetric group \mathfrak{S}_{n+1} (or the Weyl group of type A_n). We will prove the uniqueness of the VOA structure of such a VOA without assuming the simplicity. The whole arguments in Section 4 is based on induction on n .

Acknowledgement. Part of this work has been done while the second and the third named authors were staying at Shanghai Jiao Tong University in June 2016 and in November 2017. They gratefully acknowledge the hospitality there. H.Y. thanks Atsushi Matsuo for valuable comments and helpful discussions.

Notation and terminology. In this paper, vertex operator algebras (VOAs) are defined over the complex number field \mathbb{C} unless otherwise stated. A VOA V is of *CFT-type* if it has the $L(0)$ -grading $V = \bigoplus_{n \geq 0} V_n$ such that $V_0 = \mathbb{C}\mathbb{1}$, and is of *One-Zero type* or simply of *OZ-type* if it is of CFT-type and $V_1 = 0$. In this case, V is equipped with a unique invariant bilinear form such that $(\mathbb{1}|\mathbb{1}) = 1$ (cf. [Li94]). In this paper, we only consider VOAs of OZ-type. A real form $V_{\mathbb{R}}$ of V is called *compact* if the associated bilinear form is positive definite. For a subset A of V , the subalgebra generated by A is denoted by $\langle A \rangle$. For $a \in V_n$, we define its weight by $\text{wt}(a) = n$. We write $Y(a, z) = \sum_{n \in \mathbb{Z}} a_{(n)} z^{-n-1}$ for $a \in V$ and define its *zero-mode* by $o(a) := a_{(\text{wt}(a)-1)}$ if a is homogeneous, and extend it linearly. The weight two subspace V_2 carries a structure of a commutative algebra defined by the product $o(a)b = a_{(1)}b$ for $a, b \in V_2$. This algebra is called the *Griess algebra* of V . A *Virasoro vector* is an element $e \in V_2$ such that $e_{(1)}e = 2e$. It is known (cf. [Mi96]) that $L^e(n) := e_{(n+1)}$, $n \in \mathbb{Z}$ generate a Virasoro algebra of central charge $c_e = 2(e|e)$ and hence the subalgebra $\langle e \rangle$ is isomorphic to a Virasoro VOA. We denote by $\text{Vir}(e)$ the Virasoro algebra generated by e . $\text{Vir}(e)$ has a standard triangular decomposition $\text{Vir}(e)_{\pm} = \bigoplus_{n > 0} \mathbb{C}L^e(\pm n)$ and $\text{Vir}(e)_0 = \mathbb{C}L^e(0) \oplus \mathbb{C}c_e$. The universal enveloping algebra of $\text{Vir}(e)$ is denoted by $U(\text{Vir}(e))$ which is naturally graded by \mathbb{Z} such that $U(\text{Vir}(e))[j] = \{x \in U(\text{Vir}(e)) \mid [L^e(0), x] = jx\}$ for $j \in \mathbb{Z}$. Two

Virasoro vectors a and b are said to be *orthogonal* if $a_{(n)}b = 0$ for $n \geq 0$. Since V is of OZ-type, this is equivalent to $a_{(2)}b = 0$ (cf. Theorem 5.1 of [FZ92]). We denote by $L(c, h)$ the irreducible highest weight module over the Virasoro algebra with the central charge c and the highest weight h . A *simple* $c = c_e$ Virasoro vector $e \in V$ is a Virasoro vector such that $\langle e \rangle \cong L(c_e, 0)$. A Virasoro vector ω is called the *conformal vector* of V if each graded subspace V_n agrees with $\text{Ker}_V(o(\omega) - n)$ and satisfies $\omega_{(0)}a = a_{(-2)}\mathbb{1}$ for all $a \in V$. The half of the conformal vector gives the unit of the Griess algebra and hence uniquely determined. We write $\omega_{(n+1)} = L(n)$ for $n \in \mathbb{Z}$. A *sub VOA* (W, e) of V is a pair of a subalgebra W of V together with a Virasoro vector e in W such that e is the conformal vector of W . We often omit e and simply denote it by W . A sub VOA W of V is called *full* if V and W share the same conformal vector. The commutant subalgebra of (W, e) in V is defined by $\text{Com}_V W := \text{Ker}_V e_{(0)}$ (cf. [FZ92]).

2 Matsuo algebra of a 3-transposition group

We recall the definition of 3-transposition groups.

Definition 2.1. A *3-transposition group* is a pair (G, I) of a group G and a set of involutions I of G satisfying the following conditions.

- (1) G is generated by I .
- (2) I is closed under the conjugation, i.e., if $a, b \in I$ then $a^b = aba \in I$.
- (3) For any a and $b \in I$, the order of ab is bounded by 3.

A 3-transposition group (G, I) is called *indecomposable* if I is a conjugacy class of G . An indecomposable (G, I) is called *non-trivial* if I is not a singleton, i.e., G is not cyclic.

Let (G, I) be a 3-transposition group. We define a graph structure on I as follows: for $a, b \in I$, $a \sim b$ if and only if a and b are non-commutative. If $a \sim b$ then the order of ab is three and we have $a^b = b^a \in I$. We set $a \circ b := a^b = b^a$ if $a \sim b$. It is clear that I is a connected graph if and only if I is a single conjugacy class of G .

Let α, β be non-zero complex numbers. Let $B_{\alpha, \beta}(G, I) = \bigoplus_{i \in I} \mathbb{C}x^i$ be the vector space spanned by a formal basis $\{x^i \mid i \in I\}$ indexed by the set of involutions. We define a bilinear product and a bilinear form on $B_{\alpha, \beta}(G, I)$ by

$$x^i x^j := \begin{cases} 2x^i & \text{if } i = j, \\ \frac{\alpha}{2}(x^i + x^j - x^{i \circ j}) & \text{if } i \sim j, \\ 0 & \text{otherwise,} \end{cases} \quad (x^i | x^j) := \begin{cases} \frac{\beta}{2} & \text{if } i = j, \\ \frac{\alpha\beta}{8} & i \sim j, \\ 0 & \text{otherwise.} \end{cases} \quad (2.1)$$

Then $B_{\alpha, \beta}(G, I)$ is a commutative non-associative algebra with a symmetric invariant bilinear form (cf. [Ma05]). This algebra is called the *Matsuo algebra* associated with a

3-transposition group (G, I) with the accessory parameters α and β (cf. [HRS15]¹). In the following, we will simply denote $B_{\alpha,\beta}(G, I)$ by $B_{\alpha,\beta}(G)$.

Remark 2.2. For a subfield $\mathbb{Q}(\alpha, \beta) \subset K \subset \mathbb{C}$ we can define a canonical K -form $B_{\alpha,\beta}(G)_K = \bigoplus_{i \in I} Kx^i$ of $B_{\alpha,\beta}(G)$. In particular, if α, β are real, we can define a canonical \mathbb{R} -form $B_{\alpha,\beta}(G)_{\mathbb{R}} = \bigoplus_{i \in I} \mathbb{R}x^i$.

Suppose I is not a single conjugacy class. Then there is a partition $I = I_1 \sqcup I_2$ such that both I_1 and I_2 are closed under the conjugation. Then $G_1 = \langle I_1 \rangle$ and $G_2 = \langle I_2 \rangle$ are 3-transposition subgroups of G . Since $[G_1, G_2] = 1$ and $G = G_1G_2$, G is a central product of G_1 and G_2 . In this case, $B_{\alpha,\beta}(G)$ is a direct sum $B_{\alpha,\beta}(G_1) \oplus B_{\alpha,\beta}(G_2)$ of two-sided ideals which is also an orthogonal sum with respect to the bilinear form. Therefore, the structure of $B_{\alpha,\beta}(G)$ is determined by its indecomposable components.

Definition 2.3. The radical of the bilinear form on $B_{\alpha,\beta}(G)$ forms an ideal. We call the quotient algebra of $B_{\alpha,\beta}(G)$ by the radical of the bilinear form the *non-degenerate quotient*.

It is obvious that G acts on $B_{\alpha,\beta}(G)$ by conjugation. Namely, we can define a group homomorphism

$$\rho : G \rightarrow \text{Aut } B_{\alpha,\beta}(G) \quad (2.2)$$

by, for $i \in I$, letting $\rho_i x^j = x^{i \circ j}$ if $i \sim j$ and $\rho_i x^j = x^j$ otherwise. It is shown in [Ma05] that ρ is injective if and only if G is non-trivial and center-free.

Suppose $\alpha \neq 2$. Then the adjoint action of x^i has three distinct eigenvalues 2, 0 and α . For $i \in I$, let

$$B_{\alpha,\beta}(G)[i; h] = \text{Ker}_{B_{\alpha,\beta}(G)}(\text{ad}(x^i) - h) \quad \text{and} \quad B_{\alpha,\beta}^{i\pm}(G) = \{v \in B_{\alpha,\beta}(G) \mid \rho_i v = \pm v\}.$$

Then $B_{\alpha,\beta}(G)[i; 2] = \mathbb{C}x^i$ and we have

$$B_{\alpha,\beta}^{i+}(G) = \mathbb{C}x^i \oplus B_{\alpha,\beta}(G)[i; 0], \quad B_{\alpha,\beta}^{i-}(G) = B_{\alpha,\beta}(G)[i; \alpha].$$

Namely, the action of $i \in I$ on $B_{\alpha,\beta}(G)$ can be described by the adjoint action of x^i .

Suppose G is indecomposable. Then the number $|\{j \in I \mid j \sim i\}|$ is independent of the choice of $i \in I$ if I is finite. We denote this number by k . One can verify that

$$\left(\sum_{i \in I} x^i \right) \cdot x^j = \left(\frac{k\alpha}{2} + 2 \right) x^j$$

¹Strictly speaking, a Matsuo algebra is associated with a partial triple system and we are considering Matsuo algebras associated with the Fischer spaces of 3-transposition groups.

so that if $k\alpha + 4$ is non-zero then

$$\omega := \frac{4}{k\alpha + 4} \sum_{i \in I} x^i \quad (2.3)$$

satisfies $\omega v = 2v$ for $v \in B_{\alpha, \beta}(G)$ and ω gives twice the unity of $B_{\alpha, \beta}(G)$. By the invariance, one has $(x^i | \omega) = (x^i | x^i)$ and $(\omega | \omega) = \frac{2\beta |I|}{k\alpha + 4}$.

Remark 2.4. In VOA theory, a Matsuo algebra corresponds to a Griess algebra generated by $c = \beta$ Virasoro vectors having two highest weights 0 and α with binary fusion rules (cf. [Ma05]). The vector ω is the conformal vector of such a VOA and $2(\omega | \omega)$ gives the central charge.

3 VOAs generated by Ising vectors of σ -type

Recall the unitary series of the Virasoro VOAs. Let

$$\begin{aligned} c_n &:= 1 - \frac{6}{(n+2)(n+3)}, \quad n = 1, 2, 3, \dots, \\ h_{r,s}^{(n)} &:= \frac{(r(n+3) - s(n+2))^2 - 1}{4(n+2)(n+3)}, \quad 1 \leq r \leq n+1, \quad 1 \leq s \leq n+2. \end{aligned} \quad (3.1)$$

It is shown in [W93] that $L(c_n, 0)$ is rational and $L(c_n, h_{r,s}^{(n)})$, $1 \leq s \leq r \leq n+1$, are all the irreducible $L(c_n, 0)$ -modules (see also [DMZ94]). The fusion rules among $L(c_n, 0)$ -modules are computed in [W93] and given by

$$L(c_n, h_{r_1, s_1}^{(n)}) \boxtimes L(c_n, h_{r_2, s_2}^{(n)}) = \sum_{\substack{1 \leq i \leq M \\ 1 \leq j \leq N}} L(c_n, h_{|r_1 - r_2| + 2i - 1, |s_1 - s_2| + 2j - 1}^{(n)}), \quad (3.2)$$

where $M = \min\{r_1, r_2, n+2-r_1, n+2-r_2\}$ and $N = \min\{s_1, s_2, n+3-s_1, n+3-s_2\}$.

Definition 3.1. A Virasoro vector e with central charge c is called *simple* if $\langle e \rangle \cong L(c, 0)$. A simple $c = 1/2$ Virasoro vector is called an *Ising vector*.

Suppose e is an Ising vector of a VOA V of OZ-type. An Ising vector e is said to be *of σ -type on V* if there exists no irreducible $\langle e \rangle$ -submodule of V isomorphic to $L(1/2, 1/16)$. In this case, we have an isotypical decomposition

$$V = V[0]_e \oplus V[1/2]_e \quad (3.3)$$

where $V[h]_e$ is the sum of all irreducible $\langle e \rangle$ -submodules isomorphic to $L(1/2, h)$, and the Griess algebra of V decomposes into a direct sum of eigenspaces

$$V_2 = \mathbb{C}e \oplus V_2[0]_e \oplus V_2[1/2]_e \quad (3.4)$$

with $V_2[h]_e = V_2 \cap V[h]_e = \text{Ker}_{V_2}(o(e) - h)$. By the fusion rules of $L(1/2, 0)$ -modules in (3.1) and based on the decomposition (3.3), we can define an automorphism by

$$\sigma_e := (-1)^{2o(e)} = \begin{cases} 1 & \text{on } V[0]_e, \\ -1 & \text{on } V[1/2]_e. \end{cases} \quad (3.5)$$

The involution σ_e is called a *Miyamoto involution* of σ -type or a σ -*involution* (cf. [Mi96]). By the definition, we have the following conjugation.

Proposition 3.2. *Let $e \in V$ be an Ising vector of σ -type and $g \in \text{Aut}(V)$. Then we have $\sigma_{ge} = g\sigma_e g^{-1}$.*

The local structures of subalgebras generated by two Ising vectors of σ -type are completely determined in [Mi96, Ma05].

Proposition 3.3 ([Mi96, Ma05]). *Let V be a VOA of OZ-type and let a and b be distinct Ising vectors of σ -type on V . Then the Griess subalgebra B generated by a and b is one of the following.*

- (i) $(a | b) = 0$, $a_{(1)}b = 0$ and $B = \mathbb{C}a + \mathbb{C}b$. In this case, σ_a and σ_b are commutative on V .
- (ii) $(a | b) = 2^{-5}$, $\sigma_a b = \sigma_b a$, $4a_{(1)}b = a + b - \sigma_a b$ and $B = \mathbb{C}a + \mathbb{C}b + \mathbb{C}\sigma_a b$. In this case $\sigma_a \sigma_b$ has order three on V .

Conditions 1. We consider a VOA V satisfying the following conditions.

- (1) V is of OZ-type.
- (2) E_V is the set of Ising vectors of V of σ -type.
- (3) V is generated by E_V .

We set $G_V = \langle \{\sigma_e | e \in E_V\} \rangle$ and $I_V = \{\sigma_e | e \in E_V\}$. It follows from Propositions 3.2 and 3.3 that (G_V, I_V) is a 3-transposition group. In the following, we always assume that each indecomposable component of G_V is non-trivial. Then it is shown in [Ma05] that G_V is center-free and the correspondence $E_V \ni e \mapsto \sigma_e \in I_V$ is bijective. Therefore, we can index the elements of E_V by I_V so that $E_V = \{x^\alpha | \alpha \in I_V\}$ and $\sigma_{x^\alpha} = \alpha$.

Remark 3.4. If (G, I) is decomposable then we have a non-trivial partition $I = I_1 \sqcup I_2$ such that $G_i = \langle I_i \rangle$ are non-trivial 3-transposition subgroups of G and $G \cong G_1 \times G_2$. The partition $I = I_1 \sqcup I_2$ induces a partition $E = E_1 \sqcup E_2$ with $E_i = \{e \in E | \sigma_e \in I_i\}$. Let V^i be the sub VOA generated by E_i , $i = 1, 2$. Then $V \cong V^1 \otimes V^2$ and $G_i < \text{Aut}(V^i)$ (cf. [Mi96, Ma05]). Therefore, the study of V satisfying Conditions 1 is reduced to the case when G is indecomposable. The 3-transposition groups arising in this manner is classified in [Ma05] under the assumption that V has a compact real form. They are known to be finite 3-transposition groups of symplectic type (cf. [CH95]).

The following proposition is proved in Lemma 3.5 of [JL16] when G is a symmetric group but the same proof works for other G since it only depends on the local structure described in Proposition 3.3.

Proposition 3.5 ([JL16]). *Let V be a VOA satisfying Conditions 1. Then V is linearly spanned by*

$$\{x_{(-n_1)}^{\alpha_1} \cdots x_{(-n_k)}^{\alpha_k} \mathbb{1} \mid k \geq 0, \alpha_i \in I_V, n_i \geq 0\}.$$

By the proposition above, the Griess algebra of V is spanned by E_V . It follows from Proposition 3.3 that we have a Griess algebra epimorphism

$$\pi_V : B_{1/2, 1/2}(G_V) \rightarrow V_2$$

such that $\pi(x^i) = e$ if $i = \sigma_e$ in I_V . If V is simple, then the bilinear form on V_2 is non-degenerate so that the kernel of π_V is exactly the radical of the bilinear form on $B_{1/2, 1/2}(G_V)$ (cf. [Ma05]). Namely, if V is simple, then the Griess algebra of V is isomorphic to the unique simple quotient of the Matsuo algebra $B_{1/2, 1/2}(G_V)$ associated with $G_V = \langle \{\sigma_e \mid e \in E_V\} \rangle$. We will show that the whole VOA structure of V is uniquely determined by the Matsuo algebra when V is simple.

Theorem 3.6. *Let V be a simple VOA satisfying Conditions 1. Then the VOA structure of V is uniquely determined by its Griess algebra.*

Set

$$A = \bigsqcup_{k \geq 0} (I_V \times \mathbb{Z}_{\geq 0})^k = \{((\alpha_1, n_1), \dots, (\alpha_k, n_k)) \mid k \geq 0, \alpha_i \in I_V, n_i \geq 0\}, \quad (3.6)$$

$$A_N = \{((\alpha_1, n_1), \dots, (\alpha_k, n_k)) \in A \mid n_1 + \cdots + n_k + k = N\}.$$

Then $A = \bigsqcup_{N \geq 0} A_N$. We also set $A_{\leq N} = \bigsqcup_{n \leq N} A_n$. We define a map $\phi_V : A \rightarrow V$ by

$$\phi_V((\alpha_1, n_1), \dots, (\alpha_k, n_k)) = x_{(-n_1)}^{\alpha_1} \cdots x_{(-n_k)}^{\alpha_k} \mathbb{1}. \quad (3.7)$$

We will show the following proposition.

Proposition 3.7. *Let V be a VOA satisfying Conditions 1. Then the inner product $(\phi_V(a) \mid \phi_V(b))$ for $a, b \in A$ is uniquely determined by the Griess algebra of V .*

Before starting the proof of Proposition 3.7, we explain that Theorem 3.6 follows from Proposition 3.7. Suppose the proposition is true. Let V^1 and V^2 be any VOAs satisfying Conditions 1, not necessarily simple, generated by Ising vectors of σ -type. Assume that the Griess algebras of V^1 and V^2 are isomorphic. Let E_i be the set of Ising vectors of

V^i of σ -type, $I_i = \{\sigma_e \mid e \in E_i\}$ and $G_i = \langle I_i \rangle$ as before. Without loss of the generality, we may assume that G_i are indecomposable and non-trivial. Since the Griess algebras of V^1 and V^2 are isomorphic, there exists a bijection between E_1 and E_2 which induces an isomorphism $\theta : G_1 \rightarrow G_2$ such that $\theta(I_1) = I_2$. We identify I_1 with I_2 via θ and consider the index set A associated with $I_1 = I_2$ in (3.6). Since V^i is of OZ-type, V^i has a unique invariant bilinear form such that $(\mathbb{1}|\mathbb{1}) = 1$. The radical J^i of the bilinear form on V^i is the unique maximal ideal of V^i and $\tilde{V}^i = V^i/J^i$ is the unique simple quotient of V^i . We define a linear map $f : \tilde{V}^1 \rightarrow \tilde{V}^2$ by

$$f(\phi_{\tilde{V}^1}(a)) = \phi_{\tilde{V}^2}(a) \quad \text{for } a \in A.$$

This is well-defined. For, suppose we have a linear relation $\sum_j c_j \phi_{\tilde{V}^1}(a_j) = 0$ in \tilde{V}^1 . Then $\sum_j c_j \phi_{V^1}(a_j) \in J^1$. Since Griess algebras of V^1 and V^2 are isomorphic, it follows from Proposition 3.7 that $\sum_j c_j \phi_{V^1}(a_j) \in J^1$ if and only if $\sum_j c_j \phi_{V^2}(a_j) \in J^2$. Therefore, we have the same linear relation $\sum_j c_j \phi_{\tilde{V}^2}(a_j) = 0$ in \tilde{V}^2 . Then f gives an isomorphism between simple VOAs \tilde{V}^1 and \tilde{V}^2 by Proposition 5.7.9 of [LL04].

Therefore, in order to prove Theorem 3.6, it suffices to show Proposition 3.7. Namely, we will show that the inner products between elements in the spanning set $\phi_V(A)$ are uniquely determined by the Griess algebra of V under Conditions 1 and there is no need to use the whole structure of V .

The following lemma follows from Kac determinant (cf. [KR87]).

Lemma 3.8. *Let $e \in V$ be an Ising vector and $v \in V$ a highest weight vector for $\text{Vir}(e)$. Then $e_{(0)}^2 v \in \mathbb{C}e_{(-1)}v$.*

The following lemma can be proved by using the fermionic construction of the SVOA $L(1/2, 0) \oplus L(1/2, 1/2)$. The proof will be given in Appendix.

Lemma 3.9. *Let $e \in V$ be an Ising vector. Let $b \in V_2$ and $y \in V_N$ be highest weight vectors for $\text{Vir}(e)$ such that $2e_{(1)}b = b$ and $e_{(1)}y = hy$, $h \in \{0, 1/2\}$. Then for any $k \in \mathbb{Z}$, there exist $P_{h,k,j} \in \text{U}(\text{Vir}(e)_-)[j]$ such that*

$$(2e_{(i)} - \delta_{i,1}(1 - 2h)) \left(b_{(k)}y + \sum_{j>0} P_{h,k,j} b_{(k+j)}y \right) = 0$$

for $i \geq 1$. Moreover, $P_{h,k,j}$ are determined only by h, k, j and $b_{(N+1)}y \in V_0$.

Let $e \in E_V$, $a \in A$ and $k \geq 1$. We say $e_{(k)}\phi_V(a)$ has a *universal expression* if there exists an expression $e_{(k)}\phi_V(a) = \sum_j c_j \phi_V(b_j)$ in $\text{Span } \phi_V(A)$ such that if W is a VOA satisfying Conditions 1 with an isomorphism $\theta : G_V \rightarrow G_W$, then we also have the same

expression $(\theta e)_{(k)}\phi_W(a) = \sum_j c_j \phi_W(b_j)$ in $\text{Span } \phi_W(A)$ where we have identified I_W with E_W and θe is the element in E_W corresponding to $\theta\sigma_e$ in I_W .

Proposition 3.7 follows from the next lemma.

Lemma 3.10. *Let V be a VOA satisfying Conditions 1. Let $e \in E_V$ and $a \in \phi(A_N)$. Then for any $m \geq 1$, $e_{(m)}\phi_V(a)$ has a universal expression.*

Proof: We prove the lemma by induction on N . The case $N \leq 2$ is clear. Suppose the lemma holds for $a \in A_{\leq N}$ with $N \geq 2$. Suppose $a \in A_{N+1}$. Then there exist $n \geq 0$, $b \in E_V$ and $y \in \phi(A_{N-n})$ such that $\phi_V(a) = b_{(-n)}y$. Then

$$e_{(m)}\phi_V(a) = e_{(m)}b_{(-n)}y = [e_{(m)}, b_{(-n)}]y + b_{(-n)}e_{(m)}y.$$

By induction, $e_{(m)}y$ has a universal expression. On the other hand,

$$[e_{(m)}, b_{(-n)}]y = (e_{(0)}b)_{(m-n)}y + m(e_{(1)}b)_{(m-n-1)}y + \binom{m}{3}(e_{(3)}b)_{(m-n-3)}y$$

and again by induction the latter two terms have universal expressions since $e_{(1)}b \in \text{Span } E_V$ and $e_{(3)}b = (e|b)\mathbb{1}$. Therefore, we need to show that $(e_{(0)}b)_{(m-n)}y$ has a universal expression.

Case 1: $n > 0$. In this case $(e_{(0)}b)_{(m-n)}y = e_{(0)}b_{(m-n)}y - b_{(m-n)}e_{(0)}y$ and both y and $e_{(0)}y$ are in $\text{Span } \phi_V(A_{\leq N})$ so that we can apply the induction and the claim follows.

Case 2: $n = 0$ and $m > 1$. In this case $y \in \phi_V(A_N)$ and

$$\begin{aligned} (e_{(0)}b)_{(m-n)}y &= (e_{(0)}b)_{(m)}y = [e_{(1)}, b_{(m-1)}]y - (e_{(1)}b)_{(m-1)}y \\ &= e_{(1)}b_{(m-1)}y - b_{(m-1)}e_{(1)}y - (e_{(1)}b)_{(m-1)}y. \end{aligned}$$

Since $m > 1$, we can apply the induction to $b_{(m-1)}y$, $e_{(1)}y$ and $(e_{(1)}b)_{(m-1)}y$, and all of them have universal expressions in $\text{Span } \phi_V(A_{\leq N})$, and so are $e_{(1)}b_{(m-1)}y$ and $b_{(m-1)}e_{(1)}y$ again by induction.

Case 3: $n = 0$ and $m = 1$. By linearity, we may assume that $b \in V_2$ and $y \in \text{Span } \phi_V(A_N)$. We can also assume that $e_{(1)}b = tb$ with $t = 0, 1/2$ or 2 . If $t = 2$ then $b \in \mathbb{C}e$ and the claim follows since $(e_{(0)}e)_{(1)} = -e_{(0)}$. If $t = 0$ then b is a highest weight vector of highest weight 0 so that $e_{(0)}b = 0$ and the claim follows. Therefore we assume that b is a highest weight vector for $\text{Vir}(e)$ with highest weight $1/2$. In this case, we consider $e_{(1)}b_{(0)}y$ instead of $(e_{(0)}b)_{(1)}y$. Since $\langle e \rangle \cong L(1/2, 0)$ is rational, $e_{(1)}$ acts on $\phi_V(A_{\leq N})$ semisimply and y is a sum of descendants $e_{(-s_1)} \cdots e_{(-s_k)}v_\lambda$, $s_1 \geq \cdots \geq s_k \geq 0$, $k \geq 0$, of highest weight vectors $v_\lambda \in \text{Span } \phi_V(A_{\leq N})$ for $\text{Vir}(e)$. By the inductive assumption, this expression is universal, and again by linearity we may assume that $y = e_{(-s_1)} \cdots e_{(-s_k)}v_\lambda$ with $s_1 \geq \cdots \geq s_k \geq 0$ and $k \geq 0$. Moreover, by Lemma 3.8, we may assume that

$s_{k-1} \geq 1$ when $s_k = 0$. In other words, $s_1 = 0$ implies $k = 1$. Since e is of σ -type, the highest weight of v_λ is either 0 or $1/2$. We further divide Case 3 into three subcases.

Case 3-1: $k = 0$. If $y = v_\lambda \in \text{Span } \phi_V(A_N)$ is a highest weight vector for $\text{Vir}(e)$. By the inductive assumption $b_{(N+1)}y \in V_0$ has a universal expression and uniquely determined by the Griess algebra of V . Then it follows from Lemma 3.9 and the inductive assumption that $e_{(1)}b_{(0)}y$ has a universal expression.

Case 3-2: $k > 0$ and $s_1 > 0$. If $k > 0$ then $y = e_{(-s)}w$ with $s = s_1 \geq 0$ and $w = e_{(-s_2)} \cdots e_{(-s_k)}v_\lambda \in \text{Span } \phi_V(A_{N-s-1})$. Suppose $s > 0$. We have

$$e_{(1)}b_{(0)}y = e_{(1)}b_{(0)}e_{(-s)}w = e_{(1)}[b_{(0)}, e_{(-s)}]w + e_{(1)}e_{(0)}b_{(0)}w.$$

We also have $e_{(1)}e_{(0)}b_{(0)}w = e_{(0)}e_{(1)}b_{(0)}w + [e_{(1)}, e_{(0)}]b_{(0)}w = e_{(0)}e_{(1)}b_{(0)}w + e_{(0)}b_{(0)}w$. Since $b_{(0)}w \in \text{Span } \phi_V(A_{N-s})$, $e_{(1)}b_{(0)}w$ has a universal expression by induction. Therefore $e_{(1)}e_{(0)}b_{(0)}w$ has a universal expression. On the other hand, we have

$$\begin{aligned} e_{(1)}[b_{(0)}, e_{(-s)}]w &= -e_{(1)}[e_{(-s)}, b_{(0)}]w \\ &= -e_{(1)} \left((e_{(0)}b)_{(-s)} - s (e_{(1)}b)_{(-s-1)} + \binom{-s}{3} (e_{(3)}b)_{(-s-3)} \right) w \\ &= -e_{(1)}e_{(0)}b_{(-s)}w + e_{(1)}b_{(-s)}e_{(0)}w + se_{(1)} (e_{(1)}b)_{(-s-1)} w - \binom{-s}{3} e_{(1)}(e|b)\mathbb{1}_{(-s-3)}w \\ &= -e_{(0)}b_{(-s)}w - e_{(0)}e_{(1)}b_{(-s)}w + e_{(1)}b_{(-s)}e_{(0)}w + se_{(1)} (e_{(1)}b)_{(-s-1)} w. \end{aligned}$$

By the inductive assumption, $e_{(1)}b_{(-s)}w$ has a universal expression, and by Case 1, the terms $e_{(1)}b_{(-s)}e_{(0)}w$ and $e_{(1)} (e_{(1)}b)_{(-s-1)} w$ also have universal expressions. Therefore, all the terms in the above have universal expressions and the claim follows.

Case 3-3: $k = 1$ and $s_1 = 0$. Now it remains to show that $e_{(1)}b_{(0)}y$ has a universal expression when $y = e_{(0)}v_\lambda$. Note that $v_\lambda \in \text{Span } \phi_V(A_{N-1})$. We have

$$\begin{aligned} e_{(1)}b_{(0)}e_{(0)}v_\lambda &= e_{(1)}e_{(0)}b_{(0)}v_\lambda + e_{(1)}[b_{(0)}, e_{(0)}]v_\lambda \\ &= e_{(0)}b_{(0)}v_\lambda + e_{(0)}e_{(1)}b_{(0)}v_\lambda - e_{(1)}[e_{(0)}, b_{(0)}]v_\lambda \\ &= e_{(0)}b_{(0)}v_\lambda + e_{(0)}e_{(1)}b_{(0)}v_\lambda - e_{(1)}(e_{(0)}b)_{(0)}v_\lambda. \end{aligned}$$

Since $e_{(0)}b_{(0)}v_\lambda \in \text{Span } \phi_V(A_{N+1})$ and $e_{(1)}b_{(0)}v_\lambda$ has a universal expression by the inductive assumption, it suffices to consider the term $e_{(1)}(e_{(0)}b)_{(0)}v_\lambda$. We have

$$\begin{aligned} e_{(1)}(e_{(0)}b)_{(0)}v_\lambda &= (e_{(0)}b)_{(0)}e_{(1)}v_\lambda + [e_{(1)}, (e_{(0)}b)_{(0)}]v_\lambda \\ &= (e_{(0)}b_{(0)} - b_{(0)}e_{(0)})e_{(1)}v_\lambda + (e_{(0)}{}^2b)_{(1)}v_\lambda + (e_{(1)}e_{(0)}b)_{(0)}v_\lambda \\ &= e_{(0)}b_{(0)}e_{(1)}v_\lambda - b_{(0)}e_{(0)}e_{(1)}v_\lambda + (e_{(0)}{}^2b)_{(1)}v_\lambda + \frac{3}{2}(e_{(0)}b)_{(0)}v_\lambda \\ &= e_{(0)}b_{(0)}e_{(1)}v_\lambda - b_{(0)}e_{(0)}e_{(1)}v_\lambda + (e_{(0)}{}^2b)_{(1)}v_\lambda + \frac{3}{2}(e_{(0)}b_{(0)} - b_{(0)}e_{(0)})v_\lambda. \end{aligned}$$

By induction, all the term above except $(e_{(0)}^2 b)_{(1)} v_\lambda$ have universal expressions. Since $3e_{(0)}^2 b = 4e_{(-1)} b$ (cf. Lemma 3.8), it is enough to rewrite the term $(e_{(-1)} b)_{(1)} v_\lambda$. By the iterate formula, we have

$$(e_{(-1)} b)_{(1)} v_\lambda = \sum_{i \geq 0} (e_{(-1-i)} b_{(1+i)} + b_{(-i)} e_{(i)}) v_\lambda$$

and by induction all the terms in the right hand side have universal expressions. This completes the proof of Lemma 3.10. \blacksquare

Remark 3.11. The proof of Lemma 3.10 is a suitable modification of that of Lemma 3.5 of [JL16] and gives an alternative proof of Proposition 3.5.

Now by the invariance $(x_{(-n)}^\alpha \phi_V(a) | \phi_V(b)) = (\phi_V(a) | x_{(n+2)}^\alpha \phi_V(b))$ for $x^\alpha \in E_V$ and $a, b \in A$, Proposition 3.7 immediately follows from Lemma 3.10, and thus we obtain Theorem 3.6 as in the previous argument.

Proposition 3.12. *Let V be a VOA satisfying Conditions 1 and let a and b be Ising vectors such that $(a | b) = 2^{-5}$. Then the subalgebra $\langle a, b \rangle$ is simple and isomorphic to*

$$L(1/2, 0) \otimes L(7/10, 0) \oplus L(1/2, 1/2) \otimes L(7/10, 3/2).$$

Proof: By Proposition 3.3 the Griess algebra of $\langle a, b \rangle$ is isomorphic to the Matsuo algebra associated with the symmetric group \mathfrak{S}_3 of degree three. Since $\langle a, b \rangle$ is of OZ-type, and the radical J of the invariant bilinear form on $\langle a, b \rangle$ is the unique maximal ideal, it suffices to show that $J = 0$. Since $\langle a, b \rangle$ has a conformal vector

$$\eta = \frac{4}{5}(a + b + \sigma_a b) \tag{3.8}$$

given by (2.3), it follows from Theorem 5.1 of [FZ92] that we have a graded decomposition $J = \bigoplus_{n \geq 0} J_n$ with $J_n = J \cap V_n$. Let h be the top weight of J , i.e., $J_h \neq 0$ and $J_n = 0$ for $n < h$. Since a and b are of σ -type, the zero-modes $o(a)$, $o(b)$ and $o(\sigma_a b)$ act on J_h semisimply with eigenvalues in $\{0, 1/2\}$. Therefore, we have

$$\mathrm{Tr}_{J_h} o(\eta) = \frac{4}{5} \mathrm{Tr}_{J_h} o(a + b + \sigma_a b) \leq \frac{4}{5} \cdot 3 \cdot \frac{1}{2} \dim J_h = \frac{6}{5} \dim J_h. \tag{3.9}$$

On the other hand, since η is the conformal vector of $\langle a, b \rangle$, the zero mode $o(\eta)$ acts by the top weight h on J_h . Therefore $\mathrm{Tr}_{J_h} o(\eta) = h \dim J_h$ and from (3.9) we obtain $h \leq 6/5$, showing $h = 1$. This contradicts the assumption that V is of OZ-type. Therefore $J = 0$ and $\langle a, b \rangle$ is simple. Then by Theorem 3.6 its simple structure is uniquely determined as in the assertion (cf. [LY00]). \blacksquare

Proposition 3.13 ([Ma05]). *Let V be a VOA satisfying Conditions 1. Let E_V be the set of Ising vectors of V of σ -type and set $G_V = \langle \sigma_e \mid e \in E_V \rangle$. Then G_V is a 3-transposition group of symplectic type.*

Proof: It is shown in Proposition 1 of [Ma05]² that G_V is of symplectic type provided that V has a compact real form containing E_V . The key idea in the proof of [Ma05] is to find a non-zero highest weight vector for $L(7/10, 0)$ with highest weight $7/10$ when G is not of symplectic type. In (loc. cit.), the compact real form is used only to show the existence of the subalgebra isomorphic to $L(7/10, 0)$, and the existence of the highest weight vector depends only on the structure of the Griess algebra and this part is independent of the compact real form. Now by Proposition 3.12, we can obtain $L(7/10, 0)$ without the assumption on the compact real form, and we obtain the same contradiction as in [Ma05] if G_V is not of symplectic type. ■

Corollary 3.14. *Let V be a simple VOA satisfying Conditions 1 and let $V_{\mathbb{R}}$ be the real VOA generated by the set E_V of Ising vectors of V of σ -type. If the non-degenerate quotient of the real Matsuo algebra $B_{1/2, 1/2}(G_V)_{\mathbb{R}}$ associated with $G_V = \langle \sigma_e \mid e \in E_V \rangle$ is positive definite, then $V_{\mathbb{R}}$ is a compact real form of V . In this case a non-trivial indecomposable component of the 3-transposition group G_V is isomorphic to one of the groups listed in Theorem 1 of [Ma05].*

Proof: By the assumption, the invariant bilinear form of the Griess algebra of V is positive definite so that E_V is a finite set since $(a \mid b) = 2^{-5}$ or 0 for distinct $a, b \in E_V$. It follows from Proposition 3.13 and $V = \langle E_V \rangle$ that G_V is a finite 3-transposition group of symplectic type. By the positivity of the Griess algebra, it follows from [Ma05] that a non-trivial indecomposable component of G_V is one of the groups listed in Theorem 1 of [Ma05]. Since all the examples of VOAs in (loc. cit.) have compact real forms generated by Ising vectors, it follows from the uniqueness of the VOA structure of V shown in Theorem 3.6 that $V_{\mathbb{R}}$ is a compact real form of V . ■

Remark 3.15. In Theorem 1 of [Ma05], there exist non-isomorphic 3-transposition groups but realized by the same VOA. For example, $O_{10}^+(2)$ and $O_8^+(2)$ are realized by $V_{\sqrt{2}E_8}^+$. This is because $O_8^+(2) < O_{10}^+(2)$ and $B_{1/2, 1/2}(O_8^+(2))$ is a subalgebra of $B_{1/2, 1/2}(O_{10}^+(2))$ but they have isomorphic non-degenerate quotients. In our argument, we always take E_V to be the set of *all* Ising vectors of V of σ -type, and for $V = V_{\sqrt{2}E_8}^+$ we obtain the maximal one $G_V = O_{10}^+(2)$.

We propose a conjecture on positivity of a simple VOA satisfying Conditions 1.

²Proposition 3.3.8 of arXiv:math/0311400.

Conjecture 3.16. Let V be a simple VOA satisfying Conditions 1. Then the bilinear form on the \mathbb{R} -span of E_V is positive definite, i.e., the non-degenerate quotient of the real Matsuo algebra $B_{1/2,1/2}(G_V)_{\mathbb{R}}$ associated with G_V is positive definite.

If this conjecture is true, it follows from Corollary 3.14 that the classification of VOA-realizable 3-transposition groups together with VOAs generated by Ising vectors of σ -type in [Ma05] is complete without the assumption on the compact real form of a VOA.

4 Simplicity of type A_n

Let $\Phi(A_n)$ be the root system of type A_n . We fix a system of simple roots $\alpha_1, \dots, \alpha_n$ of $\Phi(A_n)$ such that $(\alpha_i | \alpha_j) = -1$ if and only if $|i - j| = 1$, and denote by $\Phi(A_n)^+$ the set of positive roots. We also fix root subsystems $\Phi(A_i)$ for $1 \leq i \leq n$ to be the root systems generated by $\{\alpha_j | 1 \leq j \leq i\}$ and set $\Phi(A_i)^+ = \Phi(A_n)^+ \cap \Phi(A_i)$. Let r_α be the reflection associated with $\alpha \in \Phi(A_n)$. The Weyl group $W(A_i)$ of $\Phi(A_i)$ is a 3-transposition group isomorphic to the symmetric group \mathfrak{S}_{i+1} of degree $i + 1$ with the set of transpositions $\{r_\alpha | \alpha \in \Phi(A_i)^+\}$.

Let M_{A_n} be the commutant subalgebra of the diagonal subalgebra $L_{\widehat{\mathfrak{sl}}_2}(n+1, 0)$ in the tensor product $L_{\widehat{\mathfrak{sl}}_2}(1, 0)^{\otimes n+1}$ (cf. [LS08, JL16]). Since $L_{\widehat{\mathfrak{sl}}_2}(1, 0)$ has a compact real form, so does $L_{\widehat{\mathfrak{sl}}_2}(1, 0)^{\otimes n+1}$ and hence M_{A_n} is a simple VOA with a compact real form. It is known (cf. [DLMN98, Ma05, LSY07, JL16]) that M_{A_n} is a rational VOA satisfying Conditions 1 with $G_{M_{A_n}} = W(A_n) = \mathfrak{S}_{n+1}$ and the Griess algebra of M_{A_n} is isomorphic to the Matsuo algebra $B_{1/2,1/2}(\mathfrak{S}_{n+1})$. In particular, $B_{1/2,1/2}(\mathfrak{S}_{n+1})$ is non-degenerate. The set of Ising vectors of M_{A_n} is in one-to-one correspondence with the set of reflections $\{r_\alpha | \alpha \in \Phi(A_n)^+\}$ so that we can index the set of Ising vectors of M_{A_n} as $E_{M_{A_n}} = \{x^\alpha | \alpha \in \Phi(A_n)^+\}$. Note that $\sigma_{x^\alpha} = r_\alpha$ in $\text{Aut}(M_{A_n}) = W(A_n)$ if $n > 1$.

Let V be a VOA satisfying Conditions 1 such that the associated 3-transposition group $G_V = \langle \{\sigma_e | e \in E_V\} \rangle$ is isomorphic to $W(A_n) \cong \mathfrak{S}_{n+1}$. In this case the Griess algebra of V is isomorphic to the non-degenerate Matsuo algebra $B_{1/2,1/2}(\mathfrak{S}_{n+1})$ and the set of Ising vectors of V is given by $E_V = \{x^\alpha | \alpha \in \Phi(A_n)^+\}$. By Theorem 3.6, V has the unique simple quotient isomorphic to M_{A_n} . In this section we will prove that V itself is automatically simple, that is, any VOA satisfying Conditions 1 whose Griess algebra is isomorphic to $B_{1/2,1/2}(\mathfrak{S}_{n+1})$ is isomorphic to M_{A_n} as a VOA.

Set $V^{[i]} := \langle x^\alpha | \alpha \in \Phi(A_i)^+ \rangle$ for $1 \leq i \leq n$. Then we have a tower of sub VOAs

$$V^{[1]} \subset V^{[2]} \subset \dots \subset V^{[n]} = V \tag{4.1}$$

such that $V^{[i]}$ satisfies Conditions 1 with $G_{V^{[i]}} = W(A_i) = \mathfrak{S}_{i+1}$. We know $V^{[1]} \cong L(1/2, 1/2)$ is simple and it follows from Proposition 3.12 that $V^{[2]}$ is also simple. So we assume

$n > 2$ and by induction we may suppose that $V^{[i]} \cong M_{A_i}$ for $1 \leq i < n$. Let ω^i be the conformal vector of $V^{[i]}$ given by (2.3). We also set $\omega^0 = 0$ for convention. Then $\eta^i := \omega^i - \omega^{i-1}$ for $1 \leq i \leq n$ are mutually orthogonal Virasoro vectors with central charges c_i in (2.3) (cf. [DLMN98, Ma05, FZ92]). By the inductive assumption, η^i , $1 \leq i < n$, are simple and $V^{[i]}$ contains a full sub VOA $\langle \eta^1, \dots, \eta^i \rangle$ isomorphic to $L(c_1, 0) \otimes \dots \otimes L(c_i, 0)$ (cf. [LS08, JL16]).

Consider the Zhu algebra $A(V) = V/O(V)$ of V (cf. [Z96]). We denote by $[a]$ the residue class $a + O(V)$ of $a \in V$ in $A(V)$. It is shown in Theorem 3.6 of [JL16] that Proposition 3.5 implies the following.

Lemma 4.1 ([JL16]). *Let V be a VOA satisfying Conditions 1. Then the Zhu algebra of V is generated by $E_V + O(V)$.*

We say a V -module N is of σ -type³ if for each $x^\alpha \in E_V$, there is no $\langle x^\alpha \rangle$ -submodule of N isomorphic to $L(1/2, 1/16)$. Let \mathfrak{a} be the two-sided ideal of $A(V)$ generated by $[x^\alpha][2x^\alpha - \mathbb{1}]$ with $\alpha \in \Phi(A_n)^+$. If $N = \bigoplus_{k \geq 0} N(k)$ is an \mathbb{N} -gradable V -module of σ -type, then the top level $N(0)$ is an $A(V)$ -module on which $[x^\alpha]$ with $\alpha \in \Phi(A_n)^+$ acts semisimply with possible eigenvalues in $\{0, 1/2\}$ so that the ideal \mathfrak{a} acts trivially on it and $N(0)$ is a module over the quotient $A(V)/\mathfrak{a}$. Since M_{A_n} is the simple quotient of V by Theorem 3.6, $A(M_{A_n})$ is a quotient algebra of $A(V)$ and we can define $A(M_{A_n})/\mathfrak{a}$ as well.

Let $\{r_\alpha \mid \alpha \in \Phi(A_n)^+\}$ be the set of transpositions of $W(A_n) = \mathfrak{S}_{n+1}$ and let T_{n+1} be the quotient algebra of $\mathbb{C}[\mathfrak{S}_{n+1}]$ by the ideal \mathfrak{b} generated by

$$\{(r_\alpha + r_\beta + r_\alpha r_\beta r_\alpha)(r_\alpha + r_\beta + r_\alpha r_\beta r_\alpha - 3) \mid \alpha, \beta \in \Phi(A_n)^+, r_\alpha \sim r_\beta\}.$$

It is shown in [JL16] that $A(M_{A_n})/\mathfrak{a}$ is isomorphic to T_{n+1} .

Lemma 4.2 ([JL16]). *There exists an isomorphism $f : T_{n+1} = \mathbb{C}[\mathfrak{S}_{n+1}]/\mathfrak{b} \rightarrow A(V)/\mathfrak{a}$ such that $r_\alpha + \mathfrak{b} \mapsto [\mathbb{1} - 4x^\alpha] + \mathfrak{a}$ for $\alpha \in \Phi(A_n)^+$.*

Proof: By the defining relation of the ideal \mathfrak{a} , we have $[\mathbb{1} - 4x^\alpha]^2 = [\mathbb{1}]$ in $A(V)/\mathfrak{a}$. Let $x^\alpha, x^\beta \in E_V$ be Ising vectors such that $r_\alpha \sim r_\beta$. Then $\langle x^\alpha, x^\beta \rangle$ is isomorphic to M_{A_2} by Proposition 3.12. We denote the Ising vector corresponding to $r_\alpha \circ r_\beta = r_\alpha r_\beta r_\alpha \in W(A_n)$ by $x^{\alpha\beta}$. It is known (cf. [LY00]) that M_{A_2} has two non-isomorphic irreducible modules of σ -type with top weights 0 and $3/5$. Since the conformal vector of $\langle x^\alpha, x^\beta \rangle$ is given by $(4/5)(x^\alpha + x^\beta + x^{\alpha\beta})$ (cf. Eq. (2.3)), it follows that

$$[x^\alpha + x^\beta + x^{\alpha\beta}][4(x^\alpha + x^\beta + x^{\alpha\beta}) - 3 \cdot \mathbb{1}] = 0$$

³Such a module is called of type I in [JL16].

in $A(V)/\mathfrak{a}$. It is shown in [JL16] that

$$[\mathbb{1} - 4x^\alpha][\mathbb{1} - 4x^\beta][\mathbb{1} - 4x^\alpha] - [\mathbb{1} - 4x^{\alpha\circ\beta}] = 0$$

in $A(\langle x^\alpha, x^\beta \rangle) = A(M_{A_2})$ and hence the relation above also holds in $A(V)$. It is also shown in Eq. (3.13) of [JL16] that if $r_\alpha r_\beta = r_\beta r_\alpha$ then $[x^\alpha][x^\beta] = [x^\beta][x^\alpha]$, implying the relation $[\mathbb{1} - 4x^\alpha][\mathbb{1} - 4x^\beta] = [\mathbb{1} - 4x^\beta][\mathbb{1} - 4x^\alpha]$. Therefore, the map f defines an algebra homomorphism from $T_{n+1} = \mathbb{C}[\mathfrak{S}_{n+1}]/\mathfrak{b}$ to $A(V)/\mathfrak{a}$. Since $A(V)$ is generated by $[x^\alpha]$ with $\alpha \in \Phi(A_n)^+$ by Lemma 4.1, f is indeed surjective. Thus we have obtained the following surjections.

$$T_{n+1} \xrightarrow{f} A(V)/\mathfrak{a} \longrightarrow A(M_{A_n})/\mathfrak{a}.$$

It is shown in Lemma 4.8 of [JL16] that T_{n+1} and $A(M_{A_n})/\mathfrak{a}$ are isomorphic. Therefore, f is also an isomorphism. \blacksquare

Corollary 4.3. *An irreducible V -module of σ -type is an irreducible module over the simple quotient M_{A_n} . In particular, there exist finitely many irreducible V -modules of σ -type.*

Since the adjoint V -module V is of σ -type, we have:

Lemma 4.4. *The adjoint V -module V has a composition series.*

Lemma 4.5. *The Virasoro vector η^i , $1 \leq i \leq n$, are simple. That is, $\langle \eta^i \rangle \cong L(c_i, 0)$.*

Proof: We prove by induction on n . Suppose $V^{[i]} \cong M_{A_i}$ for $1 \leq i < n$. Then η^i are simple for $1 \leq i < n$. It suffices to show that η^n is simple. Since $\omega^{n-1} = \eta^1 + \dots + \eta^{n-1}$ is the conformal vector of $V^{[n-1]}$, $V^{[n-1]} \otimes \langle \eta^n \rangle$ is a full sub VOA of $V^{[n]} = V$ (cf. [FZ92]). By induction, $V^{[n-1]}$ is simple and isomorphic to $M_{A_{n-1}}$. Suppose $\langle \eta^n \rangle$ is not simple. Then it follows from the structure of the Verma modules over the Virasoro algebra (cf. [IK11, W93]) that $\langle \eta^n \rangle$ has a singular vector u of weight $(n+1)(n+2)$. It follows from Lemma 4.4 that there exists a composition series $V = J_0 \supset J_1 \supset \dots \supset J_k = 0$ such that J_i/J_{i+1} is an irreducible V -module of σ -type. It follows from Corollary 4.3 that J_i/J_{i+1} are irreducible M_{A_n} -modules. Since M_{A_n} is the unique simple quotient of V by Theorem 3.6, the top quotient V/J_1 is isomorphic to M_{A_n} . It is known (cf. [LS08]) that η^i is a simple Virasoro vector in the simple quotient M_{A_n} so that u is zero in the quotient V/J_1 , i.e., $u \in J_1$. So there exists a factor J_i/J_{i+1} such that $u \in J_i$ but $u \notin J_{i+1}$. Then $u + J_{i+1}$ is a highest weight vector for the simple Virasoro vector $\eta^n + J_1$ of $V/J_1 \cong M_{A_n}$ with highest weight $(n+1)(n+2)$, which contradicts the classification of irreducible $L(c_n, 0)$ -modules (cf. (3.1)). Thus η^n is simple in V . \blacksquare

Theorem 4.6. *Let V be a VOA satisfying Conditions 1 such that $G_V = \mathfrak{S}_{n+1}$. Then V is simple and isomorphic to M_{A_n} .*

Proof: We prove the theorem by induction on $n > 2$ and we assume that $V^{[i]} \cong M_{A_i}$ for $1 \leq i < n$. By the definition, $M_{A_i} \otimes L_{\widehat{\mathfrak{sl}}_2}(i+1, 0)$ is a full sub VOA of $L_{\widehat{\mathfrak{sl}}_2}(1, 0)^{\otimes i+1}$. It is shown in [JL16] that all the irreducible M_{A_i} -modules of σ -type are given by the decomposition

$$L_{\widehat{\mathfrak{sl}}_2}(1, 0)^{\otimes i+1} = \bigoplus_{0 \leq 2j \leq i+1} M_{A_i}(2j) \otimes L_{\widehat{\mathfrak{sl}}_2}(i+1, 2j), \quad (4.2)$$

where $M_{A_i}(2j)$, $0 \leq 2j \leq i+1$, are inequivalent irreducible M_{A_i} -modules of σ -type. The top weight of $M_{A_i}(2j)$ is positive if $j > 0$ and $M_{A_i}(0)$ is isomorphic to the adjoint module M_{A_i} . By Lemma 4.5, $V = V^{[n]}$ has a full sub VOA $V^{[n-1]} \otimes \langle \eta^n \rangle$ isomorphic to $M_{A_{n-1}} \otimes L(c_n, 0)$ which is rational by [JL16, W93]. It is shown in [GKO86, LS08] that $M_{A_n}(2j)$ has the following decomposition as an $M_{A_{n-1}} \otimes L(c_n, 0)$ -module:

$$M_{A_n}(2j) = \bigoplus_{0 \leq 2k \leq n} M_{A_{n-1}}(2k) \otimes L(c_n, h_{2k+1, 2j+1}^{(n)}). \quad (4.3)$$

Let J be the maximal ideal of V . Then $V/J \cong M_{A_n}$ by Theorem 3.6. Since $V^{[n-1]} \otimes \langle \eta^n \rangle$ is rational, there is a $V^{[n-1]} \otimes \langle \eta \rangle$ -submodule X such that $V = X \oplus J$. Since the Griess algebra of V and that of M_{A_n} are isomorphic, we have $J \cap V_2 = 0$ and $V_2 \subset X$. Therefore, it suffices to show that X forms a subalgebra of V since V is generated by the Griess algebra. By Lemma 4.4, J has a composition series, and since V is of CFT-type, there is no composition factor isomorphic to M_{A_n} in J . It follows from the fusion rules in (3.2) that $L(c_n, h_{2k+1, 1}^{(n)})$, $1 \leq 2k \leq n$, are closed under the fusion product, and it follows from (3.1) that $h_{r, s}^{(n)} = h_{r', s'}^{(n)}$ if and only if $(r, s) = (r', s')$ or $(r + r', s + s') = (n + 2, n + 3)$. Therefore there is no irreducible $\langle \eta^n \rangle$ -submodule isomorphic to $L(c_n, h_{2k+1, 1}^{(n)})$ in $M_{A_n}(2j)$ if $j > 0$ by (4.3). Thus, $\langle X \rangle \cap J = 0$ and X forms a subalgebra of V . This completes the proof. \blacksquare

A Appendix

Here we prove Lemma 3.9. Consider the generating series

$$\psi(z) = \sum_{n \in \mathbb{Z}} \psi_{n+1/2} z^{-n-1}, \quad [\psi_r, \psi_s]_+ = \delta_{r+s, 0}, \quad r, s \in \mathbb{Z} + 1/2. \quad (A.1)$$

Then the SVOA $L(1/2, 0) \oplus L(1/2, 1/2)$ can be explicitly realized as follows (cf. [KR87]).

$$L(1/2, h) = \text{Span}_{\mathbb{C}} \{ \psi_{-r_1} \cdots \psi_{-r_k} \mathbb{1} \mid r_1 > \cdots > r_k > 0, k \equiv 2h \pmod{2} \}, \quad h = 0, 1/2,$$

$$Y(\mathbb{1}, z) = \text{id}, \quad Y(\psi_{-1/2} \mathbb{1}, z) = \psi(z), \quad \omega = \frac{1}{2} \psi_{-3/2} \psi_{-1/2} \mathbb{1},$$

where $\psi_r \mathbb{1} = 0$ for $r > 0$.

Let $e \in V$ be an Ising vector. Let $b \in V_2$ be a highest weight vector for $\langle e \rangle$ with highest weight $1/2$ and $y \in V_N$ a highest weight vector for $\langle e \rangle$ with highest weight h . Consider V as a $\langle e \rangle \otimes \text{Com}_V \langle e \rangle$ -module. We can write $2e = \psi_{-3/2} \psi_{1/2} \mathbb{1}$, $b = \psi_{-1/2} \mathbb{1} \otimes v$ and $y = w \otimes x$, where $\psi_{-1/2} \mathbb{1}$ and w are highest weight vectors of $L(1/2, 1/2)$ and $L(1/2, h)$, respectively. By (3.2), $L(1/2, 0)$ and $L(1/2, 1/2)$ are simple current modules. It then follows from Theorem 2.10 of [ADL05] that we have a factorization

$$Y(b, z)y = \psi(z)w \otimes J(v, z)x \quad (\text{A.2})$$

where $J(\cdot, z)$ is an intertwining operator⁴ among $\text{Com}_V \langle e \rangle$ -submodules of V .

First, we consider the case $h = 0$. Then $w = \mathbb{1}$ and as a $\langle e \rangle \otimes \text{Com}_V \langle e \rangle$ -module, we have

$$e = \frac{1}{2} \psi_{-3/2} \psi_{-1/2} \mathbb{1}, \quad b = \psi_{-1/2} \mathbb{1} \otimes v = \psi_{-1/2} v_{(-1)} \mathbb{1}, \quad y = \mathbb{1} \otimes x, \quad Y(b, z) = \psi(z) \otimes J(v, z).$$

For $n > 0$, we also have

$$\psi_{-n-1/2} \mathbb{1} = \frac{1}{n!} e_{(0)}^n \psi_{-1/2} \mathbb{1}.$$

Since $b_{(k)}y \in V_{N-k+1}^{[1/2]}_e$ (cf. Eq. (3.3)), we have $b_{(k)}y = 0$ and $v_{(k)}x = 0$ if $k \geq N$. (Note that $V_{\leq 1}^{[1/2]}_e = 0$ since V is of OZ-type.) We set $P_{1/2, k, j} = 0$ if $k \geq N$ or $k + j \geq N$. Let $i \geq 0$. Then from (A.2) with $w = \mathbb{1}$, we have

$$\begin{aligned} b_{(N-1)}y &= \psi_{-1/2} v_{(N-1)}x, \\ b_{(N-2)}y &= \psi_{-1/2} v_{(N-2)}x + e_{(0)} \psi_{-1/2} v_{(N-1)}x, \\ b_{(N-3)}y &= \psi_{-1/2} v_{(N-3)}x + e_{(0)} \psi_{-1/2} v_{(N-2)}x + \frac{e_{(0)}^2}{2!} \psi_{-1/2} v_{(N-1)}x, \\ &\vdots \\ b_{(N-i-1)}y &= \sum_{j=0}^i \frac{e_{(0)}^j}{j!} \psi_{-1/2} v_{(N-i+j-1)}x. \end{aligned}$$

Multiplying $e_{(0)}^{i-j}$ with $b_{(N-j-1)}y$ for $0 \leq j \leq i$, we obtain the following linear system:

$$\begin{aligned} e_{(0)}^i b_{(N-1)}y &= e_{(0)}^i \psi_{-1/2} v_{(N-1)}x, \\ e_{(0)}^{i-1} b_{(N-2)}y &= e_{(0)}^i \psi_{-1/2} v_{(N-1)}x + e_{(0)}^{i-1} \psi_{-1/2} v_{(N-2)}x, \\ e_{(0)}^{i-2} b_{(N-3)}y &= \frac{e_{(0)}^i}{2!} \psi_{-1/2} v_{(N-1)}x + e_{(0)}^{i-1} \psi_{-1/2} v_{(N-2)}x + e_{(0)}^{i-2} \psi_{-1/2} v_{(N-3)}x, \\ &\vdots \\ b_{(N-i-1)}y &= \sum_{j=0}^i \frac{e_{(0)}^j}{j!} \psi_{-1/2} v_{(N-i+j-1)}x. \end{aligned}$$

⁴Actually, the modes of $J(v, z)$ generate the $c = 7/10$ Neveu-Schwarz algebra. In the following argument, we do not need this fact.

This system can be formatted as follows.

$$\begin{bmatrix} e_{(0)}^i b_{(N-1)}y \\ e_{(0)}^{i-1} b_{(N-2)}y \\ e_{(0)}^{i-2} b_{(N-3)}y \\ \vdots \\ e_{(0)} b_{(N-i)}y \\ b_{(N-i-1)}y \end{bmatrix} = \exp(\mathbb{J}) \begin{bmatrix} e_{(0)}^i \psi_{-1/2} v_{(N-1)}x \\ e_{(0)}^{i-1} \psi_{-1/2} v_{(N-2)}x \\ e_{(0)}^{i-2} \psi_{-1/2} v_{(N-3)}x \\ \vdots \\ e_{(0)} \psi_{-1/2} v_{(N-i)}x \\ \psi_{-1/2} v_{(N-i-1)}x \end{bmatrix}, \quad \mathbb{J} = \begin{bmatrix} 0 & & & & & \\ 1 & 0 & & & & \\ & 1 & 0 & & & \\ & & \ddots & \ddots & & \\ & & & 1 & 0 & \\ & & & & 1 & 0 \end{bmatrix}.$$

Therefore, we obtain

$$\begin{bmatrix} e_{(0)}^i \psi_{-1/2} v_{(N-1)}x \\ e_{(0)}^{i-1} \psi_{-1/2} v_{(N-2)}x \\ e_{(0)}^{i-2} \psi_{-1/2} v_{(N-3)}x \\ \vdots \\ e_{(0)} \psi_{-1/2} v_{(N-i)}x \\ \psi_{-1/2} v_{(N-i-1)}x \end{bmatrix} = \exp(-\mathbb{J}) \begin{bmatrix} e_{(0)}^i b_{(N-1)}y \\ e_{(0)}^{i-1} b_{(N-2)}y \\ e_{(0)}^{i-2} b_{(N-3)}y \\ \vdots \\ e_{(0)} b_{(N-i)}y \\ b_{(N-i-1)}y \end{bmatrix},$$

and

$$\psi_{-1/2} v_{(N-i-1)}x = \sum_{j=0}^i \frac{(-e_{(0)})^j}{j!} b_{(N-i+j-1)}y = b_{(N-i-1)}y + \sum_{j=1}^i \frac{(-e_{(0)})^j}{j!} b_{(N-i+j-1)}y.$$

The left hand side is a highest weight vector for $\langle e \rangle$ and we obtain the lemma by setting $P_{1/2, N-i-1, j} = (-e_{(0)})^j / j!$.

Next we consider the case $h = 1/2$ and $w = \psi_{-1/2} \mathbb{1}$. Since $\psi(z)\psi_{-1/2} \mathbb{1} \in \langle e \rangle$, there exist $Q_{m+2} \in \text{U}(\text{Vir}(e_-)[m+2])$ such that $\psi_{-m-3/2} \psi_{-1/2} \mathbb{1} = Q_{m+2} \mathbb{1}$ and

$$\psi(z)\psi_{-1/2} \mathbb{1} = \mathbb{1}z^{-1} + \sum_{m \geq 0} \psi_{-m-3/2} \psi_{-1/2} \mathbb{1}z^{m+1} = \mathbb{1}z^{-1} + \sum_{m \geq 0} Q_{m+2} \mathbb{1}z^{m+1}. \quad (\text{A.3})$$

Note that $Q_{m+2} \in \text{U}(\text{Vir}(e_-)[m+2])$ is not unique but $Q_{m+2} \mathbb{1}$ is uniquely determined in $\langle e \rangle \cong L(1/2, 0)$. Since $b_{(k)}y \in V_{N-k+1} = 0$ if $k > N+1$, $b_{(N+1)}y \in V_0 = \mathbb{C}\mathbb{1}$ and $b_{(N)}y \in V_1 = 0$, we set $P_{0, k, j} = 0$ for $k \geq N$ and $j \geq 0$. We also set $P_{0, k, j} = 0$ if $k+j > N+1$. We will define $P_{0, k, j}$ with $j \geq 0$ recursively for $k < N$ as follows.

Suppose $i > 0$ and for $k > N-i$ there exist $P_{0, k, j}$, $j \geq 0$, such that

$$v_{(k-1)}x = b_{(k)}y + \sum_{j > 0} P_{0, k, j} b_{(k+j)}y. \quad (\text{A.4})$$

By (A.2) and (A.3) we have

$$b_{(N-i)}y = v_{(N-i-1)}x + \sum_{0 \leq m \leq i-1} Q_{m+2} v_{(N-i+m+1)}x$$

and

$$v_{(N-i-1)}x = b_{(N-i)}y - \sum_{0 \leq m \leq i-1} Q_{m+2} \left(b_{(N-i+m+2)}y + \sum_{j>0} P_{0,N-i+m+2,j} b_{(N-i+m+2+j)}y \right).$$

Expanding the right hand side, we obtain recursive relations for $P_{0,N-i,j}$ such that (A.4) holds for $k = N - i$. Note that all $P_{0,k,j}$ are determined only by $k, j, b_{(N+1)}y \in V_0$ and $Q_{m+2}\mathbb{1} \in \langle e \rangle$, and $Q_{m+2}\mathbb{1}$ is uniquely determined only by the structure of $L(1/2, 0)$ and independent of the structure of V itself. This completes the proof of Lemma 3.9. ■

References

- [ADL05] T. Abe, C. Dong and H. Li, Fusion rules for the vertex operator algebras $M(1)^+$ and V_L^+ . *Comm. Math. Phys.* **253** (2005), 171–219.
- [CH95] H. Cuypers and J.I. Hall, The 3-transposition groups with trivial center. *J. Algebra* **178** (1995), 149–193.
- [DLM96] C. Dong, H. Li and G. Mason, Compact automorphism groups of vertex operator algebras. *Internat. Math. Res. Notices* **18** (1996), 913–921.
- [DLMN98] C. Dong, H. Li, G. Mason and S.P. Norton, Associative subalgebras of Griess algebra and related topics. Proc. of the Conference on the Monster and Lie algebra at the Ohio State University, May 1996, ed. by J. Ferrar and K. Harada, Walter de Gruyter, Berlin - New York, 1998, pp. 27–42.
- [DMZ94] C. Dong, G. Mason and Y. Zhu, Discrete series of the Virasoro algebra and the moonshine module. Proc. Symp. Pure. Math., American Math. Soc. **56** II (1994), 295–316.
- [FZ92] I.B. Frenkel and Y. Zhu, Vertex operator algebras associated to representation of affine and Virasoro algebras. *Duke Math. J.* **66** (1992), 123–168.
- [GKO86] P. Goddard, A. Kent and D. Olive, Unitary representations of the Virasoro and super-Virasoro algebra. *Commun. Math. Phys.* **103** (1986), 105–119.
- [HRS15] J.I. Hall, F. Rehren and S. Shpectorov, Primitive axial algebras of Jordan type. *J. Algebra* **437** (2015), 79–115.
- [IK11] K. Iohara and Y. Koga, Representation Theory of the Virasoro Algebra. Springer Monographs in Mathematics. Springer-Verlag, London, 2011.
- [KR87] V. Kac and A. Raina, Highest Weight Representations of Infinite Dimensional Lie Algebras. World Scientific. Adv. Ser. In Math. Phys., Singapore, 1987.

- [JL16] C. Jiang and Z. Lin, The commutant of $L_{\widehat{\mathfrak{sl}_2}}(n, 0)$ in the vertex operator algebra $L_{\widehat{\mathfrak{sl}_2}}(1, 0)^{\otimes n}$. *Adv. Math.* **301** (2016), 227–257.
- [LS08] C.H. Lam and S. Sakuma, On a class of vertex operator algebras having a faithful S_{n+1} -action. *Taiwanese Journal of Mathematics* **12** (2008), 2465–2488.
- [LSY07] C.H. Lam, S. Sakuma and H. Yamauchi, Ising vectors and automorphism groups of commutant subalgebras related to root systems. *Math. Z.* **255**(3) (2007), 597–626.
- [LY00] C. Lam and H. Yamada, $\mathbb{Z}_2 \times \mathbb{Z}_2$ codes and vertex operator algebras. *J. Algebra* **224** (2000), 268–291.
- [LL04] J. Lepowsky and H. Li, Introduction to Vertex Operator Algebras and Their Representations. Progress in Mathematics, Vol. 227, Birkhäuser Boston, Inc., Boston, MA, 2004.
- [Li94] H. Li, Symmetric invariant bilinear forms on vertex operator algebras. *J. Pure Appl. Alg.* **96** (1994), 279–297.
- [Ma05] A. Matsuo, 3-transposition groups of symplectic type and vertex operator algebras. *J. Math. Soc. Japan* **57**(3) (2005), 639–649. [arXiv:math/0311400](https://arxiv.org/abs/math/0311400).
- [Mi96] M. Miyamoto, Griess algebras and conformal vectors in vertex operator algebras. *J. Algebra* **179** (1996), 528–548.
- [W93] W. Wang, Rationality of Virasoro vertex operator algebras. *Internat. Math. Res. Notices* **71** (1993), 197–211.
- [Z96] Y. Zhu, Modular invariance of characters of vertex operator algebras. *J. Amer. Math. Soc.* **9** (1996), 237–302.