

Unobstructedness of Galois deformation rings associated to RACSDC automorphic representations

David-A. Guiraud
IWR Heidelberg

February 28, 2017

Abstract

Let F be a CM field and let $(\bar{\tau}_{\pi,\lambda})_\lambda$ be the compatible system of residual \mathcal{G}_n -valued representations of Gal_F attached to a RACSDC automorphic representation π of $\text{GL}_n(\mathbb{A})$, as studied by Clozel, Harris and Taylor [CHT08] and others. Under mild assumptions, we prove that the fixed-determinant universal deformation rings attached to $\bar{\tau}_{\pi,\lambda}$ are unobstructed for all places λ in a subset of Dirichlet density 1, continuing the investigations of Mazur, Weston and Gamzon. During the proof, we develop a general framework for proving unobstructedness (which could be useful for other applications in future) and an $R = T$ -theorem, relating the universal crystalline deformation ring of $\bar{\tau}_{\pi,\lambda}$ and a certain unitary fixed-type Hecke algebra.

1 Introduction

This article is concerned with unobstructedness of certain Galois deformation rings. For the purpose of this introduction, let F be a number field, let k be a finite field of characteristic ℓ and fix an absolutely irreducible representation

$$\bar{\rho} : \text{Gal}_{F,S} \rightarrow \text{GL}_n(k),$$

where $S \subset \text{Pl}_F$ is a finite set of places. Then assigning to a complete Noetherian local algebra A over the ring W of Witt vectors of k the set of all $\text{GL}_n(A)$ -valued deformations of $\bar{\rho}$ defines a functor which is representable by a universal deformation ring $R_S(\bar{\rho})$, studied first by Mazur [Maz87]. It is easily seen that a vanishing of the cohomology group $H^2(\text{Gal}_{F,S}, \text{ad } \bar{\rho})$ implies that $R_S(\bar{\rho})$ is formally smooth, i.e. isomorphic to a power series ring over W . In this sense, the group $H^2(\text{Gal}_{F,S}, \text{ad } \bar{\rho})$ can be interpreted as the obstruction to the smoothness of $R_S(\bar{\rho})$, and we say that $R_S(\bar{\rho})$ is *unobstructed* if $H^2(\text{Gal}_{F,S}, \text{ad } \bar{\rho}) = 0$. We remark the following connection with a conjecture of Jannsen: Assume that $\bar{\rho}$ is the reduction of the ℓ -adic representation $\rho_{f,\ell}$ attached to a cuspidal modular eigenform f (cf. [Del71, Shi71, DS74]). Then the Frobenius eigenvalues of $\rho_{f,\ell}$ are Weil-numbers of some fixed weight w , i.e. $\rho_{f,\ell}$ is pure of weight w . A conjecture of Jannsen [Jan89, Conjecture 1] (see also [Bel09, Conjecture 5.1]) predicts the vanishing of $H^2(\text{Gal}_{F,S}, \text{ad } \rho)$. This implies that $H^2(\text{Gal}_{F,S}, \Lambda)$ is finite and torsion, where $\Lambda \subset \text{ad } \rho$ denotes an integral $\text{Gal}_{F,S}$ -stable lattice. On the other hand, our residual H^2 -vanishing implies the vanishing of $H^2(\text{Gal}_{F,S}, \Lambda)$ by Nakayama's Lemma. This, in turn, implies the vanishing of $H^2(\text{Gal}_{F,S}, \text{ad } \rho)$, as predicted by Jannsen. Besides this application, the numerous usages of Galois deformation theoretic methods in number theory indicate that the structure of universal deformation rings is of independent interest.

Unobstructedness for Galois representations attached to automorphic objects can usually not be expected for all choices of ℓ . The best we can hope for is that unobstructedness holds for almost all primes (or: for all primes in a subset of Dirichlet density 1), and this question has been studied (under different technical assumptions) in the following cases:

- For $\bar{\rho}$ the reduction of the representation $\rho_{E,\ell}$ attached to an elliptic curve E over $F = \mathbb{Q}$, cf. [Maz97];
- for $\bar{\rho}$ the reduction of the representation $\rho_{f,\ell}$ attached to a newform f of weight $k \geq 3$ over $F = \mathbb{Q}$, cf. [Wes04] (but see also [Yam04, Hat15]);
- for $\bar{\rho}$ the reduction of the representation $\rho_{f,\ell}$ attached to a Hilbert eigenform f over a totally real field F , cf. [Gam13].

Note that $n = 2$ in all these cases.

In this article, we develop a general framework for proving unobstructedness, which differs significantly from the previous approaches and which uses an $R = T$ -theorem as the main ingredient. We apply this framework to the reduction of the Galois representation attached to a regular algebraic conjugate self-dual¹ cuspidal (RACSDC) automorphic representation π of $\mathrm{GL}_n(\mathbb{A}_F)$ with ramification set S , where F is a CM field. In order to give a more precise statement, we have to recall that π gives in the first instance rise not to GL_n -valued representations, but to morphisms $r_{\pi,\lambda} : \mathrm{Gal}_{F^+} \rightarrow \mathcal{G}_n(\overline{\mathbb{Q}}_{\ell(\lambda)})$, where λ runs through the places of the coefficient field of π , where \mathcal{G}_n denotes the group scheme from [CHT08, Section 2.1] and where $\ell(\lambda)$ denotes the rational prime below λ . We make the following assumption:

Assumption 1.1. The set of the λ for which the GL_n -valued representation $\bar{r}_{\pi,\lambda}|_{\mathrm{Gal}_F}$ is absolutely irreducible has Dirichlet density 1.

We remark that this assumption is fulfilled e.g. if $n \leq 5$ or if π is extremely regular, or would follow from absolute irreducibility of the ℓ -adic system $(r_{\pi,\lambda}|_{\mathrm{Gal}_F})$, cf. Remark 8.2. For the following, we fix for each λ a lift χ of the character $\mathfrak{m} \circ \bar{r}_{\pi,\lambda}$ of Gal_F , where \mathfrak{m} is the multiplier character of the group \mathcal{G}_n , cf. Section 6.1. By $R_{S_\ell}^\chi(\bar{r}_{\pi,\lambda})$ we denote the universal ring parametrizing deformations r of $\bar{r}_{\pi,\lambda}$ which are unramified outside the places which are in S or divide $\infty, \ell(\lambda)$ and which fulfill $\mathfrak{m} \circ r = \chi$. The correct unobstructedness requirement is then the vanishing of $H^2(\mathrm{Gal}_{F,S_\ell}, \mathfrak{g}_n^{\mathrm{der}})$, where $\mathfrak{g}_n^{\mathrm{der}}$ denotes the Lie algebra of the derived subgroup of \mathcal{G}_n . Our main result is:

Theorem 1.2. *Assume that all Hodge-Tate weights of $r_{\pi,\lambda}$ (which are independent of λ , as the $r_{\pi,\lambda}$ form a compatible system) are non-consecutive: if $a, b \in \mathbb{Z}$ show up as Hodge-Tate weights, then $|a - b| \neq 1$. Then, for all λ in a set of places of Dirichlet density 1 the universal deformation ring $R_{S_\ell}^\chi(\bar{r}_{\pi,\lambda})$ is unobstructed.*

Remark that we do not require a particular splitting behavior at the places in S . We also want to stress that the developed framework is flexible and in principle applicable to Galois representations with values in other groups and can be used to establish unobstructedness of universal deformation rings with imposed deformation conditions which are more sophisticated than the fixed-determinant condition $\mathfrak{m} \circ r = \chi$. Therefore, we hope that the framework will be useful for other applications, as better modularity lifting results become available in future. We also remark that presently the

¹We remark that, in light of the results of [BLGGT14], it should be possible to weaken the conjugate self-duality assumption to an essentially self-duality assumption, thus treating RACESDC automorphic representations.

condition on the Hodge-Tate weights is necessary in order to use a local unobstructedness property at the places above $\ell(\lambda)$, a technical inconvenience we expect to weaken in future work.

We give a short outline of the article: After some remarks about notation, we start in Section 3 with a collection of the general deformation theoretic methods we will use. Moreover, we will define a suitably flexible notion of unobstructedness for conditioned deformation functors (Definition 3.27). In Section 4, we state and prove the core framework (Theorem 4.2), which uses a list of six assumptions as input and provides unobstructedness as output. This framework is presented with respect to local deformation conditions **crys**, **min**, **sm**, which have a purely formal meaning throughout Section 4. The main input is the formal smoothness of the deformation ring with respect to the conditions **min** and **crys**, which is the natural output of a suitable $R = T$ -theorem, and the desired unobstructedness is then deduced by commutative algebra arguments and comparing dimensions. It is the purpose of Section 5 to introduce and study useful local conditions which will be plugged in into the framework theorem later. After a reminder on the association of Galois deformations to automorphic forms, the additional results are provided in Section 7: We consider the deformation ring $R^{\text{min,crys}} := R_{S_\ell}^{\chi,\text{min,crys}}(\bar{\tau}_{\pi,\lambda})$ parametrizing those lifts which are minimally ramified (in the sense of Section 5.4) at all places in S and crystalline (in the Fontaine-Laffaille range) at all places dividing ℓ . Moreover, we consider a corresponding Hecke algebra \mathbb{T}^{min} which is defined as the localization of a certain endomorphism algebra of automorphic forms of the same weight and level as π , and with a certain fixed type-requirement at the places in S . Then, using the modularity lifting results of [BLGGT14], we show

Theorem 1.3. $R^{\text{min,crys}} \cong \mathbb{T}^{\text{min}}$ and, for almost all λ , $\mathbb{T}^{\text{min}} \cong W$.

This result is crucial to prove in Section 8 that, for almost all λ , there exists a suitable finite solvable extension F' of F such that the deformation ring $R_{S_\ell}^{\chi,\text{min}}(\bar{\tau}_{\pi,\lambda} | \text{Gal}_{F',S})$, parametrizing deformations of the base change of $\bar{\tau}_{\pi,\lambda}$ to F' which are minimally ramified at all places above S , is unobstructed. Moreover, we show that the minimally-ramified condition can be waived for almost all λ (Theorem 7.11). It is important to keep track of the different field extensions necessary when running through all λ , so that we are left with a set of Dirichlet density 1 to which we can apply a result on potential unobstructedness (Lemma 4.8) and finally deduce the claim of Theorem 1.2.

Acknowledgements The main part of this article grew out of my PhD thesis [Gui16] at the University of Heidelberg. I would like to thank my doctoral advisor Gebhard Böckle for his support and help, throughout my PhD studies and thereafter. Moreover, I would like to thank the Graduiertenkolleg Heidelberg (Stipendium nach dem Landesgraduiertenförderungsgesetz), MATCH (The Mathematics Center Heidelberg), the DAAD (Promos Stipendium) and the DFG (FG1920) for the funding received during my PhD studies and thereafter.

2 Notation

Before we start with the main body of this article, let us make some remarks on the used notation: If F denotes a number field, we denote by Pl_F the set of places of F and by Pl_F^{fin} the set of finite places of F . Moreover, we set $\Omega_\infty^F = \text{Pl}_F \setminus \text{Pl}_F^{\text{fin}}$ and, for a rational prime ℓ , we denote by Ω_ℓ^F the set of places of F dividing ℓ . If F is understood, we will simply write Ω_∞ and Ω_ℓ . For a place $\lambda \in \text{Pl}_F^{\text{fin}}$ we define $\ell(\lambda)$ (or ℓ , if λ is understood) as the rational prime below λ . If $S \subset \text{Pl}_F^{\text{fin}}$ and ℓ is some rational prime, we set $S_\ell := S \cup \Omega_\infty \cup \Omega_\ell$.

We denote by \widetilde{F} the Galois closure of F . When dealing with a quadratic extension $F|F^+$, we will denote by c the non-trivial element of the Galois group $\text{Gal}(F|F^+)$. Moreover, for a rational prime ℓ , we denote by $\epsilon_\ell : \text{Gal}_F \rightarrow \overline{\mathbb{Z}}_\ell^\times$ the ℓ -adic cyclotomic character and by $\bar{\epsilon}_\ell$ its mod- ℓ reduction.

If $L|F$ is a finite extension and S is a fixed set of places of F , we will write $S\langle L \rangle$ for the set $\{\nu' \in \text{Pl}_L : \nu' \text{ divides some } \nu \in S\}$. For example, if $\text{Gal}_{F,S}$ denotes the Galois group of the maximal, unramified outside S extension of F , we will write $\text{Gal}_{L,S\langle L \rangle}$ for the Galois group of the maximal, unramified outside $S\langle L \rangle$ extension of L . However, if there is no risk of confusion, we will often simplify this and write S in place of $S\langle L \rangle$ and hence $\text{Gal}_{L,S}$ instead of $\text{Gal}_{L,S\langle L \rangle}$. In a completely analogous way, if S is a subset of Pl_L , we will write $S\langle F \rangle$ (or, if there is no risk of confusion, simply S) for the set $\{\nu' \in \text{Pl}_F : \nu' \text{ is divided by some } \nu \in S\}$. If ρ is a representation of Gal_F and ν a place of F , we will use the symbol ρ_ν for the restriction of ρ to a decomposition subgroup at ν .

For a topological group Γ and a topological ring R , we denote by $\text{Rep}_R(\Gamma)$ the category of finitely generated R -modules with a continuous Γ -action. If A is a Γ -module, we denote by A^* the Pontryagin dual and by A^\vee the Tate dual of A .

We will often make statements concerning variations of deformation rings and we will shorten this using brackets. E.g., we will use the notation $R^{(\chi),[\text{min}]}(\bar{\rho}) = 0$ as a shortcut for the four statements $R(\bar{\rho}) = 0, R^\chi(\bar{\rho}) = 0, R^{\text{min}}(\bar{\rho}) = 0$ and $R^{\chi,\text{min}}(\bar{\rho}) = 0$. For cohomology groups, we abbreviate $h^i(*,*)$ for $\dim H^i(*,*)$.

Let k be a finite field of characteristic ℓ . For the valuation ring Λ of a finite extension of \mathbb{Q}_ℓ with residue field $k_\Lambda = k$, we will consider the category \mathcal{C}_Λ of complete Noetherian local Λ -algebras A fulfilling $k_A = k$.

3 Liftings and deformations

In this section, which contains nothing original, we recall the main results on deformation theory. For general background literature, we refer the reader to [Til96, Mau00, Lev13, Bal12, BC03]. Let us first fix a finite field k and denote $\ell = \text{char}(k)$. We will denote the ring of Witt vectors over k by $W(k)$, or, if k is understood, by W . Moreover, let us fix a profinite group Γ which fulfills the ℓ -finiteness condition (Φ_ℓ) of [Maz87]: For any open subgroup $H \subset \Gamma$, the maximal pro- ℓ quotient of H is topologically finitely generated.

Let G be a smooth linear algebraic group over W and fix a continuous group homomorphism $\bar{\rho} : \Gamma \rightarrow G(k)$, where $G(k)$ carries the discrete topology.

Basic facts on coefficient rings Let us first state some basic facts on the category \mathcal{C}_Λ , whose proofs we leave to the reader: The pullback in \mathcal{C}_Λ is realized by the completed tensor product $\widehat{\otimes}$, cf. [Maz97, §12]. Consequently, if $C \leftarrow A \rightarrow B$ is a diagram in \mathcal{C}_Λ , then $\text{Hom}_{\mathcal{C}_\Lambda}(B \widehat{\otimes}_A C, _)$ is the pullback of the diagram of functors $\text{Hom}_{\mathcal{C}_\Lambda}(C, _) \rightarrow \text{Hom}_{\mathcal{C}_\Lambda}(A, _) \leftarrow \text{Hom}_{\mathcal{C}_\Lambda}(B, _)$.

Consider a pushout diagram in \mathcal{C}_Λ where one arrow (say, f) is surjective. This implies that the parallel arrow (say, g) is surjective as well, so taking $I = \ker(f)$ and $J = \ker(g)$ we can extend the orthogonal arrow (say, π) to a map of short exact sequences of Λ -modules:

$$\begin{array}{ccc}
 \begin{array}{ccc} A & \xrightarrow{f} & B \\ \pi \downarrow & & \downarrow \text{dotted} \\ C & \xrightarrow{g} & P \end{array} & \sim & \begin{array}{ccccccc} 0 & \longrightarrow & I & \longrightarrow & A & \xrightarrow{f} & B & \longrightarrow & 0 \\ & & \pi|I \downarrow & & \pi \downarrow & & \downarrow & & \\ 0 & \longrightarrow & J & \longrightarrow & C & \xrightarrow{g} & P & \longrightarrow & 0 \end{array}
 \end{array}$$

If \mathfrak{J} is a finitely generated ideal of some $D \in \mathcal{C}_\Lambda$ we denote cardinality of a minimal set of generators of \mathfrak{J} by $\text{gen}_D(\mathfrak{J}) := \dim_k \mathfrak{J}/\mathfrak{m}_D \mathfrak{J}$. Then, we easily see that the following holds for the above diagram:

Proposition 3.1. $\text{gen}_C(J) \leq \text{gen}_A(I)$.

Moreover, we have the following results, which follow easily from standard facts about regular systems of parameters (cf. [Ser00, Proposition 22] and its use in Section 2 of [Gui16]) :

Lemma 3.2. *Suppose $A = \Lambda[[x_1, \dots, x_a]]$, $B = \Lambda[[x_1, \dots, x_b]] \in \mathcal{C}_\Lambda$ and let $J \subset A$ be an ideal of the form $J = (f_1, \dots, f_u)$ with $f_i \in A$ and $u \leq a$. Suppose moreover that there exists a surjective morphism $f : A/J \rightarrow B$ and denote its kernel by I . Then the following are equivalent:*

- $A/J \cong \Lambda[[x_1, \dots, x_{a-u}]]$;
- $\text{gen}_{A/J}(I) = a - u - b$;
- $\text{gen}_{A/J}(I) \leq a - u - b$.

Proposition 3.3. *Let $m \in \mathbb{N}$. Then $A \in \mathcal{C}_\Lambda$ is regular if and only if $A[[x_1, \dots, x_m]]$ is regular.*

Proposition 3.4. *Let $f : A \rightarrow B$ be a morphism in \mathcal{C}_Λ .*

1. *f is formally smooth (cf. [Gro64, §19]) if and only if B is isomorphic to a formal power series ring over A ;*
2. *Assume that A is formally smooth over Λ of relative dimension d and that f is surjective. Then f is an isomorphism if $\text{rdim}_\Lambda(B) = d + 1$ (where $\text{rdim}_\Lambda(B) = \dim_k \mathfrak{m}_B/(\mathfrak{m}_B^2, \mathfrak{m}_\Lambda)$ denotes the relative dimension over Λ).*

Lemma 3.5. *Let $A \in \mathcal{C}_\Lambda$, $m \in \mathbb{N}$ such that $\Lambda[[x_1, \dots, x_m]] \cong A \widehat{\otimes}_\Lambda \Lambda[[x]]$. Then $A \cong \Lambda[[x_1, \dots, x_{m-1}]]$.*

Lemma 3.6. *Let $\Lambda', R \in \mathcal{C}_\Lambda$, such that the structure morphism $\Lambda \rightarrow \Lambda'$ is flat. Then R is formally smooth of relative dimension $d \in \mathbb{N}$ over Λ if and only if $\Lambda' \widehat{\otimes}_\Lambda R$ is formally smooth of relative dimension d over Λ' .*

Liftings and deformations of G -valued representations

Definition 3.7. 1. A lifting of $\bar{\rho}$ to an $A \in \mathcal{C}_\Lambda$ is a continuous group homomorphism $\rho : \Gamma \rightarrow G(A)$ fulfilling $\text{mod}_{\mathfrak{m}_A} \circ \rho = \bar{\rho}$, where $\text{mod}_{\mathfrak{m}_A} : G(A) \rightarrow G(A/\mathfrak{m}_A) = G(k)$ is the canonical reduction;

2. Denote by $D_\Lambda^\square(\bar{\rho}) : \mathcal{C}_\Lambda \rightarrow \text{Sets}$ the functor which assigns to an object $A \in \mathcal{C}_\Lambda$ the set of all liftings of $\bar{\rho}$ to A .

By [Bal12, Theorem 1.2.2], $D_\Lambda^\square(\bar{\rho})$ is representable by an object $R_\Lambda^\square(\bar{\rho}) \in \mathcal{C}_\Lambda$. As an examination of its proof easily yields, we get (with respect to the ring of integers Λ' of some finite extension of $\text{Quot}(\Lambda)$ with residue field $k_{\Lambda'} = k$) an isomorphism

$$R_{\Lambda'}^\square(\bar{\rho}) \cong \Lambda' \widehat{\otimes}_\Lambda R_\Lambda^\square(\bar{\rho}). \quad (1)$$

Definition 3.8. A lifting condition is a family $\mathcal{D} = (S(A))_{A \in \mathcal{C}_\Lambda}$ of subsets $S(A) \subset D_\Lambda^\square(\bar{\rho})(A)$ s.t.

1. $\bar{\rho} \in S(k)$;
2. If $f : A \rightarrow B$ is a morphism in \mathcal{C}_Λ and $\rho \in S(A)$, then $G(f) \circ \rho \in S(B)$;
3. Let $f_1 : A_1 \rightarrow A$, $f_2 : A_2 \rightarrow A$ be morphisms in \mathcal{C}_Λ and let ρ_3 be a lifting of $\bar{\rho}$ to $A_3 := A_1 \times_A A_2$. For $i = 1, 2$ denote by $\pi_i : A_3 \rightarrow A_i$ the canonical map and by ρ_i the lifting $G(\pi_i) \circ \rho_3$ of $\bar{\rho}$ to A_i . Then, $\rho_3 \in S(A_3)$ if and only if $\rho_1 \in S(A_1)$ and $\rho_2 \in S(A_2)$.

Condition 2. guarantees that \mathcal{D} defines a subfunctor $D_\Lambda^{\square, \mathcal{D}}(\bar{\rho}) \subset D_\Lambda^{\square}(\bar{\rho})$. Condition 3. is a variation of the Mayer-Vietoris property, so a standard argument yields

Proposition 3.9. $D_\Lambda^{\square, \mathcal{D}}(\bar{\rho})$ is a relatively representable subfunctor (in the sense of [Maz97, §19]) of $D_\Lambda^{\square}(\bar{\rho})$, i.e. representable by some $R_\Lambda^{\square, \mathcal{D}}(\bar{\rho}) \in \mathcal{C}_\Lambda$. On the other hand, any representable subfunctor $F \subset D_\Lambda^{\square}(\bar{\rho})$ yields a lifting condition $\mathcal{D} = (S(A))_{A \in \mathcal{C}_\Lambda}$ via $S(A) := F(A)$.

We have the following conditioned version of (1):

$$R_{\Lambda'}^{\square, \mathcal{D}'}(\bar{\rho}) \cong \Lambda' \widehat{\otimes}_\Lambda R_\Lambda^{\square, \mathcal{D}}(\bar{\rho}), \quad (2)$$

where the condition \mathcal{D}' on the left is a truncated version of \mathcal{D} , i.e. denotes the family of those $S(A)$ as in the definition of \mathcal{D} for which $A \in \mathcal{C}_{\Lambda'}$. We will often omit this distinction and write \mathcal{D} in place of \mathcal{D}' .

Remark 3.10. Let Λ be as above and let ${}^*\mathcal{C}_\Lambda$ denote the category of complete Noetherian local Λ -algebras A such that $[k_A : k]$ is finite. Then one can extend $D_\Lambda^{\square}(\bar{\rho})$ to a functor on ${}^*\mathcal{C}_\Lambda$ by considering A -valued liftings of $\bar{\rho}$ as continuous group homomorphisms $\rho : \Gamma \rightarrow G(A)$ which fulfill $\text{mod}_{\mathfrak{m}_A} \circ \rho = \iota_{k \subset k_A} \circ \bar{\rho}$, where $\iota_{k \subset k_A} : G(k) \rightarrow G(k_A)$ is the map induced by the structure map $\Lambda \rightarrow A$. It is easy to check that this extended functor is representable by the same universal object $R_\Lambda^{\square}(\bar{\rho})$ as the functor from Definition 3.7. Moreover, if Λ' is the ring of integers of some finite extension of $\text{Quot}(\Lambda)$ such that $[k_{\Lambda'} : k] < \infty$, we have the following version of (1):

$$R_{\Lambda'}^{\square}(\iota_{k \subset k_A} \circ \bar{\rho}) \cong \Lambda' \widehat{\otimes}_\Lambda R_\Lambda^{\square}(\bar{\rho}).$$

Moreover, if \mathcal{D} is an extended lifting condition, i.e. a family $(S(A))_{A \in {}^*\mathcal{C}_\Lambda}$ fulfilling the analogue conditions of Definition 3.8 (with $A, A_i, B \in {}^*\mathcal{C}_\Lambda$), we have the following conditioned version of (2):

$$R_{\Lambda'}^{\square, \mathcal{D}}(\iota_{k \subset k_A} \circ \bar{\rho}) \cong \Lambda' \widehat{\otimes}_\Lambda R_\Lambda^{\square, \mathcal{D}}(\bar{\rho}),$$

where \mathcal{D} on the left hand side is to be understood as the Λ' -truncated version of the condition \mathcal{D} , i.e. a family indexed by ${}^*\mathcal{C}_{\Lambda'}$ instead of ${}^*\mathcal{C}_\Lambda$. Moreover, the statement of Lemma 3.6 holds if Λ' is in ${}^*\mathcal{C}_\Lambda$ instead of \mathcal{C}_Λ . (The content of this remark is strongly inspired by the treatment in [CDT99, Appendix A] and [Maz97].)

Definition 3.11. 1. A deformation of $\bar{\rho}$ to $A \in \mathcal{C}_\Lambda$ is an equivalence class of liftings to A , where two lifts are taken to be equivalent if they are conjugate by some element of $\widehat{G}(A) := \ker(\text{mod}_{\mathfrak{m}_A})$.

2. Denote by $D_\Lambda(\bar{\rho}) : \mathcal{C}_\Lambda \rightarrow \text{Sets}$ the functor which assigns to an object $A \in \mathcal{C}_\Lambda$ the set of all deformations of $\bar{\rho}$ to A .

For the following, denote by Z_G the center of G and by \mathfrak{g} (resp. by \mathfrak{z}) the Lie algebra of the special fiber of G (resp. of Z_G). We assume from now on that Z_G is formally smooth over Λ .

Theorem 3.12 ([Til96, Theorem 3.3]). *If $H^0(\Gamma, \mathfrak{g}) = \mathfrak{z}$ then $D_\Lambda(\bar{\rho})$ is representable by an object $R_\Lambda(\bar{\rho}) \in \mathcal{C}_\Lambda$.*

Observe that in the case $G = \mathrm{GL}_n$, the condition of Theorem 3.12 becomes the usual centralizer condition $\mathrm{End}_{k[\Gamma]}(\bar{\rho}) = k$. In practice, this is often deduced from absolute irreducibility of $\bar{\rho}$ by Schur's Lemma. This reasoning can be adopted to more general groups G as follows:

Definition 3.13 (Absolute Irreducibility, cf. [Ser98]). We say that $\bar{\rho}$ is absolutely irreducible if there does not exist a proper parabolic subgroup $P \subsetneq G$ over \bar{k} such that $\bar{\rho}(\Gamma) \subset P$.

Then the following can be deduced from [BMR05, Proposition 2.13]:

Lemma 3.14 (Schur's Lemma). *Assume that ℓ is very good for G (cf. [BMRT10, Section 2]) or that there exists an embedding $G \hookrightarrow \mathrm{GL}(V)$ such that $(\mathrm{GL}(V), G)$ is a reductive pair (in the sense of [BMR05, Definition 3.32]). Then $H^0(\Gamma, \mathfrak{g}) = \mathfrak{z}$ if $\bar{\rho}$ is absolutely irreducible.*

We now give an appropriate version of Definition 3.8:

Definition 3.15. A deformation condition is a lifting condition in the sense of Definition 3.8 which fulfills additionally

4. If $\rho \in S(A)$ and $g \in \widehat{G}(A)$, then $g\rho g^{-1} \in S(A)$.

This defines a relatively representable subfunctor $D_\Lambda^{\mathcal{D}}(\bar{\rho})$ of $D_\Lambda(\bar{\rho})$: If $D_\Lambda(\bar{\rho})$ is representable, the so is $D_\Lambda^{\mathcal{D}}(\bar{\rho})$ and the representing object $R_\Lambda^{\mathcal{D}}(\bar{\rho})$ is a quotient of $R_\Lambda(\bar{\rho})$. In addition to the conditions appearing in Section 5 below, we will be interested in the following conditions:

1. If $\Delta \subset \Gamma$ is a profinite subgroup and $\bar{\rho}(\Delta) = \{1\}$, then the assignment $S(A) := \{\rho \mid \rho(\Delta) = \{1\}\}$ defines a deformation condition. In the case $\Gamma = \mathrm{Gal}_K$ for a local field K and $\Delta = I_K$, we call this the unramified lifting condition and write $D_\Lambda^{(\square), \mathrm{nr}}(\bar{\rho})$ for the corresponding subfunctor.
2. Fix a representation $\chi : \Gamma \rightarrow G^{\mathrm{ab}}(\Lambda)$ such that $d(k) \circ \bar{\rho} = \bar{\chi}$, where $d : G \rightarrow G^{\mathrm{ab}}$ is the canonical projection modulo the derived subgroup G^{der} and where $\bar{\chi}$ denotes the reduction of χ . In accordance with the case $G = \mathrm{GL}_n$, we call this the fixed deformation condition and write $D_\Lambda^{(\square), \chi}(\bar{\rho})$ for the corresponding subfunctor.
3. Let $\Gamma = \mathrm{Gal}_F$ for a global field F and $\Sigma \subset \mathrm{Pl}_F$ a set of places and fix for each $\nu \in \Sigma$ a local condition D_ν of the functor $D_\Lambda^{(\square)}(\bar{\rho}_\nu)$, where $\bar{\rho}_\nu$ denotes the restriction of $\bar{\rho}$ to a decomposition group at ν . Then the assignment $S(A) = \{\rho \mid \rho_\nu \in D_\Lambda^{(\square), D_\nu}(\bar{\rho}_\nu) \forall \nu \in \Sigma\}$ defines a global condition, denoted by $\mathcal{D} = (D_\nu)_{\nu \in \Sigma}$. The afforded subfunctor of $D_\Lambda^{(\square)}(\bar{\rho})$ is denoted by $D_\Lambda^{(\square), \mathcal{D}}(\bar{\rho})$.
4. If Γ, F, Σ are as above and if $\bar{\rho}$ is unramified outside Σ , then requiring that a lift ρ is unramified outside Σ defines a global deformation condition, and we denote the corresponding subfunctor by $D_{\Sigma, \Lambda}^{(\square)}(\bar{\rho})$. It is easily seen that studying these lifts is equivalent to studying unconditioned lifts of $\bar{\rho}$, understood as a representation of the Galois group $\mathrm{Gal}_{F, \Sigma}$ of the maximal, unramified outside Σ extension F_Σ of F .

It is easily seen that decreasing multiple conditions defines another condition, i.e. it makes sense to write for example $D_\Lambda^{\square, \chi, \mathrm{nr}}(\bar{\rho})$.

Multiply framed deformations Continue to denote $\Gamma = \text{Gal}_F$ and fix finite subsets $\Sigma \subset S \subset \text{Pl}_F$ such that $\bar{\rho}$ is unramified outside S .

Definition 3.16. Following [KW09b, Section 4.1.1], we define the functor $D_\Lambda^{\square\Sigma}(\bar{\rho}) : \mathcal{C}_\Lambda \rightarrow \text{Sets}$ by

$$A \mapsto \left\{ (\rho, (\rho_\nu, \beta_\nu)_{\nu \in \Sigma}) \mid \begin{array}{l} \rho \in D_\Lambda^{\square}(\bar{\rho})(A), \rho_\nu \in D_\Lambda^{\square}(\bar{\rho}_\nu)(A), \beta_\nu \in \widehat{G}(A) \\ \text{s. t. } \rho|_{\text{Gal}(F_\nu)} = \beta_\nu \rho_\nu \beta_\nu^{-1} \end{array} \right\} / \sim$$

where $(\rho, (\rho_\nu, \beta_\nu)_{\nu \in \Sigma})$ and $(\rho', (\rho'_\nu, \beta'_\nu)_{\nu \in \Sigma})$ are taken to be equivalent if $\rho_\nu = \rho'_\nu$ for all ν and if there is a $\gamma \in \widehat{G}(A)$ such that $\rho' = \gamma \rho \gamma^{-1}$ and $\beta'_\nu = \gamma^{-1} \beta_\nu$ for all ν .

Note that specifying the ρ_ν is not strictly necessary, as they can be obtained from ρ and β_ν . We can impose a deformation condition $\mathcal{D} = (S(A))_{A \in \mathcal{C}_\Lambda}$ on multiply framed deformations in the same way we did for liftings and deformations, i.e. we allow only those triples $(\rho, (\rho_\nu, \beta_\nu)_{\nu \in \Sigma})$ for which $\rho \in S(A)$. The following assertions are immediate, cf. [KW09b, Proposition 4.1] or [Gui16, Proposition 2.62]:

Proposition 3.17. 1. $D_{[S], \Lambda}^{\square\Sigma, (\chi), \mathcal{D}}$ is representable and we denote the afforded deformation ring by

$$R_{[S], \Lambda}^{\square\Sigma, (\chi), \mathcal{D}} \text{ (if } \Sigma = \emptyset, \text{ we have to assume } H^0(\Gamma, \mathfrak{g}) = \mathfrak{z}\text{);}$$

2. If $\#\Sigma = 1$, then the functors $D_{[S], \Lambda}^{\square\Sigma, (\chi), \mathcal{D}}$ and $D_{[S], \Lambda}^{\square, (\chi), \mathcal{D}}$ are naturally isomorphic;

3. If $\Sigma \neq \emptyset$, then

$$R_{[S], \Lambda}^{\square\Sigma, (\chi), \mathcal{D}} \cong R_{[S], \Lambda}^{\square, (\chi), \mathcal{D}} \llbracket x_1, \dots, x_t \rrbracket \text{ and, if } H^0(\Gamma, \mathfrak{g}) = \mathfrak{z}, \text{ then also } R_{[S], \Lambda}^{\square, (\chi), \mathcal{D}} \cong R_{[S], \Lambda}^{(\chi), \mathcal{D}} \llbracket x_1, \dots, x_u \rrbracket$$

with $t = \dim(\mathfrak{g}) \cdot (\#\Sigma - 1)$, $u = \dim(\mathfrak{g}) - \dim(\mathfrak{z}) = \dim(\mathfrak{g}^{\text{der}})$.

From now on, let us suppose

Assumption 3.18. $H^0(\text{Gal}_{F, S}, \mathfrak{g}^{\text{der}}) = 0$.

With respect to a deformation condition $\mathcal{D} = (D_\nu)_{\nu \in \Sigma}$ as in example 4. above, we set

$$R_\Lambda^{\text{loc}\Sigma, (\chi), \mathcal{D}}(\bar{\rho}) := \widehat{\bigotimes}_{\nu \in \Sigma} R_\Lambda^{\square, (\chi_\nu), D_\nu}(\bar{\rho}_\nu).$$

The following is essentially a special case of [Bal12, Proposition 4.2.5] (which goes back to [Kis07, Proposition 4.1.5]):

Proposition 3.19. Assume that $D_\Lambda^{(\chi)}(\bar{\rho})$ is representable. Then

$$R_{S, \Lambda}^{\square\Sigma, (\chi), \mathcal{D}}(\bar{\rho}) \cong R_\Lambda^{\text{loc}\Sigma, (\chi), \mathcal{D}}(\bar{\rho}) \llbracket x_1, \dots, x_{a+b} \rrbracket / (f_1, \dots, f_a)$$

for suitable $a \in \mathbb{N}$, $f_i \in R_\Lambda^{\text{loc}\Sigma, (\chi), \mathcal{D}}(\bar{\rho}) \llbracket x_1, \dots, x_{a+b} \rrbracket$ and with $b = 0$ if the determinant is not fixed (resp. $b = (\#\Sigma - 1) \cdot \dim(\mathfrak{g}^{\text{ab}})$ if the determinant is fixed).

Corollary 3.20. Assume that each $R_\Lambda^{\square, (\chi_\nu), D_\nu}(\bar{\rho}_\nu)$ is a complete intersection ring of relative dimension d_ν over Λ . Assume moreover that $D_{[S], \Lambda}^{\mathcal{D}}(\bar{\rho})$ is representable and that $d := \sum_{\nu \in \Sigma} d_\nu > \dim(\mathfrak{g}) \cdot \#\Sigma - \dim(\mathfrak{z}) - b$ (with b as in Proposition 3.19). Then there exists a presentation

$$R_{[S], \Lambda}^{\mathcal{D}}(\bar{\rho}) \cong \Lambda \llbracket x_1, \dots, x_m \rrbracket / (f_1, \dots, f_m)$$

for a suitable $m \in \mathbb{N}$.

Proof. Using Proposition 3.19 and the assumption on \mathcal{D} , we can write

$$R_{S,\Lambda}^{\square_{\Sigma},(\chi),\mathcal{D}}(\bar{\rho}) \cong R_{\Lambda}^{\text{loc}_{\Sigma},(\chi),\mathcal{D}}(\bar{\rho})[[x_1, \dots, x_{a+b}]]/(f_1, \dots, f_a) \cong \Lambda[[x_1, \dots, x_{a+b+c+d}]]/(f_1, \dots, f_{a+c})$$

for a, b as above and for a suitably chosen $c \in \mathbb{N}_0$. On the other hand, by Cohen's structure theorem we can write $R_{S,\Lambda}^{(\chi),\mathcal{D}}(\bar{\rho}) \cong \Lambda[[x_1, \dots, x_u]]/(f_1, \dots, f_v)$ for suitable $u, v \in \mathbb{N}_0$ (and we assume that this is a minimal presentation, i.e. that the quantity $u - v$ is maximal among all ways to write $R_{S,\Lambda}^{(\chi),\mathcal{D}}(\bar{\rho})$ as a quotient of a power series ring), so by the third part of Proposition 3.17 we have

$$R_{S,\Lambda}^{\square_{\Sigma},(\chi),\mathcal{D}}(\bar{\rho}) \cong R_{S,\Lambda}^{(\chi),\mathcal{D}}(\bar{\rho})[[x_1, \dots, x_r]] \cong \Lambda[[x_1, \dots, x_{r+u}]]/(f_1, \dots, f_v)$$

with $r = \dim(\mathfrak{g}).\#\Sigma - \dim(\mathfrak{z})$. Comparing these two presentations, we get

$$u - v + \dim(\mathfrak{g}).\#\Sigma - \dim(\mathfrak{z}) \geq b + d \Rightarrow u - v \geq b + d - \dim(\mathfrak{g}).\#\Sigma + \dim(\mathfrak{z}).$$

Thus, the claim follows immediately from our assumption on d . \square

Tangent spaces and systems of local conditions With respect to a deformation condition \mathcal{D} will consider the tangent space $t_{D_{\Lambda}^{(\square),\mathcal{D}}}(k[\epsilon])$, which we consider as a (finite-dimensional) k -vector space (cf. [Gou01, Lecture 2]). There are canonical isomorphisms

$$t_{D_{\Lambda}^{\square},(\chi)} \cong Z^1(\Gamma, \mathfrak{g}^{\text{der}}), t_{D_{\Lambda}} \cong H^1(\Gamma, \mathfrak{g}) \text{ and } t_{D_{\Lambda}^{\chi}} \cong H^1(\Gamma, \mathfrak{g}^{\text{der}})' := \text{im}(H^1(\Gamma, \mathfrak{g}^{\text{der}}) \rightarrow H^1(\Gamma, \mathfrak{g})),$$

so via the embedding $D_{\Lambda}^{(\chi),\mathcal{D}}(k[\epsilon]) \hookrightarrow D_{\Lambda}^{(\square),\mathcal{D}}(k[\epsilon])$ we are provided with an assignment $\mathcal{D} \mapsto L(\mathcal{D})^{(\chi)} := D_{\Lambda}^{(\chi),\mathcal{D}}(k[\epsilon])$ from deformation conditions to subspaces of $H^1(\Gamma, \mathfrak{g})$ (resp. $H^1(\Gamma, \mathfrak{g}^{\text{der}})'$). In the case $\Gamma = \text{Gal}_F$ for a number field F and if $\mathcal{D} = (D_{\nu})_{\nu \in \Sigma}$, we call the afforded family $\mathcal{L}^{(\chi)} = (L(D_{\nu}))_{\nu \in \text{Pl}_F}$ of subspaces of $H^1(\text{Gal}_{F_{\nu}}, \mathfrak{g})$ (resp. of $H^1(\text{Gal}_{F_{\nu}}, \mathfrak{g}^{\text{der}})'$) a *system of local conditions*. Also note that there is an exact sequence

$$0 \rightarrow \mathfrak{g}/\mathfrak{g}^{\Gamma} \rightarrow t_{D_{\Lambda}^{\square},(\chi)} \rightarrow t_{D_{\Lambda}^{(\chi)}}$$

where, in case $\ell \gg 0$ (such that $\mathfrak{g} = \mathfrak{g}^{\text{der}} \oplus \mathfrak{g}^{\text{ab}}$), the object $\mathfrak{g}/\mathfrak{g}^{\Gamma}$ can be replaced by $\mathfrak{g}^{\text{der}}/(\mathfrak{g}^{\text{der}})^{\Gamma}$.

Liftings at infinity

Proposition 3.21. *Assume $\Gamma = \mathbb{Z}/2\mathbb{Z} = \{1, c\}$ and $\ell = \text{char}(\mathbb{F}) \neq 2$. Then*

$$R_{\Lambda}^{\square}(\bar{\rho}) \cong \Lambda[[x_1, \dots, x_m]] \text{ with } m = \dim(\mathfrak{g}^{c=-1}).$$

If ψ is a lift of the determinant, then the same result holds for $R_{\Lambda}^{\square,\psi}(\bar{\rho})$ after replacing \mathfrak{g} by $\mathfrak{g}^{\text{der}}$.

Proof. We use the general formula $H^2(\mathbb{Z}/n\mathbb{Z}, M) = M^{\mathbb{Z}/n\mathbb{Z}}/\text{im}(\varphi)$ with

$$\varphi : M \rightarrow M \quad m \mapsto \sum_{j=0}^{n-1} j.m.$$

Now, if $x \in \mathfrak{g}^{\{1,\tilde{c}\}}$, we see that $(\tilde{c}+1)(\frac{1}{2}x) = x \in \text{im}(\tilde{c}+1)$, hence $H^2(\{1, c\}, \mathfrak{g}) = 0$ and the lifting ring is unobstructed. To get the number of variables we have to evaluate

$$Z^1(\{1, c\}, \mathfrak{g}) = \{f : \{1, c\} \rightarrow \mathfrak{g} \mid f(xy) = f(x) + {}^x f(y)\}.$$

Looking at $x = y = c$, we see that f is uniquely determined by a vector $v = f(c)$. Looking at $x = 1, y = c$, we see that $f(1) = v + {}^c v = 0$, i. e. that $v \in \mathfrak{g}^{c=-1}$. On the other hand, any such v defines an $f \in Z^1$ via $1 \mapsto 0, c \mapsto v$.

The modifications of this argument for the fixed-determinant case are straight-forward. \square

A simple criterion for the vanishing of cohomology groups Now assume that $\Gamma = \text{Gal}_K$ for a local field K . Recall that, by local Tate duality, the Pontryagin dual of $H^2(\Gamma, \mathfrak{g})$ can be identified with $H^0(\Gamma, \mathfrak{g}^\vee) = (\mathfrak{g}^\vee)^\Gamma$. Together with the identification of $(\text{ad } \bar{\rho}^{(0)})^\vee$ and $(\text{ad } \bar{\rho}^{(0)})(1)$ via the trace pairing, this implies the following criterion for the vanishing of $H^2(\Gamma, \mathfrak{g}^{\text{der}})$ in the case $G = \text{GL}_n$:

Lemma 3.22 (Local case). *Let Γ be the absolute Galois group of a non-archimedean local field, k be a finite field of characteristic ℓ and*

$$\bar{\rho} : \Gamma \rightarrow \text{GL}_n(k)$$

a representation.

1. *If $\text{Hom}_\Gamma(\bar{\rho}, \bar{\rho}(1))$ vanishes, then $H^2(\Gamma, \text{ad } \bar{\rho})$ vanishes.*
2. *Assume that $\ell \nmid n$. Then, if $\text{Hom}_\Gamma(\bar{\rho}, \bar{\rho}(1))$ vanishes, also $H^2(\Gamma, \text{ad } \bar{\rho}^0)$ vanishes.*

In the global case, there is no such duality and we record the following:

Lemma 3.23 (Global case). *Let $\Gamma = \text{Gal}_{F,S}$ for a number field F and a (possibly empty) finite set of places of F . Let $k, \bar{\rho}$ be as in Lemma 3.22 above.*

1. *If $\text{Hom}_\Gamma(\bar{\rho}, \bar{\rho}(1))$ vanishes, then $H^0(\Gamma, (\text{ad } \bar{\rho})^\vee)$ vanishes.*
2. *Assume that $\ell \nmid n$. Then, if $\text{Hom}_\Gamma(\bar{\rho}, \bar{\rho}(1))$ vanishes, also $H^0(\Gamma, (\text{ad } \bar{\rho}^0)^\vee)$ vanishes.*

We easily deduce the following result, which also implies the vanishing of the error term δ in [Böc13a] (see Remark 5.2.3.(d) of loc.cit.) for large ℓ :

Corollary 3.24. *There exists a constant C , depending only on n and F , such that Assumption 3.18 holds if $\text{char}(k) > C, G = \text{GL}_n$ and $\bar{\rho}$ is irreducible.*

Unobstructedness

Definition 3.25. The functor $D_\Lambda^{(\square), [\chi]}(\bar{\rho})$ is called unobstructed if $h^2(\Gamma, \mathfrak{g}^{\text{der}}) = 0$.

Definition 3.26. A relatively representable subfunctor of $D_\Lambda^{(\square), [\chi]}(\bar{\rho})$ is called smooth (of dimension m) if its representing object is isomorphic to $\Lambda[[x_1, \dots, x_m]]$.

The most apparent application of the unobstructedness-property is that it implies the smoothness of the lifting/deformation ring, cf. [Böc07]: Assume that $D_\Lambda^{\square, (\chi)}(\bar{\rho})$ is smooth and (in the fixed-determinant case) that $\ell \gg 0$ and (in the non-framed case) that $D_\Lambda^{(\chi)}(\bar{\rho})$ is representable. Then

$$D_\Lambda^{\square, (\chi)}(\bar{\rho}) \cong \llbracket x_1, \dots, x_{a(+e)} \rrbracket \text{ and } D_\Lambda^{(\chi)}(\bar{\rho}) \cong \llbracket x_1, \dots, x_{b(+c)} \rrbracket$$

with $b = h^1(\Gamma, \mathfrak{g}), c = h^1(\Gamma, \mathfrak{g}^{\text{der}})' - b, a = b + \dim(\mathfrak{g}^{\text{der}}) - h^0(\Gamma, \mathfrak{g}^{\text{der}})$. The converse direction (i.e. that smoothness implies unobstructedness) is known not to hold (for general profinite groups Γ), cf. [Spr01].

In order to relax this notion to functors corresponding to deformation conditions, we restrict to the case $\Gamma = \text{Gal}_{F,S}$. Let $\mathcal{L}^{(x)} = (L_\nu^{(x)})_{\nu \in \text{Pl}_F}$ be a system of local conditions and $\mathcal{D}^{(x)} = (D_\nu^{(x)})_{\nu \in \text{Pl}_F}$ the corresponding global deformation condition.

Denote by $\mathfrak{g}^{(\text{der}),\vee}$ the Tate dual of $\mathfrak{g}^{\text{der}}$ and by $L_\nu^{(x),\perp}$ the annihilator of $L_\nu^{(x)}$ under the Tate pairing

$$H^i(F_\nu, \mathfrak{g}^{\text{der},\vee}) \times H^{2-i}(F_\nu, \mathfrak{g}^{\text{der}}) \longrightarrow H^2(F_\nu, k(1)) \cong \mathbb{Q}/\mathbb{Z}$$

for $i = 1$, cf. [NSW08, (7.2.6) Theorem]. Then we denote the corresponding dual Selmer group by

$$H_{\mathcal{L}^{(x),\perp}}^1(F, \mathfrak{g}^{(\text{der}),\vee}) := \ker \left(\bigoplus_{\nu \in \text{Pl}} \text{res}_\nu : H^1(F, \mathfrak{g}^{(\text{der}),\vee}) \longrightarrow \bigoplus_{\nu \in \text{Pl}} H^1(F, \mathfrak{g}^{(\text{der}),\vee}) / L_\nu^{(x),\perp} \right).$$

From now on, let us assume that $D_\nu^{(x)}$ for $\nu \notin S$ parametrizes unramified deformations.

Definition 3.27. We say that $D_{S,\Lambda}^{\mathcal{D}^{(x)}}(\bar{\rho})$ (or $D_{S,\Lambda}^{\square, \mathcal{D}^{(x)}}(\bar{\rho})$, or $D_{S,\Lambda}^{\square\Sigma, \mathcal{D}^{(x)}}(\bar{\rho})$ for some set of places Σ) has vanishing dual Selmer group if $H_{\mathcal{L}^{(x),\perp}}^1(F, \mathfrak{g}^{(\text{der}),\vee}) = 0$.

Definition 3.28. Let $\mathbf{m} = (m_\nu)_{\nu \in S} \in \mathbb{N}_0^S$. We say that $D_{S,\Lambda}^{\mathcal{D}^{(x)}}(\bar{\rho})$ (or $D_{S,\Lambda}^{\square, \mathcal{D}^{(x)}}(\bar{\rho})$, or $D_{S,\Lambda}^{\square\Sigma, \mathcal{D}^{(x)}}(\bar{\rho})$) is globally unobstructed (of local dimensions \mathbf{m}) if its dual Selmer group vanishes and if each $D_\Lambda^{\square, \mathcal{D}^{(x\nu)}}(\bar{\rho}_\nu)$ for $\nu \in S$ is smooth (of dimension m_ν).

We remark that if $D_{S,\Lambda}^{\mathcal{D}^{(x)}}(\bar{\rho})$ is globally unobstructed and representable, then by [Böc07, Theorem 5.2] the representing object $R_{S,\Lambda}^{\mathcal{D}^{(x)}}(\bar{\rho})$ is isomorphic to a power series ring in $h_{\mathcal{L}^{(x)}}^1(F, \mathfrak{g}^{(\text{der})})^{(l)}$ variables. The following results directly from the exact sequece of [Böc07, p. 7],

$$0 \rightarrow \text{III}_S^2(\mathfrak{g}^{\text{der}}) \rightarrow H^2(\text{Gal}_{F,S}, \mathfrak{g}^{\text{der}}) \rightarrow \bigoplus_{\nu \in S} H^2(F_\nu, \mathfrak{g}^{\text{der}}) \rightarrow H^0(F, \mathfrak{g}^{(\text{der}),\vee})^* \rightarrow 0,$$

where $H^0(F, \mathfrak{g}^{(\text{der}),\vee})^*$ vanishes for $\ell \gg 0$:

Proposition 3.29. *Assume that $D_\Lambda^{(x\nu)}(\bar{\rho}_\nu)$ is unobstructed (for all $\nu \in S$) and that $D_{S,\Lambda}^{(x)}(\bar{\rho})$ is globally unobstructed (without making an assumption on the dimension). Then $D_{S,\Lambda}^{(x)}(\bar{\rho})$ is unobstructed in the sense of Definition 3.25. For $\ell \gg 0$, also the converse is true.*

4 A general framework for unobstructedness

For this section, we take the following static point of view: Let k be a finite field with ring of Witt vectors $W = W(k)$, let S be a finite set of finite places of F . We assume $\ell := \text{char}(k) \notin S \cup \{2\}$. Then we fix a continuous representation

$$\bar{\rho} : \text{Gal}_{F,S} \rightarrow G(k)$$

together with a lift $\chi : \text{Gal}_{F,S} \rightarrow G^{\text{ab}}(k)$ of the determinant. Let us moreover fix a Borel subgroup $B \subset G$ and denote by $\mathfrak{g}^{\text{der}}$ (resp. $\mathfrak{b}^{\text{der}}$) the Lie algebra of the derived subgroup G^{der} (resp. the Lie algebra of $B \cap G^{\text{der}}$).

With respect to some choice² of local deformation conditions

- **min** of the restriction $\bar{\rho}_\nu$ of $\bar{\rho}$ to a decomposition group at $\nu \in S$,
- **sm** and **crys** of the restriction $\bar{\rho}_\nu$ of $\bar{\rho}$ to a decomposition group at $\nu|\ell$,

consider the following list of assumptions, where we leave out the W in the subscript of the occurring deformation functors and rings:

1. **(sm/k)**: For each $\nu|\ell$, the subfunctor $D^{\square, \chi_\nu, \text{sm}}(\bar{\rho}_\nu)$ of $D^{\square, \chi_\nu}(\bar{\rho}_\nu)$ is representable by a formally smooth (over W) object $R_\nu^{\square, \chi_\nu, \text{sm}}$ (and we denote the dimension by $d_\nu^{\square, \text{sm}}$).
2. **(crys)**: For each $\nu|\ell$, the subfunctor $D^{\square, \chi_\nu, \text{crys}}(\bar{\rho}_\nu)$ of $D^{\square, \chi_\nu}(\bar{\rho}_\nu)$ is representable by a formally smooth (over W) object $R_\nu^{\square, \chi_\nu, \text{crys}}$ of relative dimension

$$d_\nu^{\square, \text{crys}} = \dim(\mathfrak{g}^{\text{der}}) + (\dim(\mathfrak{g}^{\text{der}}) - \dim(\mathfrak{b}^{\text{der}}))[F_\nu : \mathbb{Q}_\ell].$$

3. **(min)**: For each $\nu \in S$, the subfunctor $D^{\square, \chi_\nu, \text{min}}(\bar{\rho}_\nu)$ of $D^{\square, \chi_\nu}(\bar{\rho}_\nu)$ is representable by a formally smooth (over W) object $R_\nu^{\square, \chi_\nu, \text{min}}$ of relative dimension

$$d_\nu^{\square, \text{min}} = \dim(\mathfrak{g}^{\text{der}}).$$

4. **(∞)**: For each $\nu|\infty$, the functor $D^{\square, \chi_\nu}(\bar{\rho}_\nu)$ is representable by an object (over W) of relative dimension $d_\nu^{\square} = \dim(\mathfrak{b}^{\text{der}})$. (As $\ell > 2 = \#\text{Gal}_{F_\nu}$, the strict ℓ -cohomological dimension $\text{scd}_\ell(\text{Gal}_{F_\nu})$ is zero, i.e. the representing object is automatically formally smooth over W .)

5. **(Presentability)**: There exists a presentation

$$R_{S_\ell}^{\square, \chi, \text{min}, \text{sm}} \cong R_{S_\ell}^{\text{loc}, \text{min}, \text{sm}}[[x_1, \dots, x_a]]/(f_1, \dots, f_b)$$

for integers a, b fulfilling $a - b = (\#S_\ell - 1) \cdot \dim(\mathfrak{g}^{\text{ab}})$. In this equation, we take

$$R_{S_\ell}^{\text{loc}, \text{min}, \text{sm}} = \widehat{\bigotimes_{\nu \in S_\ell} \tilde{R}_\nu} \text{ with } \tilde{R}_\nu = \begin{cases} R_\nu^{\square, \chi_\nu, \text{min}} & \text{if } \nu \in S; \\ R_\nu^{\square, \chi_\nu, \text{sm}} & \text{if } \nu|\ell; \\ R_\nu^{\square, \chi_\nu} & \text{if } \nu|\infty. \end{cases} \quad (3)$$

6. **($R = T$)**: The ring $R_{S_\ell}^{\square, \chi, \text{min}, \text{crys}}$ is formally smooth of relative dimension

$$r_0 := \dim(\mathfrak{g}) \cdot \#S_\ell - \dim(\mathfrak{g}^{\text{ab}}).$$

Remark 4.1 (Taylor-Wiles condition). Let $\nu|\infty$ so that $\text{scd}_\ell(\text{Gal}_{F_\nu}) = 0$, then it follows from condition (∞) , $\text{scd}_\ell(\text{Gal}_{F_\nu}) = 0$ and the remark following Definition 3.25 that

$$\dim(\mathfrak{b}^{\text{der}}) = \dim_W(R^\square) = h^1(\text{Gal}_{F_\nu}, \mathfrak{g}^{\text{der}})' + \dim(\mathfrak{g}^{\text{der}}) - h^0(\text{Gal}_{F_\nu}, \mathfrak{g}^{\text{der}}) = \dim(\mathfrak{g}^{\text{der}}) - h^0(\text{Gal}_{F_\nu}, \mathfrak{g}^{\text{der}}).$$

This implies

$$\sum_{\nu|\infty} h^0(\text{Gal}_{F_\nu}, \mathfrak{g}^{\text{der}}) = [F : \mathbb{Q}] \cdot (\mathfrak{g}^{\text{der}} - \dim(\mathfrak{b}^{\text{der}})). \quad (4)$$

²During the following applications of the presented material, we will consider for **min** the condition of Section 5.4, for **crys** the condition of Section 5.3 and for **sm** the unconditioned deformation condition. We stress, however, that for the purpose of this section we treat **min**, **crys**, **sm** purely formally as deformation conditions satisfying the listed assumptions of Definition 3.8.

We can now state the main result of this section:

Theorem 4.2. *Suppose conditions 1.-6. are met and, for $\nu|\ell$, write $d_\nu^{\square, \text{sm}} = \dim(\mathfrak{g}^{\text{der}}) \cdot ([F_\nu : \mathbb{Q}_\ell] + 1) - \delta_\nu$ for suitable numbers $\delta_\nu \in \mathbb{N}_0$.*

1. *The ring $R_{S_\ell}^{\square_{S_\ell}, \chi, \text{min}, \text{sm}}$ is formally smooth of relative dimension*

$$\#S_\ell \cdot \dim(\mathfrak{g}) - \dim(\mathfrak{g}^{\text{ab}}) + [F : \mathbb{Q}] \cdot \dim(\mathfrak{b}^{\text{der}}) - \sum_{\nu|\ell} \delta_\nu.$$

If the unframed deformation functor $D_{S_\ell}^{\chi, \text{min}, \text{sm}}$ is representable, then $R_{S_\ell}^{\chi, \text{min}, \text{sm}}$ is formally smooth of relative dimension $[F : \mathbb{Q}] \cdot \dim(\mathfrak{b}^{\text{der}}) - \sum_{\nu|\ell} \delta_\nu$.

2. *Let $\mathcal{L} := (L_\nu^\chi)_\nu$ be the system of local conditions corresponding to the deformation functor $D_{S_\ell}^{\chi, \text{min}, \text{sm}}(\bar{\rho})$. Assume*

- a) $\mathfrak{g} = \mathfrak{g}^{\text{der}} \oplus \mathfrak{g}^{\text{ab}}$ (e.g. because $\ell \gg 0$);
- b) $H^0(\text{Gal}_F, \mathfrak{g}^{\text{der}, \vee}) = 0$;
- c) For $\nu \in S$, we have $\dim(L_\nu) = h^0(\text{Gal}_{F_\nu}, \mathfrak{g}^{\text{der}})$;
- d) All δ_ν vanish.

Then $H_{\mathcal{L}^\perp}^1(\text{Gal}_{F,S}, \mathfrak{g}^{\text{der}, \vee}) = H^0(\text{Gal}_{F,S}, \mathfrak{g}^{\text{der}}) = 0$.

Remark 4.3. 1. As the deformation conditions **sm** and **crys** are relatively representable (cf. conditions 1. and 2.), $D_{S_\ell}^{\chi, \text{min}, \text{sm}}$ is representable if $D_{S_\ell}^\chi$ is representable. For example, this is the case if $\bar{\rho}$ is absolutely irreducible (in the sense of Definition 3.13).

2. For $\nu \notin S_\ell$, the equality $\dim(L_\nu) = h^0(\text{Gal}_{F_\nu}, \mathfrak{g}^{\text{der}})$ holds automatically if $\ell \gg 0$ (so that $\mathfrak{g} = \mathfrak{g}^{\text{der}} \oplus \mathfrak{g}^{\text{ab}}$).

Proof. First remark that the second claim of part 1. follows directly from Lemma 2.15, as $R_{S_\ell}^{\square_{S_\ell}, \chi, \text{min}, \text{sm}}$ is a power series ring over $R_{S_\ell}^{\chi, \text{min}, \text{sm}}$, and from the formula $\dim \mathfrak{g} = \dim \mathfrak{g}^{\text{der}} + \dim \mathfrak{g}^{\text{ab}}$.

For the first sentence of 1., we use the shorthand notation $d_T^* = \sum_{\nu \in T} d_\nu^*$ for a subset T of Pl_F . Moreover, we write d_∞^\square for $d_{\Omega_\infty}^\square$ and d_ℓ^* for $d_{\Omega_\ell}^*$. Let us consider the commutative diagram

$$\begin{array}{ccccccc} 0 & \longrightarrow & I & \longrightarrow & R_{S_\ell}^{\text{loc}, \text{min}, \text{sm}} & \xrightarrow{f} & R_{S_\ell}^{\text{loc}, \text{min}, \text{crys}} \longrightarrow 0 \\ & & \downarrow \pi & & \downarrow \pi & & \downarrow \pi' \\ 0 & \longrightarrow & J & \longrightarrow & R_{S_\ell}^{\square_{S_\ell}, \chi, \text{min}, \text{sm}} & \xrightarrow{g} & R_{S_\ell}^{\square_{S_\ell}, \chi, \text{min}, \text{crys}} \longrightarrow 0. \end{array}$$

In this diagram, the right square is a pushout square, $R_{S_\ell}^{\text{loc}, \text{min}, \text{crys}}$ is defined as in (3) (but with $\tilde{R}_\nu = R_\nu^{\square, \chi, \text{crys}}$ for $\nu|\ell$) and f, g are the canonical projections. Moreover, $\pi = \otimes_{\nu \in S_\ell} \pi_\nu$ is induced from the natural transformations

$$D_{S_\ell}^{\square_{S_\ell}, \chi, \text{min}, \text{crys}} \rightarrow \tilde{D}_\nu,$$

where \tilde{D}_ν is the deformation functor corresponding to (i.e. represented by) the ring \tilde{R}_ν in (3) and, analogously, $\pi' = \otimes_{\nu \in S_\ell} \pi'_\nu$ is defined with **crys** in place of **sm**.

Using the list of assumptions, we can rewrite the above diagram as

$$\begin{array}{ccccccc}
0 & \longrightarrow & I & \longrightarrow & W[[x_1, \dots, x_{d_\ell^{\square, \text{sm}} + d_\infty^{\square} + d_S^{\square, \text{min}}}], \dots, x_{d_\ell^{\square, \text{crys}} + d_\infty^{\square} + d_S^{\square, \text{min}}}] & \xrightarrow{f} & W[[x_1, \dots, x_{d_\ell^{\square, \text{crys}} + d_\infty^{\square} + d_S^{\square, \text{min}}}], \dots, x_{d_\ell^{\square, \text{crys}} + d_\infty^{\square} + d_S^{\square, \text{min}}}] & \longrightarrow & 0 \\
& & \downarrow \pi & & \downarrow \pi & & \downarrow \pi' & & \\
0 & \longrightarrow & J & \longrightarrow & W[[x_1, \dots, x_m]]/(f_1, \dots, f_{m-\gamma}) & \xrightarrow{g} & W[[x_1, \dots, x_{r_0}]] & \longrightarrow & 0
\end{array}$$

with $\gamma = (\#S_\ell - 1) \cdot \dim(\mathfrak{g}^{\text{ab}}) + d_\ell^{\square, \text{sm}} + d_\infty^{\square} + d_S^{\square, \text{min}}$. It is easily seen that $R_{S_\ell}^{\square, \chi, \text{min}, \text{sm}}$ is formally smooth if we can show $\text{gen}(J) \leq m - (m - \gamma) - r_0 = \gamma - r_0$. From the pushout property of the diagram, we can easily deduce that $\text{gen}(J) \leq \text{gen}(I)$. As f is a surjection of regular rings, it follows from (Serre, Local Algebra, Proposition 22) that $\text{gen}(I) = d_\ell^{\square, \text{sm}} - d_\ell^{\square, \text{crys}}$. Thus, we are left to show the inequality

$$\begin{aligned}
d_\ell^{\square, \text{sm}} - d_\ell^{\square, \text{crys}} &\leq \gamma - r_0 = (\#S_\ell - 1) \cdot \dim(\mathfrak{g}^{\text{ab}}) + d_\ell^{\square, \text{sm}} + d_\infty^{\square} + d_S^{\square, \text{min}} - \dim(\mathfrak{g}) \cdot \#S_\ell + \dim(\mathfrak{g}^{\text{ab}}) \\
&= \#S_\ell \cdot (\dim(\mathfrak{g}^{\text{ab}}) - \dim(\mathfrak{g})) + d_\ell^{\square, \text{sm}} + d_\infty^{\square} + d_S^{\square, \text{min}}.
\end{aligned}$$

By assumptions **(min)** and **(∞)** and by the identity $\dim(\mathfrak{g}^{\text{der}}) + \dim(\mathfrak{g}^{\text{ab}}) = \dim(\mathfrak{g})$, this amounts to

$$d_\ell^{\square, \text{crys}} \geq \dim(\mathfrak{g}^{\text{der}}) \cdot (\#\Omega_\ell + [F : \mathbb{Q}]) - \dim(\mathfrak{b}^{\text{der}})[F : \mathbb{Q}].$$

Assumption **(crys)** amounts precisely to the fact that this inequality is fulfilled (with equality), which implies the formal smoothness of $R_{S_\ell}^{\square, \chi, \text{min}, \text{sm}}$. Moreover, we easily check that the relative dimension of $R_{S_\ell}^{\square, \chi, \text{min}, \text{sm}}$ is

$$\begin{aligned}
&\gamma = (\#S_\ell - 1) \cdot \dim(\mathfrak{g}^{\text{ab}}) + d_\ell^{\square, \text{sm}} + d_\infty^{\square} + d_S^{\square, \text{min}} \\
&= \#S_\ell \cdot \dim \mathfrak{g}^{\text{ab}} - \dim \mathfrak{g}^{\text{ab}} + \dim \mathfrak{g}^{\text{der}} \cdot ([F : \mathbb{Q}] + \#\Omega_\ell) - \left(\sum_{\nu|\ell} \delta_\nu \right) + [F : \mathbb{Q}] \cdot \dim(\mathfrak{b}^{\text{der}}) + \#S \cdot \dim(\mathfrak{g}^{\text{der}}) \\
&= \#S_\ell \cdot \dim(\mathfrak{g}) + [F : \mathbb{Q}] \cdot \dim(\mathfrak{b}^{\text{der}}) - \dim(\mathfrak{g}^{\text{ab}}) - \sum_{\nu|\ell} \delta_\nu.
\end{aligned}$$

Concerning part 2., note that (using condition a)) we have an exact sequence

$$0 \rightarrow \mathfrak{g}/\mathfrak{g}^{\text{Gal}_{F_\nu}} = \mathfrak{g}^{\text{der}}/(\mathfrak{g}^{\text{der}})^{\text{Gal}_{F_\nu}} \rightarrow t_{D_W^{\square, \chi, \text{sm}}(\bar{\rho}_\nu)} \rightarrow t_{D_W^{\chi, \text{sm}}(\bar{\rho}_\nu)} \rightarrow 0$$

for $\nu|\ell$. Therefore, using condition 2. and the vanishing of δ_ν , we have for $\nu|\ell$ the following:

$$\dim(L_\nu) = \dim t_{D_W^{\chi, \text{sm}}(\bar{\rho}_\nu)} = h^0(\text{Gal}_{F_\nu}, \mathfrak{g}^{\text{der}}) + [F_\nu : \mathbb{Q}] \cdot \dim(\mathfrak{g}^{\text{der}}).$$

Recall the Greenberg-Wiles-Formula [NSW08, Theorem 8.7.9]:

$$\begin{aligned}
&\dim H_{\mathcal{L}}^1(\text{Gal}_{F,S}, \mathfrak{g}^{\text{der}}) - \dim H_{\mathcal{L}^\perp}^1(\text{Gal}_{F,S}, \mathfrak{g}^{\text{der}, \vee}) \\
&= h^0(\text{Gal}_{F,S}, \mathfrak{g}^{\text{der}}) - h^0(\text{Gal}_{F,S}, \mathfrak{g}^{\text{der}, \vee}) + \sum_{\nu \in S_\ell} (\dim(L_\nu) - h^0(\text{Gal}_{F_\nu}, \mathfrak{g}^{\text{der}}))
\end{aligned}$$

By [Böc07, Section 5], we know that $H_{\mathcal{L}}^1(\text{Gal}_{F,S}, \mathfrak{g}^{\text{der}})$ can be identified with the tangent space of the functor $D_{S_\ell}^{\chi, \text{min}, \text{sm}}$ and hence (by part 2.) equals $[F : \mathbb{Q}] \cdot \dim(\mathfrak{b}^{\text{der}})$. For $\nu|\infty$, we have $L_\nu \subset$

$H^1(\text{Gal}_{F,S}, \mathfrak{g}^{\text{der}}) = 0$. Thus, using the Taylor-Wiles formula (4) and assumption b), the sum evaluates to

$$\sum_{\nu \in S_\ell} (\dim(L_\nu) - h^0(\text{Gal}_{F_\nu}, \mathfrak{g}^{\text{der}})) = [F : \mathbb{Q}] \cdot \dim(\mathfrak{g}^{\text{der}}) - [F : \mathbb{Q}] \cdot (\dim(\mathfrak{g}^{\text{der}}) - \dim(\mathfrak{b}^{\text{der}})).$$

Therefore we get

$$- \dim H_{\mathcal{L}^\perp}^1(\text{Gal}_{F,S}, \mathfrak{g}^{\text{der}, \vee}) = h^0(\text{Gal}_{F,S}, \mathfrak{g}^{\text{der}}).$$

As neither quantity can be negative, they must both vanish and the result follows. \square

From the exact sequence

$$H_{\mathcal{L}^\perp}^1(\text{Gal}_{F,S}, \mathfrak{g}^{\text{der}, \vee})^* \rightarrow \text{III}_{S_\ell}^2(\mathfrak{g}^{\text{der}}) \rightarrow 0$$

(see e.g. equation (9) on p. 10 of [Boeckle07]) we can deduce:

Corollary 4.4. *Under the assumptions of part 2. of Theorem 4.2, $\text{III}_{S_\ell}^2(\mathfrak{g}^{\text{der}})$ vanishes. In particular, under these hypotheses the unrestricted deformation functor $D_{S_\ell}^{(\square_{S_\ell}), \chi}(\bar{\rho})$ is globally unobstructed precisely if the local deformation functors $D^{(\square), \chi_\nu}(\bar{\rho}_\nu)$ are relatively smooth for $\nu \in S \cup \Omega_\ell$.*

We remark that $D^{(\square), \chi_\nu}(\bar{\rho}_\nu)$ is relatively smooth for $\nu \in \Omega_\infty$ by Proposition 3.21, so Corollary 4.4 holds true with „... for $\nu \in S_\ell$ “ in place of „... for $\nu \in S \cup \Omega_\ell$ “.

Potential unobstructedness We start with the following, easy observation:

Proposition 4.5. *Let K be a local field and let K' be a finite extension of K such that ℓ does not divide the index $[K' : K]$. Let $\bar{\rho}$ be a G -valued residual representation of Gal_K and fix a lift χ of the determinant. Then unobstructedness of $D_\Lambda^{(\chi)}(\bar{\rho} | \text{Gal}_{K'})$ implies unobstructedness of $D_\Lambda^{(\chi)}(\bar{\rho})$.*

Proof. This follows immediately from the injectivity of

$$\text{res}_{K'|K} : H^2(K, \mathfrak{g}^{(\text{der})}) \rightarrow H^2(K', \mathfrak{g}^{(\text{der})}),$$

cf. [NSW08, Corollary (1.5.7)]. \square

This proof is not directly applicable to the global situation, as we have to keep track of the set of places at which we allow ramification. Therefore, we first describe a more flexible method which can also handle conditioned deformation functors:

Definition 4.6 (Dual-pre condition). Let $F'|F$ be a finite extension of number fields.

1. Let $\nu' \in \text{Pl}_{F'}, \nu \in \text{Pl}_F$ such that $\nu' | \nu$. Moreover, let $L_\nu \subset H^1(F_\nu, \mathfrak{g}^{(\text{der})}), L'_{\nu'} \subset H^1(F'_{\nu'}, \mathfrak{g}^{(\text{der})})$ be local conditions. We say that L_ν is a dual-pre- $L'_{\nu'}$ condition if $\text{res}_{\nu'}^\vee(L_\nu^\perp) \subset L'_{\nu'}^\perp$, where

$$\text{res}_{\nu'}^\vee : H^1(F_\nu, \mathfrak{g}^{(\text{der}), \vee}) \rightarrow H^1(F'_{\nu'}, \mathfrak{g}^{(\text{der}), \vee})$$

denotes the usual restriction map.

2. Let $\mathcal{L}' = (L'_{\nu'})_{\nu' \in \text{Pl}_{F'}}$ be a system of local conditions for F' . We say that a system $\mathcal{L} = (L_\nu)_{\nu \in \text{Pl}_F}$ of local conditions for F is dual-pre- \mathcal{L}' if for each pair ν, ν' as above, L_ν is a dual-pre- $L'_{\nu'}$ condition.

Example 4.7. Let F, F' be as in Definition 4.6 and fix a finite set $S \subset \text{Pl}_F$ such that $\bar{\rho}$ is unramified outside S . Take for \mathcal{L} the local system parametrizing all deformations which are unramified outside S , i.e. $L_\nu = H^1(F_\nu, \mathfrak{g}^{(\text{der})})$ if $\nu \in S$ and $L_\nu = H^1(\text{Gal}_{F_\nu}/I_{F_\nu}, \mathfrak{g}^{(\text{der})})$ otherwise. Analogously, let \mathcal{L}' the local system parametrizing all deformations which are unramified outside $S\langle F' \rangle$. Then any lift of $\bar{\rho}$ which is unramified outside S is, after restriction to $\text{Gal}_{F'}$, a lift of $\bar{\rho}|_{\text{Gal}_{F'}}$ which is unramified outside $S\langle F' \rangle$. But this implies easily that the restriction map $\text{res}_{\nu'} : H^1(F_\nu, \mathfrak{g}^{(\text{der})}) \rightarrow H^1(F'_{\nu'}, \mathfrak{g}^{(\text{der})})$ maps L_ν into $L'_{\nu'}$ for any pair of places ν, ν' with $\nu'|\nu$. Using the fact that Tate duality is given by the cup product, we see that \mathcal{L} is dual-pre- \mathcal{L}' .

Lemma 4.8. Let $\bar{\rho}, F$ and F' be as above and assume $(\ell, [F' : F]) = 1$. Let $\mathcal{L} = (L_\nu)_{\nu \in \text{Pl}_F}, \mathcal{L}' = (L'_{\nu'})_{\nu' \in \text{Pl}_{F'}}$ be systems of local conditions (with associated deformation conditions \mathcal{D} and \mathcal{D}') such that \mathcal{L} is dual-pre- \mathcal{L}' . Moreover, assume that $D^{[\square], (\chi), \mathcal{D}'}(\bar{\rho}|_{\text{Gal}_{F'}})$ has vanishing dual Selmer group. Then also $D^{[\square], (\chi), \mathcal{D}}(\bar{\rho})$ has vanishing dual Selmer group.

Proof. As above, the invertibility of $[F' : F]$ implies that the restriction map

$$H^1(\text{Gal}_F, \mathfrak{g}^{(\text{der}), \vee}) \rightarrow H^1(\text{Gal}_{F'}, \mathfrak{g}^{(\text{der}), \vee})$$

is injective. Consider the diagram

$$\begin{array}{ccccc} H^1_{\mathcal{L}^\perp}(F, \mathfrak{g}^{(\text{der}), \vee}) & \hookrightarrow & H^1(F, \mathfrak{g}^{(\text{der}), \vee}) & \longrightarrow & \bigoplus_{\nu \in \text{Pl}_F} H^1(F_\nu, \mathfrak{g}^{(\text{der}), \vee})/L_\nu^\perp \\ \downarrow \varphi & & \downarrow & & \downarrow \\ H^1_{\mathcal{L}'^\perp}(F', \mathfrak{g}^{(\text{der}), \vee}) & \hookrightarrow & H^1(F', \mathfrak{g}^{(\text{der}), \vee}) & \longrightarrow & \bigoplus_{\nu' \in \text{Pl}_{F'}} H^1(F'_{\nu'}, \mathfrak{g}^{(\text{der}), \vee})/L'_{\nu'}{}^\perp \end{array}$$

The vertical map on the right is defined because \mathcal{L} is dual-pre- \mathcal{L}' , and this implies the well-definedness of φ . A simple diagram chase implies injectivity of φ , from which the claim follows. \square

The following follows now directly from Example 4.7 and Lemma 4.8:

Corollary 4.9. Let F be a number field and let F' be a finite extension of F such that ℓ does not divide the index $[F' : F]$. Let $\bar{\rho}$ be a G -valued residual representation of Gal_F which is unramified outside a finite set of places S and fix a lift χ of the determinant. Then unobstructedness of $D_{\Lambda, S\langle F' \rangle}^{(\chi|_{\text{Gal}_{F'}})}(\bar{\rho}|_{\text{Gal}_{F'}})$ implies unobstructedness of $D_\Lambda^{(\chi)}(\bar{\rho})$.

5 Local deformation conditions for $G = \text{GL}_n$

Let K be a finite extension of \mathbb{Q}_p and let k be a finite field of characteristic ℓ . In the following, we consider deformation conditions for a continuous representation $\bar{\rho} : \text{Gal}_K \rightarrow \text{GL}_n(k)$.

5.1 Unrestricted deformations ($p \neq \ell$)

In the case $p \neq \ell$, we have the following result:

Theorem 5.1. $\text{Spec } R^\square(\bar{\rho})$ is a reduced complete intersection, flat and equidimensional of relative dimension n^2 over $\text{Spec } W$.

Proof. This is Theorem 2.5 in [Sho16]. \square

5.2 Unrestricted deformations ($p = \ell$)

For the remainder of this subsection, we assume that $\bar{\rho}$ is the semi-simplification of the reduction of a crystalline representation

$$\rho : \text{Gal}_K \rightarrow \text{GL}_n(L)$$

for a suitable finite extension L of \mathbb{Q}_p with residue field k and for $p = \ell$. Denote the set of embeddings $\tau : K \hookrightarrow \overline{\mathbb{Q}_p}$ by \mathbb{E}_K and for $\tau \in \mathbb{E}_K$ denote by $\text{HT}_\tau(\rho)$ the multiset of Hodge-Tate weights of ρ with respect to τ .

Theorem 5.2. *Assume that $K|\mathbb{Q}_p$ is unramified and that for each $\tau \in \mathbb{E}_K$*

1. *there exists an $\alpha \in \mathbb{Z}$ such that all Hodge-Tate weights in $\text{HT}_\tau(\rho)$ lie in the range $[\alpha, \alpha + \ell - 3]$;*
2. *the Hodge-Tate weights of ρ are non-consecutive, i.e. if two numbers $a, b \in \mathbb{Z}$ occur in $\text{HT}_\tau(\rho)$, then $|a - b| \neq 1$.*

Then $R^\square(\bar{\rho}) \cong W[[x_1, \dots, x_m]]$ with $m = n^2 \cdot ([K : \mathbb{Q}_\ell] + 1)$.

Before we come to the proof, recall the theory of Fontaine-Laffaille [FL82], as normalized in [CHT08] (see also [BLGGT14, Section 1.4]): We consider the category $\underline{\text{FL}}_{\mathcal{O}_K, \mathcal{O}_L}$, consisting of $\mathcal{O}_K \otimes_{\mathbb{Z}_\ell} \mathcal{O}_L$ -modules M , endowed with a decreasing filtration $(\text{Fil}^i M)_{i \in \mathbb{Z}}$ with $\text{Fil}^0 M = M$ and $\text{Fil}^{\ell-1} M = 0$ and a family of Frobenius-linear maps $\text{Fil}^i M \rightarrow M$ such that $\varphi^i | \text{Fil}^{i+1} = \ell \cdot \varphi^{i+1}$ and $\sum_i \varphi^i(\text{Fil}^i M) = M$. Let $\underline{\text{FL}}_{\mathcal{O}_K, k}$ denote the full subcategory of finite length objects which are annihilated by the maximal ideal $\varpi_L \cdot \mathcal{O}_L$. We need the following well-known facts:

- There exists an exact, fully-faithful, covariant and \mathcal{O}_L -linear functor

$$\mathbf{G}_K : \underline{\text{FL}}_{\mathcal{O}_K, \mathcal{O}_L} \rightarrow \text{Rep}_{\mathcal{O}_L}(\text{Gal}_K).$$

The essential image is closed under taking subobjects and quotients. Moreover, \mathbf{G}_K restricts to a functor

$$\underline{\text{FL}}_{\mathcal{O}_K, k} \rightarrow \text{Rep}_k(\text{Gal}_K).$$

- For $\mathbf{M} \in \underline{\text{FL}}_{\mathcal{O}_K, \mathcal{O}_L}$ projective over \mathcal{O}_L , we have

$$\text{HT}_\tau(\mathbf{G}_K(\mathbf{M} \otimes_{\mathbb{Z}_p} \mathbb{Q}_p)) = \text{FL}_\tau(\mathbf{M} \otimes_{\mathcal{O}_L} k),$$

where for $\mathbf{N} \in \underline{\text{FL}}_{\mathcal{O}_K, k}$ we denote by $\text{FL}_\tau(\mathbf{N})$ the multiset of integers i , such that

$$\text{gr}^i(\mathbf{N}^\tau) = \text{Fil}^i N \otimes_{\mathcal{O}_K \otimes_{\mathbb{Z}_p} \mathcal{O}_L, \tau \otimes 1} \mathcal{O}_L / \text{Fil}^{i+1} N \otimes_{\mathcal{O}_K \otimes_{\mathbb{Z}_p} \mathcal{O}_L, \tau \otimes 1} \mathcal{O}_L$$

does not vanish, where i is counted with multiplicity $\dim_k \text{gr}^i(\mathbf{N}^\tau)$.

- Assuming condition 1. of Theorem 5.2, any Gal_K -stable \mathcal{O}_L -lattice of ρ is in the image of \mathbf{G}_K , and so is its reduction $\Lambda / \varpi_L \cdot \Lambda$.
- Morphisms in $\underline{\text{FL}}_{\mathcal{O}_K, k}$ are strict with filtrations. If $f : \mathbf{M} \rightarrow \mathbf{N}$ is such a morphism, then $f(\text{Fil}^i M) = f(M) \cap \text{Fil}^i N$ for all $i \in \mathbb{Z}$. In particular, if $\mathbf{M}, \mathbf{N} \in \underline{\text{FL}}_{\mathcal{O}_K, k}$ fulfill

$$\text{FL}_\tau(\mathbf{M}) \cap \text{FL}_\tau(\mathbf{N}) \tag{5}$$

for all $\tau \in \mathbb{E}_K$, then $\text{Hom}_{\underline{\text{FL}}_{\mathcal{O}_K, k}}(\mathbf{M}, \mathbf{N}) = 0$.

Proof. As $h^2(K, \text{ad } \bar{\rho})$ is an upper bound on the number of generators of the kernel of a surjection $W[[x_1, \dots, x_s]] \twoheadrightarrow R^\square(\bar{\rho})$ with $s = \dim Z^1(K, \text{ad } \bar{\rho})$ (cf. [All16, Proposition 2.1.2]), we have to prove

$$H^2(K, \text{ad } \bar{\rho}) = 0. \quad (6)$$

Moreover, using the exact sequence

$$0 \rightarrow \text{ad } \bar{\rho} / (\text{ad } \bar{\rho})^{\text{Gal}_K} \rightarrow t_{D_W^\square(\bar{\rho})} \rightarrow t_{D_W(\bar{\rho})} \rightarrow 0$$

and the local Euler-Poincare formula, we can compute

$$s = h^1(K, \text{ad } \bar{\rho}) + n^2 - h^0(K, \text{ad } \bar{\rho}) = n^2 - \chi(K, \text{ad } \bar{\rho}) = n^2([K : \mathbb{Q}_p] + 1).$$

Thus, (6) implies the claim.

As the trace pairing identifies $\text{ad } \bar{\rho}^\vee$ and $\text{ad } \bar{\rho}(1)$, we are finished if we can show that

$$H^2(K, \text{ad } \bar{\rho})^* \cong H^0(K, \text{ad } \bar{\rho}^\vee) \cong H^0(K, \text{ad } \bar{\rho}(1)) \cong \text{Hom}_{\text{Gal}_K}(\bar{\rho}, \bar{\rho}(1)) = 0.$$

Because $\text{Hom}_{\text{Gal}_K}(\bar{\rho}, \bar{\rho}(1)) \cong \text{Hom}_{\text{Gal}_K}(\bar{\rho}(1 - \alpha), \bar{\rho}(2 - \alpha))$, we can assume without loss of generality that $\alpha = 1$.

It is easy to see that we can choose a Gal_K -stable \mathcal{O}_L -lattice Λ of ρ such that its reduction is semi-simple, i.e. $\Lambda / \varpi_L \Lambda \cong \bar{\rho}$ (if necessary, after replacing ρ by a base change $\rho \otimes_L L'$ to a sufficiently ramified finite extension L' of L , which does not affect the validity of (6)). By our first assumption that all weights of ρ lie in the range $[1, \ell - 2]$, it thus follows that $\bar{\rho}$ is of the form $\mathbf{G}_K(\mathbf{M})$ for a suitable $\mathbf{M} \in \underline{\text{FL}}_{\mathcal{O}_K, k}$. By the same argument, $\bar{\rho}(1) = \mathbf{G}_K(\mathbf{N})$ for a suitable $\mathbf{N} \in \underline{\text{FL}}_{\mathcal{O}_K, k}$. As the cyclotomic character shifts the weights by -1 , the second condition translates precisely into the condition (5). Thus, using that \mathbf{G}_K is fully faithful, we get

$$0 = \text{Hom}_{\underline{\text{FL}}_{\mathcal{O}_K, k}}(\mathbf{M}, \mathbf{N}) \cong \text{Hom}_{\text{Gal}_K}(\bar{\rho}, \bar{\rho}(1)). \quad \square$$

5.3 Crystalline deformations ($\ell = p$)

Consider again a representation $\rho : \text{Gal}_K \rightarrow \text{GL}_n(L)$ which fulfills the conditions of Theorem 5.2. We will also make the additional assumption that all occurring Hodge-Tate weights of $\bar{\rho}$ have multiplicity one. We will consider the deformation problem crys of $\bar{\rho}$ consisting of those lifts $\tilde{\rho} : \text{Gal}_K \rightarrow \text{GL}_n(A)$ of $\bar{\rho}$ for which $\tilde{\rho} \otimes_A A'$ lies in the essential image of \mathbf{G}_K for all Artinian quotients A' of A (cf. [CHT08], Section 2.4.1). We refer to those lifts as *FL-crystalline lifts* of $\bar{\rho}$.

That crys defines a deformation condition in the sense of Definition 3.15 was already remarked in [CHT08] and follows easily from the Ramakrishna framework, cf. [Ram93]: We remarked already in Section 5.2 that the essential image of \mathbf{G}_K is closed under subobjects and quotients. That the essential image is closed under direct sums follows immediately from the exactness of \mathbf{G}_K , since then \mathbf{G}_K preserves direct sums (see [Fre64, Theorem 3.12^(*)]). Thus we can record the following (where for part 2. we refer to the remark just below Proposition 3.9):

Lemma 5.3. *Let Λ be the ring of integers of a finite, totally ramified extension E of $\text{Quot}(W(k))$ and let Λ' be the ring of integers of a finite, totally ramified extension of E (so that we have $k = k_\Lambda = k_{\Lambda'}$). Then:*

1. The functor $D_{\Lambda}^{\square, \text{crys}}(\bar{\rho})$ is representable by a quotient $R_{\Lambda}^{\square, \text{crys}}(\bar{\rho})$ of $R_{\Lambda}^{\square}(\bar{\rho})$.
2. The functor $D_{\Lambda'}^{\square, \text{crys}}(\bar{\rho})$ is representable by

$$R_{\Lambda'}^{\square, \text{crys}}(\bar{\rho}) \cong \Lambda' \otimes_{\Lambda} R_{\Lambda}^{\square, \text{crys}}(\bar{\rho}). \quad (7)$$

We remark that the condition **crys** fulfills the extended requirements as described in Remark 3.10, so that (7) holds even if $\infty > [k_{\Lambda'} : k_{\Lambda}] > 1$.

Lemma 5.4. *Under the above hypotheses*

$$R_{\Lambda}^{\square, \text{crys}}(\bar{\rho}) \cong \Lambda[[x_1, \dots, x_m]]$$

with $m = n^2 + [K : \mathbb{Q}_{\ell}] \frac{n(n-1)}{2}$.

Proof. This is a part of the statement of [CHT08, Corollary 2.4.3]. □

We also remark that the condition **crys** defines a dual-pre-**crys** condition in the sense of Definition 4.6, cf. [Gui16, Lemma 4.15].

5.4 Minimally ramified deformations ($p \neq \ell$)

For this subsection, recall from [CHT08, Section 2.4.4] the minimal ramification condition for a lift ρ of $\bar{\rho}$. Let P_K denote the kernel of one (hence, any) surjection $I_K \twoheadrightarrow \mathbb{Z}_{\ell}$. Moreover, let $\Delta_{\bar{\rho}}$ denote the set of equivalence classes of P_K -representations over k such that $\text{Hom}_{P_K}(\tau, \bar{\rho}) \neq 0$. Then the following can easily be deduced from the material in [CHT08, Section 2.4.4], in particular [CHT08, Corollary 2.4.21]:

Proposition 5.5. *Assume that any $\tau \in \Delta_{\bar{\rho}}$ is absolutely irreducible. Then we have:*

1. *The condition of being minimally ramified defines a lifting condition, denoted **min**. The representing universal object fulfills*

$$R_{\Lambda}^{\square, \text{min}}(\bar{\rho}) \cong \Lambda[[X_1, \dots, X_{n^2}]].$$

2. *If Λ is the ring of integers of some finite extension of $\text{Quot}(\Lambda)$ with residue field $k_{\Lambda'} = k$, we have*

$$R_{\Lambda'}^{\square, \text{min}}(\bar{\rho}') \cong \Lambda' \otimes_{\Lambda} R_{\Lambda}^{\square, \text{min}}(\bar{\rho}).$$

We will be particularly interested in the case where $\bar{\rho}$ has *unipotent ramification*³, i.e. where $\bar{\rho}(P_K) = \{1\}$. In the unipotent case, we have a strong connection between minimally ramified liftings and liftings of prescribed type as considered in [Sho15]. In order to make this precise, let E denote the quotient field of Λ and \bar{E} its algebraic closure.

Definition 5.6 (Def. 2.10 of [Sho15]). Let $\tau : I_K \rightarrow \text{GL}_n(\bar{E})$ be a representation which extends to a continuous representation of the Weil group W_K of K (considered with the ℓ -adic topology). Then the isomorphism class of τ is called an *inertial type*. (*Warning:* This differs from the usual definition of an inertial type as e.g. in [GK14].)

³This notion is explained by the observation that $\bar{\rho}$ is unipotently ramified if and only if $\bar{\rho}(I_K)$ lies in a conjugate of the standard unipotent subgroup consisting of upper-triangular matrices in $\text{GL}_n(k)$ with diagonal entries all equal to 1.

Let ρ be a lift of $\bar{\rho}$ which has values in \bar{E} , then we say that ρ “is of type τ ” if $\rho|_{I_K}$ is isomorphic to τ . For the following we consider a τ which is defined over E . Then we say that a morphism $x : \text{Spec } \bar{E} \rightarrow \text{Spec } R_\Lambda^\square(\bar{\rho})$ is of type τ if the associated \bar{E} -valued representation ρ_x is of type τ . This notion depends only on the image of x (because τ is defined over E).

Definition 5.7 (Fixed type deformation ring, Def. 2.14 of [Sho15]). Let $R_\Lambda^{\square, \tau}(\bar{\rho})$ be the reduced quotient of $R_\Lambda^\square(\bar{\rho})$ which is characterized by the requirement that $\text{Spec } R_\Lambda^{\square, \tau}(\bar{\rho})$ is the Zariski closure of the \bar{E} -points of type τ in $\text{Spec } R_\Lambda^\square(\bar{\rho})$.

A general classification of inertial types is given in Section 2.2.1 of [Sho15]. Under the unipotent ramification assumption, this becomes particularly simple: The set \mathcal{I}^{uni} of the isomorphism classes of inertial types which are trivial on P_K is in bijection with the set \mathcal{Y}_n of Young diagrams of size n . The partition (l_1, \dots, l_k) (with $l_i \geq l_{i+1}$) corresponds (using the notation of [Sho15]) to the type given by the I_K -restriction of the W_K -representation

$$\bigoplus_{i=1}^k \text{Sp}(\mathbf{1}, l_i),$$

where $\text{Sp}(\bullet, \bullet)$ is defined as in [Sho16, Section 3.1]. We can express this differently: Each member of \mathcal{I}^{uni} is uniquely characterized by (the conjugacy class of) its value on the generator $\zeta := \zeta_{\text{triv}}$ of I_K/P_K , and a bijection $\nabla : \mathcal{Y}_n \rightarrow \mathcal{I}^{\text{uni}}$ is given by

$$(l_1, \dots, l_k) \xrightarrow{\nabla} \tau(\zeta) = \left[1 + \begin{pmatrix} \mathcal{B}_{l_1} & & & \\ & \mathcal{B}_{l_2} & & \\ & & \ddots & \\ & & & \mathcal{B}_{l_k} \end{pmatrix} \right] \text{ with } \mathcal{B}_m = \begin{pmatrix} 0 & 1 & & \\ & 0 & 1 & \\ & & \ddots & \ddots \\ & & & 0 & 1 \\ & & & & 0 \end{pmatrix} \in \mathbb{M}_{m \times m}(E). \quad (8)$$

On the other hand, we can associate to a $\tau \in \mathcal{I}^{\text{uni}}$ a partition of n by considering the kernel sequences:

$$\Theta : \mathcal{I}^{\text{uni}} \rightarrow \mathcal{Y}_n \quad \tau \mapsto (s_1, \dots, s_r)$$

with

$$s_i := \dim \ker(\tau(\zeta) - \mathbf{1})^i - \dim \ker(\tau(\zeta) - \mathbf{1})^{i-1}$$

and

$$r := \min\{i \mid \dim \ker(\tau(\zeta) - \mathbf{1})^i = \dim \ker(\tau(\zeta) - \mathbf{1})^{i+1}\} = \min\{i \mid \ker(\tau(\zeta) - \mathbf{1})^i = V\}.$$

(Here, V is the vector space underlying τ and we use the convention that f^0 is the identity map for any linear map f .) It follows easily from the characterization of \mathcal{I}^{uni} in (8) that $s_i \geq s_{i+1}$, i.e. that Θ has values in \mathcal{Y}_n .

It is an easy combinatorial calculation to check that τ is uniquely characterized by its value under Θ and that each Young diagram occurs as a kernel sequence (i.e. that Θ is a bijection). More precisely, we have

Lemma 5.8. *The map $\Theta \circ \nabla^{-1} : \mathcal{Y}_n \rightarrow \mathcal{Y}_n$ is given by the conjugation operation on Young diagrams (cf. [FH91, §4.1] or [HHM08, Section 2.8]). In particular, for a given $\tau \in \mathcal{I}^{\text{uni}}$, the block matrix structure of $\tau(\zeta)$ (up to reordering blocks) as in (8) determines its kernel sequence and vice versa.*

Proof. Retaining the notation used in (8), we first remark that for $i \in \mathbb{N}_0$ we have

$$\dim \ker \mathcal{B}_m^i = \min(i, m).$$

Thus, setting $\mathcal{B} = \text{diag}(\mathcal{B}_{l_1}, \dots, \mathcal{B}_{l_k})$, we get

$$\dim \ker \mathcal{B}^i = \sum_{j=1}^k \min(i, l_j).$$

Consequently the kernel sequence (s_1, \dots, s_r) associated to (l_1, \dots, l_k) is given by

$$s_i = \sum_{j=1}^k \min(i, l_j) - \min(i-1, l_j) = \#\{j | l_j \geq i\} = \max\{j | l_j \geq i\}.$$

and

$$r = \max\{l_j | j = 1 \dots k\} = l_1.$$

Hence, the transition $(l_1, \dots, l_k) \rightsquigarrow (s_1, \dots, s_r)$ is precisely the conjugation operation of reflecting a Young diagram at the main diagonal (cf. [HHM08, Section 2.8]), e.g.



□

In order to state the desired comparison result, let us recall that we consider a residual representation $\bar{\rho} : \text{Gal}_K \rightarrow \text{GL}_n(k)$ with unipotent ramification. Let $\underline{\lambda} = (l_1, \dots, l_k) \in \mathcal{Y}_n$ such that $\bar{\rho}(\zeta) \sim \mathbf{1} + \text{diag}(\mathcal{B}_{l_1}, \dots, \mathcal{B}_{l_k})$. Let $\tau = \nabla(\underline{\lambda}) \in \mathcal{T}^{\text{uni}}$.

Theorem 5.9. *Assume $\bar{\rho}$ is unipotently ramified and τ as above. Then there is an isomorphism of the quotients*

$$R_{\Lambda}^{\square, \tau}(\bar{\rho}) \cong R_{\Lambda}^{\square, \min}(\bar{\rho}) \cong \Lambda[[X_1, \dots, X_{n^2}]]$$

of $R_{\Lambda}^{\square}(\bar{\rho})$, i.e. a lifting of $\bar{\rho}$ is minimally ramified if and only if it is of type τ .

Proof. The diagram

$$\begin{array}{ccc} & R_{\Lambda}^{\square, \min}(\bar{\rho}) & \\ & \nearrow & \searrow \\ R_{\Lambda}^{\square}(\bar{\rho}) & & \bar{E} \\ & \searrow & \nearrow \\ & R_{\Lambda}^{\square, \tau}(\bar{\rho}) & \end{array}$$

allows us to consider the \bar{E} -points of $\text{Spec } R_{\Lambda}^{\square, \min}(\bar{\rho})$ and $\text{Spec } R_{\Lambda}^{\square, \tau}(\bar{\rho})$ as subsets of the \bar{E} -points of $\text{Spec } R_{\Lambda}^{\square}(\bar{\rho})$. We claim that they are equal: Translated into terms of \bar{E} -valued representations, we have to compare the sets

$$\Xi^{\min} = \left\{ \rho : \text{Gal}_K \rightarrow \text{GL}_n(\bar{E}) \mid \begin{array}{l} \rho \text{ lifts } \bar{\rho} \text{ and has values in } \mathcal{O}_{\bar{E}}, \\ \dim \ker(\rho(\zeta) - \mathbf{1})^{i-1} - \dim \ker(\rho(\zeta) - \mathbf{1})^i = l_i \ \forall i \end{array} \right\}$$

and

$$\Xi^\tau = \left\{ \rho : \text{Gal}_K \rightarrow \text{GL}_n(\overline{E}) \mid \begin{array}{l} \rho \text{ lifts } \overline{\rho} \text{ and has values in } \mathcal{O}_{\overline{E}}, \\ \rho|_{I_K} \cong \tau \end{array} \right\}.$$

Lemma 5.8 implies that $\Xi^{\min} = \Xi^\tau$.

Now by definition of the ring $R_\Lambda^{\square, \tau}(\overline{\rho})$ (as the schematic closure of the points in Ξ^τ) we have

$$\ker(R_\Lambda^{\square}(\overline{\rho}) \rightarrow R_\Lambda^{\square, \tau}(\overline{\rho})) = \bigcap_{\rho \in \Xi^\tau} \ker(\rho).$$

Moreover, we clearly have

$$\ker(R_\Lambda^{\square}(\overline{\rho}) \rightarrow R_\Lambda^{\square, \min}(\overline{\rho})) \subseteq \bigcap_{\rho \in \Xi^{\min}} \ker(\rho).$$

Hence, by $\Xi^\tau = \Xi^{\min}$ we get a factorization

$$R_\Lambda^{\square}(\overline{\rho}) \rightarrow R_\Lambda^{\square, \min}(\overline{\rho}) \xrightarrow{\varphi} R_\Lambda^{\square, \tau}(\overline{\rho})$$

where the middle and the right ring have the same spectrum as topological spaces. Now we know by Proposition 5.5 that $R_\Lambda^{\square, \min}(\overline{\rho})$ is formally smooth over Λ of relative dimension n^2 and that $\dim R_\Lambda^{\square, \tau}(\overline{\rho}) = n^2 + 1$ (combine Theorem 2.4 and Proposition 2.15 of [Sho15]). Thus, φ is an isomorphism by Proposition 3.3 and the claim follows. \square

5.5 Taylors deformation condition $(1, \dots, 1)$ ($\ell \neq p$)

We continue to consider a unipotently ramified residual representation $\overline{\rho} : \text{Gal}_K \rightarrow \text{GL}_n(k)$. If $A \in \mathcal{C}_\mathcal{O}$ is a coefficient ring, we say that an A -valued lift ρ of $\overline{\rho}$ fulfills the condition $(1, \dots, 1)$ if $\text{charPoly}(\rho(\xi)) = (T - 1)^n$ for all $\xi \in I_K$. By our assumption that $\overline{\rho}$ is unipotently ramified, it is sufficient to check the case where ξ is a topological generator of the tame inertia. This defines a deformation condition (and, in comparison to [Tay08], we don't assume that $\overline{\rho}$ is trivial, cf. [Tho12, Remark before Proposition 3.17]).

Proposition 5.10. *If a lift ρ is minimally ramified, it fulfills the Taylor condition. In particular, there is a canonical surjection*

$$R^{\square, (1, \dots, 1)}(\overline{\rho}) \twoheadrightarrow R^{\square, \min}(\overline{\rho}),$$

and a morphism $R^{\square, (1, \dots, 1)}(\overline{\rho}) \rightarrow A$ factors through this surjection if and only if the associated A -valued lift of $\overline{\rho}$ is minimally ramified.

Proof. By the unipotency assumption, we can assume that $\overline{\rho}|_{P_K}$ is trivial and $\overline{\rho}(\zeta)$ is upper-triangular with each diagonal entry equal to 1 (where ζ is a topological generator of I_K/P_K). If a lift ρ is minimal, it follows that $\rho|_{P_K}$ is trivial and that $\rho(\zeta)$ is unipotent, cf. [CHT, Lemma 2.4.15, Assertion 3. \Rightarrow 1.]. It follows that $\rho(\sigma)$ is unipotent for any $\sigma \in I_K$. This proves the claim. \square

The easy proof of the following Proposition is left to the reader (cf. [KW09a, Proof of Proposition 3.8]).

Proposition 5.11. *Let L be a finite extension of K . Let*

$$\rho^{\square, (1, \dots, 1)} : G_K \rightarrow \mathrm{GL}_n(R^{\square, (1, \dots, 1)}(\bar{\rho}))$$

be the universal lifting of $\bar{\rho}$ with respect to the condition $(1, \dots, 1)$ and let

$$\rho_L^{\square, (1, \dots, 1)} : G_L \rightarrow \mathrm{GL}_n(R^{\square, (1, \dots, 1)}(\bar{\rho}|_{G_L}))$$

be the universal lifting of $\bar{\rho}|_{G_L}$ with respect to the condition $(1, \dots, 1)$. Then there exists a unique morphism of \mathcal{C}_W -algebras $\varphi : R^{\square, (1, \dots, 1)}(\bar{\rho}|_{G_L})/(\ell) \rightarrow R^{\square, (1, \dots, 1)}(\bar{\rho})/(\ell)$ such that

$$\overline{\rho^{\square, (1, \dots, 1)}}|_{G_L} = \varphi \circ \overline{\rho_L^{\square, (1, \dots, 1)}}.$$

Lemma 5.12. *Let $\tilde{\rho}$ be an A -valued lift, where we assume that A is reduced. Write $X = \tilde{\rho}(\zeta)$. Then $\chi_X := \mathrm{charPoly}(X)$ equals $(T - 1)^n$ if $\ell \geq q^{n!}$.*

Proof. Assume first that A is an integral domain. By the condition $\varphi X \varphi^{-1} = X^q$ we see that raising to the q -th power permutes the eigenvalues of X (understood as a list of n elements). Thus, any eigenvalue of X must be a $(q^{\#S_n} - 1) = (q^{n!} - 1)$ -th root of unity. Thus, if $Q(\mu)$ denotes the decomposition field of the polynomial $f(T) = T^{q^{n!} - 1} - 1$ over the quotient field of A and $A(\mu)$ denotes the integral closure of A in $Q(\mu)$, then χ_X decomposes completely in $A(\mu)[T]$. On the other hand, each eigenvalue of X is sent to 1 by the canonical reduction map

$$\pi' : A(\mu) = A(\mu) \otimes_A A \rightarrow A(\mu) \otimes_A k.$$

As the kernel of π' is a pro- ℓ -subgroup and as $(\ell^m, q^{n!} - 1) = 1$ for any $m \in \mathbb{N}$, it follows that any eigenvalue of X is 1, i.e. that $\chi_X = (T - 1)^n$. The result for a general (reduced) A follows easily from using the embedding

$$A \hookrightarrow \prod_{\mathfrak{q}} A/\mathfrak{q},$$

where \mathfrak{q} runs through the minimal primes of A . □

Corollary 5.13. *If $\ell \geq q^{n!}$, then $R^{\square, (1, \dots, 1)}(\bar{\rho}) = R^{\square}(\bar{\rho})$. In particular, $R^{\square, (1, \dots, 1)}(\bar{\rho})$ is reduced (cf. Theorem 5.1).*

Proof. By Lemma 5.12 (together with Theorem 5.1), we see that the identity map on $R^{\square}(\bar{\rho})$ factorizes through $R^{\square, (1, \dots, 1)}(\bar{\rho})$. On the other hand, $R^{\square, (1, \dots, 1)}(\bar{\rho})$ is by definition a quotient of $R^{\square}(\bar{\rho})$. Thus, we have found a surjective endomorphism of $R^{\square, (1, \dots, 1)}(\bar{\rho})$ (which must then be an isomorphism, as the rings in question are noetherian) which factorizes via $R^{\square}(\bar{\rho})$. This proves the claim. □

6 On automorphic forms on unitary groups

6.1 The group \mathcal{G}_n

For $n \in \mathbb{N}$ recall from [CHT08, Section 2.1] the definition of the group scheme \mathcal{G}_n over \mathbb{Z} and the multiplier character $\mathfrak{m} : \mathcal{G}_n \rightarrow \mathrm{GL}_1$. We write \mathcal{G}_n^0 for the connected component of the identity and \mathfrak{g}_n for the Lie algebra of \mathcal{G}_n (where we differ in notation from [CHT08]). We have $\mathcal{G}_n^{\mathrm{der}} \cong \mathrm{GL}_n$ and $\mathcal{G}_n^{\mathrm{ab}} \cong$

$\mathrm{GL}_1 \times \mathbb{Z}/2\mathbb{Z}$. If F is a CM-field with totally real subfield F^+ , recall in particular the connection between GL_n -valued conjugate self-dual representations of Gal_F and \mathcal{G}_n -valued representations of Gal_{F^+} , cf. [CHT08, Lemma 1.1.4] or [Gee11, Lemma 5.1.1].

We will be particularly interested in deformations of \mathcal{G}_n -valued residual representations. In the local split case, there is a substantial simplification possible: Let k be a finite field and let $\bar{\rho}$ be a GL_n -valued representation of Gal_F , let $\bar{\chi}$ a character such that $\bar{\chi}\bar{\rho}^\vee \cong \bar{\rho}^c$ and let $\bar{\tau}$ be the associated $\mathcal{G}_n(k)$ -valued representation of Gal_{F^+} . Moreover, let Λ be the ring of integers of a finite extension of the quotient field of $W(k)$. The following proposition now follows easily from the definitions:

Proposition 6.1. *Let ν be a place of F^+ which splits as $\tilde{\nu}\tilde{\nu}^c$ in F . Denote $\bar{\tau}_\nu := \bar{\tau}|_{\mathrm{Gal}_{F_\nu^+}}$ and $\bar{\rho}_\nu := \bar{\rho}|_{\mathrm{Gal}_{F_\nu}}$. Fix a lift $\chi_\nu : \mathrm{Gal}_{F_\nu^+} \rightarrow \Lambda^\times$ of $\mathfrak{m} \circ \bar{\tau}_\nu$. Then*

$$R_\Lambda^{(\square), \chi_\nu}(\bar{\tau}_\nu) \cong R_\Lambda^{(\square)}(\bar{\rho}_\nu) \text{ and } H^i(F_\nu^+, \mathfrak{g}_n^{\mathrm{der}}) \cong H^i(F_\nu, \mathfrak{gl}_n), Z^1(F_\nu^+, \mathfrak{g}_n^{\mathrm{der}}) \cong Z^1(F_\nu, \mathfrak{gl}_n).$$

This observation allows us to define local conditions for deformations of $\bar{\tau}$ at split places by GL_n -valued local conditions. In order to make this precise, let $\Sigma \subset \mathrm{Pl}_{F^+}^{\mathrm{fin}}$ be a finite set of places and assume that any place in Σ splits as $\nu = \tilde{\nu}\tilde{\nu}^c$ in the extension $F|F^+$ (so, in particular, we fix a place $\tilde{\nu}$ above ν). Moreover, assume that $\bar{\tau}$ is unramified outside Σ , i.e. factorizes through $\mathrm{Gal}_{F^+, \Sigma}$. We set $\tilde{\Sigma} := \{\tilde{\nu} | \nu \in \Sigma\}$. Fix a character $\chi : \mathrm{Gal}_{F^+, \Sigma} \rightarrow \Lambda^\times$ lifting $\mathfrak{m} \circ \bar{\tau}$. Moreover, for each $\tilde{\nu} \in \tilde{\Sigma}$ fix a deformation condition D_ν of the GL_n -valued representation $\bar{\rho}_\nu$.

Definition 6.2 (Deformation problem, following [CHT08]). The collection

$$\mathcal{S} = (F|F^+, \Sigma, \tilde{\Sigma}, \Lambda, \bar{\tau}, \chi, \{D_\nu\}_{\nu \in \Sigma}),$$

parametrizing deformations r of $\bar{\tau}$ to \mathcal{C}_Λ which fulfill $\mathfrak{m} \circ r = \chi$, which are unramified outside Σ and fulfill D_ν (via Proposition 6.1) at $\nu \in \Sigma$, defines a global deformation condition.

We end this section by a remark on the conventions for multiple framings, in which we differ from [CHT08]. For this, let $T \subset \Sigma$ be a non-empty subset and recall our Definition 3.16 for the multiply framed deformation functor $D_\Lambda^{\square T, \mathcal{S}}(\bar{\tau})$ and its representing object $R_\Lambda^{\square T, \mathcal{S}}(\bar{\tau})$. Comparing this with the functor and representing object considered in [CHT08, Definition 2.2.7], which we denote by $D_\Lambda^{\square T, \mathcal{S}}(\bar{\tau})$ and $R_\Lambda^{\square T, \mathcal{S}}(\bar{\tau})$, we easily get the following observation:

Proposition 6.3. *$D_\Lambda^{\square T, \mathcal{S}}(\bar{\tau})$ is representable if and only if $D_\Lambda^{\square T, \mathcal{S}}(\bar{\tau})$ is representable, and in this case we have*

$$R_\Lambda^{\square T, \mathcal{S}}(\bar{\tau}) \cong R_\Lambda^{\square T, \mathcal{S}}(\bar{\tau})[[X_1, \dots, X_{\#T}]].$$

6.2 Automorphic forms and Hecke algebras

For this subsection, let us assume that the extension $F|F^+$ is unramified at all finite places and, in case n is even, that $\frac{n}{2}[F^+ : \mathbb{Q}]$ is even. This allows us to fix a definite unitary group H over \mathcal{O}_{F^+} , as considered in [Gue11, Section 2.11] or [Ger10, Section 1.1], whose key properties we recall here:

- The extension of scalars of H to F^+ is an outer form of GL_n/F^+ , which becomes isomorphic to GL_n/F after extending scalars to F ;
- H is quasi-split at every finite place of F^+ ;

- H is totally definite, i.e. $H(F_\infty^+)$ is compact and $H(F_\nu^+) \cong U_n(\mathbb{R})$ for all infinite places ν of F^+ ;
- For any finite place ν of F^+ which splits as $\tilde{\nu}\tilde{\nu}^c$ in F , we can choose an isomorphism $\iota_{\tilde{\nu}} : H(F_\nu^+) \rightarrow \mathrm{GL}_n(F_{\tilde{\nu}})$ whose restriction to $H(\mathcal{O}_{F_\nu^+})$ provides an isomorphism $H(\mathcal{O}_{F_\nu^+}) \cong \mathrm{GL}_n(\mathcal{O}_{F_{\tilde{\nu}}})$.

Level subgroups Let us fix a finite subset $\mathcal{T} \subset \mathrm{Pl}_{F^+}^{\mathrm{fin}}$ such that each $\nu \in \mathcal{T}$ splits as $\tilde{\nu}\tilde{\nu}^c$ in F . For the remainder of this section, the letter U will denote an open compact subgroup of $H(\mathbb{A}_{F^+}^\infty)$. For later applications, we will be particularly interested in the choice $U_{\mathcal{T}} := \prod_{\nu \in \mathrm{Pl}_{F^+}^{\mathrm{fin}}} U_\nu$ with:

- If ν is not split in $F|F^+$, then U_ν is a hyperspecial maximal compact subgroup of $H(F_\nu^+)$;
- If $\nu \notin \mathcal{T}$ splits, then $U_\nu = H(\mathcal{O}_{F_\nu^+})$;
- If $\nu \in \mathcal{T}$, then $U_\nu = \iota_{\tilde{\nu}}^{-1}(\mathrm{Iw})$, where $\mathrm{Iw} \subset \mathrm{GL}_n(\mathcal{O}_{F_{\tilde{\nu}}})$ denotes the Iwahori subgroup.

We remark that in many articles (e.g. [CHT08]) the set \mathcal{T} is enlarged by a choice of auxiliary places at which a suitable level condition is imposed. Our arguments don't require such auxiliary places.

Weights Recall the parametrization of complex and ℓ -adic representations of unitary and general linear groups, e.g. from [Gue11]:

- To a tuple $\omega = (\underline{\omega}_\tau) \in (\mathbb{Z}^{n,+})^{\mathrm{Hom}(F^+, \mathbb{R})}$ we associate the representation

$$\xi_\omega^u : H(F_\infty^+) = \prod_{\tau \in \mathrm{Hom}(F^+, \mathbb{R})} H(F_\tau^+) \cong \prod_{\tau \in \mathrm{Hom}(F^+, \mathbb{R})} U_n(\mathbb{R}) \xrightarrow{\varphi} \prod_{\tau \in \mathrm{Hom}(F^+, \mathbb{R})} \mathrm{GL}_n(W_{\underline{\omega}_\tau}^u) \subset \mathrm{GL}_n(W_\omega^u),$$

where $W_\omega^u = \otimes_\tau W_{\underline{\omega}_\tau}^u$ and where φ is the product of the highest weight representations $W_{\underline{\omega}_\tau}^u$ attached to the weight $\underline{\omega}_\tau$ (see e.g. [BC09, Gue11, Ger10]).

- Let ℓ be a rational prime such that every place ν of F^+ above ℓ splits in $F|F^+$ and fix for each such ν a place $\tilde{\nu}$ of F above ν . Let \mathcal{K} be a finite extension of \mathbb{Q}_ℓ which is F -big enough and let $\omega = (\underline{\omega}_\tau) \in (\mathbb{Z}^{n,+})^{\mathrm{Hom}(F, \mathcal{K})}$. To each $\tau \in \mathrm{Hom}(F, \mathcal{K})$ we can associate a place ν of F^+ above ℓ for which we have just fixed a place $\tilde{\nu}$. Denote this assignment $\mathrm{Hom}(F, \mathcal{K}) \rightarrow \Omega_\ell^F$ by $\tau \mapsto w_\tau$. Let

$$\begin{aligned} \xi_\omega^\mathcal{K} : \prod_{\nu \in \Omega_\ell^F} H(F_\nu^+) &\cong \prod_{\nu \in \Omega_\ell^F} \mathrm{GL}_n(F_{\tilde{\nu}}) \xrightarrow{\prod d_\nu} \prod_{\nu \in \Omega_\ell^F} \prod_{\substack{\tau \in \mathrm{Hom}(F, \mathcal{K}) \\ w_\tau = \tilde{\nu}}} \mathrm{GL}_n(F_{\tilde{\nu}}) = \prod_{\tau \in \mathrm{Hom}(F, \mathcal{K})} \mathrm{GL}_n(F_{\tilde{\nu}}) \\ &\xrightarrow{\psi} \prod_{\tau \in \mathrm{Hom}(F, \mathcal{K})} \mathrm{GL}_n(W_{\underline{\omega}_\tau}^\mathcal{K}) \subset \mathrm{GL}_n(W_\omega^\mathcal{K}) \end{aligned}$$

be the representation where each d_ν is the diagonal embedding, where $W_\omega^\mathcal{K} = \otimes_\tau W_{\underline{\omega}_\tau}^\mathcal{K}$ and where ψ is the product of the highest weight representations $W_{\underline{\omega}_\tau}^\mathcal{K}$ attached to the weight $\underline{\omega}_\tau$. The representation $\xi_\omega^\mathcal{K}$ admits an integral model over $\mathcal{O}_\mathcal{K}$, whose underlying finite free $\mathcal{O}_\mathcal{K}$ -module we denote by $M_\omega^{\mathcal{O}_\mathcal{K}}$.

Automorphic forms We denote by

$$\mathcal{A}(H) = \bigoplus_{\pi} \pi^{m(\pi)}$$

the space of (complex) automorphic forms on H , which decomposes into isomorphism classes of irreducible representations of $H(\mathbb{A}_{F^+})$, each occurring with finite multiplicity $m(\pi)$ (see e.g. [Gue11]).

Definition 6.4 (Vector-valued automorphic form). Let $\omega \in (\mathbb{Z}^{n,+})^{\text{Hom}(F^+, \mathbb{R})}$ be a weight, then we denote by \mathcal{S}_{ω} the space of locally constant functions $f : H(\mathbb{A}_{F^+}^{\infty}) \rightarrow W_{\omega}^{u, \vee}$ which fulfill

$$f(\gamma.h) = \gamma_{\infty} f(h) \quad \forall h \in H(\mathbb{A}_{F^+}^{\infty}), \gamma \in H(F^+).$$

(We denote by γ_{∞} the image of γ under the canonical embedding $H(F^+) \rightarrow H(F_{\infty}^+)$). $H(\mathbb{A}_{F^+}^{\infty})$ acts on \mathcal{S}_{ω} by right translation, and for a level subgroup U we denote by $\mathcal{S}_{\omega}(U)$ the space of U -fixed vectors.

There exists an $H(\mathbb{A}_{F^+}) = H(\mathbb{A}_{F^+, \infty}) \times H(\mathbb{A}_{F^+}^{\infty})$ -equivariant decomposition

$$\mathcal{A}(H) = \bigoplus_{\omega} W_{\omega}^u \otimes \mathcal{S}_{\omega}.$$

Thus we can associate to an $f \in \mathcal{S}_{\omega}$ the (irreducible) automorphic representation $\langle f \rangle$ which is uniquely characterized by the condition that it contains all vectors of $W_{\omega}^u \otimes f$. The main feature of the group H is the existence of avatars:

Theorem 6.5. *Let Π be a RACSDC automorphic representation of $\text{GL}_n(\mathbb{A}_F)$ of weight $\omega \in (\mathbb{Z}^{n,+})^{\text{Hom}(F, \mathbb{C})}$ in the sense of [CHT08, Section 4]. Then there exists an automorphic representation π_0 of $H(\mathbb{A}_{F^+})$ such that Π is a base change of π_0 , i.e.*

- for each archimedean place ν of F^+ and each place $\tilde{\nu}$ of F above ν , we have $\pi_{0, \nu} \cong \xi_{\omega_{\tilde{\nu}}}^u$;
- for each finite place ν of F^+ which splits as $\tilde{\nu}\tilde{\nu}^c$ in F , $\Pi_{\tilde{\nu}}$ is the local base change of $\pi_{0, \nu}$;
- if ν is a finite place of F^+ which stays inert in F and for which Π_{ν} is unramified, then π_{ν} has a fixed vector for a maximal hyperspecial compact subgroup of $H(F_{\nu}^+)$.

Proof. See [Gue11, Theorem 2.2] and [Ger10, Lemma 2.2.7]. □

Hecke algebras We continue to consider a fixed set of places \mathcal{T} as above (with corresponding level subgroup $U = U_{\mathcal{T}}$) and a weight ω . For $j \in \{1, \dots, n\}$ and for w a place of F which is split over F^+ and does not divide an element of \mathcal{T} , we consider the following Hecke operator (acting on $\mathcal{S}_{\omega}(U)$):

$$T_{F_w}^{(j)} = \left[U.l_w^{-1} \begin{pmatrix} \varpi_{F_w} \mathbf{1}_j & 0 \\ 0 & \mathbf{1}_{n-j} \end{pmatrix} .U \right]$$

For a finite set $\mathcal{T}' \subset \text{Pl}_{F^+}^{\text{fin}}$ containing \mathcal{T} and a subring \mathcal{R} of \mathbb{C} we define the Hecke algebra

$$\mathcal{R}\mathbf{T}_{\omega}^{\mathcal{T}'}(U) := \text{im} \left(\mathcal{R}[T_{F_w}^{(j)} | j \in \{1, \dots, n\}, w \in \text{Pl}_F^{\text{split}, \mathcal{T}'}] \rightarrow \text{End}_{\mathbb{C}}(\mathcal{S}_{\omega}(U)) \right),$$

where $\text{Pl}_F^{\text{split}, \mathcal{T}'}$ denotes the set of places of F which are split over F^+ and which do not divide an element of \mathcal{T}' . Besides the case $\mathcal{R} = \mathbb{Z}$ we will be interested in $\mathcal{R} = \mathcal{E}_f$ (the coefficient field of an eigenform f with respect to ${}^{\mathbb{Z}}\mathbf{T}_{\omega}^{\mathcal{T}}(U)$) and in $\mathcal{R} = \mathcal{E}(U) = \prod_f \mathcal{E}_f$, where the product (i.e. the field compositum operation) runs through all eigenforms of $\mathcal{S}_{\omega}(U)$. We note the following well-known facts: There are only finitely many (one-dimensional) eigenspaces $\mathbb{C}.f_1, \dots, \mathbb{C}.f_r$ contained in $\mathcal{S}_{\omega}(U)$, so $\mathcal{E}(U)$ is a number field. Moreover, $\mathcal{S}_{\omega}(U)$ admits a basis of eigenforms, i.e. we can choose the f_i such that

$$\mathcal{S}_{\omega}(U) \cong \mathbb{C}.f_1 \oplus \dots \oplus \mathbb{C}.f_r$$

as a $\mathbf{T}_{\omega}^{\mathcal{T}}(U)$ -module (see decomposition (3.1.1) of [Gue11]). By mapping a Hecke operator to its f -eigenvalue, any eigenform $f \in \mathcal{S}_{\omega}(U)$ gives rise to a \mathbb{Z} -algebra-homomorphism

$$\varphi_f : {}^{\mathbb{Z}}\mathbf{T}_{\omega}^{\mathcal{T}}(U) \longrightarrow \mathcal{E}(U) \quad T_{F_w}^{(j)} \longmapsto a_f(T_{F_w}^{(j)})$$

and it can be shown that $\text{im}(\varphi_f) \subset \mathcal{O}_{\mathcal{E}(U)}$. The form f is uniquely characterized by φ_f (up to \mathbb{C} -multiples).

ℓ -adic models of automorphic forms The following is based on Section 2.3 of [Gue11]. For this paragraph, we fix a rational prime ℓ which does not lie below \mathcal{T} and such that all places of F^+ above ℓ are split in the extension $F|F^+$ and consider the following setup: Let \mathcal{K} be a finite extension of \mathbb{Q}_{ℓ} which is F -big enough and fix an isomorphism $\iota : \overline{\mathcal{K}} \cong \mathbb{C}$. Moreover, we fix an ℓ -adic weight ω , i.e. an element of

$$(\mathbb{Z}^{n,+})_c^{\text{Hom}(F, \mathcal{K})} = \left\{ \omega \in (\mathbb{Z}^{n,+})^{\text{Hom}(F, \mathcal{K})} \mid \underline{\omega}_{\tau^c, i} = -\underline{\omega}_{\tau, n-i+1} \quad \forall \tau \in \text{Hom}(F, \mathcal{K}), i \in \{1, \dots, n\} \right\}.$$

Definition 6.6. For $U \subset H(\mathbb{A}_{F^+}^{\infty})$ a compact subgroup and an $\mathcal{O}_{\mathcal{K}}$ -algebra A , suppose that either the projection of U to $H(F_{\ell}^+)$ is contained in $H(\mathcal{O}_{F_{\ell}^+})$ or that A is a \mathcal{K} -algebra. Then we define

$$\mathcal{S}_{\omega}(U, A) = \{ f : H(F^+) \backslash H(\mathbb{A}_{F^+}^{\infty}) \rightarrow A \otimes_{\mathcal{O}_{\mathcal{K}}} M_{\omega}^{\mathcal{O}_{\mathcal{K}}} \mid u_{\ell}.f(hu) = f(h) \quad \forall u \in U, h \in H(\mathbb{A}_{F^+}^{\infty}) \},$$

where u_{ℓ} denotes the image of u under the projection map $H(\mathbb{A}_{F^+}^{\infty}) \rightarrow H(F_{\ell}^+)$.

We are primarily interested in the case that A is $\mathcal{O}_{\mathcal{K}}$ -flat, so that we have $\mathcal{S}_{\omega}(U, A) \cong A \otimes_{\mathcal{O}_{\mathcal{K}}} \mathcal{S}_{\omega}(U, \mathcal{O}_{\mathcal{K}})$. The main connection with complex automorphic forms is as follows (cf. also [Gue11, Section 2.3]): The isomorphism ι gives rise to a bijection $\iota_*^+ : (\mathbb{Z}^{n,+})_c^{\text{Hom}(F, \mathcal{K})} \cong (\mathbb{Z}^{n,+})^{\text{Hom}(F, \mathbb{R})}$, and the assignment $f \mapsto (h \mapsto \theta_{\omega}(h_{\ell}.f(h)))$ provides isomorphisms of $\mathbb{C}H(\mathbb{A}_{F^+}^{\infty})$ -modules

$$\bigcup_U \mathcal{S}_{\omega}(U, \mathbb{C}) \cong \mathcal{S}_{\iota_*^+(\omega)^{\vee}} \quad \text{and} \quad \mathcal{S}_{\omega}(U, \mathbb{C}) \cong \mathcal{S}_{\iota_*^+(\omega)^{\vee}}(U). \quad (9)$$

(Here, \mathbb{C} is understood as a $\mathcal{O}_{\mathcal{K}}$ -algebra via ι and $\iota_*^+(\omega)^{\vee}$ is defined by $\iota_*^+(\omega)_{\tau, i}^{\vee} = -\iota_*^+(\omega)_{\tau, n+1-i}$.)

For a place w not dividing ℓ , the operators $T_{F_w}^{(j)}$ also act on $\mathcal{S}_{\omega}(U, \mathcal{O}_{\mathcal{K}}) \subset \mathcal{S}_{\omega}(U, \mathbb{C})$, and this action commutes with the isomorphism (9). This motivates the following definition: Let \mathcal{T}' be a finite set of places of F^+ containing $\mathcal{T} \cup \Omega_{\ell}^{F^+}$ and let \mathcal{R} be a subring of $\mathcal{O}_{\mathcal{K}}$, then we define the Hecke algebra

$$\mathcal{R}\mathbf{T}_{\omega}^{\mathcal{T}'}(U) = \text{im} \left(q : \mathcal{R}[T_{F_w}^{(j)} \mid j \in \{1, \dots, n\}, w \in \text{Pl}_F^{\text{split}, \mathcal{T}'}] \longrightarrow \text{End}_{\mathcal{O}_{\mathcal{K}}}(\mathcal{S}_{\omega}(U, \mathcal{O}_{\mathcal{K}})) \right),$$

where we will often abbreviate $\mathbb{T}_\omega^{\mathcal{T}'}(U) = {}_{\mathcal{O}_K}\mathbb{T}_\omega^{\mathcal{T}'}(U)$. If $f \in S_\omega(U, \mathcal{O}_K)$ is an eigenform for this algebra, then we see, using the compatibility with the isomorphism (9), that the eigenvalue for a Hecke operator T is given by $\iota^{-1}(a_{\tilde{f}})$, where $\tilde{f} \in \mathcal{S}_{\iota_*^+(\omega)\nu}(U)$ is the corresponding complex automorphic form. In other words, we can interpret the map $\varphi_{\tilde{f}}$ from above as

$$\varphi_{\tilde{f}}^\ell : \mathbb{Z}\mathbb{T}_\omega^{\mathcal{T}_\ell}(U) \longrightarrow \iota(\mathcal{E}(U)) \cong \mathcal{E}(U).$$

Note that we use the bold symbol \mathbf{T} for complex Hecke algebras and the blackboard bold symbol \mathbb{T} for ℓ -adic Hecke algebras.

Fixed type Hecke algebras Fix a finite set $\tilde{\Sigma} \subset (\mathcal{T}' - \Omega_\ell^F)$ of places of F together with a tuple $\underline{\sigma} = (\sigma_\nu)_{\nu \in \tilde{\Sigma}}$, where each σ_ν is a finite-dimensional complex representation of $\mathrm{GL}_n(\mathcal{O}_{F_\nu})$. Let ${}_{\underline{\sigma}}S_\omega(U, \mathcal{O}_K) \subset S_\omega(U, \mathcal{O}_K)$ be the subspace of those forms f whose complex correspondents \tilde{f} fulfill the following condition for all places $\nu \in \tilde{\Sigma}$: If π_ν denotes the local component of the automorphic representation $\pi = \langle \tilde{f} \rangle$ at ν , then $\pi_\nu|_{\mathrm{GL}_n(\mathcal{O}_{F_\nu})}$ contains σ_ν as a subrepresentation. Note that the $T_{F_w}^{(j)}$ (for w in $\mathrm{Pl}_F^{\mathrm{split}, \mathcal{T}'}$) stabilize the subspace ${}_{\underline{\sigma}}S_\omega(U, \mathcal{O}_K)$, so we can define

$${}_{\underline{\sigma}}\mathbb{T}_\omega^{\mathcal{T}'}(U) = \mathrm{im} \left({}_{\underline{\sigma}}q : \mathcal{R}[T_{F_w}^{(j)} | j \in \{1, \dots, n\}, w \in \mathrm{Pl}_F^{\mathrm{split}, \mathcal{T}'}] \longrightarrow \mathrm{End}_{\mathcal{O}_K}({}_{\underline{\sigma}}S_\omega(U, \mathcal{O}_K)) \right).$$

We easily see that the assignment $q(T_{F_w}^{(j)}) \mapsto {}_{\underline{\sigma}}q(T_{F_w}^{(j)})$ defines an \mathcal{R} -algebra surjection ${}_{\underline{\sigma}}\theta$ from ${}_{\mathcal{R}}\mathbb{T}_\omega^{\mathcal{T}'}(U)$ to ${}_{\underline{\sigma}}\mathbb{T}_\omega^{\mathcal{T}'}(U)$. We note the following (for $\mathcal{R} = \mathcal{O}_K$):

- In the same way as for ${}_{\mathcal{O}_K}\mathbf{T}_\omega^{\mathcal{T}'}(U)$, we can check that ${}_{\underline{\sigma}}{}_{\mathcal{O}_K}\mathbb{T}_\omega^{\mathcal{T}'}(U)$ is free and finitely generated over \mathcal{O}_K ;
- assume that ${}_{\mathcal{O}_K}\mathbf{T}_\omega^{\mathcal{T}'}(U)_{\mathfrak{m}} \cong \mathcal{O}_K$ holds for any maximal ideal \mathfrak{m} , then ${}_{\underline{\sigma}}{}_{\mathcal{O}_K}\mathbb{T}_\omega^{\mathcal{T}'}(U)_{\mathfrak{n}}$ is a quotient of \mathcal{O}_K for any maximal ideal \mathfrak{n} . By the above bullet point, it thus follows that ${}_{\underline{\sigma}}{}_{\mathcal{O}_K}\mathbb{T}_\omega^{\mathcal{T}'}(U)_{\mathfrak{n}}$ is isomorphic to \mathcal{O}_K for any maximal ideal \mathfrak{n} .

6.3 Attaching Galois representations to automorphic forms

Retain all notation from above and let $\mathfrak{m} \subset {}_{\mathcal{O}_K}\mathbb{T}_\omega^{\mathcal{T}_\ell}(U)$ be a maximal ideal.

Proposition 6.7 ([Gue11, Proposition 3.1 and 3.2]). *There exists a representation*

$$\rho_{\mathfrak{m}} : \mathrm{Gal}_F \longrightarrow \mathrm{GL}_n({}_{\mathcal{O}_K}\mathbb{T}_\omega^{\mathcal{T}_\ell}(U)_{\mathfrak{m}})$$

with the following properties, where the first two already characterize $\rho_{\mathfrak{m}}$ uniquely:

1. $\rho_{\mathfrak{m}}$ is unramified at all but finitely many places; If a place ν of F^+ is inert and unramified in F and if U_ν is a hyperspecial maximal compact subgroup of $H(F_\nu^+)$, then $\rho_{\mathfrak{m}}$ is unramified above ν ;
2. If $\nu \in \mathrm{Pl}_{F^+}^{\mathrm{fin}} \setminus \mathcal{T}_\ell$ splits as $\tilde{\nu}\tilde{\nu}^c$ in F , then $\rho_{\mathfrak{m}}$ is unramified at $\tilde{\nu}$ and

$$\mathrm{charPoly}(\rho_{\mathfrak{m}}(\mathrm{Frob}_{\tilde{\nu}})) = X^n - T_{\tilde{\nu}}^{(1)} X^{n-1} + \dots + (-1)^j (N\tilde{\nu})^{\frac{j(j-1)}{2}} T_{\tilde{\nu}}^{(j)} X^{n-j} + \dots + (-1)^n (N\tilde{\nu})^{\frac{n(n-1)}{2}} T_{\tilde{\nu}}^{(n)}.$$

3. $\overline{\rho}_{\mathfrak{m}}^c \cong \overline{\rho}_{\mathfrak{m}} \otimes \overline{\epsilon}_\ell^{1-n}$;

4. Fix a set $\tilde{\Omega}_\ell^{F^+}$ of places of F such that $\tilde{\Omega}_\ell^{F^+} \sqcup \tilde{\Omega}_\ell^{F^+,c} = \Omega_\ell^F$ and denote by \tilde{I}_ℓ the set of embeddings $F \hookrightarrow \mathcal{K}$ which give rise to an element of $\tilde{\Omega}_\ell^{F^+}$. Suppose that $w \in \tilde{\Omega}_\ell^{F^+}$ is unramified over ℓ , that $U_{\bar{w}} = H(\mathcal{O}_{F^+, \bar{w}})$ (for $\bar{w} \in \text{Pl}_{F^+}$ the place below w) and that for each $\tau \in \tilde{I}_\ell$ above w we have

$$\ell - 1 - n \geq \omega_{\tau,1} \geq \dots \geq \omega_{\tau,n} \geq 0.$$

Then, for each open ideal $I \subset \mathcal{O}_\kappa \mathbb{T}_\omega^{\mathcal{T}_\ell}(U)$ there is an object $M_{\mathfrak{m}, I, w}$ of $\underline{\mathbf{MF}}_{\mathcal{O}_{F_w}, \mathcal{O}_\kappa}$ such that

$$(\rho_{\mathfrak{m}} \otimes_{\mathcal{O}_\kappa \mathbb{T}_\omega^{\mathcal{T}_\ell}(U)} \mathcal{O}_\kappa \mathbb{T}_\omega^{\mathcal{T}_\ell}(U)/I) | \text{Gal}_{F_w} \cong \mathbf{G}_{F_w}(M_{\mathfrak{m}, I, w}).$$

If \mathfrak{m} is non-Eisenstein in the sense of [CHT08, Definition 3.4.3], then $\rho_{\mathfrak{m}}$ and its reduction extend to

$$r_{\mathfrak{m}} : \text{Gal}_{F^+} \rightarrow \mathcal{G}_n(\mathcal{O}_\kappa \mathbb{T}_\omega^{\mathcal{T}_\ell}(U)_{\mathfrak{m}}) \quad \text{and} \quad \bar{r}_{\mathfrak{m}} : \text{Gal}_{F^+} \rightarrow \mathcal{G}_n(\mathcal{O}_\kappa \mathbb{T}_\omega^{\mathcal{T}_\ell}(U)/\mathfrak{m}).$$

Moreover, $m \circ r_{\mathfrak{m}} = \epsilon_\ell^{1-n} \delta_{F|F^+}^{\mu_{\mathfrak{m}}}$ for a suitable $\mu_{\mathfrak{m}} \in \mathbb{Z}/2\mathbb{Z}$, where $\delta_{F|F^+}$ is the non-trivial character of $\text{Gal}(F|F^+)$.

In this way we can associate to a RACSDC automorphic representation π of $\text{GL}_n(\mathbb{A}_F)$ and a finite place λ of $\mathcal{E}(U)$ a residual representation $\bar{r}_{\pi, \lambda} : \text{Gal}_{F^+} \rightarrow \mathcal{G}_n(\overline{\mathbb{F}}_{\ell(\lambda)})$. Let us assume that $\bar{r}_{\pi, \lambda}$ is absolutely irreducible for all λ in a subset of $\text{Pl}_{\mathcal{E}(U)}^{\text{fin}}$ of Dirichlet density 1. Then the set

$$\Lambda_{\mathcal{E}(U)}^1 = \{\lambda | \bar{\rho}_{\pi, \lambda'} \text{ is absolutely irreducible for all } \lambda' \text{ dividing } \ell(\lambda)\}$$

has also Dirichlet density 1. In this way, we get an association from π to the compatible systems of residual Galois representations $\mathcal{R}_\pi = (\bar{r}_{\pi, \lambda})_{\lambda \in \Lambda_{\mathcal{E}(U)}^1}$ and $\mathcal{R}'_\pi = (\bar{\rho}_{\pi, \lambda})_{\lambda \in \Lambda_{\mathcal{E}(U)}^1}$.

7 Consequences from modularity lifting theorems

Let us start with the following adaption of [KW09a, Lemma 3.6]:

Lemma 7.1. *Let k be a finite field of characteristic ℓ , G a profinite group satisfying the ℓ -finiteness condition and $\eta : G \rightarrow \mathcal{G}_n(k)$ be an absolutely irreducible continuous representation. Let $\mathcal{F}_n(G)$ be a subcategory of deformations of η in k -algebras which defines a deformation condition. Let $\eta_{\mathcal{F}} : G \rightarrow \text{GL}_n(R_{\mathcal{F}})$ be the universal deformation of η in $\mathcal{F}_n(G)$. Then $R_{\mathcal{F}}$ is finite if and only if $\eta_{\mathcal{F}}(G)$ is finite.*

Proof of the Lemma. The proof of Lemma 3.6 of Khare-Wintenberger goes through verbatim, except that we have to refer to [CHT08, Lemma 2.1.12] instead of Carayol's Lemma. \square

7.1 A minimal $R = T$ -theorem

Our starting point is a RACSDC automorphic representation $\pi = \langle f \rangle \subset \mathcal{S}_\omega(U)$ (where $U = U(S)$ for a finite set of places S of F) and a place $\lambda \in \Lambda_{\mathcal{E}(U)}^1$. Fix a finite F -big enough extension \mathcal{K} of $\mathcal{E}(U)_\lambda$. We abbreviate $r, \bar{r}, \rho, \bar{\rho}$ for the associated Galois representations via Proposition 6.7 for the unique maximal ideal \mathfrak{m} containing $\mathcal{O}_\kappa \otimes_{\mathbb{Z}} \ker \varphi_f^\ell$. We assume furthermore the following:

- All places of S_ℓ split in the extension $F|F^+$;

- all ramification of ρ is unipotent (this can always be achieved by a finite solvable base change);
- ρ is a minimally ramified (at all places in S) and FL-crystalline (at all places dividing ℓ) lift of $\bar{\rho}$;
- $\bar{\rho}$ is absolutely irreducible;
- $\bar{\rho}(\text{Gal}_{F(\zeta_\ell)})$ is adequate.

Let us abbreviate $R^{(\min),[\text{crys}]} := R_{\mathcal{O}_K, S_\ell}^{\chi, (\min), [\text{crys}]}(\bar{\tau})$ for the ring parametrizing fixed-determinant deformations of $\bar{\tau}$ which are unramified outside S_ℓ , (minimally ramified in S) and [FL-crystalline at places dividing ℓ]. Moreover, let \mathbb{T} (resp. \mathbb{T}^{\min}) denote the Hecke algebra $\mathcal{O}_K \mathbb{T}_\omega^{S_\ell}(U)_\mathfrak{m}$ (resp. $\mathcal{O}_K \mathbb{T}_\omega^{S_\ell}(U)_{\underline{\sigma}\theta(\mathfrak{m})}$), where \mathfrak{m} is the maximal ideal such that $\bar{\tau}_\mathfrak{m} \cong \bar{\tau}$ and where $\underline{\sigma} = (\sigma_\nu)_{\nu \in \tilde{S}}$ is defined as follows:

For each $\nu \in \tilde{S}$ we can associate an inertial type τ_ν in the same way as we did just before Theorem 5.9. To each τ_ν one can associate a representation $\sigma_\nu = \sigma(\tau_\nu)$ of $K = \text{GL}_n(\mathcal{O}_{F_\nu})$ (which is then the K -type of the $\text{GL}_n(F_\nu)$ -representation associated to an extension of τ_ν to Gal_{F_ν} .) For more details on the construction of $\sigma(\tau)$, see [Sho15, Section 4.6], [BC09, Section 6.5.2], [SZ99] and the following remark:

Remark 7.2. Recall the following:

- Consider the finite group $\mathfrak{G} = \text{GL}_n(\ell(\nu))$ and its standard Borel subgroup $\mathfrak{B} \subset \mathfrak{G}$. Then the irreducible constituents of the (complex) representation $\text{ind}_{\mathfrak{B}}^{\mathfrak{G}}(1)$ are called the unipotent representations of \mathfrak{G} . These representations can (canonically) be parametrized by the irreducible representations of the Weyl group $\mathcal{W}(\mathfrak{G}) \cong S_n$, see e.g. [Pra14, Corollary 4.4]. The irreducible representations of S_n in turn can be parametrized by partitions of n in terms of Specht modules, cf. [JK81]. In other words, we get a canonical bijection $h : \mathcal{Y}_n \cong \text{Rep}(\mathfrak{G})^{\text{uni}}$, where $\text{Rep}(\mathfrak{G})^{\text{uni}}$ denotes the set of all unipotent representations of \mathfrak{G} up to isomorphism. The map h can be explicitly described in terms of induction from Levi subgroups (cf. [Sho15, Definition 4.34]) and sends $(1, \dots, 1)$ to the trivial representation and (n) to the Steinberg representation.
- Under the unipotent ramification assumption, the set of inertial types \mathcal{I}^{uni} is in bijection with the set \mathcal{Y}_n of partitions of n via ∇ from Section 5.4.

Now, we have a decomposition

$$\text{ind}_I^K(1) \cong \text{infl}_{\mathfrak{G}}^K \text{ind}_{\mathfrak{B}}^{\mathfrak{G}}(1) \cong \bigoplus_{\Pi \in \text{Rep}(\mathfrak{G})^{\text{uni}}} m_\Pi \text{infl}_{\mathfrak{G}}^K(\Pi),$$

where $I \subset K$ denotes the Iwahori subgroup, $\text{infl}_{\mathfrak{G}}^K$ denotes the inflation along the pro- $\ell(\nu)$ -radical of K and where the $m_\Pi \geq 1$ are suitable multiplicities. Analogous to [BC09, Remark 6.5.2 iii)], one can thus check that the assignment $\tau \mapsto \sigma(\tau)$ is described in terms of partitions as $\tau \mapsto \text{infl}_{\mathfrak{G}}^K(h \circ \nabla(\tau))$. Observe that the special case $n = 2$ is precisely [BC09, Remark 6.5.2 iii)] and [Sho15, Example 2.17].

We stress that the notions \mathbb{T} and \mathbb{T}^{\min} depend on the choice of the place λ .

Proposition 7.3. *A map $h : \mathbb{T} \rightarrow \overline{\mathbb{Q}}_\ell$ factorizes through \mathbb{T}^{\min} if and only if the concatenation $h' : R^{\text{crys}} \rightarrow \mathbb{T} \rightarrow \overline{\mathbb{Q}}_\ell$ factors through $R^{\text{min, crys}}$.*

Proof. The map h corresponds to an automorphic form $g \in S_\omega(U, \mathcal{O}_K)$ such that $\bar{r}_{\langle g \rangle} \cong \bar{r}$. By the above and Theorem 5.9, $\rho_{\langle g \rangle, \nu}$ (for $\nu \in S$) is a minimally ramified lift of $\bar{\rho}_\nu$ if and only if $\langle g \rangle_\nu$ is of type σ_ν . Thus, h factorizes through \mathbb{T}^{\min} if and only if $g \in \underline{\sigma} S_\omega(U, \mathcal{O}_K)$, if and only if $r_{\langle g \rangle}$ is (as a lift of \bar{r}) minimally ramified in S , if and only if the associated map $h' : R^{\text{crys}} \rightarrow \overline{\mathbb{Q}}_\ell$ factorizes through $R^{\min, \text{crys}}$. \square

Theorem 7.4. $R^{\min, \text{crys}}$ is finite flat over \mathcal{O}_K , so in particular there exists a characteristic-0 point of $\text{Spec } R^{\min, \text{crys}}$. Moreover, we have isomorphisms

$$R^{\min, \text{crys}} \cong R_{\text{red}}^{\min, \text{crys}} \cong \mathbb{T}^{\min}.$$

Proof. We first remark that $R^{\min, \text{crys}}/(\ell)$ is of finite cardinality, or, equivalently (by Nakayama's Lemma), that $R^{\min, \text{crys}}$ is finitely generated as a \mathcal{O}_K -module. This follows directly from [BLGGT14, Theorem 2.3.2], as we know that the local deformation rings $R^{\square, \chi_\nu, \text{crys}}(\bar{\rho}_\nu)$ and $R^{\square, \chi_\nu, \min}(\bar{\rho}_\nu)$ are smooth, hence correspond to irreducible components of $\text{Spec } R^{\square, \chi_\nu}(\bar{\rho}_\nu)$ on which the local lifts ρ_ν live.

Next, we remark that by Corollary 3.20 (together with the smoothness-results Lemma 5.4, Proposition 5.5, Proposition 3.21, the identity $\dim(\mathfrak{gl}_n^{c_\nu = -1}) = \frac{n(n+1)}{2}$ and Remark 7.6 below) there exists a presentation

$$R^{\min, \text{crys}} \cong \mathcal{O}_K[[X_1, \dots, X_m]]/(f_1, \dots, f_m).$$

for some $m \in \mathbb{N}_0$.

Using this presentation and the finiteness of $R^{\min, \text{crys}}/(\ell)$, it follows as in the proof of Theorem 3.7 of [KW09a] or of Lemma 2 in Böckle's Appendix to [Kha03] that $R^{\min, \text{crys}}$ is finite flat over \mathcal{O}_K , hence free and finitely generated over \mathcal{O}_K . This proves the first claim.

As a second step, we remark that any morphism $f : R^{\min, \text{crys}} \rightarrow \overline{\mathbb{Q}}_\ell$ factorizes over \mathbb{T}^{\min} : By [BLGGT14, Thm 2.3.1], such an f factorizes over the non-minimal Hecke algebra \mathbb{T} . Therefore Proposition 7.3 applies.

Now, $R^{\min, \text{crys}}[\frac{1}{\ell}] = R^{\min, \text{crys}} \otimes_{\mathcal{O}_K} \mathcal{K}$ is a finite \mathcal{K} -algebra, hence $R^{\min, \text{crys}}[\frac{1}{\ell}]$ is Artinian, see e.g. [AM69, Exercise 8.3]. Therefore, $R^{\min, \text{crys}}[\frac{1}{\ell}]$ can be decomposed into a product of finitely many local Artinian rings

$$R^{\min, \text{crys}} \left[\frac{1}{\ell} \right] \cong \bigoplus_{\mathfrak{p}} R^{\min, \text{crys}} \left[\frac{1}{\ell} \right]_{\mathfrak{p}}$$

and by [All16, Theorem 3.1.3] the tangent space $\mathfrak{p}/\mathfrak{p}^2$ of each $R^{\min, \text{crys}}[\frac{1}{\ell}]_{\mathfrak{p}}$ vanishes. Hence, $\mathfrak{p} = \mathfrak{p}^2$, and it follows from Nakayama's lemma that $\mathfrak{p} = 0$, i.e. that $R^{\min, \text{crys}}[\frac{1}{\ell}]_{\mathfrak{p}}$ is a field. Thus, $R^{\min, \text{crys}}[\frac{1}{\ell}]$ is a finite product of fields. The same is true for $\mathbb{T}^{\min}[\frac{1}{\ell}]$: As $\mathbb{T}^{\min}[\frac{1}{\ell}]$ is finitely-generated, its Jacobson radical equals its nilradical, which vanishes because \mathbb{T}^{\min} is reduced. Hence $\mathbb{T}^{\min}[\frac{1}{\ell}]$ is semisimple, i.e. a product of finitely many fields.

Consider the exact sequence

$$0 \rightarrow \ker(\varphi) \rightarrow R^{\min, \text{crys}} \xrightarrow{\varphi} \mathbb{T}^{\min} \rightarrow 0, \tag{10}$$

where φ denotes the canonical projection. It follows from the above observation about $R^{\min, \text{crys}}[\frac{1}{\ell}]$ and $\mathbb{T}^{\min}[\frac{1}{\ell}]$ together with Proposition 7.3 that $\varphi[\frac{1}{\ell}]$ is an isomorphism. Moreover, as both $R^{\min, \text{crys}}$ and \mathbb{T}^{\min} are finite free over \mathcal{O}_K , (10) is a split exact sequence of free \mathcal{O}_K -modules. Hence, $\ker(\varphi) = 0$ since $\varphi[\frac{1}{\ell}]$ is an isomorphism. This completes the proof of the theorem. \square

Corollary 7.5.

$$\begin{array}{ccc}
 R^{\text{crys}} & \longrightarrow & R^{\text{min,crys}} \\
 \downarrow & & \downarrow \\
 \mathbf{T} & \longrightarrow & \mathbf{T}^{\text{min}}
 \end{array}$$

is a pushout diagram.

Remark 7.6. We remark that for each $\nu \in \Omega_\infty$ the local deformation ring $L_\nu^+ R_{W(k_\lambda)}^{\square, \chi_\nu}(\bar{\tau}_{\lambda, \nu})$ is formally smooth of relative dimension $d_\nu^\square = \dim(\mathfrak{b}_n^{\text{der}})$: We get from Proposition 3.21 that $L_\nu^+ R_{W(k_\lambda)}^{\square, \chi_\nu}(\bar{\tau}_{\lambda, \nu})$ is formally smooth of relative dimension $\dim((\mathfrak{g}_n)^{c_\nu = -1}) = \dim(\mathfrak{g}_n^{c_\nu = -1})$, where c_ν is the non-trivial element of the decomposition group at ν . By construction (see Lemma 2.1.4 and Proposition 3.4.4 of [CHT08]), the image of $\bar{\tau}_\lambda(c_\nu)$ is not contained in $\text{GL}_n \times \text{GL}_1$. Moreover,

$$\mathfrak{m} \circ \bar{\tau}_\lambda(c_\nu) = \bar{\epsilon}_\ell^{1-n}(c_\nu) \delta_{F|F^+}^{\mu_m}(c_\nu) = (-1)^{\mu_m + p},$$

where $p = n + 1 \pmod{2} \in \mathbb{Z}/2\mathbb{Z}$, where ϵ_ℓ denotes the cyclotomic character (sending c_ν to -1), where $\delta_{F|F^+}$ denotes the non-trivial character of $\text{Gal}(F|F^+)$ and where μ_m is a suitable element of $\mathbb{Z}/2\mathbb{Z}$. As in [Tho12, Corollary 6.9], we get $\mu_m \equiv n \pmod{2}$, so we have $\mathfrak{m} \circ \bar{\tau}_\lambda(c_\nu) = -1$, independent of the parity of n . Using [CHT08, Lemma 2.1.3], this implies $\dim(\mathfrak{g}_n^{c_\nu = -1}) = \frac{n(n+1)}{2} = \dim(\mathfrak{b}_n^{\text{der}})$.

7.2 A $T = O$ -theorem

Now, let $\mathcal{E} \supset \mathcal{E}(U)$ be a number field with ring of integers $\mathcal{O}_\mathcal{E}$. For each $\lambda \in \text{Pl}_{\mathcal{E}}^{\text{fin}}$ such that $\ell := \ell(\lambda) \gg 0$, let us fix an F -big enough extension \mathcal{K}_λ of \mathcal{E}_λ and let us abbreviate

$$\mathbf{T} := {}^{\mathcal{O}_\mathcal{E}}\mathbf{T}_\omega^\mathcal{T}(U) \text{ and } \mathbf{T}_\lambda := \mathcal{O}_{\mathcal{K}_\lambda} \otimes_{\mathcal{O}_\mathcal{E}} \mathbf{T} \cong {}^{\mathcal{O}_{\mathcal{K}_\lambda}}\mathbf{T}_{\omega_\ell}^{\mathcal{T}_\ell}(U).$$

Observe the following about the isomorphism on the right hand side: Using that $\mathcal{S}_\omega(U)$ admits a basis of eigenforms, we can embed \mathbf{T} into a product of finitely many $\mathcal{O}_{\mathcal{E}(U)}$. Hence, \mathbf{T} is finitely generated as a \mathbb{Z} -module, hence as a \mathbb{Z} -algebra. It follows that there exists a Sturm-like bound $C \in \mathbb{N}$ such that \mathbf{T} is already generated by those $T_{F_w}^{(j)}$ with $\ell(w) \leq C$. Hence, using the compatibility from (9), we get

$$\mathcal{O}_{\mathcal{K}_\lambda} \otimes_{\mathcal{O}_\mathcal{E}} \mathbf{T} \cong \mathcal{O}_{\mathcal{K}_\lambda} \otimes_{\mathcal{O}_\mathcal{E}} {}^{\mathcal{O}_\mathcal{E}}\mathbf{T}_\omega^\mathcal{T}(U) \cong {}^{\mathcal{O}_{\mathcal{K}_\lambda}}\mathbf{T}_{\omega_\ell}^{\mathcal{T}_\ell}(U)$$

as long as $\ell > C$. Then we have:

Lemma 7.7. *For almost all λ (the failure set depending only on \mathbf{T}), \mathbf{T}_λ decomposes as a product of finitely many complete discrete valuation rings, finite over \mathbb{Z}_ℓ .*

Proof. First, we see that \mathbf{T} is an order in $\mathbf{T} \otimes_{\mathcal{O}_\mathcal{E}} \mathcal{E} \cong k_1 \times \dots \times k_m$, where the k_i denote suitable number fields (containing E) and the decomposition follows because $\mathbf{T} \otimes_{\mathcal{O}_\mathcal{E}} \mathcal{E}$ is reduced (as already remarked). Hence, \mathbf{T} is contained in the maximal order $\bigoplus_{i=1}^m \mathcal{O}_{k_i}$. It follows that there exists a suitable $N \in \mathbb{N}$ such that

- $\mathbf{T}[\frac{1}{N}] \cong \bigoplus_{i=1}^m \mathcal{O}_{k_i}[\frac{1}{N}]$;
- for any λ with $\ell(\lambda) \not\equiv N$, we have $\mathbf{T}_\lambda \cong \mathbf{T}[\frac{1}{N}]_\lambda := \mathcal{O}_{\mathcal{K}_\lambda} \otimes_{\mathcal{O}_\mathcal{E}} \mathbf{T}[\frac{1}{N}]$.

Thus, for those λ we get an isomorphism $\mathbf{T}_\lambda \cong \bigoplus_{i=1}^m \mathcal{O}_{\mathcal{K}_\lambda} \otimes_{\mathcal{O}_\mathcal{E}} \mathcal{O}_{k_i}$. As each factor itself is a product of complete discrete valuation rings (cf. e.g. [Ser79, Ch.2, §3, Theorem 1(ii)]), the lemma follows. \square

Because we assumed that \mathcal{E} contains all Hecke eigenvalues, in fact all the fields k_i in the above proof are equal to \mathcal{E} . Hence, for almost all ℓ , the above lemma implies that \mathbf{T}_λ is isomorphic to a product of finitely many copies of $\mathcal{O}_{\mathcal{K}_\lambda}$. Thus, we get

Corollary 7.8. *For almost all λ and all maximal ideals $\mathfrak{m} \subset \mathbf{T}_\lambda$, we have an isomorphism $\mathbf{T}_{\lambda, \mathfrak{m}} \cong \mathcal{O}_{\mathcal{K}_\lambda}$.*

7.3 An $R = R^{\min}$ -theorem

We retain all notation from the above and start with a preparatory corollary (to Corollary 5.13):

Corollary 7.9. *For almost all λ for which $\bar{\rho}_\lambda$ is absolutely irreducible, we have $R^{\text{crys}, (1, \dots, 1)} = R^{\text{crys}}$.*

Proof. Let $m := \max\{p \in \mathbb{N} \mid p \text{ prime}, \nu|p \text{ for some } \nu \in S\}$. Then, for all λ with $\ell(\lambda) > m^{n!}$, the claim follows directly from Corollary 5.13. \square

Moreover, we need a congruence argument: First, recall that the Hecke algebra ${}^{\mathcal{O}_\mathcal{K}}\mathbb{T}_\omega^{S_\ell}(U)$ acts semisimply on $\mathcal{S}_\omega(U)$, so the space $\mathcal{S}_\omega(U)$ decomposes into finitely many eigenspaces. For the following, let us consider *congruences*, by what we mean triples (H_1, H_2, ℓ) , where $H_1 \neq H_2$ are two Hecke eigenspaces and where ℓ is a rational prime such that there exists an isomorphism $\bar{\rho}_{f_1, \lambda_1} \otimes \bar{\mathbb{F}}_\ell \cong \bar{\rho}_{f_2, \lambda_2} \otimes \bar{\mathbb{F}}_\ell$ for some choice of forms $f_i \in H_i$ and of places λ_i of the corresponding coefficient fields fulfilling $\ell(\lambda_i) = \ell$.

Proposition 7.10. *There exist only finitely many such congruences in $\mathcal{S}_\omega(U)$.*

Proof. We easily see that a congruence (H_1, H_2, ℓ) corresponds to two distinct minimal prime ideals $\mathfrak{p}_{f_1}, \mathfrak{p}_{f_2}$ of \mathbf{T} for which there exists a maximal ideal $\mathfrak{m} \subset \mathbf{T}$ which contains ℓ , \mathfrak{p}_{f_1} and \mathfrak{p}_{f_2} . It follows from the finite flatness of \mathbf{T} over \mathbb{Z} that for given eigenforms f_1, f_2 there exist only finitely many maximal ideals containing \mathfrak{p}_{f_1} and \mathfrak{p}_{f_2} . Thus, the claim follows immediately from the finite-dimensionality of the space of automorphic representations of given level and weight. \square

Theorem 7.11. *For almost all λ for which $\bar{\rho}_\lambda$ is absolutely irreducible, we have*

$$R^{\text{crys}} \cong R^{\min, \text{crys}}.$$

Proof. We apply the proof of Theorem 7.4, where we replace $R^{\min, \text{crys}}$ by R^{crys} and \mathbb{T}^{\min} by $R^{\min, \text{crys}}$. In the following, we only point out those steps which need additional explanation:

The part that $R^{\text{crys}}/(\ell)$ is finite follows as above, but we have to assure that Guerberoff's Theorem 4.1 remains applicable, at least for almost all ℓ . This is a direct consequence of Corollary 7.9.

The existence of a presentation

$$R^{\text{crys}} \cong \mathcal{O}_\mathcal{K}[[X_1, \dots, X_m]]/(f_1, \dots, f_m);$$

follows by Corollary 3.20 (as in the proof of Theorem 7.4, but referring to Theorem 5.1 instead of Proposition 5.5).

Moreover, we claim that for almost all λ , any morphism $R^{\text{crys}} \rightarrow \bar{\mathbb{Q}}_\ell$ factorizes over $R^{\min, \text{crys}}$. Using automorphy lifting, this claim can be restated as follows: For almost all λ , the following holds: For

any automorphic form g whose associated Galois representation $\rho_{g,\lambda}$ reduces to $\bar{\rho}_\lambda$, $\rho_{g,\lambda}$ is a minimally ramified lift of $\bar{\rho}_\lambda$. Now, let λ be a place such that this statement fails. Then, as there always exists a minimally ramified lift of $\bar{\rho}_{g,\lambda}$ with a corresponding automorphic form f (cf. Theorem 7.4), we get a congruence $(\mathcal{O}_K \mathbb{T}_\omega^{S_\ell}(U) \cdot f, \mathcal{O}_K \mathbb{T}_\omega^{S_\ell}(U) \cdot g, \ell(\lambda))$. Thus, the claim follows from Proposition 7.10.

The rest of the proof goes through verbatim. \square

We need a technical lemma:

Lemma 7.12. *Let $R_1, R_2 \in \mathcal{C}_\Lambda$ and assume that there exist suitable $\alpha, \beta, m, h \in \mathbb{N}_0$ such that*

1. $R_1 \cong \Lambda[[y_1, \dots, y_{\beta+h}]]/(g_1, \dots, g_\beta)$;
2. $R_2 \cong \Lambda[[z_1, \dots, z_{m+h}]]$;
3. $R_1[[x_1, \dots, x_{\alpha+m}]]/(f_1, \dots, f_\alpha) \cong R_2$.

Then, R_1 has only one irreducible component.

Proof. Let us rewrite the last condition as

$$\Lambda[[x_1, \dots, x_{\alpha+m}, y_1, \dots, y_{\beta+h}]]/(f_1, \dots, f_\alpha, g_1, \dots, g_\beta) \cong \Lambda[[z_1, \dots, z_{m+h}]].$$

It follows immediately by [Ser79, Proposition 22, Part c) \Rightarrow a)], that $f_1, \dots, f_\alpha, g_1, \dots, g_\beta$ is a subset of regular system of parameters of $\Lambda[[x_1, \dots, x_{\alpha+m}, y_1, \dots, y_{\beta+h}]]$. It follows easily that then g_1, \dots, g_h is a subset of a regular system of parameters of $\Lambda[[y_1, \dots, y_{\beta+h}]]$, and so (by [Ser79, Proposition 22, Part a) \Rightarrow c)], it follows that R_1 is regular. As any regular local ring is an integral domain, the claim follows. \square

Let us close with the following corollary (to Theorem 7.11), giving a local $R = R^{\min}$ result:

Corollary 7.13. *For almost all λ , $R^{\square, \chi_\nu, \min}(\bar{\rho}_{\lambda, \nu}) \cong R^{\square, \chi_\nu}(\bar{\rho}_{\lambda, \nu})$ holds for any $\nu \in S$.*

Proof. Assume that λ is such that Theorem 7.11 holds. Moreover, fix a place $\nu_0 \in S$ and write $R^{\square_{S_\ell}, (S-\nu_0)-\min, \text{crys}}$ for the ring parametrizing the same (global) lifts as $R^{\square_{S_\ell}, \min, \text{crys}}$, except that we don't impose any condition at ν_0 at all. Then we have isomorphisms

$$R^{\text{loc}, \chi, (S-\nu_0)-\min, \text{crys}}[[x_1, \dots, x_a]]/(f_1, \dots, f_b) \cong R^{\square_{S_\ell}, (S-\nu_0)-\min, \text{crys}} \cong R^{\square_{S_\ell}, \min, \text{crys}}$$

where the first isomorphism (as well as the identity $a - b = \#S_\ell - 1$) follows from Proposition 3.19. We know that all local deformation rings occurring as tensor factors in the definition $R^{\text{loc}, \chi, (S-\nu_0)-\min, \text{crys}}$ (except for possibly at ν) are formally smooth, and we also know their dimension:

$$\dim_W(R_\nu^{\square, \chi_w, \mathcal{D}_w}) = \begin{cases} \frac{n(n+1)}{2} & \text{if } w|\infty, \mathcal{D}_w = \emptyset; \\ n^2 + [F : \mathbb{Q}] \frac{n(n-1)}{2} & \text{if } w|\ell, \mathcal{D}_w = \text{crys}; \\ n^2 & \text{if } w \in (S - \nu), \mathcal{D}_w = \min; \end{cases}$$

By our $R^{\min} \cong T^{\min} \cong \mathcal{O}$ -results from the previous two sections, which hold again after ruling out finitely many λ , also $R^{\square_{S_\ell}, \min, \text{crys}}$ is formally smooth of dimension $(n^2 + 1)\#S_\ell - 1$. Thus, we get that

$$R^{\square, \chi_{\nu_0}}(\bar{\rho}_{\nu_0})[[x_1, \dots, x_{\alpha+m}]]/(f_1, \dots, f_\alpha) = R^{\square, \min, \text{crys}} = \mathcal{O}_K[[x_1, \dots, x_{(n^2+1)\#S_\ell-1}]]$$

with $m = (n^2 + 1)\#\mathcal{S}_\ell - 1 - n^2$. Now, by Theorem 5.1, $R^{\square, \chi_{\nu_0}}(\bar{\rho}_{\nu_0})$ is a complete intersection ring, i.e. we can write

$$R^{\square, \chi_{\nu_0}}(\bar{\rho}_{\nu_0}) = \mathcal{O}_K[[y_1, \dots, y_{\beta+h}]]/(g_1, \dots, g_\beta)$$

with $h = \dim R^{\square, \chi_{\nu_0}}(\bar{\rho}_{\nu_0}) = n^2$. By Lemma 7.12 applied to $\Lambda = \mathcal{O}_K$, $R_1 = R^{\square, \chi_{\nu_0}}(\bar{\rho}_{\nu_0})$ and $R_2 = R^{\square, \min, \text{crys}}$, we see that $R^{\square, \chi_{\nu_0}}(\bar{\rho}_{\nu_0})$ has only one irreducible component and hence we must have $R^{\square, \chi_{\nu_0, \min}}(\bar{\rho}_{\nu_0}) \cong R^{\square, \chi_{\nu_0}}(\bar{\rho}_{\nu_0})$. \square

8 Unobstructedness for RACSDC automorphic representations

We are now in a position to state and prove our main result. For this, let π be a RACSDC automorphic representation of $\text{GL}_n(\mathbb{A}_F)$ with ramification set S . By passing to a unitary avatar $\Pi \subset \mathcal{S}_\omega(U)$ (for a suitable weight ω and level $U = U_S$), we can attach the compatible system $\mathcal{R}_\pi = (\bar{r}_\lambda)_{\lambda \in \Lambda_{\mathcal{E}_\Pi}^1}$. Here, \mathcal{E}_Π denotes the number field generated by all Hecke eigenvalues of Π , $\Lambda_{\mathcal{E}_\Pi}^1 \subset \text{Pl}_{\mathcal{E}_\Pi}$ denotes the set of places for which \bar{r}_λ is absolutely irreducible and we assume the following:

Assumption 8.1. 1. **(Irreducibility):** The set $\Lambda_{\mathcal{E}_\Pi}^1 \subset \text{Pl}_{\mathcal{E}_\Pi}$ has Dirichlet density 1;

2. **(No consecutive weights):** The multisets of Hodge-Tate weights HT_τ of the system \mathcal{R}_π fulfill (for all embeddings τ) the condition from Theorem 5.2: If two numbers a, b occur in HT_τ , then $|a - b| \neq 1$.

We stress that we understand the first part as a general conjecture on Galois representations attached to RACSDC representations (so, in particular, we assume that this is correct independently of the choice of F or π), while the second part puts a constraint on our choice of π . We also have the following:

Remark 8.2. The first part of Assumption 8.1 is known to hold e.g. if π is extremely regular [BLGGT14] or if $n \leq 5$ [CG13]. Results in this direction are also contained in [PT15], but they are not directly applicable to our situation. We also remark that all entries in the ℓ -adic system $(\rho_{\pi, \lambda})_{\lambda \in \text{Pl}_{\mathcal{E}(U)}}$ are expected (by cuspidality of π) to be absolutely irreducible and that this, using suitable modularity lifting theorems, is expected to imply absolute irreducibility of $\bar{\rho}_{\pi, \lambda}$ for almost all λ . An established result in this direction is that absolute irreducibility of the ℓ -adic system implies absolute irreducibility of $\bar{\rho}_{\pi, \lambda}$ except for a failure set of Dirichlet density 0, see [PSW16].

Our main result is now as follows:

Theorem 8.3. *Presuming Assumption 8.1, there exists a subset $\Lambda_{\mathcal{E}_\Pi}^0 \subset \Lambda_{\mathcal{E}_\Pi}^1$ of Dirichlet density 1 such that the functor $D_{S_\ell, W(k_\lambda)}^{\square_{S_\ell, \chi}}(\bar{r}_\lambda)$ is globally unobstructed whenever $\lambda \in \Lambda_{\mathcal{E}_\Pi}^0$.*

As a first step towards the proof, let us consider the following simplifying assumption:

Assumption 8.4. 1. $F|F^+$ is unramified at all finite places and, in case n is even, then also $\frac{n}{2}[F^+ : \mathbb{Q}]$ is even;

2. each place ν of F^+ which lies below S splits in $F|F^+$ as, say, $\tilde{\nu}\tilde{\nu}^c$; (For archimedean places, this condition is automatically fulfilled, so we can replace S by $S \sqcup \Omega_\infty$ without loss of generality.)

3. for each place ν of F^+ which lies below S , the Weil-Deligne representation (r_ν, W_ν) attached to Π has unramified underlying Weil-representation r_ν .

Remark that the third part can be characterized as follows: The ℓ -adic representation $r_{\Pi,\lambda}$ is at ν a minimally ramified deformation of $\bar{\tau}_{\Pi,\lambda}$. (As the system associated to Π is *compatible*, this does not depend on the choice of $\lambda \in \Lambda_{\mathcal{E}\Pi}^1$.) Now, consider the following (seemingly weaker) variation of Theorem 8.3:

Theorem 8.5. *Presuming Assumptions 8.1 and 8.4, there exists a subset $\Lambda_{\mathcal{E}\Pi}^0 \subset \Lambda_{\mathcal{E}\Pi}^1$ of Dirichlet density 1 such that the functor $D_{S_\ell, W(k_\lambda)}^{\square_{S_\ell, \chi}}(\bar{\tau}_\lambda)$ is globally unobstructed whenever $\lambda \in \Lambda_{\mathcal{E}\Pi}^0$.*

Proof (that Theorem 8.5 implies Theorem 8.3). It is a standard argument (cf. e.g. the proof of [CHT08, Theorem 4.4.2]) that there exists a finite solvable extension $F_1^+|F^+$ of totally real fields such that the extension $F_1 = F_1^+.F|F_1^+$ and the compatible family associated to the base change Π_{F_1} of Π to F_1 fulfill Assumption 8.4. Thus, referring to Lemma 4.8 and eliminating the finitely many places λ which divide the index $[F_1^+ : F^+]$, we see that Assumption 8.4 can be included in the statement of Theorem 8.3 without causing loss of generality. \square

Consequently, the remainder of this section is devoted to the proof of Theorem 8.5. For better comprehension, let us give an overview of the strategy of the proof: We want to arrange for a situation where the framework of Section 4 is applicable, i.e. we want to consider suitable field extensions $L_{(\lambda)}^+$ for as many λ as possible such that Theorem 4.2 implies the vanishing of the dual Selmer groups of the base changed functors ${}^{L_{(\lambda)}^+}D_{S_\ell \langle L_{(\lambda)}^+ \rangle, W(k_\lambda)}^{\square_{S_\ell \langle L_{(\lambda)}^+ \rangle, \chi}}(\bar{\tau}_\lambda)$. This application of Theorem 4.2 happens in Theorem 8.12 below. By a careful choice of the extensions $L_{(\lambda)}^+$, we ensure that the potential unobstructedness arguments of Section 4 apply and yield the vanishing of the dual Selmer groups of the non-base changed functor. The local parts of the unobstructedness-condition then follow directly from the material in Section 5.2, allowing us to conclude the statement of Theorem 8.5. The crucial property we have to impose on the extension $L_{(\lambda)}^+$ is *procurability* (Definition 8.7), i.e. that the deformation ring $R_{S_\ell \langle L_{(\lambda)}^+ \rangle, \mathcal{O}_{K_\lambda}}^{\chi|\text{Gal}_{L^+}, \text{crys}}(\bar{\tau}_\lambda|\text{Gal}_{L^+})$ is isomorphic to \mathcal{O}_{K_λ} (corresponding to condition $(R = T)$ in Section 4). It is the content of Theorem 8.8 that for a set of places of Dirichlet density 1 we can find such suitable procurable extensions. This, in turn, is established by studying the seemingly weaker condition of \star -procurability (see the list $\star_1 - \star_5$ below), which is proved to imply procurability almost everywhere (see Claim 1 below). By an argument based on Chebotarev's density theorem (and postponed to Appendix A), we can conclude that for a density-1 set we can find such \star -procurable extensions of 2-power degree.

8.1 Proof of Theorem 8.5

Let us begin with some preparatory definitions:

Definition 8.6. A totally real, finite extension L^+ of F^+ is called *pre-admissible* if the extension $L^+|F^+$ is Galois and solvable and if $L := F.L^+$ is unramified over L^+ at every finite place.

We remark that these conditions are designed to capture the following: If L^+ is pre-admissible, then there exists a unitary group H over L^+ (as considered in Section 6.2) and a unitary avatar Π_L on $H(\mathbb{A}_{L^+})$ of the base change π_L of π to L .

For the following, let \mathcal{E} be a number field containing $\mathcal{E}(U)$ and let L^+ be pre-admissible.

Definition 8.7. A place $\lambda \in \Lambda_{\mathcal{E}}^1$ is L^+ -*procurable* if the following two conditions are fulfilled:

- P.1) The restriction of $\bar{\rho}_\lambda$ to Gal_L remains absolutely irreducible;
- P.2) There exists an L -big enough extension \mathcal{K}_λ of \mathcal{E}_λ such that there is an isomorphism

$$L^+ R_{S_\ell, \mathcal{O}_{\mathcal{K}_\lambda}}^{\chi, \text{crys}}(\bar{\tau}_\lambda) := R_{S_\ell(L^+), \mathcal{O}_{\mathcal{K}_\lambda}}^{\chi | \text{Gal}_{L^+}, \text{crys}}(\bar{\tau}_\lambda | \text{Gal}_{L^+}) \cong \mathcal{O}_{\mathcal{K}_\lambda}. \quad (11)$$

We remark that the first condition is rather harmless and affects only a failure set of Dirichlet density 0, cf. Assumption 8.1. We also remark that in the second condition, we consider $\bar{\tau}_\lambda$ as a representation with values in the residue field $k_{\mathcal{O}_{\mathcal{K}_\lambda}}$ of $\mathcal{O}_{\mathcal{K}_\lambda}$ instead of k_λ .

With respect to a pre-admissible extension L^+ , define $\text{Proc}(L^+) \subset \Lambda_{\mathcal{E}}^1$ as the subset of those λ which are L^+ -procurable. Then we have:

Theorem 8.8. *There exists a nested sequence $F^+ = L_0^+ \subset L_1^+ \subset \dots$ of pre-admissible extensions of F^+ such that*

$$\lim_{i \rightarrow \infty} \delta \left(\bigcup_{j=1}^i \text{Proc}(L_j^+) \right) = 1, \quad (12)$$

where $\delta(\Delta)$ denotes the density of those rational primes q for which each $\lambda \in \text{Pl}_{\mathcal{E}}$ above q fulfills $\lambda \in \Delta$.

Proof. Let us first introduce another notation: With respect to a pre-admissible extension L^+ , we say that $\lambda \in \Lambda_{\mathcal{E}}^1$ is L^+ - \star -*procurable*, if the following list is met (where, as usual, we abbreviate $\ell = \ell(\lambda)$):

- \star_1) ℓ is not divisible by any element of S ;
- \star_2) ℓ is unramified in the extension $L|\mathbb{Q}$;
- \star_3) all places of L above S_ℓ are split in the extension $L|L^+$;
- \star_4) the base change π_L of π to L remains cuspidal;
- \star_5) if $\nu \in \text{Pl}_A$ lies above S , then π_L admits a non-trivial Iwahori fix-vector.

As above, this defines a subset $\text{Proc}^*(L^+) \subset \Lambda_{\mathcal{E}}^1$. (Observe that condition \star_4 does not depend on λ , but we intentionally include it in the list. So, if Π_L fails to be cuspidal, we have $\text{Proc}^*(L^+) = \emptyset$.)

Claim 1: $\text{Proc}^*(L^+) - \text{Proc}(L^+)$ is finite.

Proof of Claim 1. We can suppose that $\text{Proc}^*(L^+)$ is not empty (otherwise the claim is trivially true), so in particular that π_L is a RACSDC representation and there exists a unitary group and an avatar Π_L over L . Now, for each $\lambda \in \text{Proc}^*(L^+)$ we pick an L -big enough field extension \mathcal{K}_λ of \mathcal{E}_λ . We consider the complex Hecke algebra ${}^{\mathcal{O}_{\mathcal{E}}}\mathbf{T}_{\omega}^S(U)$ and the ℓ -adic model $\mathbb{T} := {}^{\mathcal{O}_{\mathcal{E}}}\mathbb{T}_{\omega}^{S_\ell}(U)$.

Write $\Pi_L = \langle f \rangle$ for the unitary avatar of the base change of π to L and for a suitable choice $f \in \mathcal{S}_{\omega}(U)$. We see that $\bar{\rho}_\lambda | \text{Gal}_L$ equals the reduction of the representation attached to the maximal ideal $\mathfrak{m} = \ker(\varphi_{f^{(\lambda)}}) \subset \mathbb{T}$ by Proposition 6.7, where $f^{(\lambda)}$ is the ℓ -adic model of f .

Recalling that we presume Assumption 8.1, we see easily that the preconditions of Theorem 7.4 hold for almost all of these choice of $L|L^+$, $\ell = \ell(\lambda)$, $U, \omega, \mathcal{E}(U), \mathcal{K}_\lambda$ and \mathfrak{m} : The main issue is the adequateness of $\bar{\rho}(\text{Gal}_{F(\zeta_\ell)})$, which follows from [BLGGT14, Proposition 2.1.2] as long as $\ell > 2(n+1)$. Thus, the

desired isomorphism (11) follows for almost all λ in $\text{Proc}^*(L^+)$ by Corollary 7.8. This completes the proof of the claim. \clubsuit


Consequently, it suffices to show that there exists a nested sequence $F^+ = L_0^+ \subset L_1^+ \subset \dots$ of pre-admissible extensions of F^+ such that equation (12) holds with Proc^* instead of Proc . For the construction of these extensions, we define the set

$$\Theta_F := \{d \in \mathbb{N} \mid \sqrt{d} \notin F, \text{ the base change } \pi \rightsquigarrow \pi_{F(\sqrt{d})} \text{ remains cuspidal} \}.$$

This set is not empty (cf. [AC89, Thm 4.2]), so choose a $d_1 \in \Theta_F$ and take $L_1^+ = F^+(\sqrt{d_1})$.

Claim 2: L_1^+ is pre-admissible.

Proof of Claim 2. The extension $L_1^+|F^+$ is automatically Galois and solvable because $[L_1^+ : F^+] = 2$. Thus we are left to check that $L_1|L_1^+$ is unramified everywhere. For this, we use the identity of the discriminants $\Delta_{L_1|F^+} = \Delta_{L_1^+|F^+} \Delta_{F|F^+} = \Delta_{L_1^+|F^+}$ (cf. [Jan96, Exercise 3 on p. 51]). Consider the diagram of field extensions



If now w is a prime of L_1^+ that ramifies in (3), then the prime v of F^+ which lies below w must ramify in the extension $L_1|F^+$. But then v divides $\Delta_{L_1^+|F^+} = \Delta_{L_1|F^+}$, i.e. v ramifies in (1). This implies that v has ramification index 4 in the extension $L_1|F^+$. But in (2), v is unramified by the prerequisites, so it can at most ramify in (4), yielding a ramification index of 2 in $L_1|F^+$. This contradicts the assumption that w ramifies in (3). \clubsuit

Claim 3: $\delta(\text{Proc}^*(F_1^+)) \geq \frac{1}{2}$.

Proof of Claim 3. We check which λ fail the list $\star_1 - \star_5$:

- Concerning \star_1 and \star_2 , we have to exclude the finitely many places λ for which $\ell(\lambda)$ is not coprime to S or ramifies in $L_1^+|\mathbb{Q}$;
- By an estimation based on Chebotarev's density theorem (postponed as Lemma A to the appendix), the density of those ℓ which fulfill the condition that all primes of L_1^+ above ℓ are split in the extension $L_1|L_1^+$ is at least $\frac{1}{2}$;
- Condition \star_4 is universally fulfilled by our choice of L_1^+ ;
- Concerning condition \star_5 , we remark that by local-global compatibility (cf. [CH13, Theorem 1.4] and the references therein) π_L admits an Iwahori-fixed vector if $\rho|_{\text{Gal}_L}$ has unipotent ramification at ν [Wed08, (4.3.6) Proposition]. Thus, condition \star_5 follows immediately from Assumption 8.4.

\clubsuit

For the next tower step we take $F_2^+ = F_1^+(\sqrt{d_2})$ for some $d_2 \in \Theta_{F_2}$. It is again easy to check that $\Theta_{F_2} \neq \emptyset$ and that F_2^+ is pre-admissible. As in the proof of Claim 3, the statement of Lemma A implies $\delta(\text{Proc}^*(F_2^+)) \geq \frac{3}{4}$. Iterating this construction of quadratic extensions we end up with a nested sequence of pre-admissible fields F_j^+ such that

$$\delta \left(\bigcup_{j=1}^i \text{Proc}^*(L_j^+) \right) \geq \delta(\text{Proc}^*(L_i^+)) \geq 1 - \frac{1}{2^i} \xrightarrow{i \rightarrow \infty} 1.$$

Together with Claim 1, this concludes the proof of Theorem 8.8. \square

We now give a slight variant of the above:

Definition 8.9. With regard to a pre-admissible extension L^+ of F^+ , we say that $\lambda \in \Lambda_{\mathcal{E}}^1$ is L^+ - $\#$ -procurable if the restriction of $\bar{\rho}_\lambda$ to Gal_L (with $L = F.L^+$) remains absolutely irreducible and if there is an isomorphism

$$L^+ R_\lambda^{\square, \chi, \text{crys}} \cong W(k_\lambda)[[x_1, \dots, x_u]], \quad (13)$$

where $L^+ R_\lambda^{\square, \chi, \text{crys}} = L^+ R_{S_\ell, W(k_\lambda)}^{\square, \chi, \text{crys}}(\bar{\rho}_\lambda)$ and $u = \dim(\mathfrak{g}_n^{\text{der}}) = n^2$. The set of all λ which are L^+ - $\#$ -procurable is denoted by $\text{Proc}^\#(L^+)$.

Corollary 8.10. *There exists a nested sequence $F^+ = L_0^+ \subset L_1^+ \subset \dots$ of pre-admissible extensions of F^+ such that*

$$\lim_{i \rightarrow \infty} \delta \left(\bigcup_{j=1}^i \text{Proc}^\#(L_j^+) \right) = 1.$$

Proof. For $i \in \mathbb{N}$, denote $\Delta_i = \bigcup_{j \leq i} \text{Proc}(L_j^+)$. Also fix for each $\lambda \in \Delta_i$ some $j \leq i$ such that $\lambda \in \text{Proc}(L_j^+)$. Denote the corresponding field extension from the proof of Theorem 8.8 by $L_{(\lambda)} = L_{(\lambda)}^+.F$. By Theorem 8.8, for such a $\lambda \in \Delta_i$ we have the identity (11) for a suitable extension $\mathcal{O}_{\mathcal{K}_\lambda}$ of $W(k_\lambda)$. The third part of Proposition 3.17 then yields

$$L_{(\lambda)}^+ R_{S_\ell, \mathcal{O}_{\mathcal{K}_\lambda}}^{\square, \chi, \text{crys}}(\bar{\rho}_\lambda) \cong \mathcal{O}_{\mathcal{K}_\lambda}[[x_1, \dots, x_u]].$$

Thus, we can use Lemma 3.6 (and, if necessary, Remark 3.10) to deduce the desired isomorphism (13). \square

Corollary 8.11. *There exists a subset $\Lambda_{\mathcal{E}}^2 \subset \Lambda_{\mathcal{E}}^1$ of Dirichlet density 1 such that for each $\lambda \in \Lambda_{\mathcal{E}}^2$ there exists a finite, totally real extension $L_{(\lambda)}^+$ of F and an isomorphism*

$$L_{(\lambda)}^+ R_{S_\ell \langle L_{(\lambda)}^+ \rangle, W(k_\lambda)}^{\square_{S_\ell \langle L_{(\lambda)}^+ \rangle}, \chi, \text{crys}}(\bar{\rho}_\lambda) \cong W(k_\lambda)[[x_1, \dots, x_{w(\lambda)}]]$$

with $w(\lambda) = (n^2 + 1) \cdot \#S_\ell \langle L_{(\lambda)}^+ \rangle - 1$.

Proof. This follows directly from Proposition 3.17. \square

Next, we will apply the framework of Section 4 to the attained λ :

Theorem 8.12. *There exists a cofinite subset $\Lambda_{\mathcal{E}}^3 \subset \Lambda_{\mathcal{E}}^2$ such that the following holds: Let $\lambda \in \Lambda_{\mathcal{E}}^3$ and $L_{(\lambda)}^+$ the corresponding extension from Corollary 8.11. Then the functors*

$$L_{(\lambda)}^+ D_{S_{\ell}(L_{(\lambda)}^+), W(k_{\lambda})}^{\square, \chi, \min}(\bar{\tau}_{\lambda}) \quad \text{and} \quad L_{(\lambda)}^+ D_{S_{\ell}(L_{(\lambda)}^+), W(k_{\lambda})}^{\square, \chi}(\bar{\tau}_{\lambda})$$

have vanishing dual Selmer group.

Proof. We start with the **min**-case: When applying the framework, we take for **sm** the condition parametrizing arbitrary deformations, for **crys** the condition parametrizing FL-crystalline deformations (cf. Section 5.3) and for **min** the condition parametrizing minimally ramified deformations (cf. Section 5.4). Moreover, we take $\chi = \epsilon_{\ell}^{1-n} \delta_{F|F^+}^{n(\bmod 2)}$. Let us now check the following list of conditions (and we abbreviate $L^+ = L_{(\lambda)}^+$ as we check this for a fixed $\lambda \in \Lambda_{\mathcal{E}}^3$):

1. (**sm**/ k): As we took for **sm** the unrestricted deformation condition, we have to check that for each $\nu \in \Omega_{\ell}$ the functor $L_{W(k_{\lambda})}^+ D_{W(k_{\lambda})}^{\square, \chi_{\nu}}(\bar{\tau}_{\lambda, \nu})$ is representable and that the representing object is formally smooth of relative dimension

$$d_{\nu}^{\square, \text{sm}} = \dim(\mathfrak{g}_n^{\text{der}})([L_{\bar{\nu}} : \mathbb{Q}_{\ell}] + 1) = n^2([L_{\bar{\nu}} : \mathbb{Q}_{\ell}] + 1) = n^2([L_{\nu}^+ : \mathbb{Q}_{\ell}] + 1).$$

(This also amounts to the vanishing of the error terms δ_{ν} in Theorem 4.2.)

Check: Representability was already remarked in Section 3. For the remaining claim, we first refer to Proposition 6.1 in order to get an isomorphism

$$L_{\bar{\nu}}^+ R_{W(k_{\lambda})}^{\square, \chi_{\nu}}(\bar{\tau}_{\lambda, \nu}) \cong L_{\bar{\nu}} D_{W(k_{\lambda})}^{\square, \chi_{\nu}}(\bar{\rho}_{\lambda, \nu}).$$

Now the claim follows from Theorem 5.2.

2. (**crys**): For each $\nu \in \Omega_{\ell}$, the subfunctor

$$L_{W(k_{\lambda})}^+ D_{W(k_{\lambda})}^{\square, \chi_{\nu}, \text{crys}}(\bar{\tau}_{\lambda, \nu}) \hookrightarrow L_{W(k_{\lambda})}^+ D_{W(k_{\lambda})}^{\square, \chi_{\nu}}(\bar{\tau}_{\lambda, \nu})$$

is relatively representable and the representing object is formally smooth of relative dimension $d_{\nu}^{\square, \text{crys}} = \dim(\mathfrak{g}_n^{\text{der}}) + (\dim(\mathfrak{g}_n^{\text{der}}) - \dim(\mathfrak{b}_n^{\text{der}}))[L_{\nu}^+ : \mathbb{Q}_{\ell}]$, where \mathfrak{b}_n denotes the Lie algebra of a Borel subgroup of \mathcal{G}_n .

Check: By definition, we have

$$L_{\bar{\nu}}^+ R_{W(k_{\lambda})}^{\square, \chi_{\nu}, \text{crys}}(\bar{\tau}_{\lambda, \nu}) \cong L_{\bar{\nu}} D_{W(k_{\lambda})}^{\square, \chi_{\nu}, \text{crys}}(\bar{\rho}_{\lambda, \nu}).$$

Thus, the claim follows from Lemma 5.4.

3. (**min**): For each $\nu \in S$, the subfunctor

$$L_{W(k_{\lambda})}^+ D_{W(k_{\lambda})}^{\square, \chi_{\nu}, \min}(\bar{\tau}_{\lambda, \nu}) \hookrightarrow L_{W(k_{\lambda})}^+ D_{W(k_{\lambda})}^{\square, \chi_{\nu}}(\bar{\tau}_{\lambda, \nu})$$

is relatively representable and the representing object is formally smooth of relative dimension $d_{\nu}^{\square, \min} = \dim(\mathfrak{g}_n^{\text{der}})$.

Check: Again, by definition, we have

$$L_{\bar{\nu}}^+ R_{W(k_{\lambda})}^{\square, \chi_{\nu}, \min}(\bar{\tau}_{\lambda, \nu}) \cong L_{\bar{\nu}} D_{W(k_{\lambda})}^{\square, \chi_{\nu}, \min}(\bar{\rho}_{\lambda, \nu}).$$

Thus, the claim follows from Proposition 5.5.

4. (∞): For each $\nu \in \Omega_\infty$, the local deformation ring $L_\nu^+ R_{W(k_\lambda)}^{\square, \chi_\nu}(\bar{\tau}_{\lambda, \nu})$ is formally smooth of relative dimension $d_\nu^\square = \dim(\mathfrak{b}_n^{\text{der}})$.

Check: This was already used, see Remark 7.6.

5. (**Presentability**): Consider the ring

$$L^+ R^{\text{loc, min}} := \widehat{\bigotimes_{\nu \in S_\ell \langle L^+ \rangle} L^+ \tilde{R}_\nu}$$

with $L^+ \tilde{R}_\nu = L_\nu^+ D_{W(k_\lambda)}^{\square, \chi_\nu, \text{min}}(\bar{\tau}_{\lambda, \nu})$ if $\nu \in S$ and $L_\nu^+ D_{W(k_\lambda)}^{\square, \chi_\nu}(\bar{\tau}_{\lambda, \nu})$ otherwise. Then, there exists a presentation

$$L^+ R_{S_\ell, W(k_\ell)}^{\square, \chi}(\bar{\tau}_\lambda) \cong L^+ R^{\text{loc, min}}[[X_1, \dots, X_a]]/(f_1, \dots, f_b)$$

with $a - b = (\#S_\ell \langle L^+ \rangle - 1) \cdot \dim(\mathfrak{g}_n^{\text{ab}})$.

Check: This is the content of Proposition 3.19, but we have to check Assumption 3.18. As $\mathfrak{g}_n^{\text{der}} = \mathfrak{gl}_n$, this condition holds by Corollary 3.24 for almost all λ .

6. ($R = T$): The ring $L^+ R_{S_\ell, W(k_\ell)}^{\square, \chi, \text{min, crys}}(\bar{\tau}_\lambda)$ is formally smooth of relative dimension $r_0 = \dim(\mathfrak{g}) \cdot \#S_\ell \langle L^+ \rangle - \dim(\mathfrak{g}^{\text{ab}})$.

Check: This follows from Corollary 8.11.

We see that the general requirements of Theorem 4.2 are met, so let us check the additional requirements of part 2. of Theorem 4.2:

a) The condition $\ell \gg 0$ can be achieved by leaving out finitely many λ ;

b) The vanishing of $H^0(\text{Gal}_{L^+}, \mathfrak{g}_n^{\text{der}, \vee})$ can be checked by observing

$$H^0(\text{Gal}_{L^+}, \mathfrak{g}_n^{\text{der}, \vee}) \subset H^0(\text{Gal}_L, \mathfrak{g}_n^{\text{der}, \vee}) \cong H^0(\text{Gal}_L, \mathfrak{gl}_n^\vee),$$

as the adjoint representation of Gal_L on $\mathfrak{g}_n^{\text{der}}$ (via $\bar{\tau}_\lambda$) corresponds to the adjoint representation of Gal_L on \mathfrak{gl}_n (via $\bar{\rho}_\lambda$), cf. [CHT08, Section 2.1]. Thus, the desired vanishing follows for almost all λ by Corollary 3.24.

c) For $\nu \in S \langle L^+ \rangle$, $\dim(L_{\lambda, \nu}) = h^0(\text{Gal}_{L_\nu^+}, \mathfrak{g}_n^{\text{der}})$: As ν is split, Proposition 6.1 yields $h^0(\text{Gal}_{L_\nu^+}, \mathfrak{g}_n^{\text{der}}) = h^0(\text{Gal}_{L_{\bar{\nu}}}, \mathfrak{gl}_n^{\text{der}})$, where the action on \mathfrak{gl}_n is via $\bar{\rho}_{\lambda, \bar{\nu}}$. The claim thus follows from [CHT08, Corollary 2.4.21].

The finitely many exclusions which occurred in the above items are now the places we must exclude from $\Lambda_\mathcal{E}^2$ to get $\Lambda_\mathcal{E}^3$. This finishes the first part, i.e. that $L_{(\lambda)}^+ D_{S_\ell \langle L_{(\lambda)}^+ \rangle, W(k_\lambda)}^{\square, \chi, \text{min}}(\bar{\tau}_\lambda)$ has vanishing dual Selmer group.

Concerning the second statement (i.e. the claimed vanishing of the non-minimal dual Selmer group) we first note that on each level $L_{(\lambda)}^+$ we can apply the $R = R^{\text{min}}$ -result of Corollary 7.13, yielding the desired vanishing except for a finite failure set. In other words: Fix a place λ' , then we have

$$L_{(\lambda)}^+ D_{S_\ell \langle L_{(\lambda)}^+ \rangle, W(k_\lambda)}^{\square, \chi, \text{min}}(\bar{\tau}_\lambda) = L_{(\lambda)}^+ D_{S_\ell \langle L_{(\lambda)}^+ \rangle, W(k_\lambda)}^{\square, \chi}(\bar{\tau}_\lambda)$$

for all λ with $L_{(\lambda)}^+ = L_{(\lambda')}^+$, except for a finite failure set \mathfrak{F}_λ . We should check that the occurrence of these failure sets at each step in the tower of field extensions does not disturb the desired result. For this, recall that the $L_{(\lambda)}^+$ show up in the tower $F^+ = L_0^+ \subset L_1^+ \subset \dots$ and that, by the first statement, we have

$$\lim_{i \rightarrow \infty} \delta \left\{ \lambda \mid L_{(\lambda)}^+ D_{S_{\ell(L_{(\lambda)}^+)}, W(k_\lambda)}^{\square, \chi, \min}(\bar{r}_\lambda) \text{ has vanishing dual Selmer group, } L_{(\lambda)}^+ \subset L_i^+ \right\} = 1.$$

But this clearly implies

$$\lim_{i \rightarrow \infty} \delta \left\{ \lambda \mid L_{(\lambda)}^+ D_{S_{\ell(L_{(\lambda)}^+)}, W(k_\lambda)}^{\square, \chi}(\bar{r}_\lambda) \text{ has vanishing dual Selmer group, } L_{(\lambda)}^+ \subset L_i^+, \lambda \notin \mathfrak{F}_\lambda \right\} = 1,$$

completing the proof. \square

Proof of Theorem 8.5. The „has vanishing dual Selmer group“-part of Theorem 8.5 follows immediately from Theorem 8.12 and the potential unobstructedness result of Lemma 4.8, as each $[L_{(\lambda)}^+ : F^+]$ is a power of 2 (and k_λ has odd characteristic for $\lambda \in \Lambda_\mathcal{E}^2$).

It remains to show that for all $\nu \in \Omega_\ell^{F^+}$ the local lifting ring $R_{W(k_\lambda)}^{\square, \chi_\nu}(\bar{r}_{\lambda, \nu})$ is relatively smooth. By Theorem 5.2 we know that

$$\begin{aligned} & \lim_{i \rightarrow \infty} \delta \left\{ \lambda \mid R_{W(k_\lambda)}^{\square, \chi_\nu}(\bar{r}_\lambda | \text{Gal}_{L_{(\lambda)}^+, \nu'}) \text{ is unobstructed for all } \nu' \in \Omega_{\ell(L_{(\lambda)}^+)}^+, L_{(\lambda)}^+ \subset L_i^+ \right\} \\ &= \lim_{i \rightarrow \infty} \delta \left\{ \lambda \mid \text{any } \nu' \in \Omega_{\ell(L_{(\lambda)}^+)}^+ \text{ is split in the extension } L_{(\lambda)} | L_{(\lambda)}^+, L_{(\lambda)}^+ \subset L_i^+ \right\} = 1. \end{aligned}$$

Using Proposition 4.5 and Corollary 4.4, the claim follows. \square

Appendix A: A lemma on prime densities in non-Galois extensions


Let us consider a CM-field F with totally real subfield F^+ and we denote by $L^+ = F^+(\sqrt{d_1}, \dots, \sqrt{d_k})$ the totally real extension of F^+ of degree 2^k , obtained by adjoining the square roots of some choice of elements $d_1, \dots, d_k \in \mathbb{N}$ such that each d_i is a non-square in the Galois closure \tilde{F}^+ of F^+ . Set $L = L^+.F$. Then we have:

Lemma A. *Let $\Xi_\mathbb{Q}$ be the set of all those rational primes ℓ with the following property: For any place \wp of L^+ ,*

$$[\wp \text{ divides } \ell] \implies [\wp \text{ splits in } L|L^+].$$

Then the density $\delta(\Xi_\mathbb{Q})$ of $\Xi_\mathbb{Q}$ in the set of all rational primes is at least $1 - \frac{1}{2^k}$.

Proof. Consider the following diagram of fields



with corresponding Galois groups $\Delta = \mathbb{Z}/2\mathbb{Z}$, $\Omega = (\mathbb{Z}/2\mathbb{Z})^k$ and Γ, H (for which we don't make an assumption). By our initial assumption that the d_i are not squares we have

$$\mathrm{Gal}(\tilde{L}^+|\mathbb{Q}) \cong \Gamma \times \Omega \quad \text{and, hence,} \quad \mathrm{Gal}(\tilde{L}|\mathbb{Q}) \cong \Gamma \times \Omega \times \Delta.$$

Now, let \mathfrak{P} be a place of \tilde{L} with corresponding Frobenius element $(\gamma, \omega, \delta) \in \mathrm{Gal}(\tilde{L}|\mathbb{Q})$. As Ω and Δ are abelian, the conjugacy class of \mathfrak{P} can be written as $\{(u\gamma u^{-1}, \omega, \delta) | u \in \Gamma\}$ and consists precisely of the Frobenii of the places of L lying over the same rational prime p as \mathfrak{P} . Let \wp be the place of L below \mathfrak{P} . Its Frobenius element is given by

$$(\gamma, \omega, \delta)^{e_{\gamma, \omega, \delta}} \in H \times \{1\} \times \Delta = \mathrm{Gal}(\tilde{L}|L^+)$$

for $e_{\gamma, \omega, \delta}$ minimal such that $(\gamma, \omega, \delta)^{e_{\gamma, \omega, \delta}} \in H \times \{1\} \times \Delta$. The condition that \wp splits in $L|L^+$ then amounts precisely to $(\gamma, \omega, \delta)^{e_{\gamma, \omega, \delta}} \in H \times \{1\} \times \{1\}$, or, written in a more sophisticated way, that $q((\gamma, \omega, \delta)^{e_{\gamma, \omega, \delta}}) = 1$, where

$$q : \mathrm{Gal}(\tilde{L}|L^+) \rightarrow \mathrm{Gal}(\tilde{L}|L^+) / \mathrm{Gal}(\tilde{L}|\tilde{L}^+)$$

is the quotient map. If $\omega \neq 1$, we clearly must have $2|e_{\gamma, \omega, \delta}$, which implies that \wp splits in $L|L^+$. It is also important to note that the condition $\omega \neq 1$ is not destroyed by conjugation inside $\mathrm{Gal}(\tilde{L}|\mathbb{Q})$. Now, set

$$\Xi^* = \{(\gamma, \omega, \delta) \in \mathrm{Gal}(\tilde{L}|\mathbb{Q}) \mid q((\gamma, \omega, \delta)^{e_{\gamma, \omega, \delta}}) = 1\}$$

and consider the subset $\Xi \subset \Xi^*$ which consists of those $g \in \Xi^*$ for which the complete conjugacy class is contained in Ξ^* , i.e. $\Xi = \{g \in \Xi^* \mid \langle g \rangle \subset \Xi^*\}$.

We can give another characterization of this set: Ξ is the union of all conjugacy classes $\langle g \rangle \subset \mathrm{Gal}(\tilde{L}|\mathbb{Q})$ with the following property: If \mathbf{P}_g denotes the set of all places \mathfrak{P} of \tilde{L} such that $\mathrm{Frob}_{\mathfrak{P}} \in \langle g \rangle$, then for any place \wp of L^+ the following holds:

$$[\exists \mathfrak{P} \in \mathbf{P}_g \text{ such that } \mathfrak{P} \text{ divides } \wp] \implies [\wp \text{ splits in } L|L^+].$$

Then we have

$$\#\Xi \geq \#\{(\gamma, \omega, \delta) \in \mathrm{Gal}(\tilde{L}|\mathbb{Q}) \mid \omega \neq 1\} = (2^k - 1) \cdot 2 \cdot \#\Gamma.$$

As $\Xi_{\mathbb{Q}} = \{\ell \in \mathrm{Pl}_{\mathbb{Q}} \mid \exists g \in \Xi \text{ such that } \mathfrak{P}|\ell \text{ for all } \mathfrak{P} \in \mathbf{P}_g\}$, it follows from Chebotarev's density theorem that

$$\delta(\Xi_{\mathbb{Q}}) \geq \frac{(2^k - 1) \cdot 2 \cdot \#\Gamma}{\#\mathrm{Gal}(\tilde{L}|\mathbb{Q})} = 1 - \frac{1}{2^k}. \quad \square$$

9 References

- [All16] P. ALLEN, *On automorphic points in polarized deformation rings*, preprint, 2016.
- [AC89] J. ARTHUR AND L. CLOZEL, *Simple algebras, base change, and the advanced theory of the trace formula*, Annals of Mathematics Studies **120**, Princeton University Press, Princeton, NJ, 1989.
- [AM69] M. F. ATIYAH AND I. G. MACDONALD, *Introduction to commutative algebra*, Addison-Wesley Publishing Co., Reading, Mass.-London-Don Mills, Ont., 1969.

- [BLGGT14] T. BARNET-LAMB, T. GEE, D. GERAGHTY AND R. TAYLOR, *Potential automorphy and change of weight*, Ann. of Math. (2) **179** (2014), no. 2, pp. 501-609.
- [BMR05] M. BATE, B. MARTIN AND G. RÖHRLE, *A geometric approach to complete reducibility*, Invent. Math. **161** (2005), no. 1, 177-218.
- [BMRT10] M. BATE, B. MARTIN, G. RÖHRLE AND R. TANGE, *Complete reducibility and separability*, Trans. Amer. Math. Soc. **362** (2010), no. 8, 4283-4311.
- [Bal12] S. BALAJI, *G-valued potentially semi-stable deformation rings*, Ph.D. thesis, The University of Chicago, Chicago, IL, 2012.
- [Bel09] J. BELLAÏCHE, *Introduction into the conjecture of Bloch-Kato*, unpublished notes for the Clay Summer School on Galois representations, Honolulu, 2009.
- [BC09] J. BELLAÏCHE AND G. CHENEVIER, *Families of Galois Representations and Selmer Groups*, Astérisque **324**, Publications de la SMF, Paris, 2009.
- [BC03] F. BLEHER AND T. CHINBURG, *Deformations with respect to an algebraic group*, Illinois J. Math. **47** (2003), no. 3, pp. 899-919.
- [Böc07] G. BÖCKLE, *Presentations of universal deformation rings*, in L-functions and Galois representations, London Math. Soc. Lecture Note Ser. **320**, Cambridge Univ. Press, Cambridge, 2007, pp. 24-58.
- [Böc13a] G. BÖCKLE, *Deformations of Galois representations*, in Elliptic curves, Hilbert modular forms and Galois deformations, Adv. Courses Math. CRM Barcelona, Basel, 2013, pp. 21-115.
- [CG13] F. CALEGARI AND T. GEE, *The conjectural connections between automorphic representations and Galois representations*, to appear in the Proceedings of the LMS Durham Symposium 2011.
- [CH13] G. CHENEVIER AND M. HARRIS, *Construction of automorphic Galois representations, II*, Camb. J. Math. **1**, 2013, pp. 53–73.
- [CHT08] L. CLOZEL, M. HARRIS AND R. TAYLOR, *Automorphy for some l -adic lifts of automorphic mod l Galois representations*, Publ. Math. Inst. Hautes Études Sci. No. **108** (2008), 1-181.
- [CDT99] B. CONRAD, F. DIAMOND AND R. TAYLOR, *Modularity of certain potentially Barsotti-Tate Galois Representations*, J. Amer. Math. Soc. **12** (1999), no. 2, pp. 521-567.
- [Del71] P. DELIGNE, *Formes modulaires et représentations de $GL(2)$* , Lecture Notes in Math. **179** (1971), pp. 139–172.
- [DS74] P. DELIGNE AND J.-P. SERRE, *Formes modulaires de poids 1*, Ann. Sci. ENS **7** (1974), pp. 507–530.
- [FL82] J.-M. FONTAINE AND G. LAFFAILLE, *Construction de représentations p -adiques*, Ann. Sci. École Norm. Sup. (4) **15** (1982), no. 4, 547-608.
- [Fre64] P. FREYD, *Abelian categories. An introduction to the theory of functors*, Harper's Series in Modern Mathematics, Harper & Row, Publishers, New York, 1964.

- [FH91] W. FULTON AND J. HARRIS, *Representation Theory. A first course*, Graduate Texts in Mathematics **129**, Readings in Mathematics, Springer-Verlag, New York, 1991.
- [Gam13] A. GAMZON, *Unobstructed Hilbert modular deformation problems*, 2013, to appear in Journal de Théorie des Nombres des Bordeaux.
- [Gee11] T. GEE, *Automorphic lifts of prescribed types*, Math. Ann. **350** (2011), no. 1, 107-144.
- [GK14] T. GEE AND M. KISIN, *The Breuil-Mézard conjecture for potentially Barsotti-Tate representations*, Forum Math. Pi **2** (2014), e1, 56 pp.
- [Ger10] D. GERAGHTY, *Modularity lifting theorems for ordinary Galois representations*, Ph.D. thesis, Harvard University, Cambridge, MA, 2010.
- [Gou01] F. Q. GOUVÊA, *Deformations of Galois representations*, IAS/Park City Math. Ser. **9**, Arithmetic algebraic geometry (Park City, UT, 1999), Amer. Math. Soc., Providence, RI, 2001, pp. 233-406.
- [Gue11] L. GUERBEROFF, *Modularity lifting theorems for Galois representations of unitary type*, Compos. Math. **147** (2011), no. 4, 1022-1058.
- [Gui16] D.-A. GUIRAUD, *A framework for unobstructedness of Galois deformation rings*, PhD thesis, Heidelberg University, 2016. Available at <http://www.ub.uni-heidelberg.de/archiv/20248>.
- [Gro64] A. GROTHENDIECK, *Eléments de géométrie algébrique. IV. Étude locale des schémas et des morphismes de schémas. I.*, Inst. Hautes Études Sci. Publ. Math. No. 20, 1964.
- [HHM08] J. HARRIS, J. HIST AND M. MOSSINGHOFF, *Combinatorics and graph theory. Second edition*, Undergraduate Texts in Mathematics. Springer, New York, 2008.
- [Hat15] J. HATLEY, *Obstruction criteria for modular deformation problems*, PhD thesis, University of Massachusetts Amherst, Amherst, MA, 2015.
- [JK81] G. JAMES AND A. KERBER, *The representation theory of the symmetric group*, Encyclopedia of Mathematics and its applications **16**, Addison-Wesley Publishing Co., Reading, Mass., 1981.
- [Jan89] U. JANNSEN, *On the l -adic cohomology of varieties over number fields and its Galois cohomology*, in: Galois groups over \mathbf{Q} (Berkeley, CA, 1987), Math. Sci. Res. Inst. Publ., **16**, Springer, New York, 1989, pp. 315–360.
- [Jan96] G. JANUSZ, *Algebraic number fields. Second edition*, Graduate Studies in Mathematics 7, American Mathematical Society, Providence, RI, 1996.
- [Kha03] C. KHARE, *On isomorphisms between deformation rings and Hecke rings*, with an appendix by G. Böckle, Invent. Math. **154**, no. 1, 2003, pp. 199–222.
- [KW09a] C. KHARE AND J.-P. WINTENBERGER, *On Serre's conjecture for 2-dimensional mod p representations of $\text{Gal}(\overline{\mathbf{Q}}/\mathbf{Q})$* , Ann. of Math. (2), (**169**), 2009, pp. 229–253.
- [KW09b] C. KHARE AND J.-P. WINTENBERGER, *Serre's modularity conjecture (II)*, Invent. Math. **178**, 2009, pp. 505-586.

- [Kis07] M. KISIN, *Modularity of 2-dimensional Galois representations*, Current Developments in Mathematics **2005**, 2007, pp. 191-230.
- [Lev13] B. LEVIN, *G-valued flat deformations and local models*, Ph.D. thesis, Stanford University, Stanford, CA, 2013.
- [Mau00] D. MAUGER, *Algèbre de Hecke quasi-ordinaire universelle d'un groupe réductif*, Ph.D. thesis, Université Paris-Nord (Paris 13), Paris, 2000.
- [Maz87] B. MAZUR, *Deforming Galois Representations*, in Galois groups over \mathbf{Q} , Y. Ihara, K. Ribet, J.-P. Serre (editors), MSRI Publ. 16 (1987), Springer-Verlag, New York-Berlin, 1989, 385-437.
- [Maz97] B. MAZUR, *An introduction to the deformation theory of Galois representations*, in Modular Forms and Fermat's Last Theorem, G. Cornell, J. Silverman, and G. Stevens (editors), Springer-Verlag, New York-Berlin, 1997, pp. 243-311.
- [NSW08] J. NEUKIRCH, A. SCHMIDT, K. WINGBERG, *Cohomology of Number Fields, Second edition*, Grundlehren der Mathematischen Wissenschaften 323, Springer-Verlag, Berlin, 2008.
- [PT15] S. PATRIKIS AND R. TAYLOR, *Automorphy and irreducibility of some l -adic representations*, Composition Math. **151**, 2015, p. 207.
- [PSW16] S. PATRIKIS, A. SNOWDEN AND A. WILES, *Residual irreducibility of compatible systems*, preprint, 2016.
- [Pra14] D. PRASAD, *Notes on representations of finite groups of Lie type*, unpublished lecture notes, 2014.
- [Ram93] R. RAMAKRISHNA, *On a variation of Mazur's deformation functor*, Compositio Math. **87** (1993), 269-286.
- [SZ99] P. SCHNEIDER AND E.-W. ZINK, *K-types for the tempered components of a p -adic general linear group*, J. Reine Angew. Math. **517** (1999), pp. 161-208.
- [Ser79] J.-P. SERRE, *Local Fields*, Graduate Texts in Mathematics **67**, Springer-Verlag, New York-Berlin, 1979.
- [Ser98] J.-P. SERRE, *The notion of complete reducibility in group theory*, Part II of the Moursund lectures, available at <http://math.uoregon.edu/resources/serre>, 1998.
- [Ser00] J.-P. SERRE, *Local Algebra*, Springer Monographs in Mathematics, Springer-Verlag, Berlin, 2000.
- [Shi71] G. SHIMURA, *Introduction to the arithmetic theory of automorphic functions*, Iwanami Shoten Publishers, Tokyo and Princeton University Press, Princeton, NJ, 1971.
- [Sho15] J. SHOTTON, *The Breuil-Mézard conjecture when l is not equal to p* , Ph.D. thesis, Imperial College, London, 2015.
- [Sho16] J. SHOTTON, *The Breuil-Mézard conjecture when $l \neq p$* , preprint, 2016.
- [Spr01] J. SPRANG, *A universal deformation ring with unexpected Krull dimension*, Math. Z. **275** (2013), no. 1-2, pp. 647-652.

- [Tay08] R. TAYLOR, *Automorphy for some l -adic lifts of automorphic mod l Galois representations. II*, Publ. Math. Inst. Hautes Études Sci. **(108)**, 2008, pp. 183–239.
- [Tho12] J. THORNE, *On the automorphy of l -adic Galois representations with small residual image*, J. Inst. Math. Jussieu **(12)**, no. 4, 2012, pp. 855–920.
- [Til96] J. TILOUINE, *Deformations of Galois representations and Hecke algebras*, Publ. Mehta Res. Inst., Narosa Publ., New Delhi, 1996.
- [Wed08] T. WEDHORN, *The local Langlands correspondence for $GL(n)$ over p -adic fields*, In: School on Automorphic Forms on $GL(n)$, (ed. by L. Göttsche, G. Harder, M.S. Raghunathan), ICTP Lecture Notes **21** (2008), pp. 237–320.
- [Wes04] T. WESTON, *Unobstructed modular deformation problems*, American J. Math. **126** (2004), pp. 1237–1252.
- [Yam04] A. YAMAGAMI, *On the Unobstructedness of the Deformation Problems of Residual Modular Representations*, Tokyo J. of Math. **27**, no. 2 (2004), pp. 443–445.