

MODULAR LINEAR DIFFERENTIAL EQUATIONS OF FOURTH ORDER AND MINIMAL \mathcal{W} -ALGEBRAS

KAZUYA KAWASETSU AND YUICHI SAKAI

ABSTRACT. A characterization of the minimal \mathcal{W} -algebras associated with the Deligne exceptional series at level $-h^\vee/6$ is obtained by using one-parameter family of modular linear differential equations of order 4. In particular, the characters of the Ramond-twisted modules of minimal \mathcal{W} -algebras related to the Deligne exceptional series satisfy one of these differential equations. In order to obtain the characterization, the differential equations in the one parameter family which have solutions of ‘‘CFT type’’ are classified, whose solutions are explicitly described.

1. INTRODUCTION

The modular linear differential equations (MLDEs) play an important role in the study of 2D conformal field theories, the theory of vertex operator algebras (VOAs) and the theory of modular forms (see e.g. [MMS1], [KZ2], [DLM], [Mil], [T] and [AKNS]). For example, they have appeared in attempts to classify C_2 -cofinite, rational VOAs and rational conformal field theories (RCFTs) from their characters.

S. Mathur, S. Mukhi and A. Sen classified the unitary RCFTs whose characters satisfy the second order MLDE of weight 0

$$(\#_s) \quad f'' - \frac{1}{6}E_2 f' - sE_4 f = 0 \quad (s \in \mathbb{C}),$$

where f is a function on the complex upper-half plane \mathbb{H} and E_k is the normalized Eisenstein series of weight k ([MMS1] and [MMS2]). Here

$$f' = D(f) = \frac{1}{2\pi\sqrt{-1}} \frac{df}{d\tau} = q \frac{df}{dq}, \quad (q = e^{2\pi i\tau}, \tau \in \mathbb{H}).$$

(The eq. $(\#_{\mu(t)})$, where $\mu(t) = t(t+2)/144$ for $t \in \mathbb{Q}$, is known as the *Kaneko-Zagier equation* [KZ2].)

In [MMS1] and [MMS2], they showed that such RCFTs coincide with the Wess-Zumino-Witten models (WZW models) at level 1 associated to the Lie algebras in the sequence $A_1 \subset A_2 \subset G_2 \subset D_4 \subset F_4 \subset E_6 \subset E_7 \subset E_8$, which is called the *Deligne exceptional series* ([D]). Note that the chiral algebra of the WZW model at level $k \in \mathbb{Z}_{\geq 0}$ associated to a simple Lie algebra \mathfrak{g} coincides with the *affine VOA* at level k associated with the same Lie algebra \mathfrak{g} .

The main theme of this paper is the following relations between MLDEs and VOAs associated with the Deligne exceptional series. We first prove a vertex operator algebraic analogue of the

2010 *Mathematics Subject Classification.* Primary 17B69; Secondary 17B67, 17B25, 17B68.

Key words and phrases. Modular linear differential equations, Modular forms, Vertex operator algebras, minimal \mathcal{W} -algebras, Deligne exceptional series, Modular invariance of characters.

above result in [MMS2]. Let V be a self-dual simple, C_2 -cofinite, rational VOA of CFT type such that the conformal weights of the V -modules are all non-negative and that the characters of modules over V satisfy (\sharp_s) for some $s \in \mathbb{C}$. We show that V is isomorphic to the affine VOA at level 1 associated with a Lie algebra in the Deligne exceptional series (Theorem 6.1 and Remark 6.1 (ii)).

Moreover, we find other MLDEs related to the Deligne exceptional series: we establish a characterization of \mathcal{W} -algebras associated with the Deligne exceptional series in terms of the following one-parameter family of MLDEs of order 4. Let s be a complex number. We study the MLDE of order 4 and weight 0 which has the form

$$(b_s) \quad \begin{aligned} & D^4(f) - E_2 D^3(f) + (3D(E_2) + \alpha_1(s)E_4)D^2(f) \\ & - \left(D^2(E_2) + \frac{\alpha_1(s)}{2}D(E_4) - \alpha_2(s)E_6 \right) D(f) + \alpha_3(s)E_8 f = 0, \end{aligned}$$

where

$$\alpha_1 = \frac{-25s^2 + 120s + 1332}{7200}, \quad \alpha_2 = \frac{(5s + 6)^2}{14400}, \quad \alpha_3 = \frac{(s - 18)(s + 6)(5s + 6)^2}{8294400}.$$

In § 3, we give a necessary condition of $s \in \mathbb{C}$ such that (b_s) has a solution of CFT type, where a function f is called of CFT type if f has a q -expansion $f(\tau) = q^\alpha(1 + \sum_{n=1}^{\infty} a_n q^n)$, where α is a rational number and a_n is a non-negative integer for each positive integer n . Moreover, we construct exact solutions of (b_s) for these values of s in Appendices B–C.

The space spanned by the characters of modules over a simple, C_2 -cofinite, rational VOA of CFT type is invariant under the slash action with weight 0 of $SL_2(\mathbb{Z})$, and is contained in the space of solutions of a MLDE ([Zhu]). The modular invariance property for $\frac{1}{m}\mathbb{Z}$ -graded, simple, C_2 -cofinite, \mathbb{Z}_m -rational, conical VOAs V is proved in [Va] (see also [DLM]), where $m \in \mathbb{Z}_{>0}$. Here a $\frac{1}{m}\mathbb{Z}$ -graded VOA $V = \bigoplus_{n \in \frac{1}{m}\mathbb{Z}} V_n$ is called *conical* if $\dim V_0 = 1$ and $\dim V_n = 0$ for all $n < 0$. (Therefore, the conical \mathbb{Z} -graded VOAs are the same as the VOAs of CFT type.)

Let W be a $\frac{1}{2}\mathbb{Z}_{\geq 0}$ -graded VOA. Let $W^{(0)}$ and $W^{(1)}$ be the subspaces of W spanned by the homogeneous vectors of weight \mathbb{Z} and $1/2 + \mathbb{Z}$, respectively. We say that W is *proper* $\frac{1}{2}\mathbb{Z}_{\geq 0}$ -graded if $W^{(1)} \neq 0$. The space spanned by the characters of the modules over a proper $\frac{1}{2}\mathbb{Z}_{\geq 0}$ -graded VOA is not modular invariant, but the space spanned by the characters of the *Ramond-twisted modules* is modular invariant (see § 5 for the details).

The \mathcal{W} -algebras are $\frac{1}{2}\mathbb{Z}_{\geq 0}$ -graded VOAs constructed using the *Drinfeld-Sokolov reduction* from affine VOAs (see [KRW]). The characters of Ramond-twisted modules over \mathcal{W} -algebras at admissible levels are determined in [KW2]. Moreover, several examples of \mathcal{W} -algebras at non-admissible levels are found to be C_2 -cofinite and \mathbb{Z}_2 -rational in [K2]. Among them, the characters of the Ramond-twisted modules over the \mathcal{W} -algebra $\mathcal{W}_{-5}(E_8, f_\theta)$ satisfy the second order MLDE $(\sharp_{\mu(19/5)})$ ([K2]). It was proved that there do not exist RCFTs whose characters satisfy $(\sharp_{\mu(19/5)})$ ([MMS2]). Thus we are interested in studying \mathcal{W} -algebras and $\frac{1}{2}\mathbb{Z}_{\geq 0}$ -graded VOAs with connections of the MLDEs.

In § 5–§ 8, we show that the characters of the Ramond-twisted modules over the \mathcal{W} -algebra $\mathcal{W}_{-h^\vee/6}(\mathfrak{g}, f_\theta)$ at level $-h^\vee/6$ associated with a Lie algebra \mathfrak{g} in the Deligne exceptional series

satisfy (b_s) with

$$(1.1) \quad s = \frac{6(7h^\vee - 18)}{5(h^\vee + 6)},$$

where h^\vee is the dual Coxeter number of \mathfrak{g} . The \mathcal{W} -algebras $W = \mathcal{W}_{-h^\vee/6}(\mathfrak{g}, f_\theta)$ satisfy the conditions (a)–(h) (*Main Conditions*) below:

Main Conditions:

(a) There is a complex number s such that the space of characters of Ramond-twisted W -modules is contained in the space of solutions of (b_s) and such that any pair of indicial roots of (b_s) does not have a non-zero integral difference.

(b) The vector spaces $W^{(0)}$ and $W^{(1)}$ have the forms

$$W^{(0)} \cong V \otimes L(-3/5, 0), \quad W^{(1)} \cong P \otimes L(-3/5, 3/4)$$

with a simple, C_2 -cofinite, rational VOA V and simple current V -module P . Here $L(-3/5, h)$ is the Virasoro minimal model of central charge $-3/5$ and conformal weight h .

(c) The conformal weight of the simple current V -module P is non-negative.

(d) The conformal weight of each non-vacuum irreducible V -module is non-zero.

(e) Let M and N be irreducible V -modules with the same conformal weights. Then the characters of M and N coincide.

(f) Let M and N be irreducible Ramond-twisted W -modules with the same conformal weights. Then M and N have the same character.

(g) Let S and T be irreducible Ramond-twisted W -modules with the minimum conformal weight. Let $S \cong \bigoplus_{i=1}^n S^i$ be the irreducible decomposition of S as a $W^{(0)}$ -module. Then as a $W^{(0)}$ -module, T decomposes into the direct sum $T \cong \bigoplus_{i=1}^n T^i$ such that for each $i = 1, \dots, n$, either $T^i = S^i$ or $T^i = (S^i)'$ holds, where $(S^i)'$ is the restricted dual of S^i over $W^{(0)}$.

(h) Any irreducible V -module is a summand of either an untwisted or Ramond-twisted W -module M as a V -module.

We show that the proper $\frac{1}{2}\mathbb{Z}_{\geq 0}$ -graded VOA W satisfying (a)–(h) is the \mathcal{W} -algebras $\mathcal{W}_{-h^\vee/6}(\mathfrak{g}, f_\theta)$ associated with a Lie algebra \mathfrak{g} in the Deligne exceptional series (see Theorem 8.1). For the moment, we cannot omit any of these 8 conditions. Although condition (a) provides information on the characters of Ramond-twisted W -modules as (\sharp_s) gives information on the characters of RCFTs in [MMS1] and [MMS2], we note that determining a proper $\frac{1}{2}\mathbb{Z}_{\geq 0}$ -graded VOA W by using the characters of Ramond-twisted W -modules may not be easy. It is because the vacuum module of W is not Ramond-twisted. Note also that characters of minimal \mathcal{W} -algebras can be determined by using the Kazhdan-Lusztig polynomials in principle ([A]), but it is not easy to compute the general Kazhdan-Lusztig polynomials.

The paper is organized as follows. In § 2, we give a brief explanation of a proof of Theorem 8.1 by using Main Conditions (the proof is given in § 8). In § 3, candidates of $s \in \mathbb{Q}$ such that (b_s) has a solution of CFT type are found by solving Diophantus equations. Fundamental systems of the space of solutions of such (b_s) 's are given in Appendices B–C. In § 4, we explain how to construct these solutions of (b_s) . Moreover, we deduce some properties of the space of solutions

of (b_s) , which will be used in the proof of Theorem 8.1. In § 5 we briefly recall the notion for VOAs. In § 6.1 we review affine VOAs and give a characterization of the affine VOAs associated with the Deligne exceptional series in terms of (\sharp_s) . In § 6.2 we study the modules over an extension of the minimal model $L(-3/5, 0)$. In § 7 we show that the characters of the Ramond-twisted modules over minimal \mathcal{W} -algebras associated with the Deligne exceptional series satisfy (b_s) for s given by (1.1). Theorem 8.1 is proved in § 8. In § 9 we give several comments on the relation between the characters of VOAs and (b_s) . In Appendix A, we introduce modular forms which are necessary to represent explicit solutions of (b_s) which have solutions of CFT type as explained in Appendices B–C. We divide such a set of s into two parts: (I) the number s such that (b_s) has a solution of CFT type and each solution of CFT type is not quasimodular of positive depth and (II) the number s such that (b_s) has a quasimodular solution of CFT type of positive depth. The quasimodular forms (of positive depth) can be expressed in terms of homogenous polynomials that have modular forms as coefficients with respect to E_2 (cf. [KZ1]). The solutions of (b_s) of case (I) are studied in Appendix B, and that of (b_s) of case (II) are studied in Appendix C. In Appendix D, we define polynomials which are used to represent solutions of MLDEs in Appendices B–C.

2. OVERVIEW OF THE PROOF OF A MAIN RESULT

In this section, we briefly explain how to prove Theorem 8.1 in § 8.

By condition (a), we have a restriction on the conformal weights of the Ramond-twisted irreducible W -modules. Then by (b) and (h), we obtain restrictions on the conformal weights of the irreducible V -modules. (The condition (h) is related with the orbifold theory (cf. [CM]).) By using (a) and (c)–(g), we can study the V -modules and Ramond-twisted W -modules. (The condition (c) is satisfied by all known simple C_2 -cofinite rational VOAs to the best of authors' knowledge.) Suppose that there are only two inequivalent irreducible V -modules up to isomorphisms. It then follows that the space of characters of Ramond-twisted W -modules is two-dimensional. By using the basis of MLDEs given in Appendix B, we see that $s = 32/5$ and the characters of V satisfies $(\sharp_{\mu(\tau/2)})$. It then follows by Theorem 6.1 that $V \cong V_1(E_7)$, where $V_k(\mathfrak{g})$ is the affine VOA at level k associated with a finite-dimensional simple Lie algebra \mathfrak{g} . Then we see that $W \cong \mathcal{W}_{-5}(E_8, f_\theta)$ by Lemma 7.1. Suppose that the number of the irreducible V -modules up to isomorphisms is more than 2. Then we can construct the modules of the VOA $R = V \otimes V_1(A_1) \oplus P \otimes V_1(A_1; \alpha/2)$, and we see that the characters of the R -modules satisfy $(\sharp_{\mu(t)})$ with $t \in \mathbb{Q}$. It then follows by Theorem 6.1 that $R \cong V_1(\mathfrak{g})$ with a Lie algebra \mathfrak{g} in the Deligne exceptional series. Then we see that $W \cong \mathcal{W}_{-h\nu/6}(\mathfrak{g}, f_\theta)$ by Lemma 7.1. (See § 8 for the detail of the proof.)

3. MODULAR LINEAR DIFFERENTIAL EQUATIONS WITH SOLUTIONS OF CFT TYPE

In this section we show that if (b_s) has a solution of CFT type, then s belongs to (3.12) on § 3.6.

3.1. Indicial equations. Before studying MLDEs with solutions of CFT type, we recall the notion of indicial equations of MLDEs, which is used to classify such MLDEs.

Let f be a function of the form $f(\tau) = q^\alpha \sum_{n=0}^{\infty} a_n q^n$ with $\alpha \in \mathbb{Q}$, $a_0 = 1$ and $a_n \in \mathbb{C}$ for each $n \geq 1$. We call the number α the *exponent* of f . Let n be a non-negative integer, and suppose that $\alpha + n \neq 0$ for any $n > 0$. Let s be a complex number, and suppose that f is a solution

of (b_s) . By substituting the q -expansion of $f(\tau)$ into (b_s) and comparing the coefficients of $q^{\alpha+n}$ in the both-sides, we obtain a relation

$$(3.1) \quad (\alpha + n)^4 a_n + \sum_{i=0}^n \left(-e_{2,i}(n-i+\alpha)^3 a_{n-i} + (3ie_{2,i} + \alpha_1(s)e_{4,i})(n-i+\alpha)^2 a_{n-i} \right. \\ \left. - \left(i^2 e_{2,i} + \frac{\alpha_1(s)}{2} i e_{4,i} - \alpha_2(s)e_{6,i} \right) (n-i+\alpha) a_{n-i} + \alpha_3(s)e_{8,i} a_{n-i} \right) = 0,$$

where $E_k(\tau) = \sum_{n=0}^{\infty} e_{k,n} q^n$ is the Fourier expansion of the normalized Eisenstein series of weight k .

By substituting $n = 0$ into (3.1), we see that α satisfies the relation $(120\alpha - 5s - 6)(24\alpha - s - 6)(24\alpha + s - 18)(120\alpha + 5s + 6) = 0$, which is called the *indicial equation* of (b_s) . The roots $\alpha_1 = -s/24 - 1/20$, $\alpha_2 = -s/24 + 3/4$, $\alpha_3 = s/24 + 1/4$ and $\alpha_4 = s/24 + 1/20$ of the indicial equation are called the *indicial roots (indices)* of (b_s) . Since $\alpha \in \mathbb{Q}$ and α coincides with one of $\alpha_1, \dots, \alpha_4$, we have $s \in \mathbb{Q}$. It is well known that if there is no integral differences among the indicial roots, then for each indicial root α , there exists a solution of (b_s) of the form $f(\tau) = q^\alpha(1 + O(q))$.

In the following, suppose that f is of CFT type, so that $a_n \in \mathbb{Z}_{\geq 0}$ for $n \geq 1$. We call -24α a *formal central charge* of (b_s) . For each $1 \leq i \leq 4$, we call $\alpha_i - \alpha$ a *formal conformal weight* of (b_s) .

3.2. The case $\alpha = -s/24 - 1/20$. It follows from (3.1) with $n = 1$ that $(5s - 54)(25s^2 + 5sa_1 + 120s - 42a_1 + 108) = 0$. Assuming that $s \neq 54/5$, we have

$$(3.2) \quad 25s^2 + 5sa_1 + 120s - 42a_1 + 108 = 0.$$

By setting $t = 5s$, we see that (3.2) becomes

$$(3.3) \quad t^2 + ta_1 + 24t - 42a_1 + 108 = 0,$$

which is a monic quadratic in t with integral coefficients. Therefore, any rational solution t is an integer. Since $(t - 42)(t + a_1 + 66) = -2880$ by (3.3) and both $t - 42$ and $t + a_1 + 66$ are integers, we see that s is an element of the set

$$(3.4) \quad \left\{ -\frac{2838}{5}, -\frac{1398}{5}, -\frac{918}{5}, -\frac{678}{5}, -\frac{534}{5}, -\frac{438}{5}, -\frac{318}{5}, -\frac{278}{5}, -\frac{246}{5}, -\frac{198}{5}, -30, -\frac{138}{5}, -\frac{118}{5}, \right. \\ \left. -\frac{102}{5}, -\frac{78}{5}, -\frac{54}{5}, -\frac{48}{5}, -\frac{38}{5}, -6, -\frac{22}{5}, -\frac{18}{5}, -\frac{6}{5}, -\frac{3}{5}, \frac{2}{5}, \frac{6}{5}, 2, \frac{12}{5}, \frac{18}{5}, \frac{22}{5}, \frac{24}{5}, \frac{26}{5}, \frac{27}{5}, \frac{32}{5}, \right. \\ \left. \frac{33}{5}, \frac{34}{5}, \frac{36}{5}, \frac{37}{5}, \frac{38}{5}, \frac{39}{5}, 8, \frac{41}{5} \right\}.$$

For each value s in (3.4), we see that any formal conformal weight of (b_s) is not a positive integer by evaluating the value $\alpha_j - \alpha_1$ ($j = 2, 3, 4$). Therefore, the Fourier expansion of the solution f is determined by (3.1).

By substituting $n = 2$ into (3.1), we have

$$-386208 - 720360s - 255500s^2 - 15750s^3 + 3125s^4 - 72792a_1 - 22140sa_1 - 26250s^2a_1 \\ + 1375s^3a_1 + 139536a_2 - 12960sa_2 + 300s^2a_2 = 0.$$

Since a_2 is a non-negative integer, (3.4) reduces to

$$\left\{ -\frac{438}{5}, -\frac{318}{5}, -\frac{198}{5}, -30, -\frac{138}{5}, -\frac{78}{5}, -\frac{48}{5}, -\frac{38}{5}, -\frac{18}{5}, -\frac{6}{5}, -\frac{3}{5}, \frac{2}{5}, \frac{6}{5}, \frac{12}{5}, \frac{18}{5}, \frac{22}{5}, \frac{27}{5}, 6, \frac{32}{5}, \frac{39}{5} \right\}.$$

The similar discussions above for $n = 3$ and 4 show that this set reduces to

$$(3.5) \quad \left\{ -\frac{318}{5}, -\frac{198}{5}, -\frac{138}{5}, -\frac{78}{5}, -\frac{48}{5}, -\frac{38}{5}, -\frac{18}{5}, -\frac{6}{5}, -\frac{3}{5}, \frac{2}{5}, \frac{6}{5}, \frac{12}{5}, \frac{18}{5}, \frac{22}{5}, \frac{27}{5}, 6, \frac{32}{5} \right\}.$$

Hence, if f is a solution of (b_s) of CFT type, then s is either $54/5$ or a value in (3.5).

3.3. The case $\alpha = -s/24 + 3/4$. In this case, we have $(s - 18)(75s^2 + 15sa_1 + 100s - 306a_1 + 348) = 0$ by substituting $n = 1$ into (3.1). Assuming $s \neq 18$ and setting $t = 15s$, we have $t^2 + 3ta_1 + 20t - 918a_1 + 1044 = 0$, which is a monic quadratic equation in t with integral coefficients. Therefore any rational solution t is an integer. Since $(t - 306)(t + 3a_1 + 326) = 100800$, it follows that s is an element of the set

$$(3.6) \quad \left\{ -\frac{16698}{5}, -\frac{6618}{5}, -\frac{4098}{5}, -\frac{2298}{5}, -\frac{5294}{15}, -\frac{1578}{5}, -\frac{948}{5}, -\frac{858}{5}, -\frac{1934}{15}, -114, -\frac{498}{5}, -\frac{1094}{15}, \right. \\ \left. -\frac{318}{5}, -\frac{494}{15}, -\frac{138}{5}, -\frac{254}{15}, -\frac{66}{5}, -\frac{48}{5}, -\frac{44}{15}, -\frac{14}{15}, -\frac{3}{5}, \frac{6}{5}, \frac{82}{15}, \frac{106}{15}, \frac{42}{5}, \frac{166}{15}, 12, \frac{226}{15}, \frac{78}{5}, \frac{50}{3}, \right. \\ \left. \frac{256}{15}, \frac{87}{5}, \frac{271}{15}, \frac{274}{15}, \frac{286}{15}, \frac{96}{5}, \frac{292}{15}, \frac{298}{15}, \frac{301}{15}, \frac{304}{15} \right\}.$$

For each value s in the above set, we see that any formal conformal weight of (b_s) is not a positive integer by evaluating the value $\alpha_j - \alpha_2$ ($j = 1, 3, 4$). Therefore, the Fourier expansion of the solution f is determined by (3.1).

By substituting $n = 2$ into (3.1), we have

$$625s^4 - 1350s^3 + 475a_1s^3 - 155340s^2 - 13650a_1s^2 + 140a_2s^2 - 385128s - 1836a_1s \\ - 8736a_2s - 474336 + 161352a_1 + 136080a_2 = 0.$$

For an s in (3.10), the following one gives non-negative integer a_2

$$\left\{ -\frac{498}{5}, -\frac{318}{5}, -\frac{138}{5}, -\frac{3}{5}, \frac{6}{5}, \frac{42}{5}, \frac{87}{5}, \frac{96}{5} \right\}.$$

Similarly, by considering $n = 3, 4, \dots, 32$, the set reduces to

$$(3.7) \quad \left\{ -\frac{3}{5}, \frac{6}{5}, \frac{42}{5} \right\}.$$

Hence, if f is a solution of (b_s) of CFT type, then s is either 18 or a value in (3.7).

3.4. **The case $\alpha = \mathbf{s}/24 + \mathbf{1}/4$.** It follows from (3.1) with $n = 1$ that $(s+6)(25s^2 - 5sa_1 + 130s - 78a_1 + 144) = 0$. Assuming $s \neq -6$ and setting $t = 5s$, we see that $t^2 - ta_1 + 26t - 78a_1 + 144 = 0$, which is a monic quadratic equation in t with integral coefficients. It then follows that any rational solution t is an integer. Since $(t+78)(t-a_1-52) = 4200$, we see that s belongs to the set

$$(3.8) \quad \left\{ -\frac{77}{5}, -\frac{76}{5}, -15, -\frac{74}{5}, -\frac{73}{5}, -\frac{72}{5}, -\frac{71}{5}, -14, -\frac{68}{5}, -\frac{66}{5}, -\frac{64}{5}, -\frac{63}{5}, -\frac{58}{5}, -\frac{57}{5}, -\frac{54}{5}, -\frac{53}{5}, \right. \\ \left. -10, -\frac{48}{5}, -\frac{43}{5}, -\frac{38}{5}, -\frac{36}{5}, -\frac{28}{5}, -\frac{22}{5}, -\frac{18}{5}, -\frac{8}{5}, -\frac{3}{5}, \frac{6}{5}, \frac{22}{5}, \frac{27}{5}, \frac{42}{5}, \frac{62}{5}, \frac{72}{5}, 18, \frac{97}{5}, \frac{122}{5}, \frac{132}{5}, \right. \\ \left. \frac{202}{5}, \frac{222}{5}, \frac{272}{5}, \frac{342}{5}, \frac{447}{5}, \frac{522}{5}, \frac{622}{5}, \frac{762}{5}, \frac{972}{5}, \frac{1322}{5}, \frac{2022}{5}, \frac{4122}{5} \right\}.$$

For each value s in the above set, we see that any formal conformal weight of (b_s) is not a positive integer by evaluating the value $\alpha_j - \alpha_3$ ($j = 1, 2, 4$). Therefore, the Fourier expansion of the solution f is uniquely determined by (3.1).

By substituting $n = 2$ into (3.1), we have

$$\begin{aligned} & -245592 - 295812s - 124830s^2 - 9975s^3 + 625s^4 + 15120a_1 + 9216sa_1 - 6180s^2a_1 \\ & - 400s^3a_1 + 54648a_2 + 5016sa_2 + 110s^2a_2 = 0. \end{aligned}$$

For s in (3.8), the following one gives non-negative integer a_2 :

$$\left\{ -15, -\frac{73}{5}, -\frac{68}{5}, -\frac{66}{5}, -\frac{63}{5}, -\frac{58}{5}, -\frac{54}{5}, -\frac{48}{5}, -\frac{38}{5}, -\frac{18}{5}, -\frac{8}{5}, -\frac{3}{5}, \frac{6}{5}, \frac{22}{5}, \frac{42}{5}, \frac{62}{5}, \frac{72}{5}, \frac{122}{5}, \frac{132}{5}, \right. \\ \left. \frac{222}{5}, \frac{342}{5}, \frac{447}{5}, \frac{762}{5}, \frac{2022}{5} \right\}.$$

Similarly, by considering $n = 3, 4, \dots, 23$, the set reduces to

$$(3.9) \quad \left\{ -\frac{66}{5}, -\frac{18}{5}, -\frac{8}{5}, -\frac{3}{5}, \frac{6}{5} \right\}.$$

Hence, if f is a solution of (b_s) of CFT type, then s is either -6 or a value in (3.9).

3.5. **The case $\alpha = \mathbf{s}/24 + \mathbf{1}/20$.** By substituting $n = 1$ into (3.1), we have $(5s+66)(25s^2 - 5sa_1 + 45s - 18a_1 + 18) = 0$. Assuming $s \neq -66/5$ and setting $t = 5s$, we see that $t^2 - ta_1 + 9t - 18a_1 + 18 = 0$, which is a monic quadratic equation in t with integral coefficients. Therefore the number t is an integer. Since $(t+18)(t-a_1-9) = 180$, we see that

$$(3.10) \quad s = -\frac{17}{5}, -\frac{16}{5}, -3, -\frac{14}{5}, -\frac{13}{5}, -\frac{12}{5}, -\frac{9}{5}, -\frac{8}{5}, -\frac{6}{5}, -\frac{3}{5}, 0, \frac{2}{5}, \frac{12}{5}, \frac{18}{5}, \frac{27}{5}, \frac{42}{5}, \frac{72}{5}, \frac{162}{5}.$$

For each s in (3.10), we see that any formal conformal weight of (b_s) is not a positive integer by evaluating the value $\alpha_j - \alpha_4$ ($j = 1, 2, 3$). Therefore, the Fourier expansion of the solution f is uniquely determined by (3.1).

By substituting $n = 2$ into (3.1), we have

$$\begin{aligned} & -661608 - 1138860s - 551250s^2 - 47625s^3 + 3125s^4 - 41472a_1 + 11160sa_1 - 24000s^2a_1 \\ & - 1750s^3a_1 + 176904a_2 + 18360sa_2 + 450s^2a_2 = 0. \end{aligned}$$

Then among the values of s in (3.10), the following give non-negative integer a_2 :

$$s = -3, -\frac{8}{5}, -\frac{6}{5}, -\frac{3}{5}, \frac{2}{5}, \frac{12}{5}, \frac{42}{5}.$$

Similarly, by considering $n = 3$, we see that s is one of the following:

$$(3.11) \quad s = -\frac{8}{5}, -\frac{6}{5}, -\frac{3}{5}, \frac{2}{5}, \frac{12}{5}, \frac{42}{5}.$$

Hence, if f is a solution of (b_s) of CFT type, then s is either $-66/5$ or a value in (3.11).

3.6. A necessary condition of s such that (b_s) has a solution of CFT type. Let s be a complex number and suppose that (b_s) has a solution of CFT type. By combining (3.5), (3.7), (3.9), (3.11) and $s = 54/5, 18, -6, -66/5$, we see that s is one of the following 23 numbers:

$$(3.12) \quad \begin{aligned} & -\frac{318}{5}, -\frac{198}{5}, -\frac{138}{5}, -\frac{78}{5}, -\frac{48}{5}, -\frac{38}{5}, -\frac{18}{5}, -\frac{6}{5}, -\frac{3}{5}, \frac{2}{5}, \frac{6}{5}, \frac{12}{5}, \frac{18}{5}, \frac{22}{5}, \frac{27}{5}, 6, \frac{32}{5}, \frac{54}{5}, \\ & \frac{42}{5}, 18, -\frac{66}{5}, -6, -\frac{8}{5}. \end{aligned}$$

4. SOLUTIONS OF MODULAR LINEAR DIFFERENTIAL EQUATIONS

Let s be a number in (3.12). In Appendices B–C, we give solutions of the MLDE (b_s) of (quasi)modular forms. In this section, we explain our method to construct such solutions of (b_s) . We first find a certain rational number k and a positive integer N , and construct solutions f of a MLDE (b_s^k) defined below, assuming that the MLDE (b_s^k) has solutions of (quasi)modular forms of weight k and level N . It then follows that the function $g = \eta^{-2k}f$ is a solution of (b_s) . Thus, we find solutions of (b_s) of quasimodular forms.

Moreover, we give some properties of the space of solutions of (b_s) which is used in § 8.

4.1. Construction of solutions. In this subsection, we explain our method to construct solutions of MLDEs listed in Appendices B–C.

First, we introduce a MLDE (b_s^k) . Let k and s be complex numbers. The differential equation

$$(b_s^k) \quad \vartheta_k^4(f) + \left(\alpha_1(s) - \frac{11}{36}\right)E_4\vartheta_k^2(f) + \frac{36\alpha_1(s) + 216\alpha_2(s) - 5}{216}E_6\vartheta_k(f) + \alpha_3(s)E_8f = 0$$

is a MLDE of order 4 with weight k , where $\vartheta_k(f) = D(f) - k(E_2/12)f$ is the Serre derivation and ϑ_k^i is the iterated Serre derivation defined by $\vartheta_k^i(f) = \vartheta_{k+2i-2} \circ \vartheta_{k+2i-4}^{i-1}(f)$ for $i \geq 1$ and $\vartheta_k^0(f) = f$. By setting $k = 0$ and using relations $24\eta' = E_2\eta$, $12E_2' = E_2^2 - E_4$, $3E_4' = E_2E_4 - E_6$ and $2E_6' = E_2E_6 - E_4^2$ (and $E_4^2 = E_8$), we see that the equation (b_s^0) is equal to the MLDE (b_s) . Because $\vartheta_k(\eta^{2\ell} \cdot f) = \eta^{2\ell}\vartheta_{k-\ell}(f)$, f is a solution of (b_s) if and only if $\eta^{2k} \cdot f$ is a solution of (b_s^k) .

Now we explain how to construct solutions of (b_s) of (quasi)modular forms. Let s be a number in (3.12) and $\alpha_1, \dots, \alpha_4$ the indicial roots of (b_s) introduced in § 3.1, which are rational numbers.

Set $k = -12 \min\{\alpha_1, \dots, \alpha_4\}$. Let N be the least common multiple of the denominators of the numbers $\alpha_1 - k/12, \dots, \alpha_4 - k/12$. Let F be a solution of (b_s^k) and suppose that F is a modular form of weight k and level N . The modular forms of level N and weight k are homogeneous polynomials in specific modular forms of level N given in Table 2 in Appendix A. Therefore, by calculating sufficiently many Fourier coefficients of F , we can determine F as a polynomial in the modular forms. In order to show that F does satisfy the MLDE (b_s^k) , we just substitute F to the left-hand side of (b_s^k) and show that it is zero by using differential relations and functional equations of modular forms given in Appendix A. We can also use a similar method to construct solutions of (b_s^k) of quasimodular forms.

We here perform this method for a typical case $\mathbf{s} = \mathbf{6}/\mathbf{5}$. Since the set of indicial roots of $(b_{6/5})$ is given by $S = \{-1/10, 1/10, 3/10, 7/10\}$, we have $k = 6/5$ and $N = 5$. Let α be any indicial root of $(b_{6/5})$. Let F_α be a solution of $(b_{6/5}^{\alpha})$ of the form $F_\alpha(\tau) = q^{\alpha+6/5}(1 + O(q))$. We suppose that F_α is a modular form of weight $6/5$ and level 5. It then follows that F_α has the form $F_\alpha = \sum_{n=0}^6 \beta_n \psi_1^n \psi_2^{6-n}$ with $\beta_0, \dots, \beta_6 \in \mathbb{C}$, where ψ_1 and ψ_2 are modular forms of weight $1/5$ and level 5 given in Table 2 in Appendix A:

$$\psi_1(\tau) = \eta(\tau)^{2/5} q^{-1/60} \prod_{\substack{n>0 \\ n \neq 0, \pm 2 \pmod{5}}} \frac{1}{1 - q^n}, \quad \psi_2(\tau) = \eta(\tau)^{2/5} q^{11/60} \prod_{\substack{n>0 \\ n \neq 0, \pm 1 \pmod{5}}} \frac{1}{1 - q^n}$$

(Rogers-Ramanujan functions). Since $F_\alpha/\eta^{12/5}$ is a solution of $(b_{6/5})$ of exponent α , we can compute any Fourier coefficient of F_α by using (3.1). Therefore we can determine β_0, \dots, β_6 (by computing finite number of Fourier coefficients of F_α .) In this way, we see that

$$(4.1) \quad \begin{aligned} F_{-1/10} &= \psi_1(\psi_1^5 + 2\psi_2^5), & F_{1/10} &= \psi_2(2\psi_1^5 - \psi_2^5)/2, \\ F_{3/10} &= \psi_1^4 \psi_2^2, & F_{7/10} &= \psi_1^2 \psi_2^4. \end{aligned}$$

Thus we find an explicit description of $\{F_\alpha | \alpha \in S\}$ under the assumption that F_α is a modular form of weight $6/5$ and level 5 for any $\alpha \in S$. By using the differential relations of ψ_1 and ψ_2 written in Appendix A (d), we can show that the right-hand sides of (4.1) do satisfy the MLDE $(b_{6/5}^{\alpha})$. Since $\{f_{\alpha+1/10} := F_\alpha/\eta^{12/5} | \alpha \in S\}$ is a fundamental system of solutions of $(b_{6/5})$, we have an explicit description of solutions of $(b_{6/5})$ in terms of ψ_1 and ψ_2 , as written in Appendix B (f).

In a similar way, we can construct solutions of (b_s) listed in Appendices B–C. By using the list of solutions, we have the following theorem.

Theorem 4.1. *Let s be a complex number and suppose that (b_s) has a solution f of CFT type. Suppose that f is not a quasimodular form of positive depth. Then s is one of the following 17 numbers:*

$$(4.2) \quad -\frac{48}{5}, -\frac{38}{5}, -\frac{6}{5}, -\frac{3}{5}, \frac{2}{5}, \frac{6}{5}, \frac{12}{5}, \frac{18}{5}, \frac{22}{5}, \frac{27}{5}, 6, \frac{32}{5}, \frac{54}{5}, 18, -\frac{66}{5}, -6, -\frac{8}{5}.$$

4.2. Modular invariant subspaces. In this subsection we find some properties of the space of solutions of (b_s) which is used in § 8.

A function f on \mathbb{H} is called of *character type* if f has the form $f(\tau) = q^\beta \sum_{n=0}^{\infty} a_n q^n$ with $\beta \in \mathbb{Q}$, $a_n \in \mathbb{Z}_{\geq 0}$ ($n \in \mathbb{Z}_{\geq 0}$) and $a_0 \geq 1$.

Proposition 4.1. *Let s be one of the numbers $-3/5, 2/5, 6/5, 12/5, 18/5, 22/5, 27/5, 6$ and $32/5$. Let f and g be linearly-independent solutions of (\mathfrak{b}_s) with exponents α and $\beta \in \mathbb{Q}$. Suppose that $\beta - \alpha = 4/5$, f is of CFT type, g is of character type, and the space spanned by f and g is modular invariant. Then $s = 32/5$.*

Proof. We show the assertion by contradiction in case by case bases. Suppose that $s = -3/5$. Then the set of indices of $(\mathfrak{b}_{-3/5})$ is $\{31/40, 9/40, 1/40, -1/40\}$. Since $\beta - \alpha = 4/5$, we have $\alpha = -1/40$ and $\beta = 31/40$. Therefore, $f = f_0$ and $g = c \cdot f_{4/5}$ with $c \in \mathbb{C}^\times$, where the solutions f_h with formal conformal weights h of $(\mathfrak{b}_{-3/5})$ are defined in Appendix B (d). By using the expression of f_0 in Appendix B (d), we can show that $f_0(-1/\tau) = \sqrt{2/5}(\sin(2\pi/5)f_0(\tau) + \sin(\pi/5)f_{4/5}(\tau) + \sin(\pi/5)f_{1/20}(\tau) + \sin(2\pi/5)f_{1/4}(\tau))$. Therefore, the space spanned by f_0 and $f_{4/5}$ is not closed under $SL_2(\mathbb{Z})$, which contradicts the assumption. We can reduce the other cases outside $s = 32/5$ to contradiction in a similar way. Thus we have $s = 32/5$. \square

The *modular Wronskian* $W(F)$ of a vector-valued modular function $F = {}^t(f_1, \dots, f_n)$ ($n \geq 1$) is defined by

$$W(F) = \det(F, \vartheta_0(F), \vartheta_0^2(F), \dots, \vartheta_0^{n-1}(F)).$$

Proposition 4.2. *If either $s = -48/5$ or $-38/5$, then there is no modular invariant subspace U of the space of solutions of (\mathfrak{b}_s) such that U is spanned by functions of character type and such that a function of CFT type belongs to U .*

Proof. We here only show the case of $s = -48/5$. The case of $s = -38/5$ is similarly proved. Since $f_0, f_{4/5}, f_{-1/2}$ and $f_{-7/10}$ defined in (a) in Appendix B form a fundamental system of solutions of $(\mathfrak{b}_{-48/5})$, it follows that

$$f_0(-1/\tau) = c_1 f_0(\tau) + c_2 f_{4/5}(\tau) + c_3 f_{-1/2}(\tau) + c_4 f_{-7/10}(\tau) \quad (c_i \in \mathbb{C}).$$

As any scalar multiple of $f_{-7/10}$ is not of character type, it suffices to show $c_4 \neq 0$. Suppose that $c_4 = 0$. Since $\eta^{42/5} f_0$ is not a modular form (with a multiplier system) on $SL_2(\mathbb{Z})$, one of c_2 and c_3 is non-zero. Therefore, one of

$$F_1 := \begin{pmatrix} f_0 \\ f_{4/5} \\ f_{-1/2} \end{pmatrix}, \quad F_2 := \begin{pmatrix} f_0 \\ f_{-1/2} \end{pmatrix}, \quad F_3 := \begin{pmatrix} f_0 \\ f_{4/5} \end{pmatrix}$$

is a vector-valued modular function on $SL_2(\mathbb{Z})$. Let λ_i be the sum of exponents of all components of the vector F_i . Since the exponents of $f_0, f_{4/5}$ and $f_{-1/2}$ are $7/20, 23/20$ and $-3/20$, respectively, it follows that $\lambda_1 = 27/20, \lambda_2 = 1/5$ and $\lambda_3 = 3/2$. Suppose that F_i is a vector-valued modular function on $SL_2(\mathbb{Z})$. It then follows from [Ma, Theorem 3.7] that $W(F_i)/\eta^{24\lambda_i}$ is a non-zero classical modular form on $SL_2(\mathbb{Z})$ of weight $w = 6 - 12\lambda_i$. Then we see that w is negative, it contradicts to the fact that there does not exist a non-zero classical modular form on $SL_2(\mathbb{Z})$ of negative weight. Hence, we have $c_4 \neq 0$, as desired. \square

5. VERTEX OPERATOR ALGEBRAS

In this section we recall some properties for $\frac{1}{2}\mathbb{Z}_{\geq 0}$ -graded VOAs.

Let $W = \bigoplus_{n \in \frac{1}{2}\mathbb{Z}_{\geq 0}} W_n$ be a simple $\frac{1}{2}\mathbb{Z}_{\geq 0}$ -graded conical VOA with central charge $c \in \mathbb{C}$. Set $W^{(0)} = \bigoplus_{n \in \mathbb{Z}} W_n$ and $W^{(1)} = \bigoplus_{n \in 1/2 + \mathbb{Z}} W_n$. Then there exists an automorphism σ of W defined by $\sigma(a) = a$ and $\sigma(v) = -v$ for each $a \in W^{(0)}$ and $v \in W^{(1)}$. If $W^{(1)} \neq 0$, then the order of σ is 2, otherwise $\sigma = \text{id}_W$. We set $G = \langle \sigma \rangle$. The VOA W is called \mathbb{Z}_2 -rational if W is rational and σ -rational. The W -module W is called the *vacuum module*.

Suppose that W is C_2 -cofinite and \mathbb{Z}_2 -rational. The σ -twisted W -modules are called the *Ramond-twisted modules*. It then follows that any Ramond-twisted irreducible W -module M has the form $M = \bigoplus_{n=0}^{\infty} M_{n+h}$, where $h \in \mathbb{C}$ and $M_{n+h} = \{v \in M \mid L_0 v = (n+h)v\}$ such that $M_h \neq 0$. The number h is called the *conformal weight* of M . If $W^{(1)} = 0$, then W is a $\mathbb{Z}_{\geq 0}$ -graded VOA, and then the Ramond-twisted W -modules coincide with the untwisted W -modules. The space spanned by the characters of the Ramond-twisted modules is $SL_2(\mathbb{Z})$ -invariant (see [Va, Theorem 1.3]).

Let M be a Ramond-twisted irreducible W -module with a conformal weight h . The *character* of M is defined by $\chi_M(\tau) = q^{h-c/24} \sum_{n=0}^{\infty} \dim M_{n+h} q^n$. The exponent of χ_M is also called an exponent of M . Suppose that the conformal weight of any Ramond-twisted irreducible W -module is a rational number. The minimum number h_{\min} among the conformal weights of all Ramond-twisted irreducible W -modules is called the *minimum conformal weight* of W . The number $\tilde{c} = c - 24h_{\min}$ is called the *effective central charge* of $\frac{1}{2}\mathbb{Z}_{\geq 0}$ -graded VOA W ([DM1]). For a proof of the following lemma, refer to [K2].

Lemma 5.1. *Let W be a vertex operator superalgebra, V a vertex operator subalgebra of W and suppose that $W = V \oplus P$ with a simple current V -module P . Let w be an element of P and a an integer such that $z^a Y(w, z)w \in W[[z]]$ and $z^a Y(w, z)w|_{z=0} \neq 0$. If $a \in 2\mathbb{Z}$, then W is a vertex operator algebra.*

6. AFFINE VERTEX OPERATOR ALGEBRAS AND VIRASORO MINIMAL MODELS

In this section we recall the notion of affine VOAs and give a characterization of the Deligne exceptional series by using the MLDEs (\sharp_s). Moreover, we study extensions of the Virasoro minimal model $L(-3/5, 0)$, which will be used to study \mathcal{W} -algebras.

6.1. Affine vertex operator algebras and second order modular linear differential equations. Let \mathfrak{g} be a finite-dimensional simple Lie algebra and $\widehat{\mathfrak{g}}$ the affine Kac-Moody Lie algebra associated with \mathfrak{g} . The simple affine VOA at level $k \in \mathbb{C}$ associated with \mathfrak{g} is denoted by $V_k(\mathfrak{g})$. Let λ be a weight of \mathfrak{g} . If the simple $\widehat{\mathfrak{g}}$ -module $L(\lambda + k\Lambda_0)$ is a $V_k(\mathfrak{g})$ -module, we write $V_k(\mathfrak{g}; \lambda) = L(\lambda + k\Lambda_0)$. Recall the MLDEs of second order (\sharp_s) and the number $\mu(t) = t(t+2)/144$ defined in Introduction.

Let V be a C_2 -cofinite, rational VOA of CFT type with the central charge $c \in \mathbb{C}$ such that the characters of the modules over V satisfy (\sharp_s) and that the conformal weights of the irreducible V -modules are positive. In [MMS1], it is shown that the character of V coincides with the character of $V_1(\mathfrak{g})$ with a Lie algebra \mathfrak{g} in the Deligne exceptional series and $s = \mu(c/2)$. Moreover, the

unitary RCFTs whose characters satisfy (\sharp_s) are classified in [MMS2]: they are the WZW models at level 1 associated to the Deligne exceptional series. We here prove a vertex operator algebraic analogue of this result.

Theorem 6.1. *Let $V = \bigoplus_{n=0}^{\infty} V_n$ be a self-dual vertex operator algebra with the central charge $c \in \mathbb{C}$. Suppose that V satisfies the following 3 conditions: (M1) V is simple, C_2 -cofinite, rational and of CFT type, (M2) the characters of the modules over V satisfy modular linear differential equation (\sharp_s) of order 2 and weight 0 for some $s \in \mathbb{C}$, (M3) $\dim V_1 \geq 3$. Then V is isomorphic to the simple affine vertex operator algebra at level 1 associated with a Lie algebra in the Deligne exceptional series and $s = \mu(c/2)$.*

As we shall see in Remark 6.1 below, we can replace the condition (M3) in the theorem with the condition (M4): the conformal weights of all irreducible modules over V are non-negative. Note that the conformal weights of the non-vacuum irreducible modules over the chiral algebra (VOA) of a unitary RCFT are positive. In order to prove the theorem, we first show a lemma.

Lemma 6.1. *Let $L(f) = 0$ be a modular linear differential equation of order $n \leq 4$ and weight 0. Let $\alpha_1, \dots, \alpha_n$ be all indices of $L(f) = 0$ and suppose that $\alpha_i \in \mathbb{Q}$ and $\alpha_i < \alpha_j$ for $1 \leq i \neq j \leq n$. Let f_1, \dots, f_n be holomorphic functions on \mathbb{H} such that f_i has the form $f_i(\tau) = q^{\alpha_i}(1 + O(q))$ for $1 \leq i \leq n$, and that $F = {}^t(f_1, \dots, f_n)$ forms a vector-valued modular form of weight 0. Then (f_1, \dots, f_n) is a fundamental system of solutions of $L(f) = 0$.*

Proof. Since F forms a vector-valued modular form of weight 0, we have a non-zero classical modular form G of weight $n^2 - n - 12 \sum_{i=1}^n \alpha_i$ such that $W(F) = G \cdot \eta^{24 \sum_{i=1}^n \alpha_i}$ ([Ma, Theorem 3.7]). By evaluating the coefficient of $E_2 D^{n-1}(f)$ in $L(f) = 0$, which is equal to $-n(n-1)/12$, we find that $n(n-1)/12$ is equal to the sum of all exponents. Thus we find $n^2 - n - 12 \sum_{i=1}^n \alpha_i = 0$ so that G is a non-zero constant. Therefore (f_1, \dots, f_n) is a fundamental system of solutions of $L(f) = 0$ by [Ma, Theorem 4.3]. \square

Recall that the indices of $(\sharp_{\mu(t)})$ are given by $-t/12$ and $(t+2)/12$ ([KNS]). We now prove Theorem 6.1.

Proof of Theorem 6.1. Put $d = \dim V_1$ and let \tilde{c} be the effective central charge of V . By using (M1) and (M2), we see that $(\tilde{c}, d) = (2/5, 0), (2/5, 1), (1, 1), (1, 3), (2, 3), (2, 8), (14/5, 14), (4, 28), (4, 8), (26/5, 52), (6, 78), (6, 14), (7, 133), (38/5, 190)$ or $(8, 248)$, and $s = \mu(\tilde{c}/2)$ ([MMS1] and [KNS]). The assumption (M3) forces $(\tilde{c}, d) \neq (2/5, 0), (2/5, 1)$ and $(1, 1)$. Since V_1 is a reductive Lie algebra and $\text{rank } V_1 \leq \tilde{c}$ by [DM1, Theorem 1.1 and Theorem 1.2], we see that V_1 is one of the Lie algebras listed in Table 1.

TABLE 1. Possibility of \mathfrak{g} .

(\tilde{c}, d)		$(2, 8)$	$(14/5, 14)$		$(4, 8)$				
d	3			28		52	78	133	248
V_1	A_1	A_2	G_2	G_2^2, D_4	$A_2, A_1^2 \times \mathbb{C}^2$	F_4	B_6, C_6, E_6	E_7	E_8

It follows from [DM2, Theorem 1.1] that the tensor product affine VOA $U = \bigotimes_{i=1}^n V_{k_i}(\mathfrak{g}_i)$ with $k_1, \dots, k_n \in \mathbb{Z}_{>0}$ is embedded in V , where $n \in \mathbb{Z}_{>0}$ and $\mathfrak{g}_1, \dots, \mathfrak{g}_n$ are simple ideals of \mathfrak{g} such that

\mathfrak{g} decomposes into the sum $\mathfrak{g} = \mathfrak{a} \oplus \bigoplus_{i=1}^n \mathfrak{g}$ with an abelian Lie algebra \mathfrak{a} . By comparing $\dim V_2$ with $\dim U_2$, we see that $n = 1$, $k_1 = 1$ and \mathfrak{g} is a Lie algebra in the Deligne exceptional series. (The dimension of V_2 can be computed by using the MLDE ($\sharp_{\mu(\tilde{c}/2)}$). We here only give a proof of the case $d = 28$. The other cases are proved in a similar way.

Since $\chi_V(\tau)$ is a solution of CFT type of ($\sharp_{\mu(2)}$) and $\dim V_1 = 28$, it follows that $\chi_V(\tau) = q^\alpha(1 + 28q + 134q^2 + O(q^3))$ with $\alpha \in \mathbb{Q}$, and hence we have $\dim V_2 = 134$. Suppose that $V_1 \cong G_2^2$. It then follows that $U \cong V_{k_1}(G_2) \otimes V_{k_2}(G_2)$ with $k_1, k_2 \in \mathbb{Z}_{>0}$. We see that $\dim U_2 = 280$ if $k_1 = k_2 = 1$, $\dim U_2 = 434$ if $k_1 \neq 1$ and $k_2 \neq 1$ and $\dim U_2 = 357$ otherwise, which contradicts $\dim V_2 = 134$. Hence, $V_1 \cong D_4$. It then follows that $U \cong V_k(D_4)$ with $k \in \mathbb{Z}_{>0}$. Then we have $k = 1$ since $\dim U_2 = 434$ otherwise. Hence $V_1(D_4)$ is embedded in V . Since the space of the characters of $V_1(D_4)$ -modules is 2-dimensional and the exponents of $V_1(D_4)$ -modules are $-1/6$ and $1/3$, it follows that the characters of $V_1(D_4)$ satisfy ($\sharp_{\mu(2)}$) by Lemma 6.1. Therefore, as $V_1(D_4) \subset V$, we have $V \cong V_1(D_4)$, and since the central charge of $V_1(D_4)$ is 4, we see that $s = \mu(c/2)$. \square

Remark 6.1. (i) In [MMS1] and [MMS2], it is also proved that a RCFT whose characters satisfy ($\sharp_{\mu(1/5)}$) coincides with the Virasoro minimal model of central charge $-22/5$.

(ii) In Theorem 6.1, we can replace condition (M3) with (M4). (The condition (M4) is stated just after Theorem 6.1.) It is because in the proof of Theorem 6.1, we use (M3) in order to exclude the cases $(\tilde{c}, d) = (2/5, 0)$, $(2/5, 1)$ and $(1, 1)$. However, (M4) also excludes the cases of $(\tilde{c}, d) = (2/5, 0)$ and $(1, 1)$. Since $\tilde{c} \geq \text{rank}(V_1)$, we also have $(\tilde{c}, d) \neq (2/5, 1)$. Hence Theorem 6.1 holds even if we replace (M3) with (M4).

6.2. Extensions of a Virasoro minimal model. We denote by $L(-3/5, 0)$ the Virasoro minimal model of central charge $-3/5$. In this subsection we study extensions of $L(-3/5, 0)$, which will be used to study \mathcal{W} -algebras in the subsequent sections § 7 and § 8.

All irreducible $L(-3/5, 0)$ -modules up to the isomorphisms are given by $L(-3/5, 0)$, $L(-3/5, -1/20)$, $L(-3/5, 3/4)$ and $L(-3/5, 1/5)$. The module $L(-3/5, 3/4)$ is a simple current with the fusion products $L(-3/5, 3/4) \boxtimes L(-3/5, 3/4) = L(-3/5, 0)$ and $L(-3/5, 3/4) \boxtimes L(-3/5, -1/20) = L(-3/5, 1/5)$.

Let W be a $\frac{1}{2}\mathbb{Z}_{\geq 0}$ -graded VOA of the form $W^{(0)} = V \otimes L(-3/5, 0)$ and $W^{(1)} = P \otimes L(-3/5, 0)$ with a simple, rational, C_2 -cofinite VOA V and a simple current V -module P . It then follows that $P \boxtimes P = V$. Let M be an irreducible V -module and h an element of $\{0, -1/20, 3/4, 1/5\}$.

Definition 6.1. Define the $V \otimes L(-3/5, 0)$ -modules $L(M; h)$ by

$$L(M; h) = M \otimes L(-3/5, h) \oplus (M \boxtimes P) \otimes (L(-3/5, h) \boxtimes L(-3/5, 3/4)).$$

Lemma 6.2. *Suppose that $h_{M \boxtimes P} - h_M \in 3/4 + \mathbb{Z}$ (resp., $1/4 + \mathbb{Z}$). Then the following hold. (1) $L(M; -1/20)$ and $L(M; 3/4)$ are irreducible Ramond-twisted W -modules (resp., untwisted W -modules). (2) $L(M \boxtimes P; -1/20)$ and $L(M \boxtimes P; 3/4)$ are irreducible untwisted W -modules (resp., Ramond-twisted W -modules).*

Proof. (1) Suppose $h_{M \boxtimes P} - h_M \in 3/4 + \mathbb{Z}$. We show that $L(M; -1/20)$ is an irreducible Ramond-twisted W -module. The other statements can be proved in a similar way. We have $L(M; -1/20) =$

$M \otimes L(-3/5, -1/20) \oplus (M \boxtimes P) \otimes L(-3/5, 1/5)$. The conformal weight of the first summand is $h_M - 1/20$, and that of the second summand is $h_{M \boxtimes P} + 1/5 \in h_M + 19/20 + \mathbb{Z}$. Hence, the conformal weights of $L(M; -1/20)$ lie in $h_M - 1/20 + \mathbb{Z}$. Therefore, $L(M; -1/20)$ is Ramond-twisted. The irreducibility follows from the construction of $L(M; h)$.

(2) Since $(M \boxtimes P) \boxtimes P \cong M$ by the associativity of the fusion products and $P \boxtimes P \cong V$, the statement follows from (1). \square

7. IDENTIFICATION AND \mathcal{W} -ALGEBRAS

In this section we show the following theorem.

Theorem 7.1. *If \mathfrak{g} is a Lie algebra in the Deligne exceptional series, then the characters of the Ramond-twisted modules over the \mathcal{W} -algebra $\mathcal{W}_{-h^\vee/6}(\mathfrak{g}, f_\theta)$ satisfy (b_s) , where s is given by (1.1).*

We first introduce some notations and show a lemma, which is needed to give a proof of Theorem 7.1.

Let \mathfrak{g} be a Lie algebra in the Deligne exceptional series and $W(\mathfrak{g}) = \mathcal{W}_{-h^\vee/6}(\mathfrak{g}, f_\theta)$ the simple \mathcal{W} -algebra at level $k = -h^\vee/6$ associated with \mathfrak{g} and a minimal nilpotent element f_θ of \mathfrak{g} . If $\mathfrak{g} = A_1$, then it is well known that $W(A_1) \cong L(-3/5, 0)$ (see e.g. [KRW]). Therefore, we see that the characters of $W(A_1)$ satisfy $(b_{-3/5})$.

We now suppose that $\mathfrak{g} \neq A_1$. Then $W(\mathfrak{g})$ is a proper $\frac{1}{2}\mathbb{Z}_{\geq 0}$ -graded conical VOA. Define a VOA $U_{\mathfrak{g}}$ and simple current V -module $P_{\mathfrak{g}}$ by the formula $V_1(\mathfrak{g}) \cong U_{\mathfrak{g}} \otimes V_1(A_1(\theta)) \oplus P_{\mathfrak{g}} \otimes V_1(A_1(\theta); \theta/2)$, where θ is the highest root of \mathfrak{g} and $A_1(\theta)$ is the simple Lie algebra of type A_1 generated by the highest and lowest root vectors of \mathfrak{g} .

Lemma 7.1. [K2] *The vertex operator algebra $W(\mathfrak{g})$ is C_2 -cofinite and \mathbb{Z}_2 -rational. Moreover, $W(\mathfrak{g})$ is isomorphic to the simple current extension of $U_{\mathfrak{g}} \otimes L(-3/5, 0)$ by $P_{\mathfrak{g}} \otimes L(-3/5, 3/4)$.*

We denote by Mod and Mod_{tw} the sets of the equivalence classes of the irreducible untwisted and Ramond-twisted modules over $W(\mathfrak{g})$, respectively. Let S be the space of characters of Ramond-twisted $W(\mathfrak{g})$ -modules and T the space of characters of $U_{\mathfrak{g}}$ -modules.

Proof of Theorem 7.1. For $\mathfrak{g} = A_2, G_2, D_4, F_4, E_6, E_7$ and E_8 , the numbers s in (1.1) are $s = 2/5, 6/5, 12/5, 18/5, 22/5, 27/5$ and $32/5$, respectively. We show the assertion in a case by case basis.

1. $\mathfrak{g} = A_2$. Since $U_{A_2} = V_{\sqrt{3}A_1}$ and $P_{A_2} = V_{\sqrt{3}A_1 + \sqrt{3}\alpha/2}$, the complete set of the irreducible modules over $V_{\sqrt{3}A_1}$ up to isomorphisms is given by $N_k := V_{\sqrt{3}A_1 + k\sqrt{3}\alpha/6}$ with $k \in \mathbb{Z}_6$. The conformal weights of N_i ($0 \leq i \leq 5$) are $0, 1/12, 1/3, 3/4, 1/3$ and $1/12$, respectively. The space T has a basis $(\chi_{N_i})_{0 \leq i \leq 5}$. Since $P_{A_2} \boxtimes N_k = N_{k+3}$ for each $k \in \mathbb{Z}_6$, it follows by Lemma 6.2 that $\text{Mod} = \{L(N_k; -1/20), L(N_k; 3/4) \mid k \in \mathbb{Z}_6, k \text{ is odd}\}$ and $\text{Mod}_{\text{tw}} = \{L(N_k; -1/20), L(N_k; 3/4) \mid k \in \mathbb{Z}_6, k \text{ is even}\}$. The space S has a basis $\chi_{L(N_0; -1/20)}, \chi_{L(N_4; 3/4)}, \chi_{L(N_2; -1/20)}, \chi_{L(N_0; 3/4)}$ with the conformal weights $-1/20, 1/12, 17/60, 3/4$, respectively. Since the central charge of $W(A_2)$ is $2/5$, the exponents of this basis of S are $-1/15, 1/15, 4/15, 11/15$. As they correspond to the indices of MLDE $(b_{2/5})$, we see that each element of S satisfies $(b_{2/5})$ by Lemma 6.1.

2. $\mathfrak{g} = G_2$. Since $U_{G_2} = V_3(A_1)$ and $P_{G_2} = V_3(A_1; 3\alpha/2)$, the complete set of the irreducible modules over $V_3(A_1)$ up to the isomorphisms is given by $N_k := V_3(A_1; k\alpha/2)$ with $k \in \mathbb{Z}_4$. The conformal weights of N_i ($0 \leq i \leq 3$) are $0, 3/20, 2/5$ and $3/4$. Since $P_{G_2} \boxtimes N_k = N_{3-k}$ for each $k \in \mathbb{Z}_4$, it follows that $\text{Mod} = \{L(N_k; -1/20), L(N_k; 3/4) \mid k \in \mathbb{Z}_4, k \text{ is odd}\}$ and $\text{Mod}_{\text{tw}} = \{L(N_k; -1/20), L(N_k; 3/4) \mid k \in \mathbb{Z}_4, k \text{ is even}\}$. The space S has a basis $\chi_{L(N_0; -1/20)}, \chi_{L(N_2; 3/4)}, \chi_{L(N_2; -1/20)}, \chi_{L(N_0; 3/4)}$ with the conformal weights $-1/20, 3/20, 7/20, 3/4$, respectively. Since the central charge of $W(G_2)$ is $6/5$, the exponents of the basis of S are $-1/10, 1/10, 3/10, 7/10$, which shows that each element of S satisfies $(b_{6/5})$ by Lemma 6.1.

3. $\mathfrak{g} = D_4$. Since $U_{D_4} = V_1(A_1)^{\otimes 3}$ and $P_{D_4} = V_1(A_1; \alpha/2)^{\otimes 3}$, the complete set of the irreducible modules over $V_1(A_1)^{\otimes 3}$ up to the isomorphisms is given by $N_{k_1, k_2, k_3} := V_1(A_1; k_1\alpha/2) \otimes V_1(A_1; k_2\alpha/2) \otimes V_1(A_1; k_3\alpha/2)$ with $k_1, k_2, k_3 \in \mathbb{Z}_2$. The conformal weight of N_{k_1, k_2, k_3} is $(k_1^2 + k_2^2 + k_3^2)/4$. The space T has a basis $\chi_{N_{0,0,0}}, \chi_{N_{1,0,0}}, \chi_{N_{1,1,0}}, \chi_{N_{1,1,1}}$. Since $P_{D_4} \boxtimes N_{k_1, k_2, k_3} = N_{1-k_1, 1-k_2, 1-k_3}$ for each $k_1, k_2, k_3 \in \mathbb{Z}_2$, it follows that $\text{Mod} = \{L(N_{k_1, k_2, k_3}; -1/20), L(N_{k_1, k_2, k_3}; 3/4), \mid k_1, k_2, k_3 \in \mathbb{Z}_2, k_1 + k_2 + k_3 \text{ is odd}\}$ and $\text{Mod}_{\text{tw}} = \{L(N_{k_1, k_2, k_3}; -1/20), L(N_{k_1, k_2, k_3}; 3/4), \mid k_1, k_2, k_3 \in \mathbb{Z}_2, k_1 + k_2 + k_3 \text{ is even}\}$. The space S has a basis $\chi_{L(N_{0,0,0}; -1/20)}, \chi_{L(N_{0,1,1}; 3/4)}, \chi_{L(N_{1,1,0}; -1/20)}, \chi_{L(N_{0,0,0}; 3/4)}$ with the conformal weights $-1/20, 1/4, 9/20, 3/4$, respectively. Since the central charge of $W(D_4)$ is $12/5$, the exponents of the basis of S are $-3/20, 3/20, 7/20, 13/20$, which shows that each element of S satisfies $(b_{12/5})$ by Lemma 6.1.

4. $\mathfrak{g} = F_4$. Since $U_{F_4} = V_1(C_3)$ and $P_{F_4} = V_1(C_3; \varpi_3)$, the complete set of the irreducible modules over $V_1(C_3)$ is given by $N_k := V_1(C_3; \varpi_k)$ with $k \in \mathbb{Z}_4$, where $\varpi_0 = 0$. The conformal weights of N_k ($0 \leq k \leq 3$) are $0, 7/20, 3/5$ and $3/4$, respectively. Since $P_{F_4} \boxtimes N_k = N_{3-k}$ for each $k \in \mathbb{Z}_4$, it follows that $\text{Mod} = \{L(N_k; -1/20), L(N_k; 3/4) \mid k \in \mathbb{Z}_4, k \text{ is odd}\}$ and $\text{Mod}_{\text{tw}} = \{L(N_k; -1/20), L(N_k; 3/4) \mid k \in \mathbb{Z}_4, k \text{ is even}\}$. The space S has a basis $\chi_{L(N_0; -1/20)}, \chi_{L(N_2; 3/4)}, \chi_{L(N_2; -1/20)}, \chi_{L(N_0; 3/4)}$ with the conformal weights $-1/20, 7/20, 11/20, 3/4$, respectively. Since the central charge of $W(F_4)$ is $18/5$, the exponents of the basis of S are $-1/5, 1/5, 2/5, 3/5$, which shows that each element of S satisfies $(b_{18/5})$ by Lemma 6.1.

5. $\mathfrak{g} = E_6$. Since $U_{E_6} = V_1(A_5) \cong V_{A_5}$ and $P_{E_6} = V_1(A_5; \varpi_3) \cong V_{A_5+3\varpi_1}$, the complete set of the irreducible modules over V_{A_5} is given by $N_k := V_{A_5+k\varpi_1}$ with $0 \leq k \leq 5$. The conformal weights of N_i ($0 \leq i \leq 5$) are $0, 5/12, 2/3, 3/4, 2/3$ and $5/12$, respectively. The space T has a basis $(\chi_{N_i})_{0 \leq i \leq 3}$. Since $P_{E_6} \boxtimes N_k = N_{k+3}$ for each $k \in \mathbb{Z}_6$, it follows that $\text{Mod} = \{L(N_k; -1/20), L(N_k; 3/4) \mid k \in \mathbb{Z}_6, k \text{ is odd}\}$ and $\text{Mod}_{\text{tw}} = \{L(N_k; -1/20), L(N_k; 3/4) \mid k \in \mathbb{Z}_6, k \text{ is even}\}$. The space S has a basis $\chi_{L(N_0; -1/20)}, \chi_{L(N_4; 3/4)}, \chi_{L(N_2; -1/20)}, \chi_{L(N_0; 3/4)}$ with the conformal weights $-1/20, 5/12, 37/60, 3/4$, respectively. Since the central charge of $W(E_6)$ is $22/5$, the exponents of the basis of S are $-7/30, 7/30, 13/30, 17/30$, which shows that each element of S satisfies $(b_{22/5})$ by Lemma 6.1.

6. $\mathfrak{g} = E_7$. Since $U_{E_7} = V_1(D_6) \cong V_{D_6}$ and $P_{E_7} = V_1(D_6; \varpi_6) \cong V_{D_6+\varpi_6}$, the complete set of the irreducible modules over V_{D_6} up to the isomorphisms is given by $N_0 := V_{D_6}, N_1 := V_{D_6+k\varpi_1}, N_2 := V_{D_6+\varpi_5}$ and $N_3 := V_{D_6+\varpi_6}$. The conformal weights of N_k ($0 \leq k \leq 3$) are $0, 1/2, 3/4$ and $3/4$, respectively. The space T has a basis $(\chi_{N_k})_{0 \leq k \leq 2}$. Since $P_{E_7} \boxtimes N_k = N_{3-k}$ for each $k \in \mathbb{Z}_4$, it follows that $\text{Mod} = \{L(N_k; -1/20), L(N_k; 3/4) \mid k \in \mathbb{Z}_4, k \text{ is odd}\}$ and $\text{Mod}_{\text{tw}} = \{L(N_k; -1/20), L(N_k; 3/4) \mid k \in \mathbb{Z}_4, k \text{ is even}\}$. The space S has a basis $\chi_{L(N_0; -1/20)}, \chi_{L(N_2; 3/4)}$,

$\chi_{L(N_2; -1/20)}$, $\chi_{L(N_0; 3/4)}$ with the conformal weights $(-1/20, 1/2, 7/10, 3/4)$, respectively. Since the central charge of $W(E_7)$ is $27/5$, the exponents of the basis of S are $-11/40, 11/40, 19/40, 21/40$, which shows that each element of S satisfies $(b_{27/5})$ by Lemma 6.1.

7. $\mathfrak{g} = E_8$. Since $U_{E_8} = V_1(E_7) \cong V_{E_7}$ and $P_{E_8} = V_1(E_7; \varpi_7) \cong V_{E_7 + \varpi_7}$, the complete set of the irreducible modules over V_{E_7} is $N_k := V_{E_7 + k\varpi_7}$ with $k \in \mathbb{Z}_2$. The conformal weights of N_0 and N_1 are 0 and $3/4$, respectively. Since $P_{E_8} \boxtimes N_k = N_{1+k}$ for each $k \in \mathbb{Z}_2$, it follows that $\text{Mod} = \{L(N_k; -1/20), L(N_k; 3/4) \mid k \in \mathbb{Z}_2, k \text{ is odd}\}$ and $\text{Mod}_{\text{tw}} = \{L(N_k; -1/20), L(N_k; 3/4) \mid k \in \mathbb{Z}_2, k \text{ is even}\}$. The space S has a basis $\chi_{L(N_0; -1/20)}$, $\chi_{L(N_0; 3/4)}$ with the conformal weights $-1/20, 3/4$, respectively. Since the central charge of $W(E_8)$ is $32/5$, the exponents of the basis of S are $-19/60, 29/60$, which shows that each element of S satisfies $(\sharp_{\mu(19/5)})$ by Lemma 6.1. Since the left-hand side of $(b_{32/5})$ is rewritten as a linear combination of derivatives of the left-hand side of $(\sharp_{\mu(19/5)})$ as in (B.1) in Appendix B, each element of S satisfies $(b_{32/5})$.

These proves the statements. \square

Remark 7.1. (a) If we formally set $h^\vee = 24$ (the number 24 is not the dual Coxeter number of any Lie algebra in the Deligne exceptional series) into (1.1), we see that (b_6) also has a solution of CFT type. The number $h^\vee = 24$ appears in many studies related with the Deligne exceptional series (see e.g. [CdM], [MMS1], [LM] and [K1]). A similar phenomenon is observed in the study of the second order MLDEs (see [MMS1] and [K1]). We now explain this phenomenon. Let \mathfrak{g} be a Lie algebra in the Deligne exceptional series. The characters of the affine VOA $V_1(\mathfrak{g})$ associated with \mathfrak{g} satisfy $(\sharp_{\mu(c_{\mathfrak{g}}/2)})$, where $c_{\mathfrak{g}}$ is the central charge $c_{\mathfrak{g}} = \dim \mathfrak{g} / (1 + h^\vee)$ of $V_1(\mathfrak{g})$. Since the Lie algebra \mathfrak{g} in the Deligne exceptional series satisfies $\dim \mathfrak{g} = 2(5h^\vee - 6)(h^\vee + 1)/(h^\vee + 6)$ by the *Deligne dimension formula* (see e.g. [CdM] and [D]), the number $c_{\mathfrak{g}}$ is a rational function in h^\vee . If we formally substitute $h^\vee = 24$ into the rational function $c_{\mathfrak{g}}$ in h^\vee , we have $c_{\mathfrak{g}} = 38/5$, and $(\sharp_{\mu(19/5)})$ has solutions of CFT type and character type. A vertex algebra $V_{E_{7+1/2}}$ associated with the intermediate Lie algebra $E_{7+1/2}$ ([LM, W]) was constructed in [K1], and modular invariant characters of $V_{E_{7+1/2}}$ were shown to satisfy $(\sharp_{\mu(19/5)})$ (see [K1]). If we formally substitute $h^\vee = 3/2$ into the rational functions $c_{\mathfrak{g}}$ and (1.1), we have $c_{\mathfrak{g}} = 2/5$ and $s = -6/5$. The MLDE $(\sharp_{\mu(1/5)})$ has a solution of CFT type with the exponent $-c_{\mathfrak{g}}/24 (= -1/60)$, and $(b_{-6/5})$ has a solution $f = 1$.

(b) Let \mathfrak{g} be a Lie algebra in the Deligne exceptional series and M a unique Ramond-twisted irreducible $\mathcal{W}_{-h^\vee/6}(\mathfrak{g}, f_\theta)$ -module with the minimum conformal weight. Since the character of M is of CFT type, we can obtain for any given $n \geq 0$ a formula for $\dim M_n$ in terms of the variable h^\vee by using (3.1) with $a_0 = 1$ (cf. [T]).

8. CHARACTERIZATION OF \mathcal{W} -ALGEBRAS ASSOCIATED WITH THE DELIGNE EXCEPTIONAL SERIES

Recall the **Main Conditions** given in Introduction. In this section we prove the following theorem.

Theorem 8.1. *Let W be a simple, C_2 -cofinite, \mathbb{Z}_2 -rational, $\frac{1}{2}\mathbb{Z}_{\geq 0}$ -graded, conical vertex operator algebra. Suppose that W satisfies the Main Conditions. Then W is isomorphic to $\mathcal{W}_{-h^\vee/6}(\mathfrak{g}, f_\theta)$ associated with a Lie algebra $\mathfrak{g} \neq A_1$ in the Deligne exceptional series.*

In order to prove Theorem 8.1, we will show several propositions. Let $W = \bigoplus_{n \in (1/2)\mathbb{Z}} W_n$ be a simple, C_2 -cofinite, \mathbb{Z}_2 -rational, $\frac{1}{2}\mathbb{Z}_{\geq 0}$ -graded, conical VOA with central charge $c \in \mathbb{C}$. Assume that W satisfies the Main Conditions. Then we have a complex number s which satisfies (a) in the Main Conditions. Moreover, we have a simple, C_2 -cofinite, rational VOA V of CFT type, and a simple current V -module P satisfying (b).

We first give candidates of the number s .

Proposition 8.1. *The number s is one of $-3/5, 2/5, 6/5, 12/5, 18/5, 22/5, 27/5, 6$ and $32/5$.*

Proof. We first show that s belongs to (4.2) and that $s \neq -48/5$ nor $-38/5$. By using (b), we have an irreducible Ramond-twisted W -module $L(V; -1/20)$ (this notation is defined in Definition 6.1). Since $c+3/5$ is the central charge of V , we have $c \in \mathbb{Q}$ ([AM]). Since $\chi_{L(V; -1/20)}(\tau) = q^{-c/24-1/20}(1+O(q))$ by (c), the character of $L(V; -1/20)$ is of CFT type. Since W is \mathbb{Z}_2 -rational, the space of characters of Ramond-twisted W -modules is $SL_2(\mathbb{Z})$ -invariant ([Va]), which shows that $\chi_{L(V; -1/20)}$ is not a quasimodular form of positive depth. It then follows from Theorem 4.1 that s belongs to (4.2). The above $SL_2(\mathbb{Z})$ -invariance of characters also shows that $s \neq -48/5$ nor $-38/5$ by Proposition 4.2.

We finally show $s \neq -6/5, 54/5, 18, -66/5, -6$ nor $-8/5$. Since the indices of (b_s) do not have non-zero integral differences by (a), we have $s \neq 54/5, 18, -66/5$ nor -6 . By using (c) again, we have $\chi_{L(V; -1/20)} \geq \chi_{L(-3/5, -1/20)}$. Here we say that two functions f and g of character type satisfy $f \geq g$ if $a_n \geq b_n$ for any $n \in \mathbb{Z}_{\geq 0}$, where $f(\tau) = \sum_{n=0}^{\infty} a_n q^{n+\alpha}$ and $g(\tau) = \sum_{n=0}^{\infty} b_n q^{n+\beta}$ denote the Fourier expansions of f and g with $a_0, b_0 \neq 0$. However, the solutions of CFT type of $(b_{-6/5})$ and $(b_{-8/5})$, which can be found in Appendix B (c) and (q), are not greater than nor equal to $\chi_{L(-3/5, -1/20)}$ because $\chi_{L(-3/5, -1/20)}(\tau) = q^{1/40}(1+q+q^2+2q^3+O(q))$. Thus $s \neq -6/5$ nor $-8/5$, which completes the proof. \square

We next construct a VOA R from V and P . We will later show that the characters of R satisfy a specific MLDE of order 2 if $s \neq 32/5$. (Then by Theorem 7.1, we specify R , which enables us to determine V and P .) Let A_1 be a simple Lie algebra of type A_1 and α a positive root of A_1 . The affine VOA $V_1(A_1)$ has a unique non-vacuum irreducible module $V_1(A_1; \alpha/2)$. It is well known that $V_1(A_1; \alpha/2)$ is a simple current with $V_1(A_1; \alpha/2) \boxtimes V_1(A_1; \alpha/2) = V_1(A_1)$ and the conformal weight of $V_1(A_1; \alpha/2)$ is $1/4$.

Proposition 8.2. *There is a unique vertex operator algebra R of the form $R \cong V \otimes V_1(A_1) \oplus P \otimes V_1(A_1; \alpha/2)$, which is called the simple current extension of $V \otimes V_1(A_1)$ by $P \otimes V_1(A_1; \alpha/2)$.*

Proof. Put $A = V \otimes V_1(A_1)$ and $B = P \otimes V_1(A_1; \alpha/2)$. Since the conformal weight of B is an integer, we have a super VOA R such that $R = A \oplus B$, which is unique up to isomorphisms of super VOAs (see [C]). Let u be a non-zero element of P and v a highest weight vector of $L(-3/5, 3/4)$. Since W is simple, $w = u \otimes v \in W$ satisfies $Y(w, z)w \neq 0$. Take $a \in \mathbb{Z}$ such that $z^a Y(w, z)w \in W[[z]]$ and $z^a Y(w, z)w|_{z=0} \neq 0$. As W is a VOA, it follows from Lemma 5.1 that $e^{-\pi\sqrt{-1}a} = 1$. Since v satisfies $v(1/2)v = k|0\rangle$ with non-zero $k \in \mathbb{C}$, where $|0\rangle$ is the vacuum vector of $L(-3/5, 0)$, it follows that $z^{3/2}Y(v, z)v \in L(-3/5, 0)[[z]]$ and $z^{3/2}Y(v, z)v|_{z=0} \neq 0$. Therefore, the element u satisfies $z^{a-3/2}Y(u, z)u \in V[[z]]$ and $z^{a-3/2}Y(u, z)u|_{z=0} \neq 0$. A highest weight vector $e^{\alpha/2} \in V_1(A_1; \alpha/2)$

of \mathfrak{h} -weight $\alpha/2$ satisfies $e^{\alpha/2}(-3/2)e^{\alpha/2} = k_1 e^\alpha$ with a non-zero number $k_1 \in \mathbb{C}$, where e^α is a homogeneous element of \mathfrak{h} -weight α . It then follows that $z^{-1/2}Y(e^{\alpha/2}, z)e^{\alpha/2} \in V_1(A_1)[[z]]$ and $z^{-1/2}Y(e^{\alpha/2}, z)e^{\alpha/2}|_{z=0} \neq 0$. Therefore, $u_1 = u \otimes e^{\alpha/2} \in R$ satisfies $z^{a-2}Y(u_1, z)u_1 \in R[[z]]$ and $z^{a-2}Y(u_1, z)u_1|_{z=0} \neq 0$. It follows from Lemma 5.1 that R is a VOA since $e^{-\pi\sqrt{-1}(a-2)} = 1$. \square

We next describe the irreducible modules over V . It turns out in the following proposition that the space of characters of V -modules is at most 4-dimensional, which will show that the space of characters of R -modules is at most 2-dimensional. We denote by h_M the conformal weight of an irreducible V -module M . Let \mathcal{A} denote the set of isomorphism classes of irreducible V -modules M which satisfy $h_M \notin \mathbb{Z}$ and $h_{M \boxtimes P} - h_M \in 3/4 + \mathbb{Z}$.

Proposition 8.3. *The set $\{V, P\} \sqcup \mathcal{A} \sqcup (\mathcal{A} \boxtimes P)$ is the complete set of the isomorphism classes of irreducible V -modules, where $\mathcal{A} \boxtimes P = \{M \boxtimes P | M \in \mathcal{A}\}$. Moreover, for any $M, N \in \mathcal{A}$, both equalities $\chi_M = \chi_N$ and $\chi_{M \boxtimes P} = \chi_{N \boxtimes P}$ hold.*

To show the proposition, we will prove several lemmas.

We write the conformal weight of each irreducible Ramond-twisted W -module L as r_L .

Lemma 8.1. (I) *For any irreducible V -module M , the conformal weights of M and $M \boxtimes P$ satisfy either $h_{M \boxtimes P} - h_M \in 3/4 + \mathbb{Z}$ or $1/4 + \mathbb{Z}$. (II) Let M be an irreducible V -module such that $h_{M \boxtimes P} - h_M \in 3/4 + \mathbb{Z}$. Then the following assertions (1)–(6) hold: (1) If $h_{M \boxtimes P} - h_M \geq -1/4$, then $r_{L(M; -1/20)} = h_M - 1/20$. (2) If $h_{M \boxtimes P} - h_M \leq -1/4$, then $r_{L(M; -1/20)} = h_{M \boxtimes P} + 1/5$. (3) If $h_{M \boxtimes P} - h_M \geq 3/4$, then $r_{L(M; 3/4)} = h_M + 3/4$. (4) If $h_{M \boxtimes P} - h_M \leq 3/4$, then $r_{L(M; 3/4)} = h_{M \boxtimes P}$. (5) $r_{L(M; 3/4)} - r_{L(M; -1/20)} \in 4/5 + \mathbb{Z}$. (6) $r_{L(V; -1/20)} = -1/20$ and $r_{L(V; 3/4)} = 3/4$.*

Proof. The assertion (I) follows from condition (h). Since $r_{L(M; -1/20)} = \min\{h_M - 1/20, h_{M \boxtimes P} + 1/5\}$ and $r_{L(M; 3/4)} = \min\{h_M + 3/4, h_{M \boxtimes P}\}$, we have assertions (1)–(5). The assertion (6) follows from (e). \square

Lemma 8.2. *Let M be an irreducible V -module such that $h_{M \boxtimes P} - h_M \in 3/4 + \mathbb{Z}$. If $h_M \in \mathbb{Z}$, then M is isomorphic to V .*

Proof. If $h_M = 0$, then $M \cong V$ by (d). We show $h_M = 0$ by contradiction. Suppose on the contrary that $h_M \in \mathbb{Z}_{>0}$ and let k be an element of $\{-1/20, 3/4\}$. Since both $L(M; k)$ and $L(V; k)$ are Ramond-twisted by Lemma 6.2, the number $r_{L(M; k)} - r_{L(V; k)}$ is an integer.

We now show $r_{L(M; k)} = r_{L(V; k)}$ by contradiction. Suppose that $r_{L(M; k)} \neq r_{L(V; k)}$, which implies that $\chi_{L(M; k)}$ and $\chi_{L(V; k)}$ are linearly-independent. Then we have $i, j \in \{1, \dots, 4\}$ such that $i \neq j$, $\alpha_i - \alpha_j \in \mathbb{Z}$ and $r_{L(V; k)} \in \alpha_i + \mathbb{Z}$. Here $\alpha_1, \dots, \alpha_4$ are the indices of (b_s) . It follows by (a) that $\alpha_i = \alpha_j$. Since (b_s) has at most one pair of multiple indicial roots, we have a solution with a logarithmic term, and the space spanned by the solutions of (b_s) of character type whose exponents belong to $\alpha_i + \mathbb{Z}$ is one-dimensional. It contradicts the linearly-independence of $\chi_{L(M; k)}$ and $\chi_{L(V; k)}$. Hence, $r_{L(M; k)} = r_{L(V; k)}$.

The equality $r_{L(M; -1/20)} = r_{L(V; -1/20)}$, Lemma 8.1 and $h_M \neq 0$ show that $h_{M \boxtimes P} + 1/5 = -1/20$ and $h_{M \boxtimes P} - h_M < -1/4$. Therefore, $h_{M \boxtimes P} = -1/4$ and $h_M > 0$. By using Lemma 8.1 again, we have $r_{L(M; 3/4)} = -1/4$. As $r_{L(V; 3/4)} = \min\{3/4, h_P\}$, condition (c) leads to $r_{L(V; 3/4)} \geq 0$, which contradicts $r_{L(M; 3/4)} = r_{L(V; 3/4)}$. Thus we have $h_M = 0$, which completes the proof. \square

Lemma 8.3. *Let M and N be elements of \mathcal{A} and suppose that (A) $r_{L(M;-1/20)} = r_{L(N;-1/20)}$ and (B) $r_{L(M;3/4)} = r_{L(N;3/4)}$ hold. Then $\chi_M = \chi_N$ and $\chi_{M \boxtimes P} = \chi_{N \boxtimes P}$.*

Proof. We first show that either

$$(8.1) \quad h_M = h_N \quad \text{or} \quad h_{M \boxtimes P} = h_{N \boxtimes P}$$

holds, whose proof is divided into 4 cases: **1.** $h_{M \boxtimes P} - h_M < -1/4$, **2.** $h_{M \boxtimes P} - h_M = -1/4$, **3.** $h_{M \boxtimes P} - h_M = 3/4$, and **4.** $h_{M \boxtimes P} - h_M > 3/4$.

1. By Lemma 8.1, we have $r_{L(M;-1/20)} = h_{M \boxtimes P} + 1/5$ and $r_{L(M;3/4)} = h_{M \boxtimes P}$. Suppose that $h_{N \boxtimes P} - h_N \leq 3/4$, which implies $r_{L(N;3/4)} = h_{N \boxtimes P}$. Then we have $h_{M \boxtimes P} = h_{N \boxtimes P}$ by assumption (2). Suppose that $h_{N \boxtimes P} - h_N > 3/4$. It then follows that $r_{L(N;-1/20)} = h_N - 1/20$ and $r_{L(N;3/4)} = h_N + 3/4$, which contradict (A) or (B). Thus, we have (8.1) in this case.

2. Suppose $h_{M \boxtimes P} - h_M = -1/4$. Then $r_{L(M;-1/20)} = h_M - 1/20 = h_{M \boxtimes P} + 1/5$ by Lemma 8.1. Since $r_{L(N;-1/20)} = \min\{h_N - 1/20, h_{N \boxtimes P} + 1/5\}$, the assumption (A) forces (8.1).

The assertions **3–4** are proved in a similar way, which completes the proof of (8.1).

We now prove the lemma. Suppose first that $h_M = h_N$. Then we have $\chi_M = \chi_N$ by (e). It then follows from (A) and (f) that $\chi_{L(M;-1/20)} = \chi_{L(N;-1/20)}$, which shows $\chi_{M \boxtimes P} = \chi_{N \boxtimes P}$. In a similar way, we have $\chi_M = \chi_N$ and $\chi_{M \boxtimes P} = \chi_{N \boxtimes P}$ when $h_{M \boxtimes P} = h_{N \boxtimes P}$. Thus we have the lemma. \square

Set $C_s = \{(i, j) \mid 1 \leq i, j \leq 4, \alpha_j - \alpha_i \in 4/5 + \mathbb{Z}\}$. We have completed the preliminaries on the proof of Proposition 8.3.

Proof of Proposition 8.3. Let M and N be elements of \mathcal{A} . Since $P \boxtimes P \cong V$, by Lemma 8.1 (I) and Lemmas 8.2–8.3, it suffices to show $r_{L(M;k)} = r_{L(N;k)}$ for each $k = -1/20, 3/4$. We divide the proof into 3 parts: **1.** $s \neq 6$ nor $6/5$, **2.** $s = 6$, and **3.** $s = 6/5$.

1. Suppose that $s \neq 6$ nor $6/5$. It then follows by Proposition 8.1 that $C_s = \{(1, 2), (3, 4)\}$. Since $h_M, h_N \notin \mathbb{Z}$, Lemma 8.1 (5) shows that the following (A) or (B) holds:

- (A) $r_{L(V;-1/20)} = \alpha_1$, $r_{L(V;3/4)} = \alpha_2$, $r_{L(M;-1/20)} = r_{L(N;-1/20)} = \alpha_3$, and $r_{L(M;3/4)} = r_{L(N;3/4)} = \alpha_4$;
(B) $r_{L(V;-1/20)} = \alpha_3$, $r_{L(V;3/4)} = \alpha_4$, $r_{L(M;-1/20)} = r_{L(N;-1/20)} = \alpha_1$, and $r_{L(M;3/4)} = r_{L(N;3/4)} = \alpha_2$.
Hence we have $r_{L(M;k)} = r_{L(N;k)}$ for $k = -1/20, 3/4$.

2. If $s = 6$, then there exists a unique pair of $1 \leq i < j \leq 4$ such that $\alpha_i = \alpha_j$. Therefore, there is a solution of (b_s) with a logarithmic term, and the space of characters of Ramond-twisted W -modules is at most 3-dimensional. Hence, there are at most 3 conformal weights of W -modules. Since $h_M, h_N \notin \mathbb{Z}$, the numbers $r_{L(M;-1/20)}$ and $r_{L(N;-1/20)}$ do not belong to $r_{L(V;-1/20)} + \mathbb{Z}$. Hence, $r_{L(M;-1/20)} = r_{L(N;-1/20)} = r_{L(V;3/4)}$, and eventually, $r_{L(M;3/4)} = r_{L(N;3/4)}$.

3. If $s = 6/5$, then we see that $(\alpha_i)_{1 \leq i \leq 4} = (-1/10, 7/10, 3/10, 1/10)$ and $C_{6/5} = \{(3, 4), (4, 1), (1, 2)\}$. Since there are no integral differences between the indices, the exponent of any irreducible Ramond-twisted W -module belongs to $\{\alpha_i \mid 1 \leq i \leq 4\}$. Let N^0, N^1 and N^2 be irreducible V -modules such that $h_{N^i \boxtimes P} - h_{N^i} \in 3/4 + \mathbb{Z}$ for each $0 \leq i \leq 2$. It then follows that $r_{L(N^i;-1/20)} \in \{\alpha_1, \alpha_3, \alpha_4\}$. Hence, it suffices to show that $\#\{r_{L(N^i;-1/20)} \mid i = 0, 1, 2\} \leq 2$. Suppose on the contrary that the exponents of $L(N^0;-1/20)$, $L(N^1;-1/20)$ and $L(N^2;-1/20)$ are α_3, α_4 and α_1 , respectively. It then follows that the exponent of $L(N^1;3/4)$ is α_1 . Therefore, (g) implies that the $W^{(0)}$ -module

$N^2 \otimes L(-3/5, -1/20)$ is isomorphic to one of the modules $N^1 \otimes L(-3/5, 3/4)$, $(N^1 \boxtimes P) \otimes L(-3/5, 0)$, $(N^1)' \otimes L(-3/5, 3/4)'$ and $(N^1 \boxtimes P)' \otimes L(-3/5, 0)'$. It contradicts $L(-3/5, 0)' \cong L(-3/5, 0)$ and $L(-3/5, 3/4)' \cong L(-3/5, 3/4)$. Thus, $\#\{r_{L(N^i; -1/20)} | i = 0, 1, 2\} \leq 2$, which shows $r_{L(M; k)} = r_{L(N; k)}$ for $k = -1/20$ and $3/4$.

The discussions 1–3 show the lemma. \square

We now give a proof of Theorem 8.1.

Proof of Theorem 8.1. We divide a proof into two parts (A): $\mathcal{A} = \emptyset$ and (B): $\mathcal{A} \neq \emptyset$.

The case (A) Suppose that $\mathcal{A} = \emptyset$. It then follows from Proposition 8.3 and (h) that the space of characters of Ramond-twisted W -modules has a basis $(\chi_{L(V; -1/20)}, \chi_{L(V; 3/4)})$. Then we see that $s = 32/5$ by Propositions 4.1 and 8.1. Since any solution of $(b_{32/5})$ of CFT type has an exponent $-19/60$ by § 3, we see that $\chi_{L(V; -1/20)}$ has the exponent $-19/60$ and $\chi_{L(V; 3/4)}$ is not of CFT type. Therefore, the central charge of V is 7 and $h_P = 3/4$. Then Lemma 6.1 shows that the characters of V satisfy $(\sharp_{\mu(7/2)})$, which implies $\dim V_1 = 133$. By using (d), we see that the VOA V is self-dual. Therefore, by Theorem 6.1 we have $V \cong V_1(E_7)$ and $P \cong V_1(E_7; \varpi_7)$, and then $V \cong U_{E_8}$ and $P \cong P_{E_8}$. By Lemma 7.1, we see that $W \cong \mathcal{W}_{-5}(E_8, f_\theta)$.

The case (B) Suppose that $\mathcal{A} \neq \emptyset$. Then we have an element M of \mathcal{A} . Set $L^0 = L(V; -1/20)$, $L^1 = L(V; 3/4)$, $L^2 = L(M; -1/20)$ and $L^3 = L(M; 3/4)$ with the characters χ_0, χ_1, χ_2 and χ_3 , respectively. It then follows from Proposition 8.3 that the space of characters of Ramond-twisted W -modules has a spanning set $\{\chi_0, \chi_1, \chi_2, \chi_3\}$. Define $m \in \mathbb{Z}$ by $h_{M \boxtimes P} - h_M = 3/4 + m$. By Lemma 8.1, the conformal weights r_i of L^i are

$$(r_0, r_1, r_2, r_3) = \begin{cases} (-1/20, 3/4, h_M - 1/20, h_M + 3/4) & \text{if } m \geq 0, \\ (-1/20, 3/4, h_M + m + 19/20, h_M + m + 3/4) & \text{if } m \leq -1. \end{cases}$$

By comparing the exponents of L^0, L^1, L^2 and L^3 and indicial roots of (b_s) , we see that $m \leq -1$, $s = c$ and $c = 12h_M + 12m + 42/5$.

By Proposition 8.2, we have the simple current extension R of $V \otimes V_1(A_1)$ by $P \otimes V_1(A_1; \alpha/2)$. It follows from Proposition 8.3 that the space of characters of R -modules has a basis which consists of the characters of R itself and $K = M \otimes V_1(A_1) \oplus (M \boxtimes P) \otimes V_1(A_1; \alpha/2)$. Since $m \leq -1$, the conformal weight of K is $h_M + m + 1$. As the central charge of V is $12h_M + 12m + 9$, that of R is $12h_M + 12m + 10$. Since the exponents of the characters of R coincide with the indices of $(\sharp_{\mu(6h_M + 6m + 5)})$, the characters of R satisfy $(\sharp_{\mu(6h_M + 6m + 5)})$ by Lemma 6.1. It follows from (d) that R is self-dual. As $A_1 \subset R_1$, Theorem 6.1 implies that $R \cong V_1(\mathfrak{g})$ with a Lie algebra \mathfrak{g} in the Deligne exceptional series. Since $P \neq 0$ and R has two irreducible modules, it follows that $\mathfrak{g} \neq A_1$ and E_8 . Therefore, we have $V = U_{\mathfrak{g}}$, $P = P_{\mathfrak{g}}$ and that W is isomorphic to $\mathcal{W}_{-h^\vee/6}(\mathfrak{g}, f_\theta)$ by Lemma 7.1. Thus we have proved Theorem 8.1. \square

9. THE OTHER IDENTIFICATION PROBLEMS

In Theorem 7.1, we find rational VOAs whose characters satisfy (b_s) when s has the form (1.1) with the dual Coxeter number h^\vee of one of the Lie algebras in the Deligne exceptional series. Moreover, we discuss the case of $s = 6$ and $s = -6/5$ (these numbers are given by (1.1) with

$h^\vee = 24$ and $3/2$) in Remark 7.1. In this section we discuss about the problem how to find rational VOAs whose characters satisfy (b_s) , where s is one of the remaining numbers appearing in (4.2).

(1) $s = -48/5$. By Proposition 4.2, the characters of a simple C_2 -cofinite rational VOA of CFT type do not satisfy $(b_{-48/5})$. The affine VOA $V = V_{-3/2}(G_2)$ at admissible level $-3/2$ is not C_2 -cofinite but semi-simple in the category \mathcal{O} . The VOA V has two ordinary irreducible modules with the conformal weights 0 and $4/5$. The characters of ordinary modules over affine VOAs at admissible levels satisfy MLDEs ([AK]). Since the central charge of $V_{-3/2}(G_2)$ is $-42/5$, the irreducible characters have the forms $q^{7/20}(1+14q+\sum_{n=2}^{\infty} a_n q^n)$ and $q^{23/20}(7+\sum_{n=1}^{\infty} b_n q^n)$ with non-negative integers a_n and b_n . Since the solutions f_0 and $7f_{4/5}$ of (b_s) defined in **(a)** of Appendix B also have this form, we propose a conjecture below.

Conjecture 1. *The characters of the irreducible ordinary modules over $V_{-3/2}(G_2)$ coincide with the solutions f_0 and $7f_{4/5}$ of $(b_{-48/5})$.*

Note that if this conjecture is true, The solution f_0 in **(a)** in Appendix B gives a character formula of $V_{-3/2}(G_2)$ in terms of modular forms in Table 2.

(2) $s = -38/5$. The same argument as in **(1)** suggests the following conjecture.

Conjecture 2. *There exists an extension V of the admissible affine vertex operator algebra $V_{-4/3}(A_2)$ of the form $V \cong V_{-4/3}(A_2) \oplus V_{-4/3}(A_2; 2\varpi_1) \oplus V_{-4/3}(A_2; 2\varpi_2)$. Moreover, the characters of the ordinary irreducible modules over V coincide with the solutions $f_{-8/15}$ and $f_{-1/3}$ of $(b_{-38/5})$ up to a scalar multiple.*

(3) Let s be an element of $\{54/5, 18, -66/5, -6\}$. Let V be a rational VOA and suppose that the characters of V satisfy (b_s) . There is a solution of vacuum type of (b_s) of exponent $-s/24$. However, since $1 - s/24$ is the exponent of a solution of (b_s) , it is difficult to know the dimension of the Lie algebra V_1 . If we know $\dim V_1$, we have finite candidates of the structure of V_1 , and it often enables us to find affine sub VOA of V . Then by using the representation theory of affine VOA's, there is a chance to determine the structure of V . However, since it is difficult to determine $\dim V_1$ in this case, we can not use this discussion, which makes it difficult to find V . When $s = 18$ or -6 , the character of the trivial VOA $V = \mathbb{C}$ satisfies (b_s) since $f = 1$ is a solution of (b_s) .

(4) $s = -8/5$. The characters of the Virasoro minimal model $L(-22/5, 0)$ coincide with the solutions f_0 and $f_{-1/5}$ of $(b_{-8/5})$.

APPENDIX A. MODULAR FORMS, DIFFERENTIAL RELATIONS AND FUNCTIONAL EQUATIONS

In this appendix we introduce some modular forms of levels $N = 2-5$ and 15 to describe solutions of (b_s) in Appendices B–C. The modular forms are listed in Table 2. Moreover, we list some differential relations and functional equations of modular forms in **(a)**–**(g)** below. By using these, we can prove that the functions listed in Appendices B–C are solutions of (b_s) .

The first column of Table 2 is the names of the modular forms f . The second column shows the Fourier expansion $f(\tau) = \sum_{n=0}^{\infty} a_n q^{n+\alpha}$ of f , where $\tau \in \mathbb{H}$ and $q = e^{2\pi i\tau}$. The third and fourth columns are the weight and level of f , respectively.

TABLE 2. Modular Forms

Name	Fourier Expansion	Weight	Level
H_2	$1 + 24 \sum_{n=1}^{\infty} \left(\sum_{d n, d:\text{odd}} d \right) q^n$	2	2
Δ_2	$\eta(q^2)^{16}/\eta(q)^8$	2	2
I_3	$1 + 6 \sum_{n=1}^{\infty} \left(\sum_{d n} \left(\frac{d}{3} \right) \right) q^n$	1	3
Δ_3	$\eta(q^3)^3/\eta(q)$	1	3
θ	$\sum_{n \in \mathbb{Z}} q^{n^2}$	1/2	4
Δ_4	$\eta(q^4)^2/\eta(q^2)$	1/2	4
ψ_1	$\eta(q)^{2/5} \left\{ q^{-1/60} \prod_{\substack{n>0 \\ n \neq 0, \pm 2 \pmod{5}}} (1 - q^n)^{-1} \right\}$	1/5	5
ψ_2	$\eta(q)^{2/5} \left\{ q^{11/60} \prod_{\substack{n>0 \\ n \neq 0, \pm 1 \pmod{5}}} (1 - q^n)^{-1} \right\}$	1/5	5
I_{15}	$\eta(q^3)^2 \eta(q^5)^2 / \eta(q) \eta(q^{15})$	1	15
Δ_{15}	$\eta(q)^2 \eta(q^{15})^2 / \eta(q^3) \eta(q^5)$	1	15

These modular forms and Eisenstein series satisfy the following differential relations and functional equations.

(a) Level 2

$$\begin{aligned} 6H_2' &= E_2 H_2 - H_2^2 + 192\Delta_2^2, & E_4 &= H_2^2 + 192\Delta_2^2, \\ 6\Delta_2' &= (E_2 + 2H_2)\Delta_2, & E_6 &= (H_2^2 - 576\Delta_2^2)H_2. \end{aligned}$$

(b) Level 3

$$\begin{aligned} 12I_3' &= E_2 I_3 - I_3^3 + 108\Delta_3^3, & E_4 &= I_3(I_3^3 + 216\Delta_3^3), \\ 12\Delta_3' &= (E_2 + 3I_3^2)\Delta_3, & E_6 &= I_3^6 - 540I_3^3\Delta_3^3 - 5832\Delta_3^6. \end{aligned}$$

(c) Level 4

$$\begin{aligned} 24\theta' &= (E_2 - \theta^4 + 80\Delta_4^4)\theta, & E_4 &= \theta^8 + 224\theta^4\Delta_4^4 + 256\Delta_4^8, \\ 24\Delta_4' &= (E_2 + 5\theta^4 - 16\Delta_4^4)\Delta_4, & E_6 &= (\theta^4 + 16\Delta_4^4)(\theta^8 - 544\theta^4\Delta_4^4 + 256\Delta_4^8). \end{aligned}$$

(d) Level 5

$$\begin{aligned} 60\psi_1' &= (E_2 - \psi_1^{10} + 66\psi_1^5\psi_2^5 + 11\psi_2^{10})\psi_1, & 60\psi_2' &= (E_2 + 11\psi_1^{10} - 66\psi_1^5\psi_2^5 - \psi_2^{10})\psi_2, \\ E_4 &= \psi_1^{20} + 228\psi_1^{15}\psi_2^5 + 494\psi_1^{10}\psi_2^{10} - 228\psi_1^5\psi_2^{15} + \psi_2^{20}, \\ E_6 &= (\psi_1^{10} + \psi_2^{10})(\psi_1^{20} - 522\psi_1^{15}\psi_2^5 - 10006\psi_1^{10}\psi_2^{10} + 522\psi_1^5\psi_2^{15} + \psi_2^{20}). \end{aligned}$$

(e) Level 10

$$\begin{aligned}
6H_2(q^5)' &= 5\{E_2(q^5)H_2(q^5) - H_2(q^5)^2 + 192\Delta_2(q^5)^2\}, \\
6\Delta_2(q^5)' &= 5\Delta_2(q^5)\{E_2(q^5) + H_2(q^5)\}, \\
5E_2(q^5) &= E_2(q) + 4\{\psi_1(q)^{10} + \psi_2(q)^{10}\}, \\
55H_2(q^5) &= -3\{5\psi_2(q)^{10} + 7\psi_1(q^2)^5\psi_1(q)^5 - 21\psi_1(q^2)^5\psi_2(q)^5 + 30\psi_2(q^2)^5\psi_1(q)^5\} \\
&\quad + 76\psi_1(q^2)^{10} - 78\psi_1(q^2)^5\psi_2(q^2)^5 + 70\psi_2(q^2)^{10}, \\
3000\Delta_2(q^5)^2 &= 2\{H_2(q)^2 - 60\Delta_2(1)^2\} - H_2(q)\psi_1(q)^{10} \\
&\quad - H_2(q^5)\{38\psi_1(q)^{10} + 132\psi_1(q)^5\psi_2(q)^5 + 37\psi_2(q)^{10}\} - 720\Delta_2(q)\Delta_2(q^5) \\
&\quad + 2H_2(q)\{8\psi_1(q^2)^{10} - 26\psi_1(q^2)^5\psi_2(q^2)^5 + 3\psi_2(q^2)^{10}\} \\
&\quad + 10H_2(q^5)\{2\psi_1(q^2)^{10} - 2\psi_1(q^2)\psi_2(q^2)^5 + 3\psi_2(q^2)^{10}\}
\end{aligned}$$

(f) Level 15

$$\begin{aligned}
12I_{15}' &= (E_2 - 5I_{15}^2 - 2I_{15}\Delta_{15} - 13\Delta_{15}^2 + I_3^2)I_{15}, \\
12\Delta_{15}' &= (E_2 + 13I_{15}^2 - 2I_{15}\Delta_{15} + 5\Delta_{15}^2 - 2I_3^2)\Delta_{15}. \\
12I_3(q^5)' &= E_2(q)I_3(q^5) + I_3(q)\{4I_{15}(q) + 4I_{15}(q)\Delta_{15}(q) + 2\Delta_{15}(q)^2\} \\
&\quad - I_3(q^5)\{5I_{15}(q)^2 + 2I_{15}(q)\Delta_{15}(q) - 5\Delta_{15}(q)^2\}, \\
I_3(q)^2 &= 6I_{15}(q)^2 + 6\Delta_{15}(q)^2 - 5I_3(q^5)^2, \\
I_3(q)I_3(q^5) &= I_{15}(q)^2 + 4I_{15}(q)\Delta_{15}(q) - \Delta_{15}(q)^2, \\
I_3(q)^3 &= 6I_3(q)\{I_{15}(q)^2 + \Delta_{15}(q)^2\} - 5I_3(q^5)\{I_{15}(q)^2 + 4I_{15}(q)\Delta_{15}(q) - \Delta_{15}(q)^2\}, \\
108\Delta_3(q)^3 &= I_3(q)\{25I_{15}(q)^2 - 2I_{15}(q)\Delta_{15}(q) - \Delta_{15}(q)^2\} \\
&\quad - 5I_3(q^5)\{5I_{15}(q)^2 + 8I_{15}(q)\Delta_{15}(q) + \Delta_{15}(q)^2\}, \\
120\psi_2(q)^{10} &= 45\{I_{15}(q)^2 - 5\Delta_{15}(q)^2\} - 6I_3(q)\{14I_{15}(q) - 2I_{15}(q)\Delta_{15}(q) - 5\Delta_{15}(q)^2\} \\
&\quad + 3I_3(q^5)\{8I_{15}(q) + 64\Delta_{15}(q) - 5I_3(q)\}.
\end{aligned}$$

(g) Level 20

$$\begin{aligned}
\theta(q^5)' &= \theta(q^5)\{E_2(q) + 4\psi_1(q)^{10} + 4\psi_2(q)^{10} - 5\theta(q^5)^4 + 400\Delta_4(q^5)^4\}, \\
\Delta_4(q^5)' &= \Delta_4(q^5)\{E_2(q) + 4\psi_1(q)^{10} + 4\psi_2(q)^{10} + 25\theta(q^5)^4 - 80\Delta_4(q^5)^4\}, \\
\theta(q)^5 &= 80\theta(q)\Delta_4(q)^4 - 5\theta(q)\{\psi_1(q)^5 + \psi_2(q)^5\} \\
&\quad + 6\theta(q^5)\{\psi_1(q)^{10} - 11\psi_1(q)^5\psi_2(q)^5 - \psi_2(q)^{10}\}, \\
\Delta_4(q)^5 &= 5\Delta_4(q)\{\theta(q)^4 - \psi_1(q)^{10} - \psi_2(q)^{10}\} \\
&\quad + 6\Delta_4(q^5)\{\psi_1(q)^{10} - 11\psi_1(q)^5\psi_2(q)^5 - \psi_2(q)^{10}\}, \\
40\Delta_4(q)^3\Delta_4(q^5) &= 5\theta(q)^3\theta(q^5) + \psi_1(q)^{10} - 36\psi_1(q)^5\psi_2(q)^5 - \psi_2(q)^{10}
\end{aligned}$$

$$\begin{aligned}
& - 6\{\psi_1(q^4)^{10} - 36\psi_1(q^4)^5\psi_2(q^4)^5 - \psi_2(q^4)^{10}\}, \\
48\psi_1(q^4)^{10} &= 2\{5\psi_1(q)^{10} + 36\psi_1(q)^5\psi_2(q)^5 + \psi_2(q)^{10}\} + 45\theta^2(q^5)\{\theta(q^5)^2 + \theta(q)^2\} \\
& - 2\theta(q)\theta(q^5)\{135\theta(q^5)^2 + 71\theta(q)^2\} \\
& - 160\Delta_4(q)^3\Delta_4(q^5) + 360\psi_1(q^4)^5\psi_1(q)^5 - 504\psi_2(q^4)^5\psi_2(q)^5, \\
48\psi_2(q^4)^{10} &= 2\{\psi_1(q)^{10} - 36\psi_1(q)^5\psi_2(q)^5 + 5\psi_2(q)^{10}\} + 45\theta^2(q^5)\{\theta(q^5)^2 + \theta(q)^2\} \\
& + 2\theta(q)\theta(q^5)\{135\theta(q^5)^2 + 71\theta(q)^2\} \\
& + 160\Delta_4(q)^3\Delta_4(q^5) - 504\psi_1(q^4)^5\psi_1(q)^5 + 360\psi_2(q^4)^5\psi_2(q)^5.
\end{aligned}$$

By using the above (a)–(g), we can prove that the functions listed in Appendices B–C are solutions of (b_s).

APPENDIX B. MODULAR LINEAR DIFFERENTIAL EQUATIONS WITH A SOLUTION OF CFT TYPE

In this appendix we list a fundamental system of solutions of (b_s) in a case by case basis when (b_s) has a solution of CFT type and does not have a quasimodular solution of positive depth. Among s in (3.12), the numbers in (4.2) have the required conditions. The solutions of the remaining cases are given in Appendix C. For $r \in \mathbb{Q}$, the solution f_r in this section has the form $f_r(\tau) = q^{r-\alpha/24}(1 + O(q))$, where α is a formal central charge of (b_s).

(a) $s = -48/5$. A fundamental system of solutions of (b_{-48/5}) is given by

$$\begin{aligned}
f_0 &= \frac{\psi_2(q)\Delta_2(q)(11\psi_1(q)^{10} - 66\psi_1(q)^5\psi_2(q)^5 - \psi_2(q)^{10} + H_2(q))}{12\eta(q)^{42/5}} \\
&= q^{7/20}(1 + 14q + 119q^2 + 770q^3 + 4088q^4 + 18676q^5 + \dots), \\
f_{4/5} &= \frac{\psi_1(q)\Delta_2(q)(-\psi_1(q)^{10} + 66\psi_1(q)^5\psi_2(q)^5 + 11\psi_2(q)^{10} + H_2(q))}{84\eta(q)^{42/5}} \\
&= q^{23/20}\left(1 + 14q + \frac{769}{7}q^2 + 642q^3 + 3103q^4 + 13078q^5 + 49616q^6 + \dots\right), \\
f_{-1/2} &= \frac{\psi_2(q)\{-H_2(q)^2 + 192\Delta_2(q)^2 + H_2(q)(22\psi_1(q)^{10} - 132\psi_1(q)^5\psi_2(q)^5 - 2\psi_2(q)^{10})\}}{21\eta(q)^{42/5}} \\
&= q^{-3/20}\left(1 + 40q + 381q^2 + 2865q^3 + \frac{115789}{7}q^4 + 81261q^5 + 348612q^6 + \dots\right), \\
f_{-7/10} &= \frac{\psi_1(q)\{H_2(q)^2 - 192\Delta_2(q)^2 + H_2(q)(2\psi_1(q)^{10} - 132\psi_1(q)^5\psi_2(q)^5 - 22\psi_2(q)^{10})\}}{3\eta(q)^{42/5}} \\
&= q^{-7/20}(1 - 63q - 1883q^2 - 18403q^3 - 122388q^4 - 645036q^5 - 2896215q^6 + \dots).
\end{aligned}$$

(b) $\mathbf{s} = -38/5$. A fundamental system of solutions of $(b_{-38/5})$ is given by

$$\begin{aligned}
f_{-8/15} &= \frac{\psi_1(q)G_1(I_{15}(q), \Delta_{15}(q), I_3(q), I_3(q^5))}{16\eta(q)^{32/5}} \\
&= q^{-4/15} (1 - 56q - 776q^2 - 5088q^3 - 24932q^4 - \dots), \\
f_{-1/3} &= \frac{\psi_2(q)G_1(I_{15}(q), \Delta_{15}(q), -I_3(q), -I_3(q^5))}{128\eta(q)^{32/5}} \\
&= q^{-1/15} \left(1 + 15q + 100q^2 + \frac{4629}{8}q^3 + 2635q^4 + \dots\right), \\
f_{4/5} &= \frac{\psi_1(q)G_2(I_{15}(q), \Delta_{15}(q), I_3(q), I_3(q^5), \psi_2(q)^5)}{936493073280\Delta_3(q)^2\eta(q)^{32/5}} \\
&= q^{16/15} \left(1 + \frac{28}{3}q + \frac{164}{3}q^2 + \frac{752}{3}q^3 + \frac{1955}{2}q^4 + \dots\right), \\
f_0 &= \frac{\psi_2(q)G_3(I_{15}(q), \Delta_{15}(q), I_3(q), I_3(q^5), \psi_2(q)^5)}{31216435776\Delta_3(q)^2\eta(q)^{32/5}} \\
&= q^{4/15} (1 + 8q + 56q^2 + 288q^3 + 1254q^4 + \dots),
\end{aligned}$$

where $G_1(u, x, y)$, $G_2(u, v, x, y, w)$ and $G_3(u, v, x, y, w)$ are homogeneous polynomials of degree 3, 5 and 5, respectively (they are given in Appendix D).

(c) $\mathbf{s} = -6/5$. The following functions are solutions of $(b_{-6/5})$:

$$\begin{aligned}
f_0 &= 1, \\
f_{1/5} &= \frac{1}{5} \int_0^q \psi_1(q)^4 \psi_2(q) (\psi_1(q)^5 - 3\psi_2(q)^5) \frac{dq}{q} = q^{1/5} \left(1 + \frac{1}{3}q + \frac{12}{11}q^2 + \frac{11}{16}q^3 + \frac{4}{7}q^4 + \dots\right), \\
f_{4/5} &= \frac{1}{15} \int_0^q \psi_1(q) \psi_2(q)^4 (12\psi_1(q)^5 + 4\psi_2(q)^5) \frac{dq}{q} = q^{4/5} \left(1 + \frac{28}{27}q + \frac{4}{7}q^2 + \frac{80}{57}q^3 + \frac{5}{9}q^4 + \dots\right).
\end{aligned}$$

It seems unlikely that $f_{1/5}$ and $f_{4/5}$ are modular forms (since denominators of the Fourier coefficients of them looks like unbounded). Since $(b_{-6/5})$ has the index 0 as a double root, $(b_{-6/5})$ has a solution with logarithmic terms.

However, modular functions

$$\frac{f'_{1/5}}{\eta^4} = \frac{\psi_1(q)^4 \psi_2(q) (\psi_1(q)^5 - 3\psi_2(q)^5)}{5\eta(q)^4}, \quad \frac{f'_{4/5}}{\eta^4} = \frac{\psi_1(q) \psi_2(q)^4 (12\psi_1(q)^5 + 4\psi_2(q)^5)}{60\eta(q)^4}$$

on $\Gamma(30)$ satisfy a third order MLDE

$$f''' - \frac{1}{2}E_2 f'' + \left(\frac{1}{2}E_2' - \frac{9}{100}E_4\right) f' + \frac{19}{5400}E_6 f = 0.$$

The other solution of this MLDE is given by

$$\frac{\psi_1(q)^{10} - 36\psi_1(q)^5 \psi_2(q)^5 - \psi_2(q)^{10}}{\eta(q)^4} = q^{-1/6} (1 - 26q - 126q^2 - 500q^3 - \dots).$$

(d) $\mathbf{s} = -\mathbf{3}/\mathbf{5}$. A fundamental system of solutions of $(b_{-3/5})$ is given by

$$\begin{aligned} f_0 &= \frac{\theta(q) + \theta(q^5)}{2\eta(q)^{3/5}\psi_1(q)} = q^{-1/40}(1 + q + q^2 + 2q^3 + 3q^4 + \cdots), \\ f_{4/5} &= \frac{\theta(q) - \theta(q^5)}{2\eta(q)^{3/5}\psi_2(q)} = q^{31/40}(1 + q + q^2 + 2q^3 + 2q^4 + \cdots), \\ f_{1/4} &= \frac{\Delta_4(q) + \Delta_4(q^5)}{\eta(q)^{3/5}\psi_1(q)} = q^{9/40}(1 + q + 2q^2 + 2q^3 + 3q^4 + \cdots), \\ f_{1/20} &= \frac{\Delta_4(q) - \Delta_4(q^5)}{\eta(q)^{3/5}\psi_2(q)} = q^{1/40}(1 + q^2 + q^3 + 2q^4 + 2q^5 + \cdots). \end{aligned}$$

(e) $\mathbf{s} = \mathbf{2}/\mathbf{5}$. A fundamental system of solutions of $(b_{2/5})$ is given by

$$\begin{aligned} f_0 &= \frac{I_{15}(q) - \Delta_{15}(q) + I_3(q)}{2\eta(q)^{8/5}\psi_1(q)} = q^{-1/15}(1 + 4q + 8q^2 + 20q^3 + 37q^4 + \cdots), \\ f_{4/5} &= \frac{-I_{15}(q) + \Delta_{15}(q) + I_3(q)}{6\eta(q)^{8/5}\psi_2(q)} = q^{11/15}\left(1 + \frac{4}{3}q + \frac{10}{3}q^2 + \frac{20}{3}q^3 + \frac{38}{3}q^4 + \cdots\right), \\ f_{1/3} &= \frac{G(I_{15}(q), \Delta_{15}(q), I_3(q), I_3(q^5))}{864\eta(q)^{8/5}\psi_1(q)} = q^{4/15}\left(1 + \frac{5}{2}q + 6q^2 + \frac{23}{2}q^3 + 23q^4 + \cdots\right), \\ f_{2/15} &= \frac{G(-I_{15}(q), -\Delta_{15}(q), I_3(q), I_3(q^5))}{432\eta(q)^{8/5}\psi_2(q)} = q^{1/15}(1 + 2q + 7q^2 + 12q^3 + 26q^4 + \cdots), \end{aligned}$$

where

$$G(u, v, x, y) = 60(3u^3 + v^3) + 81(u - v)(u + v)x - (5u + 9v - 12x)x^2 - 108(u^2 + v^2)y - 25(7u + 3v)y^2 + 15y^3.$$

(f) $\mathbf{s} = \mathbf{6}/\mathbf{5}$. A fundamental system of solutions of $(b_{6/5})$ is given by

$$\begin{aligned} f_0 &= \frac{\psi_1(q)(\psi_1(q)^5 + 2\psi_2(q)^5)}{\eta(q)^{12/5}} = q^{-1/10}(1 + 8q + 23q^2 + 68q^3 + \cdots), \\ f_{1/5} &= \frac{\psi_2(q)(2\psi_1(q)^5 - \psi_2(q)^5)}{2\eta(q)^{12/5}} = q^{1/10}\left(1 + \frac{9}{2}q + 16q^2 + 38q^3 + \cdots\right), \\ f_{2/5} &= \frac{\psi_1(q)^4\psi_2(q)^2}{\eta(q)^{12/5}} = q^{3/10}(1 + 4q + 12q^2 + 30q^3 + \cdots), \\ f_{4/5} &= \frac{\psi_1(q)^2\psi_2(q)^4}{\eta(q)^{12/5}} = q^{7/10}(1 + 2q + 7q^2 + 16q^3 + \cdots). \end{aligned}$$

(g) $\mathbf{s} = 12/5$. A fundamental system of solutions of $(b_{12/5})$ is given by

$$\begin{aligned}
f_0 &= \frac{1}{40\eta(q)^{18/5}\psi_1(q)} \{8\psi_2(q)^5(18\psi_1(q)^5 + \psi_2(q)^5) + 16(\psi_1(q^2)^{10} + 21\psi_1(q^2)^5\psi_2(q^2)^5 - 2\psi_2(q^2)^{10}) \\
&\quad + 5H_2(q) + 19H_2(q^5)\} \\
&= q^{-3/20}(1 + 18q + 81q^2 + 306q^3 + 909q^4 + \dots), \\
f_{4/5} &= \frac{1}{360\eta(q)^{18/5}\psi_2(q)} \{-8\psi_2(q)^5(18\psi_1(q)^5 + \psi_2(q)^5) - 16(\psi_1(q^2)^{10} + 21\psi_1(q^2)^5\psi_2(q^2)^5 \\
&\quad - 2\psi_2(q^2)^{10}) + 21H_2(q) - 5H_2(q^5)\} \\
&= q^{13/20}\left(1 + \frac{34}{9}q + 17q^2 + 50q^3 + \frac{428}{3}q^4 + \dots\right), \\
f_{1/2} &= \frac{1}{6\eta(q)^{18/5}\psi_1(q)^6} \{\Delta_2(q)(5\psi_1(q)^5 + \psi_2(q)^5 + \psi_1(q^2)^5 - \psi_2(q^2)^5) + \Delta_2(q^5)(7\psi_1(q)^5 - \psi_2(q)^5 \\
&\quad - \psi_1(q^2)^5 - 7\psi_2(q^2)^5)\} \\
&= q^{7/20}\left(1 + \frac{20}{3}q + 27q^2 + 89q^3 + \frac{766}{3}q^4 + \dots\right), \\
f_{3/10} &= \frac{1}{2\eta(q)^{18/5}\psi_2(q)^6} \{\Delta_2(q)(-\psi_1(q)^5 + 5\psi_2(q)^5 + \psi_1(q^2)^5 + \psi_2(q^2)^5) + \Delta_2(q^5)(\psi_1(q)^5 \\
&\quad + 7\psi_2(q)^5 + 7\psi_1(q^2)^5 - \psi_2(q^2)^5)\} \\
&= q^{3/20}(1 + 9q + 39q^2 + 131q^3 + 387q^4 + \dots).
\end{aligned}$$

(h) $\mathbf{s} = 18/5$. A fundamental system of solutions of $(b_{18/5})$ is given by

$$\begin{aligned}
f_0 &= \frac{\psi_1(q)^2(\psi_1(q)^{10} + 24\psi_1(q)^5\psi_2(q)^5 - 6\psi_2(q)^{10})}{\eta(q)^{24/5}} = q^{-1/5}(1 + 36q + 240q^2 + 1144q^3 + \dots), \\
f_{2/5} &= \frac{\psi_2(q)^2(6\psi_1(q)^{10} + 24\psi_1(q)^5\psi_2(q)^5 - \psi_2(q)^{10})}{6\eta(q)^{24/5}} = q^{1/5}\left(1 + 14q + \frac{461}{6}q^2 + 330q^3 + \dots\right), \\
f_{3/5} &= \frac{\psi_1(q)^4\psi_2(q)^3(4\psi_1(q)^5 + 3\psi_2(q)^5)}{4\eta(q)^{24/5}} = q^{2/5}\left(1 + \frac{39}{4}q + 51q^2 + \frac{417}{2}q^3 + \dots\right), \\
f_{4/5} &= \frac{\psi_1(q)^3\psi_2(q)^4(3\psi_1(q)^5 - 4\psi_2(q)^5)}{3\eta(q)^{24/5}} = q^{3/5}\left(1 + \frac{20}{3}q + 36q^2 + 136q^3 + \dots\right).
\end{aligned}$$

(i) $\mathbf{s} = 22/5$. A fundamental system of solutions of $(b_{22/5})$ is given by

$$\begin{aligned}
f_0 &= \frac{G_4(I_{15}(q), \Delta_{15}(q), I_3(q), I_3(q^5))}{24\psi_1(q)\eta(q)^{28/5}} = q^{-7/30}(1 + 56q + 476q^2 + 2632q^3 + 11270q^4 + \dots), \\
f_{4/5} &= \frac{G_4(-I_{15}(q), -\Delta_{15}(q), I_3(q), I_3(q^5))}{504\psi_2(q)\eta(q)^{28/5}}
\end{aligned}$$

$$\begin{aligned}
&= q^{17/30} \left(1 + \frac{28}{3}q + \frac{1196}{21}q^2 + \frac{752}{3}q^3 + \frac{2851}{3}q^4 + \cdots \right), \\
f_{2/3} &= \frac{G_5(I_{15}(q), \Delta_{15}(q), I_3(q), I_3(q^5), \psi_2(q)^5)}{26309472\psi_1(q)\Delta_3(q)\eta(q)^{28/5}} = q^{13/30} \left(1 + 12q + 73q^2 + 338q^3 + \frac{9070}{7}q^4 + \cdots \right), \\
f_{7/15} &= \frac{G_6(I_{15}(q), \Delta_{15}(q), I_3(q), I_3(q^5), \psi_2(q)^5)}{7516992\psi_2(q)\Delta_3(q)} \\
&= q^{7/30} \left(1 + \frac{35}{2}q + 112q^2 + \frac{1099}{2}q^3 + 2163q^4 + \cdots \right),
\end{aligned}$$

where $G_4(u, v, x, y)$, $G_5(u, v, x, y, w)$ and $G_6(u, v, x, y, w)$ are homogeneous polynomials of degree 3, 4 and 4, respectively (they are written in Appendix D).

(j) $\mathbf{s} = 27/5$. A fundamental system of solutions of $(b_{27/5})$ is given by

$$\begin{aligned}
f_0 &= \frac{G_7(\theta(q), \theta(q^5), \psi_1(q)^5, \psi_2(q)^5)}{10\psi_1(q)\eta(q)^{33/5}} = q^{-11/40} \left(1 + 99q + 1122q^2 + 7425q^3 + 37191q^4 + \cdots \right), \\
f_{4/5} &= \frac{G_8(\theta(q), \theta(q^5), \psi_1(q)^5, \psi_2(q)^5)}{330\psi_2(q)\eta(q)^{33/5}} = q^{21/40} \left(1 + \frac{41}{3}q + 98q^2 + 513q^3 + 2214q^4 + \cdots \right), \\
f_{3/4} &= \frac{G_9(\theta(q), \theta(q^5), \psi_1(q^4)^5, \psi_2(q^4)^5, \Delta_4(q)^3\Delta_4(q^5))}{9641984\psi_1(q)\Delta_4(q)\eta(q)^{33/5}} \\
&= q^{19/40} \left(1 + 15q + \frac{1191}{11}q^2 + 577q^3 + 2505q^4 + \cdots \right), \\
f_{11/20} &= \frac{G_{10}(\theta(q), \theta(q^5), \psi_1(q^4)^5, \psi_2(q^4)^5, \Delta_4(q)^3\Delta_4(q^5))}{7888896\psi_2(q)\Delta_4(q)\eta(q)^{33/5}} \\
&= q^{11/40} \left(1 + 22q + \frac{506}{3}q^2 + 957q^3 + 4279q^4 + \cdots \right),
\end{aligned}$$

where $G_7(u, v, x, y)$, $G_8(u, v, x, y)$, $G_9(u, v, x, y, w)$ and $G_{10}(u, v, x, y, w)$ are homogeneous polynomials of degree 7, 7, 8 and 8, respectively, which are given in Appendix D.

(k) $\mathbf{s} = 6$. The following functions are solutions of (b_6) :

$$\begin{aligned}
f_0 &= \frac{\psi_1(q)^3(\psi_1(q)^{15} + 126\psi_1(q)^{10}\psi_2(q)^5 + 117\psi_1(q)^5\psi_2(q)^{10} - 12\psi_2(q)^{15})}{\eta(q)^{36/5}} \\
&= q^{-3/10} \left(1 + 144q + 1926q^2 + 14160q^3 + 77499q^4 + \cdots \right), \\
f_{3/5} &= \frac{\psi_2(q)^3(12\psi_1(q)^{15} + 117\psi_1(q)^{10}\psi_2(q)^5 - 126\psi_1(q)^5\psi_2(q)^{10} + \psi_2(q)^{15})}{12\eta(q)^{36/5}} \\
&= q^{3/10} \left(1 + \frac{99}{4}q + 210q^2 + \frac{7739}{6}q^3 + 6195q^4 + \cdots \right), \\
f_{4/5} &= \frac{\psi_1(q)^4\psi_2(q)^4(9\psi_1(q)^{10} + 26\psi_1(q)^5\psi_2(q)^5 - 9\psi_2(q)^{10})}{9\eta(q)^{36/5}}
\end{aligned}$$

$$= q^{1/2} \left(1 + \frac{152}{9}q + 134q^2 + 772q^3 + \frac{10778}{3}q^4 + \dots \right).$$

The function $\eta(q)^{12} = q^{1/2} + \dots$ is a modular form of weight 6 on $\Gamma(2)$. Then we have

$$\eta(q)^{12} = \eta(q)^{36/5} \psi_1(q) \psi_2(q) (\psi_1(q)^{10} - 11\psi_1(q)^5 \psi_2(q)^5 - \psi_2(q)^{10})$$

since we see that the both-sides of this equality are solutions of $\vartheta_6(f) = 0$ by using functional equations given in Appendix A (d), and the leading coefficients of both q -series are one. Hence, f_r is a modular function on $\Gamma(10)$ for each $r = 0, 3/5, 4/5$.

We have the MLDE (b_6) by applying the Serre derivation $\vartheta_6(F) = F' - (E_2/2)F$ to the third order MLDE

$$f''' - \frac{1}{2}E_2 f'' + \left(\frac{1}{2}E_2' - \frac{9}{100}E_4 \right) f' + \frac{9}{200}E_6 f = 0,$$

which have the formal conformal weights 0, 3/5, 4/5 and formal central charge 6.

There is a solution of (b_6) with a logarithmic term of the form

$$(2\pi\sqrt{-1}\tau) \cdot f_{4/5} - q^{3/2} \left(\frac{2530}{81} + \frac{191600}{693}q + \frac{8906965}{4788}q^2 + \frac{5783927675}{632016}q^3 + \frac{385857740243}{9927918}q^4 + O(q^5) \right).$$

(I) $s = 32/5$. There exist solutions of ($b_{32/5}$) of the forms

$$\begin{aligned} f_0 &= \frac{\psi_1(q)^4 (\psi_1(q)^{15} + 171\psi_1(q)^{10}\psi_2(q)^5 + 247\psi_1(q)^5\psi_2(q)^{10} - 57\psi_2(q)^{15})}{\eta(q)^{38/5}} \\ &= q^{-19/60} (1 + 190q + 2831q^2 + 22306q^3 + 129276q^4 + 611724q^5 + \dots), \\ f_{4/5} &= \frac{\psi_2(q)^4 (57\psi_1(q)^{15} + 247\psi_1(q)^{10}\psi_2(q)^5 - 171\psi_1(q)^5\psi_2(q)^{10} + \psi_2(q)^{15})}{57\eta(q)^{38/5}} \\ &= q^{29/60} \left(1 + \frac{58}{3}q + \frac{493}{3}q^2 + \frac{57362}{57}q^3 + \frac{14761}{3}q^4 + 20734q^5 + \dots \right). \end{aligned}$$

The functions f_0 and $f_{4/5}$ satisfy ($\sharp_{\mu(19/5)}$) and the MLDE ($b_{32/5}$) is rewritten as

$$(B.1) \quad \vartheta_6 \circ \vartheta_4(L(f)) - \frac{11}{3600}E_4 L(f) = 0,$$

where $L(f)$ is the left-hand side of ($\sharp_{\mu(19/5)}$). Hence we have other two solutions which are given by

$$\begin{aligned} f_{5/6} &= \frac{5}{144} \left(30f_{4/5}(q) \frac{\psi_1(q)^4 \psi_2(q) (\psi_1(q)^5 - 3\psi_2(q)^5)}{\eta(q)^4} - f_0(q) \int_0^q f_{4/5}(q_0) \eta(q_0)^{18/5} \psi_2(q_0) \frac{dq_0}{q_0} \right) \\ &= q^{31/60} \left(1 + \frac{200}{11}q + \frac{28647}{187}q^2 + \frac{3989341}{4301}q^3 + \frac{562835919}{124729}q^4 + \dots \right), \\ f_{19/30} &= \frac{19}{144} \left(10f_0(q) \frac{\psi_1(q) \psi_2(q)^4 (3\psi_1(q)^5 + \psi_2(q)^5)}{19\eta(q)^4} - f_{4/5}(q) \int_0^q f_0(q_0) \eta(q_0)^{18/5} \psi_1(q_0) \frac{dq_0}{q_0} \right) \\ &= q^{19/60} \left(1 + \frac{133}{5}q + \frac{13243}{55}q^2 + \frac{1454051}{935}q^3 + \frac{168154408}{21505}q^4 + \dots \right). \end{aligned}$$

The latter two solutions are most likely not modular functions (since the denominators of coefficients of q -series look like unbounded).

(m) $s = 54/5$. A fundamental system of solutions of $(b_{54/5})$ is given by

$$\begin{aligned} f_0 &= \frac{P(\psi_1(q), \psi_2(q))}{\eta(q)^{12}} = q^{-1/2}(1 + 36q + 2490q^2 + 38360q^3 + 398715q^4 + \dots), \\ f_{4/5} &= \frac{Q(\psi_1(q), \psi_2(q))}{3\eta(q)^{22}} = q^{3/10} \left(1 + \frac{212}{3}q + 1312q^2 + 14480q^3 + \frac{350635}{3}q^4 + \dots \right), \\ f_1 &= \frac{R(\psi_1(q), \psi_2(q))}{132\eta(q)^{12}} = q^{1/2} \left(1 + \frac{95}{2}q + \frac{25360}{33}q^2 + \frac{346965}{44}q^3 + \frac{666770}{11}q^4 + \dots \right), \\ f_{6/5} &= \frac{S(\psi_1(q), \psi_2(q))}{22\eta(q)^{12}} = q^{7/10} \left(1 + \frac{372}{11}q + \frac{10779}{22}q^2 + \frac{51626}{11}q^3 + \frac{379482}{11}q^4 + \dots \right), \end{aligned}$$

where

$$\begin{aligned} P(x, y) &= x^{30} + 6x^{25}y^5 + 1875x^{20}y^{10} - 6080x^{15}y^{15} + 18135x^{10}y^{20} - 1038x^5y^{25} - 3y^{30}, \\ Q(x, y) &= x^6y^4(3x^{20} + 134x^{15}y^5 + 57x^{10}y^{10} + 216x^5y^{15} - 22y^{20}), \\ R(x, y) &= y^5(132x^{30} + 2970x^{25}y^5 - 1520x^{20}y^{10} + 7035x^{15}y^{15} - 390x^5y^{25} - y^{30}), \\ S(x, y) &= x^4y^6(22x^{20} + 216x^{15}y^5 - 57x^{10}y^{10} + 134x^5y^{15} - 3y^{20}). \end{aligned}$$

(n) $s = 18$. A fundamental system of solutions of $(b_{18/5})$ is given by

$$\begin{aligned} f_{-4/5} &= \frac{G_{11}(\psi_1(q), \psi_2(q))}{\eta(q)^{12}} = q^{-4/5}(1 - 216q - 90984q^2 - 4550240q^3 - 107053506q^4 + \dots), \\ f_0 &= 1, \\ f_{4/5} &= \frac{G_{12}(\psi_1(q), \psi_2(q))}{4959\eta(q)^{96/5}} = q^{4/5} \left(1 + \frac{248}{3}q + \frac{22360}{9}q^2 + \frac{837856}{19}q^3 + \frac{1680020}{3}q^4 + \dots \right), \\ f_1 &= \frac{G_{13}(\psi_1(q), \psi_2(q))}{4408\eta(q)^{96/5}} = q \left(1 + 63q + \frac{31596}{19}q^2 + \frac{4150739}{152}q^3 + \frac{181085301}{551}q^4 + \dots \right), \end{aligned}$$

where $G_{11}(x, y)$, $G_{12}(x, y)$ and $G_{13}(x, y)$ are homogeneous polynomials of degree 23, 23 and 48, respectively. They are given in Appendix D.

(o) $\mathbf{s} = \mathbf{66/5}$. A fundamental system of solutions of $(b_{66/5})$ is given by

$$\begin{aligned} f_{-1/5} &= -\frac{P(\psi_1(q), \psi_2(q))}{4\eta(q)^{12}} = q^{-1/2} \left(1 - \frac{315}{4}q - 11570q^2 - \frac{456545}{2}q^3 - 2506845q^4 - \dots \right), \\ f_0 &= \frac{Q(\psi_1(q), \psi_2(q))}{\eta(q)^{12}} = q^{-3/10} (1 + 232q + 4902q^2 + 57276q^3 + 490507q^4 + \dots), \\ f_{4/5} &= \frac{R(\psi_1(q), \psi_2(q))}{1653\eta(q)^{12}} = q^{1/2} \left(1 + \frac{80}{3}q + \frac{1010}{3}q^2 + \frac{57840}{19}q^3 + \frac{414330}{19}q^4 + \dots \right), \\ f_{8/5} &= \frac{S(\psi_1(q), \psi_2(q))}{551\eta(q)^{12}} = q^{13/10} \left(1 + 24q + \frac{5458}{19}q^2 + \frac{45800}{19}q^3 + \frac{8847495}{551}q^4 + \dots \right), \end{aligned}$$

where

$$\begin{aligned} P(x, y) &= x^5 (4x^{25} - 435x^{20}y^5 - 37265x^{15}y^{10} - 44080x^{10}y^{15} - 2755x^5y^{20} + 1653y^{25}), \\ Q(x, y) &= x^9y (x^{20} + 203x^{15}y^5 - 406x^{10}y^{10} - 1653x^5y^{15} + 551y^{20}), \\ R(x, y) &= y^5 (1653x^{25} + 2755x^{20}y^5 - 44080x^{15}y^{10} + 37265x^{10}y^{15} - 435x^5y^{20} - 4y^{25}), \\ S(x, y) &= xy^9 (551x^{20} + 1653x^{15}y^5 - 406x^{10}y^{10} - 203x^5y^{15} + y^{20}). \end{aligned}$$

(p) $\mathbf{s} = -\mathbf{6}$. A fundamental system of solutions of (b_{-6}) is given by

$$\begin{aligned} f_{-1/5} &= \frac{\psi_1(q)^2 (\psi_1(q)^{10} - 66\psi_1(q)^5\psi_2(q)^5 - 11\psi_2(q)^{10})}{\eta(q)^{24/5}} \\ &= q^{-1/5} (1 - 54q - 395q^2 - 1836q^3 - 6950q^4 - \dots), \\ f_0 &= \frac{\psi_1(q)^6\psi_2(q) (2\psi_1(q)^5 + 11\psi_2(q)^5)}{2\eta(q)^{24/5}} = 1 + \frac{33}{2}q + 100q^2 + \frac{893}{2}q^3 + 1629q^4 + \dots, \\ f_{1/5} &= \frac{\psi_2(q)^2 (11\psi_1(q)^{10} - 66\psi_1(q)^5\psi_2(q)^5 - \psi_2(q)^{10})}{11\eta(q)^{24/5}} \\ &= q^{1/5} \left(1 + 4q + \frac{296}{11}q^2 + 110q^3 + \frac{4344}{11}q^4 + \dots \right), \\ f_1 &= \frac{\psi_1(q)\psi_2(q)^6 (11\psi_1(q)^5 - 2\psi_2(q)^5)}{11\eta(q)^{24/5}} = q \left(1 + \frac{68}{11}q + \frac{299}{11}q^2 + \frac{1102}{11}q^3 + \frac{3511}{11}q^4 + \dots \right), \end{aligned}$$

(q) $\mathbf{s} = -\mathbf{8/5}$. The following two functions satisfy $(b_{-8/5})$:

$$\begin{aligned} f_0 &= \frac{\psi_2(q)}{\eta(q)^{2/5}} = q^{11/60} (1 + q^2 + q^3 + q^4 + q^5 + \dots), \\ f_{-1/5} &= \frac{\psi_1(q)}{\eta(q)^{2/5}} = q^{-1/60} (1 + q + q^2 + q^3 + 2q^4 + 2q^5 + \dots). \end{aligned}$$

We see that f_0 and $f_{-1/5}$ also satisfy $(\sharp_{\mu(1/5)})$, and $(b_{-8/5})$ is rewritten as $\vartheta_6 \circ \vartheta_4(L_{1/5}(f)) - (551/3600)E_4L_{1/5}(f) = 0$, where $L_{1/5}(f)$ denotes the left-hand side of $(\sharp_{\mu(1/5)})$. Hence two other

solutions are given by

$$\begin{aligned}
& f_{-1/6} \\
&= \frac{1}{36} \left\{ 30 \frac{\psi_1(q)^7 \psi_2(q)^3 (\psi(q)_1^5 - 3\psi_2(q)^5) (\psi_1(q)^{10} - 11\psi_1(q)^5 \psi_2(q)^5 - \psi_2(q)^{10})^2}{\eta(q)^{14}} \right. \\
&\quad \left. - f_0(q) \int_0^q \frac{\psi_1(q_0)^5 (\psi_1(q_0)^{15} + 171\psi_1(q_0)^{10} \psi_2(q_0)^5 + 247\psi_1(q_0)^5 \psi_2(q_0)^{10} - 57\psi_2(q_0)^{15}) dq_0}{\eta(q_0)^4} \frac{dq_0}{q_0} \right\} \\
&= q^{1/60} \left(1 - \frac{2}{5}q + \frac{1}{11}q^2 + \frac{26}{85}q^3 + \frac{434}{1265}q^4 + \frac{9824}{27115}q^5 + \cdots \right),
\end{aligned}$$

$$\begin{aligned}
& f_{19/30} \\
&= \frac{5}{36} \left\{ 10 \frac{\psi_1(q)^3 \psi_2(q)^7 (3\psi_1(q)^5 + \psi_2(q)^5) (\psi_1(q)^{10} - 11\psi_1(q)^5 \psi_2(q)^5 - \psi_2(q)^{10})^2}{\eta(q)^{14}} \right. \\
&\quad \left. - f_{-1/5}(q) \int_0^q \frac{\psi_1(q_0)^5 (57\psi_1(q_0)^{15} + 247\psi_1(q_0)^{10} \psi_2(q_0)^5 - 171\psi_1(q_0)^5 \psi_2(q_0)^{10} + \psi_2(q_0)^{15}) dq_0}{3\eta(q_0)^4} \frac{dq_0}{q_0} \right\} \\
&= q^{49/60} \left(1 + \frac{38}{33}q + \frac{371}{561}q^2 + \frac{22558}{12903}q^3 + \frac{383219}{374187}q^4 + \frac{938830}{374187}q^5 + \cdots \right).
\end{aligned}$$

It is most likely that the last two solutions are not modular functions (since the denominators of coefficients of q -series look like unbounded).

APPENDIX C. BASES OF MODULAR LINEAR DIFFERENTIAL EQUATIONS WITH SOLUTIONS OF QUASIMODULAR FORMS OF POSITIVE DEPTHS

In this appendix we construct a fundamental system of solutions of (b_s) when (b_s) has a quasimodular solution of positive depth of CFT type. Among the candidates for s given in (3.12), the six numbers $-318/5$, $-198/5$, $-138/5$, $-78/5$, $-18/5$ and $42/5$ give such cases.

C.1. Quasimodular forms and solutions with logarithmic terms. If a MLDE $L(f) = 0$ of weight k has solutions of quasimodular forms of weight $k + r$ and depth $r (> 0)$, then there are solutions of $L(f) = 0$ with logarithmic terms (see [KNS]).

For a congruence subgroup $\Gamma \subset \mathrm{SL}_2(\mathbb{Z})$, we now suppose that a quasimodular form $F_k := A_k E_2 + B_k$ of weight $k + 1$ and depth 1 on Γ is a solution of a MLDE (of weight k on $\mathrm{SL}_2(\mathbb{Z})$), where A_k and B_k are modular forms on Γ of weight $k - 1$ and $k + 1$, respectively. Then the function $G_k = G(F_k) = (2\pi\sqrt{-1})^{-1}(F_k \log q + 12A_k)$ is a solution because

$$F_k \Big|_k \begin{pmatrix} a & b \\ c & d \end{pmatrix} = c G_k + d F_k \quad \text{for } \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \Gamma$$

is a solution by modular invariance property. We find that

$$\begin{pmatrix} F_k \\ G_k \end{pmatrix} \Big|_k \begin{pmatrix} a & b \\ c & d \end{pmatrix} = \begin{pmatrix} d & c \\ b & a \end{pmatrix} \begin{pmatrix} F_k \\ G_k \end{pmatrix} \quad \text{for each } \begin{pmatrix} a & b \\ c & d \end{pmatrix} \in \Gamma.$$

C.2. Solutions of modular linear differential equations of quasimodular forms of positive depths. In this subappendix we list solutions of (b_s) of quasimodular forms, where s is listed in the beginning of Appendix C. By using construction in Subappendix C.1, we have a fundamental system of the space of solutions of (b_s) .

(a) $s = -318/5$. Let ψ_1 and ψ_2 be the modular functions defined in Appendix A. A set of two linearly independent solutions of $(b_{-318/5})$ is given by

$$\begin{aligned} f_0 &= \frac{F_1(\psi_1(q), \psi_2(q))'}{50841895104\eta(q)^{192/5}} + \frac{F_2(\psi_1(q), \psi_2(q))}{419325701671800\eta(q)^{312/5}} \\ &= q^{13/5}(1 + 260q + 30056q^2 + 2119676q^3 + 104823121q^4 + \dots), \\ f_{4/5} &= \frac{F_1(\psi_2(q), -\psi_1(q))'}{28346417433013680\eta(q)^{312/5}} + \frac{F_2(\psi_2(q), -\psi_1(q))}{5451234121733400\eta(q)^{312/5}} \\ &= q^{17/5}(1 + 236q + 25306q^2 + 1680916q^3 + 79143742q^4 + \dots), \end{aligned}$$

where $F_1(x, y)$ and $F_2(x, y)$ are homogeneous polynomials of degree 151 and 161, respectively. (They are given in Appendix D). The solutions f_0 and $f_{4/5}$ are quasimodular forms of weight 1 and depth 1. The system $(f_0, f_{4/5}, G(f_0), G(f_{4/5}))$ is a fundamental system of solutions of $(b_{-318/5})$. Here $G(*)$ is a function with a logarithmic term defined in Subappendix C.1.

(b) $s = -198/5$. A set of two linearly independent solutions of $(b_{-198/5})$ is given by

$$\begin{aligned} f_0 &= \frac{F_3(\psi_1(q), \psi_2(q))'}{50841895104\eta(q)^{192/5}} + \frac{F_4(\psi_1(q), \psi_2(q))}{15888092220\eta(q)^{192/5}} \\ &= q^{8/5}(1 + 144q + 8880q^2 + 331840q^3 + 8770284q^4 + \dots), \\ f_{4/5} &= \frac{F_3(\psi_2(q), -\psi_1(q))'}{4236824592\eta(q)^{192/5}} + \frac{F_4(\psi_2(q), -\psi_1(q))}{1324007685\eta(q)^{192/5}} \\ &= q^{12/5}\left(1 + \frac{380}{3}q + 7164q^2 + 251344q^3 + \frac{18958205}{3}q^4 + \dots\right), \end{aligned}$$

where $F_3(x, y)$ and $F_4(x, y)$ are homogeneous polynomials of degree 91 and 101, respectively (they are given in Appendix D). The solutions f_0 and $f_{4/5}$ are quasimodular forms of weight 1 and depth 1. The system $(f_0, f_{4/5}, G(f_0), G(f_{4/5}))$ is a fundamental system of solutions of $(b_{-198/5})$.

(c) $s = -138/5$. A set of two linearly independent solutions of $(b_{-138/5})$ is given by

$$\begin{aligned} f_0 &= \frac{P(\psi_1(q), \psi_2(q))'}{48360312\eta(q)^{132/5}} + \frac{Q(\psi_1(q), \psi_2(q))}{21981960\eta(q)^{132/5}} \\ &= q^{11/10}(1 + 88q + 3256q^2 + 74360q^3 + 1232814q^4 + \dots), \\ f_{4/5} &= \frac{P(\psi_2(q), -\psi_1(q))'}{4396392\eta(q)^{132/5}} + \frac{Q(\psi_2(q), -\psi_1(q))}{1998360\eta(q)^{132/5}} \\ &= q^{19/10}(1 + 76q + 2584q^2 + 55568q^3 + 876329q^4 + \dots), \end{aligned}$$

where

$$\begin{aligned}
P(x, y) &= x(2x^{60} - 423x^{55}y^5 + 62386x^{50}y^{10} + 17130550x^{45}y^{15} + 257495595x^{40}y^{20} \\
&\quad + 449723758x^{35}y^{25} + 110860458x^{30}y^{30} + 274470804x^{25}y^{35} - 44451050x^{20}y^{40} \\
&\quad - 31757435x^{15}y^{45} + 1691208x^{10}y^{50} - 3542x^5y^{55} - 11y^{60}), \\
Q(x, y) &= xy^5(159x^{65} - 51748x^{60}y^5 - 1801195x^{55}y^{10} - 11218651x^{50}y^{15} + 431478190x^{45}y^{20} \\
&\quad - 348592259x^{40}y^{25} + 1688812058x^{35}y^{30} + 402533490x^{30}y^{35} + 852888611x^{25}y^{40} \\
&\quad - 3573970x^{20}y^{45} - 4304879x^{15}y^{50} + 269233x^{10}y^{55} + 1760x^5y^{60} + y^{65}).
\end{aligned}$$

The solutions f_0 and $f_{4/5}$ are quasimodular forms of weight 1 and depth 1. Moreover, the system $(f_0, f_{4/5}, G(f_0), G(f_{4/5}))$ is a fundamental system of solutions of $(b_{-138/5})$.

(d) $\mathbf{s} = -78/5$. A set of linearly independent solutions of $(b_{-78/5})$ is given by

$$\begin{aligned}
f_0 &= \frac{P(\psi_1(q), \psi_2(q))'}{2604\eta(q)^{72/5}} - \frac{Q(\psi_1(q), \psi_2(q))}{2170\eta(q)^{72/5}} = q^{3/5}(1 + 36q + 576q^2 + 6312q^3 + 53739q^4 + \dots), \\
f_{4/5} &= -\frac{P(\psi_2(q), -\psi_1(q))}{23436\eta(q)^{72/5}} + \frac{Q(\psi_2(q), -\psi_1(q))}{19530\eta(q)^{72/5}} \\
&= q^{7/5}\left(1 + \frac{284}{9}q + 476q^2 + 4888q^3 + \frac{117116}{3}q^4 + \dots\right),
\end{aligned}$$

where

$$\begin{aligned}
P(x, y) &= y(6x^{30} + 1642x^{25}y^5 - 1440x^{20}y^{10} - 20615x^{15}y^{15} + 12665x^{10}y^{20} - 123x^5y^{25} - y^{30}), \\
Q(x, y) &= x^5y(x^{10} - 11x^5y^5 - y^{10})(x^{25} - 435x^{20}y^5 - 6670x^{15}y^{10} - 3335x^5y^{20} + 87y^{25}).
\end{aligned}$$

The solutions f_0 and $f_{4/5}$ are quasimodular forms of weight 1 and depth 1. Moreover, the system $(f_0, f_{4/5}, G(f_0), G(f_{4/5}))$ is a fundamental system of solutions of $(b_{-78/5})$.

(e) $\mathbf{s} = -18/5$. A set of linearly independent solutions of $(b_{-18/5})$ is given by

$$\begin{aligned}
f_0 &= \frac{5\psi_2(q)'}{\eta(q)^{12/5}} = q^{1/10}(1 + 6q^2 + 16q^3 + 36q^4 + 72q^5 + \dots), \\
f_{4/5} &= \frac{5\psi_1(q)'}{3\eta(q)^{12/5}} = q^{9/10}\left(1 + \frac{8}{3}q + 6q^2 + 16q^3 + \frac{101}{3}q^4 + 72q^5 + \dots\right).
\end{aligned}$$

The solutions f_0 and $f_{4/5}$ are quasimodular forms of weight 1 and depth 1. Moreover, the system $(f_0, f_{4/5}, G(f_0), G(f_{4/5}))$ is a fundamental system of solutions of $(b_{-18/5})$.

(f) $s = 42/5$. A set of linearly independent solutions of $(b_{42/5})$ is given by

$$\begin{aligned} f_0 &= \frac{5 \{ \psi_2(q)^4 (57\psi_1(q)^{15} + 247\psi_1(q)^{10}\psi_2(q)^5 - 171\psi_1(q)^5\psi_2(q)^{10} + \psi_2(q)^{15}) \}' }{28\eta(q)^{48/5}} \\ &= q^{2/5} (1 + 36q + 436q^2 + 3536q^3 + 21912q^4 + 113760q^5 + \dots), \\ f_{1/5} &= \frac{5 \{ \psi_1(q)^4 (\psi_1(q)^{15} + 171\psi_1(q)^{10}\psi_2(q)^5 + 247\psi_1(q)^5\psi_2(q)^{10} - 57\psi_2(q)^{15}) \}' }{912\eta(q)^{48/5}} \\ &= q^{3/5} \left(1 + 25q + 276q^2 + \frac{8379}{4}q^3 + 12481q^4 + 62859q^5 + \dots \right). \end{aligned}$$

The solutions f_0 and $f_{1/5}$ are quasimodular forms of weight 1 and depth 1. Moreover, the system $(f_0, f_{4/5}, G(f_0), G(f_{4/5}))$ is a fundamental system of solutions of $(b_{-42/5})$.

Remark C.1. The numbers $s = -318/5, -198/5, -138/5, -78/5, -48/5$ and $-38/5$ in (3.12) has the form $s = -2(5h^\vee + 9)/5$ with the dual Coxeter numbers h^\vee of the Lie algebras E_8, E_7, E_6, D_4, A_2 and A_1 (the simply-laced Lie algebras in the Deligne exceptional series). When we substitute $h^\vee = 3/2, 4, 9$ and 24 (the dual Coxeter numbers of the non simply-laced Lie algebras G_2 and F_4 in the Deligne exceptional series and formal dual Coxeter numbers of $A_{1/2}$ and $E_{7+1/2}$ [LM, K1]) in the above expression of s , we have $s = -33/5, -58/5, -108/5$ and $-258/5$. Let s be one of the numbers $-33/5, -58/5, -108/5$ and $-258/5$. We believe that the modular linear differential equation (b_s) has a solution of the form $q^{\alpha_1} (5 + \sum_{n=1}^{\infty} a_n q^n)$ with $a_n \in \mathbb{Z}_{\geq 0}$. We checked it for small values of n : we see that $a_n \in \mathbb{Z}_{\geq 0}$ for $1 \leq n < 100$.

The MLDEs $(b_{282/5})$ and $(b_{132/5})$ can also have solutions of the form $q^{\alpha_1} (5 + \sum_{n=1}^{\infty} a_n q^n)$ with $a_n \in \mathbb{Z}_{\geq 0}$.

APPENDIX D. THE POLYNOMIALS WHICH EXPRESSES THE SOLUTIONS OF MODULAR LINEAR DIFFERENTIAL EQUATIONS

In this appendix we define the polynomials appearing in the solutions of (b_s) in Appendices B–C.

$$\begin{aligned} G_1(u, v, x, y) &= 180(7u^3 + v^3) - 9(11u^2 - 31v^2)x \\ &\quad - x^2(213u + 39v + 34x) + 162(u^2 + v^2)y - 5y^2(195u - 15v + 17y), \end{aligned}$$

$$\begin{aligned} G_2(u, v, x, y, w) &= -720(223958344u^5 + 406176133v^5) \\ &\quad + (3184319633u^4 - 197068307233v^4 + 33362124000w^4)x \\ &\quad + 60(1353945577u^3 + 878294370v^3)x^2 + 6(6929323501v^2 - 1198443520w^2)x^3 \\ &\quad - x^4(1858938780v + 2043544673x) + 15(2134148833u^4 - 1674476673v^4)y \\ &\quad - 1500(154243777u^3 + 15399378v^3)y^2 - 10(11725793962u^2 + 3577812017v^2)y^3 \\ &\quad + 7500(43261956u + 12682717v)y^4 + 71019200725y^5, \end{aligned}$$

$$\begin{aligned} G_3(u, v, x, y, w) &= 144(699752744u^5 - 2362467v^5) \\ &\quad + (361371889u^4 - 1097126369v^4 + 1211988000w^4)x \end{aligned}$$

$$\begin{aligned}
& + 12(498413977u^3 + 838363630v^3)x^2 + 6(574859825v^2 + 326289904w^2)x^3 \\
& + (1446719244v - 436791181x)x^4 + 15(158488301u^4 + 365080659v^4)y \\
& - 300(475629839u^3 + 83270830v^3)y^2 - 10(940591436u^2 + 521391871v^2)y^3 \\
& + 1500(32993592u + 10616473v)y^4 + 5519145425y^5,
\end{aligned}$$

$$\begin{aligned}
G_4(u, v, x, y) &= 180u^3 + 27u^2(3x - 4y) + 7u(x^2 - 25y^2) \\
&+ 3(20v^3 + 8x^3 + 5y^3 - 9v^2(3x + 4y) - v(7x^2 + 25y^2)),
\end{aligned}$$

$$\begin{aligned}
G_5(u, v, x, y, w) &= 927276u^4 + 27002724v^4 - 1508220u^2y^2 + 9998940v^2y^2 \\
&- 8u^3(719719x + 1768043y) - 8v^3(1664689x + 9604649y) \\
&+ 4u(1080301x^3 + 3999985y^3) + 4v(390017x^3 + 7292255y^3) \\
&- 49(120000w^4 - 137104w^2x^2 + 7621x^4 - 10925y^4),
\end{aligned}$$

$$\begin{aligned}
G_6(u, v, x, y, w) &= 927276u^4 + 27002724v^4 - 1508220u^2y^2 + 9998940v^2y^2 \\
&- 8u^3(257897x + 1184773y) - 8v^3(733055x + 6576439y) \\
&+ 4u(548167x^3 + 2441935y^3) + 4v(222227x^3 + 2618105y^3) \\
&- 49(120000w^4 - 137104w^2x^2 + 7621x^4 - 10925y^4),
\end{aligned}$$

$$\begin{aligned}
G_7(u, v, x, y) &= 12u^5v^2 + 60u^4v^3 + 3u^2v(4v^4 + 61x^2 - 150xy - 39y^2) \\
&- 60uv^2(5v^4 + 5x^2 + 5xy + 3y^2) - 12v^3(5v^4 - 4x^2 + 44xy + 4y^2) \\
&+ u^3(300v^4 + 55x^2 - 18xy + 15y^2),
\end{aligned}$$

$$\begin{aligned}
G_8(u, v, x, y) &= 12u^5v^2 - 60u^4v^3 + 12v^3(5v^4 + 4x^2 - 44xy - 4y^2) \\
&- 60uv^2(5v^4 + 3x^2 - 5xy + 5y^2) + u^3(300v^4 + 15x^2 + 18xy + 55y^2) \\
&- 3u^2v(4v^4 - 39x^2 + 150xy + 61y^2),
\end{aligned}$$

$$\begin{aligned}
G_9(u, v, x, y, w) &= 738713u^8 + 105264u^7v - 109300u^6v^2 - 339760u^5v^3 \\
&+ 2u^4(764115v^4 + 1803136w - 708480x^2) + 80u^3(33729v^5 - 349760vw) \\
&- 20u^2(136625v^6 - 763776v^2w) - 2546000uw^7 + 3415625v^8 \\
&+ 6400v^4(695w + 1216x^2 - 109y^2) \\
&+ 64\{2023440w^2 - 8w(8049x^2 + 2936xy + 32151y^2) \\
&- 142563x^4 - 33071418x^2y^2 - 228288xy^3 - 34771y^4\},
\end{aligned}$$

$$\begin{aligned}
G_{10}(u, v, x, y, w) &= 1564813u^8 - 315792u^7v - 127492u^6v^2 + 1019280u^5v^3 \\
&- 6u^4(194875v^4 + 1803136w - 708480x^2) - 16u^3(505935v^5 + 2039872vw) \\
&- 20u^2(159365v^6 + 2291328v^2w) + 7638000uw^7 + 3984125v^8 \\
&- 19200v^4(695w + 1216x^2 - 109y^2)
\end{aligned}$$

$$\begin{aligned}
& + 64\{(1558352w^2 + 40w(29311x^2 + 301704xy - 5191y^2) \\
& \quad + 277889x^4 - 65865346x^2y^2 + 684864xy^3 - 45487y^4)\}, \\
G_{11}(x, y) &= x^8(x^{15} + 171x^{10}y^5 + 247x^5y^{10} - 57y^{15})(x^{25} - 435x^{20}y^5 - 6670x^{15}y^{10} \\
& \quad - 3335x^5y^{20} + 87y^{25}), \\
G_{12}(x, y) &= y^8(57x^{15} + 247x^{10}y^5 - 171x^5y^{10} + y^{15})(87x^{25} + 3335x^{20}y^5 + 6670x^{10}y^{15} \\
& \quad - 435x^5y^{20} - y^{25}) \\
G_{13}(x, y) &= x^4y^9(4408x^{35} + 105792x^{30}y^5 + 255432x^{25}y^{10} - 39585x^{20}y^{15} + 224280x^{15}y^{20} \\
& \quad - 110124x^{10}y^{25} + 4144x^5y^{30} - 6y^{35}).
\end{aligned}$$

$$\begin{aligned}
F_1(x, y) &= y(26x^{150} - 14027x^{145}y^5 + 4302519x^{140}y^{10} - 1108131323x^{135}y^{15} + 352658709221x^{130}y^{20} \\
& \quad + 410367539820195x^{125}y^{25} + 40445085344188305x^{120}y^{30} + 1299280606676985415x^{115}y^{35} \\
& \quad + 15619328979178395195x^{110}y^{40} + 55612071446837334235x^{105}y^{45} \\
& \quad - 87018500083008224035x^{100}y^{50} - 533269157005860756105x^{95}y^{55} \\
& \quad - 788818605648335112065x^{90}y^{60} - 1013858166164551300020x^{85}y^{65} \\
& \quad - 296448112259269579710x^{80}y^{70} - 1142271481069396890795x^{75}y^{75} \\
& \quad + 1168196293014645641265x^{70}y^{80} - 2113215283984202178330x^{65}y^{85} \\
& \quad + 2233117451140029299360x^{60}y^{90} - 1786660513705581667220x^{55}y^{95} \\
& \quad + 791868758851817951430x^{50}y^{100} - 151112565855362432510x^{45}y^{105} \\
& \quad + 12466416846335713220x^{40}y^{110} - 431194652630175240x^{35}y^{115} \\
& \quad + 5161178221901730x^{30}y^{120} - 5636816703218x^{25}y^{125} \\
& \quad - 22864978164x^{20}y^{130} - 122736542x^{15}y^{135} - 623536x^{10}y^{140} - 2403x^5y^{145} - 5y^{150}),
\end{aligned}$$

$$\begin{aligned}
F_2(x, y) &= x^5y(-x^{155} + 2795x^{150}y^5 - 1610905x^{145}y^{10} + 626661790x^{140}y^{15} - 272934105110x^{135}y^{20} \\
& \quad + 5921185002218x^{130}y^{25} + 2403876207873450x^{125}y^{30} + 76784270584164950x^{120}y^{35} \\
& \quad + 772577150281220900x^{115}y^{40} - 2601547321924922600x^{110}y^{45} \\
& \quad - 132050328432197323010x^{105}y^{50} - 761932811447226146950x^{100}y^{55} \\
& \quad - 646462771950208704200x^{95}y^{60} - 2253288292561296417525x^{90}y^{65} \\
& \quad - 1120714651267256252025x^{85}y^{70} - 4859622107798545477335x^{80}y^{75} \\
& \quad + 1377006603588253977975x^{75}y^{80} - 5767727038509251934525x^{70}y^{85} \\
& \quad + 2918064034868992877075x^{65}y^{90} - 2471409054146940922300x^{60}y^{95} \\
& \quad + 203085067837087672160x^{55}y^{100} - 317362540657481937650x^{50}y^{105} \\
& \quad + 61492360597697519350x^{45}y^{110} + 350110675044144200x^{40}y^{115} \\
& \quad - 419469534314529300x^{35}y^{120} + 21455464506696458x^{30}y^{125}
\end{aligned}$$

$$\begin{aligned}
& -543093587090560x^{25}y^{130} + 5212276736290x^{20}y^{135} + 16214520280x^{15}y^{140} \\
& + 62528105x^{10}y^{145} + 202921x^5y^{150} + 375y^{155}), \\
F_3(x, y) = & -x(3x^{90} - 903x^{85}y^5 + 159961x^{80}y^{10} - 28650647x^{75}y^{15} - 13575756015x^{70}y^{20} \\
& - 474499473257x^{65}y^{25} - 4010788885483x^{60}y^{30} - 7747481577039x^{55}y^{35} - 5318572158667x^{50}y^{40} \\
& - 5669746887385x^{45}y^{45} + 1236811165611x^{40}y^{50} - 3418602513421x^{35}y^{55} + 633550350937x^{30}y^{60} \\
& + 416362598871x^{25}y^{65} - 52095354335x^{20}y^{70} \\
& + 1196042149x^{15}y^{75} - 1716384x^{10}y^{80} - 6032x^5y^{85} - 16y^{90}), \\
F_4(x, y) = & -xy^5(231x^{95} - 87604x^{90}y^5 + 25209980x^{85}y^{10} + 1278608345x^{80}y^{15} + 28385930062x^{75}y^{20} \\
& - 156161181524x^{70}y^{25} - 3918089501554x^{65}y^{30} + 2542402356410x^{60}y^{35} - 21700694333270x^{55}y^{40} \\
& - 5108101637858x^{50}y^{45} - 21968910753568x^{45}y^{50} - 1511305888168x^{40}y^{55} \\
& - 8123129966420x^{35}y^{60} + 467241056080x^{30}y^{65} + 38837423474x^{25}y^{70} \\
& - 5177696692x^{20}y^{75} + 111522673x^{15}y^{80} + 1101610x^{10}y^{85} + 2000x^5y^{90} + y^{95}).
\end{aligned}$$

Acknowledgment. The authors would like to express their thanks to Prof. K. Nagatomo for helpful advice and continuous encouragement. K.K. would like to thank Prof. C.H. Lam, Y. Arike and N. Genra for useful discussions. K.K.'s research is partly supported by the Australian Research Council Discovery Project DP160101520. Y.S. was supported in part by Japan Society for the Promotion of Science, Grant-in-Aid for Scientific Research (S) 16H06336.

REFERENCES

- [AM] G. Anderson, G. Moore, Rationality in conformal field theory., *Comm. Math. Phys.* 117 (3) (1988) 441–450.
- [A] T. Arakawa, Representation theory of superconformal algebras and the Kac-Roan-Wakimoto conjecture', *Duke Math. J.* 130 (3) (2005) 435–478.
- [AK] T. Arakawa and K. Kawasetsu, “Quasi-lisse vertex algebras and modular linear differential equations.” (to appear in Kostant memorial volume), arXiv:1610.05865.
- [AKNS] Y. Arike, M. Kaneko, K. Nagatomo, Y. Sakai, Affine Vertex Operator Algebras and Modular Linear Differential Equations. *Lett. Math. Phys.* 106 (5) (2016) 693–718.
- [C] S. Carnahan, Building vertex algebras from parts, arXiv preprint arXiv:1408.5215 (2014).
- [CM] S. Carnahan, M. Miyamoto, Regularity of fixed-point vertex operator subalgebras, arXiv preprint arXiv:1603.05645 (2016).
- [CdM] A.M. Cohen, R. de Man, Computational evidence for Deligne’s conjecture regarding exceptional Lie groups, *Comp. Rend. Acad. Sci. Ser. I Math.* 322 (1996) 427–432.
- [D] P. Deligne, La série exceptionnelle de groupes de Lie, *Comp. Rend. Acad. Sci. Ser. I Math.* 322 (1996) 321–326.
- [DLM] C. Dong, H. Li, G. Mason, Modular-Invariance of Trace Functions in Orbifold Theory and Generalized Moonshine, *Comm. Math. Phys.* 214 (1) (2000) 1–56.
- [DM1] C. Dong, G. Mason, Rational vertex operator algebras and the effective central charge, *Int. Math. Res. Not.* 2004.56 (2004) 2989–3008.
- [DM2] C. Dong, G. Mason, Integrability of C_2 -cofinite vertex operator algebras, *Int. Math. Res. Not.* 2006 (2006) 80468.

- [KRW] V.G. Kac, S. Roan, M. Wakimoto, Quantum reduction for affine superalgebras, *Comm. Math. Phys.* 241 (2-3) (2003) 307–342.
- [KW2] V.G. Kac, M. Wakimoto, On rationality of W -algebras, *Transf. Groups* 13 (3-4) (2008) 671–713.
- [KNS] M. Kaneko, K. Nagatomo, Y. Sakai, Modular forms and second order ordinary differential equations: applications to vertex operator algebras, *Lett. Math. Phys.* 103 (4) (2013) 439–453.
- [KZ1] M. Kaneko, D. Zagier, A generalized Jacobi theta function and quasimodular forms, *The moduli space of curves* (Texel Island, 1994), *Progr. Math.*, vol. 129, Birkhäuser Boston, Boston, MA (1995) 165–172.
- [KZ2] M. Kaneko, D. Zagier, Supersingular j -invariants, hypergeometric series, and Atkin’s orthogonal polynomials, *AMS IP STUD. MATH.* 7 (1998) 97–126.
- [K1] K. Kawasetsu, The intermediate vertex subalgebras of the lattice vertex operator algebras, *Lett. Math. Phys.* 104 (2) (2014) 157–178.
- [K2] K. Kawasetsu, W -algebras with non-admissible levels and the Deligne exceptional series, *Int. Math. Res. Not.* (2016) rnw240.
- [LM] J.M. Landsberg, L. Manivel, The sextonions and $E_{7\frac{1}{2}}$, *Adv. Math.* 201 (1) (2006) 143–179.
- [Ma] G. Mason, Vector-valued modular forms and linear differential operators, *Int. J. Num. Th.* 3 (03) (2007) 377–390.
- [MMS1] S. Mathur, S. Mukhi, A. Sen, On the classification of rational conformal field theories, *Phys. Lett. B* 213 (3) (1988) 303–308.
- [MMS2] S. Mathur, S. Mukhi, A. Sen, Reconstruction of conformal field theories from modular geometry on the torus, *Nucl. Phys. B* 318 (2) (1989) 483–540.
- [Mil] A. Milas, On certain automorphic forms associated to rational vertex operator algebras, (English summary) *Moonshine: the first quarter century and beyond* (Edinburgh 2004), Vol. 372, *London Math. Soc. Lecture Note Ser.*, Cambridge Univ. Press, Cambridge, 2010, pp. 330–357.
- [NS] K. Nagatomo Y. Sakai, Vertex operator algebras with central charge $1/2$ and $-68/7$, *Proc. Japan Acad. Ser. A, Math. Sci.* 92 (2) (2016) 33–37.
- [T] M.P. Tuite, Exceptional vertex operator algebras and the Virasoro algebra, *Contemp. Math.* 497 (2009) 213–225.
- [Va] J. van Ekeren, Modular invariance for twisted modules over a vertex operator superalgebra, *Comm. Math. Phys.* 322 (2) (2013) 333–371.
- [W] B.W. Westbury, Sextonions and the magic square, *J. London Math. Soc.* 73 (02) (2006) 455–474.
- [Zhu] Y. Zhu, Modular invariance of characters of vertex operator algebras, *J. Amer. Math. Soc.* 9 (1) (1996) 237–302.

SCHOOL OF MATHEMATICS AND STATISTICS, THE UNIVERSITY OF MELBOURNE, 3010, AUSTRALIA AND MULTIPLE ZETA RESEARCH CENTER, KYUSHU UNIVERSITY, 744, MOTOOKA, NISHI-KU, FUKUOKA 819-0395, JAPAN.

E-mail address: kazuya.kawasetsu@unimelb.edu.au and dynamixaxs@gmail.com