

EFFECTIVE MULTI-SCALE APPROACH TO THE SCHRÖDINGER COCYCLE OVER A SKEW SHIFT BASE

R. HAN, M. LEMM, W. SCHLAG

ABSTRACT. We prove a conditional theorem on the positivity of the Lyapunov exponent for a Schrödinger cocycle over a skew shift base with a cosine potential and the golden ratio as frequency. For coupling below 1, which is the threshold for Herman's subharmonicity trick, we formulate three conditions on the Lyapunov exponent in a finite but large volume and on the associated large deviation estimates at that scale. Our main results demonstrate that these finite-size conditions imply the positivity of the infinite volume Lyapunov exponent. This paper shows that it is possible to make the techniques developed for the study of Schrödinger operators with deterministic potentials, based on large deviation estimates and the avalanche principle, effective.

1. INTRODUCTION

The study of Lyapunov exponents occupies a central role in ergodic theory and dynamical systems. They arise in a multitude of distinct settings, such as diffeomorphisms on a manifold, chaotic dynamics in nonlinear systems as exhibit by the standard map, cocycles defined over some base, and the theory of localization. Perhaps the most fundamental question about Lyapunov exponents relates to their simplicity. Or more quantitatively, to the gaps between them. In the case of $SL_2(\mathbb{R})$ cocycles this amounts to the question of positivity of the top Lyapunov exponent. Another much studied property of these exponents concerns their continuity relative to external parameters. For a beautiful introduction to this field see the textbook [Via].

This paper studies Schrödinger cocycles

$$(x, v) \in X \times \mathbb{R}^2 \mapsto (Tx, A_\lambda(x, E)v)$$

$$A_\lambda(x, E) = \begin{bmatrix} \lambda f(x) - E & -1 \\ 1 & 0 \end{bmatrix} \in SL_2(\mathbb{R})$$

where (X, μ, T) is some ergodic system, $\lambda, E \in \mathbb{R}$ and $f : X \rightarrow \mathbb{R}$ is measurable. These cocycles arise in the spectral analysis of the operators

$$(H_{\lambda, x}\psi)_n = \psi_{n+1} + \psi_{n-1} + \lambda f(T^n x)\psi_n, \quad n \in \mathbb{Z}$$

The authors thank the Institute for Advanced Study, Princeton, for its hospitality during the 2017-18 academic year. The third author was partially supported by the NSF, DMS-1500696. The authors thank James Maynard and Silviu Klein for helpful conversations, and Jean Bourgain and Thomas Spencer for their interest in this work.

Indeed, solutions of $H_{\lambda,x}\psi = E\psi$ are given by

$$(1.1) \quad \begin{pmatrix} \psi_{n+1} \\ \psi_n \end{pmatrix} = M_n(x; \lambda, E) \begin{pmatrix} \psi_1 \\ \psi_0 \end{pmatrix},$$

$$M_n(x; \lambda, E) = \prod_{j=n}^1 A_\lambda(T^j x, E), \quad n \geq 1$$

The growth of solutions to (1.1) μ -a.e. in x is governed by the Lyapunov exponent

$$L(\lambda, E) = \lim_{n \rightarrow \infty} n^{-1} \int_X \log \|M_n(x; \lambda, E)\| \mu(dx)$$

which always exists by subadditivity. By unimodularity of the matrices, $L(\lambda, E) \geq 0$. The main issue is then to determine strict positivity. We remark that by classical ergodic theory (Fürstenberg-Kesten theorem, Kingman's subadditive ergodic theorem [Via]),

$$n^{-1} \log \|M_n(x; \lambda, E)\| \rightarrow L(\lambda, E) \quad \mu \text{ a.s.}$$

as $n \rightarrow \infty$. Fürstenberg's theorem [Fur], shows that $L > 0$ for all λ, E for T the Bernoulli shift and μ a nontrivial probability distribution. Herman's subharmonicity argument [Her], which is recalled in Section 6, shows that $L(\lambda, E) \geq \log \lambda > 0$ if $\lambda > 1$, $X = \mathbb{T}$, $f(x) = 2 \cos(2\pi x)$, and $Tx = x + \omega$ a rotation (for general analytic f and large λ see [SorSpe]). On the other hand, one has $L(\lambda, E) = 0$ for all $0 < \lambda < 1$ and $E \in \text{spec}(H_{\lambda,x})$. The latter is the spectrum of the Harper or almost Mathieu operator

$$(H_{\lambda,x}\psi)_n = \psi_{n+1} + \psi_{n-1} + 2\lambda \cos(2\pi(x + n\omega))\psi_n, \quad n \in \mathbb{Z}$$

which does not depend on x (assuming ω irrational). In particular, $L(\lambda, 0) = 0$, cf. [BelSim, Dam].

In contrast to the Harper operator, its analog over the skew-shift base is conjectured to exhibit positive Lyapunov exponents for all $\lambda > 0$ and E . To be specific, let $X = \mathbb{T}^2$, $T(x, y) = (x + y, y + \omega)$, where ω is irrational (or Diophantine). Iterating T yields

$$(1.2) \quad M_n(x, y; \lambda, E) = \prod_{j=n}^1 \begin{bmatrix} 2\lambda f(x + jy + j(j-1)\omega/2, y + j\omega) - E & -1 \\ 1 & 0 \end{bmatrix}$$

The presence of $j^2\omega/2$ in these matrices appears to be the origin of the conjectured exponential growth of the norm of these matrices for all E (assuming $\partial_x f(x, y) \not\equiv 0$, with f analytic). In fact, the distribution of the fractional parts of $\{j^2\omega\}_{j=1}^N$ is known to be "random" in some sense as $N \rightarrow \infty$ for generic ω , see the Poissonian conjecture in [RudSarZah], as well as [MarStr, D.R.H-B]. Note that this is in stark contrast to the distribution of $\{j\omega\}_{j=1}^N$.

However, not only is this randomness property in and of itself delicate (see some negative results to this effect in [RudSarZah]), but how to use it in the context of (1.2) is entirely unclear. As far as rigorous results are concerned, Bourgain [Bou2]

proved that for all $\lambda > 0$ there exists a set of $\omega \in \mathbb{T}$ with positive measure (which decreases to 0 as $\lambda \rightarrow 0$), so that the operator

$$(H\psi)_n = \psi_{n+1} + \psi_{n-1} + \lambda \cos(n(n-1)\omega/2)\psi_n$$

exhibits point spectrum whose closure has positive measure. This was the first result of its kind which showed that for small λ the skew shift leads to completely different behavior than the shift, i.e., potentials $\cos(n\omega)$.

In this paper we present an effective multi-scale machinery aiming at positivity of the Lyapunov exponent for the matrices

$$(1.3) \quad M_n(x, y; \lambda, E) = \prod_{j=n}^1 \begin{bmatrix} 2\lambda \cos(2\pi(x + jy + j(j-1)\omega/2)) - E & -1 \\ 1 & 0 \end{bmatrix}$$

uniformly in E , and in the range $0 < \lambda \leq 1$. We fix ω to be the golden ratio. By the aforementioned estimate by Herman, one has $L(\lambda, E) \geq \log \lambda > 0$ for $\lambda > 1$. So only $\lambda \leq 1$ is of interest here. The basis of our analysis is the inductive argument from [BouGolSch], which established Anderson localization for large λ for the skew shift model, at the expense of removing a small set (in measure) of frequencies ω and phases (x, y) (the largeness of λ depended on the smallness of the measure of excluded parameters). The proof in [BouGolSch] is not effective, and it was not possible to explicitly determine the size of admissible λ in relation to the other parameters.

To formulate our main results, recall the finite-volume Lyapunov exponents

$$L_N(\lambda, E) := \int_{\mathbb{T}^2} \frac{1}{N} \log \|M_N(x, y; \lambda, E)\| \, dx dy,$$

and their limits $L = \lim_{N \rightarrow \infty} L_N$. We quantify the failure of the Fürstenberg-Kesten theorem via the following level sets \mathcal{B}_N ,

$$\mathcal{B}_N := \left\{ (x, y) \in \mathbb{T}^2 : \left| \frac{1}{N} \log \|M_N(x, y; \lambda, E)\| - L_N(\lambda, E) \right| > \frac{1}{10} L_N(\lambda, E) \right\}.$$

The machinery developed in this paper establishes a method for checking the positivity of the Lyapunov exponent $L(\lambda, E)$ by verifying information on a finite, initial scale. We could have formulated a very general “finite-size criterion” which establishes $L(\lambda, E) > 0$ under appropriate assumptions on the initial scale and for appropriate values of various other parameters. Instead, we have opted to present three representative theorems that can be obtained from the machinery developed in this paper by making specific choices.

These representative theorems differ by the precise assumptions (i)-(iii) made at the initial scale. We comment on this further after the first theorem. Moreover, the various other parameters appearing in our proof are identical in all three cases. These parameters are only chosen in the final part of the proof, Section 9, so they can easily be modified.

For a Borel set, $|\cdot|$ denotes the Lebesgue measure.

Theorem 1.1. *Consider the skew-shift cocycle given by (1.3). Let ω be the golden ratio and let $\lambda \in [1/2, 1]$. Let $N_0 := 2 \times 10^{37}$. Assume that for some energy $E \in [-2 - 2\lambda, 2 + 2\lambda]$ the following hold:*

- (i). $L_{N_0}(\lambda, E) \geq 2 \times 10^{-4}$,
- (ii). $L_{N_0}(\lambda, E) - L_{2N_0}(\lambda, E) \leq L_{N_0}(\lambda, E)/8$,
- (iii). $\max(|\mathcal{B}_{N_0}|, |\mathcal{B}_{2N_0}|) \leq N_0^{-21}$.

Then we have

$$L(\lambda, E) \geq \frac{1}{2}L_{N_0}(\lambda, E) > 0.$$

Before we give the two alternative theorems, we comment on conditions (i)-(iii).

Remark 1.2. (i). First, one might expect that $L(\lambda, E) > c\lambda^2$ holds for small λ , by analogy with the Figotin-Pastur asymptotics in the random case. Numerical experimentation suggests that is indeed the case for our model with $c > 10^{-2}$ (with a generous margin of error). Therefore, we would expect to have $L_{N_0}(\lambda, E) \geq 2 \times 10^{-3}$. Condition (i) was chosen to allow for an even wider margin. We remark that we can lower the number 2×10^{-4} to basically any positive constant, at the expense of increasing N_0 .

(ii). Condition (ii) is known to hold if the Lyapunov exponent is positive and N_0 is large enough. Indeed, it follows from the methods in [GolSch] that

$$L_{N_0}(\lambda, E) - L_{2N_0}(\lambda, E) \leq cL_{N_0}(\lambda, E)/N_0$$

with some absolute constant $c \sim 1$, see Section 8 below for the details. Given the size of N_0 , condition (ii) is indeed asking for very little.

(iii). Finally, condition (iii) is some weak form of a *large deviation estimate* as in [BouGol, GolSch, BouGolSch]. In fact, analogy with these references suggests that a bound of the form $|\mathcal{B}_N| < \exp(-N^{\frac{1}{10}})$ should hold for large N (and perhaps a much stronger bound, say with $N^{\frac{1}{2}}$ or larger). For (iii) to hold in this case would then require $N > 8 \cdot 10^{31}$, which is within our range. It is important to note that condition (iii) differs strongly from (i) and (ii). Indeed, while the latter conditions are intimately related to the $L > 0$, (iii) is not. For the shift dynamics it is known that the large deviation estimates hold apriori, i.e., without any reference to the positivity of the Lyapunov exponent, see [BouGol, GolSch, Bou1]. For the skew shift, however, such apriori derivations are currently not known. Rather, we rely an inductive procedure that uses lower bounds on L_n , the Lyapunov exponents in finite volume.

We now state two further representative theorems. These alternative finite-size criteria both involve much smaller initial scales N_0 , at the price of having a more restrictive assumption (iii) on the measure of the set \mathcal{B}_{N_0} .

Theorem 1.3. *Consider the skew-shift cocycle given by (1.3). Let ω be the golden ratio and let $\lambda \in [1/2, 1]$. Let $N_0 := 3 \times 10^5$. Assume that for some energy $E \in [-2 - 2\lambda, 2 + 2\lambda]$ the following hold:*

- (i). $L_{N_0}(\lambda, E) \geq 2 \times 10^{-4}$,
- (ii). $L_{N_0}(\lambda, E) - L_{2N_0}(\lambda, E) \leq L_{N_0}(\lambda, E)/8$,
- (iii). $\max(|\mathcal{B}_{N_0}|, |\mathcal{B}_{2N_0}|) \leq N_0^{-141}$.

Then we have

$$L(\lambda, E) \geq \frac{1}{2}L_{N_0}(\lambda, E) > 0.$$

The upper bound in assumption (iii) is more restrictive than in Theorem 1.3. Importantly, it is still polynomial in nature. Hence, in view of Remark 1.2 (iii), it may hold depending on the precise kind of exponential decay that is presumably exhibited by the true $|\mathcal{B}_N|$.

In the next representative result, Theorem 1.4, we strengthen assumption (i) somewhat (in a way that is compatible with the numerics described in Remark 1.2 (i) above). This allows to reduce the initial scale even further, to the value $N_0 = 3 \times 10^4$, which may be amenable to numerical investigation.

Theorem 1.4. *Consider the skew-shift cocycle given by (1.3). Let ω be the golden ratio and let $\lambda \in [1/2, 1]$. Let $N_0 := 3 \times 10^4$. Assume that for some energy $E \in [-2 - 2\lambda, 2 + 2\lambda]$ the following hold:*

- (i). $L_{N_0}(\lambda, E) \geq 2 \times 10^{-3}$,
- (ii). $L_{N_0}(\lambda, E) - L_{2N_0}(\lambda, E) \leq L_{N_0}(\lambda, E)/8$,
- (iii). $\max(|\mathcal{B}_{N_0}|, |\mathcal{B}_{2N_0}|) \leq N_0^{-165}$.

Then we have

$$L(\lambda, E) \geq \frac{1}{2}L_{N_0}(\lambda, E) > 0.$$

Regarding assumption (iii), the comment made after Theorem 1.3 still applies. In particular, the relatively small value of N_0 in this result elevates the problem of finding an **analytical proof** of (iii). Indeed, (i) and (ii) are accessible numerically by a Figotin-Pastur expansion, for example, but it seems completely unreasonable to ask for a computer assisted proof of (iii).

The restriction $\lambda \in [1/2, 1]$ was chosen for convenience. In fact, our methods apply to any given interval of the form $[\lambda_0, 1]$, $\lambda_0 > 0$, albeit with increasing N_0 as $\lambda_0 \rightarrow 0$. Similarly, the golden ratio was chosen for simplicity. One can replace it by a class of Diophantine frequencies obeying an explicit Diophantine condition.

It remain to be seen what the true range of applicability of our methods are, and to which extent they can also be refined. It may be possible to verify assumptions (i) and (ii) of Theorem 1.4 numerically. However, the measure estimates (iii) would seem the most delicate to check reliably.

The methods in this paper are an adaptation of those in [GolSch, BouGolSch, Bou1]. One of our motivations was to obtain an effective rendition of the techniques based on harmonic analysis (subharmonic functions, Riesz representation theorem, John-Nirenberg type estimates for BMO functions) in combination with linear algebra and the geometry of matrix products (“avalanche principle”, [GolSch]). This had never been attempted before, but we show that it is possible to do so.

2. EFFECTIVE RIESZ REPRESENTATION

It is of fundamental importance to the entire method to make the underlying potential theory effective. To this end it is most convenient to remain on the disk since other geometries will lead to complicated Green functions. The disk will suffice for our purposes, thanks to a variant of Herman's regularization [Her], which we present in Section 6.

Definition 2.1. Given $R > 0$, we write D_R for the open disk of radius R around the origin in \mathbb{C} . Let $z = re(\phi)$, we write $P_z(\theta)$ for the Poisson kernel

$$P_z(\theta) := \frac{1 - r^2}{1 - 2r \cos(2\pi(\phi - \theta)) + r^2}.$$

The following constants will be used throughout, with $1 < R_2 < R_1 < R$:

(2.1)

$$B_0(R, R_1, R_2) := \frac{1}{2 \log(R/R_1)} \left(\frac{R_1 + R_2}{R_1 - R_2} \right) \times \begin{cases} \log R & \text{if } R^2 - R_1^2 > R, \\ \log \left(\frac{R^2}{R^2 - R_1^2} \right), & \text{if } R^2 - R_1^2 < R. \end{cases}$$

$$B_1(R, R_1, R_2) := B_0(R, R_1, R_2) \frac{8R_2}{R_2^2 - 1},$$

$$B_2(R, R_1, R_2) := B_0(R, R_1, R_2)$$

$$\frac{16\pi(R_2^2 - 1) \sqrt{16R_2^2 - \left(\sqrt{R_2^4 + 34R_2^2 + 1} - 1 - R_2^2 \right)^2}}{\left(3R_2^2 + 3 - \sqrt{R_2^4 + 34R_2^2 + 1} \right)^2}$$

as well as

$$(2.2) \quad B_3(R, R_1, R_2) := \sqrt{5B_2(R, R_1, R_2) + \frac{10\pi}{\log R/R_1}}.$$

The main result of this section is the following Riesz representation theorem for subharmonic functions. The essential feature here are the explicitly computable constants. Recall that a subharmonic function in some domain $\Omega \subset \mathbb{C}$ is an upper semicontinuous function $u : \Omega \rightarrow \mathbb{R} \cup \{-\infty\}$ which satisfies the sub-mean value property in Ω .

Theorem 2.2. *Let $1 < R_2 < R_1 < R$ and let $v : \overline{D_R} \rightarrow \mathbb{R} \cup \{-\infty\}$ be a subharmonic function satisfying*

$$(2.3) \quad v(z) \leq B, \quad v(0) = m.$$

Then, for all $w \in D_{R_1}$, we have the Riesz representation

$$(2.4) \quad v(w) = \int_{D_{R_1}} \log |z - w| \mu(dz) + h(w),$$

where

(i) μ is a positive measure satisfying the bound

$$(2.5) \quad \mu(D_{R_1}) \leq \frac{B - m}{\log(R/R_1)},$$

(ii) h is harmonic on D_{R_1} and satisfies the following bounds

$$(2.6) \quad \min_{c \in \mathbb{R}} \max_{|w| \leq R_2} |h(w) - c| \leq B_0(R, R_1, R_2)(B - m),$$

$$\left| \frac{d^k}{d\varphi^k} h(e(\varphi)) \right| \leq B_k(R, R_1, R_2)(B - m), \quad (k = 1, 2),$$

The proof of this theorem will occupy the rest of this section.

Proof of Theorem 2.2. The basic idea of the proof is that the equation $\mu = \frac{1}{2\pi} \Delta v$ holds in the distributional sense, with μ a positive measure. Without loss of generality, we may assume that v is smooth. If this is not the case, we convolve v with a radial nonnegative mollifier. The submean property then guarantees monotone convergence. We skip these technical details.

2.1. Riesz representation. Rescaling the unit disk yields the Green function on any disk.

Lemma 2.3 (Green's function for the disk). *The function $G : D_R \times D_R \rightarrow \mathbb{R}$ given by*

$$G(z, w) := \frac{1}{2\pi} \log \left| \frac{R(z - w)}{R^2 - z\bar{w}} \right|$$

satisfies $\Delta_z G(z, w) = \delta_w$ and $G(z, w) = 0$ when $|z| = R$.

Proof. To see this, notice that $G(z, w) = G_1(z/R, w/R)$ where

$$G_1(z, w) := \frac{1}{2\pi} \log \left| \frac{z - w}{1 - z\bar{w}} \right|$$

is the Green function of the unit disk. □

Let $w \in D_{R_1}$. By Green's second identity for the domain D_R , we have

$$v(w) - \int_{D_R} G(z, w) \Delta v(z) \text{Vol}(dz) = \int_{\partial D_R} v(z) \frac{\partial G}{\partial n_z}(z, w) \sigma(dz),$$

where Vol is the standard volume measure and σ is the (unnormalized) arclength measure on the circle ∂D_R . Since v is smooth and subharmonic, Δv is a non-negative, continuous function, call it $2\pi\mu$. Therefore

$$(2.7) \quad v(w) = \int_{D_R} 2\pi G(z, w) \mu(dz) + h_0(w),$$

where

$$(2.8) \quad h_0(w) := \int_{\partial D_R} v(z) \frac{\partial G}{\partial n_z}(z, w) \sigma(dz).$$

By Lemma 2.3, we then have Riesz representation with the functions

$$(2.9) \quad v(w) = \int_{D_{R_1}} \log |z - w| \mu(dz) + h(w),$$

where

$$(2.10) \quad h(w) := \int_{D_R \setminus D_{R_1}} \log \left| \frac{R(z-w)}{R^2 - z\bar{w}} \right| \mu(dz) + \int_{D_{R_1}} \log \left| \frac{R}{R^2 - z\bar{w}} \right| \mu(dz) + h_0(w)$$

Lemma 2.4. *$h(w)$ is harmonic in D_{R_1} .*

Proof. Write $w = re^{2\pi i\varphi}$. The first and second term in (2.10) are harmonic because they are real parts of analytic functions on D_{R_1} . For the third term, recall that $\frac{\partial G}{\partial n_z}(z, w)$ is the Poisson kernel whence

$$(2.11) \quad h_0(w) = \int_0^1 v(Re(\theta)) P_{r/R}(\varphi - \theta) d\theta.$$

The Poisson kernel is harmonic in all of D_R and this proves the lemma. \square

2.2. Control of the Riesz mass.

Lemma 2.5. *We have the following bound on the Riesz mass:*

$$(2.12) \quad \mu(D_{R_1}) \leq \frac{B - m}{\log(R/R_1)}.$$

Proof. Taking $w = 0$ in (2.7), we see

$$(\log R/R_1)\mu(D_{R_1}) \leq \int_{D_R} \log \frac{R}{|z|} \mu(dz) = h_0(0) - v(0) \leq B - m,$$

in which we used

$$(2.13) \quad h_0(0) \leq B,$$

which comes from the maximum principle and the fact that $h_0(w)$ is the harmonic function on D_R with boundary values $v(\partial D_R)$ by (2.11). \square

2.3. Control of the harmonic part. We have the following estimate for the harmonic part.

Lemma 2.6. *Let $1 < R_2 < R_1 < R$.*

$$\min_{c \in \mathbb{R}} \max_{|w| \leq R_2} |h(w) - c| \leq B_0(R, R_1, R_2)(B - m),$$

with constant $B_0(R, R_1, R_2)$ given by (2.1).

Proof. We will first prove an upper bound and then use Harnack's inequality to conclude a lower bound. From (2.10), and $G(z, w) \leq 0$ on $D_R \times D_R$,

$$h(w) \leq \int_{D_{R_1}} \log \left| \frac{R}{R^2 - z\bar{w}} \right| \mu(dz) + h_0(w).$$

From (2.13), we infer that for all $w \in D_{R_1}$,

$$h(w) \leq \log \left| \frac{R}{R^2 - R_1^2} \right| \mu(D_{R_1}) + B.$$

Now we distinguish cases. On the one hand, if $R < R^2 - R_1^2$, then the logarithm is negative and (2.14) implies

$$h(w) \leq B.$$

On the other hand, if $R > R^2 - R_1^2$, then we use Lemma 2.5 to obtain

$$\begin{aligned} h(w) &\leq \frac{1}{\log R/R_1} \left(B \log \frac{R^2}{R_1(R^2 - R_1^2)} - m \log \frac{R}{R^2 - R_1^2} \right) \\ &= \frac{m \log R - B \log R_1}{\log R/R_1} + \frac{\log R^2/(R^2 - R_1^2)}{\log R/R_1} (B - m). \end{aligned}$$

Combining the two cases, we arrive at the upper bound (2.14)

$$h(w) \leq \alpha, \quad \alpha := \begin{cases} B, & \text{if } R^2 - R_1^2 > R, \\ \frac{m \log R - B \log R_1}{\log R/R_1} + \frac{\log R^2/(R^2 - R_1^2)}{\log R/R_1} (B - m), & \text{if } R^2 - R_1^2 < R. \end{cases}$$

Consider the non-negative harmonic function $\alpha - h(w)$ on D_{R_1} . By Harnack's inequality,

$$\alpha - h(w) \leq \frac{R_1 + |w|}{R_1 - |w|} (\alpha - h(0)),$$

which implies the following lower bound

$$(2.15) \quad h(w) \geq \frac{R_1 + |w|}{R_1 - |w|} h(0) - \frac{2|w|}{R_1 - |w|} \alpha.$$

By (2.9) with $w = 0$ and (2.5), we have

$$h(0) = v(0) - \int_{D_{R_1}} \log |z| \mu(dz) \geq m - \frac{\log R_1}{\log R/R_1} (B - m).$$

Together with (2.15), this yields

$$(2.16) \quad h(w) \geq \frac{R_1 + |w|}{R_1 - |w|} \frac{m \log R - B \log R_1}{\log R/R_1} - \frac{2|w|}{R_1 - |w|} \alpha.$$

Based on (2.14) and (2.16), we obtain

$$\begin{aligned} &\min_{c \in \mathbb{R}} \max_{|w| \leq R_2} |h(w) - c| \\ &\leq \frac{1}{2} \left(\alpha - \min_{|w| \leq R_2} \left(\frac{R_1 + |w|}{R_1 - |w|} \frac{m \log R - B \log R_1}{\log R/R_1} - \frac{2|w|}{R_1 - |w|} \alpha \right) \right) \\ &= \frac{1}{2} \max_{|w| \leq R_2} \left(\frac{R_1 + |w|}{R_1 - |w|} \right) \left(\alpha - \frac{m \log R - B \log R_1}{\log R/R_1} \right) \\ &= \frac{1}{2} \left(\frac{R_1 + R_2}{R_1 - R_2} \right) \left(\alpha - \frac{m \log R - B \log R_1}{\log R/R_1} \right) \\ &= B_0(R, R_1, R_2)(B - m). \end{aligned}$$

This proves Lemma 2.6. □

Lemma 2.7. *For $k = 1, 2$, we have*

$$\left| \frac{d^k}{d\varphi^k} h(e^{2\pi i\varphi}) \right| \leq B_k(R, R_1, R_2)(B - m)$$

with constants B_1, B_2 given by (2.1).

Proof. Since h is harmonic in D_{R_1} , we have that for any constant c ,

$$h(e^{2\pi i\varphi}) - c = \int_0^1 (h(R_2 e^{2\pi i\theta}) - c) P_{1/R_2}(\varphi - \theta) d\theta.$$

We take a derivative in φ and estimate h using Lemma 2.6. This gives

$$\begin{aligned} \left| \frac{d}{d\varphi} h(e^{2\pi i\varphi}) \right| &\leq \left(\min_{c \in \mathbb{R}} \max_{|w| \leq R_2} |h(w) - c| \right) \int_0^1 \left| \frac{\partial}{\partial \varphi} P_{1/R_2}(\varphi - \theta) \right| d\theta \\ &\leq B_0(R, R_1, R_2)(B - m) \int_0^1 \left| \frac{\partial}{\partial \theta} P_{1/R_2}(\theta) \right| d\theta. \end{aligned}$$

We recall that

$$P_{1/R_2}(\theta) = \frac{R_2^2 - 1}{R_2^2 - 2R_2 \cos(2\pi\theta) + 1}$$

and therefore

$$\frac{\partial}{\partial \theta} P_{1/R_2}(\theta) = -\frac{4\pi R_2(R_2^2 - 1) \sin(2\pi\theta)}{(R_2^2 - 2R_2 \cos(2\pi\theta) + 1)^2}.$$

Since $\sin(2\pi\theta)$ changes sign at $\theta = 1/2$, we conclude that

$$\begin{aligned} \int_0^1 \left| \frac{\partial}{\partial \theta} P_{1/R_2}(\theta - \varphi) \right| d\theta &= -\int_0^{1/2} \frac{\partial}{\partial \theta} P_{1/R_2}(\theta) d\theta + \int_{1/2}^1 \frac{\partial}{\partial \theta} P_{1/R_2}(\theta) d\theta \\ &= 2P_{1/R_2}(0) - 2P_{1/R_2}(1/2) \\ &= \frac{8R_2}{R_2^2 - 1}. \end{aligned}$$

This proves the claim for $k = 1$.

For the second derivative, we argue similarly. We have

$$\left| \frac{d^2}{d\varphi^2} h(e^{2\pi i\varphi}) \right| \leq B_0(R, R_1, R_2)(B - m) \int_0^1 \left| \frac{\partial^2}{\partial \theta^2} P_{1/R_2}(\theta) \right| d\theta,$$

where

$$\frac{\partial^2}{\partial \theta^2} P_{1/R_2}(\theta) = \frac{-8\pi^2 R_2(R_2^2 - 1)(2R_2 \cos^2(2\pi\theta) + (R_2^2 + 1) \cos(2\pi\theta) - 4R_2)}{(R_2^2 - 2R_2 \cos(2\pi\theta) + 1)^3}.$$

By symmetry, we may restrict our attention to $\theta \in [0, 1/2]$ from now on. On that interval, the function $\frac{\partial^2}{\partial \theta^2} P_{1/R_2}$ has exactly one zero. Its location, call it $\theta_0 \in [0, 1/2]$, is given by

$$(2.17) \quad \theta_0 = \frac{1}{2\pi} \arccos \left(\frac{\sqrt{R_2^4 + 34R_2^2 + 1} - (R_2^2 + 1)}{4R_2} \right).$$

It is easy to see that $\frac{\partial^2}{\partial\theta^2}P_{1/R_2}$ is negative on $[0, \theta_0)$, and hence positive on $(\theta_0, 1/2]$. Therefore

$$\begin{aligned} \int_0^1 \left| \frac{\partial^2}{\partial\theta^2}P_{1/R_2}(\theta) \right| d\theta &= -2 \int_0^{\theta_0} \frac{\partial^2}{\partial\theta^2}P_{1/R_2}(\theta) d\theta + 2 \int_{\theta_0}^{1/2} \frac{\partial^2}{\partial\theta^2}P_{1/R_2}(\theta) d\theta \\ &= 2 \frac{\partial}{\partial\theta}P_{1/R_2}(0) - 4 \frac{\partial}{\partial\theta}P_{1/R_2}(\theta_0) + 2 \frac{\partial}{\partial\theta}P_{1/R_2}(1/2) \\ &= \frac{16\pi R_2(R_2^2 - 1) \sin(2\pi\theta_0)}{(R_2^2 - 2R_2 \cos(2\pi\theta_0) + 1)^2}. \end{aligned}$$

since $\frac{\partial}{\partial\theta}P_{1/R_2}(0) = \frac{\partial}{\partial\theta}P_{1/R_2}(1/2) = 0$. When we evaluate the last expression using the definition (2.17) of θ_0 , we obtain the quantity

$$\frac{16\pi(R_2^2 - 1) \sqrt{16R_2^2 - \left(1 + R_2^2 - \sqrt{R_2^4 + 34R_2^2 + 1}\right)^2}}{\left(\sqrt{R_2^4 + 34R_2^2 + 1} - 3R_2^2 - 3\right)^2}.$$

This proves the claim for $k = 2$. \square

3. \mathbb{T}^1 SPLITTING LEMMA

For any $f \in L^1(\mathbb{T})$, let $\langle f \rangle = \int_{\mathbb{T}} f(x) dx$. For a function f on \mathbb{C} , let us denote $f(e(x))$ by $f(x)$ for simplicity. For a Borel set U , let $|U|$ be its Lebesgue measure.

Lemma 3.1. *Let v be as in Theorem 2.2. Assume that for some constant c*

$$(3.1) \quad v(x) = v_1(x) + v_0(x) + c$$

with $\|v_1\|_{L^1(\mathbb{T})} < \varepsilon_1$ and $\|v_0\|_{L^\infty(\mathbb{T})} < \varepsilon_0$. Then we have

$$(3.2) \quad \begin{aligned} \int_{\mathbb{T}} \exp\left(\frac{\pi}{4}\delta_0^{-1}|v(x) - c|\right) dx &\leq C_0 \\ \delta_0 &:= \frac{9}{2}\varepsilon_0 + 2B_3(R, R_1, R_2)\sqrt{\varepsilon_1(B - m)} \\ C_0 &:= 2\sqrt{2} \exp\left(\pi\left[\frac{17}{144} + \frac{B_1}{16B_3^2}\right]\right) \end{aligned}$$

with constants given by Definition 2.1.

As a corollary of the exponential integrability, we have the following estimate on the level sets from Markov's inequality.

Corollary 3.2. *For any $\varepsilon_2 > 0$, we have*

$$|\{x \in \mathbb{T} : |v(x)| > \varepsilon_2\}| \leq 2\sqrt{2} \exp\left(\frac{\pi}{4}\left[\frac{17}{144} + \frac{B_1}{16B_3^2} - \varepsilon_2\delta_0^{-1}\right]\right)$$

with δ_0 as in (3.2).

Note that this level set estimate is only useful if $\varepsilon_2 \gg \varepsilon_0$ and $\varepsilon_2^2 \gg \varepsilon_1(B - m)$.

Proof of Lemma 3.1. For simplicity, we will denote $B_3(R, R_1, R_2)$ by B_3 throughout the proof. We will first show the following special form of the Riesz representation, valid only on the unit circle. The idea is simply to reflect the part of the disk outside the circle back inside of it.

Lemma 3.3. *Let $1 < R_2 < R_1 < R$ and v be defined as in Theorem 2.2. Then there exists a positive measure $\tilde{\mu}$ and a harmonic function \tilde{h} on D_R such that*

$$(3.3) \quad v(e(\varphi)) = \int_{D_1} \log |z - e(\varphi)| \tilde{\mu}(dz) + \tilde{h}(e(\varphi)),$$

with the following estimates

$$(3.4) \quad \tilde{\mu}(\overline{D_1}) \leq \frac{B - m}{\log R/R_1}.$$

and \tilde{h} satisfies the bound Lemma 2.6 on the circle as well as

$$(3.5) \quad \left| \frac{d^k}{d\varphi^k} \tilde{h}(e(\varphi)) \right| \leq B_k(R, R_1, R_2)(B - m), \quad k = 1, 2,$$

where B_1, B_2 are the same constants as those in Theorem 2.2.

Proof. By Theorem 2.2, we have

$$(3.6) \quad v(w) = \int_{D_{R_1}} \log |z - w| \mu(dz) + h(w),$$

with $\mu(D_{R_1}) \leq \frac{B-m}{\log R/R_1}$, and $\left| \frac{d^k}{d\varphi^k} h(e(\varphi)) \right| \leq B_k(R, R_1, R_2)(B - m)$, $k = 1, 2$.

Let us define μ^* by reflection, i.e.,

$$(3.7) \quad \mu^*(E) = \mu(E^*),$$

where

$$E^* = \{z^{-1} : z \in E\}$$

for any measurable set $E \subset \mathbb{C}$. Then for any $|w| = 1$,

$$\begin{aligned} & \int_{D_{R_1}} \log |z - w| \mu(dz) \\ &= \int_{D_1} \log |z - w| \mu(dz) + \int_{D_{R_1} \setminus \overline{D_1}} \log |z - w| \mu(dz) \\ &= \int_{D_1} \log |z - w| \mu(dz) + \int_{D_{R_1} \setminus \overline{D_1}} \log |w - z^{-1}| \mu(dz) + \int_{D_{R_1} \setminus \overline{D_1}} \log |z| \mu(dz) \\ &= \int_{D_1} \log |z - w| \mu(dz) + \int_{D_1 \setminus \overline{D_1/R_1}} \log |w - z| \mu^*(dz) + \int_{D_{R_1} \setminus \overline{D_1}} \log |z| \mu(dz) \\ (3.8) \quad &= \int_{D_1} \log |z - w| \tilde{\mu}(dz) + \int_{D_{R_1} \setminus \overline{D_1}} \log |z| \mu(dz), \end{aligned}$$

where, for any $E \subset \overline{D_1}$,

$$(3.9) \quad \tilde{\mu}(E) = \mu(E) + \mu^*(E \cap (D_1 \setminus \overline{D_1/R_1})) = \mu(E) + \mu(E^* \cap (D_{R_1} \setminus \overline{D_1})).$$

By (3.9), it is clear that we have the following estimate for $\tilde{\mu}$

$$(3.10) \quad \tilde{\mu}(\overline{D_1}) = \mu(D_{R_1}) \leq \frac{B-m}{\log(R/R_1)}.$$

By (3.6) and (3.8), we have for $|w| = 1$,

$$(3.11) \quad v(w) = \int_{D_1} \log |z-w| \tilde{\mu}(dz) + h(w) + \int_{D_{R_1} \setminus \overline{D_1}} \log |z| \mu(dz),$$

in which the third term is a constant. Let us take $\tilde{h} = h + \int_{D_{R_1} \setminus \overline{D_1}} \log |z| \mu(dz)$. Since \tilde{h} only differs from h by a constant, the estimates on the derivatives still hold. \square

The Riesz representation (3.3) allows us to give upper bounds on the parameters ε_0 and ε_1 in Lemma 3.1 in terms of $B-m$. This will be relevant in the proof of that lemma.

Corollary 3.4. *We may always assume in Lemma 3.1 that*

$$(3.12) \quad \varepsilon_0 \leq B_0(R, R_1, R_2)(B-m), \quad \varepsilon_1 \leq \frac{13}{20} \frac{B-m}{\log(R/R_1)}$$

Alternatively, we can assume that $\varepsilon_0 = 0$ and

$$(3.13) \quad \varepsilon_1 = \left(B_0(R, R_1, R_2) + \frac{13}{20 \log(R/R_1)} \right) (B-m)$$

Proof. In view of (3.3) we set

$$v_0(\varphi) := \tilde{h}(e(\varphi)), \quad v_1(\varphi) = \int_{D_1} \log |z - e(\varphi)| \tilde{\mu}(dz)$$

Then ε_0 is the constant from Lemma 2.6 and we claim that

$$\varepsilon_1 := \|\log |1 - e(\varphi)|\|_{L^1_\varphi} \|\tilde{\mu}\|$$

is an admissible choice. Indeed,

$$\begin{aligned} \|v_1\|_{L^1(\mathbb{T})} &\leq \int_{D_1} \|\log |z - e(\varphi)|\|_{L^1(\mathbb{T}_\varphi)} \tilde{\mu}(dz) \\ &= \int_{D_1} \|\log ||z| - e(\varphi)|\|_{L^1(\mathbb{T}_\varphi)} \tilde{\mu}(dz) \\ &\leq \max_{0 \leq r \leq 1} \|\log |r - e(\varphi)|\|_{L^1(\mathbb{T}_\varphi)} \|\tilde{\mu}\| \end{aligned}$$

Set $h(r) := \|\log |r - e(\varphi)|\|_{L^1(\mathbb{T}_\varphi)}$ with $0 \leq r \leq 1$. In order to establish the claim, it suffices to verify that $h(r)$ is nondecreasing. First,

$$\int_0^1 \log |r - e(\varphi)| d\varphi = \int_0^1 \log |1 - re(\varphi)| d\varphi = 0$$

since $\log|1 - r\zeta|$ is harmonic in ζ for $|\zeta| < 1$ and any fixed $0 \leq r \leq 1$. Therefore, if $0 < r < 1$ and $0 < \varphi_0(r) < \frac{1}{2}$ is the unique solution of $|r - e(\varphi_0)| = 1$, then

$$h(r) = 2 \int_{\varphi_0(r)}^{1-\varphi_0(r)} \log|r - e(\varphi)| d\varphi = \int_{\varphi_0(r)}^{1-\varphi_0(r)} \log(1 + r^2 - 2r \cos(2\pi\varphi)) d\varphi$$

Consequently,

$$\begin{aligned} h'(r) &= \int_{\varphi_0(r)}^{1-\varphi_0(r)} \frac{2(r - \cos(2\pi\varphi))}{1 + r^2 - 2r \cos(2\pi\varphi)} d\varphi \\ &\geq \int_{\varphi_0(r)}^{1-\varphi_0(r)} \frac{r}{1 + r^2 - 2r \cos(2\pi\varphi)} d\varphi \geq 0 \end{aligned}$$

In the second line we used that on the domain of integration

$$|r - e(\varphi)|^2 = 1 + r^2 - 2r \cos(2\pi\varphi) \geq 1$$

whence $2r - 2 \cos(2\pi\varphi) \geq r$. Therefore, indeed $h(r) \leq h(1)$, justifying our choice of ε_1 above. Finally,

$$\begin{aligned} h(1) &= \|\log|1 - e(\varphi)|\|_{L^1_{\varphi}} = -2 \int_0^1 \min(\log|1 - e(\varphi)|, 0) d\varphi \\ &= -2 \int_{-\frac{1}{6}}^{\frac{1}{6}} \log|1 - e(\varphi)| d\varphi = -4 \int_0^{\frac{1}{6}} \log(2 \sin(\pi\varphi)) d\varphi < 13/20. \end{aligned}$$

and $\|\tilde{\mu}\|$ is controlled by (3.4). □

Definition 3.5. Henceforth we impose the condition that

$$(3.14) \quad 289(B_0(R, R_1, R_2) + \frac{13}{20 \log R/R_1}) < B_3^2(R, R_1, R_2)$$

where the constants are those from Definition 2.1.

Returning to the proof of Lemma 3.1, we denote the first term in (3.3) by u , viz.

$$(3.15) \quad u(x) = \int_{\mathcal{D}_1} \log|z - e(x)| \tilde{\mu}(dz).$$

Then $v = u + \tilde{h}$. For any $f \in L^1(\mathbb{T})$ with $\langle f \rangle = 0$, the anti-derivative $D^{-1}f$ is uniquely defined as the absolutely continuous function

$$(3.16) \quad (D^{-1}f)(t) = \int_0^t f(x) dx + m(f), \quad \langle D^{-1}f \rangle = 0,$$

for arbitrary $t \in \mathbb{T}$. The constant $m(f)$ is chosen to ensure the vanishing mean. For $e_n(x) = \exp(2\pi inx)$, one has $D^{-1}(e_n) = (2\pi in)^{-1}e_n$ for all $n \neq 0$, whereas $D^{-1}e_0 = 0$. In the distributional sense, D^{-1} also applies to (complex) measures. For example, with δ_0 now being the Dirac delta,

$$D^{-1}(\delta_0 - 1)(x) = -(x + \frac{1}{2})\mathbf{1}_{[-\frac{1}{2} < x < 0]}(x) + (\frac{1}{2} - x)\mathbf{1}_{[0 < x < \frac{1}{2}]}(x).$$

For any $z = |z|e(y) \in D_1$, one has the elementary relation

$$(3.17) \quad \begin{aligned} \frac{d}{dx} \log |e(x) - z| &= \frac{2\pi|z| \sin(2\pi(x-y))}{1 - 2|z| \cos(2\pi(x-y)) + |z|^2} \\ &= \pi Q_z(x) \\ &= \pi(\mathcal{H}[P_z])(x), \end{aligned}$$

where \mathcal{H} denotes the Hilbert transform and Q_z is the standard notation for the conjugate function of the Poisson kernel. In particular,

$$(3.18) \quad \log |e(x) - z| = \pi(D^{-1}\mathcal{H}[P_z])(x)$$

holds for any $z \in D_1$. We thus have

$$(3.19) \quad u(x) = (D^{-1}\mathcal{H}[\nu])(x),$$

where

$$\frac{d\nu}{dx}(x) = \pi \int_{D_1} P_z(x) \tilde{\mu}(dz)$$

is a positive measure, with

$$(3.20) \quad \nu(\mathbb{T}) = \pi \tilde{\mu}(\overline{D_1}).$$

Set

$$(3.21) \quad \epsilon := \frac{1}{B_3} \sqrt{\frac{\epsilon_1}{B-m}},$$

and define $J_\epsilon(x) := \frac{1}{2\epsilon} \mathbb{1}_{[-\epsilon, \epsilon]}(x)$ to be the box kernel. Because of the upper bound in (3.13) on ϵ_1 and (3.14), one has

$$\epsilon_1 \leq (B_0(R, R_1, R_2) + \frac{13}{20 \log R/R_1})(B-m) < \frac{B_3^2}{289}(B-m)$$

which ensures that $\epsilon < \frac{1}{17}$. We will use this smallness property for the remainder of the proof. For example, it guarantees that J_ϵ is in fact well-defined on the circle (less than $\frac{1}{2}$ is enough here, but below we will need this sharper bound). Then

$$\begin{aligned} v &= v - J_\epsilon * v + J_\epsilon * v_1 + J_\epsilon * v_0 \\ &= (u - J_\epsilon * u) + (\tilde{h} - J_\epsilon * \tilde{h}) + J_\epsilon * v_1 + J_\epsilon * v_0. \end{aligned}$$

The last three terms have small L^∞ norms, in the sense that

$$\left\{ \begin{array}{l} \|J_\epsilon * v_1\|_{L^\infty(\mathbb{T})} \leq \|J_\epsilon\|_{L^\infty(\mathbb{T})} \|v_1\|_{L^1(\mathbb{T})} < \frac{\epsilon_1}{2\epsilon} = \frac{B_3}{2} \sqrt{\epsilon_1(B-m)}, \\ \|J_\epsilon * v_0\|_{L^\infty(\mathbb{T})} \leq \|J_\epsilon\|_{L^1(\mathbb{T})} \|v_0\|_{L^\infty(\mathbb{T})} < \epsilon_0, \\ \|\tilde{h} - J_\epsilon * \tilde{h}\|_{L^\infty(\mathbb{T})} \leq \frac{\epsilon}{2} \|\tilde{h}'\|_{L^\infty(\mathbb{T})} \leq \frac{1}{2} \epsilon B_1 (B-m) = \frac{B_1}{2B_3} \sqrt{\epsilon_1(B-m)}. \end{array} \right.$$

Hence

$$(3.22) \quad |(u - J_\epsilon * u)(x)| \geq |v(x) - C| - \epsilon_0 - \left(\frac{B_3}{2} + \frac{B_1}{2B_3} \right) \sqrt{\epsilon_1(B-m)}.$$

By (3.19), we have

$$(3.23) \quad \begin{aligned} (u - J_\epsilon * u)(x) &= (D^{-1}\mathcal{H}[\nu - J_\epsilon * \nu])(x) \\ &= \mathcal{H}[D^{-1}(\nu - J_\epsilon * \nu)](x). \end{aligned}$$

Next, we control the pointwise size of the term in brackets in (3.23). Since the Hilbert transform eliminates constants, the integration constant in (3.16) drops out.

Lemma 3.6. *Modulo additive constants the function $D^{-1}(\nu - J_\epsilon * \nu)$ satisfies*

$$(3.24) \quad \|D^{-1}(\nu - J_\epsilon * \nu)\|_{L^\infty(\mathbb{T})} \leq \frac{9}{2}\varepsilon_0 + 2B_3\sqrt{\varepsilon_1(B-m)}$$

Proof. We begin with the observation that (recall ν is a positive measure)

$$(3.25) \quad \begin{aligned} |(\nu - J_\epsilon * \nu)([a, b])| &= \left| \int_{\mathbb{T}} (\mathbb{1}_{[a, b]} - \mathbb{1}_{[a, b]} * J_\epsilon)(x) \nu(dx) \right| \\ &\leq \sup_{\theta \in \mathbb{T}} \nu([\theta - \epsilon, \theta + \epsilon]), \end{aligned}$$

uniformly in $[a, b] \subset \mathbb{T}$. If $b - a \geq 2\epsilon$, then on the one hand,

$$|(\mathbb{1}_{[a, b]} - \mathbb{1}_{[a, b]} * J_\epsilon)(x)| \leq \frac{1}{2}(\mathbb{1}_{[a-\epsilon, a+\epsilon]} + \mathbb{1}_{[b-\epsilon, b+\epsilon]})(x)$$

On the other hand, if $b - a < 2\epsilon$ then by translation invariance it suffices to consider the symmetric expression

$$f(x) := \mathbb{1}_{[-d, d]} - \mathbb{1}_{[-d, d]} * J_\epsilon, \quad 2d = b - a, \quad d < \epsilon$$

For any $0 < d < \epsilon/2$ this function satisfies

$$\begin{aligned} |f| &\leq \frac{d}{\epsilon} \mathbb{1}_{(-\epsilon-d, \epsilon+d)} + (1 - 2d/\epsilon) \mathbb{1}_{[-d, d]} \\ &\leq \frac{d}{\epsilon} \mathbb{1}_{(-\epsilon-d, \epsilon-d]} + \frac{d}{\epsilon} \mathbb{1}_{(\epsilon-d, 2\epsilon-d)} + (1 - 2d/\epsilon) \mathbb{1}_{(-\epsilon, \epsilon)} \end{aligned}$$

whereas for $\epsilon/2 < d < \epsilon$ one has

$$|f| \leq \frac{1}{2} \mathbb{1}_{(-\epsilon-d, \epsilon+d)} \leq \frac{1}{2} \mathbb{1}_{(-\epsilon-d, \epsilon-d]} + \frac{1}{2} \mathbb{1}_{(\epsilon-d, 3\epsilon-d)}$$

In either case (3.25) holds.

It therefore suffices to estimate $\sup_{\theta \in \mathbb{T}} \nu([\theta - \epsilon, \theta + \epsilon])$. Next, we define an atom τ' in the Hardy space $H^1(\mathbb{T})$ as follows:

$$(3.26) \quad \tau'(x) = \begin{cases} (x - (a - 3\epsilon))/\epsilon^2, & a - 3\epsilon \leq x \leq a - 2\epsilon, \\ ((a - \epsilon) - x)/\epsilon^2, & a - 2\epsilon \leq x \leq a - \epsilon, \\ ((a + \epsilon) - x)/\epsilon^2, & a + \epsilon \leq x \leq a + 2\epsilon, \\ (x - (a + 3\epsilon))/\epsilon^2, & a + 2\epsilon \leq x \leq a + 3\epsilon, \\ 0, & \text{otherwise.} \end{cases}$$

Note that this is well-defined on the circle since $\epsilon < \frac{1}{6}$. By construction, $\langle \tau' \rangle = 0$. Set $\tau(x) := \int_{a-\frac{1}{2}}^x \tau'(t) dt$. Moreover, $\tau \geq 0$, $\langle \tau \rangle = 4\epsilon$, and $\tau(x) = 1$ on $[a - \epsilon, a + \epsilon]$.

Thus

$$(3.27) \quad \nu([a - \epsilon, a + \epsilon]) \leq \int_{\mathbb{T}} \tau(x) \nu(dx) = \langle \tau, \nu \rangle.$$

Let us consider

$$(3.28) \quad \begin{aligned} & |(\tau - \langle \tau \rangle, \nu)| \\ &= |(\frac{d}{dx} \mathcal{H}[\tau], D^{-1} \mathcal{H}[\nu])| = |(\mathcal{H}[\tau'], u)| \\ &= |(\mathcal{H}[\tau'], v - \tilde{h})| = |(\mathcal{H}[\tau'], v_0 + v_1 - \tilde{h})| \\ &\leq |(\mathcal{H}[\tau'], v_0)| + |(\mathcal{H}[\tau'], v_1)| + |(\tau, \mathcal{H}[\tilde{h}'])| \\ &\leq \|\mathcal{H}[\tau']\|_{L^1(\mathbb{T})} \|v_0\|_{L^\infty(\mathbb{T})} + \|\mathcal{H}[\tau']\|_{L^\infty(\mathbb{T})} \|v_1\|_{L^1(\mathbb{T})} + \|\tau\|_{L^1(\mathbb{T})} \|\mathcal{H}[\tilde{h}']\|_{L^\infty(\mathbb{T})} \\ &\leq \varepsilon_0 \|\mathcal{H}[\tau']\|_{L^1(\mathbb{T})} + \varepsilon_1 \|\mathcal{H}[\tau']\|_{L^\infty(\mathbb{T})} + 2\epsilon \|\frac{d^2}{dx^2} \tilde{h}\|_{L^\infty(\mathbb{T})} \end{aligned}$$

In the last line, we used the following lemma on the third term.

Lemma 3.7. *For any $f \in C^1(\mathbb{T})$ one has $\|\mathcal{H}[f]\|_\infty \leq \frac{1}{2} \|f'\|_\infty$.*

Proof. Since $\sin(\pi x) \geq 2x$ for all $0 < x \leq \frac{1}{2}$, one has

$$\|\mathcal{H}[f]\|_\infty = \sup_{y \in \mathbb{T}} \left| \int_{\mathbb{T}} \frac{f(x) - f(y)}{\sin(\pi(x-y))} \cos(\pi(x-y)) dx \right| \leq \frac{1}{2} \|f'\|_\infty$$

as claimed. \square

In order to bound the the other terms in the last line of (3.28) we prove two lemmas.

Lemma 3.8. *Let τ' be defined by (3.26) and assume that $0 < \epsilon \leq \frac{1}{17}$. Then we have*

$$(3.29) \quad \|\mathcal{H}[\tau']\|_{L^1(\mathbb{T})} \leq \frac{9}{2}.$$

Remark 3.9. The upper bound $\frac{1}{17}$ is a particular choice which we have found to be convenient in the last section of the paper. A more restrictive assumption on ϵ will slightly improve the bound; e.g., assuming $\epsilon < \frac{1}{36}$ yields the value 4.2 for $\frac{9}{2}$. Such improvements are mainly due to the lower bound in (3.40) approaching π as $\epsilon \rightarrow 0$.

Proof. By translation symmetry, we may assume $a = 0$. We let $0 < \epsilon \leq \frac{1}{17}$ and $I = [-r\epsilon, r\epsilon]$ with $r = 3.6 > 3$. Notice that $r\epsilon < \frac{1}{2}$. We decompose $\|\mathcal{H}[\tau']\|_{L^1(\mathbb{T})}$ into the two parts

$$\|\mathcal{H}[\tau']\|_{L^1(\mathbb{T})} = \|\mathcal{H}[\tau']\|_{L^1(I)} + \|\mathcal{H}[\tau']\|_{L^1(I^c)}.$$

By Cauchy-Schwarz and the fact that the Hilbert transform is an isometry on $L^2(\mathbb{T})$, we have

$$\|\mathcal{H}[\tau']\|_{L^1(I)} \leq \sqrt{|I|} \|\mathcal{H}[\tau']\|_{L^2(I)} \leq \sqrt{|I|} \|\tau'\|_{L^2(I)} = \frac{2\sqrt{2}}{\sqrt{3}} \sqrt{r}.$$

In the remainder of the proof, we bound $\|\mathcal{H}[\tau']\|_{L^1(I^\epsilon)}$. By symmetry, we have $\|\mathcal{H}[\tau']\|_{L^1(I^\epsilon)} = 2\|\mathcal{H}[\tau']\|_{L^1([r\epsilon, \frac{1}{2}])}$. Hence, it suffices to consider the interval $[r\epsilon, \frac{1}{2}]$. For all $x \in [r\epsilon, 1/2]$, we have

$$\begin{aligned}
|\mathcal{H}[\tau'](x)| &= \left| \int_{\text{supp } \tau'} \tau'(y) \cot(\pi(x-y)) dy \right| \\
&= \left| \int_{\text{supp } \tau'} \tau'(y) (\cot(\pi(x-y)) - \cot(\pi x)) dy \right| \\
(3.30) \quad &= \left| \int_{\text{supp } \tau'} \tau'(y) \frac{\sin(\pi y)}{\sin(\pi(x-y)) \sin(\pi x)} dy \right| \\
&\leq \int_{\text{supp } \tau'} |\tau'(y)| \frac{|\sin(\pi y)|}{\sin(\pi(x-y)) \sin(\pi x)} dy.
\end{aligned}$$

Here we used that $x-y \in [(r-3)\epsilon, \frac{1}{2} + 3\epsilon] \subset [0, 1]$. To estimate this expression further, we decompose $\text{supp } \tau'$ into the four intervals

$$I_1 := [2\epsilon, 3\epsilon], \quad I_2 := [\epsilon, 2\epsilon], \quad I_3 := [-2\epsilon, -\epsilon], \quad I_4 := [-3\epsilon, -2\epsilon].$$

We write $\rho_j(x) \in I_j$ for the point in I_j that is nearest to x in the toroidal distance, i.e., $\|x - \rho_j(x)\|_{\mathbb{T}} = \text{dist}_{\mathbb{T}}(x, I_j)$. (That point is not unique if x is ‘‘antipodal’’ to the center of I_j ; in this case we define ρ_j as the right endpoint of I_j for definiteness.) Notice that $\rho_j(x)$ is constant for $j \in \{1, 2\}$ with

$$(3.31) \quad \rho_1(x) = \rho_{1,1} := 3\epsilon, \quad \rho_2(x) = \rho_{2,1} := 2\epsilon,$$

and piecewise constant for $j \in \{3, 4\}$, i.e.,

$$\rho_j(x) = \begin{cases} \rho_{j,1}, & \text{if } x \in [0, t_j], \\ \rho_{j,2}, & \text{if } x \in (t_j, \frac{1}{2}]. \end{cases}$$

Specifically, we have

$$\begin{aligned}
(3.32) \quad \rho_3(x) &= \begin{cases} -\epsilon, & \text{if } x \in [0, \frac{1}{2} - \frac{3\epsilon}{2}], \\ -2\epsilon, & \text{if } x \in (\frac{1}{2} - \frac{3\epsilon}{2}, \frac{1}{2}], \end{cases} \\
\rho_4(x) &= \begin{cases} -2\epsilon, & \text{if } x \in [0, \frac{1}{2} - \frac{5\epsilon}{2}], \\ -3\epsilon, & \text{if } x \in (\frac{1}{2} - \frac{5\epsilon}{2}, \frac{1}{2}], \end{cases}
\end{aligned}$$

From (3.30) and $|\sin(\pi y)| \leq \pi|y|$, we obtain

$$|\mathcal{H}[\tau'](x)| \leq \pi \sum_{j=1}^4 \frac{1}{\sin(\pi(x - \rho_j(x))) \sin(\pi x)} \int_{I_j} |\tau'(y)| |y| dy.$$

We integrate both sides over $x \in [r\epsilon, \frac{1}{2}]$ and find

$$(3.33) \quad \|\mathcal{H}[\tau']\|_{L^1([r\epsilon, \frac{1}{2}])} \leq \pi \sum_{j=1}^4 f_j(\epsilon) \int_{I_j} |\tau'(y)| |y| dy,$$

where we introduced the notation

$$f_j(\epsilon) := \int_{r\epsilon}^{r\frac{1}{2}} \frac{1}{\sin(\pi(x - \rho_j(x))) \sin(\pi x)} dx,$$

for $j \in \{1, 2, 3, 4\}$. The following lemma gives a bound on these integrals.

Lemma 3.10. *For $j \in \{1, 2\}$, we have*

$$(3.34) \quad f_j(\epsilon) \leq \frac{1}{2.98\pi\rho_{j,1}} \log \left(1 + \frac{\pi}{2.98} \frac{\rho_{j,1}}{r\epsilon - \rho_{j,1}} \right).$$

and for $j \in \{3, 4\}$, we have

$$(3.35) \quad f_j(\epsilon) \leq \frac{2}{5\pi|\rho_{j,1}|} \log \left(1 + \frac{2\pi}{5} \frac{|\rho_{j,1}|}{r\epsilon} \right) + 0.21.$$

We postpone the proof of Lemma 3.10 for now. To continue the proof of Lemma 3.8, recall (3.33). We perform the integration in y and find

$$(3.36) \quad \begin{aligned} \int_{I_1} |\tau'(y)||y|dy &= \int_{I_4} |\tau'(y)||y|dy = \frac{7\epsilon}{6}, \\ \int_{I_2} |\tau'(y)||y|dy &= \int_{I_3} |\tau'(y)||y|dy = \frac{5\epsilon}{6}. \end{aligned}$$

Then we apply Lemma 3.10 and recall Definitions (3.31) and (3.32) of $\rho_{j,1}$. This gives

$$(3.37) \quad \begin{aligned} \|\mathcal{H}[\tau']\|_{L^1([r\epsilon, \frac{1}{2}])} &\leq \pi \sum_{j=1}^4 f_j(\epsilon) |\tau'(y)||y|dy \\ &\leq \frac{1}{17.88} \left(\frac{7}{3} \log \left(1 + \frac{\pi}{2.98} \frac{3}{r-3} \right) + \frac{5}{2} \log \left(1 + \frac{\pi}{2.98} \frac{2}{r-2} \right) \right) \\ &\quad + \frac{1}{15} \left(5 \log \left(1 + \frac{2\pi}{5} \frac{1}{r} \right) + \frac{7}{2} \log \left(1 + \frac{2\pi}{5} \frac{2}{r} \right) \right) + (0.42)\pi\epsilon. \end{aligned}$$

Notice that $(0.42)\pi\epsilon < 0.08$. We write $\nu(r)$ for the expression in the last line. Altogether, we have shown that

$$\|\mathcal{H}[\tau']\|_{L^1(\mathbb{T})} \leq \frac{2\sqrt{2}}{\sqrt{3}}\sqrt{r} + 2\nu(r) < 4.5.$$

In the second step, we evaluated the expression at $r = 3.6$. This proves the main claim of Lemma 3.8. It remains to give the

Proof of Lemma 3.10. We begin by observing that

$$(3.38) \quad \frac{d}{dx} \left(\frac{1}{\sin x'} \log \left(\frac{\sin(x-x')}{\sin(x)} \right) \right) = \frac{1}{\sin x \sin(x-x')},$$

whenever x, x' are such that the logarithm is well-defined.

Let $j \in \{1, 2\}$, which implies that $\rho_j(x) = \rho_{j,1} > 0$. By (3.38) and $\sin, \cos \leq 1$, we have

$$\begin{aligned}
 (3.39) \quad f_j(\epsilon) &= \frac{1}{\pi \sin(\pi \rho_{j,1})} \left[\log \left(\frac{\sin(\pi(x - \rho_{j,1}))}{\sin(\pi x)} \right) \right]_{x=r\epsilon}^{x=\frac{1}{2}} \\
 &\leq \frac{1}{\pi \sin(\pi \rho_{j,1})} \log \left(\frac{\sin(\pi r\epsilon)}{\sin(\pi(r\epsilon - \rho_{j,1}))} \right) \\
 &\leq \frac{1}{\pi \sin(\pi \rho_{j,1})} \log \left(1 + \frac{\sin(\pi \rho_{j,1})}{\sin(\pi(r\epsilon - \rho_{j,1}))} \right)
 \end{aligned}$$

Next, we estimate the sin's by linear functions. While the upper bound $\sin(\pi x) \leq \pi x$ is valid for all x (and is sharp for small x), a linear lower bound on $\sin(\pi x)$ depends directly on the allowed range of x values. This is conveniently expressed via the quotient

$$\inf_{x \in [0, 3\epsilon]} \frac{\sin(\pi x)}{x} = \frac{\sin(3\pi\epsilon)}{3\epsilon} > 2.98$$

In the last step, we used that $\epsilon < \frac{1}{17}$. We may verify that all the arguments of $\sin(\pi \cdot)$ in the last line of (3.39) are located in the interval $[0, 3\epsilon]$. Therefore

$$\begin{aligned}
 f_j(\epsilon) &\leq \frac{1}{\pi \sin(\pi \rho_{j,1})} \log \left(1 + \frac{\sin(\pi \rho_{j,1})}{\sin(\pi(r\epsilon - \rho_{j,1}))} \right) \\
 &\leq \frac{1}{2.98\pi \rho_{j,1}} \log \left(1 + \frac{\pi}{2.98} \frac{\rho_{j,1}}{r\epsilon - \rho_{j,1}} \right).
 \end{aligned}$$

This proves (3.34).

Next, let $j \in \{3, 4\}$, so that $\rho_{j,1}, \rho_{j,2} < 0$. We have $r\epsilon < t_j$ by our assumptions on r, ϵ and (3.38) yields

$$\begin{aligned}
 f_j(\epsilon) &\leq \frac{1}{\pi \sin(\pi \rho_{j,1})} \left[\log \left(\frac{\sin(\pi(x - \rho_{j,1}))}{\sin(\pi x)} \right) \right]_{x=r\epsilon}^{x=t_j} \\
 &\quad + \left(\frac{1}{2} - t_j \right) \max_{x \in (t_j, \frac{1}{2}]} \frac{1}{\sin(\pi(x - \rho_{j,2})) \sin(\pi x)}.
 \end{aligned}$$

The second term is an error term (it vanishes as $\epsilon \rightarrow 0$). Indeed, recalling the definition of t_j and $\rho_{j,2}$ from (3.32), we see that for $j \in \{3, 4\}$,

$$\left(\frac{1}{2} - t_j \right) \frac{1}{\sin(\pi(\frac{1}{2} - \rho_{j,2})) \sin(\pi t_j)} \leq \frac{5\epsilon}{2 \cos^2(3\pi\epsilon)} \leq \frac{5\epsilon}{2(1 - \frac{9}{2}\pi^2\epsilon^2)^2} \leq 0.21,$$

where the last estimate used that $\epsilon \leq \frac{1}{17}$. Therefore, we have

$$\begin{aligned}
& \int_{r\epsilon}^{\frac{1}{2}} \frac{1}{\sin(\pi(x - \rho_j(x))) \sin(\pi x)} dx \\
& \leq \frac{1}{\pi \sin(\pi \rho_{j,1})} \left[\log \left(\frac{\sin(\pi(x - \rho_{j,1}))}{\sin(\pi x)} \right) \right]_{x=r\epsilon}^{x=t_j} + 0.21 \\
& \leq \frac{1}{\pi \sin(\pi |\rho_{j,1}|)} \log \left(\frac{\sin(\pi(r\epsilon - \rho_{j,1}))}{\sin(\pi r\epsilon)} \right) + 0.21 \\
& \leq \frac{2}{5\pi |\rho_{j,1}|} \log \left(1 + \frac{2\pi}{5} \frac{|\rho_{j,1}|}{r\epsilon} \right) + 0.21.
\end{aligned}$$

In the second step, we used that $0 \leq t_j < t_j - \rho_{j,1} \leq \frac{1}{2}$ and monotonicity properties of \sin . In the last step, we used $\cos \leq 1$ and

$$(3.40) \quad \inf_{x \in [0, (r+2)\epsilon]} \frac{\sin(\pi x)}{x} = \frac{\sin(\pi(r+2)\epsilon)}{(r+2)\epsilon} > \frac{5}{2}.$$

This bound holds because $(r+2)\epsilon < \frac{1}{3}$, which may be verified from $r = 3.6$ and $\epsilon \leq \frac{1}{17}$. This shows (3.35) and concludes the proof of Lemma 3.10, and hence also of Lemma 3.8. \square

Next, we control the pointwise size of $\mathcal{H}[\tau']$.

Lemma 3.11. *We have*

$$\|\mathcal{H}[\tau']\|_{L^\infty(\mathbb{T})} \leq \frac{5}{2\epsilon}$$

Proof. By translation invariance, we can set $a = 0$. Let us first consider

$$\int_{\text{supp}(\tau')} \tau'(x) \cot \pi(x - y) dy.$$

If $x \notin \text{supp}(\tau')$, then $\int_{\text{supp}(\tau')} \tau'(x) \cot \pi(x - y) dy = 0$. Thus, we can assume without loss of generality that $x \in [-3\epsilon, -\epsilon]$. On the one hand,

$$\begin{aligned}
\left| \int_{[\epsilon, 3\epsilon]} \tau'(x) \cot \pi(x - y) dy \right| & \leq 2\epsilon |\tau'(x)| \sup_{y \in [\epsilon, 3\epsilon]} |\cot \pi(x - y)| \\
& \leq 2 \sup_{u \in [2\epsilon, 6\epsilon]} |\cot(\pi u)| \leq 2 \cot(2\pi\epsilon) \leq \frac{1}{2\epsilon},
\end{aligned}$$

since $6\epsilon\pi \leq \pi - 2\pi\epsilon$ and $\sin(2\pi\epsilon) \geq 4\epsilon$ (recall that $\epsilon < \frac{1}{8}$). On the other hand, for the negative support of τ' , we can further assume by symmetry that $x \in [-3\epsilon, -2\epsilon]$.

Thus,

$$\begin{aligned}
& \left| \int_{[-3\epsilon, -\epsilon]} \tau'(x) \cot \pi(x-y) dy \right| \\
&= \left| \tau'(x) \int_{2x+3\epsilon}^{-\epsilon} \cot \pi(y-x) dy \right| = \left| \tau'(x) \int_{x+3\epsilon}^{-\epsilon-x} \cot(\pi y) dy \right| \\
(3.41) \quad &= \frac{1}{\epsilon^2} (x+3\epsilon) \int_{x+3\epsilon}^{-\epsilon-x} \frac{1}{2y} dy = \frac{1}{\epsilon} \frac{x+3\epsilon}{2\epsilon} \log \left(\frac{2\epsilon}{x+3\epsilon} - 1 \right) \\
&\leq \frac{1}{\epsilon} \sup_{t \in [0, 1/2]} t \log \left(\frac{1}{t} - 1 \right) < \frac{1}{4\epsilon}.
\end{aligned}$$

Hence, overall we have

$$(3.42) \quad \left| \int_{\text{supp}(\tau')} \tau'(x) \cot \pi(x-y) dy \right| < \frac{3}{4\epsilon}.$$

Now let us consider with $x \in [-3\epsilon, -\epsilon]$,

$$\begin{aligned}
& \left| \mathcal{H}[\tau'](x) - \int_{\text{supp}(\tau')} \tau'(x) \cot \pi(x-y) dy \right| \\
&= \left| \int_{\text{supp}(\tau')} (\tau'(y) - \tau'(x)) \cot \pi(x-y) dy \right| \\
(3.43) \quad &\leq \int_{-3\epsilon}^{-\epsilon} \left| \frac{\tau'(y) - \tau'(x)}{\sin \pi(y-x)} \right| dy + \int_{\epsilon}^{3\epsilon} \frac{|\tau'(y)| + |\tau'(x)|}{|\sin \pi(y-x)|} dy \\
&\leq \int_{-3\epsilon}^{-\epsilon} \frac{|x-y|\epsilon^{-2}}{2|x-y|} dy + \left(\frac{1}{4} + \frac{1}{2} \log 2 \right) \epsilon^{-1} \leq \frac{1.6}{\epsilon}.
\end{aligned}$$

In summary,

$$(3.44) \quad \|\mathcal{H}[\tau']\|_{L^\infty(\mathbb{T})} < \frac{5}{2\epsilon}$$

as claimed. \square

Combining (3.20), (3.27), (3.28), with Lemmas 2.5, 3.8 and 3.11, we have

$$\begin{aligned}
(\tau, \nu) &\leq \frac{9}{2}\epsilon_0 + 5\frac{\epsilon_1}{2\epsilon} + 2\epsilon B_2(B-m) + \langle \tau \rangle \nu(\mathbb{T}) \\
&\leq \frac{9}{2}\epsilon_0 + 5\frac{\epsilon_1}{2\epsilon} + \epsilon \left(2B_2 + \frac{4\pi}{\log R/R_1} \right) (B-m) \\
&= \frac{9}{2}\epsilon_0 + 2B_3 \sqrt{\epsilon_1(B-m)},
\end{aligned}$$

Note that our choice (3.21) of ϵ minimizes the contribution of ϵ_1 . Finally, (3.25) concludes the proof of Lemma 3.6. \square

In order to prove the exponential integrability of $v-c$, and thus complete the proof of Lemma 3.1, we invoke the following classical result about the Hilbert transform of bounded functions on the circle, see for example [Kat].

Lemma 3.12. *Let f be a real-valued function on \mathbb{T} such that $|f| \leq 1$. Then for any $0 \leq \alpha < \frac{1}{2}\pi$,*

$$\int_{\mathbb{T}} e^{\alpha|\mathcal{H}[f](x)|} dx \leq \frac{2}{\cos \alpha} = 2 \sec \alpha.$$

Applying Lemma 3.12 to $f = (D^{-1}(\nu - J_\epsilon * \nu))/\|(D^{-1}(\nu - J_\epsilon * \nu))\|_{L^\infty(\mathbb{T})}$, by (3.23), we have

$$(3.45) \quad \int_{\mathbb{T}} \exp(\beta|(u - J_\epsilon * u)(x)|) dx \leq 2 \sec \alpha,$$

where $\alpha = \beta(\frac{9}{2}\epsilon_0 + 2B_3\sqrt{\epsilon_1(B-m)}) < \frac{\pi}{2}$. Taking $\alpha = \frac{\pi}{4}$ in (3.45), then (3.22) yields (absorbing the constant c into v for simplicity)

$$(3.46) \quad \begin{aligned} & \int_{\mathbb{T}} \exp\left(\frac{\pi|v(x)|}{18\epsilon_0 + 8B_3\sqrt{\epsilon_1(B-m)}}\right) dx \\ & \leq 2\sqrt{2} \exp\left(\frac{\pi\epsilon_0 + \pi(B_3 + B_1/B_3)\sqrt{\epsilon_1(B-m)}/2}{18\epsilon_0 + 8B_3\sqrt{\epsilon_1(B-m)}}\right) \\ & \leq 2\sqrt{2} \exp\left(\pi\left[\frac{17}{144} + \frac{B_1}{16B_3^2}\right]\right), \end{aligned}$$

which concludes the proof of Lemma 3.1. \square

Decay of the Fourier coefficients. We conclude this section with an important decay estimate on the Fourier coefficients of the subharmonic function v . This lemma will be used in Sec.7.

Lemma 3.13. *Let v be as in Theorem 2.2, then the Fourier coefficients of v satisfy*

$$|\hat{v}(k)| \leq \frac{C(R, R_1, R_2)}{|k|}(B-m), \quad \text{for any } k \neq 0,$$

in which

$$(3.47) \quad C(R, R_1, R_2) = \frac{1}{2 \log R/R_1} + \frac{1}{2\pi} B_1(R, R_1, R_2).$$

Proof. For any $k \neq 0$, we have

$$(3.48) \quad |\hat{v}(k)| = \left| \int_{\mathbb{T}} (u(x) + \tilde{h}(x)) e^{-2\pi i k x} dx \right| \leq \frac{1}{2\pi|k|} \left(|\hat{u}'(k)| + |\hat{\tilde{h}}'(k)| \right).$$

By Lemma 3.3, we have $|\tilde{h}'(x)| \leq B_1(R, R_1, R_2)(B-m)$, hence

$$(3.49) \quad |\hat{\tilde{h}}'(k)| \leq B_1(R, R_1, R_2)(B-m).$$

By (3.19), (3.20) and Theorem 2.2, we have

$$(3.50) \quad |\hat{u}'(k)| = |\widehat{\mathcal{H}[\nu]}(k)| = |\hat{v}(k)| \leq \pi \tilde{\mu}(\overline{D_1}) \leq \frac{\pi(B-m)}{\log R/R_1}.$$

In view of (3.48), (3.49) and (3.50) we infer that

$$|\hat{v}(k)| \leq \frac{1}{2\pi|k|} \left(B_1(R, R_1, R_2)(B - m) + \frac{\pi(B - m)}{\log R/R_1} \right).$$

as claimed. \square

4. \mathbb{T}^2 SPLITTING LEMMA

Our applications to the skew-shift dynamics on \mathbb{T}^2 require a version of the splitting lemma in two variables. First, we formalize the class of plurisubharmonic functions that we will be working with.

Definition 4.1. Let $v(z, w)$ be a continuous pluri-subharmonic function on $D_R \times D_R$, satisfying the following estimates for $R > 1$,

$$(4.1) \quad \begin{cases} v(z, w) \leq B_4(R) \text{ for } \forall (z, w) \in D_R \times \partial D_1, \text{ and } v(0, e(y)) \geq m_4 \text{ for } \forall y \in \mathbb{T}, \\ v(z, w) \leq B_5(R) \text{ for } \forall (z, w) \in \partial D_1 \times D_R, \text{ and } v(e(x), 0) \geq m_5 \text{ for } \forall x \in \mathbb{T}, \\ |v(e(x), e(y))| \leq B_6 \text{ for } \forall (x, y) \in \mathbb{T}^2 \end{cases}$$

For a function f defined on a polydisk in \mathbb{C}^2 which contains \mathbb{T}^2 , let us denote $f(e(x), e(y))$ by $f(x, y)$ for simplicity. In particular, we will write $v(x, y)$ on \mathbb{T}^2 . The average is denoted by $\langle f \rangle_{\mathbb{T}^2} := \int_{\mathbb{T}^2} f(x, y) dx dy$.

Below, we will analyze a particular Schrödinger cocycle over a skew shift base. At that opine, we will specify the constants in Definitions 2.1 and 4.1. But for now we develop more analytical machinery with these constants as parameters. Recall that for a Borel set U , the Lebesgue measure will be written as $|U|$.

Lemma 4.2. *Let v be as in Definition 4.1, and assume (3.14). Let $0 < r < 1$, $0 < \varepsilon_1$ and $0 < \varepsilon_0 < \varepsilon_3 < \varepsilon_2$. Assume*

$$v(x, y) = v_0(x, y) + v_1(x, y) + \langle v \rangle_{\mathbb{T}^2},$$

where $\|v_0\|_{L^\infty(\mathbb{T}^2)} < \varepsilon_0$, and $\|v_1\|_{L^1(\mathbb{T}^2)} < \varepsilon_1$. Then

$$\begin{aligned} |\{(x, y) \in \mathbb{T}^2 : |v(x, y) - \langle v \rangle_{\mathbb{T}^2}| > \varepsilon_2\}| &< 2(2C_0)^{\frac{1}{2}} \exp\left(-\frac{\pi}{8\delta_0^{(1)}}\varepsilon_3\right) \\ &+ C_0 \exp\left(-\frac{\pi}{4\delta_0^{(2)}}\varepsilon_2\right) \end{aligned}$$

in which $\frac{9}{2}$ and $B_3 = B_3(R, R_1, R_2)$ are defined as in (2.2), C_0 in (3.2), and

$$(4.2) \quad \begin{aligned} \delta_0^{(1)} &:= \frac{9}{2}\varepsilon_0 + 2B_3\sqrt{\varepsilon_1^r(B_4 - m_4)} \\ \delta_0^{(2)} &:= \frac{9}{2}\varepsilon_3 + 4B_3\sqrt{B_6}\sqrt{\varepsilon_1^{1-r}(B_5 - m_5)} \end{aligned}$$

Proof. Fix $0 < r < 1$. Let

$$(4.3) \quad A_1 := \left\{ y \in \mathbb{T} : \int_{\mathbb{T}} |v_1(x, y)| \, dx < \varepsilon_1^r \right\}.$$

By Markov's inequality, we have

$$(4.4) \quad |A_1^c| < \varepsilon_1^{1-r}.$$

For any fixed $y \in A_1$, we have $\|v_0(\cdot, y)\|_{L^\infty(\mathbb{T})} < \varepsilon_0$ and $\|v_1(\cdot, y)\|_{L^1(\mathbb{T})} < \varepsilon_1^r$. Applying Lemma 3.1 in the x variable, we then have

$$\int_{\mathbb{T}} \exp\left(\frac{\pi}{4\delta_0^{(1)}} |v(x, y) - \langle v \rangle_{\mathbb{T}^2}| \right) dx \leq C_0$$

Integrating over $y \in A_1$, and interchanging the integrations, yields

$$(4.5) \quad \int_{\mathbb{T}} \int_{A_1} \exp\left(\frac{\pi}{4\delta_0^{(1)}} |v(x, y) - \langle v \rangle_{\mathbb{T}^2}| \right) dy dx \leq C_0$$

For $\gamma > 0$, let us define

$$(4.6) \quad A_2 := \left\{ x \in \mathbb{T} : \int_{A_1} \exp\left(\frac{\pi}{4\delta_0^{(1)}} |v(x, y) - \langle v \rangle_{\mathbb{T}^2}| \right) dy \leq C_0 \gamma^{-1} \right\}.$$

By Markov's inequality,

$$(4.7) \quad |A_2^c| < \gamma.$$

For $x \in A_2$ and $\varepsilon_3 > \varepsilon_0$, let us define

$$(4.8) \quad A_3 := \{y \in A_1 : |v(x, y) - \langle v \rangle_{\mathbb{T}^2}| < \varepsilon_3\},$$

Again, by Markov's inequality,

$$(4.9) \quad |A_3^c| < C_0 \gamma^{-1} \exp\left(-\frac{\pi\varepsilon_3}{4\delta_0^{(1)}}\right)$$

Thus for $x \in A_2$, $\{y \in \mathbb{T} : |v(x, y) - \langle v \rangle| > \varepsilon_3\} \subseteq A_1^c \cup A_3^c$, with the following measure estimate

$$(4.10) \quad |\{y \in \mathbb{T} : |v(x, y) - \langle v \rangle_{\mathbb{T}^2}| > \varepsilon_3\}| \leq \varepsilon_1^{1-r} + C_0 \gamma^{-1} \exp\left(-\frac{\pi\varepsilon_3}{4\delta_0^{(1)}}\right).$$

Here we divide into two different cases, depending on which term on the right-hand side dominates.

Case 1: $\varepsilon_1^{1-r} < C_0 \gamma^{-1} \exp\left(-\frac{\pi\varepsilon_3}{4\delta_0^{(1)}}\right)$. Then (4.10) directly implies that for any $x \in A_2$,

$$(4.11) \quad |\{y \in \mathbb{T} : |v(x, y) - \langle v \rangle_{\mathbb{T}^2}| > \varepsilon_3\}| \leq 2C_0 \gamma^{-1} \exp\left(-\frac{\pi\varepsilon_3}{4\delta_0^{(1)}}\right)$$

Together with (4.7), we conclude that

$$(4.12) \quad |\{(x, y) \in \mathbb{T}^2 : |v(x, y) - \langle v \rangle_{\mathbb{T}^2}| > \varepsilon_3\}| \leq \gamma + 2C_0 \gamma^{-1} \exp\left(-\frac{\pi\varepsilon_3}{4\delta_0^{(1)}}\right)$$

Case 2: $\varepsilon_1^{1-r} \geq C_0 \gamma^{-1} \exp\left(-\frac{\pi\varepsilon_3}{4\delta_0^{(1)}}\right)$. Then for any $x \in A_2$,

$$(4.13) \quad |\{y \in \mathbb{T} : |v(x, y) - \langle v \rangle_{\mathbb{T}^2}| > \varepsilon_3\}| \leq 2\varepsilon_1^{1-r}.$$

For $x \in A_2$, let

$$(4.14) \quad \begin{aligned} \tilde{v}_{x,0}(y) &= (v(x, y) - \langle v \rangle_{\mathbb{T}^2}) \mathbf{1}_{\{y \in \mathbb{T} : |v(x, y) - \langle v \rangle_{\mathbb{T}^2}| \leq \varepsilon_3\}} \\ \tilde{v}_{x,1}(y) &= (v(x, y) - \langle v \rangle_{\mathbb{T}^2}) \mathbf{1}_{\{y \in \mathbb{T} : |v(x, y) - \langle v \rangle_{\mathbb{T}^2}| > \varepsilon_3\}}. \end{aligned}$$

Then (4.13) implies, assuming $x \in A_2$,

$$(4.15) \quad \begin{aligned} v(x, y) &= \tilde{v}_{x,0}(y) + \tilde{v}_{x,1}(y) + \langle v \rangle_{\mathbb{T}^2}, \\ \|\tilde{v}_{x,0}(\cdot)\|_{L^\infty(\mathbb{T})} &\leq \varepsilon_3, \\ \|\tilde{v}_{x,1}(\cdot)\|_{L^1(\mathbb{T})} &\leq 2\varepsilon_1^{1-r} \|v(x, \cdot) - \langle v \rangle_{\mathbb{T}^2}\|_{L^\infty(\mathbb{T})} \leq 4B_6 \varepsilon_1^{1-r}. \end{aligned}$$

Applying Corollary 3.2 in the y variable, we obtain that for any $x \in A_2$ and any $\varepsilon_2 > \varepsilon_3$,

$$(4.16) \quad |\{y \in \mathbb{T} : |v(x, y) - \langle v \rangle_{\mathbb{T}^2}| > \varepsilon_2\}| \leq C_0 \exp\left(-\frac{\pi\varepsilon_2}{4\delta_0^{(2)}}\right)$$

Together with (4.7), we then get

$$(4.17) \quad |\{(x, y) \in \mathbb{T}^2 : |v(x, y) - \langle v \rangle_{\mathbb{T}^2}| > \varepsilon_2\}| \leq \gamma + C_0 \exp\left(-\frac{\pi\varepsilon_2}{4\delta_0^{(2)}}\right).$$

Finally, we choose γ to equalize the terms in (4.12):

$$\gamma = (2C_0)^{\frac{1}{2}} \exp\left(-\frac{\pi\varepsilon_3}{8\delta_0^{(1)}}\right)$$

Then the estimate of Case 1, namely (4.12), yields

$$(4.18) \quad \begin{aligned} &|\{(x, y) \in \mathbb{T}^2 : |v(x, y) - \langle v \rangle_{\mathbb{T}^2}| > \varepsilon_2\}| \\ &\leq |\{(x, y) \in \mathbb{T}^2 : |v(x, y) - \langle v \rangle_{\mathbb{T}^2}| > \varepsilon_3\}| \\ &\leq 2(2C_0)^{\frac{1}{2}} \exp\left(-\frac{\pi\varepsilon_3}{8\delta_0^{(1)}}\right) \end{aligned}$$

The estimate of Case 2, namely (4.17), becomes

$$(4.19) \quad \begin{aligned} &|\{(x, y) \in \mathbb{T}^2 : |v(x, y) - \langle v \rangle_{\mathbb{T}^2}| > \varepsilon_2\}| \\ &\leq (2C_0)^{\frac{1}{2}} \exp\left(-\frac{\pi\varepsilon_3}{8\delta_0^{(1)}}\right) + C_0 \exp\left(-\frac{\pi\varepsilon_2}{4\delta_0^{(2)}}\right). \end{aligned}$$

Combining (4.18) with (4.19), we conclude that

$$\begin{aligned} &|\{(x, y) \in \mathbb{T}^2 : |v(x, y) - \langle v \rangle_{\mathbb{T}^2}| > \varepsilon_2\}| \\ &\leq 2(2C_0)^{\frac{1}{2}} \exp\left(-\frac{\pi\varepsilon_3}{8\delta_0^{(1)}}\right) + C_0 \exp\left(-\frac{\pi\varepsilon_2}{4\delta_0^{(2)}}\right). \end{aligned}$$

as claimed. \square

5. AVALANCHE PRINCIPLE

The Avalanche Principle (AP) is a device to compare the logarithm of the norm of a long product $A_n A_{n-1} \dots A_2 A_1$ of matrices to the sum of the logarithms of the norms of shorter sections of the product. In the original formulation from [GolSch] for $\mathrm{SL}_2(\mathbb{R})$ matrices the length of the chain was limited depending on the norms of the individual matrices A_j . The same restriction applied to the extension of the AP to $\mathrm{SL}_d(\mathbb{R})$ matrices in [Sch]. Later, Duarte and Klein [DuaKle] found a different proof of the AP which does not use impose any restriction on the length of the chain. Even though the older version [GolSch] would suffice for our purposes, we present the argument from [DuaKle] with explicit constants. (These are not provided in [DuaKle].)

Thus, this section is devoted to making the constants in Chapter 2 of [DuaKle2] effective (we mostly follow [DuaKle2] instead of [DuaKle] for the sake of simplicity). We use the same notation as [DuaKle2], which we first recall. Although we only need the results for $\mathrm{SL}_2(\mathbb{R})$ matrices in this paper, we aim at proving more general results which are of independent interest.

Let $\mathrm{GL}_d(\mathbb{R})$ be the general linear group of real $d \times d$ matrices.

Definition 5.1. Given matrices $g_0, g_1, \dots, g_{n-1} \in \mathrm{GL}_d(\mathbb{R})$, the expansion rift is the ratio

$$\rho(g_0, g_1, \dots, g_{n-1}) := \frac{\|g_{n-1} \cdots g_1 g_0\|}{\|g_{n-1}\| \cdots \|g_1\| \|g_0\|} \in (0, 1].$$

Given $g \in \mathrm{GL}_d(\mathbb{R})$, let

$$s_1(g) \geq s_2(g) \geq \dots \geq s_d(g) > 0$$

denote the sorted singular values of g . The first singular value $s_1(g)$ is the operator norm

$$s_1(g) = \max_{x \in \mathbb{R}^d \setminus \{0\}} \frac{\|gx\|}{\|x\|} := \|g\|.$$

The last singular value of g is the least expansion factor of f , regarded as a linear transformation, and it can be characterized by

$$s_d(g) = \min_{x \in \mathbb{R}^d \setminus \{0\}} \frac{\|gx\|}{\|x\|} = \|g^{-1}\|^{-1}.$$

Definition 5.2. The gap (or the singular gap) of $g \in \mathrm{GL}_d(\mathbb{R})$ is the ratio between its first and second singular values.

$$\mathrm{gr}(g) := \frac{s_1(g)}{s_2(g)}.$$

Remark 5.3. If $g \in \mathrm{SL}_2(\mathbb{R})$, then $\mathrm{gr}(g) = \|g\|^2$.

Let $\mathbb{P}(\mathbb{R}^d)$ denote the projective space. Points in $\mathbb{P}(\mathbb{R}^d)$ are equivalence classes \hat{x} of non-zero vectors $x \in \mathbb{R}^d$. We consider the projective distance $\delta : \mathbb{P}(\mathbb{R}^d) \times \mathbb{P}(\mathbb{R}^d) \rightarrow [0, 1]$,

$$\delta(\hat{x}, \hat{y}) := \sin(\angle(x, y)),$$

where \angle is the length of the arc connecting x and y .

Definition 5.4. Given $g \in \mathrm{GL}_d(\mathbb{R})$ such that $\mathrm{gr}(g) > 1$, the most expanding direction of g is the singular direction $\hat{\mathbf{v}} \in \mathbb{P}(\mathbb{R}^d)$ associated with the first singular value $s_1(g)$ of g . Let $\mathbf{v}(g)$ be any of the two unit vector representatives of the projective point $\hat{\mathbf{v}}(g)$. We set $\hat{\mathbf{v}}^*(g) := \hat{\mathbf{v}}(g^*)$ and $\mathbf{v}^*(g) := \mathbf{v}(g^*)$.

Any matrix $g \in \mathrm{GL}_d(\mathbb{R})$ maps the most expanding direction of g to the most expanding direction of g^* , multiplying vectors by the factor $s_1(g) = \|g\|$.

$$g\mathbf{v}(g) = \pm s_1(g)\mathbf{v}^*(g).$$

The matrix g also induces a projective map $\hat{g} : \mathbb{P}(\mathbb{R}^d) \rightarrow \mathbb{P}(\mathbb{R}^d)$, $\hat{g}(\hat{x}) := \hat{g}x$, for which one has

$$\hat{g}(\hat{\mathbf{v}}(g)) = \hat{\mathbf{v}}^*(g) \quad \text{and} \quad \hat{g}^*(\hat{\mathbf{v}}^*(g)) = \hat{\mathbf{v}}(g).$$

Theorem 5.5. Let $n \geq 1$ and $0 < \epsilon \leq \frac{1}{10}$. Given $0 < \kappa \leq \frac{1}{10}\epsilon^2$ and $g_0, g_1, \dots, g_{n-1} \in \mathrm{GL}_d(\mathbb{R})$, if

- (G) $\mathrm{gr}(g_i) \geq \kappa^{-1}$ for $j = 0, 1, \dots, n-1$,
- (A) $\rho(g_{j-1}, g_j) \geq \epsilon$ for $j = 1, 2, \dots, n-1$,

then, writing $g^j := g_{j-1} \cdots g_1 g_0$, we have

- (i) $\max \{ \delta(\hat{\mathbf{v}}(g^n), \hat{\mathbf{v}}(g_0)), \delta(\hat{\mathbf{v}}^*(g^n), \hat{\mathbf{v}}^*(g_{n-1})) \} \leq 3\kappa\epsilon^{-1}$,
- (ii) $e^{-5n\kappa/\epsilon^2} \leq \frac{\rho(g_0, g_1, \dots, g_{n-1})}{\rho(g_0, g_1) \cdots \rho(g_{n-2}, g_{n-1})} \leq e^{11n\kappa/\epsilon^2}$.

The proof follows the general line of argumentation in [DuaKle2], keeping track of the effective constants throughout.

Staging the proof. The projective distance $\delta : \mathbb{P}(\mathbb{R}^d) \times \mathbb{P}(\mathbb{R}^d) \rightarrow [0, 1]$ determines a complementary angle function $\alpha : \mathbb{P}(\mathbb{R}^d) \times \mathbb{P}(\mathbb{R}^d) \rightarrow [0, 1]$, defined by

$$\alpha(\hat{x}, \hat{y}) := |\cos(\angle(x, y))|.$$

Let us also introduce the algebraic operation

$$a \oplus b := a + b - ab.$$

For properties of $a \oplus b$, one may refer to Proposition 2.1 of [DuaKle2].

Lemma 5.6. Given $g \in \mathrm{GL}_d(\mathbb{R})$ with $\mathrm{gr}(g) > 1$, $\hat{x} \in \mathbb{P}(\mathbb{R}^d)$ and a unit vector $x \in \hat{x}$, writing $\alpha = \alpha(\hat{x}, \hat{\mathbf{v}}(g))$ we have

- (a) $\alpha \leq \frac{\|gx\|}{\|x\|} \leq \sqrt{\alpha^2 \oplus \mathrm{gr}(g)^{-2}}$,
- (b) $\delta(\hat{g}(\hat{x}), \hat{\mathbf{v}}^*(g)) \leq \alpha^{-1} \mathrm{gr}(g)^{-1} \delta(\hat{x}, \hat{\mathbf{v}}(g))$,
- (c) The restriction of the map $\hat{g} : \mathbb{P}(\mathbb{R}^d) \rightarrow \mathbb{P}(\mathbb{R}^d)$ to the disk $\{\hat{x} \in \mathbb{P}(\mathbb{R}^d) : \delta(\hat{x}, \hat{\mathbf{v}}(g)) \leq r\}$ has Lipschitz constant $\leq \frac{\pi}{2} \frac{r + \sqrt{1-r^2}}{\mathrm{gr}(g)(1-r^2)}$ with respect to the δ -metric.

Proof. The factor $\frac{\pi}{2}$ in the Lipschitz constant is already explicit in the proof of Lemma 2.2 of [DuaKle2]. \square

Corollary 5.7. Given $g \in \mathrm{GL}_d(\mathbb{R})$ such that $\mathrm{gr}(g) \geq \kappa^{-1}$, define

$$\Sigma_\epsilon := \{\hat{x} \in \mathbb{P}(\mathbb{R}^d) : \alpha(\hat{x}, \hat{\mathbf{v}}(g)) \geq \epsilon\} = B(\hat{\mathbf{v}}(g), \sqrt{1 - \epsilon^2}).$$

Given a point $\hat{x} \in \Sigma_\epsilon$,

- (a) $\delta(\hat{g}(\hat{x}), \hat{g}(\hat{\mathbf{v}}(g))) \leq \kappa \epsilon^{-1} \delta(\hat{x}, \hat{\mathbf{v}}(g))$,
- (b) The map $\hat{g}|_{\Sigma_\epsilon} \rightarrow \mathbb{P}(\mathbb{R}^d)$ has Lipschitz constant $\leq \frac{\sqrt{2}\pi}{2} \kappa \epsilon^{-2}$.

Proof. (a) follows directly from (b) of Lemma 5.6. (b) follows from (c) of Lemma 5.6 and the fact that $\epsilon + \sqrt{1 - \epsilon^2} \leq \sqrt{2}$. \square

Definition 5.8. Given $g, g' \in \mathrm{GL}_d(\mathbb{R})$ with $\mathrm{gr}(g), \mathrm{gr}(g') > 1$, we define their lower angle as

$$\alpha(g, g') := \alpha(\hat{\mathbf{v}}^*(g), \hat{\mathbf{v}}(g')).$$

The upper angle between g and g' is

$$\beta(g, g') := \sqrt{\mathrm{gr}(g)^{-2} \oplus \alpha(g, g')^2 \oplus \mathrm{gr}(g')^{-2}}.$$

Lemma 5.9. Given $g, g' \in \mathrm{GL}_d(\mathbb{R})$, if $\mathrm{gr}(g), \mathrm{gr}(g') > 1$, then

$$\alpha(g, g') \leq \rho(g, g') \leq \beta(g, g').$$

This lemma has the following immediate corollary. It shows how the assumptions (G) and (A) in Theorem 5.5 will be used.

Corollary 5.10. Given $g, g' \in \mathrm{GL}_d(\mathbb{R})$, if $\mathrm{gr}(g), \mathrm{gr}(g') \geq \kappa^{-1}$ and $\rho(g, g') \geq \epsilon$, then

$$\delta(\hat{\mathbf{v}}^*(g), \hat{\mathbf{v}}(g')) \leq \sqrt{1 - \frac{\epsilon^2}{1 + 2\frac{\kappa^2}{\epsilon^2}}}.$$

We recall that $g^j := g_{j-1} \cdots g_1 g_0$.

Lemma 5.11. If $\mathrm{gr}(g_j) > 1$ for $j = 0, 1, \dots, n-1$, and $\mathrm{gr}(g^j) > 1$ for $j = 1, 2, \dots, n$, then

$$\prod_{j=1}^{n-1} \alpha(g^j, g_j) \leq \rho(g_0, g_1, \dots, g_{n-1}) \leq \prod_{j=1}^{n-1} \beta(g^j, g_j).$$

Proof of Theorem 5.5. To simplify the notation, we will write $c_0 = \frac{1}{10}$, $\hat{\mathbf{v}}_j := \hat{\mathbf{v}}(g_j)$ and $\hat{\mathbf{v}}_j^* := \hat{\mathbf{v}}^*(g_j)$ for $j = 0, 1, \dots, n-1$. We also let

$$g_j = g_{2n-1-j}^*, \quad \hat{\mathbf{v}}_j = \hat{\mathbf{v}}_{2n-1-j}^*, \quad \text{and} \quad \hat{\mathbf{v}}_j^* = \hat{\mathbf{v}}_{2n-1-j} \quad \text{for } j = n, n+1, \dots, 2n-1.$$

For each $i = 0, 1, \dots, 2n-1$ and $j = 0, 1, \dots, 2n-i$, set

$$\hat{\mathbf{v}}_i^j := \hat{g}_{i+j-1} \cdots \hat{g}_{i+1} \hat{g}_i \hat{\mathbf{v}}_i.$$

In terms of the notation above, we have $(\widehat{g^n})^* \hat{\mathbf{v}}(g_0) = \hat{\mathbf{v}}_0^{2n}$ and $\hat{\mathbf{v}}_0 = \hat{\mathbf{v}}_{2n-1}^1$.

By Assumption (A), we have $\rho(g_{j-1}, g_j) \geq \epsilon$ for $1 \leq j \leq n-1$. Hence for $n+1 \leq j \leq 2n-1$,

$$(5.1) \quad \rho(g_{j-1}, g_j) = \rho(g_{2n-j}^*, g_{2n-j-1}^*) = \rho(g_{2n-j-1}, g_{2n-j}) \geq \epsilon.$$

Clearly, we also have

$$(5.2) \quad \rho(g_{n-1}, g_n) = \rho(g_{n-1}, g_{n-1}^*) = \frac{\|g_{n-1}^* g_{n-1}\|}{\|g_{n-1}\|^2} = 1.$$

Therefore combining Assumption (A) with (5.1) and (5.2), we have

$$(5.3) \quad \rho(g_{j-1}, g_j) \geq \epsilon \quad \text{for } j = 1, 2, \dots, 2n - 1.$$

We begin with the proof of statement (i). We will prove $\delta(\hat{\mathbf{v}}(g^n), \hat{\mathbf{v}}(g_0)) \leq 3\kappa\epsilon^{-1}$. The other bound can be proved in exactly the same way.

First, we will show that for $\tilde{\epsilon} = t\epsilon$, $t = 2/3$, we have

Lemma 5.12. *For any $1 \leq j \leq 2n - 1$,*

$$\hat{g}_{j-1}(B(\hat{\mathbf{v}}_{j-1}, \sqrt{1 - \tilde{\epsilon}^2})) \subseteq B(\hat{\mathbf{v}}_j, \sqrt{1 - \tilde{\epsilon}^2}).$$

Proof. Taking any $\hat{x} \in B(\hat{\mathbf{v}}_{j-1}, \sqrt{1 - \tilde{\epsilon}^2})$, we have

$$\delta(\hat{x}, \hat{\mathbf{v}}_{j-1}) = \sin(\angle(\hat{x}, \hat{\mathbf{v}}_{j-1})) \leq \sqrt{1 - \tilde{\epsilon}^2}.$$

By (a) of Corollary 5.7,

$$(5.4) \quad \begin{aligned} \delta(\hat{g}_{j-1}\hat{x}, \hat{g}_{j-1}\hat{\mathbf{v}}_{j-1}) &= \delta(\hat{g}_{j-1}\hat{x}, \hat{\mathbf{v}}_{j-1}^1) = \sin(\angle(\hat{g}_{j-1}\hat{x}, \hat{\mathbf{v}}_{j-1}^1)) \\ &\leq \kappa\tilde{\epsilon}^{-1}\sqrt{1 - \tilde{\epsilon}^2} \leq \frac{c_0\epsilon}{t}\sqrt{1 - t^2\epsilon^2}. \end{aligned}$$

By (5.3) and Corollary 5.10, we have

$$(5.5) \quad \begin{aligned} \delta(\hat{\mathbf{v}}^*(g_{j-1}), \hat{\mathbf{v}}(g_j)) &= \delta(\hat{\mathbf{v}}_{j-1}^1, \hat{\mathbf{v}}_j) = \sin(\angle(\hat{\mathbf{v}}_{j-1}^1, \hat{\mathbf{v}}_j)) \\ &\leq \sqrt{1 - \frac{\epsilon^2}{1 + 2\frac{\kappa^2}{\epsilon^2}}} \leq \sqrt{1 - \frac{\epsilon^2}{1 + 2c_0^2\epsilon^2}}. \end{aligned}$$

Let $\theta_1 = \angle(\hat{g}_{j-1}\hat{x}, \hat{\mathbf{v}}_{j-1}^1)$ and $\theta_2 = \angle(\hat{\mathbf{v}}_{j-1}^1, \hat{\mathbf{v}}_j)$. Then

$$(5.6) \quad \begin{aligned} \delta(\hat{g}_{j-1}\hat{x}, \hat{\mathbf{v}}_j) &\leq |\cos\theta_1|\sin\theta_2 + \sin\theta_1|\cos\theta_2| \\ &= \sqrt{1 - \sin^2\theta_1}\sin\theta_2 + \sin\theta_1\sqrt{1 - \sin^2\theta_2} := f(\sin\theta_1, \sin\theta_2). \end{aligned}$$

With $f(x, y) = y\sqrt{1 - x^2} + x\sqrt{1 - y^2}$, it is easy to see that both $\frac{\partial f}{\partial x}$ and $\frac{\partial f}{\partial y}$ have the same sign as $\sqrt{1 - x^2}\sqrt{1 - y^2} - xy$. Thus both $\frac{\partial f}{\partial x}$ and $\frac{\partial f}{\partial y}$ are positive if $x^2 + y^2 < 1$.

By (5.4) and (5.5), we have

$$(5.7) \quad \sin^2\theta_1 + \sin^2\theta_2 \leq \frac{c_0^2\epsilon^2}{t^2}(1 - t^2\epsilon^2) + 1 - \frac{\epsilon^2}{1 + 2c_0^2\epsilon^2} < 1.$$

Here it is enough to have that for $\tilde{\epsilon} = t\epsilon$,

$$\frac{c_0^2}{t^2} + c_0^2\epsilon^2 < 1.$$

Then (5.6) implies

$$(5.8) \quad \begin{aligned} \delta(\hat{g}_{j-1}\hat{x}, \hat{\mathbf{v}}_j) &\leq f\left(\frac{c_0\epsilon}{t}\sqrt{1 - t^2\epsilon^2}, \sqrt{1 - \frac{\epsilon^2}{1 + 2c_0^2\epsilon^2}}\right) \\ &< \left(1 - \frac{c_0^2\epsilon^2}{2t^2}(1 - t^2\epsilon^2)\right) \left(1 - \frac{\epsilon^2}{2 + 4c_0^2\epsilon^2}\right) + \frac{c_0\epsilon^2}{t}\left(1 - \frac{c_0^2\epsilon^2}{2}\right) \\ &< \sqrt{1 - t^2\epsilon^2} = \sqrt{1 - \tilde{\epsilon}^2}. \end{aligned}$$

(By our choice of $c_0 = \frac{1}{10}$ and $t = \frac{2}{3}$, the ϵ^2 coefficients of (5.8) corresponds to $-\frac{9}{800} - \frac{1}{2} + \frac{3}{20} < -\frac{2}{9}$.) \square

This lemma has the following intermediate corollary.

Corollary 5.13. *For any $1 \leq j \leq 2n - 1$ and $1 \leq m \leq 2n - j - 1$, we have*

$$\hat{g}_{j+m-1} \cdots \hat{g}_j \hat{g}_{j-1} B(\hat{\mathbf{v}}_{j-1}, \sqrt{1 - \tilde{\epsilon}^2}) \subseteq B(\hat{\mathbf{v}}_{j+m}, \sqrt{1 - \tilde{\epsilon}^2}).$$

Next, let us show

Lemma 5.14. *For any $0 \leq j \leq 2n - 1$, for any $\hat{x} \in B(\hat{\mathbf{v}}_j, \sqrt{1 - \tilde{\epsilon}^2})$,*

$$\delta(\hat{g}_{2n-1} \cdots \hat{g}_{j+2} \hat{g}_j \hat{x}, \hat{\mathbf{v}}_j^{2n-j}) \leq \kappa \tilde{\epsilon}^{-1} \left(\frac{\sqrt{2}\pi}{2} \kappa \tilde{\epsilon}^{-2} \right)^{2n-j-1}$$

Proof. By Corollary 5.13, for any $0 \leq m \leq 2n - j - 1$, we have that both the two elements $\hat{g}_{j+m-1} \cdots \hat{g}_{j+1} \hat{g}_j \hat{x}$ and $\hat{\mathbf{v}}_j^m$ belong to $B(\hat{\mathbf{v}}_{j+m}, \sqrt{1 - \tilde{\epsilon}^2})$. Hence by (a) of 5.7, we have that for $m = 0$,

$$(5.9) \quad \delta(\hat{g}_j \hat{x}, \hat{\mathbf{v}}_j^1) = \delta(\hat{g}_j \hat{x}, \hat{g}_j \hat{\mathbf{v}}_j) \leq \kappa \tilde{\epsilon}^{-1} \delta(\hat{x}, \hat{\mathbf{v}}_j) < \kappa \tilde{\epsilon}^{-1}.$$

For $1 \leq m \leq 2n - j - 1$, by (b) of 5.7, we have

$$(5.10) \quad \begin{aligned} \delta(\hat{g}_{j+m} \hat{g}_{j+m-1} \cdots \hat{g}_{j+1} \hat{g}_j \hat{x}, \hat{\mathbf{v}}_j^{m+1}) &= \delta(\hat{g}_{j+m} \hat{g}_{j+m-1} \cdots \hat{g}_{j+1} \hat{g}_j \hat{x}, \hat{g}_{j+m} \hat{\mathbf{v}}_j^m) \\ &\leq \frac{\sqrt{2}\pi}{2} \kappa \tilde{\epsilon}^{-2} \delta(\hat{g}_{j+m-1} \cdots \hat{g}_{j+1} \hat{g}_j \hat{x}, \hat{\mathbf{v}}_j^m). \end{aligned}$$

(5.9) and (5.10) imply that

$$(5.11) \quad \delta(\hat{g}_{2n-1} \cdots \hat{g}_{j+2} \hat{g}_j \hat{x}, \hat{\mathbf{v}}_j^{2n-j}) \leq \kappa \tilde{\epsilon}^{-1} \left(\frac{\sqrt{2}\pi}{2} \kappa \tilde{\epsilon}^{-2} \right)^{2n-j-1}$$

as desired. \square

In particular, combining Corollary 5.13 with Lemma 5.14, we have the following corollary

Corollary 5.15. *For any $1 \leq j \leq 2n - 1$,*

$$\delta(\hat{\mathbf{v}}_{j-1}^{2n-j+1}, \hat{\mathbf{v}}_j^{2n-j}) \leq \kappa \tilde{\epsilon}^{-1} \left(\frac{\sqrt{2}\pi}{2} \kappa \tilde{\epsilon}^{-2} \right)^{2n-j-1}$$

Next, we will show

Lemma 5.16. *For any $\hat{x} \in B(\hat{\mathbf{v}}_0, \sqrt{1 - \tilde{\epsilon}^2})$, we have*

$$\delta(\widehat{(g^n)^* g^n \hat{x}}, \hat{\mathbf{v}}_0) \leq 3\kappa \tilde{\epsilon}^{-1}, \quad \delta(\hat{\mathbf{v}}_0^{2n}, \hat{\mathbf{v}}_{2n-1}^1) \leq 3\kappa \tilde{\epsilon}^{-1}.$$

Proof. By Corollary 5.15, we have

$$(5.12) \quad \delta(\hat{\mathbf{v}}_0^{2n}, \hat{\mathbf{v}}_{2n-1}^1) \leq \sum_{j=1}^{2n-1} \delta(\hat{\mathbf{v}}_{j-1}^{2n-j+1}, \hat{\mathbf{v}}_j^{2n-j}) \leq \kappa \tilde{\epsilon}^{-1} \sum_{j=0}^{2n-2} \left(\frac{\sqrt{2}\pi}{2} \kappa \tilde{\epsilon}^{-2} \right)^j.$$

By Lemma 5.14,

$$(5.13) \quad \delta(\hat{g}_{2n-1} \cdots \hat{g}_1 \hat{g}_0 \hat{x}, \hat{\mathbf{v}}_0^{2n}) \leq \kappa \tilde{\epsilon}^{-1} \left(\frac{\sqrt{2}\pi}{2} \kappa \tilde{\epsilon}^{-2} \right)^{2n-1}.$$

Hence, combining (5.12) with (5.13), we conclude that

$$\delta(\widehat{(g^n)^* g^n} \hat{x}, \hat{\mathbf{v}}(g_0)) \leq \kappa \tilde{\epsilon}^{-1} \sum_{j=0}^{2n-1} \left(\frac{\sqrt{2}\pi}{2} \kappa \tilde{\epsilon}^{-2} \right)^j \leq \frac{\kappa \tilde{\epsilon}^{-1}}{1 - \frac{\sqrt{2}\pi}{2} \kappa \tilde{\epsilon}^{-2}} \leq \frac{1}{t - \frac{\pi \epsilon_0}{\sqrt{2}t}} \kappa \epsilon^{-1} < 3\kappa \epsilon^{-1}.$$

□

We are now ready to give the

Proof of (i). Lemma 5.16 shows that $\widehat{(g^n)^* g^n}$ maps the ball $B(\hat{\mathbf{v}}_0, \sqrt{1 - \tilde{\epsilon}^2})$ into itself. By Corollary 5.7, it has the contracting Lipschitz factor $\leq (\frac{\sqrt{2}\pi}{2} \kappa \tilde{\epsilon}^{-2})^{2n} \ll 1$. Therefore the map $\widehat{(g^n)^* g^n}$ has a unique fixed point in $B(\hat{\mathbf{v}}_0, \sqrt{1 - \tilde{\epsilon}^2})$, call it x_* . Lemma 5.16 implies that

$$(5.14) \quad \delta(x_*, \hat{\mathbf{v}}(g_0)) < 3\kappa \epsilon^{-1}.$$

The claim will follow once we prove that $x_* = \hat{\mathbf{v}}(g^n)$. Since $\hat{\mathbf{v}}(g^n)$ is a fixed point of $\widehat{(g^n)^* g^n}$, it suffices to prove that $\hat{\mathbf{v}}(g^n) \in B(\hat{\mathbf{v}}_0, \sqrt{1 - \tilde{\epsilon}^2})$. (Notice that $\widehat{(g^n)^* g^n}$ has several fixed points, one for every eigenvalue of $(g^n)^* g^n$.)

Let $\delta_* := \delta(\hat{\mathbf{v}}(g^n), x_*)$. We will show that $\delta_* = 0$. For any unit vector v , we have

$$|\langle \mathbf{v}(g^n), v \rangle| = \frac{1}{(s_1(g^n))^2} |\langle (g^n)^* g^n \mathbf{v}(g^n), v \rangle| \leq \left| \langle \mathbf{v}(g^n), \frac{(g^n)^* g^n v}{|(g^n)^* g^n v|} \rangle \right|,$$

where we used that $|(g^n)^* g^n v| \leq (s_1(g^n))^2$. This lifts to a relation on projective space:

$$(5.15) \quad \delta(\hat{\mathbf{v}}(g^n), \widehat{(g^n)^* g^n} \hat{v}) \leq \delta(\hat{\mathbf{v}}(g^n), \hat{v}).$$

We apply this with $\hat{v} = \hat{v}_*$ as the “halfway point” between $\hat{\mathbf{v}}(g^n)$ and x_* , i.e., \hat{v}_* satisfies

$$\delta(\hat{\mathbf{v}}(g^n), \hat{v}_*) = \delta(x_*, \hat{v}_*) = \frac{\delta_*}{2}.$$

(This \hat{v}_* can be constructed by following the arc that connects $\hat{\mathbf{v}}(g^n)$ with x_* , assuming that $\mathbf{v}(g^n) \neq x_*$.)

Notice that $\hat{v}_* \in B(\hat{\mathbf{v}}_0, \sqrt{1 - \tilde{\epsilon}^2})$ because (5.14) gives

$$\delta(\hat{\mathbf{v}}_0, \hat{v}_*) \leq \delta(\hat{\mathbf{v}}_0, x_*) + \frac{\delta_*}{2} \leq \frac{3\kappa}{\epsilon} + \frac{\delta_*}{2} \leq \frac{3\epsilon}{10} + \frac{1}{2} < \sqrt{1 - \tilde{\epsilon}^2}.$$

Recall that $\widehat{(g^n)^* g^n}$ maps the ball $B(\hat{\mathbf{v}}_0, \sqrt{1 - \tilde{\epsilon}^2})$ into itself with Lipschitz factor $\leq L_0 := (\frac{\sqrt{2}\pi}{2} \kappa \tilde{\epsilon}^{-2})^{2n} \ll 1$. Since $\widehat{(g^n)^* g^n} x_* = x_*$, we have

$$\delta(\widehat{(g^n)^* g^n} \hat{v}_*, x_*) \leq L_0 \delta(\hat{v}_*, x_*) = L_0 \frac{\delta_*}{2}.$$

We combine this bound and (5.15), with $\hat{v} = \hat{v}_*$, to conclude that

$$\delta_* = \delta(\hat{\mathbf{v}}(g^n), x_*) \leq \delta(\hat{\mathbf{v}}(g^n), (\widehat{g^n})^* \widehat{g^n} \hat{v}_*) + \delta((\widehat{g^n})^* \widehat{g^n} \hat{v}_*, x_*) \leq \left(1 + \frac{L_0}{2}\right) \delta_*.$$

Since $L_0 < 1$, this implies $\delta_* = 0$, i.e., $x_* = \hat{\mathbf{v}}(g^n)$. Consequently, (5.14) reads $\delta(\hat{\mathbf{v}}(g^n), \hat{\mathbf{v}}(g_0)) \leq 3\kappa\epsilon^{-1}$ as claimed in (i) of Theorem 5.5. The other bound in (i) can be proved in exactly the same way.

Proof of (ii) By Lemma 5.11, we have

$$(5.16) \quad \prod_{j=1}^{n-1} \frac{\alpha(g^j, g_j)}{\beta(g_{j-1}, g_j)} \leq \frac{\rho(g_0, \dots, g_{n-1})}{\prod_{j=1}^{n-1} \rho(g_{j-1}, g_j)} \leq \prod_{j=1}^{n-1} \frac{\beta(g^j, g_j)}{\alpha(g_{j-1}, g_j)}.$$

We will show that the factors

$$\frac{\alpha(g^j, g_j)}{\beta(g_{j-1}, g_j)} \quad \text{and} \quad \frac{\beta(g^j, g_j)}{\alpha(g_{j-1}, g_j)}$$

are very close to 1, with logarithms of order $\kappa\epsilon^{-2}$. From conclusion (i), applied to the sequence of matrices g_0, g_1, \dots, g_{j-1} , we have

$$(5.17) \quad \max\{\delta(\hat{\mathbf{v}}^*(g^j), \hat{\mathbf{v}}^*(g_{j-1})), \delta(\hat{\mathbf{v}}(g^j), \hat{\mathbf{v}}(g_0))\} \leq 3\kappa\epsilon^{-1}$$

for all $1 \leq j \leq n$. From (5.17), we deduce that

$$(5.18) \quad \left| \log \frac{\alpha(g^j, g_j)}{\alpha(g_{j-1}, g_j)} \right| \leq \frac{|\alpha(g^j, g_j) - \alpha(g_{j-1}, g_j)|}{\min\{\alpha(g^j, g_j), \alpha(g_{j-1}, g_j)\}} \\ \leq \frac{\delta(\hat{\mathbf{v}}^*(g^j), \hat{\mathbf{v}}^*(g_{j-1}))}{\min\{\alpha(g^j, g_j), \alpha(g_{j-1}, g_j)\}} \leq \frac{3\kappa\epsilon^{-1}}{\min\{\alpha(g^j, g_j), \alpha(g_{j-1}, g_j)\}}.$$

We estimate the minimum as follows, using (5.17) and Corollary 5.10,

$$(5.19) \quad \min\{\alpha(g^j, g_j), \alpha(g_{j-1}, g_j)\} \geq \alpha(g_{j-1}, g_j) - |\alpha(g^j, g_j) - \alpha(g_{j-1}, g_j)| \\ \geq \alpha(g_{j-1}, g_j) - 3\kappa\epsilon^{-1} \\ \geq \frac{\epsilon}{\sqrt{1 + 2\frac{\kappa^2}{\epsilon^2}}} - 3\frac{\kappa}{\epsilon} \\ \geq \frac{2\epsilon}{3}.$$

In the last step, we used $\frac{\kappa}{\epsilon} \leq \frac{\epsilon}{10}$. Returning to (5.18), we have shown

$$(5.20) \quad \left| \log \frac{\alpha(g^j, g_j)}{\alpha(g_{j-1}, g_j)} \right| \leq \frac{9}{2} \frac{\kappa}{\epsilon^2}.$$

From the definition of the upper angle β and Corollary 5.10, we also have

$$(5.21) \quad \left| \log \frac{\beta(g_{j-1}, g_j)}{\alpha(g_{j-1}, g_j)} \right| \leq \log \sqrt{1 + 2\frac{\kappa^2}{\alpha(g_{j-1}, g_j)}} \leq \log \sqrt{1 + 2\frac{\kappa^2}{\epsilon^2}} \leq \frac{\kappa^2}{\epsilon^2}.$$

Hence (5.20) and (5.21) yield

$$\left| \log \frac{\alpha(g^j, g_j)}{\beta(g_{j-1}, g_j)} \right| \leq \frac{9}{2} \frac{\kappa}{\epsilon^2} + \frac{\kappa^2}{\epsilon^2} < 5 \frac{\kappa}{\epsilon^2}.$$

Together with (5.16), this implies the lower bound in (ii), i.e.,

$$e^{-5n\kappa/\epsilon^2} \leq \frac{\rho(g_0, g_1, \dots, g_{n-1})}{\rho(g_0, g_1) \cdots \rho(g_{n-2}, g_{n-1})}.$$

For the upper bound, we argue similarly. The only difference occurs in the analog of (5.21), i.e., the estimate on

$$\left| \log \frac{\beta(g^j, g_j)}{\alpha(g^j, g_j)} \right|.$$

To bound this quantity, we need to control the gap ratio $(\text{gr}(g^j))^{-1}$. This control is provided by the following lemma.

Lemma 5.17. *We have $(\text{gr}(g^j))^{-1} \leq \kappa' := 20 \frac{\kappa}{\epsilon}$.*

We postpone the proof of the lemma for now. It gives

$$\left| \log \frac{\beta(g^j, g_j)}{\alpha(g^j, g_j)} \right| \leq \log \sqrt{1 + 2 \frac{(\kappa')^2}{\alpha(g^j, g_j)}} \leq 600 \frac{\kappa^2}{\epsilon^3}.$$

In the second step, we used that $\alpha(g^j, g_j) \geq 2\epsilon/3$ by (5.19). Recalling (5.20), one has

$$\left| \log \frac{\beta(g^j, g_j)}{\alpha(g_{j-1}, g_j)} \right| \leq \frac{9}{2} \frac{\kappa}{\epsilon^2} + 600 \frac{\kappa^2}{\epsilon^3} < 11 \frac{\kappa}{\epsilon^2}.$$

In the last estimate, we used that $600\kappa/\epsilon \leq 60\epsilon \leq 6$.

By (5.16), this proves the upper bound in (ii), i.e.,

$$\frac{\rho(g_0, g_1, \dots, g_{n-1})}{\rho(g_0, g_1) \cdots \rho(g_{n-2}, g_{n-1})} \leq e^{11n\kappa/\epsilon^2}.$$

It remains to prove Lemma 5.17. For this part, we follow page 71 in [DuaKle] and make the constants precise. From Proposition 2.28 in [DuaKle], we see that

$$(\text{gr}(g^j))^{-1} = \|(D\hat{g}^j)_{\hat{\mathbf{v}}(g^j)}\|.$$

where $(D\hat{g}^j)_{\hat{\mathbf{v}}(g^j)}$ is the derivative of $\hat{g}^j : \mathbb{P}(\mathbb{R}^d) \rightarrow \mathbb{P}(\mathbb{R}^d)$, evaluated at $\hat{\mathbf{v}}(g^j)$. The norm of this derivative is bounded by the Lipschitz constant in a neighborhood of $\hat{\mathbf{v}}(g^j)$. Since the Lipschitz constant is $\leq L^j$, with $L := \frac{\sqrt{2\pi}}{2} \kappa \tilde{\epsilon}^{-2}$, everywhere on the ball $B(\hat{\mathbf{v}}_0, \sqrt{1 - \tilde{\epsilon}^2})$, we immediately obtain the bound $(\text{gr}(g^j))^{-1} \leq L^j$. However, this is not good enough for our purposes (note that L is an order one quantity in general).

We may improve the estimate as follows: Applying statement (i) of the theorem with $n = j$, we obtain that

$$\delta(\hat{\mathbf{v}}(g^j), \hat{\mathbf{v}}(g_0)) < 3 \frac{\kappa}{\epsilon}.$$

Now, a calculation based on Proposition 2.28 in [DuaKle] shows that

$$\|(D\hat{g}_0)_{\hat{v}(g^j)} - (D\hat{g}_0)_{\hat{v}(g_0)}\| \leq 12\pi \frac{\kappa}{\epsilon}$$

and therefore

$$\|(D\hat{g}_0)_{\hat{v}(g^j)}\| \leq \|(D\hat{g}_0)_{\hat{v}(g_0)}\| + 12\pi \frac{\kappa}{\epsilon} = \text{gr}(g_0)^{-1} + 12\pi \frac{\kappa}{\epsilon} \leq \kappa + 12\pi \frac{\kappa}{\epsilon} \leq \left(12\pi + \frac{1}{10}\right) \frac{\kappa}{\epsilon}.$$

Finally, we apply the chain rule and estimate the derivative of the product $g_{j-1} \dots g_1$ by its Lipschitz constant L^{j-1} , which satisfies $L^{j-1} \leq L \leq \frac{9\pi}{40\sqrt{2}}$ for $j \geq 2$. Therefore,

$$\begin{aligned} (\text{gr}(g^j))^{-1} &= \|(D\hat{g}^j)_{\hat{v}(g^j)}\| \leq \|(D\hat{g}_{j-1} \dots \hat{g}_1)_{\hat{v}(g^j)} g\| \|(D\hat{g}_0)_{\hat{v}(g^j)}\| \\ &\leq L \left(12\pi + \frac{1}{10}\right) \frac{\kappa}{\epsilon} < 20 \frac{\kappa}{\epsilon}. \end{aligned}$$

This proves Lemma 5.17 and hence completes the proof of Theorem 5.5. \square

6. HERMAN'S REGULARIZATION

6.1. Monodromy matrices. One has $T_\omega^n(x, y) = (x + ny + \frac{n(n-1)}{2}\omega, y + n\omega)$ for any positive integer n , where T is the skew-shift with frequency ω . Denote the projection of \mathbb{T}^2 onto the first coordinate by \mathcal{P} , viz. $\mathcal{P}(x, y) = x$.

We consider the Schrödinger operator

$$(H_{\lambda, \omega, x, y} \psi)_n = \psi_{n+1} + \psi_{n-1} + 2\lambda \cos(2\pi \mathcal{P}(T_\omega^n(x, y))) \psi_n$$

with $\lambda > 0$. This equation has the following cocycle reformulation

$$\begin{aligned} (6.1) \quad \begin{pmatrix} \psi_{n+1} \\ \psi_n \end{pmatrix} &= \begin{pmatrix} E - 2\lambda \cos(2\pi \mathcal{P}(T_\omega^n(x, y))) & -1 \\ 1 & 0 \end{pmatrix} \begin{pmatrix} \psi_n \\ \psi_{n-1} \end{pmatrix} \\ &=: M(\lambda, E; T_\omega^n(x, y)) \begin{pmatrix} \psi_n \\ \psi_{n-1} \end{pmatrix}. \end{aligned}$$

Define the transfer matrices $M_n(\lambda, E; x, y)$ to be

$$(6.2) \quad M_n(\lambda, E; x, y) = \begin{cases} \prod_{j=n}^1 M(\lambda, E; T_\omega^j(x, y)), & n \geq 1, \\ \text{Id}, & n = 0, \\ (M_{-n}(\lambda, E; T_\omega^{n+1}(x, y)))^{-1}, & n < 0. \end{cases}$$

Then

$$\begin{pmatrix} \psi_{n+1} \\ \psi_n \end{pmatrix} = M_n(E; x, y) \begin{pmatrix} \psi_1 \\ \psi_0 \end{pmatrix},$$

The following function on \mathbb{T}^2 plays a fundamental role in our analysis:

$$(6.3) \quad \begin{aligned} u_n(\lambda, E; x, y) &:= \frac{1}{n} \log \|M_n(\lambda, E; x, y)\| \\ &= \frac{1}{n} \log \left\| \prod_{j=n}^1 \begin{pmatrix} E - \lambda e^{2\pi i(x+jy+\frac{j(j-1)}{2}\omega)} & -\lambda e^{-2\pi i(x+jy+\frac{j(j-1)}{2}\omega)} \\ 1 & 0 \end{pmatrix} \right\|. \end{aligned}$$

Let $z = e^{2\pi i x}$, $w = e^{2\pi i y}$, $a = e^{\pi i \omega}$, as well as

$$(6.4) \quad A_\lambda(\lambda, E, z, w, a) := \begin{pmatrix} Ezw - \lambda z^2 w^2 a - \lambda \bar{a} & -zw \\ zw & 0 \end{pmatrix}.$$

Then for $(z, w) \in \partial D_1 \times \partial D_1$,

$$(6.5) \quad \begin{aligned} u_n(\lambda, E; x, y) &= \frac{1}{n} \log \left\| \prod_{j=n}^1 \begin{pmatrix} E - \lambda z w^j a^{j(j-1)} - \lambda z^{-1} w^{-j} \bar{a}^{j(j-1)} & -1 \\ 1 & 0 \end{pmatrix} \right\| \\ &= \frac{1}{n} \log \left\| \prod_{j=n}^1 \begin{pmatrix} E z w^j - \lambda z^2 w^{2j} a^{j(j-1)} - \lambda \bar{a}^{j(j-1)} & -z w^j \\ z w^j & 0 \end{pmatrix} \right\| \\ &= \frac{1}{n} \log \left\| \prod_{j=n}^1 A_\lambda(E, z, w^j, a^{j(j-1)}) \right\| \\ &=: v_n(\lambda, E; z, w). \end{aligned}$$

Note that $v_n(\lambda, E; z, w)$ is a pluri-subharmonic function on \mathbb{C}^2 . *Herman's regularization* refers to the transition from the first to the second line in (6.5), which removes the singularities z^{-1} and w^{-1} . Note that

$$(6.6) \quad v_n(\lambda, E; 0, w) = v_n(\lambda, E; z, 0) = \log \lambda$$

For simplicity, we will write A instead of A_λ , since λ will be a fixed parameter within some range. As a general rule for the arguments of the matrix function A , the complex variables z, w will belong to some disk D_R , whereas $|a| = 1$ and E will be real-valued within some range. We will also keep $0 < \lambda < 1$.

6.2. Explicit bounds on the monodromy matrices. As a first step towards obtaining the explicit constants in Definition 2.1 and 4.1 we prove the following bounds on v_n .

Lemma 6.1. *Let $0 < \lambda < 1$, and $R_3 \geq 1$. Define*

$$(6.7) \quad U(\lambda, R_3) := \frac{1}{2} \log \left(\left(\lambda \left(1 + \frac{1}{R_3^2} \right)^2 + \frac{2}{R_3} \right)^2 + \frac{2}{R_3^2} \right).$$

Then for any E with $|E| \leq 2 + 2\lambda$, $v_n(\lambda, E, z, w)$ from (6.5) satisfies the following estimates

- for any $w \in \partial D_1$,

$$(6.8) \quad v_n(\lambda, E; z, w) \leq 2 \log R_3 + U(\lambda, R_3) \text{ for } \forall z \in \overline{D_{R_3}}, \text{ and } v_n(\lambda, E; 0, w) = \log \lambda.$$

- for any $(z, w) \in \partial D_1 \times \overline{D_{R_3}}$, we have upper bound

$$(6.9) \quad v_n(\lambda, E; z, w) \leq (n+1) \log R_3 + U(\lambda, R_3)$$

We also have $v_n(\lambda, E; z, 0) = \log \lambda$ for any $z \in \partial D_1$.

- for any $(z, w) \in \partial D_1 \times \partial D_1$,

$$(6.10) \quad |v_n(\lambda, E; z, w)| \leq U(\lambda, 1).$$

Remark 6.2. Let us note that

$$(6.11) \quad 4U(\lambda, 1) \geq 2 \log 6 > 1.$$

Proof. Clearly (6.10) follows from (6.8) with $R_3 = 1$. We will use that for any complex-valued matrix

$$\|A\|^2 = \|A^*A\| \leq \text{Tr}(A^*A)$$

For A as in (6.4) this means that

$$\begin{aligned} \|A(E, z, w, a)\|^2 &\leq |Ez w - \lambda z^2 w^2 a - \lambda \bar{a}|^2 + 2|z|^2 |w|^2 \\ &\leq (|E||z w| + \lambda |z w|^2 + \lambda)^2 + 2|z|^2 |w|^2 \end{aligned}$$

whence

$$(6.12) \quad \begin{aligned} |v_n(E; z, w)| &\leq \frac{1}{n} \sum_{j=1}^{2n} \log \|A(E, z, w^j, a^{j(j-1)})\| \\ &\leq \frac{1}{2n} \sum_{j=1}^n \log \left((\lambda(|z|^2 |w|^{2j} + 1) + |E||z||w|^j)^2 + 2|z|^2 |w|^{2j} \right). \end{aligned}$$

For $w \in \partial D_1$ and $|z| \leq R_3$, (6.12) yields

$$(6.13) \quad \begin{aligned} |v_n(E; z, w)| &\leq \frac{1}{2n} \sum_{j=1}^n \log \left((\lambda(R_3^2 + 1) + |E|R_3)^2 + 2R_3^2 \right) \\ &\leq 2 \log R_3 + \frac{1}{2} \log \left(\left(\lambda \left(1 + \frac{1}{R_3^2}\right) + \frac{2+2\lambda}{R_3} \right)^2 + \frac{2}{R_3^2} \right) \\ &= 2 \log R_3 + U(\lambda, R_3). \end{aligned}$$

This proves (6.8).

Next, we turn to (6.9). For $z \in \partial D_1$ and $|w| \leq R_3$, (6.12) yields

$$\begin{aligned} |v_n(E; z, w)| &\leq \frac{1}{2n} \sum_{j=1}^n \log \left((\lambda(R_3^{2j} + 1) + |E|R_3^j)^2 + 2R_3^{2j} \right) \\ &\leq (n+1) \log R_3 + \frac{1}{2n} \sum_{j=1}^n \log \left(\left(\lambda \left(1 + \frac{1}{R_3^{2j}}\right) + (2+2\lambda) \frac{1}{R_3^j} \right)^2 + \frac{2}{R_3^{2j}} \right). \end{aligned}$$

Note that the summands are maximized at $j = 1$, which gives us the constant $2U(\lambda, R_3)$. Hence, in total

$$|v_n(E; z, w)| \leq (n+1) \log R_3 + U(\lambda, R_3)$$

as claimed. \square

7. LONG SUMS OF SKEW-SHIFT FUNCTIONS

In this section we establish a key large-deviation estimate on the ergodic averages of a pluri-subharmonic function, as defined above, over a long skew-shift orbit. The argument is based on [BouGolSch, Lemma 2.6], but deviates from that reference in ways which are essential for our purposes. The precise dependence on all parameters is made explicit and effective. This leads to a somewhat cumbersome formulation which is, however, absolutely necessary for the main application. We wish to point out one technical feature of our version of this argument, namely that we only use a trivial bound on the number theoretic divisor function, see the constant C^* below. We have found this to lead to the best constants. We also remark that significant gains in the following proposition would lead to dramatic improvements in the inductive machinery that we use to control the Lyapunov exponent, cf. the next two sections. At this point, however, it is not clear how to obtain such gains.

Recall constants $B_1(R, R_1, R_2)$ is as in (2.1), $B_3(R, R_1, R_2)$ is as in (2.2), and $B_4(R), B_5(R), m_4, m_5$ are as in (8.8). In the following we will write B_1, B_3, B_4, B_5 and omit the dependence on the radii.

Proposition 7.1. *Let $\omega = \frac{\sqrt{5}-1}{2}$ be the golden ratio. Let v be defined as in the beginning of Section 4, let $C = C(R, R_1, R_2)$ be the constant as in (3.47), and impose Definition 3.5. Let $\delta \in (0, 1/2)$ and $\delta_2, \delta_3 > 0$ be constants. Assuming*

- (i). $C(B_5 - m_5) \leq K^\delta$,
- (ii). $K \geq 38$,
- (iii). $\exp\left(4(\log K)^{\delta_2}\right) \geq K + 1$,
- (iv). $21K^{-\frac{9}{10} + \frac{9}{5}\delta}(\log K)^{\frac{9}{10} + \frac{9}{5}\delta_2} + 4C(B_4 - m_4) \leq K^\delta(\log K)^{\delta_2}$.

Then for any positive parameter $C_2 > 0$, we have

$$\begin{aligned} & \left| \left\{ (x, y) \in \mathbb{T}^2 : \left| \frac{1}{K} \sum_{k=1}^K v \circ T_\omega^k(x, y) - \langle v \rangle \right| > \varepsilon_4 \right\} \right| \\ & \leq 2\sqrt{2} \exp\left(\frac{\pi}{4} \left[\frac{17}{36} + \frac{B_1}{4B_3^2} - \varepsilon_5 \right]\right) \\ & \quad + \sqrt{2}(C(B_4 - m_4))^{-1} K^{\frac{1}{5} - \frac{2\delta}{5}} (\log K)^{-\frac{1}{5} - \frac{2\delta_2}{5}} \exp\left(-2(\log K)^{\delta_2}\right), \end{aligned}$$

where

$$(7.1) \quad \begin{aligned} \varepsilon_4 &= C_2 K^{-\frac{1}{10} + \frac{\delta}{5}} (\log K)^{\frac{1}{10} + \frac{\delta_2}{5} + \delta_3}, \\ \varepsilon_5 &= C_2 \left(472.5 + 3.2B_3(B_4 - m_4)\sqrt{C} \right)^{-1} (\log K)^{\delta_3}. \end{aligned}$$

Before proving Proposition 7.1, we will review some background of continued fractions.

7.1. Continued fractions. Each $\omega \in [0, 1)$ has the following unique expansion

$$(7.2) \quad \omega = \frac{1}{a_1 + \frac{1}{a_2 + \frac{1}{a_3 + \dots}}},$$

where $a_i \in \mathbb{N}_+$. We will denote this expansion by $\omega = [a_1, a_2, \dots]$. If $\omega \in \mathbb{Q}$, the expansion is finite, while it is infinite for irrational ω .

Let $\omega \in [0, 1) \setminus \mathbb{Q}$, let

$$(7.3) \quad \frac{p_n}{q_n} = \frac{1}{a_1 + \frac{1}{a_2 + \frac{1}{\dots + \frac{1}{a_n}}}}$$

be the continued fraction approximants of ω . These approximants satisfy the following three properties:

$$(7.4) \quad q_{n+1} = a_{n+1}q_n + q_{n-1} \quad \text{with } q_0 := 1.$$

$$(7.5) \quad \|k\omega\|_{\mathbb{T}} \geq \|q_n\omega\|_{\mathbb{T}} \quad \text{for any } q_n \leq |k| < q_{n+1}.$$

$$(7.6) \quad \frac{1}{q_{n+1} + q_n} \leq \|q_n\omega\|_{\mathbb{T}} \leq \frac{1}{q_{n+1}}.$$

A very special number we are interested in is the golden ratio $\omega = \frac{\sqrt{5}-1}{2}$, it is well-known that ω has continued fraction expansion with $a_i \equiv 1$ for any $i \geq 1$. Then by (7.4), we have $q_{n+1} = q_n + q_{n-1}$ with $q_0 = q_1 = 1$. It is then easy to find out that for any $n \geq 0$, Hence for any $n \geq 0$, $q_{n+1} \leq 2q_n$. Then by (7.5) and (7.6), we have the following property of the golden ratio:

Proposition 7.2. *The golden ratio satisfies*

$$(7.7) \quad \|k\omega\|_{\mathbb{T}} \geq \frac{1}{3|k|} \quad \text{for any } k \neq 0.$$

The optimal bound here is $\frac{\sqrt{5}+1}{2}$, but the constant 3 is sufficient. We will use the following corollary of (7.7) repeatedly.

Corollary 7.3. *The golden ratio satisfies the following two properties:*

- Let ℓ be a positive integer such that $\|\ell\omega\|_{\mathbb{T}} \leq \sigma$, then $\ell \geq \frac{1}{3\sigma}$.
- Let $\ell, \tilde{\ell}$ be two distinct positive integers such that $\max(\|\ell\omega\|_{\mathbb{T}}, \|\tilde{\ell}\omega\|_{\mathbb{T}}) \leq \sigma$, then $|\ell - \tilde{\ell}| \geq \frac{1}{6\sigma}$.

In order to control small divisors we will rely on the following two propositions.

Lemma 7.4. *For $\theta \in \mathbb{R}$, we have*

$$(7.8) \quad \left| \sin\left(\frac{\pi}{2}\theta\right) \right| \geq \|\theta\|_{\mathbb{T}}.$$

Proof. If $\theta \in \mathbb{Z} + \frac{1}{2}$, $|\sin(\frac{\pi}{2}\theta)| = \sin(\frac{\pi}{4}) > \frac{1}{2} = \|\theta\|_{\mathbb{T}}$.

If $\theta \notin \mathbb{Z} + \frac{1}{2}$, then there exists a unique $k \in \mathbb{Z}$, such that $\theta = k + \|\theta\|_{\mathbb{T}}$ (if $\theta \in [k, k + \frac{1}{2})$), or $\theta = k - \|\theta\|_{\mathbb{T}}$ (if $\theta \in (k - \frac{1}{2}, k)$). If k is an even number, then

$|\sin(\frac{\pi}{2}\theta)| = \sin(\frac{\pi}{2}\|\theta\|_{\mathbb{T}}) \geq \|\theta\|_{\mathbb{T}}$, in which we used $\sin x \geq \frac{2}{\pi}x$ for $0 \leq x \leq \frac{\pi}{2}$. If k is odd, then $|\sin(\frac{\pi}{2}\theta)| = \cos(\frac{\pi}{2}\|\theta\|_{\mathbb{T}}) \geq \cos(\frac{\pi}{4}) > \frac{1}{2} \geq \|\theta\|_{\mathbb{T}}$. \square

We will also use the following two estimates:

Lemma 7.5. *For any positive integer R ,*

$$(7.9) \quad \left| \sum_{k=1}^R e(k\ell\omega) \right| \leq \min \left(R, \frac{2}{2 \sin(\pi\|\ell\omega\|_{\mathbb{T}})} \right) \leq \min \left(R, \frac{1}{2\|\ell\omega\|_{\mathbb{T}}} \right),$$

and

$$(7.10) \quad \left| \sum_{k=1}^R e\left(\frac{1}{2}k\ell\omega\right) \right| \leq \min \left(R, \frac{1}{\|\ell\omega\|_{\mathbb{T}}} \right).$$

Proof. For $\theta \notin \mathbb{Z}$, we have

$$\left| \sum_{k=1}^R e(k\theta) \right| = \min \left(R, \left| \frac{e(\theta) - e((k+1)\theta)}{1 - e(\theta)} \right| \right) \leq \min \left(R, \frac{1}{|\sin(\pi\theta)|} \right),$$

then (7.9) follows from taking $\theta = \ell\omega$, and using $\sin(\pi x) \geq 2x$ for $0 \leq x \leq \frac{1}{2}$. (7.10) follows from taking $\theta = \frac{1}{2}\ell\omega$ and employing (7.8). \square

7.2. Proof of Proposition 7.1. Let $\hat{v}(\ell, y)$ and $\hat{v}(x, \ell)$ denote the Fourier coefficients relative to the first and second variables, respectively, and by $\hat{v}(\ell_1, \ell_2)$ we mean the Fourier transform in both variables. For simplicity, let us omit the dependence of $C(R, R_1, R_2), B_3(R, R_1, R_2)$ on the radii.

First, we note the following estimate as a corollary of Lemma 3.13.

Corollary 7.6. *For any $\ell \neq 0$, we have*

$$(7.11) \quad \begin{aligned} \sup_{x \in \mathbb{T}} |\hat{v}(x, \ell)| &\leq \frac{C}{|\ell|} (B_5 - m_5), \\ \sup_{y \in \mathbb{T}} |\hat{v}(\ell, y)| &\leq \frac{C}{|\ell|} (B_4 - m_4), \end{aligned}$$

and

$$(7.12) \quad \begin{aligned} \left(\sum_{\ell_1 \in \mathbb{Z}} |\hat{v}(\ell_1, \ell_2)|^2 \right)^{\frac{1}{2}} &\leq \frac{C}{|\ell_2|} (B_5 - m_5), \quad \text{for any } \ell_2 \neq 0, \\ \left(\sum_{\ell_2 \in \mathbb{Z}} |\hat{v}(\ell_1, \ell_2)|^2 \right)^{\frac{1}{2}} &\leq \frac{C}{|\ell_1|} (B_4 - m_4), \quad \text{for any } \ell_1 \neq 0, \end{aligned}$$

Proof. Note that (7.11) follows directly from Lemma 3.13. On the other hand,

$$\left(\sum_{\ell_1 \in \mathbb{Z}} |\hat{v}(\ell_1, \ell_2)|^2 \right)^{\frac{1}{2}} = \|\hat{v}(\cdot, \ell_2)\|_{L^2(\mathbb{T})} \leq \sup_{x \in \mathbb{T}} |\hat{v}(x, \ell_2)|.$$

Hence, (7.12) reduces to (7.11). \square

With some positive integer p_1 to be determined, let

$$(7.13) \quad \begin{aligned} v(x, y) &= \sum_{|\ell_1| \leq p_1} \hat{v}(\ell_1, y) e(\ell_1 x) + \sum_{|\ell_1| > p_1} \hat{v}(\ell_1, y) e(\ell_1 x) \\ &=: v_1(x, y) + \tilde{v}_1(x, y), \end{aligned}$$

where v_1 and \tilde{v}_1 are the low and high frequency parts, respectively.

By Corollary 7.6,

$$(7.14) \quad \begin{aligned} \sup_{y \in \mathbb{T}} \|\tilde{v}_1(\cdot, y)\|_{L^1(\mathbb{T})} &\leq \left(\sum_{|\ell_1| > p_1} \sup_{y \in \mathbb{T}} |\hat{v}(\ell_1, y)|^2 \right)^{\frac{1}{2}} \\ &\leq C_5 \left(\sum_{|\ell_1| > p_1} \frac{1}{\ell_1^2} \right)^{\frac{1}{2}} (B_4 - m_4) \\ &\leq \sqrt{2} C (B_4 - m_4) p_1^{-\frac{1}{2}}. \end{aligned}$$

Next we further decompose v_1 into low and high frequency parts in the y variable. With some positive integer p_2 to be determined, let

$$(7.15) \quad \begin{aligned} v_1(x, y) &= \sum_{\substack{|\ell_1| \leq p_1 \\ |\ell_2| > p_2}} \hat{v}(\ell_1, \ell_2) e(\ell_1 x + \ell_2 y) + \sum_{\substack{|\ell_1| \leq p_1 \\ |\ell_2| \leq p_2}} \hat{v}(\ell_1, \ell_2) e(\ell_1 x + \ell_2 y) \\ &=: v_2(x, y) + v_3(x, y). \end{aligned}$$

By Corollary 7.6, we have

$$(7.16) \quad \begin{aligned} \|v_2(x, y)\|_{L^1(\mathbb{T}^2)} &\leq \|v_2(x, y)\|_{L^2(\mathbb{T}^2)} = \left(\sum_{|\ell_1| \leq p_1, |\ell_2| > p_2} |\hat{v}(\ell_1, \ell_2)|^2 \right)^{\frac{1}{2}} \\ &\leq \left(\sum_{\ell_1 \in \mathbb{Z}, |\ell_2| > p_2} |\hat{v}(\ell_1, \ell_2)|^2 \right)^{\frac{1}{2}} \leq \left(\sum_{|\ell_2| > p_2} \frac{C^2}{\ell_2^2} (B_5 - m_5)^2 \right)^{\frac{1}{2}} \\ &< \sqrt{2} C (B_5 - m_5) p_2^{-\frac{1}{2}}. \end{aligned}$$

Hence, by Markov's inequality,

$$(7.17) \quad \begin{aligned} &\left| \left\{ y \in \mathbb{T} : \frac{1}{K} \int_{\mathbb{T}} \left| \sum_{k=1}^K v_2 \circ T_\omega^k(x, y) \right| dx > t \right\} \right| \\ &\leq \sqrt{2} C (B_5 - m_5) p_2^{-\frac{1}{2}} t^{-1}. \end{aligned}$$

We denote the set on the left-hand side of 7.17 by $\mathcal{A}(t)$.

Now let us consider v_3 , which will lead to small divisor problems. By Corollary 7.6 and the fact that $\hat{v}(0, 0) = \langle v \rangle$, separating the cases $\ell_1 = 0$, $\ell_2 = 0$, and $\ell_1 \ell_2 \neq 0$,

yields

$$\begin{aligned}
& \sup_{(x,y) \in \mathbb{T}^2} \left| \frac{1}{K} \sum_{k=1}^K v_3 \circ T_\omega^k(x,y) - \langle v \rangle \right| \\
& \leq \frac{1}{K} \sum_{\substack{|\ell_1| \leq p_1 \\ |\ell_2| \leq p_2 \\ |\ell_1| + |\ell_2| \neq 0}} |\hat{v}(\ell_1, \ell_2)| \left| \sum_{k=1}^K e \left(\ell_1 \left(ky + \frac{k(k-1)\omega}{2} \right) + \ell_2 k\omega \right) \right| \\
(7.18) \quad & \leq \frac{C(B_5 - m_5)}{K} \sum_{1 \leq |\ell_2| \leq p_2} \frac{1}{|\ell_2|} \left| \sum_{k=1}^K e(\ell_2 k\omega) \right| \\
& + \frac{C(B_4 - m_4)}{K} \sum_{1 \leq |\ell_1| \leq p_1} \frac{1}{|\ell_1|} \left| \sum_{k=1}^K e \left(\ell_1 \left(ky + \frac{k(k-1)\omega}{2} \right) \right) \right| \\
& + \frac{1}{K} \sum_{1 \leq |\ell_2| \leq p_2} \sum_{1 \leq |\ell_1| \leq p_1} |\hat{v}(\ell_1, \ell_2)| \left| \sum_{k=1}^K e \left(\ell_1 \left(ky + \frac{k(k-1)\omega}{2} \right) + \ell_2 k\omega \right) \right|
\end{aligned}$$

We now separately consider the sums appearing on the previous three lines. First,

$$S_1 := \frac{1}{K} \sum_{1 \leq |\ell_2| \leq p_2} \frac{1}{|\ell_2|} \left| \sum_{k=1}^K e(\ell_2 k\omega) \right|$$

Second, by Cauchy-Schwarz,

$$\begin{aligned}
& K^{-1} \sum_{1 \leq |\ell_1| \leq p_1} \frac{1}{|\ell_1|} \left| \sum_{k=1}^K e \left(\ell_1 \left(ky + \frac{k(k-1)\omega}{2} \right) \right) \right| \\
(7.19) \quad & \leq K^{-1} \left(\sum_{1 \leq |\ell_1| \leq p_1} \frac{1}{\ell_1^2} \right)^{\frac{1}{2}} \left(\sum_{1 \leq |\ell_1| \leq p_1} \left| \sum_{k=1}^K e \left(\ell_1 \left(ky + \frac{k(k-1)\omega}{2} \right) \right) \right|^2 \right)^{\frac{1}{2}} \\
& \leq 2K^{-1} \left(\sum_{1 \leq |\ell_1| \leq p_1} \left| \sum_{k=1}^K e \left(\ell_1 \left(ky + \frac{k(k-1)\omega}{2} \right) \right) \right|^2 \right)^{\frac{1}{2}} =: S_2
\end{aligned}$$

And, finally,

$$\begin{aligned}
& K^{-1} \sum_{1 \leq |\ell_2| \leq p_2} \sum_{1 \leq |\ell_1| \leq p_1} |\hat{v}(\ell_1, \ell_2)| \left| \sum_{k=1}^K e \left(\ell_1 \left(ky + \frac{k(k-1)\omega}{2} \right) + \ell_2 k\omega \right) \right| \\
& \leq K^{-1} \sum_{1 \leq |\ell_2| \leq p_2} \left(\sum_{1 \leq |\ell_1| \leq p_1} |\hat{v}(\ell_1, \ell_2)|^2 \right)^{\frac{1}{2}} \left(\sum_{1 \leq |\ell_1| \leq p_1} \left| \sum_{k=1}^K e \left(\ell_1 \left(ky + \frac{k(k-1)\omega}{2} \right) + \ell_2 k\omega \right) \right|^2 \right)^{\frac{1}{2}}
\end{aligned}$$

$$\begin{aligned} &\leq C(B_5 - m_5)K^{-1} \sum_{1 \leq |\ell_2| \leq p_2} \frac{1}{|\ell_2|} \left(\sum_{1 \leq |\ell_1| \leq p_1} \left| \sum_{k=1}^K e \left(\ell_1 \left(ky + \frac{k(k-1)\omega}{2} \right) + \ell_2 k \omega \right) \right|^2 \right)^{\frac{1}{2}} \\ &=: C(B_5 - m_5)S_3 \end{aligned}$$

Returning to (7.18) we conclude that

$$(7.20) \quad \sup_{(x,y) \in \mathbb{T}^2} \left| \frac{1}{K} \sum_{k=1}^K v_3 \circ T_\omega^k(x,y) - \langle v \rangle \right| \leq C(B_5 - m_5)S_1 + C(B_4 - m_4)S_2 + C(B_5 - m_5)S_3.$$

7.2.1. *Estimate of S_1 .* Applying (7.9) to S_1 , we infer that

$$(7.21) \quad \begin{aligned} S_1 &< 2 \sum_{1 \leq \ell_2 \leq p_2} \frac{1}{\ell_2} \min \left(1, \frac{1}{2K \|\ell_2 \omega\|_{\mathbb{T}}} \right) \\ &= 2 \sum_{1 \leq \ell_2 \leq p_2} \mathbf{1}_{\{\ell_2: \|\ell_2 \omega\| \leq \frac{1}{2K}\}} \frac{1}{\ell_2} \\ &\quad + 2 \sum_{j=1}^{2^j \leq 2K} \sum_{1 \leq \ell_2 \leq p_2} \mathbf{1}_{\{\ell_2: \frac{2^j-1}{2K} < \|\ell_2 \omega\|_{\mathbb{T}} \leq \frac{2^j}{2K}\}} \frac{1}{\ell_2} \min \left(1, \frac{1}{2K \|\ell_2 \omega\|_{\mathbb{T}}} \right) \\ &=: S_{1,1} + S_{1,2}. \end{aligned}$$

Estimate of $S_{1,1}$.

$$(7.22) \quad S_{1,1} = 2 \sum_{1 \leq \ell_2 \leq p_2} \mathbf{1}_{\{\ell_2: \|\ell_2 \omega\| \leq \frac{1}{2K}\}} \frac{1}{\ell_2}.$$

By Corollary 7.3, if for some $\ell \geq 1$, $\|\ell \omega\|_{\mathbb{T}} \leq \frac{1}{2K}$, then

$$(7.23) \quad \ell \geq \frac{2}{3}K.$$

Moreover, if for some distinct $\ell, \tilde{\ell} \geq 1$, $\max(\|\ell \omega\|_{\mathbb{T}}, \|\tilde{\ell} \omega\|_{\mathbb{T}}) \leq \frac{1}{2K}$, then

$$(7.24) \quad |\ell - \tilde{\ell}| \geq \frac{1}{3}K.$$

By (7.22), (7.23) and (7.24), we have

$$(7.25) \quad S_{1,1} \leq 2 \sum_{\ell=1}^{\lfloor 3p_2/K \rfloor - 1} \frac{3}{\ell+1} \frac{1}{K} < \frac{6}{K} \log \frac{3p_2}{K}.$$

For this estimate, and from this point on, we assume that $p_2 \geq K$.

Estimate of $S_{1,2}$.

$$(7.26) \quad \begin{aligned} S_{1,2} &\leq \frac{1}{K} \sum_{j=1}^{2^j \leq 2K} \sum_{1 \leq \ell_2 \leq p_2} \mathbf{1}_{\{\ell_2: \frac{2^{j-1}}{2K} < \|\ell_2 \omega\|_{\mathbb{T}} \leq \frac{2^j}{2K}\}} \frac{1}{\ell_2 \|\ell_2 \omega\|_{\mathbb{T}}} \\ &\leq 2 \sum_{j=1}^{2^j \leq 2K} \sum_{1 \leq \ell_2 \leq p_2} \mathbf{1}_{\{\ell_2: \frac{2^{j-1}}{2K} < \|\ell_2 \omega\|_{\mathbb{T}} \leq \frac{2^j}{2K}\}} \frac{1}{2^{j-1} \ell_2} \end{aligned}$$

By Corollary 7.3, if for some $\ell \geq 1$, $\|\ell \omega\|_{\mathbb{T}} \leq \frac{2^j}{2K}$, then we have

$$(7.27) \quad \ell \geq \frac{2}{2^j 3} K.$$

Moreover, if for some distinct $\ell, \tilde{\ell} \geq 1$, $\max(\|\ell \omega\|_{\mathbb{T}}, \|\tilde{\ell} \omega\|_{\mathbb{T}}) \leq \frac{2^j}{2K}$, then

$$(7.28) \quad |\ell - \tilde{\ell}| \geq \frac{1}{2^j 3} K.$$

By (7.26), (7.27) and (7.28), we have

$$(7.29) \quad \begin{aligned} S_{1,2} &\leq 2 \sum_{j=1}^{2^j \leq 2K} \sum_{\ell=1}^{\lfloor 2^j 3 p_2 / K \rfloor - 1} \frac{6}{\ell + 1} \frac{1}{K} \\ &\leq \frac{12}{K} \sum_{j=1}^{2^j \leq 2K} \log \frac{2^j 3 p_2}{K}. \\ &\leq \frac{12}{K} \left(\frac{\log 2}{2} (\log_2 K + 1)^2 + (\log_2 K + 1) \log \frac{3 p_2}{K} \right). \end{aligned}$$

Putting (7.21), (7.25) and (7.29) together, we have

$$(7.30) \quad S_1 \leq \frac{6}{K} \left(\log 2 (\log_2 K + 2)^2 + (2 \log_2 K + 3) \log \frac{3 p_2}{K} \right).$$

Henceforth we assume $p_2 \geq K > 23$, which allows us to simplify (7.30) into

$$(7.31) \quad S_1 \leq 26 \frac{\log K}{K} \log p_2.$$

7.2.2. Estimate of S_2 and S_3 . In order to estimate S_2 and S_3 , we use the well-known method of Weyl-differencing, cf. for example [Mon]. As a first step,

$$\begin{aligned} &\left| \sum_{k=1}^K e \left(\ell_1 \left(ky + \frac{k(k-1)\omega}{2} \right) + \ell_2 k \omega \right) \right|^2 \\ &= \left(\sum_{k=1}^K e \left(\ell_1 \left(ky + \frac{k(k-1)\omega}{2} \right) + \ell_2 k \omega \right) \right) \left(\sum_{j=1}^K e \left(-\ell_1 \left(jy + \frac{j(j-1)\omega}{2} \right) - \ell_2 j \omega \right) \right) \\ &= \sum_{j,k=1}^K e \left(\ell_1 \left((y - \frac{\omega}{2})(k-j) + \frac{\omega}{2}(k^2 - j^2) \right) + \ell_2 \omega(k-j) \right) \end{aligned}$$

Let $\ell = k + j$ and $m = k - j$, hence $\ell \equiv m \pmod{2}$. Let us denote the sum over even integers j 's by $\sum_{\mathcal{E},j}$, and odd integers by $\sum_{\mathcal{O},j}$.

$$\begin{aligned}
& \left| \sum_{k=1}^K e \left(\ell_1 \left(ky + \frac{k(k-1)\omega}{2} \right) + \ell_2 k\omega \right) \right|^2 \\
(7.32) \quad &= \sum_{m=1-K}^{K-1} \sum_{\ell=2+|m|}^{2K-|m|} e \left(\ell_1 m \left(y + \frac{(\ell-1)\omega}{2} \right) + \ell_2 m\omega \right) \\
& \quad + \sum_{m=1-K}^{K-1} \sum_{\ell=2+|m|}^{2K-|m|} e \left(\ell_1 m \left(y + \frac{(\ell-1)\omega}{2} \right) + \ell_2 m\omega \right),
\end{aligned}$$

In which by (7.9), with $m = 2\tilde{m}$ and $\ell = 2\tilde{\ell}$,

$$\begin{aligned}
& \left| \sum_{m=1-K}^{K-1} \sum_{\ell=2+|m|}^{2K-|m|} e \left(\ell_1 m \left(y + \frac{(\ell-1)\omega}{2} \right) + \ell_2 m\omega \right) \right| \\
(7.33) \quad &= \left| \sum_{\tilde{m}=-\lfloor \frac{K-1}{2} \rfloor}^{\lfloor \frac{K-1}{2} \rfloor} \sum_{\tilde{\ell}=1+|\tilde{m}|}^{K-|\tilde{m}|} e \left(\ell_1 \tilde{m} \left(2y + (2\tilde{\ell}-1)\omega \right) + 2\ell_2 \tilde{m}\omega \right) \right| \\
& \leq K + 2 \sum_{\tilde{m}=1}^{\lfloor \frac{K-1}{2} \rfloor} \min \left(K - 2\tilde{m}, \frac{1}{2\|2\ell_1 \tilde{m}\omega\|_{\mathbb{T}}} \right) \\
& \leq K + 2K \sum_{\tilde{m}=1}^{\lfloor \frac{K-1}{2} \rfloor} \min \left(1, \frac{1}{2K\|2\ell_1 \tilde{m}\omega\|_{\mathbb{T}}} \right),
\end{aligned}$$

and with $m = 2\tilde{m} - 1$, $\ell = 2\tilde{\ell} - 1$,

$$\begin{aligned}
& \left| \sum_{m=1-K}^{K-1} \sum_{\ell=2+|m|}^{2K-|m|} e \left(\ell_1 m \left(y + \frac{(\ell-1)\omega}{2} \right) + \ell_2 m\omega \right) \right| \\
(7.34) \quad & \leq 2 \sum_{m=1}^{K-1} \left| \sum_{\ell=2+|m|}^{2K-|m|} e \left(\ell_1 m \left(y + \frac{(\ell-1)\omega}{2} \right) + \ell_2 m\omega \right) \right|
\end{aligned}$$

which is further equal to

$$\begin{aligned}
& 2 \sum_{\tilde{m}=1}^{\lfloor \frac{K}{2} \rfloor} \left| \sum_{\tilde{\ell}=\tilde{m}+1}^{K-\tilde{m}+1} e \left(\ell_1(2\tilde{m}-1) \left(y + \tilde{\ell}\omega \right) + \ell_2(2\tilde{m}-1)\omega \right) \right| \\
& \leq 2 \sum_{\tilde{m}=1}^{\lfloor \frac{K}{2} \rfloor} \min \left(K - 2\tilde{m} + 1, \frac{1}{2\|2\ell_1\tilde{m}\omega\|_{\mathbb{T}}} \right) \\
& \leq 2K \sum_{\tilde{m}=1}^{\lfloor \frac{K}{2} \rfloor} \min \left(1, \frac{1}{2K\|2\ell_1\tilde{m}\omega\|_{\mathbb{T}}} \right).
\end{aligned}$$

Plugging the estimates of (7.33) and (7.34) into (7.32), yields

$$\left| \sum_{k=1}^K e \left(\ell_1 \left(ky + \frac{k(k-1)\omega}{2} \right) + \ell_2 k\omega \right) \right|^2 \leq K + 4K \sum_{m=1}^{\lfloor \frac{K}{2} \rfloor} \min \left(1, \frac{1}{2K\|2\ell_1 m\omega\|_{\mathbb{T}}} \right).$$

Hence we have

$$\begin{aligned}
(7.35) \quad S_2 & \leq \frac{2}{K} \left(2p_1K + 4K \sum_{1 \leq |\ell_1| \leq p_1} \sum_{m=1}^{\lfloor \frac{K}{2} \rfloor} \min \left(1, \frac{1}{2K\|2\ell_1 m\omega\|_{\mathbb{T}}} \right) \right)^{\frac{1}{2}} \\
& \leq \frac{2}{K} \left(2p_1K + 8K \sum_{\ell_1=1}^{p_1} \sum_{m=1}^{\lfloor \frac{K}{2} \rfloor} \min \left(1, \frac{1}{2K\|2\ell_1 m\omega\|_{\mathbb{T}}} \right) \right)^{\frac{1}{2}} \\
& \leq \frac{2}{K} \left(2p_1K + 8KC_{p_1K}^* \sum_{j=1}^{p_1K} \min \left(1, \frac{1}{2K\|j\omega\|_{\mathbb{T}}} \right) \right)^{\frac{1}{2}}.
\end{aligned}$$

and similarly

$$(7.36) \quad S_3 \leq \frac{2}{K} (\log(p_2) + 1) \left(2p_1K + 8KC_{p_1K}^* \sum_{j=1}^{p_1K} \min \left(1, \frac{1}{2K\|j\omega\|_{\mathbb{T}}} \right) \right)^{\frac{1}{2}}.$$

The constant $C_{p_1K}^*$ comes from over-counting.

Next, we will need to bound $C_{p_1K}^*$ and $\sum_{j=1}^{p_1K} \min \left(1, \frac{1}{2K\|j\omega\|_{\mathbb{T}}} \right)$ separately.

Estimate of $C_{p_1K}^$.* We first note the following simple bound on $C_{p_1K}^*$

$$(7.37) \quad C_{p_1K}^* \leq \min(p_1, \tau^*(p_1K)),$$

where $\tau^*(p_1K) := \max_{1 \leq n \leq p_1K} \tau(n)$ with $\tau(n)$ be the divisor function of n . Standard divisor bound yields the following estimates

Lemma 7.7.

$$(7.38) \quad \tau^*(p_1 K) \leq \begin{cases} (p_1 K)^{\frac{1.06602}{\log \log(p_1 K)}}, \\ C(\epsilon)(p_1 K)^\epsilon. \end{cases}$$

The second inequality above holds for any integer $p_1 K \geq 1$ with explicit constants $C(\frac{1}{2}) = 2$ and $C(\frac{1}{8}) = 42000$. It also holds for $p_1 K \leq 327680000$ with constant $C(\frac{1}{50}) = 702$.

Combining (7.37) with Lemma 7.7, we have that for any $0 \leq \alpha \leq 1$,

$$(7.39) \quad C_{p_1 K}^* \leq (C(\epsilon))^\alpha (p_1 K)^{\alpha\epsilon} p_1^{1-\alpha}.$$

We will only use the case when $\alpha = 0$, but we keep this as a reference for the sake of completeness.

Estimate of $\sum_{\ell=1}^{p_1 K} \min\left(1, \frac{1}{2K\|\ell\omega\|_{\mathbb{T}}}\right)$.

In analogy with (7.21), we will split the term $\sum_{\ell=1}^{p_1 K} \min\left(1, \frac{1}{2K\|\ell\omega\|_{\mathbb{T}}}\right)$ appearing in (7.36) as follows:

$$(7.40) \quad \begin{aligned} & \sum_{\ell=1}^{p_1 K} \min\left(1, \frac{1}{2K\|\ell\omega\|_{\mathbb{T}}}\right) \\ &= \sum_{\ell=1}^{p_1 K} \mathbb{1}_{\{\ell: \|\ell\omega\|_{\mathbb{T}} < \frac{1}{2K}\}} + \sum_{j=1}^{2^j < 2K} \sum_{\ell=1}^{p_1 K} \mathbb{1}_{\{\ell: \frac{2^{j-1}}{2K} \leq \|\ell\omega\|_{\mathbb{T}} < \frac{2^j}{2K}\}} \frac{1}{2K\|\ell\omega\|_{\mathbb{T}}} \\ &\leq \sum_{\ell=1}^{p_1 K} \mathbb{1}_{\{\ell: \|\ell\omega\|_{\mathbb{T}} < \frac{1}{2K}\}} + \sum_{j=1}^{2^j < 2K} \sum_{\ell=1}^{p_1 K} \mathbb{1}_{\{\ell: \frac{2^{j-1}}{2K} \leq \|\ell\omega\|_{\mathbb{T}} < \frac{2^j}{2K}\}} \frac{1}{2^{j-1}} \\ &=: S_4 + S_5. \end{aligned}$$

By Corollary 7.3, if for some $\ell \geq 1$, $\|\ell\omega\|_{\mathbb{T}} < \frac{2^j}{2K}$, then

$$(7.41) \quad \ell \geq \frac{2}{2^j 3} K.$$

By Corollary 7.3, if for some distinct $\ell, \tilde{\ell} \geq 1$, $\max(\|\ell\omega\|_{\mathbb{T}}, \|\tilde{\ell}\omega\|_{\mathbb{T}}) < \frac{2^j}{K}$, then

$$(7.42) \quad |\ell - \tilde{\ell}| \geq \frac{1}{2^j 3} K.$$

Combining (7.41) with (7.42), we have

$$(7.43) \quad S_4 \leq 3p_1,$$

and

$$(7.44) \quad S_5 \leq \sum_{j=1}^{2^j < 2K} 6p_1 \leq 6(\log_2 K + 1)p_1.$$

In view of (7.35), (7.36), (7.40), (7.43) and (7.44) together, one has

$$\begin{aligned}
(7.45) \quad S_2 &< \frac{2}{K} (2p_1K + 24p_1K C_{p_1K}^* (2 \log_2 K + 3))^{\frac{1}{2}} \\
&< 8\sqrt{3} (C_{p_1K}^*)^{\frac{1}{2}} p_1^{\frac{1}{2}} K^{-\frac{1}{2}} (\log_2 K + 2)^{\frac{1}{2}} \\
&< 20 (C_{p_1K}^*)^{\frac{1}{2}} p_1^{\frac{1}{2}} K^{-\frac{1}{2}} (\log K)^{\frac{1}{2}}, \quad \text{for } K \geq 23,
\end{aligned}$$

and

$$\begin{aligned}
(7.46) \quad S_3 &< 8\sqrt{3} (C_{p_1K}^*)^{\frac{1}{2}} (\log(p_2) + 1) p_1^{\frac{1}{2}} K^{-\frac{1}{2}} (\log_2 K + 2)^{\frac{1}{2}} \\
&< 25 (\log p_2) (C_{p_1K}^*)^{\frac{1}{2}} p_1^{\frac{1}{2}} K^{-\frac{1}{2}} (\log K)^{\frac{1}{2}}, \quad \text{for } p_2 \geq K \geq 38.
\end{aligned}$$

7.2.3. *Combining S_1, S_2, S_3 .* Taking $p_1 = \lfloor K^{\delta_1} \rfloor$ and $p_2 = \lfloor e^{4(\log K)^{\delta_2}} \rfloor$, the estimate of S_1 , namely (7.31), becomes

$$(7.47) \quad S_1 < 105K^{-1} (\log K)^{\delta_2+1}.$$

Recall from the preceding that we impose the conditions $K \geq 38$ and $\exp(4(\log K)^{\delta_2}) \geq K + 1$, note that these are our assumptions (ii) and (iii). The estimate of S_2 , (7.45), becomes

$$(7.48) \quad S_2 < 20 (C_{K^{1+\delta_1}}^*)^{\frac{1}{2}} K^{-\frac{1-\delta_1}{2}} (\log K)^{\frac{1}{2}}.$$

The estimate of S_3 , (7.46), becomes

$$(7.49) \quad S_3 < 100 (C_{K^{1+\delta_1}}^*)^{\frac{1}{2}} K^{-\frac{1-\delta_1}{2}} (\log K)^{\delta_2+\frac{1}{2}}.$$

Combining (7.47), (7.48), (7.49), (7.20) with our assumption (i) that $C(B_5 - m_5) \leq K^\delta$, yields

$$\begin{aligned}
(7.50) \quad &\sup_{(x,y) \in \mathbb{T}^2} \left| \frac{1}{K} \sum_{k=1}^K v_3 \circ T_\omega^k(x, y) - \langle v \rangle \right| \\
&< 105K^{-1+\delta} (\log K)^{\delta_2+1} + 20C(B_4 - m_4) (C_{K^{1+\delta_1}}^*)^{\frac{1}{2}} K^{-\frac{1-\delta_1}{2}} (\log K)^{\frac{1}{2}} \\
&+ 100 (C_{K^{1+\delta_1}}^*)^{\frac{1}{2}} K^{-\frac{1-\delta_1}{2}+\delta} (\log K)^{\delta_2+\frac{1}{2}}.
\end{aligned}$$

By (7.39), with $\alpha = 0$, we have

$$\begin{aligned}
&\sup_{(x,y) \in \mathbb{T}^2} \left| \frac{1}{K} \sum_{k=1}^K v_3 \circ T_\omega^k(x, y) - \langle v \rangle \right| \\
&< 105K^{-1+\delta} (\log K)^{\delta_2+1} + 20C(B_4 - m_4) K^{-\frac{1-2\delta_1}{2}} (\log K)^{\frac{1}{2}} \\
&+ 100K^{-\frac{1-2\delta_1}{2}+\delta} (\log K)^{\delta_2+\frac{1}{2}}.
\end{aligned}$$

By condition (iv) in our statement of the proposition, we have

$$21K^{-\frac{1}{2}+\delta-\delta_1} (\log K)^{\delta_2+\frac{1}{2}} + 4C(B_4 - m_4) \leq K^\delta (\log K)^{\delta_2}.$$

which implies

$$(7.51) \quad \sup_{(x,y) \in \mathbb{T}^2} \left| \frac{1}{K} \sum_{k=1}^K v_3 \circ T_\omega^k(x, y) - \langle v \rangle \right| < 105K^{-\frac{1-2\delta_1}{2} + \delta} (\log K)^{\delta_2 + \frac{1}{2}} =: \varepsilon_0.$$

Let

$$(7.52) \quad t = C(B_4 - m_4)K^{-\frac{1}{2}\delta_1}$$

in (7.17), then for any $y \notin \mathcal{A}(t)$, with (7.14) we have,

$$(7.53) \quad \begin{aligned} \left\| \frac{1}{K} \sum_{k=1}^K (\tilde{v}_1 + v_2) \circ T_\omega^k(\cdot, y) \right\|_{L^1(\mathbb{T})} &\leq \sqrt{2}C(B_4 - m_4)K^{-\frac{1}{2}\delta_1} + t \\ &= (\sqrt{2} + 1)C(B_4 - m_4)K^{-\frac{1}{2}\delta_1} =: \varepsilon_1. \end{aligned}$$

Recall that $v = \tilde{v}_1 + v_2 + v_3$. For any fixed $y \notin \mathcal{A}(t)$, consider the subharmonic function

$$v_y(z) := \frac{1}{K} \sum_{k=1}^K v \circ T_\omega^k(z, y) \quad \text{with } z \in D_R.$$

This subharmonic function is going to satisfy the following bounds

$$v_y(z) \leq B_4 \text{ for } \forall z \in D_R, \quad \text{and } v_y(0) \geq m_4.$$

By (7.51) and (7.53), we know $v_y(x) - \langle v \rangle$ can be decomposed into two parts, one with small L^∞ norm ε_0 , the other with small L^1 norm ε_1 . We will choose δ_1 such that $\varepsilon_0 \sim \sqrt{\varepsilon_1}$, in the sense that

$$(7.54) \quad K^{-\frac{1-2\delta_1}{2} + \delta} (\log K)^{\delta_2 + \frac{1}{2}} = K^{-\frac{1}{4}\delta_1},$$

which yields

$$(7.55) \quad K^{\delta_1} = \frac{K^{\frac{2}{5} - \frac{4\delta}{5}}}{(\log K)^{\frac{2}{5} + \frac{4\delta_2}{5}}}, \quad \text{with } 0 < \delta_1 < \frac{2}{5} - \frac{4\delta}{5}.$$

Then

$$(7.56) \quad \varepsilon_0 \leq 105K^{-\frac{1}{10} + \frac{\delta}{5}} (\log K)^{\frac{1}{10} + \frac{\delta_2}{5}},$$

and

$$(7.57) \quad \varepsilon_1 = (\sqrt{2} + 1)C(B_4 - m_4)K^{-\frac{1}{5} + \frac{2\delta}{5}} (\log K)^{\frac{1}{5} + \frac{2\delta_2}{5}}.$$

Applying Corollary 3.2 to $v_y - \langle v \rangle$, we obtain that for

$$(7.58) \quad \varepsilon_4 = C_2 K^{-\frac{1}{10} + \frac{\delta}{5}} (\log K)^{\frac{1}{10} + \frac{\delta_2}{5} + \delta_3}$$

with some constant $C_2 > 0$,

$$(7.59) \quad \begin{aligned} \sup_{y \notin \mathcal{A}(t)} \left| \left\{ x \in \mathbb{T} : \left| \frac{1}{K} \sum_{k=1}^K v \circ T_\omega^k(x, y) - \langle v \rangle \right| > \varepsilon_4 \right\} \right| \\ \leq 2\sqrt{2} \exp \left(\frac{\pi}{4} \left[\frac{17}{36} + \frac{B_1}{4B_3^2} - \varepsilon_2 \delta_0^{-1} \right] \right), \end{aligned}$$

where

$$\delta_0 = \left(472.5 + 2B_3(B_4 - m_4)\sqrt{(\sqrt{2} + 1)C} \right) K^{-\frac{1}{10} + \frac{\delta}{5}} (\log K)^{\frac{1}{10} + \frac{\delta_2}{5}},$$

and hence

$$\begin{aligned} \varepsilon_4 \delta_0^{-1} &= C_2 \left(472.5 + 2B_3(B_4 - m_4)\sqrt{(\sqrt{2} + 1)C} \right)^{-1} (\log K)^{\delta_3} \\ &\geq C_2 \left(472.5 + 3.2B_3(B_4 - m_4)\sqrt{C} \right)^{-1} (\log K)^{\delta_3} =: \varepsilon_5. \end{aligned}$$

(7.59) together (7.52) and (7.17) imply

$$\begin{aligned} &\left| \left\{ (x, y) \in \mathbb{T}^2 : \left| \frac{1}{K} \sum_{k=1}^K v \circ T_\omega^k(x, y) - \langle v \rangle \right| > \varepsilon_4 \right\} \right| \\ &\leq 2\sqrt{2} \exp\left(\frac{\pi}{4} \left[\frac{17}{36} + \frac{B_1}{4B_3^2} - \varepsilon_5 \right]\right) + |\mathcal{A}(t)| \\ &\leq 2\sqrt{2} \exp\left(\frac{\pi}{4} \left[\frac{17}{36} + \frac{B_1}{4B_3^2} - \varepsilon_5 \right]\right) \\ &\quad + \sqrt{2}(C(B_4 - m_4))^{-1} K^{\frac{1}{5} - \frac{2\delta}{5}} (\log K)^{-\frac{1}{5} - \frac{2\delta_2}{5}} \exp\left(-2(\log K)^{\delta_2}\right), \end{aligned}$$

as claimed. \square

8. MULTI-SCALE ESTIMATES

In this section we commence with the inductive arguments in our multi-scale Lyapunov exponent machinery. In analogy with [GolSch, Bou1] we proceed by combining the large deviation estimates with the avalanche principle. We begin with the basic induction step, which provides a lower bound for the Lyapunov exponent at a large scale from information on the Lyapunov exponents at smaller scales, in combination with level-set estimates. In Proposition 8.4, which is the main result of this section, we will also invoke the quantitative control on the Birkhoff averages over the skew shift from the previous section in order to derive large deviation estimates at the larger scale.

The following subsection will serve as an abstract multi-scale scheme to providing a lower bound of the (maximal) Lyapunov exponent, assuming large deviation estimates. In our application to the skew shift, the large deviation estimates will come from Proposition 7.1, see Section 8.2.

8.1. Abstract multi-scale scheme.

8.1.1. *Lyapunov exponent.* Let (X, μ, S) be an ergodic dynamical system. A linear cocycle over (X, μ, S) is a skew-product map

$$F_A : X \times \mathbb{R}^d \rightarrow X \times \mathbb{R}^d,$$

given by

$$X \times \mathbb{R}^d \ni (x, v) \rightarrow (Sx, A(x)v) \in X \times \mathbb{R}^d,$$

where

$$A : X \rightarrow \mathrm{SL}_d(\mathbb{R})$$

is a measurable function.

The forward iterates F_A^n of a linear cocycle F_A are given by $F_A^n(x, v) = (S^n x, M_n(x)v)$, where

$$M_n(x) := A(S^{n-1}x) \cdots A(Sx)A(x) \quad (n \in \mathbb{N}).$$

A linear cocycle A is said to be μ -integrable if

$$\int_X \log \|A(x)\| \, d\mu < +\infty.$$

Due to the fact that norms are sub-multiplicative with respect to matrix products, the sequence of functions $\log \|A^{(n)}(x)\|$ are subadditive. The Fürstenberg-Kesten theorem (or Kingman's ergodic theorem) implies that for a μ -integrable linear cocycle, the following μ -a.e. limit exists

$$L(A) := \lim_{n \rightarrow \infty} \frac{1}{n} \log \|M_n(x)\|,$$

and it is called the (maximal) Lyapunov exponent of A . Moreover,

$$L(A) := \lim_{n \rightarrow \infty} \int_X \frac{1}{n} \log \|M_n(x)\| \, \mu(dx) = \inf_{n \geq 1} \int_X \frac{1}{n} \log \|M_n(x)\| \, \mu(dx).$$

We point out the since $A \in \mathrm{SL}_d(\mathbb{R})$, we have $\|M_n(x)\| \geq 1$, hence $L(A) \geq 0$.

8.1.2. Inductive scheme.

Let us denote

$$L_n(A) := \int_X \frac{1}{n} \log \|M_n(x)\| \, d\mu(x).$$

For simplicity, we may omit the dependence of $L(A), L_n(A)$ on A , and simply write L and L_n .

Let us further assume that there exists a constant $C_3 > 0$, such that

$$(8.1) \quad \frac{1}{n} \log \|M_n(x)\| \leq C_3 < +\infty,$$

for μ -a.e. x , uniformly in n . We point out that in our application to the skew-shift model, C_3 can be taken as $U(\lambda, 1)$, see (6.5) and (6.10).

Definition 8.1. In our multi-scale scheme, we quantify the failure of the Fürstenberg-Kesten theorem via the following sets \mathcal{B}_n :

$$\mathcal{B}_n := \left\{ x \in X : \left| \frac{1}{n} \log \|M_n(x)\| - L_n \right| > \frac{1}{10} L_n \right\}.$$

The lemma below shows how to inductively obtain estimates of L_N at a larger scale N , based on information at a smaller scale n . The key ingredient is the Avalanche Principle, Theorem 5.5.

Lemma 8.2. *Let $n, N/n \in \mathbb{N}$ be positive integers, and $\delta \in (0, 1/2)$. Let C_3 be as in (8.1). Assume the following three conditions:*

(a).

$$nL_n \geq 7,$$

(b).

$$L_n - L_{2n} \leq \frac{1}{8}L_n,$$

(c).

$$\max(\mu(\mathcal{B}_n), \mu(\mathcal{B}_{2n})) \leq N^{-\frac{12}{5} + \frac{4\delta}{5}}.$$

Then we have

$$(8.2) \quad L_N \geq L_n - \left(2 - \frac{2n}{N}\right)(L_n - L_{2n}) - \frac{11}{n}e^{-\frac{1}{2}nL_n} - 8C_3N^{-\frac{7}{5} + \frac{4\delta}{5}},$$

and

$$(8.3) \quad L_N - L_{2N} \leq \frac{n}{N}(L_n - L_{2n}) + \frac{22}{n}e^{-\frac{1}{2}nL_n} + 24C_3N^{-\frac{7}{5} + \frac{4\delta}{5}}.$$

8.1.3. Multi-scale scheme.

Lemma 8.3 below shows how information on a sequence of larger and larger scales determines the limit L .

Lemma 8.3. *Let $\delta \in (0, 1/2)$ be a constant, and C_3 be as in (8.1). Let $\{N_m\}_{m=0}^\infty \in \mathbb{N}$ be a sequence of positive integers, such that $10 \leq N_m/N_{m-1} \in \mathbb{N}$ for $1 \leq m$. Assume that the following hold for an integer $j \geq 0$ (note that (2)-(4) below are empty conditions for $j = 0$):*

(1)

$$N_0L_{N_0} \geq 7, \quad \text{and} \quad L_{N_0} - L_{2N_0} \leq \frac{1}{8}L_{N_0},$$

(2)

$$\sum_{m=0}^{j-1} \frac{1}{N_m} e^{-\frac{1}{2}N_mL_{N_m}} < \frac{1}{512}L_{N_0},$$

(3)

$$\sum_{m=1}^j N_m^{-\frac{7}{5} + \frac{4\delta}{5}} < \frac{1}{1280C_3}L_{N_0},$$

(4)

$$\max(\mu(\mathcal{B}_{N_m}), \mu(\mathcal{B}_{2N_m})) \leq N_{m+1}^{-\frac{12}{5} + \frac{4\delta}{5}}, \quad \text{for } 0 \leq m \leq j-1.$$

Then we have the following four estimates for $j \geq 0$.

First,

$$(8.4) \quad \begin{aligned} L_{N_j} &\geq L_{N_0} - \left(2 - \frac{2N_0}{N_j}\right) (L_{N_0} - L_{2N_0}) \\ &\quad - \sum_{m=1}^j \left(\frac{11}{N_{m-1}} e^{-\frac{1}{2}N_{m-1}L_{N_{m-1}}} + 8C_3 N_m^{-\frac{7}{5} + \frac{4\delta}{5}} \right) \\ &\quad - \sum_{m=1}^{j-1} \left(2 - \frac{2N_m}{N_j}\right) \left(\frac{22}{N_{m-1}} e^{-\frac{1}{2}N_{m-1}L_{N_{m-1}}} + 24C_3 N_m^{-\frac{7}{5} + \frac{4\delta}{5}} \right), \end{aligned}$$

in which $\sum_{m=1}^0 = \sum_{m=1}^{-1} := 0$.

Second,

$$(8.5) \quad L_{N_j} - L_{2N_j} \leq \frac{N_0}{N_j} (L_{N_0} - L_{2N_0}) + \sum_{m=1}^j \frac{N_m}{N_j} \left(\frac{22}{N_{m-1}} e^{-\frac{1}{2}N_{m-1}L_{N_{m-1}}} + 24C_3 N_m^{-\frac{7}{5} + \frac{4\delta}{5}} \right),$$

in which $\sum_{m=1}^0 := 0$.

Third,

$$(8.6) \quad L_{N_j} - L_{2N_j} \leq \frac{1}{8} L_{N_j}$$

Fourth,

$$(8.7) \quad L_{2N_j} \geq \frac{1}{2} L_{N_0}, \quad \text{and} \quad N_j L_{N_j} \geq 7.$$

8.2. Application to the skew-shift model. The two cornerstones of the abstract multi-scale scheme are:

- Initial scale N_0 estimates, including (1). $N_0 L_{N_0} \geq 7$, (2). $L_{N_0} - L_{2N_0} \leq \frac{1}{8} L_{N_0}$ and (3). Large deviation estimates of $\mu(\mathcal{B}_{N_0})$ and $\mu(\mathcal{B}_{2N_0})$.
- Large deviation estimates of $\mu(\mathcal{B}_{N_j})$ and $\mu(\mathcal{B}_{2N_j})$ for $j \geq 1$.

In this subsection, we will present a machinery that inductively provides large deviation estimates for scales N_j , $j \geq 1$, thus reducing the problem to the initial scale only. The key ingredients are the Avalanche Principle and the quantitative control of the ergodic averages of plurisubharmonic functions over a skew-shift orbit, Proposition 7.1.

Let us recall some notations from Section 6: $u_n(\lambda, E; x, y) = \frac{1}{n} \log \|M_n(\lambda, E; x, y)\|$, and $v_n(\lambda, E; z, w)$ be the complexification of u_n from \mathbb{T}^2 to \mathbb{C}^2 , as in (6.5). The constant $U(\lambda, 1)$, as in (6.10), is a uniform (in n and E) L^∞ upper bound on $u_n(\lambda, E; x, y)$. For simplicity, we will omit the dependence of $u_n(x, y)$, $v_n(z, w)$, $\log \|M_n(x, y)\|$ and L_n on λ, E , since λ will be fixed, and our estimates are uniform in $E \in [-2 - 2\lambda, 2 + 2\lambda]$. Recall also from Lemma 6.1 with $R_3 = R$ that the bounds with respect to v_n satisfy

$$(8.8) \quad \begin{aligned} B_4 - m_4 &= 2 \log R + U(\lambda, R) - \log \lambda, \\ B_5^{(n)} - m_5^{(n)} &= (n+1) \log R + U(\lambda, R) - \log \lambda. \end{aligned}$$

Let us finally also recall the constants $B_3(R, R_1, R_2)$ as in (2.2), $C(R, R_1, R_2)$ as in (3.47), and $C_0(R, R_1, R_2)$ is as in (3.2). In the following we will write B_3, C, C_0 for simplicity.

Proposition 8.4. *Let $\omega = \frac{\sqrt{5}-1}{2}$ be the golden ratio. Let $\delta \in (0, 1/2)$ and*

$$\delta_2, \delta_3, \delta_4, C_2, C_4, C_5 > 0$$

be constants. Let $n, N \in \mathbb{N}$ be positive integers and assume that n divides N . In addition to the conditions (a)-(c) in Lemma 8.2 and Definition 3.5, assume further that the following properties hold for both $\tilde{N} = N$ and $2N$:

- (I). $C((2n+1)\log R + U(\lambda, R) - \log \lambda) \leq \tilde{N}^\delta,$
- (II). $\tilde{N} \geq 38,$
- (III). $\exp\left(4(\log \tilde{N})^{\delta_2}\right) \geq \tilde{N} + 1,$
- (IV). $21\tilde{N}^{-\frac{9}{10} + \frac{9}{5}\delta}(\log \tilde{N})^{\frac{9}{10} + \frac{9}{5}\delta_2} + 4C(B_4 - m_4) \leq \tilde{N}^\delta(\log \tilde{N})^{\delta_2},$
- (V). $2n\tilde{N}^{-1}(L_n - L_{2n}) + 8U(\lambda, 1)\tilde{N}^{-\frac{7}{5} + \frac{4\delta}{5}} + 5U(\lambda, 1)n\tilde{N}^{-1} < C_2\tilde{N}^{-\frac{1}{10} + \frac{\delta}{5}}(\log \tilde{N})^{\frac{1}{10} + \frac{\delta_2}{5} + \delta_3},$
- (VI). $22n^{-1}\exp(-nL_n/2) < C_2\tilde{N}^{-\frac{1}{10} + \frac{\delta}{5}}(\log \tilde{N})^{\frac{1}{10} + \frac{\delta_2}{5} + \delta_3},$
- (VII). $4\sqrt{2}\exp\left(\frac{\pi}{4}\left[\frac{17}{36} + \frac{B_1}{4B_3^2} - C_2(472.5 + 3.2B_3(B_4 - m_4)\sqrt{C})^{-1}(\log \tilde{N})^{\delta_3}\right]\right) \leq \tilde{N}^{-\frac{7}{5} + \frac{4\delta}{5}},$
- (VIII). $2\sqrt{2}(C(B_4 - m_4))^{-1}\tilde{N}^{\frac{1}{5} - \frac{2\delta}{5}}(\log \tilde{N})^{-\frac{1}{5} - \frac{2\delta_2}{5}}\exp\left(-2(\log \tilde{N})^{\delta_2}\right) \leq \tilde{N}^{-\frac{7}{5} + \frac{4\delta}{5}},$
- (IX). $\tilde{N} > (\log R + U(\lambda, R) - \log \lambda)(\log R)^{-1},$
- (X). $C_4(\log \tilde{N})^{\delta_4} > 4,$
- (XI). $C_5(\log \tilde{N})^{\delta_4} > C_4.$

Then the following holds for both $\tilde{N} = N$ and $2N$:

$$\begin{aligned} & \left| \left\{ (x, y) \in \mathbb{T}^2 : |v_{\tilde{N}}(x, y) - L_{\tilde{N}}| > C_2C_5\tilde{N}^{-\frac{1}{10} + \frac{\delta}{5}}(\log \tilde{N})^{\frac{1}{10} + \frac{\delta_2}{5} + \delta_3 + 2\delta_4} \right\} \right| \\ & \leq 2(2C_0)^{\frac{1}{2}} \exp\left(\frac{-\pi C_2C_4(\log \tilde{N})^{\delta_4}}{144C_2 + 48B_3\sqrt{2}U(\lambda, 1)(B_4 - m_4)(\log \tilde{N})^{-\frac{1}{10} - \frac{\delta_2}{5} - \delta_3}}\right) \\ & \quad + C_0 \exp\left(\frac{-\pi C_2C_5(\log \tilde{N})^{\delta_4}}{18C_2C_4 + 96B_3U(\lambda, 1)\sqrt{\log \tilde{R}}(\log \tilde{N})^{-\frac{1}{10} - \frac{\delta_2}{5} - \delta_3 - \delta_4}}\right). \end{aligned}$$

Remark 8.5. Note that our conditions (I)-(IV) correspond to (i)-(iv) of Proposition 7.1. In particular, (I) is (i) of Proposition 7.1 with $B_5^{(2n)} - m_5^{(2n)}$ given in (8.8).

8.3. Proofs.

Before proving Lemma 8.2 and Proposition 8.4, we will first give a quick proof of Lemma 8.3 based on Lemma 8.2.

Proof of Lemma 8.3. For $m \geq 1$, let us denote

$$\begin{aligned} \alpha_m & := 11N_{m-1}^{-1}e^{-\frac{1}{2}N_{m-1}L_{N_{m-1}}} + 8C_3N_m^{-\frac{7}{5} + \frac{4\delta}{5}}, \\ \beta_m & := 22N_{m-1}^{-1}e^{-\frac{1}{2}N_{m-1}L_{N_{m-1}}} + 24C_3N_m^{-\frac{7}{5} + \frac{4\delta}{5}}. \end{aligned}$$

Note that in terms of α and β , our conditions (2) and (3) in the statement of the lemma become

$$(8.9) \quad \sum_{m=1}^j \alpha_m \leq \frac{11}{512} L_{N_0} + \frac{8}{1280} L_{N_0} = \frac{71}{2560} L_{N_0},$$

and

$$(8.10) \quad \sum_{m=1}^j \beta_m \leq \frac{22}{512} L_{N_0} + \frac{24}{1280} L_{N_0} = \frac{79}{1280} L_{N_0}.$$

Our proof is based on induction on j . Note that for the induction base case $j = 0$: (8.5), (8.6) and (8.7) follow directly from condition (1). (8.4) follows from the fact that $L_{N_0} - L_{2N_0} \geq 0$.

Now let us suppose Lemma 8.3 holds for $j = J$ for some $J \geq 0$. Note that conditions (2)-(4) with $j = J + 1$ already imply those with $j = J$. Hence by our inductive assumption, (8.4), (8.5), (8.6) and (8.7) hold for $j = J$, whence

$$(8.11) \quad L_{N_J} \geq L_{N_0} - \left(2 - \frac{2N_0}{N_J}\right) (L_{N_0} - L_{2N_0}) - \sum_{m=1}^J \alpha_m - \sum_{m=1}^{J-1} \left(2 - \frac{2N_m}{N_J}\right) \beta_m,$$

$$(8.12) \quad L_{N_J} - L_{2N_J} \leq \frac{N_0}{N_J} (L_{N_0} - L_{2N_0}) + \sum_{m=1}^J \frac{N_m}{N_J} \beta_m,$$

$$(8.13) \quad L_{N_J} - L_{2N_J} \leq \frac{1}{8} L_{N_J},$$

and

$$(8.14) \quad N_J L_{N_J} \geq 7.$$

Note that (8.13), (8.14) and our condition (4) in the statement of the lemma with $m = J$ verify the conditions of Lemma 8.2 for $n = N_J$ and $N = N_{J+1}$. Therefore Lemma 8.2 implies

$$(8.15) \quad L_{N_{J+1}} \geq L_{N_J} - \left(2 - \frac{2N_J}{N_{J+1}}\right) (L_{N_J} - L_{2N_J}) - \alpha_{J+1},$$

and

$$(8.16) \quad L_{N_{J+1}} - L_{2N_{J+1}} \leq \frac{N_J}{N_{J+1}} (L_{N_J} - L_{2N_J}) + \beta_{J+1}$$

Plugging (8.12) into (8.16), we obtain

$$(8.17) \quad \begin{aligned} L_{N_{J+1}} - L_{2N_{J+1}} &\leq \frac{N_0}{N_{J+1}} (L_{N_0} - L_{2N_0}) + \frac{N_J}{N_{J+1}} \left(\sum_{m=1}^J \frac{N_m}{N_J} \beta_m \right) + \beta_{J+1} \\ &= \frac{N_0}{N_{J+1}} (L_{N_0} - L_{2N_0}) + \sum_{m=1}^{J+1} \frac{N_m}{N_{J+1}} \beta_m, \end{aligned}$$

this proves (8.5) for $j = J + 1$.

Plugging (8.11) and (8.12) with $j = J + 1$ into (8.15), we have
(8.18)

$$\begin{aligned} L_{N_{J+1}} &\geq L_{N_0} - \left(2 - \frac{2N_0}{N_J}\right) (L_{N_0} - L_{2N_0}) - \sum_{m=1}^J \alpha_m - \sum_{m=1}^{J-1} \left(2 - \frac{2N_m}{N_J}\right) \beta_m \\ &\quad - \left(2 - \frac{2N_J}{N_{J+1}}\right) \left(\frac{N_0}{N_J} (L_{N_0} - L_{2N_0}) + \sum_{m=1}^J \frac{N_m}{N_J} \beta_m\right) - \alpha_{J+1} \\ &= L_{N_0} - \left(2 - \frac{2N_0}{N_{J+1}}\right) (L_{N_0} - L_{2N_0}) - \sum_{m=1}^{J+1} \alpha_m - \sum_{m=1}^J \left(2 - \frac{2N_m}{N_{J+1}}\right) \beta_m. \end{aligned}$$

This proves (8.4) for $j = J + 1$.

Combining (8.17), (8.18) with the fact that $10 \leq N_{j+1}/N_j$ for any $j \geq 0$, yields

$$\begin{aligned} &8(L_{N_{J+1}} - L_{2N_{J+1}}) - L_{N_{J+1}} \\ &\leq -L_{N_0} + \left(2 + \frac{6N_0}{N_{J+1}}\right) (L_{N_0} - L_{2N_0}) + \sum_{m=1}^{J+1} \alpha_m + \sum_{m=1}^{J+1} \left(2 + \frac{6N_m}{N_{J+1}}\right) \beta_m \\ &\leq -L_{N_0} + \left(2 + \frac{6}{10}\right) (L_{N_0} - L_{2N_0}) + \sum_{m=1}^{J+1} \alpha_m + 8 \sum_{m=1}^{J+1} \beta_m. \end{aligned}$$

Using (8.9) and (8.10), and the fact that $L_{N_0} - L_{2N_0} \leq \frac{1}{8}L_{N_0}$, we conclude that

$$(8.19) \quad 8(L_{N_{J+1}} - L_{2N_{J+1}}) - L_{N_{J+1}} \leq -\frac{393}{2560}L_{N_0} < 0.$$

This proves (8.6) for $j = J + 1$.

By (8.18) and the fact that $N_{m+1} \geq 10N_m$ for any $m \geq 0$, we have

$$(8.20) \quad L_{N_{J+1}} \geq L_{N_0} - \left(2 - \frac{1}{5}\right) (L_{N_0} - L_{2N_0}) - \sum_{m=1}^{J+1} \alpha_m - \left(2 - \frac{1}{5}\right) \sum_{m=1}^J \beta_m.$$

Plugging (8.9) and (8.10) with $j = J + 1$ into (8.20), and using that $L_{N_0} - L_{2N_0} \leq \frac{1}{8}L_{N_0}$, yields

$$(8.21) \quad L_{N_{J+1}} \geq \frac{8143}{12800}L_{N_0},$$

which also implies $N_J L_{N_J} \geq 7$. (8.7) with $j = J + 1$ then follows from (8.19) and (8.21), indeed,

$$L_{2N_{J+1}} \geq \frac{7}{8}L_{N_{J+1}} \geq \frac{7}{8} \times \frac{8143}{12800}L_{N_0} \geq \frac{1}{2}L_{N_0},$$

as desired. \square

Proof of Lemma 8.2. Let $\tilde{N} = N$ or $2N$. Let us define

$$(8.22) \quad \mathcal{B}^{(\tilde{N})} := \left(\bigcup_{j=0}^{\tilde{N}-1} S^{-j} \mathcal{B}_n \right) \cup \left(\bigcup_{j=0}^{\tilde{N}-n-1} S^{-j} \mathcal{B}_{2n} \right).$$

We have the following measure estimates for $\mathcal{B}^{(\tilde{N})}$ by condition (c) of the statement of the lemma,

$$(8.23) \quad \mu(\mathcal{B}^{(N)}) \leq \frac{2N - n}{N^{\frac{12}{5} - \frac{4\delta}{5}}} \leq 2N^{-\frac{7}{5} + \frac{4\delta}{5}}, \quad \text{and} \quad \mu(\mathcal{B}^{(2N)}) \leq 4N^{-\frac{7}{5} + \frac{4\delta}{5}}.$$

Taking any $x \notin \mathcal{B}^{(\tilde{N})}$, by our definitions of \mathcal{B}_n and \mathcal{B}_{2n} , we have

$$(8.24) \quad e^{\frac{11}{10}nL_n} \geq \|M_n(S^j x)\| \geq e^{\frac{9}{10}nL_n} =: \kappa^{-\frac{1}{2}}, \quad \text{for any } 0 \leq j \leq \tilde{N} - 1,$$

and

$$(8.25) \quad \|M_{2n}(S^j x)\| \geq e^{\frac{9}{5}nL_{2n}}, \quad \text{for any } 0 \leq j \leq \tilde{N} - n - 1.$$

Hence for any $0 \leq j \leq \tilde{N} - n - 1$, (8.24) and (8.25) imply that

$$\frac{\|M_{2n}(S^j x)\|}{\|M_n(S^{j+n} x)\| \|M_n(S^j x)\|} \geq \exp\left(2n(L_{2n} - L_n) - \frac{1}{5}n(L_n + L_{2n})\right) =: \epsilon.$$

We now need to verify the assumptions of the Avalanche Principle, Theorem 5.5. First, by sub-additivity of $\log \|M_n(x)\|$, we have $L_{2n} \leq L_n$. This together with our assumptions (a) and (b) yield

$$(8.26) \quad \epsilon = \exp\left(\frac{9}{5}nL_{2n} - \frac{11}{5}nL_n\right) \leq e^{-\frac{2}{5}nL_n} \leq e^{-\frac{14}{5}} < \frac{1}{10},$$

and

$$\kappa\epsilon^{-2} = \exp\left(-2nL_n + \frac{3}{5}nL_n + \frac{2}{5}nL_{2n} + 4n(L_n - L_{2n})\right) \leq e^{-\frac{1}{2}nL_n} \leq e^{-\frac{7}{2}} < \frac{1}{10}.$$

Applying Theorem 5.5 to $x \notin \mathcal{B}^{(\tilde{N})}$, we conclude that for each $0 \leq k \leq n - 1$,

$$\begin{aligned} & \left| \frac{1}{\tilde{N}} \log \|M_{\tilde{N}}(S^k x)\| + \sum_{j=1}^{\tilde{N}-2n} \log \|M_n(S^{j+n+k} x)\| \right. \\ & \quad \left. - \sum_{j=0}^{\tilde{N}-2n} \log \|M_{2n}(S^{j+n+k} x)\| \right| \leq \frac{11}{n} \kappa\epsilon^{-2} \leq \frac{11}{n} e^{-\frac{1}{2}nL_n}. \end{aligned}$$

Summing over $k \in [0, n - 1]$ and dividing by n , and finally applying the triangle inequality yields

$$(8.27) \quad \left| \frac{1}{n} \sum_{k=0}^{n-1} \frac{1}{\tilde{N}} \log \|M_{\tilde{N}}(S^k x)\| + \frac{1}{\tilde{N}} \sum_{j=n}^{\tilde{N}-n-1} \frac{1}{n} \log \|M_n(S^j x)\| \right. \\ \left. - \frac{2}{\tilde{N}} \sum_{j=0}^{\tilde{N}-n-1} \frac{1}{2n} \log \|M_{2n}(S^j x)\| \right| \leq \frac{11}{n} e^{-\frac{1}{2}nL_n}.$$

Integrating over $x \in X$, and using our definition of C_3 (8.1), we infer the following due to (8.23):

$$(8.28) \quad \left| L_N + \frac{N-2n}{N}L_n - 2\frac{N-n}{N}L_{2n} \right| \leq \frac{11}{n}e^{-\frac{1}{2}nL_n} + 4C_3\mu(\mathcal{B}^{(N)}) \\ < \frac{11}{n}e^{-\frac{1}{2}nL_n} + 8C_3N^{-\frac{7}{5}+\frac{4\delta}{5}},$$

and

$$(8.29) \quad \left| L_{2N} + \frac{N-n}{N}L_n - \frac{2N-n}{N}L_{2n} \right| \leq \frac{11}{n}e^{-\frac{1}{2}nL_n} + 4C_3\mu(\mathcal{B}^{(2N)}) \\ < \frac{11}{n}e^{-\frac{1}{2}nL_n} + 16C_3N^{-\frac{7}{5}+\frac{4\delta}{5}}.$$

From (8.28), we conclude that

$$L_N \geq L_n - \left(2 - \frac{2n}{N}\right)(L_n - L_{2n}) - \frac{11}{n}e^{-\frac{1}{2}nL_n} - 8C_3N^{-\frac{7}{5}+\frac{4\delta}{5}}.$$

This proves (8.2).

Taking the difference between (8.28) and (8.29), we obtain

$$L_N - L_{2N} \leq \frac{22}{n}e^{-\frac{1}{2}nL_n} + 24C_3N^{-\frac{7}{5}+\frac{4\delta}{5}} + \frac{n}{N}(L_n - L_{2n}).$$

This proves (8.3).

Proof of Proposition 8.4. This will be a continuation of the proof of Lemma 8.2. Note that all the constants C_3 's will be replaced by $U(\lambda, 1)$. Let \tilde{N} be either N or $2N$. Let us consider the first term in (8.27):

$$(8.30) \quad \left| \frac{1}{n} \sum_{k=0}^{n-1} \frac{1}{\tilde{N}} \log \|M_{\tilde{N}}(T_\omega^k(x, y))\| - \frac{1}{\tilde{N}} \log \|M_{\tilde{N}}(x, y)\| \right| \\ \leq \frac{1}{n\tilde{N}} \sum_{k=0}^{n-1} \left| \log \|M_{\tilde{N}}(T_\omega^k(x, y))\| - \log \|M_{\tilde{N}}(x, y)\| \right| \\ \leq \frac{1}{n\tilde{N}} \sum_{k=0}^{n-1} \left(\log \|M_k(x, y)\| + \log \|M_k(T_\omega^{\tilde{N}}(x, y))\| \right) \\ \leq \frac{1}{n\tilde{N}} \sum_{k=0}^{n-1} 2kU(\lambda, 1) \\ \leq \frac{U(\lambda, 1)n}{\tilde{N}}.$$

Hence (8.27) leads to

$$(8.31) \quad \left| \frac{1}{\tilde{N}} \log \|M_{\tilde{N}}(x, y)\| + \frac{1}{\tilde{N}} \sum_{j=n}^{\tilde{N}-n-1} \frac{1}{n} \log \|M_n(T_{\omega}^j(x, y))\| \right. \\ \left. - \frac{2}{\tilde{N}} \sum_{j=0}^{\tilde{N}-n-1} \frac{1}{2n} \log \|M_{2n}(T_{\omega}^j(x, y))\| \right| \leq \frac{11}{n} e^{-\frac{1}{2}nL_n} + \frac{U(\lambda, 1)n}{\tilde{N}},$$

holds for $(x, y) \notin \mathcal{B}^{(\tilde{N})}$. This implies

$$(8.32) \quad \left| \frac{1}{\tilde{N}} \log \|M_{\tilde{N}}(x, y)\| + \frac{1}{\tilde{N}} \sum_{j=0}^{\tilde{N}-1} \frac{1}{n} \log \|M_n(T_{\omega}^j(x, y))\| \right. \\ \left. - \frac{2}{\tilde{N}} \sum_{j=0}^{\tilde{N}-1} \frac{1}{2n} \log \|M_{2n}(T_{\omega}^j(x, y))\| \right| \leq \frac{11}{n} e^{-\frac{1}{2}nL_n} + \frac{5U(\lambda, 1)n}{\tilde{N}}.$$

Now we will apply Proposition 7.1 to v_n and v_{2n} with $K = \tilde{N}$. Note that conditions (I)-(IV) ensure the applicability of that proposition. Therefore following (7.1), we define

$$(8.33) \quad \varepsilon_4 = C_2 \tilde{N}^{-\frac{1}{10} + \frac{\delta}{5}} (\log \tilde{N})^{\frac{1}{10} + \frac{\delta_2}{5} + \delta_3}, \\ \varepsilon_5 = C_2 \left(472.5 + 3.2B_3(B_4 - m_4)\sqrt{C} \right)^{-1} (\log \tilde{N})^{\delta_3}.$$

For $\tilde{n} = n$ or $2n$, denote

$$\mathcal{C}_{\tilde{n}} := \left\{ (x, y) \in \mathbb{T}^2 : \left| \frac{1}{\tilde{N}} \sum_{j=0}^{\tilde{N}-1} v_{\tilde{n}} \circ T_{\omega}^j(x, y) - L_{\tilde{n}} \right| > \varepsilon_4 \right\}.$$

Then Proposition 7.1 implies that

$$(8.34) \quad \max(|\mathcal{C}_n|, |\mathcal{C}_{2n}|) \leq 2\sqrt{2} \exp\left(\frac{\pi}{4} \left[\frac{17}{36} + \frac{B_1}{4B_3^2} - \varepsilon_5 \right]\right) \\ + \sqrt{2} (C(B_4 - m_4))^{-1} \tilde{N}^{\frac{1}{5} - \frac{2\delta}{5}} (\log \tilde{N})^{-\frac{1}{5} - \frac{2\delta_2}{5}} \exp\left(-2(\log \tilde{N})^{\delta_2}\right).$$

Let

$$\mathcal{E} := \mathcal{C}_n \cup \mathcal{C}_{2n} \cup \mathcal{B}^{(\tilde{N})}.$$

For $(x, y) \notin \mathcal{E}$, by (8.32) we have that,

$$(8.35) \quad \left| \frac{1}{\tilde{N}} \log \|M_{\tilde{N}}(x, y)\| + L_n - 2L_{2n} \right| \leq \frac{11}{n} e^{-\frac{1}{2}nL_n} + \frac{5U(\lambda, 1)n}{\tilde{N}} + 2\varepsilon_4.$$

Together with (8.28), this implies that for any $(x, y) \notin \mathcal{E}$,

$$\begin{aligned}
& \left\| \mathbb{1}_{\mathcal{E}^c}(x, y) \left(\frac{1}{\tilde{N}} \log \|M_{\tilde{N}}(x, y)\| - L_{\tilde{N}} \right) \right\|_{L^\infty(\mathbb{T}^2)} \\
(8.36) \quad & \leq |L_{\tilde{N}} + L_n - 2L_{2n}| + \frac{11}{n} e^{-\frac{1}{2}nL_n} + \frac{5U(\lambda, 1)n}{\tilde{N}} + 2\varepsilon_4 \\
& \leq \frac{2n}{\tilde{N}}(L_n - L_{2n}) + \frac{22}{n} e^{-\frac{1}{2}nL_n} + 8U(\lambda, 1)\tilde{N}^{-\frac{7}{5} + \frac{4\delta}{5}} + \frac{5U(\lambda, 1)n}{\tilde{N}} + 2\varepsilon_4 \\
& =: \varepsilon_0.
\end{aligned}$$

Recall (8.23) states that

$$|\mathcal{B}^{(N)}| < 2N^{-\frac{7}{5} + \frac{4\delta}{5}}, \quad \text{and} \quad |\mathcal{B}^{(2N)}| \leq 4N^{-\frac{7}{5} + \frac{4\delta}{5}}.$$

Since $\delta > 0$, this clearly leads to

$$(8.37) \quad |\mathcal{B}^{(\tilde{N})}| \leq 2^{\frac{17}{5} - \frac{4\delta}{5}} \tilde{N}^{-\frac{7}{5} + \frac{4\delta}{5}} < 16\tilde{N}^{-\frac{7}{5} + \frac{4\delta}{5}}.$$

Combining (8.37) with (8.34), we obtain

$$\begin{aligned}
(8.38) \quad & \left\| \mathbb{1}_{\mathcal{E}}(x, y) \left(\frac{1}{\tilde{N}} \log \|M_{\tilde{N}}(x, y)\| - L_{\tilde{N}} \right) \right\|_{L^1(\mathbb{T}^2)} \\
& \leq U(\lambda, 1) |\mathcal{C}_n \cup \mathcal{C}_{2n} \cup \mathcal{B}^{(\tilde{N})}| \\
& \leq U(\lambda, 1) \left\{ 4\sqrt{2} \exp\left(\frac{\pi}{4} \left[\frac{17}{36} + \frac{B_1}{4B_3^2} - \varepsilon_5 \right]\right) + 16\tilde{N}^{-\frac{7}{5} + \frac{4\delta}{5}} \right. \\
& \quad \left. + 2\sqrt{2}(C(B_4 - m_4))^{-1} \tilde{N}^{\frac{1}{5} - \frac{2\delta}{5}} (\log \tilde{N})^{-\frac{1}{5} - \frac{2\delta_2}{5}} \exp\left(-2(\log \tilde{N})^{\delta_2}\right) \right\} =: \varepsilon_1.
\end{aligned}$$

By our conditions (V)-(VIII), we have

$$\begin{aligned}
(8.39) \quad & \varepsilon_0 \leq 4\varepsilon_4, \\
& \varepsilon_1 \leq 18U(\lambda, 1)\tilde{N}^{-\frac{7}{5} + \frac{4\delta}{5}} =: 18U(\lambda, 1)\tilde{N}^\eta,
\end{aligned}$$

Indeed, note that the right-hand-sides of (V) and (VI) are precisely ε_4 , which allow us to bound ε_0 by $4\varepsilon_4$. On the other hand, (VII) and (VIII) simply state that the sum of the first two terms in the braces defining ε_1 are bounded by the third term, viz. $2\tilde{N}^{-\frac{7}{5} + \frac{4\delta}{5}}$.

Let

$$\begin{aligned}
(8.40) \quad & \varepsilon_3 := C_4 \varepsilon_4 (\log \tilde{N})^{\delta_4}, \\
& \varepsilon_2 := C_5 \varepsilon_4 (\log \tilde{N})^{2\delta_4}, \\
& r = \frac{1 - 2\delta}{7 - 4\delta} \in (0, 1).
\end{aligned}$$

Our condition (X) and (XI) ensure that $\varepsilon_0 \leq 4\varepsilon_4 < \varepsilon_3 < \varepsilon_2$. Recall that $B_4 - m_4$ and $B_5^{(\tilde{N})} - m_5^{(\tilde{N})}$ are as in (8.8). Therefore, by our condition (IX) we have

$$(8.41) \quad \left(B_5^{(\tilde{N})} - m_5^{(\tilde{N})} \right) \tilde{N}^{-1} < 2 \log R.$$

Applying Lemma 4.2 to $v_{\tilde{N}}$, and taking (8.41) into account, we obtain

$$(8.42) \quad \begin{aligned} & \left| \left\{ (x, y) \in \mathbb{T}^2 : |v_{\tilde{N}}(x, y) - L_{\tilde{N}}| > \varepsilon_2 \right\} \right| \\ & \leq 2(2C_0)^{\frac{1}{2}} \exp \left(-\pi \left(36\varepsilon_0 + 16B_3 \sqrt{\varepsilon_1^r (B_4 - m_4)} \right)^{-1} \varepsilon_3 \right) \\ & \quad + C_0 \exp \left(-\pi \left(18\varepsilon_3 + 16B_3 \sqrt{U(\lambda, 1)} \sqrt{\varepsilon_1^{1-r} (B_5^{(\tilde{N})} - m_5^{(\tilde{N})})} \right)^{-1} \varepsilon_2 \right) \\ & \leq 2(2C_0)^{\frac{1}{2}} \exp \left(-\pi \left(36\varepsilon_0 + 16B_3 \sqrt{\varepsilon_1^r (B_4 - m_4)} \right)^{-1} \varepsilon_3 \right) \\ & \quad + C_0 \exp \left(-\pi \left(18\varepsilon_3 + 16B_3 \sqrt{U(\lambda, 1)} \sqrt{2\varepsilon_1^{1-r} \tilde{N} \log R} \right)^{-1} \varepsilon_2 \right). \end{aligned}$$

Note that we changed B_6 into $U(\lambda, 1)$ in this expression. Inserting our estimates of $\varepsilon_0, \varepsilon_1$, see (8.39), and choices of $\varepsilon_2, \varepsilon_3, r$, see (8.40), into (8.42), we arrive at

$$\begin{aligned} & \left| \left\{ (x, y) \in \mathbb{T}^2 : |v_{\tilde{N}}(x, y) - L_{\tilde{N}}| > C_5 \varepsilon_4 (\log \tilde{N})^{2\delta_4} \right\} \right| \\ & \leq 2(2C_0)^{\frac{1}{2}} \exp \left(\frac{-\pi C_4 \varepsilon_4 (\log \tilde{N})^{\delta_4}}{144\varepsilon_4 + 16B_3 \sqrt{(18U(\lambda, 1))^r \tilde{N}^{\eta r} (B_4 - m_4)}} \right) \\ & \quad + C_0 \exp \left(\frac{-\pi C_5 \varepsilon_4 (\log \tilde{N})^{2\delta_4}}{18C_4 \varepsilon_4 (\log \tilde{N})^{\delta_4} + 16B_3 \sqrt{U(\lambda, 1)} \sqrt{2(18U(\lambda, 1))^{1-r} \tilde{N}^{\eta(1-r)+1} \log R}} \right). \end{aligned}$$

Using (6.11), and $0 \leq r \leq 1$, we estimate

$$(18U(\lambda, 1))^r \leq 18U(\lambda, 1), \quad \text{and} \quad (18U(\lambda, 1))^{1-r} \leq 18U(\lambda, 1),$$

respectively. Hence we have

$$\begin{aligned} & \left| \left\{ (x, y) \in \mathbb{T}^2 : |v_{\tilde{N}}(x, y) - L_{\tilde{N}}| > C_5 \varepsilon_4 (\log \tilde{N})^{2\delta_4} \right\} \right| \\ & \leq 2(2C_0)^{\frac{1}{2}} \exp \left(\frac{-\pi C_4 \varepsilon_4 (\log \tilde{N})^{\delta_4}}{144\varepsilon_4 + 48B_3 \sqrt{2U(\lambda, 1)} \tilde{N}^{\eta r} (B_4 - m_4)} \right) \\ & \quad + C_0 \exp \left(\frac{-\pi C_5 \varepsilon_4 (\log \tilde{N})^{2\delta_4}}{18C_4 \varepsilon_4 (\log \tilde{N})^{\delta_4} + 96B_3 U(\lambda, 1) \sqrt{\tilde{N}^{\eta(1-r)+1} \log R}} \right). \end{aligned}$$

Plugging in our choice of ε_4 , see (8.33), and noting that the powers of \tilde{N} in numerators and denominators cancel out due to our choice of r , we infer that

$$\begin{aligned}
& \left| \left\{ (x, y) \in \mathbb{T}^2 : |v_{\tilde{N}}(x, y) - L_{\tilde{N}}| > C_5 \varepsilon_4 (\log \tilde{N})^{2\delta_4} \right\} \right| \\
& \leq 2(2C_0)^{\frac{1}{2}} \exp \left(\frac{-\pi C_2 C_4 (\log \tilde{N})^{\frac{1}{10} + \frac{\delta_2}{5} + \delta_3 + \delta_4}}{144 C_2 (\log \tilde{N})^{\frac{1}{10} + \frac{\delta_2}{5} + \delta_3} + 48 B_3 \sqrt{2U(\lambda, 1)} (B_4 - m_4)} \right) \\
& \quad + C_0 \exp \left(\frac{-\pi C_2 C_5 (\log \tilde{N})^{\frac{1}{10} + \frac{\delta_2}{5} + \delta_3 + 2\delta_4}}{18 C_2 C_4 (\log \tilde{N})^{\frac{1}{10} + \frac{\delta_2}{5} + \delta_3 + \delta_4} + 96 B_3 U(\lambda, 1) \sqrt{\log R}} \right) \\
& = 2(2C_0)^{\frac{1}{2}} \exp \left(\frac{-\pi C_2 C_4 (\log \tilde{N})^{\delta_4}}{144 C_2 + 48 B_3 \sqrt{2U(\lambda, 1)} (B_4 - m_4) (\log \tilde{N})^{-\frac{1}{10} - \frac{\delta_2}{5} - \delta_3}} \right) \\
& \quad + C_0 \exp \left(\frac{-\pi C_2 C_5 (\log \tilde{N})^{\delta_4}}{18 C_2 C_4 + 96 B_3 U(\lambda, 1) \sqrt{\log R} (\log \tilde{N})^{-\frac{1}{10} - \frac{\delta_2}{5} - \delta_3 - \delta_4}} \right),
\end{aligned}$$

as desired. \square

The readers will note that the constants were chosen in such a way that in the final steps of the proof only powers of $\log N$ remained inside of the exponential. We have found this to be more efficient over intermediate scales. The following, final, section of this paper will show how our work up to this point allows for such concrete estimates with specific numbers.

9. EXPLICIT NUMBERS AND PROOF OF THEOREM 1.1

Our goal here is to make concrete choices of our parameters so as to arrive at an actual multi-scale scheme for the skew-shift operator from Section 6. Let \mathcal{B}_n be as in Definition 8.1. The values below were found to be convenient ones, but clearly many other choices could have been made.

Definition 9.1. Set

$$R := 4, R_1 := 3, R_2 := 2,$$

in Definition 2.1. The coupling constant in (6.1) is required to obey $\lambda \in [\frac{1}{2}, 1]$. Further, in Proposition 8.4 set

$$\delta := \frac{1}{8}, \delta_2 := 1, \delta_3 := 2, \delta_4 := \frac{3}{2}.$$

as well as

$$C_2 := 203, C_4 := \frac{145}{\pi}, C_5 := \frac{850}{\pi}.$$

By an explicit computation, the condition in Definition 3.5 is satisfied. In fact, one has

$$B_3^2 - 289 \left(B_0 + \frac{13}{20 \log(R/R_1)} \right) > 61 > 0$$

Proposition 9.2. *Let $\omega = \frac{\sqrt{5}-1}{2}$ be the golden ratio, and consider the model (6.1) with $\lambda \in [\frac{1}{2}, 1]$ arbitrary but fixed. Let $a \geq 7$ and let n, N be positive integers such that $N \geq 10^{12}$, n divides N , and*

$$(9.1) \quad 10^{13}(n+1)^8 \leq N, \quad \frac{N}{(\log N)^{\frac{92}{3}}} < \frac{1}{2} \left(\frac{203}{22} e^{a/2} n \right)^{\frac{40}{3}}.$$

Impose the conditions

(a).

$$nL_n \geq a,$$

(b).

$$L_n - L_{2n} \leq \frac{1}{8}L_n,$$

(c).

$$\max(|\mathcal{B}_n|, |\mathcal{B}_{2n}|) \leq N^{-\frac{23}{10}}.$$

Then we have

$$(9.2) \quad \left| \left\{ (x, y) \in \mathbb{T}^2 : |v_{\tilde{N}}(x, y) - L_{\tilde{N}}| > 5.5 \times 10^4 \tilde{N}^{-\frac{3}{40}} (\log \tilde{N})^{\frac{53}{10}} \right\} \right| \leq 10 \exp \left(-(\log \tilde{N})^{\frac{3}{2}} \right),$$

holds for $\tilde{N} = N$ and $2N$.

Remark 9.3. We will choose the constant $a = 7$ along the inductive multiscale procedure. The only exception is the first step of the induction, which goes from the scale N_0 to N_1 , where for some of our main results we use a larger value of a . This is made possible by assumption (i) on the Lyapunov exponent at the initial scale and it is the reason behind the relatively small values of N_0 in Theorems 1.3 and 1.4.

Proof. We need to check the hypotheses of Proposition 8.4. We already verified (3.14), and the conditions of Lemma 8.2 hold by assumption. Let $\tilde{N} = N$ or $2N$. The function

$$[0.5, 1] \rightarrow \mathbb{R} : \lambda \mapsto U(\lambda, 4) - \log \lambda$$

is decreasing and positive. Hence

$$(9.3) \quad 0.5 < U(1, 4) \leq U(\lambda, 4) - \log \lambda \leq U\left(\frac{1}{2}, 4\right) - \log \frac{1}{2} < 1.$$

Further, the constant C in (I) satisfies $C < 11.97$. So that condition is implied by the stronger one

$$12^8(4 \log(2)n + 2 \log(2) + 1)^8 \leq \tilde{N}$$

which we may further strengthen to

$$36^8(n+1)^8 < 10^{13}(n+1)^8 \leq N,$$

which is the left-hand side of (9.1). Condition (II) holds, as does (III) since $\exp(4(\log \tilde{N})^{\delta_2}) = \tilde{N}^4 \geq \tilde{N} + 1$. Condition (IX) is implied by the stronger one

$$\tilde{N} > \frac{1 + \log R}{\log R} = \frac{2 \log(2) + 1}{2 \log(2)} \sim 1.721$$

which clearly holds. In view of (8.8) and (9.3), we have

$$(9.4) \quad 4 \log(2) + 0.5 \leq B_4 - m_4 \leq 4 \log(2) + 1.$$

Condition (IV) will therefore hold provided

$$\tilde{N}^{\frac{1}{8}} \log(\tilde{N}) - 21 \tilde{N}^{-\frac{27}{40}} (\log(\tilde{N}))^{\frac{27}{10}} - 181 > 0$$

The left-hand side is increasing in \tilde{N} , and one checks by explicit computation that it is positive if $N \geq 10^8$. So this condition holds as well. Condition (VIII) is implied by the following one

$$\tilde{N}^{-\frac{13}{10}} > \frac{2\sqrt{2}}{C(4 \log(2) + 0.5)} \tilde{N}^{-\frac{37}{20}} (\log(\tilde{N}))^{-\frac{3}{5}}$$

Simplifying this, one obtains the stronger condition

$$\tilde{N}^{\frac{1}{5}} (\log(\tilde{N}))^{\frac{3}{5}} > 0.08$$

which holds provided $N \geq 2$. So condition (VIII) holds.

Next, we look at condition (VI). Using the assumed lower bound $nL_n \geq a$ we find the condition

$$22e^{-a/2} < 203 n \tilde{N}^{-\frac{3}{40}} (\log(\tilde{N}))^{\frac{23}{10}}.$$

We recall that $\tilde{N} \in \{N, 2N\}$ and estimate $\log(\tilde{N}) \geq \log(N)$. This inequality follows from the upper bound in (9.1). For condition (VII), one checks that it follows from the slightly stronger

$$\tilde{N}^{-\frac{13}{10}} - 5.66 \exp(0.374 - 0.05(\log(\tilde{N}))^2) > 0$$

which holds for $\tilde{N} \geq 10^{12}$ (but it fails for 10^{11}). Hence we impose the second lower bound in (9.1). For condition (V), we use

$$L_n - L_{2n} \leq \frac{1}{8} L_n \leq \frac{U(\lambda, 1)}{8} \leq \frac{1}{4}$$

and so it suffices to check that

$$\frac{21n}{2\tilde{N}} + 16\tilde{N}^{-\frac{13}{10}} < 203\tilde{N}^{-\frac{3}{40}} (\log(\tilde{N}))^{\frac{23}{10}}$$

Bounding n in terms of N via (9.1) and discarding the $\log \tilde{N}$ on the right-hand side reduces us to

$$\frac{21}{2 \cdot 10^{\frac{13}{8}}} \cdot N^{-\frac{7}{8}} + 16\tilde{N}^{-\frac{13}{10}} < 203\tilde{N}^{-\frac{3}{40}}$$

This holds for all $N \geq 1$ so we are done with (V). Finally, we turn to (X) and (XI). Using $N \geq 10^{12}$ they hold provided

$$C_4 \geq 0.03, \quad C_5 \geq 0.007 \cdot C_4$$

Our actual values assigned to these constants satisfy

$$46 < C_4 < 47, \quad C_5 > 270$$

and so all conditions of Proposition 8.4 hold.

As for the conclusion of that proposition, we first compute $C_2C_5 < 5.5 \times 10^4$. Thus, the size of the deviations satisfy

$$C_2C_5\tilde{N}^{-\frac{1}{10}+\frac{\delta}{5}}(\log \tilde{N})^{\frac{1}{10}+\frac{\delta_2}{5}+\delta_3+2\delta_4} < 5.5 \times 10^4\tilde{N}^{-\frac{3}{40}}(\log \tilde{N})^{\frac{53}{10}}$$

as stated in (9.2). As for the measure bound, we calculate that

$$\begin{aligned} 2(2C_0)^{\frac{1}{2}} + C_0 &< 10, \\ U(\lambda, 1) &= \frac{1}{2} \log \left((4\lambda + 2)^2 + 2 \right) \leq \frac{1}{2} \log 38. \end{aligned}$$

Thus, in view of (9.4), one has, $48B_3\sqrt{2U(\lambda, 1)(B_4 - m_4)} < 11518$, and

$$144 + 11518C_2^{-1}(\log \tilde{N})^{-\frac{23}{10}} \leq 144 + \frac{11518}{203}(12 \log 10)^{-\frac{23}{10}} < 145.$$

Hence the first exponential in the measure bound of Proposition 8.4 contributes less than

$$\exp \left(-\pi C_4(\log \tilde{N})^{\frac{3}{2}}/145 \right) < \exp \left(-(\log \tilde{N})^{\frac{3}{2}} \right).$$

For the second exponential, we have $18C_4 < 831$, $96B_3U(\lambda, 1)\sqrt{2 \log 2} < 13317$, and

$$831 + 13317C_2^{-1}(\log \tilde{N})^{-\frac{38}{10}} \leq 831 + \frac{13317}{203}(12 \log 10)^{-\frac{38}{10}} < 850.$$

Hence, the second exponential contributes less than

$$\exp \left(-\pi C_5(\log \tilde{N})^{\frac{3}{2}}/850 \right) < \exp \left(-(\log \tilde{N})^{\frac{3}{2}} \right),$$

and we are done. \square

9.1. Proof of Theorem 1.1. Let $N_0 := 2 \times 10^{37}$. We define a sequence of scales $N_j := N_{j-1}^9$ for $j \geq 1$. In particular, $N_1 > 5 \times 10^{335}$. The proof is based on an induction on scales, where at every step we first apply Lemma 8.3 to control the Lyapunov exponent at the next scale. Afterwards, we apply Proposition 9.2 to obtain the large deviation estimate at the next scale and then we continue the induction.

For later purposes, we note some properties of this choice of scales. The last inequality is the main reason why we need to choose the scale so that N_1 is large.

Lemma 9.4. *Recall that we defined $N_{j+1} := N_j^9$ with $N_0 = 2 \times 10^{37}$. For all $j \geq 1$, we have the following bounds:*

$$(9.5) \quad 10^{13}(N_{j-1} + 1)^8 \leq N_j, \quad \frac{N_j}{(\log N_j)^{\frac{92}{3}}} < \frac{1}{2} \left(\frac{203}{22} e^{7/2} N_{j-1} \right)^{\frac{40}{3}}$$

as well as

$$(9.6) \quad 10 \exp \left(-(\log N_j)^{\frac{3}{2}} \right) \leq (N_j^9)^{-2.3} = N_{j+1}^{-2.3},$$

and

$$(9.7) \quad 5.5 \times 10^4 N_j^{-\frac{3}{40}} (\log N_j)^{\frac{53}{10}} \leq \frac{1}{20} L_{N_0}.$$

Proof. From the definition of the N_j , we have

$$(9.8) \quad 10^{13} (N_{j-1} + 1)^8 \leq N_j < N_{j-1}^{13},$$

and this implies (9.5). Notice that we have $N_j \geq N_1 > 5 \times 10^{335}$ for all $j \geq 1$. Then (9.6) follows from the inequality

$$(9.9) \quad 10 \exp\left(-(\log x)^{\frac{3}{2}}\right) \leq (x^9)^{-2.3},$$

which holds for all $x \geq 2.06 \times 10^{186}$. For (9.7), we note

$$5.5 \times 10^4 x^{-\frac{3}{40}} (\log x)^{\frac{53}{10}} \leq 10^{-5} = \frac{1}{20} \times 2 \times 10^{-4} \leq \frac{1}{20} L_{N_0},$$

where the first inequality holds for all $x \geq 10^{334}$. This proves the lemma. \square

We will inductively apply Lemma 8.3 to $j = 1, 2, 3, \dots$. We begin with $j = 1$. Condition (1) of Lemma 8.3 follows from our assumptions (i) and (ii) and the fact that $N_0 = 2 \times 10^{37}$. Condition (2) with $j = 1$ is fulfilled since

$$\frac{1}{N_0} \exp\left(-\frac{1}{2} N_0 L_{N_0}\right) \leq \frac{1}{2 \times 10^{37}} < \frac{2}{512} 10^{-4} \leq \frac{1}{512} L_{N_0}.$$

For condition (3), recall that $N_1 > 5 \times 10^{335}$, $\delta = \frac{1}{8}$ and $C_3 = U(\lambda, 1) \leq \frac{1}{2} \log 38$. Then we have

$$N_1^{-\frac{13}{10}} < (5 \times 10^{335})^{-1.3} < \frac{1}{320 \log 38} 10^{-4} \leq \frac{1}{1280 U(\lambda, 1)} L_{N_0}.$$

For condition (4),

$$(9.10) \quad \max(|\mathcal{B}_{N_0}|, |\mathcal{B}_{2N_0}|) \leq N_0^{-21} = (N_1)^{-7/3} \leq N_1^{-2.3}.$$

Hence Lemma 8.3 applies to $j = 1$, and yields

$$(9.11) \quad N_1 L_{N_1} \geq 7, \quad L_{N_1} - L_{2N_1} \leq \frac{1}{8} L_{N_1},$$

and

$$(9.12) \quad L_{2N_1} \geq \frac{1}{2} L_{N_0}.$$

We would like to apply Lemma 8.3 for $j = 2$. This requires measure estimates for \mathcal{B}_{N_1} and \mathcal{B}_{2N_1} . To this end, we invoke Proposition 9.2 with $n = N_0$, $N = N_1$ and $a = 7$. Let us check that its conditions are satisfied. First, we have (9.1) by applying (9.5) with $j = 0$. Moreover, conditions (a)-(c) hold by assumptions (i)-(iii) and (9.10). Hence, we can apply Proposition 9.2 and obtain that, for $\tilde{N} = N_1$ and $2N_1$,

$$\left| \left\{ (x, y) \in \mathbb{T}^2 : |v_{\tilde{N}}(x, y) - L_{\tilde{N}}| > 5.5 \times 10^4 \tilde{N}^{-\frac{3}{40}} (\log \tilde{N})^{\frac{53}{10}} \right\} \right| \leq 10 \exp\left(-(\log N_1)^{2.3}\right) \leq N_2^{-2.3}.$$

In the second step, we used (9.6) with $j = 1$. To turn this into measure estimates for $\mathcal{B}_{\tilde{N}}$, notice that (9.7) and (9.12) imply

$$5.5 \times 10^4 \tilde{N}^{-\frac{3}{40}} (\log \tilde{N})^{\frac{53}{10}} \leq \frac{1}{20} L_{N_0} \leq \frac{1}{10} L_{\tilde{N}}.$$

Therefore,

$$(9.13) \quad \max(|\mathcal{B}_{N_1}|, |\mathcal{B}_{2N_1}|) \leq N_2^{-2.3}.$$

We have shown how to pass from scale N_0 to N_1 via Lemma 8.3 and Proposition 9.2, by using the properties (9.5)-(9.7).

We can now iterate this procedure: We apply Lemma 8.3 with $j = 2$. The main input is the measure estimate (9.13), which verifies condition (4). The remaining conditions hold by our choice of scales, (9.11), (9.12) and assumptions (i) and (ii). (Notice that the sums in conditions (2) and (3) are rapidly convergent.) From Lemma 8.3, we obtain estimates of L_{N_2} and L_{2N_2} , in particular $L_{2N_2} \geq \frac{1}{2} L_{N_0}$. Then Proposition 9.2 yields the measure estimates for \mathcal{B}_{N_2} and \mathcal{B}_{2N_2} , which is the key input for Lemma 8.3 with $j = 3$, etc. We conclude that, after k steps of this procedure, we have

$$L_{2N_k} \geq \frac{1}{2} L_{N_0}.$$

This yields

$$L \geq \frac{1}{2} L_{N_0},$$

by taking $k \rightarrow \infty$, and we have proved Theorem 1.1. \square

9.2. Proof of Theorems 1.3. We follow the general line of argumentation of Theorem 1.1. The only difference is that in the sequence of scales N_j , we take the first step to be very large. Namely, while $N_0 = 3 \times 10^5$, we define

$$(9.14) \quad N_1 := 3 \times 10^{334}, \quad N_{j+1} := N_j^9, \quad \forall j \geq 1.$$

Notice that for $j \geq 1$, the scales N_j are essentially the ones used in the proof of Theorem 1.1 above. Therefore we have the following analog of Lemma 9.4, in which (9.6) for $j = 1$ is replaced

Lemma 9.5. *We have*

$$(9.15) \quad 10^{13}(N_0 + 1)^8 \leq N_1, \quad \frac{N_1}{(\log N_1)^{\frac{92}{3}}} < \frac{1}{2} \left(\frac{203}{22} e^{30} N_0 \right)^{\frac{40}{3}}.$$

Moreover, for all $j \geq 1$, we have the bounds

$$(9.16) \quad 10^{13}(N_j + 1)^8 \leq N_{j+1}, \quad \frac{N_{j+1}}{(\log N_{j+1})^{\frac{92}{3}}} < \frac{1}{2} \left(\frac{203}{22} e^{7/2} N_j \right)^{\frac{40}{3}},$$

as well as (9.6) and (9.7).

Except for (9.15), the bounds are only concerned with N_j , $j \geq 1$ and therefore follow in the same way as for Lemma 9.5. The new bound (9.15) follows from

$$\left(2 \frac{N_1}{(\log N_1)^{\frac{92}{3}}}\right)^{\frac{3}{40}} \frac{1}{\frac{203}{22} e^{30}} < 29974 < N_0.$$

This establishes Lemma 9.5. As before, we will successively apply Lemma 8.3 and Proposition 9.2 and iterate. We begin by applying Lemma 8.3 with $j = 1$. Condition (1) is immediate from assumption (i) and $N_0 = 3 \times 10^5$; indeed:

$$(9.17) \quad N_0 L_{N_0} \geq 2N_0 10^{-4} = 60.$$

We can use this inequality to verify condition (2) as well:

$$(9.18) \quad \frac{1}{N_0} e^{-\frac{N_0 L_{N_0}}{2}} \leq \frac{1}{3} 10^{-5} e^{-30} < 10^{-18} < \frac{2}{512} 10^{-4} < \frac{L_{N_0}}{512}.$$

Condition (3) holds by our choice of N_1 . Finally, condition (4) holds by assumption (iii):

$$(9.19) \quad \max(|\mathcal{B}_{N_0}|, |\mathcal{B}_{2N_0}|) \leq N_0^{-141} < (3 \times 10^{334})^{-2.3} = N_1^{-2.3}.$$

Hence, Lemma 8.3 applies and yields (9.11) and (9.12) as before. Next, we verify the assumption of Proposition 9.2 with $n = N_0$ and $N = N_1$. The key difference is that we now take $a = 60$. This is made possible by (9.17), since it verifies condition (a) of Proposition 9.2. Condition (b) is immediate from assumption (ii) and condition (c) was checked in (9.19). The bounds (9.1) hold by (9.15). Therefore, we can apply Proposition 9.2. Combining the resulting estimate with (9.7) for $j = 1$, (9.12) and (9.6) for $j = 1$, we obtain the measure estimate

$$\max(|\mathcal{B}_{N_1}|, |\mathcal{B}_{2N_1}|) \leq N_2^{-2.3}.$$

At this point, we have moved completely from scale N_0 to scale N_1 and can follow the argument from Theorem 1.1 verbatim. In particular, we take $a = 7$ in every subsequent application of Proposition 9.2. The only difference is the $m = 0$ term in condition (2) of Lemma 8.3, which now involves $N_0 = 3 \times 10^{-5}$. By (9.18), we can replace condition (2) by the stronger bound

$$\sum_{m=1}^{j-1} N_m^{-1} e^{-\frac{1}{2} N_m L_{N_m}} < \frac{10^{-4}}{256} - 10^{-18}$$

and this holds by our choice of scales and the estimates (9.12) along the induction (notice again the rapid convergence of the series). We conclude that

$$L \geq \frac{1}{2} L_{N_0}$$

and this proves Theorem 1.3. □

9.3. Proof of Theorem 1.4. Again, we follow the same steps for a different sequence of scales. We have $N_0 = 3 \times 10^4$. We define the sequence of scales N_j , $j \geq 1$ by

$$N_1 := 3 \times 10^{320}, \quad N_{j+1} := N_j^9, \quad \forall j \geq 1.$$

We still have Lemma 9.5 for this choice of scales. Indeed, (9.16) and (9.6) still follow from the inequalities (9.8) and (9.9) given in the proof of Lemma 9.4. For (9.7), we now use assumption (i) to find

$$5.5 \times 10^4 x^{-\frac{3}{40}} (\log x)^{\frac{53}{10}} \leq 10^{-4} = \frac{1}{20} \times 2 \times 10^{-3} \leq \frac{1}{20} L_{N_0},$$

where the first inequality holds for all $x \geq 10^{320}$, so in particular for all N_j with $j \geq 1$. Finally, (9.15) follows from

$$\left(2 \frac{N_1}{(\log N_1)^{\frac{92}{3}}} \right)^{\frac{3}{40}} \frac{1}{\frac{203}{22} e^{30}} < 2938 < N_0.$$

This establishes Lemma 9.5 for the new choice of scales.

Next we check the hypotheses for Lemma 8.3 with $j = 1$. Condition (1) is immediate from assumption (i) and $N_0 = 3 \times 10^4$:

$$(9.20) \quad N_0 L_{N_0} \geq 2N_0 10^{-3} = 60$$

(Compare this to (9.17).) Condition (2) holds by

$$(9.21) \quad \frac{1}{N_0} e^{-\frac{N_0 L_{N_0}}{2}} \leq \frac{1}{3} 10^{-4} e^{-30} < 10^{-17} < \frac{2}{512} 10^{-3} < \frac{L_{N_0}}{512},$$

where we used (9.20). Condition (3) holds by our choice of scales N_j , $j \geq 1$, and condition (4) holds by assumption (iii):

$$(9.22) \quad \max(|\mathcal{B}_{N_0}|, |\mathcal{B}_{2N_0}|) \leq N_0^{-165} < (3 \times 10^{320})^{-2.3} = N_1^{-2.3}.$$

Therefore we can apply Lemma 8.3 and obtain (9.11) and (9.12). As in the proof of Theorem 1.3, the first application of Proposition 9.2 utilizes $a = 60$. This is made possible by (9.17), since it verifies condition (a) of Proposition 9.2. Now we iterate the argument in the same way as was done for Theorem 1.1 and 1.3. (Notice that the series in conditions (2) and (3) of Lemma 8.3 are still rapidly convergent.) The end result is the lower bound

$$L \geq \frac{1}{2} L_{N_0}$$

and Theorem 1.4 is proved. \square

REFERENCES

- [BelSim] Béllissard, J., Simon, B. *Cantor spectrum for the almost Mathieu equation.* J. Funct. Anal. 48 (1982), no. 3, 408–419.
- [Bou1] Bourgain, J. *Green's function estimates for lattice Schrödinger operators and applications.* Annals of Mathematics Studies, 158. Princeton University Press, Princeton, NJ, 2005.
- [Bou2] Bourgain, J. *On the spectrum of lattice Schrödinger operators with deterministic potential.* Dedicated to the memory of Thomas H. Wolff. J. Anal. Math. 87 (2002), 37–75.

- [BouGol] Bourgain, J., Goldstein, M. *On nonperturbative localization with quasi-periodic potential*. Ann. of Math. (2) 152 (2000), no. 3, 835–879.
- [BouGolSch] Bourgain, J., Goldstein, M., Schlag, W. *Anderson localization for Schrödinger operators on \mathbb{Z} with potentials given by the skew-shift*. Comm. Math. Phys. 220 (2001), no. 3, 583–621.
- [Dam] Damanik, D. *Lyapunov exponents and spectral analysis of ergodic Schrödinger operators: a survey of Kotani theory and its applications*. Spectral theory and mathematical physics: a Festschrift in honor of Barry Simon’s 60th birthday, 539–563, Proc. Sympos. Pure Math., 76, Part 2, Amer. Math. Soc., Providence, RI, 2007.
- [DuaKle] Duarte, P., Klein, S. *Lyapunov exponents of linear cocycles. Continuity via large deviations*. Atlantis Studies in Dynamical Systems, 3. Atlantis Press, Paris, 2016.
- [DuaKle2] Duarte, P., Klein, S. *Continuity of the Lyapunov Exponents of Linear Cocycles* Publicações Matemáticas do IMPA, 2017.
https://impa.br/wp-content/uploads/2017/08/31CBM_02.pdf
- [Fur] Fürstenberg, H. *Noncommuting random products*. Trans. Amer. Math. Soc. 108 (1963), 377–428.
- [GolSch] Goldstein, M., Schlag, W. *Hölder continuity of the integrated density of states for quasi-periodic Schrödinger equations and averages of shifts of subharmonic functions*. Ann. of Math. (2) 154 (2001), no. 1, 155–203.
- [D.R.H-B] Heath-Brown, D. R. *Pair correlation for fractional parts of cn^2* . Math. Proc. Cambridge Philos. Soc. 148 (2010), no. 3, 385–407.
- [Her] Herman, Michael-R. *Une méthode pour minorer les exposants de Lyapounov et quelques exemples montrant le caractère local d’un théorème d’Arnol’d et de Moser sur le tore de dimension 2*. Comment. Math. Helv. 58 (1983), no. 3, 453–502.
- [Kat] Katznelson, Y. *An introduction to harmonic analysis*. Third edition. Cambridge Mathematical Library. Cambridge University Press, Cambridge, 2004.
- [MarStr] Marklof, J., Strömbergsson, A. *Equidistribution of Kronecker sequences along closed horocycles*. Geom. Funct. Anal. 13 (2003), no. 6, 1239–1280.
- [Mon] Montgomery, Hugh L. *Ten lectures on the interface between analytic number theory and harmonic analysis*. CBMS Regional Conference Series in Mathematics, 84. American Mathematical Society, Providence, RI, 1994.
- [RudSarZah] Rudnick, Z., Sarnak, P., Zaharescu, A. *The distribution of spacings between the fractional parts of $n^2\alpha$* . Invent. Math. 145 (2001), no. 1, 37–57.
- [Sch] Schlag, W. *Regularity and convergence rates for the Lyapunov exponents of linear cocycles*. J. Mod. Dyn. 7 (2013), no. 4, 619–637.
- [SorSpe] Sorets, E., Spencer, T. *Positive Lyapunov exponents for Schrödinger operators with quasi-periodic potentials*. Comm. Math. Phys. 142 (1991), no. 3, 543–566.
- [Via] Viana, M. *Lectures on Lyapunov exponents*. Cambridge Studies in Advanced Mathematics, 145. Cambridge University Press, Cambridge, 2014.

INSTITUTE FOR ADVANCED STUDY, SCHOOL OF MATH, 1 EINSTEIN DRIVE, PRINCETON, NJ 08540, U.S.A.