

IMAGINARY QUADRATIC NUMBER FIELDS WITH CLASS GROUPS OF SMALL EXPONENT

ANDREAS-STEPHAN ELSENHANS, JÜRGEN KLÜNERS, AND FLORIN NICOLAE

ABSTRACT. Let $D < 0$ be a fundamental discriminant and denote by $E(D)$ the exponent of the ideal class group $\text{Cl}(D)$ of $K = \mathbb{Q}(\sqrt{D})$. Under the assumption that no Siegel zeros exist we compute all such D with $E(D)$ is a divisor of 8. We compute all D with $|D| \leq 3.1 \cdot 10^{20}$ such that $E(D) \leq 8$.

1. INTRODUCTION

Let D be a fundamental discriminant, i.e. the discriminant of a quadratic number field. For $D < 0$ let $E(D)$ be the exponent of the ideal class group $\text{Cl}(D)$ of the imaginary quadratic field $K = \mathbb{Q}(\sqrt{D})$. Under the Extended Riemann Hypothesis it is known (see [3], [13]) that $E(D) \gg \frac{\log |D|}{\log \log |D|}$. Without any unproved hypothesis it is not even known that $E(D) \rightarrow \infty$. In [13], Theorem 1, it is shown that there is at most one imaginary quadratic field with $|D| > 5460$ and $E(D) = 2$. In [3], [13] it is shown (ineffectively) that there are finitely many imaginary quadratic fields with $E(D) = 3$. In [9], Theorem 2, it is observed that for given $r \geq 0$ there are finitely many imaginary quadratic fields with $E(D) = 2^r$ or $E(D) = 3 \cdot 2^r$. (See also [6].) Finally, there are finitely many imaginary quadratic fields with $E(D) = 5$ ([9], Theorem 1).

In this note we are interested to determine all D such that the class group has exponent at most 8, i.e. $\text{Cl}(D)^c$ is the trivial group for some $c \leq 8$. In other words we want that the class group is of type $C_2^r \times C_3^s, C_5^r, C_7^r$, or $C_2^r \times C_4^s \times C_8^t$, where C_i denotes the cyclic group of order i . For any given r, s it is known that there are infinitely many $D < 0$ such that $C_2^r \times C_4^s$ is a subgroup of $\text{Cl}(D)$.

Our computations show the following:

Theorem 1. *There are exactly 1555 imaginary quadratic fields with discriminant $|D| \leq 3.1 \cdot 10^{20}$ and class group of exponent ≤ 8 .*

Exponent	Number of fields found	Field with largest discriminant
1	9	$\mathbb{Q}(\sqrt{-163})$
2	56	$\mathbb{Q}(\sqrt{-5460})$
3	17	$\mathbb{Q}(\sqrt{-4027})$
4	203	$\mathbb{Q}(\sqrt{-435435})$
5	27	$\mathbb{Q}(\sqrt{-37363})$
6	432	$\mathbb{Q}(\sqrt{-5761140})$
7	33	$\mathbb{Q}(\sqrt{-118843})$
8	778	$\mathbb{Q}(\sqrt{-430950520})$

TABLE 1. Number of imaginary quadratic fields with small exponent

The discriminants with more than 7 decimal places are -11148180, -12517428, -15337315, -15898740, -17168515, -28663635, -29493555, -31078723, -430950520.

Theorem 2. *Assuming ERH, our computations found all fields for exponent up to 5 and 8. Assuming the non-existence of Siegel-zeros, our computation found all fields with exponents 2,4,8.*

Without assuming any unproven statement we can conclude that there is at most one missing field with exponent 2,4 or 8.

We remark that for exponent $2 \leq c \leq 8, c \neq 7$ it is (unconditionally) known that there are only finitely many imaginary fields with exponent c . The explicit lists are only known using suitable conjectures.

Let

$$2 = p_1 < p_2 < p_3 < \dots$$

be the sequence of prime numbers. For $n \geq 1$ let

$$d_n := p_1 \cdot \dots \cdot p_n$$

be the product of the first n prime numbers. In Section 4 we determine for given $r \geq 1$ a number N_{2^r} such that there is at most one imaginary quadratic field with $|D| \geq d_{N_{2^r}}$ and $E(D) = 2^r$. For $r = 1$ we have $N_2 = 11$ and $d_{N_2} \leq 2.01 \cdot 10^{11}$. For $r = 2$ we have $N_4 = 24$ and $d_{N_4} \leq 2.38 \cdot 10^{34}$. For $r = 3$ we have $N_8 = 58$ and $d_{N_8} \leq 3.17 \cdot 10^{110}$.

2. SOME THEORETICAL ESTIMATES

As before let $D < 0$ be a fundamental discriminant. In the following we want to use the knowledge of a small splitting prime p , i.e. a prime $p \nmid D$ such that $p\mathcal{O}_K = \mathfrak{p} \cdot \bar{\mathfrak{p}}$ splits into two different prime ideals in K . We are interested to give some lower bound of the order of the ideal \mathfrak{p} in the class group $\text{Cl}(D)$. The following lemma appears in different forms at least in [5] p. 174-175, [3] Lemma 2, [13] Lemma 5.

Lemma 1. *Let $c > 0$ be an integer, let p be a split prime in K , and let \mathfrak{p} be a prime divisor of p in K . If $p^c < |D|/4$, then the order of \mathfrak{p} in the class group $\text{Cl}(D)$ is strictly larger than c .*

As shown in Lemma 1 a small split prime in an imaginary quadratic number field already gives a good lower bound for the exponent of the class group. Here we want to use the extended Riemann hypothesis (ERH). Using this, we can prove:

Theorem 3. *Let K be a quadratic number field such that the absolute value of the discriminant D is larger than $e^{25} \approx 7.2 \cdot 10^{10}$. Assume the extended Riemann hypothesis. Then there exists a split prime p such that*

$$p \leq (1.881 \log(|D|) + 2 \cdot 0.34 + 5.5)^2.$$

Proof. This is the result of Table 3 on page 1731 in [1]. □

We remark that in this paper there are similar statements with weaker constants for small discriminants. We used those in the following table for exponent smaller or equal to 3.

Exponent	Bound for $ D $
1	$1.7 \cdot 10^3$
2	$6 \cdot 10^6$
3	$9.7 \cdot 10^{10}$
4	$3.4 \cdot 10^{15}$
5	$2.3 \cdot 10^{20}$
6	$2.5 \cdot 10^{25}$
7	$3.9 \cdot 10^{30}$
8	$8.9 \cdot 10^{35}$

TABLE 2. ERH based bound (Theorem 3) for various exponents

3. USING SIEGEL-TATUZAWA BOUNDS

In the already cited paper by Weinberger [13], it is suggested to use efficient bounds based on Siegel zeros. Compared to the original Siegel bounds they have the advantage that the constants can be explicitly computed. Weinberger used this approach to determine all (assuming that there are no Siegel zeros) imaginary quadratic number fields of exponent 2. If we do not assume any unproven conjecture it is shown that at most one field is missing.

For a fundamental discriminant D we define the character

$$\chi(n) := \chi_D(n) := \left(\frac{D}{n} \right),$$

where the symbol denotes the Kronecker symbol. We associate to it the following L -series:

$$L(s, \chi) = \sum_{n=1}^{\infty} \frac{\chi(n)}{n^s}, \Re(s) > 0.$$

Let $h(D) := |\text{Cl}(D)|$ be the class number of the field $K = \mathbb{Q}(\sqrt{D})$. For $D < -4$ it is well known that

$$h(D) = \frac{\sqrt{|D|} L(1, \chi)}{\pi}.$$

We are interested in good lower bounds for the value $L(1, \chi)$. It is known that there are no zeros of $L(s, \chi)$ for $\Re(s) \geq 1$. We get efficient lower bounds, if we assume that there are no real zeros of $L(s, \chi)$ which are close to 1. Tatuzawa [12, Lemma 9 and Theorem 2] proved:

Lemma 2. *Let $0 < \varepsilon < 1/2$. There is at most one $|D| \geq \max(e^{1/\varepsilon}, e^{11.2})$ such that*

$$L(1, \chi) \leq \frac{0.655\varepsilon}{|D|^\varepsilon}.$$

In this case $L(s, \chi)$ has a real zero s with $1 - \varepsilon/4 < s < 1$.

As consequences we obtain

Lemma 3. *Let $|D| > e^{11.2}$. If $L(s, \chi) \neq 0$ for $1 - \frac{1}{4 \log |D|} \leq s < 1$ then*

$$h(D) > \frac{0.655}{\pi e} \cdot \frac{\sqrt{|D|}}{\log |D|}.$$

Proof. We apply Lemma 2 with $\varepsilon = 1/\log(|D|)$ and obtain

$$L(1, \chi) > \frac{0.655\varepsilon}{|D|^\varepsilon} = \frac{0.655}{e \log |D|}.$$

Using the class number formula $L(1, \chi) = \frac{\pi h(D)}{\sqrt{|D|}}$, we get the assertion. \square

Lemma 4. *Let $A \geq e^{11.2}$. For all $|D| \geq A$ with at most one exception it holds that*

$$h(D) > \frac{0.655}{\pi \cdot \log A} \cdot |D|^{\frac{1}{2} - \frac{1}{\log A}}.$$

For $|D| = A^m$ with $m \geq 1$ we get:

$$h(D) > \frac{m \cdot 0.655 \cdot \sqrt{|D|}}{\pi \cdot e^m \cdot \log |D|}$$

with at most one exception.

Proof. We apply Lemma 2 with $\varepsilon = 1/\log A$ and obtain

$$L(1, \chi) > \frac{0.655\varepsilon}{|D|^\varepsilon},$$

$$h(D) = \frac{\sqrt{|D|}L(1, \chi)}{\pi} > \frac{0.655}{\pi \cdot \log A} \cdot |D|^{\frac{1}{2} - \frac{1}{\log A}}$$

for all $|D| \geq A$ with at most one exception. The second statement is a straightforward computation. \square

The following lemma gives some improvement for the case with one exception.

Lemma 5. *Let $A \geq 10^6$, $\varepsilon := 1/\log A$ and $m := \frac{\log |D|}{\log A}$. Then for all D with $|D| \geq A$ we have with at most one exception:*

$$L(1, \chi) \geq \min \left(\frac{1}{7.732 \log |D|}, 1.5 \cdot 10^6 \frac{\varepsilon}{|D|^\varepsilon} \right)$$

$$(1) \quad h(D) \geq \min \left(\frac{\sqrt{|D|}}{\pi \cdot 7.732 \cdot \log |D|}, \frac{m \cdot 1.5 \cdot 10^6 \cdot \sqrt{|D|}}{\pi \cdot e^m \cdot \log |D|} \right).$$

Note that for $m \leq 19.2$ the first number is the minimum.

Proof. This is the main result of [4]. \square

Let us compare the result of Lemma 5 with Lemmata 3 and 4. When comparing Lemma 3 with Lemma 5, then we see that Lemma 3 gives a lower class group bound which is about a factor 2 better. But here we have to assume that there are no Siegel zeros. When we compare Lemma 4 with Lemma 5, then we see that for $m \geq 2.6$ Lemma 5 is better. Note that the second number in the minimum is always better than the bound in Lemma 4. Furthermore it is important to note that $(10^6)^{19.2} > 10^{115}$ which is sufficient for all our computations for exponent 8.

4. FIELDS WITH EXPONENT A POWER OF TWO

Weinberger proved in [13], Theorem 1, that there is at most one imaginary quadratic field with $|D| > 5460$ and $E(D) = 2$. We want to generalize his result to exponent 2^r . Let

$$2 = p_1 < p_2 < p_3 < \dots$$

be the sequence of prime numbers. For $n \geq 1$ let

$$d_n := p_1 \cdot \dots \cdot p_n$$

be the product of the first n prime numbers. In this section we determine for given $r \geq 1$ a number N_{2^r} such that there is at most one imaginary quadratic field with $|D| \geq d_{N_{2^r}}$ and $E(D) = 2^r$.

Lemma 6. *Let $D < 0$ be a fundamental discriminant. If $E(D) = 2^r$ with $r \geq 1$ then*

$$h(D) \leq 2^{r(\omega(D)-1)}.$$

Proof. If $E(D) = 2^r$ then the class group $\text{Cl}(D)$ is isomorphic to $C_2^{a_1} \times C_{2^2}^{a_2} \times \dots \times C_{2^r}^{a_r}$, $a_1 \geq 0, \dots, a_{r-1} \geq 0, a_r > 0$, and we have

$$h(D) = 2^{a_1+2a_2+\dots+ra_r}.$$

By genus theory we have that

$$a_1 + \dots + a_r = \omega(D) - 1,$$

hence

$$h(D) = 2^{a_1+2a_2+\dots+ra_r} \leq 2^{r(a_1+\dots+a_r)} = 2^{r(\omega(D)-1)}.$$

□

Lemma 7. *If D is a fundamental discriminant then*

$$|D| \geq d_{\omega(D)}.$$

Proof. We have that

$$D = \prod_{p|D} p^*$$

with $2^* \in \{-4, -8, 8\}$ and $p^* = (-1)^{\frac{p-1}{2}} p$ for $p \neq 2$, hence

$$|D| \geq p_1 \cdot \dots \cdot p_{\omega(D)} = d_{\omega(D)}.$$

□

Theorem 4. *Let $r \geq 1$ be an integer. Let N_{2^r} be the smallest of the integers N such that*

$$\begin{aligned} d_N &\geq e^{11.2}, \\ p_N^{\frac{1}{2} - \frac{1}{\log d_N}} &\geq 2^r, \end{aligned}$$

and

$$\frac{0.655}{\pi \cdot e} \cdot \frac{\sqrt{d_N}}{\log d_N} \geq 2^{r(N-1)}.$$

Let $D < 0$ be a fundamental discriminant with $|D| \geq d_{N_{2^r}}$. If $L(s, \chi) \neq 0$ for $1 - \frac{1}{4 \log |D|} \leq s < 1$ then $E(D) \neq 2^r$.

Without any assumption on zeros of L -functions, there is at most one D with $|D| \geq d_{N_{2^r}}$ and $E(D) = 2^r$.

Proof. Let $N \geq 1$ be an integer with the three properties from the hypothesis. Assume that $L(s, \chi) \neq 0$ for $1 - \frac{1}{4 \log |D|} \leq s < 1$. We apply Lemma 3 and obtain

$$h(D) > \frac{0.655}{\pi e} \cdot \frac{\sqrt{|D|}}{\log |D|}.$$

Suppose that $E(D) = 2^r$. Lemma 6 implies

$$2^{r(\omega(D)-1)} > \frac{0.655}{\pi e} \cdot \frac{\sqrt{|D|}}{\log |D|} \geq \frac{0.655}{\pi e} \cdot \frac{\sqrt{d_N}}{\log d_N} \geq 2^{r(N-1)},$$

hence

$$\omega(D) > N.$$

By Lemma 7 we have that

$$|D| \geq d_{\omega(D)} \geq d_N \cdot p_N^{\omega(D)-N},$$

so

$$\begin{aligned} 2^{r(\omega(D)-1)} &> \frac{0.655}{\pi e} \cdot \frac{\sqrt{|D|}}{\log |D|} \geq \frac{0.655}{\pi e} \cdot \frac{\sqrt{d_N} \cdot \sqrt{p_N}^{\omega(D)-N}}{\log d_N + (\omega(D) - N) \log p_N} = \\ &= \frac{0.655}{\pi e} \cdot \frac{\sqrt{d_N} \cdot 2^{r(\omega(D)-N)} \cdot \left(\frac{\sqrt{p_N}}{2^r}\right)^{\omega(D)-N}}{\log d_N + (\omega(D) - N) \log p_N}, \\ 2^{r(N-1)} &> \frac{0.655}{\pi e} \cdot \frac{\sqrt{d_N} \cdot \left(\frac{\sqrt{p_N}}{2^r}\right)^{\omega(D)-N}}{\log d_N + (\omega(D) - N) \log p_N} \geq \frac{0.655}{\pi e} \cdot \frac{\sqrt{d_N}}{\log d_N}, \end{aligned}$$

since the hypothesis $p_N^{\frac{1}{2} - \frac{1}{\log d_N}} \geq 2^r$ implies $\sqrt{p_N} > 2^r$ so the function

$$x \mapsto \frac{\left(\frac{\sqrt{p_N}}{2^r}\right)^{x-N}}{\log d_N + (x - N) \log p_N}, x \geq N$$

is increasing. This contradicts the hypothesis

$$\frac{0.655}{\pi \cdot e} \cdot \frac{\sqrt{d_N}}{\log d_N} \geq 2^{r(N-1)}.$$

So $E(D) \neq 2^r$.

We make now no assumption on zeros of L -functions. We apply Lemma 4 with $A = d_N$ and obtain

$$h(D) > \frac{0.655}{\pi \cdot \log d_N} \cdot |D|^{\frac{1}{2} - \frac{1}{\log d_N}}$$

for all discriminants $D < 0$ such that $|D| \geq d_N$ with at most one exception. Let $D < 0$ be a discriminant such that $|D| \geq d_N$ and

$$h(D) > \frac{0.655}{\pi \cdot \log d_N} \cdot |D|^{\frac{1}{2} - \frac{1}{\log d_N}}.$$

Suppose that $E(D) = 2^r$. By Lemma 6 and the choice of N we have that

$$2^{r(\omega(D)-1)} \geq h(D) > \frac{0.655}{\pi \cdot \log d_N} \cdot |D|^{\frac{1}{2} - \frac{1}{\log d_N}} \geq 2^{r(N-1)}$$

hence

$$\omega(D) > N.$$

By Lemma 7 we have that

$$|D| \geq d_{\omega(D)} > d_N \cdot p_N^{\omega(D)-N},$$

so

$$\begin{aligned} 2^{r(\omega(D)-1)} &\geq h(D) > \frac{0.655}{\pi \cdot \log d_N} \cdot |D|^{\frac{1}{2} - \frac{1}{\log d_N}} > \\ &> \frac{0.655}{\pi \cdot \log d_N} \cdot d_N^{\frac{1}{2} - \frac{1}{\log d_N}} \cdot p_N^{(\omega(D)-N) \cdot (\frac{1}{2} - \frac{1}{\log d_N})} = \\ &= \frac{0.655}{\pi} \cdot \frac{d_N^{\frac{1}{2} - \frac{1}{\log d_N}}}{\log d_N} \cdot 2^{r(\omega(D)-N)} \cdot \left(\frac{p_N^{\frac{1}{2} - \frac{1}{\log d_N}}}{2^r} \right)^{\omega(D)-N} \\ &\geq \frac{0.655}{\pi} \cdot \frac{d_N^{\frac{1}{2} - \frac{1}{\log d_N}}}{\log d_N} \cdot 2^{r(\omega(D)-N)}, \end{aligned}$$

since

$$\left(\frac{p_N^{\frac{1}{2} - \frac{1}{\log d_N}}}{2^r} \right)^{\omega(D)-N} \geq 1$$

by the hypothesis $p_N^{\frac{1}{2} - \frac{1}{\log d_N}} \geq 2^r$. It follows that

$$2^{r(N-1)} > \frac{0.655}{\pi} \cdot \frac{d_N^{\frac{1}{2} - \frac{1}{\log d_N}}}{\log d_N} = \frac{0.655}{\pi \cdot e} \cdot \frac{\sqrt{d_N}}{\log d_N},$$

in contradiction with the hypothesis

$$\frac{0.655}{\pi \cdot e} \cdot \frac{\sqrt{d_N}}{\log d_N} \geq 2^{r(N-1)}.$$

So $E(D) \neq 2^r$. □

Example. For $r = 1$ we have $N_2 = 11$ and $d_{N_2} \leq 2.01 \cdot 10^{11}$. For $r = 2$ we have $N_4 = 24$ and $d_{N_4} \leq 2.38 \cdot 10^{34}$. For $r = 3$ we have $N_8 = 58$ and $d_{N_8} \leq 3.17 \cdot 10^{110}$.

5. ALGORITHM USING SIEGEL BOUNDS

In this section we want to use the estimates from Lemmata 3 and 5 in order to compute all imaginary quadratic number fields with exponent 4 and 8. When we use the bounds from Lemma 3, we potentially miss fields such that the corresponding L -series has a Siegel zero close to 1. The estimate from Lemma 5 is a little bit weaker. This leads to more expensive computations, but it has the advantage that we can prove that we miss at most one field. These estimates are not valid for fields with small discriminant. This is not a big problem, since the class groups of fields with small discriminants are known. E.g. the web-page [2] provides a table of all quadratic fields up to absolute discriminant 10^7 . We checked the small fields independently by computations in Magma we do not describe here.

Let $c \in \{4, 8\}$ be the exponent we are looking for. Then we split our problem by looking at discriminants with k different prime factors. The estimates in Section 4 show that we can bound the maximal number of prime factors. In the following we write our discriminants as a product of k fundamental discriminants p^* . For every odd prime p it holds that $p^* = (-1)^{\frac{p-1}{2}} p$, and $2^* \in \{-4, -8, 8\}$.

In the following let

$$D = p_1^* \cdots p_k^*, \text{ where we assume that } p_1 < \cdots < p_k.$$

From genus theory it is well known that the 2-rank of the class group $\text{Cl}(D)$ is exactly $k - 1$. When we assume that the class group is of exponent c , then the maximal possible class group is $(\mathbb{Z}/c\mathbb{Z})^{k-1}$ and therefore of order c^{k-1} . Using the estimates of Lemma 3 or 5 we can compute an upper bound for the absolute value of D . The basic idea of the algorithm is to test all D smaller than this bound consisting of exactly k prime factors. Especially, when k is large it is not practical to list and test all those D . Therefore we would like to reduce this list further. The 4-rank of class groups of quadratic number fields is well studied and there are nice formulas to compute it. Furthermore, it is known that the average 4-rank is small [7, 8]. Assume that we know the 4-rank r_4 of $\text{Cl}(D)$ and denote by $r_2 := k - 1$ the 2-rank. Then the maximal possible class group of exponent c improves to

$$(\mathbb{Z}/c\mathbb{Z})^{r_4} \times (\mathbb{Z}/2\mathbb{Z})^{r_2 - r_4} \text{ of order } 2^{r_2 - r_4} c^{r_4}.$$

This gives an improvement by factor $(c/2)^{r_2 - r_4}$.

5.1. Redei matrices. In this section we want to study the 4-rank of the class group of quadratic number fields. These things are well known and based on the works of Redei. In the following we use the Kronecker symbols $\left(\frac{D}{p}\right)$ and corresponding to $D = p_1^* \cdots p_k^*$ we define the matrix $M = (c_{ij}) \in \mathbb{F}_2^k$ via

$$(-1)^{c_{ij}} := \left(\frac{p_j^*}{p_i}\right) \text{ for } 1 \leq j \neq i \leq k \text{ and } c_{ii} := \sum_{j=1, j \neq i}^k c_{ij}, 1 \leq i \leq k.$$

Theorem 5 (Redei). *Let $D = p_1^* \cdots p_k^*$. Then $\text{rk}_4(\text{Cl}(D)) = k - 1 - \text{Rank}(M)$.*

See [10] 10.b, [8] (2.7).

The relation $c_{ii} := \sum_{j=1, j \neq i}^k c_{ij}$ for $1 \leq i \leq k$ shows that the last column of M is the sum of the first $k - 1$ columns and therefore dependent from the first $k - 1$ columns. Furthermore we see by the multiplicativity of the Kronecker symbol that

$$(-1)^{c_{ii}} = \left(\frac{\prod_{j=1, j \neq i}^k p_j^*}{p_i}\right).$$

In order to understand this matrix, the following lemma is helpful. It deals also with the prime 2 except when $2^* = -4$.

Lemma 8. *Let p_i and p_j different prime numbers such $p_i^* \neq -4 \neq p_j^*$. Then*

$$(2) \quad \left(\frac{p_i^*}{p_j}\right) \left(\frac{p_j^*}{p_i}\right) = \begin{cases} 1 & p_i^* > 0 \text{ or } p_j^* > 0 \\ -1 & p_i^* < 0 \text{ and } p_j^* < 0. \end{cases}$$

The proof is straightforward from the reciprocity law. From this we see that if we have two different primes congruent to 3 modulo 4 dividing D then we get an entry 1 in our matrix. Furthermore the matrix is not symmetric.

In our algorithm we compute a lot of Redei matrices and the corresponding ranks. In order to simplify these computations we look when the last row is the sum of the first $k - 1$ rows.

Lemma 9. *Let $D = p_1^* \cdots p_k^* < 0$ be a product of fundamental discriminants with $p_1 < p_2 < \dots < p_k$ and denote by M the Redei matrix and by the vector (d_1, \dots, d_k) the sum of the rows of M .*

Assume that $p_1^ \neq -4$. Then the vector (d_1, \dots, d_k) is zero. In the case $p_1^* = -4$ we define $\tilde{D} := -D/4$ and we get*

$$\left((-1)^{d_1}, \dots, (-1)^{d_k}\right) = \left(\left(\frac{2}{\tilde{D}}\right), \left(\frac{2}{p_2}\right), \dots, \left(\frac{2}{p_k}\right)\right).$$

Proof. The sum of the j -th column is 0 if and only if the product of the corresponding symbols in the exponents are 1. We get for this product:

$$(3) \quad \prod_{i=1, i \neq j}^k \left(\frac{p_j^*}{p_i}\right) \cdot \left(\frac{\prod_{i=1, i \neq j}^k p_i^*}{p_j}\right) = \prod_{i=1, i \neq j}^k \left(\frac{p_j^*}{p_i}\right) \left(\frac{p_i^*}{p_j}\right).$$

Let us assume that $p_1^* \neq -4$. Therefore we are able to apply Lemma 8. If $p_j^* > 0$ then all factors are 1. If $p_j^* < 0$ then the number of negative $p_i^* \neq p_j^*$ is even and therefore the product is 1.

It remains to study the case $p_1^* = -4$. If we multiply the product (3) with $\prod_{i=2}^k \left(\frac{2}{p_i}\right)$ if $j = 1$ and with $\left(\frac{2}{p_j}\right)$ for $j \neq 1$ we get the product in the situation that $2^* = -8$ and we know that this product is 1. Therefore the product is like expected. \square

Note that in the case $p_1^* = -4$ we only get row sum 0, if all odd prime divisors are congruent to $\pm 1 \pmod{8}$.

Using Theorem 5 we know that $\text{rk}_4(\text{Cl}(D)) = k - 1 - \text{Rank}(M)$. Therefore we are interested to get good lower bounds for the rank of the Redei matrix M . Denote by t the number of negative p_i^* . Then we get $\text{Rank}(M) \geq (t - 1)/2$, see [11]. This paper also discusses the cases where this bound is sharp.

5.2. Using the Redei matrix. In this section we describe an algorithm to compute all fields with exponent $c = 2^r$, where we focus on the cases $c = 4, 8$. In order to use the Redei matrices we restrict to discriminants with exactly k prime factors. Using Lemma 5 we can give an upper bound for the number of prime factors for fields of exponent 2^r which is valid for all fields with at most one exception. In the example after Theorem 4 we get 11, 24, and 58 for the exponents 2, 4, 8, respectively. We remark that these bounds can be improved, but the following algorithm is very efficient for the cases close to the bound. For a given exponent $c = 2^r$ we call the following algorithm for all $k \geq 1$ up to the computed upper bound.

For the algorithm we have to make the decision if we want to use the lower bound of Lemma 3 or of Lemma 5. The first lemma has the advantage that the bound is better and therefore the computation will be faster. In this case we only compute all wanted fields which have no Siegel zero. If we assume that no Siegel zeros exist, then this computation is complete. The bound of the second lemma is weaker, but it has the advantage that we miss at most one field (if it exists, it has a Siegel zero). In our range for exponent 8, this bound is about a factor 2 weaker than the first bound. In order to simplify the presentation we only give the description of the algorithm using the bound of Lemma 5.

The main algorithm to call is Algorithm 4. This algorithm computes the global variables B_0, \dots, B_k which will be used in Algorithms 1 and 3. The main idea of the following algorithms is that the knowledge of a factor of D gives some partial information on the Redei matrix. This information can be used to give an upper bound on the 4-rank of the class group which then gives improved bounds on the maximal possible discriminant. In theory we expect that the average 4-rank is close to 1 [8] and therefore the upper bound c^{k-1} for the class number of a field with exponent c is quite pessimistic. The described approach improves the upper bound of the class number, and therefore the maximal possible discriminant, when the lower rank bound of the Redei matrix increases, and therefore the possible 4-rank decreases.

The goal of the first algorithm is to give a quick check, if the fundamental discriminant $D = p_1^* \cdots p_k^*$ has an exponent which is a divisor of c . The correctness of the algorithm is obvious by using Lemma 1. If possible, we try to avoid the actual computation of the class group.

Algorithm 1. (Check(c, p_1^*, \dots, p_k^*))

- Input: Exponent $c = 2^r$, prime fundamental discriminants p_1^*, \dots, p_k^* .
- Output: Return true, iff $D = p_1^* \cdots p_k^* < 0$ and the exponent of the class group of $\mathbb{Q}(\sqrt{D})$ is a divisor of c .
- Step 1: If $D := p_1^* \cdots p_k^* > 0$ then return false.
- Step 2: Compute the smallest prime q which is split in $K := \mathbb{Q}(\sqrt{D})$.
- Step 3: If $q^c < |D|/4$ then return false (see Lemma 1).
- Step 4: If the c -th power of a prime ideal above q is not principal, then return false. This test can be done most efficiently by using binary quadratic forms.
- Step 5: Repeat the test of Step 4 with the 2nd smallest splitting prime.
- Step 6: Compute the rank s of the Redei matrix of D . If $|D| > B_s$ then return false (B_s is a global variable computed in Algorithm 4).
- Step 7: Compute the class group of K . If the exponent divides c then return true, otherwise return false.

Let $D = p_1^* \cdots p_k^*$ be a negative fundamental discriminant and assume that p_1^*, \dots, p_ℓ^* are known to us. Let M be the Redei matrix of D defined in Section 5.1. Denote by N the minor defined by the first ℓ rows and first ℓ columns. Trivially, we get that $\text{Rank}(N) \leq \text{Rank}(M)$. Since p_1^*, \dots, p_ℓ^* are known to us, we can compute all entries of N except the diagonal. The following algorithm tests all possible combinations for the diagonal and therefore computes a lower bound for the rank of N and M .

Algorithm 2. (LowerRedeiBound(p_1^*, \dots, p_ℓ^*))

- Input: Prime fundamental discriminants p_1^*, \dots, p_ℓ^* with $\ell < k$.
- Output: Returns a lower bound for the rank of the Redei matrix of all D where D has exactly k prime factors and $p_1^* \cdots p_\ell^* \mid D$.

- Step 1: For all $1 \leq i \neq j \leq \ell$ compute c_{ij} via $(-1)^{c_{ij}} := \left(\frac{p_j^*}{p_i}\right)$.
- Step 2: For all $(a_1, \dots, a_\ell) \in \mathbb{F}_2^\ell$ compute the rank of the matrix $D = (d_{ij})$, where $d_{ij} = c_{ij}$ for $i \neq j$ and $d_{ii} = a_i$.
- Step 3: Return the minimal rank computed in Step 2.

In the following algorithm we denote by $P[i]$ the i -th prime number. We update the lower bound of the rank of the Redei matrix and append the next fundamental prime discriminant. This function will call itself recursively. The number m is the index of the smallest prime that can be used next.

Algorithm 3. (NextTuple(m,Discs,k,c))

Input: Number m of the next prime to use, list Discs = $[p_1^*, \dots, p_\ell^*]$ of fundamental prime discs, k , exponent $c = 2^r$

Output: List of all discriminants D with k prime factors such that $p_1^* \cdots p_\ell^* \mid D$ (with at most one exception)

- Step 1: $s := \text{LowerRedeiBound}(p_1^*, \dots, p_\ell^*)$.
- Step 2: $B := B_s$, $\text{res} := []$ (empty list), $\text{bound} := B/(|p_1^* \cdots p_\ell^*|)$, $i := m$, $C := P[i] \cdots P[i + k - \ell - 1]$.
- Step 3: While $C \leq \text{bound}$ do
- 1 $p_{\ell+1}^* := (-1)^{(P[i]-1)/2} P[i]$.
 - 2 If $k = \ell + 1$ then call $\text{Check}(c, p_1^*, \dots, p_k^*)$ and append $D = p_1^* \cdots p_k^*$ to res , if the check is successful.
 - 3 If $k > \ell + 1$ then call $\text{NextTuple}(i, \{p_1^*, \dots, p_{\ell+1}^*\}, k, c)$ and append the computed D 's to res .
 - 4 $i := i + 1$, $C := P[i] \cdots P[i + k - \ell - 1]$.

Step 4: Return res .

In the main algorithm we compute the global variables B_0, \dots, B_{k-1} and we split the computation into four parts, depending on the behavior at 2.

Algorithm 4. (Computation of fields with exponent 2^r)

Input: Exponent 2^r , number of prime factors k

Output: All fields with exponent $c = 2^r$ with at most one exception

- Step 1: By numerical approximation compute a bound B_ℓ for all $0 \leq \ell \leq k - 1$ such that for all $|D| > B_\ell$ we get that the minimum of (1) is greater than $2^\ell \cdot c^{k-1-\ell}$.
- Step 2: Call $\text{res1} := \text{NextTuple}(2, \{1\}, k, c)$.
- Step 3: Call $\text{res2} := \text{NextTuple}(2, \{-4\}, k, c)$.
- Step 4: Call $\text{res3} := \text{NextTuple}(2, \{-8\}, k, c)$.
- Step 5: Call $\text{res4} := \text{NextTuple}(2, \{8\}, k, c)$.
- Step 6: Return the discriminants from res1 , res2 , res3 , res4 .

We remark that there are obvious improvements in the implementation which we have not described here for simplification. Note that there might be one missing example for the overall algorithm by using Lemma 5. The reason is that we compute all fields with exponent dividing c assuming the bounds of Lemma 5. Therefore missing examples have the property that the bounds of Lemma 5 are wrong and this can happen at most one time.

The algorithm described above only computes discriminants D such that $|D| > 10^6$. We could easily reduce this lower bound, but this lower bound is not an issue since there are known tables of class groups for all quadratic fields of small discriminant. The web-page [2] gives all fields up to 10^7 .

The overall running time of our algorithm on one core is about 17 hours when we use the estimates from Lemma 4. The cases $k \in \{30, \dots, 58\}$ take about 50 seconds. Only for small k we need to compute class groups. The cases $k \leq 6$ take about half an hour. The most expensive cases are $k \in \{8, \dots, 12\}$ which take more one hour each, the worst case being $k = 10$ which takes almost 3 hours.

When we use the estimates from Lemma 5 which are sufficient to prove that we miss at most one example, then the running time is about 60 hours. In case we can take a bound which is a factor 2 better than the bound in Lemma 4, the running time improves to less than 3 hours.

6. DIRECT SEARCHING FOR SMALL DISCRIMINANTS

Lemma 10. *For a prime p and an exponent c there are less than $2p^{c/2}$ imaginary quadratic fields $K = \mathbb{Q}(\sqrt{D})$, such that p splits in K and the ideal above p has order divisible by c in the class group.*

Proof. The assumption implies that the equation $4p^c = x^2 + |D|y^2$ has an integer solution with $x \cdot y \neq 0$. Thus, the field K is one of $\mathbb{Q}(\sqrt{-(4p^c - x^2)})$ for $x \in \mathbb{Z}$ and $4p^c - x^2 > 0$. This shows $1 \leq x < 2p^{c/2}$. Thus, there are less than $2p^{c/2}$ fields. \square

Remarks 1. 1 *The lemma above results in an algorithm to enumerate all imaginary quadratic fields such that p is a split prime and the exponent of the class group is a divisor of c . It's complexity is $O(p^{c/2})$. More precisely, we first compute a finite list of fields that is a superset of the fields we are searching. The superfluous fields can easily be removed in a second step.*

2 *As a slight variation, we can enumerate all quadratic fields such that p is the smallest split prime and the exponent of the class group is a divisor of c . For this we just have to sift out all those fields that have a smaller split prime. Note that this can be done without factoring $Dy^2 = 4p^c - x^2$.*

3 *A C-implementation of this approach lists all the fields with smallest split prime ≤ 197 and $c = 8$ in less than 4 minutes. We find 268 fields with exponent 1,2,4 and 778 fields with exponent 8. The largest one is $\mathbb{Q}(\sqrt{-430950520})$.*

4 *Doing the same computation with exponent 3, 5, 6, 7 takes less than a minute.*

5 *An imaginary quadratic field with class group exponent ≤ 8 not listed has smallest split prime > 193 .*

6 *Searching for imaginary quadratic fields with $|D|$ up to a given bound and smallest split prime > 193 can be done by a multiply-focused enumeration similar to [14]. I.e. we have to sift out all those D that have a small split prime.*

Remarks 2. We want to search for fields with no split prime $p \leq 193$ and $|D| < 3 \cdot 10^{20}$. This can be done by sieving. To maximize the speed we have to use bit-level operations and tables of pre-computed data. Further, non negative integers $< 2^{64}$ have the fastest arithmetic. Thus, the main loop should be restricted to this.

This results in the following approach:

- We want to search for all imaginary quadratic fields $\mathbb{Q}(\sqrt{D})$ with smallest split prime > 193 . This implies $\left(\frac{D}{p}\right) \neq 1$ for all primes $p \leq 193$.
- Using the chinese remainder theorem with the module $m = 16 \cdot 3 \cdot 5 \cdots 47 \approx 4.9 \cdot 10^{18}$ the number of possible residue classes for $|D|$ is $220172127436800 \approx 2.2 \cdot 10^{14}$.
- Let r be the smallest non-negative representative of a feasible residue class mod m . We have to test the fields with $|D| = r, r + m, r + 2m, \dots, r + 63m$ in parallel.
- The prime 53 rules out a field if $\left(\frac{D}{53}\right) = 1$. For $|D| = r, r + m, r + 2m, \dots, r + 63m$, we can encode this in a sequence of 64 bits. The k -th bit is 1 iff the field for $|D| = r + km$ is not ruled out by $\left(\frac{-r - km}{53}\right) = 1$.

We tabulate these bit-sequences for each residue class of $r \pmod{53}$.

- Similarly each other prime $p \leq 193$ and each residue class $r \pmod{p}$ we get a sequence of 64 bits. The k -th bit is 1 iff the field for $|D| = r + km$ is not ruled out by $\left(\frac{-r - km}{p}\right) = 1$.

This gives us further tables of bit-sequences. One table for each prime p with one entry of 64 bits for each residue class of $r \pmod{p}$.

- To combine the information modulo the various primes, we have to pick those $|D|$, that are not ruled out by one of the primes up to 193. In the language of bit sequences this means that we have to do logical **and** of the sequences.
- If a result bit of the **and** is 0, the corresponding field is ruled out.
- If a result bit is 1, the field $\mathbb{Q}(\sqrt{-r - km})$ needs a more detailed inspection.

This approach leads to the following algorithm.

Algorithm 5. (Multifocused bit-vector sieve)

Input: No input.

Output: Print all imaginary quadratic fields $\mathbb{Q}(\sqrt{D})$ without smallest split prime $p > 193$, $|D| < 3.1 \cdot 10^{20}$ and $4p^8 > |D|$.

Step 1: Set the modules $m_1 = 3 \cdot 5 \cdot 7 \cdot 11 \cdot 13 \cdot 17 = 255255$, $m_2 = 19 \cdot 23 \cdot 29 \cdot 31 = 392863$, $m_3 = 37 \cdot 41 \cdot 43 \cdot 47 = 3065857$, and $m = 16m_1m_2m_3 \approx 4.9 \cdot 10^{18}$.

Step 2: For each module m_i compute a list of the integers r in $\{0..m_i - 1\}$ such that $\left(\frac{-r}{p}\right) \neq 1$ for all primes p dividing m_i .

Step 3: For each prime $p \leq 193$ not dividing m set up a list $(l_p[0], \dots, l_p[p-1])$ of p bit-vectors of length 64. The k -th bit in $l_p[i]$ is 0 if and only if $\left(\frac{-(i+km)}{p}\right) = 1$.

Step 4: In a quadruple loop run over the cartesian product of the 3 lists computed in step 2 and $\{3, 4, 8, 11 \pmod{16}\}$ and do the following:

- a) Use the chinese remainder theorem to find the unique integer $0 \leq r < m$ congruent to the given residues modulo $m_1, m_2, m_3, 16$.
- b) Do a logical **and** of the bit-vectors $l_p[r \bmod p]$ for all prime p with $53 \leq p \leq 193$.
- c) If the k -th bit of the resulting bit-vector is 1, the field $\mathbb{Q}(\sqrt{D})$ with $D := -r - k \cdot m$ is suspicious.
- d) Compute the smallest split prime p for each suspicious field $\mathbb{Q}(\sqrt{D})$. If $4p^8 > |D|$ then print the field.

Remark 1. The algorithm above finds all the imaginary quadratic fields $\mathbb{Q}(\sqrt{D})$ with $|D| < 3.14 \cdot 10^{20}$, no split-prime ≤ 193 . We print out only those fields that may have a class group exponent ≤ 8 . The run time on a single core on an Intel i5 processor is about 40 days. However, we can the loop over the cartesian product in parallel. The result is as follows:

- The algorithm above results in 1002279 imaginary quadratic fields.
- None of them has a class group of exponent ≤ 100 .

REFERENCES

- [1] E. Bach and J. Sorenson, Explicit bounds for primes in residue classes. *Math. Comput.* 65(216), 1717–1735, 1996.
- [2] M. Boy and G. Malle, A Database for Class Groups of Number Fields. http://www.mathematik.uni-kl.de/~numberfieldtables/KT_2/doc.html
- [3] D. W. Boyd and H. Kisilevsky, On the exponent of the ideal class groups of complex quadratic fields. *Proc. Amer. Math. Soc.* 31, 433–436, 1972.
- [4] Y.-G. Chen, On the Siegel-Tatuzawa-Hoffstein theorem. *Acta Arith.* 130, 361–367, 2007.
- [5] M. Deuring, Imaginäre quadratische Zahlkörper mit der Klassenzahl Eins. *Inv. Math.* 5, 169–179, 1968.
- [6] A. G. Earnest and O. H. Körner, On ideal class groups of 2-power exponent. *Proc. Amer. Math. Soc.* 86(2), 196–198, 1982.
- [7] E. Fouvry and J. Klüners, On the 4–rank of quadratic number fields. *Inv. Math.* 167, 455–516, 2007.
- [8] F. Gerth, The 4–class ranks of quadratic fields. *Inv. Math.* 77, 489–515, 1989.
- [9] D. R. Heath-Brown, Imaginary quadratic fields with class group exponent 5. *Forum Math.* 20(2), 275–283, 2008.
- [10] L. Rédei, Arithmetischer Beweis des Satzes über die Anzahl der durch vier teilbaren Invarianten der absoluten Klassengruppe im quadratischen Zahlkörper. *J. Reine Angew. Math.* 171, 55–60, 1938.
- [11] Y. Sueyoshi, On Rédei matrices with minimal rank. *Far East J. Math. Sci. (FJMS)*, 3, 121–128, 2001.
- [12] T. Tatuzawa, On a theorem of Siegel. *Jap. J. Math* 21, 163–178, 1952.
- [13] P. J. Weinberger, Exponents of the class groups of complex quadratic fields. *Acta Arith.* 22, 117–124, 1973.
- [14] K. Wooding and H. C. Williams, Doubly-focused enumeration of pseudosquares and pseudocubes. *LNCS* 4076, 208 – 221, 2006.

UNIVERSITÄT PADERBORN, FAKULTÄT EIM, INSTITUT FÜR MATHEMATIK, WARBURGER STR. 100,
33098 PADERBORN, DEUTSCHLAND

E-mail address: elsenhan@math.uni-paderborn.de

UNIVERSITÄT PADERBORN, FAKULTÄT EIM, INSTITUT FÜR MATHEMATIK, WARBURGER STR. 100,
33098 PADERBORN, DEUTSCHLAND

E-mail address: klueners@math.uni-paderborn.de

“SIMION STOILOW” INSTITUTE OF MATHEMATICS OF THE ROMANIAN ACADEMY, P.O.BOX 1-
764,RO-014700 BUCHAREST, ROMANIA

E-mail address: florin.nicolae@imar.ro