

A QUANTUM PROBABILISTIC APPROACH TO HECKE ALGEBRAS FOR p -ADIC PGL_2

TAKEHIRO HASEGAWA, HAYATO SAIGO, SEIKEN SAITO, AND SHINGO SUGIYAMA

ABSTRACT. The subject of the present paper is an application of quantum probability to p -adic objects. We give a quantum-probabilistic interpretation of the spherical Hecke algebra for $\mathrm{PGL}_2(F)$, where F is a p -adic field. As a byproduct, we obtain a new proof of the Fourier inversion formula for $\mathrm{PGL}_2(F)$.

INTRODUCTION

Hecke operators acting on the space of modular forms are central objects in number theory. There are stochastic problems of eigenvalues of Hecke operators. As a remarkable example, the Sato-Tate conjecture (proved in [7], [6] and [3]) is known. This states that, eigenvalues of all Hecke operators $\{T_p\}_p$ with respect to a fixed simultaneous eigenvector are equidistributed in the interval $[-2, 2]$ with respect to the probability measure

$$\frac{\sqrt{4-x^2}}{2\pi} dx.$$

Before the Sato-Tate conjecture was resolved, as a deduction from the Eichler-Selberg trace formula, Serre [14] had found out the vertical Sato-Tate law: For a fixed p -th Hecke operator T_p for a prime number p , all eigenvalues of T_p are equidistributed in the interval $[-2, 2]$ with respect to the probability measure

$$\frac{p+1}{(p^{1/2}+p^{-1/2})^2-x^2} \frac{\sqrt{4-x^2}}{2\pi} dx.$$

Each of two measures as above is called the Wigner semicircle distribution and the Kesten distribution with parameter $p+1$, respectively. The Kesten distribution with parameter $p+1$ is understood as the spherical Plancherel measure for $\mathrm{PGL}_2(\mathbb{Q}_p)$ and appears in studies of modular forms and their L -functions (e.g., [11], [5], [21], [16], [17], [18] and [19]).

The Wigner semicircle distribution and the Kesten distributions have been studied in a framework of quantum probability theory, which captures a classical random variable as an element of a unital $*$ -algebra not necessarily commutative. A quantum probability space is a pair (\mathcal{A}, φ) of a unital $*$ -algebra \mathcal{A} and a state $\varphi : \mathcal{A} \rightarrow \mathbb{C}$. The state φ is a substitute of expectation, and (\mathcal{A}, φ) recovers classical probability theory when \mathcal{A} is commutative. It is well-known that the m -th moment $\varphi(a^m)$ for $a \in \mathcal{A}$ is described by that of a probability measure and an interacting Fock space (cf. [1]), on which an analogue of decomposition of the quantum harmonic oscillator into the creation and the annihilation operators are

2010 *Mathematics Subject Classification.* Primary 46L53; Secondary 33D80.

Key words and phrases. Quantum probability, spherical Hecke algebras, Fourier inversion formulas.

discussed. Such a decomposition is applied to analysis of asymptotic spectral distributions of graphs (cf. [2], [8] and [10]) and to quantum-classical correspondence asymptotically controlled by the arcsine law (cf. [12] and [13]).

The consideration above inspires us to give a new insight into the Hecke operator T_p in terms of quantum probability theory. In this paper, we treat the p -adic group $G = \mathrm{PGL}_2(\mathbb{Q}_p)$ for a prime number p and its spherical Hecke algebra $\mathcal{H}(G, K)$ consisting of all $\mathrm{PGL}_2(\mathbb{Z}_p)$ -biinvariant \mathbb{C} -valued functions on G with compact support, keeping in mind that the Hecke operator T_p on modular forms corresponds to an action to adelicized modular forms of an element $T(p\mathbb{Z}_p)$ of $\mathcal{H}(G, K)$ (cf. [4, §3.6]). Our new contribution in this paper is to give an interacting Fock space structure to $\mathcal{H}(G, K)$ and a quantum decomposition of the $T(p\mathbb{Z}_p)$ -multiple (see Theorem 4). As a corollary, we give a new proof of the Fourier inversion formula for $\mathrm{PGL}_2(\mathbb{Q}_p)$ originally given by Macdonald [9]. The key of his proof is a concrete calculation of integrals concerned with the explicit formula (3.1) of spherical function. In contrast to his proof, we use quantum probability theory and intrinsic properties of spherical functions, and prove the inversion formula without explicit forms of those functions. Throughout this paper, we describe our results for any non-archimedean local field F in place of \mathbb{Q}_p .

1. PRELIMINARIES

Let \mathbb{N} denote the set of all positive integers and set $\mathbb{N}_0 = \mathbb{N} \cup \{0\}$. For a set X and its subset A , ch_A denotes the characteristic function of A on X .

Let F be a non-archimedean local field, \mathfrak{o} its integer ring and \mathfrak{p} the maximal ideal of \mathfrak{o} . The cardinality of the residue field $\mathfrak{o}/\mathfrak{p}$ is denoted by q . We fix a uniformizer ϖ of F . For example, the p -adic field $F = \mathbb{Q}_p$ for a prime number p is a non-archimedean local field with characteristic 0. In this case, we have $\mathfrak{o} = \mathbb{Z}_p$, $\mathfrak{p} = p\mathbb{Z}_p$ and $q = p$. The prime number p is a uniformizer of \mathbb{Q}_p .

Set $G = \mathrm{PGL}(2, F)$ and $K = \mathrm{PGL}(2, \mathfrak{o})$. The unit element of G is denoted by 1_2 . Then K is a maximal compact subgroup of G and the Cartan decomposition $G = \coprod_{n=0}^{\infty} K \begin{bmatrix} \varpi^n & 0 \\ 0 & 1 \end{bmatrix} K$ holds (cf. [4, Proposition 4.6.2]). We take a Haar measure dg on G such that the volume of K equals 1.

Let B be the Borel subgroup of G consisting of the upper triangular matrices. For $s \in \mathbb{C}$, we denote by $\pi_s = \mathrm{Ind}_B^G(|\cdot|^s \boxtimes |\cdot|^{-s})$ the normalized parabolic induction whose representation space V_s consists of all \mathbb{C} -valued locally constant functions f on G satisfying $f(\begin{bmatrix} a & b \\ 0 & d \end{bmatrix} g) = |\frac{a}{d}|^{s+1/2} f(g)$ for all $g \in G$ and $\begin{bmatrix} a & b \\ 0 & d \end{bmatrix} \in B$ and the G -action is the right translation. Set $\mathfrak{X} = i[0, 2\pi(\log q)^{-1}]$. If $s \in \mathfrak{X}$, then π_s is an irreducible unitarizable representation of G and the K -invariant subspace of V_s is one dimensional. We remark that the representations π_s ($s \in \mathfrak{X}$) exhaust the unitary representations appearing in the spectral decomposition of $L^2(G/K)$.

Let $\mathcal{H}(G)$ be the Hecke algebra of G , that is, the space of all \mathbb{C} -valued locally constant functions with compact support. This is a non-commutative associative algebra over \mathbb{C} without unit, where the product $f_1 \circ f_2$ for $f_1, f_2 \in \mathcal{H}(G)$ is the convolution defined by

$$(f_1 \circ f_2)(g) = \int_G f_1(gx^{-1})f_2(x)dx, \quad g \in G.$$

Then $\mathcal{H}(G)$ has a $*$ -algebra structure by $f^*(g) = \overline{f(g^{-1})}$. The $*$ -algebra $\mathcal{H}(G)$ acts on $C(G)$ by right translation R :

$$(R(f)\varphi)(g) = \int_G f(x)\varphi(gx)dx, \quad f \in \mathcal{H}(G), \varphi \in C(G).$$

Let $\mathcal{H}(G, K)$ denote the subalgebra of $\mathcal{H}(G)$ consisting of all K -biinvariant functions in $\mathcal{H}(G)$. It has a $*$ -subalgebra of $\mathcal{H}(G)$ and commutative (cf. [4, Theorem 4.6.1]). Furthermore, $\mathcal{H}(G, K)$ has a unit ch_K . Remark that $f(g) = \overline{f(g^{-1})}$ for any $f \in \mathcal{H}(G, K)$ because of $K[\varpi_0^{-n} \ 0]K = K[\varpi_0^n \ 0]K$. In particular, we have $f_1^* = \overline{f_1}$ and $f_1 \circ f_2 = R(f_2)f_1$ for any $f_1, f_2 \in \mathcal{H}(G, K)$. The L^2 -inner product on $L^2(G, dg)$ is denoted by $\langle \cdot, \cdot \rangle_G$.

2. HECKE OPERATORS

For any $n \in \mathbb{N}_0$, let $T(\mathfrak{p}^n)$ be the characteristic function of

$$Z \backslash Z\{g \in M_2(\mathfrak{o}) \mid \det(g)\mathfrak{o} = \mathfrak{p}^n\},$$

where Z is the center of $\text{GL}_2(F)$. Then, $T(\mathfrak{p}^0) = \text{ch}_K$ holds. The action of $T(\mathfrak{p}^n)$ as $R(T(\mathfrak{p}^n))$ is called the Hecke operator at \mathfrak{p}^n . The following is a well-known recurrence equation (cf. [4, Proposition 4.6.4]).

Lemma 1. *For any $n \in \mathbb{N}$, we have*

$$T(\mathfrak{p}) \circ T(\mathfrak{p}^n) = T(\mathfrak{p}^{n+1}) + qT(\mathfrak{p}^{n-1}).$$

Let Ψ_n be the characteristic function of $\mathbf{K}[\varpi_0^n \ 0]\mathbf{K}$ on G for any $n \in \mathbb{N}_0$. Then we have the relation $T(\mathfrak{p}^n) = \sum_{r=0}^{\lfloor n/2 \rfloor} \Psi_{n-2r}$ in the same way as [15, Lemma 6]. By solving this on Ψ_n , we have the following (cf. [15, Remark 7]).

Lemma 2. *We have $\Psi_0 = \text{ch}_K$, $\Psi_1 = T(\mathfrak{p})$, and $\Psi_n = T(\mathfrak{p}^n) - T(\mathfrak{p}^{n-2})$ for $n \geq 2$.*

Set $T'(\mathfrak{p}) = q^{-1/2}T(\mathfrak{p})$ and $\Phi_n = q^{-n/2}\Psi_n$ ($n \in \mathbb{N}_0$).

Lemma 3. *We have the relations*

$$T'(\mathfrak{p}) \circ \Phi_n = \Phi_{n+1} + \Phi_{n-1} \quad (n \geq 2),$$

$$T'(\mathfrak{p}) \circ \Phi_1 = \Phi_2 + \frac{q+1}{q}\Phi_0.$$

Proof. We use Lemmas 1 and 2. For $n = 1$, a direct computation gives us

$$T'(\mathfrak{p})\Phi_1 = \frac{1}{q}T(\mathfrak{p})^2 = \frac{1}{q}(T(\mathfrak{p}^2) + qT(\mathfrak{p}^0)) = \frac{1}{q}(T(\mathfrak{p}^2) - T(\mathfrak{p}^0)) + \frac{1+q}{q}T(\mathfrak{p}^0) = \Phi_2 + \frac{q+1}{q}\Phi_0.$$

For any $n \geq 3$, we obtain

$$\begin{aligned} T'(\mathfrak{p})\Phi_n &= q^{-1/2}T(\mathfrak{p}) \circ q^{-n/2}(T(\mathfrak{p}^n) - T(\mathfrak{p}^{n-2})) \\ &= q^{-(n+1)/2}(T(\mathfrak{p}^{n+1}) + qT(\mathfrak{p}^{n-1}) - T(\mathfrak{p}^{n-1}) - qT(\mathfrak{p}^{n-3})) \\ &= q^{-(n+1)/2}(T(\mathfrak{p}^{n+1}) - T(\mathfrak{p}^{n-1})) + q^{-(n-1)/2}(T(\mathfrak{p}^{n-1}) - T(\mathfrak{p}^{n-3})) \\ &= \Phi_{n+1} + \Phi_{n-1}. \end{aligned}$$

The case $n = 2$ is similarly computed. Thus we are done. □

Recall that $\mathcal{H}(G, K)$ is a pre-Hilbert space with respect to the L^2 -inner product $\langle \cdot, \cdot \rangle_G$. The Cartan decomposition of G yields that $\{\Phi_n\}_n$ is an orthogonal system and generates $\mathcal{H}(G, K)$ because of $f(g) = \sum_{n=0}^{\infty} f\left(\begin{smallmatrix} \varpi_0^n & 0 \\ 0 & 1 \end{smallmatrix}\right) \Psi_n$.

Therefore we can define B^+ and B^- in $\text{End}(\mathcal{H}(G, K))$ by

$$B^+ \Phi_0 = \sqrt{\frac{q+1}{q}} \Phi_1, \quad B^+ \Phi_n = \Phi_{n+1} \quad (n \geq 1),$$

$$B^- \Phi_0 = 0, \quad B^- \Phi_1 = \sqrt{\frac{q+1}{q}} \Phi_0, \quad B^- \Phi_n = \Phi_{n-1} \quad (n \geq 2).$$

Set $B^0 = 0 \in \text{End}(\mathcal{H}(G, K))$. Combining these with Lemma 3, we have the following.

Theorem 4. *The quintuple $(\mathcal{H}(G, K), \{\Phi_n\}_n, B^+, B^-, B^0)$ is an interacting Fock space. The associated Jacobi coefficient $(\{\omega_n\}_n, \{\alpha_n\}_n)$ is given by $\omega_1 = \frac{q+1}{q}$, $\omega_n = 1$ for $n \geq 2$, and $\alpha_n = 0$ for all $n \in \mathbb{N}_0$. Furthermore, we have the quantum decomposition*

$$R(T'(\mathfrak{p})) = B^+ + B^- + B^0 \in \text{End}(\mathcal{H}(G, K)).$$

The operators B^+ , B^- and B^0 are called the creation, the annihilation, and the preservation operators, respectively.

Let $\varphi : \mathcal{H}(G, K) \rightarrow \mathbb{C}$ be the vector state for Φ_0 :

$$\varphi(f) = \langle R(f)\Phi_0, \Phi_0 \rangle_G, \quad f \in \mathcal{H}(G, K).$$

Then $(\mathcal{H}(G, K), \varphi)$ is a quantum probability space and $T'(\mathfrak{p})$ is a real random variable in $(\mathcal{H}(G, K), \varphi)$. This yields the following equality of m -th moments.

Corollary 5. *For any $m \in \mathbb{N}_0$, we have*

$$\varphi(T'(\mathfrak{p})^m) = \langle R(T'(\mathfrak{p}))^m \Phi_0, \Phi_0 \rangle_G = \int_{-\infty}^{\infty} x^m \frac{q+1}{(q^{1/2} + q^{-1/2})^2 - x^2} \frac{\sqrt{4-x^2}}{2\pi} \text{ch}_{[-2,2]}(x) dx.$$

Proof. This is a consequence of [10, Theorem 4.11]. To complete the proof, it suffices to determine a probability measure μ on \mathbb{R} associated with $(\{\omega_n\}_n, \{\alpha_n\}_n)$. In this case, as in [10, p.96], μ is the free Meixner distribution with parameter $(\frac{q+1}{q}, 1, 0)$, i.e.,

$$\int_{-\infty}^{\infty} \frac{1}{z-x} d\mu(x) = \frac{1}{z - \frac{\frac{q+1}{q}}{z - \frac{1}{z - \frac{1}{z - \frac{1}{\ddots}}}}}}, \quad \text{Im}(z) \neq 0.$$

Since the continued fraction as above is periodic, the right-hand side is computed as

$$\frac{(2 - \frac{q+1}{q})z - \frac{q+1}{q} \sqrt{z^2 - 4}}{2(1 - \frac{q+1}{q})z^2 + 2(\frac{q+1}{q})^2},$$

where the square root is taken so that $\sqrt{z} > 0$ for any $z > 0$. Hence the inverse of the Stieltjes transformation yields

$$(2.1) \quad d\mu(x) = \frac{q+1}{(q^{1/2} + q^{-1/2})^2 - x^2} \frac{\sqrt{4-x^2}}{2\pi} \text{ch}_{[-2,2]}(x) dx.$$

Hence, we obtain the desired formula. \square

There exists an orthogonal polynomial system $\{P_n\}_n$ associated with $(\{\omega_n\}_n, \{\alpha_n\}_n)$. By the proof of [10, Theorem 4.11], we have a unitary mapping $\mathcal{U} : \mathcal{H}(G, K) \rightarrow L^2(\mathbb{R}, d\mu)$ by $\Phi_n \mapsto (\prod_{j=1}^n \omega_j)^{-1/2} P_n(x)$ and $R(T'(\mathfrak{p}))$ corresponds to the x -multiple on $L^2(\mathbb{R}, d\mu)$ by \mathcal{U} . The mapping \mathcal{U} induces an isometry $L^2(K \backslash G / K, dg) \cong L^2(\mathbb{R}, d\mu)$.

Remark : In the proof of Corollary 5, we can determine μ also in the following way. By Lemma 1, $T(\mathfrak{p}^n)$ is expressed by $T'(\mathfrak{p})$ as $q^{-n/2} T(\mathfrak{p}^n) = U_n(2^{-1} T'(\mathfrak{p}))$ for any $n \in \mathbb{N}_0$, where U_n is the n -th Chebyshev polynomial of 2nd kind. Combining this with the definition of Φ_n , the family $\{P_n\}_n$ is described as $P_0(x) = 1$, $P_1(x) = U_1(2^{-1}x)$ and $P_n(x) = U_n(2^{-1}x) - q^{-1} U_{n-2}(2^{-1}x)$ for $n \geq 2$. Then, from the computation in [14, §2.3], μ is given by (2.1).

3. SPHERICAL FUNCTIONS AND FOURIER TRANSFORMS

In this section, we show an application of Corollary 5 to the Fourier inversion formula for G/K .

3.1. Fourier inversion formula. Let us review the Fourier inversion formula for G . For $s \in \mathfrak{X}$, there exists $\Omega_s : G \rightarrow \mathbb{C}$ with properties

- (i) $R(T'(\mathfrak{p}))\Omega_s = (q^{-s} + q^s)\Omega_s$,
- (ii) $\Omega_s(k_1 g k_2) = \Omega_s(g)$ for all $k_1, k_2 \in K$ and $g \in G$,
- (iii) $\Omega_s(1_2) = 1$.

Then Ω_s is uniquely determined by three properties, and called a spherical function. The spherical function Ω_s can be given as follows. Let ϕ_0 be the K -invariant vector of π_s determined by $\phi_0 \in V_s^K$ and $\phi_0(1_2) = 1$. The representation π_s has a G -invariant inner product given by $\langle \phi_1, \phi_2 \rangle_K = \int_K \phi_1(k) \overline{\phi_2(k)} dk$ ($\phi_1, \phi_2 \in V_s$). Note $\|\phi_0\|_K = 1$. Then, Ω_s is expressed as $\Omega_s(g) = \langle \pi_s(g)\phi_0, \phi_0 \rangle_K$ (cf. [4, Proposition 4.6.6]). In particular, $s \mapsto \Omega_s(g)$ for any fixed $g \in G$ is analytic and $|\Omega_s(g)| \leq \|\pi_s(g)\phi_0\|_K \|\phi_0\|_K = \|\phi_0\|_K^2 = 1$ holds.

We define the Fourier transformation $\mathcal{F} : \mathcal{H}(G, K) \rightarrow C(\mathfrak{X})$ by

$$\mathcal{F}(f)(s) = \int_G \Omega_s(g) f(g) dg.$$

Let $\mathcal{A} = \mathbb{C}[q^{-s} + q^s] \subset C(\mathfrak{X})$ be the ring of polynomial functions on \mathfrak{X} with respect to $q^{-s} + q^s$. Similarly to \mathcal{F} , we define $\mathcal{F}^* : \mathcal{A} \rightarrow C(G)$ by

$$\mathcal{F}^*(\alpha)(g) = \int_{\mathfrak{X}} \Omega_s(g) \alpha(s) d\nu(s).$$

Here $d\nu(s)$ is a positive Radon measure on \mathfrak{X} given by

$$d\nu(s) = \frac{1 + q^{-1}}{4\pi i} \frac{|1 - q^{-2s}|^2}{|1 - q^{-2s-1}|^2} (\log q) ds.$$

This measure is called the spherical Plancherel measure for G . Remark that the space \mathfrak{X} parametrizes the spherical tempered unitary dual of G . The Fourier inversion formula is stated as follows.

Theorem 6. *The image of \mathcal{F} and of \mathcal{F}^* is contained in \mathcal{A} and $\mathcal{H}(G, K)$, respectively. Both $\mathcal{F} : \mathcal{H}(G, K) \rightarrow \mathcal{A}$ and $\mathcal{F}^* : \mathcal{A} \rightarrow \mathcal{H}(G, K)$ are isometries and each of these is the inverse of the other. Furthermore, both \mathcal{F} and \mathcal{F}^* are extended to isometries between L^2 -spaces;*

$$L^2(K \backslash G / K, dg) \cong L^2(\mathfrak{X}, d\nu).$$

This result as above is a special case of [9, Theorem 5.1.2] (see also [20, Theorem 4.7] and [22, Theorem 3.1]).

3.2. Proof of the Fourier inversion formula. By Corollary 5, we obtain the isometry $\mathcal{U} : \mathcal{H}(G, K) \rightarrow \bigoplus_{n=0}^{\infty} \mathbb{C}P_n$, as was explained in the last of §2. We have a homeomorphism $\mathfrak{X} \cong [-2, 2]$ by $s \mapsto q^{-s} + q^s$, under which $d\nu$ is transformed into $d\mu$ given by (2.1). This leads us naturally to the isometry $\iota : L^2(\mathbb{R}, d\mu) \cong L^2(\mathfrak{X}, d\nu)$.

We give integral representations of \mathcal{U} and of \mathcal{U}^{-1} . The following gives us an alternative proof of Theorem 6.

Theorem 7. *We have $\iota \circ \mathcal{U} = \mathcal{F}$ and $\mathcal{U}^{-1} \circ \iota^{-1} = \mathcal{F}^*$. Namely, we have $\text{Im}(\mathcal{F}) \subset \mathcal{A}$, $\text{Im}(\mathcal{F}^*) \subset \mathcal{H}(G, K)$,*

$$\iota \circ \mathcal{U}(f) = \int_G \Omega_s(g) f(g) dg$$

for any $f \in \mathcal{H}(G, K)$ and

$$\mathcal{U}^{-1} \circ \iota^{-1}(\alpha) = \int_{\mathfrak{X}} \Omega_s(g) \alpha(s) d\nu(s)$$

for any $\alpha \in \mathcal{A}$.

Proof. We have only to compute $\iota \circ \mathcal{U}(T'(\mathfrak{p})^m)$ and $\mathcal{U}^{-1} \circ \iota^{-1}((q^{-s} + q^s)^m)$ for any $m \in \mathbb{N}_0$. By definition, we see

$$\iota \circ \mathcal{U}(T'(\mathfrak{p})^m)(s) = (q^{-s} + q^s)^m = R(T'(\mathfrak{p})^m) \Omega_s(1_2) = \int_G \Omega_s(1_2 g) T'(\mathfrak{p})^m(g) dg$$

for $m \in \mathbb{N}_0$, which leads the assertion for $\iota \circ \mathcal{U}$. Let $\langle \cdot, \cdot \rangle_{\mathfrak{X}}$ denote the L^2 -inner product on $L^2(\mathfrak{X}, d\nu)$. For $m, n \in \mathbb{N}_0$, we see

$$\begin{aligned} & \langle \mathcal{U}^{-1} \circ \iota^{-1}((q^{-s} + q^s)^m) - \int_{\mathfrak{X}} \Omega_s(\cdot) (q^{-s} + q^s)^m d\nu(s), T'(\mathfrak{p})^n \rangle_G \\ &= \langle \mathcal{U}^{-1} \circ \iota^{-1}((q^{-s} + q^s)^m), T'(\mathfrak{p})^n \rangle_G - \langle \int_{\mathfrak{X}} \Omega_s(\cdot) (q^{-s} + q^s)^m d\nu(s), T'(\mathfrak{p})^n \rangle_G \\ &= \langle (q^{-s} + q^s)^m, \iota \circ \mathcal{U}(T'(\mathfrak{p})^n) \rangle_{\mathfrak{X}} - \int_G \int_{\mathfrak{X}} \Omega_s(g) (q^{-s} + q^s)^m d\nu(s) T'(\mathfrak{p})^n(g) dg. \end{aligned}$$

Here we use that both \mathcal{U} and ι are isometries. By the integral representation of $\iota \circ \mathcal{U}$, the first term of the last line of the equalities above is transformed into

$$\int_{\mathfrak{X}} (q^{-s} + q^s)^m \int_G \Omega_s(g) T'(\mathfrak{p})^n(g) dg d\nu(s).$$

Since both \mathfrak{X} and $\text{supp}(T'(\mathfrak{p})^n)$ are compact, Fubini's theorem is applied and the last line of the equalities above vanishes. Since n is arbitrary, the equality

$$\mathcal{U}^{-1} \circ \iota^{-1}((q^{-s} + q^s)^m)(g) = \int_{\mathfrak{X}} \Omega_s(g)(q^{-s} + q^s)^m d\nu(s)$$

holds, from which we obtain the assertion for $\mathcal{U}^{-1} \circ \iota^{-1}$. \square

Remark : By the Macdonald formula [9, Theorem 4.1.2], we have

$$(3.1) \quad \Omega_s\left(\begin{smallmatrix} \varpi^n & 0 \\ 0 & 1 \end{smallmatrix}\right) = \frac{q^{-n/2}}{1+q^{-1}} \left(\frac{1-q^{-1+2s}}{1-q^{2s}} q^{-ns} + \frac{1-q^{-1-2s}}{1-q^{-2s}} q^{ns} \right), \quad n \in \mathbb{N}_0$$

(see also [4, Theorem 4.6.6]). His proof of Theorem 6, which is valid for a general p -adic group, relies on explicit calculations of the integrals $\mathcal{F}(\Psi_m)$ and $\mathcal{F}^*((q^{-s} + q^s)^m)$ (cf. [20] and [22, §5.4]). Contrary to Macdonald's method, we use properties (i) and (iii) of the spherical function Ω_s but not its explicit formula (3.1). Indeed, we first obtain isometries $\iota \circ \mathcal{U}$ and $\mathcal{U}^{-1} \circ \iota^{-1}$ by virtue of quantum probability theory, and finally obtain integral representations \mathcal{F} and \mathcal{F}^* of them.

ACKNOWLEDGEMENTS

The authors would like to thank Hiroki Sako for fruitful discussion. Takehiro Hasegawa was partially supported by JSPS KAKENHI (grant number 15K17508). Hayato Saigo was partially supported by JSPS KAKENHI (grant number 26870696). Seiken Saito was partially supported by JSPS KAKENHI (grant number 16K05259).

REFERENCES

- [1] L. Accardi and M. Bożejko, *Interacting Fock spaces and Gaussianization of probability measures*, Infin. Dimens. Anal. Quantum. Probab. Relat. Top. Vol. **01**, No. 04, 663–670, 1998.
- [2] L. Accardi and N. Obata, *Foundations of Quantum Probability Theory* (in Japanese), Makino Shoten, 2003.
- [3] T. Barnet-Lamb, D. Geraghty, M. Harris and R. Taylor, *A family of Calabi-Yau varieties and potential automorphy II*, Publ. Res. Inst. Math. Sci. **47**, no. 1, 29–98, 2011.
- [4] D. Bump, *Automorphic Forms and Representations*, Cambridge University Press, 1998.
- [5] B. Feigon and D. Whitehouse, *Averages of central L -values of Hilbert modular forms with an application to subconvexity*, Duke Math. J. **149**, 347–410, 2009.
- [6] T. Gee, *The Sato-Tate conjecture for modular forms of weight 3*, Doc. Math. **14**, 771–800, 2009.
- [7] M. Harris, N. Shepherd-Barron and R. Taylor, *A family of Calabi-Yau varieties and potential automorphy*, Ann. of Math.(2) **171**, no.2, 779–813, 2010.
- [8] A. Hora and N. Obata, *Quantum Probability and Spectral Analysis of Graphs*, Theoretical and Mathematical Physics, Springer, Berlin Heidelberg-Verlag, 2007.
- [9] I.G. Macdonald, *Spherical functions on a group of p -adic type*, Uspekhi Mat. Nauk, Vol. **28**, Issue 5(173), 155–224, 1973.
- [10] N. Obata, *Spectral analysis of growing graphs. A quantum probability point of view*, SpringerBriefs in Mathematical Physics, 20. Springer, Singapore, 2017.
- [11] D. Ramakrishnan and J. Rogawski, *Average values of modular L -series via the relative trace formula*, Pure Appl. Math. Quart. **1**, 701–735, 2005.
- [12] H. Saigo, *A new look at the arcsine law and “quantum-classical correspondence”*, Infin. Dimens. Anal. Quantum Probab. Relat. Top. **15**, no. 3, 1250021, 5pp, 2012.
- [13] H. Saigo and H. Sako, *The Arcsine law and an asymptotic behavior of orthogonal polynomials*, Ann. Inst. H. Poincaré D, Volume 3, Issue 4, 405–427, 2016.

- [14] J.P. Serre, *Répartition asymptotique des valeurs propres de l'opérateur de Hecke T_p* , J. Amer. Math. Soc., 10, No.1, 75–102, 1997.
- [15] S. Sugiyama, *Regularized periods of automorphic forms on $GL(2)$* , Tohoku Math. J. (2) **65**, Number 3, 373–409, 2013.
- [16] S. Sugiyama, *Asymptotic behaviors of means of central values of automorphic L -functions for $GL(2)$* , J. Number Theory, **156**, 195–246, 2015.
- [17] S. Sugiyama and M. Tsuzuki, *Relative trace formulas and subconvexity estimates for L -functions of Hilbert modular forms*, Acta Arith. **176**, 1–63 2016.
- [18] S. Sugiyama and M. Tsuzuki, *Existence of Hilbert cusp forms with non-vanishing L -values*, Canad J. Math. **68** No.4, 908–960, 2016.
- [19] S. Sugiyama and M. Tsuzuki, *An explicit trace formula of Jacquet-Zagier type for Hilbert modular forms*, preprint. <https://arxiv.org/pdf/1710.02789.pdf>
- [20] M. Tadić, *Harmonic analysis of spherical functions on reductive groups over p -adic fields*, Pac. J. Math., Vol. **109**, No. 1, 215–235, 1983.
- [21] M. Tsuzuki, *Spectral means of central values of automorphic L -functions for $GL(2)$* , Mem. Amer. Math. Soc., **235**(1110), 2015.
- [22] M. Tsuzuki, *Orbital Integrals on Unitary Hyperbolic Spaces Over p -adic Fields*, Tokyo J. of Math., **39**, Number 3, 923–975, 2017.

Takehiro HASEGAWA

Shiga University, Otsu, Shiga 520-0862, Japan

E-mail : thasegawa3141592@yahoo.co.jp

Hayato SAIGO

Nagahama Institute of Bio-Science and Technology, 1266, Tamura, Nagahama 526-0829, Japan

E-mail : h_saigoh@nagahama-i-bio.ac.jp

Seiken SAITO

Waseda University, Shinjuku, Tokyo 169-8050, Japan

E-mail : seiken.saito@aoni.waseda.jp

Shingo SUGIYAMA

Institute of Mathematics for Industry, Kyushu University, 744, Motoooka, Nishi-ku, Fukuoka 819-0395, Japan

E-mail : s-sugiyama@imi.kyushu-u.ac.jp