

SINGULARITIES OF THE PROJECTIONS
OF n -DIMENSIONAL KNOTS

EIJI OGASA

Department of Mathematical Sciences
University of Tokyo, Komaba
Tokyo 153, Japan

pqr100pqr100@yahoo.co.jp

This paper is published in

Mathematical Proceedings of the Cambridge Philosophical Society, 126, 1999, 511-519.

This manuscript is not the published version.

Abstract. Let $n \geq 5$. There is a smoothly knotted n -dimensional sphere in $(n + 2)$ -space such that the singular point set of its projection in $(n + 1)$ -space consists of double points and that the components of the singular point set are two. (The sphere is *knotted* in the sense that it does not bound any embedded $(n + 1)$ -ball in $(n + 2)$ -space.) Furthermore, the projection is not the projection of any unknotted sphere in $(n + 2)$ -space. There are two inequivalent embeddings of an n -manifold in $(n + 2)$ -space such that the projection of one of these in $(n + 1)$ -space has no double points and the projection of the other has a connected embedded double point set.

This research was partially supported by Research Fellowships of the Promotion of Science for Young Scientists.

1. INTRODUCTION AND MAIN RESULTS

In the study of classical knots, the projections of a knot into \mathbb{R}^2 plays an important role (see e.g. [A], [BL], [C], [CF], [J], [Kf1], [Ko], [Re], [V], [W], for example). For 2-dimensional knots in \mathbb{R}^4 , the projection in \mathbb{R}^3 is considered (see [CS2], [Km], [KSS], [Su], for example.). Projections of codimension two submanifolds of \mathbb{R}^n into \mathbb{R}^{n-1} is studied by [CS1], [CS3], [G], [R], etc. Projections of p -dimensional submanifolds of \mathbb{R}^n into \mathbb{R}^{n-1} , for $p < n - 2$, is studied by [SS], etc.

In this paper, we consider the projection (into \mathbb{R}^{n+1}) of n -dimensional embeddings in \mathbb{R}^{n+2} where $n \geq 5$. We work in the smooth category throughout. We consider those embeddings for which the projection has relatively simple self-intersections. We show that there are embeddings that are truly knotted, but whose projections have simple self intersections. First we introduce some notation.

We work in the smooth category.

An (*oriented*) n -(*dimensional*) *knot* K is a smooth oriented submanifold of $\mathbb{R}^{n+1} \times \mathbb{R}$ which is diffeomorphic to the standard n -sphere. We say that n -knots K_1 and K_2 are *equivalent* if there exists an orientation preserving diffeomorphism $f : \mathbb{R}^{n+1} \times \mathbb{R} \rightarrow \mathbb{R}^{n+1} \times \mathbb{R}$ such that $f(K_1) = K_2$ and $f|_{K_1} : K_1 \rightarrow K_2$ is an orientation preserving diffeomorphism.

Note. In many other papers, including the author's, the definition of n -knot is a smooth oriented submanifold of $\mathbb{R}^{n+1} \times \mathbb{R}$ which is PL homeomorphic to the standard n -sphere. But in this paper, we adopt the former one and reject the latter one.

Let T be the unit n -sphere of $\mathbb{R}^{n+1} \times \{0\} \subset \mathbb{R}^{n+1} \times \mathbb{R}$. Then T is an n -knot. An n -knot K is said to be *unknotted* if K is equivalent to T . If K is not unknotted, then we say that K is *truly knotted*.

Let $\pi : \mathbb{R}^{n+1} \times \mathbb{R} \rightarrow \mathbb{R}^{n+1} \times \{0\}$ be the natural projection map. We suppose $\pi|_K$ is a self-transverse immersion. The *projection* P of an n -knot K is $\pi|_K(K)$ of \mathbb{R}^{n+1} . We give P an orientation by using the orientation of K naturally. The *singular point set* of the projection of an n -knot K is the set $\{x \in \pi|_K(K) \mid \# \{(\pi|_K)^{-1}(x)\} \geq 2\}$. Let $\mu(P)$ denote the number of the connected components of the singular point set of the projection P .

Let K be an n -knot with a projection P . Then the number $\mu(P)$ measures the complexity of K as follows.

Let $n=1$. If $\mu(P) \leq 2$, then K is unknotted. (It is proved by checking all possible projections concretely.)

Let $n=2$. Suppose the singular point set of P consists of double points. If $\mu(P) \leq 2$, then K is unknotted.

Let n be any natural number. There is an n -knot K with a projection P with the following properties. (1) $\mu(K)=3$ (2) K is truly knotted. (3)The singular point set of P consists of double points. Proof. Let K_1 be the trefoil knot. Let K_n be the (0-twist) spun knot of K_{n-1} ($n \geq 2$). (See [Z] for twist spun knots.)

It is natural to consider the following problem.

Problem A. Let K be an n -knot with a Projection P (thus the underlying manifold K

is an n -sphere). Suppose the singular point set of P consists of double points. Suppose that $\mu(P) \leq 2$. Then, is K unknotted?

Of course, if $n = 1$ or 2 , as mentioned above, then the answer is affirmative. But for general $n \geq 5$, we prove the answer to Problem A is negative in §3.

We prove:

Theorem 1. *Let $n \geq 5$. There is an n -knot K with a projection P with the following properties.*

- (1) K is diffeomorphic to the standard n -sphere.
- (2) The singular point set of P consists of double points.
- (3) $\mu(P) = 2$.
- (4) K is truly knotted.

In §4, furthermore, we prove the projection of the n -knot constructed in the proof of Theorem 1 has the following property.

Theorem 2. *Let $n \geq 5$. There is an n -knot with a projection P such that P is not the projection of any knot which is unknotted.*

Note. (1) It is well-known that the projection of any 1-dimensional knot is the projection of a 1-knot which is unknotted. The fact is used in definitions of the Jones polynomial and the Conway-Alexander polynomial. See [Kf1] and [Kf2].

(2) The author proved the $n \geq 3$ case of Theorem 2 is true in [O]. But $\mu(P)$ of the examples are greater than two.

In the case of codimension two submanifolds of \mathbb{R}^{n+2} which are diffeomorphic to a connected closed manifold and which are not spheres, we have the following Problem B corresponding to Problem A.

Let M be a connected closed n -manifold. Let K be a submanifold of \mathbb{R}^{n+2} which is diffeomorphic to M . Suppose $\pi|_K$ is transverse. Put $P = \pi(K)$. The number $\mu(P)$ is defined similarly.

Submanifolds K_1 and K_2 in \mathbb{R}^{n+2} are said to be *equivalent* if there is a diffeomorphism $f : \mathbb{R}^{n+2} \rightarrow \mathbb{R}^{n+2}$ such that $f(K_1) = K_2$ and that $f|_{K_1}$ is an orientation preserving diffeomorphism if M is oriented.

Problem B. Let M, K, P and $\mu(\)$ be as above. Suppose the singular point set of P consists of double points. Suppose that $\mu(P) \leq 2$ (resp. ≤ 1). Then, is an equivalence class of submanifolds determined uniquely? In particular, is it determined uniquely when M is embedded in \mathbb{R}^{n+1} ?

[Sh] shows that, when $M \cong T^2$, then the equivalence class of submanifolds is determined provided $\mu(\) \leq 2$. On the other hand, for high dimensional case we have the following.

In §2 we prove:

Theorem 3. *Let $n \geq 5$. There is a closed connected oriented n -dimensional manifold M as follows. There are submanifolds K_i with a projection P_i ($i = 0, 1$) which are diffeomorphic to M with the following properties.*

- (1) $\mu(P_0)=0$.
- (2) $\mu(P_1)=1$.
- (3) *The singular point set of P_i consists of double points.*
- (4) K_0 is equivalent to neither K_1 , $-K_1$, K_1^* nor $-K_1^*$.
- (5) M is embedded in \mathbb{R}^{n+1} .

The construction of the manifold M in Theorem 3 will be used in the proofs of Theorem 1 and 2.

2. THE PROOF OF THEOREM 3

We first prove the case of $n = 5$.

We define submanifolds K_0 and $K_1 \subset \mathbb{R}^7 = \mathbb{R}^6 \times \mathbb{R}^1$ which are diffeomorphic to $S^3 \times S^2$. Of course $S^3 \times S^2$ is embedded in \mathbb{R}^6 .

We define $K_0 \subset \mathbb{R}^7 = \mathbb{R}^6 \times \mathbb{R}^1$. Let A_0 be a trivially embedded 3-sphere in $\mathbb{R}^6 \times \{0\}$. Take the tubular neighborhood N_0 of A_0 in $\mathbb{R}^6 \times \{0\}$. Then ∂N_0 is diffeomorphic to $S^3 \times S^2$. Define K_0 to be ∂N_0 . The projection P_0 of K_0 coincides with K_0 . Obviously $\mu(P_0)=0$.

We define $K_1 \subset \mathbb{R}^7 = \mathbb{R}^6 \times \mathbb{R}^1$. Take a self-transverse immersion $g : S^3 \looparrowright \mathbb{R}^6 \times \{0\}$ such that the singular point set is one point p . Then $\sharp\{g^{-1}(p)\} = 2$. We suppose that there is a subset V of $\mathbb{R}^6 \times \mathbb{R}^1$ with the following properties.

- (1) $V = \{(x_1, x_2, x_3, y_1, y_2, y_3, z) \mid x_1^2 + x_2^2 + x_3^2 < 1, y_1^2 + y_2^2 + y_3^2 < 1, z \in \mathbb{R}\}$.
- (2) $V \cap g(S^3)$ is a union of two open 3-discs D_x^3 and D_y^3 .
- (3) $D_x^3 = \{(x_1, x_2, x_3, y_1, y_2, y_3, z) \mid x_1^2 + x_2^2 + x_3^2 < 1, y_1 = y_2 = y_3 = 0, z = 0\}$.
- (4) $D_y^3 = \{(x_1, x_2, x_3, y_1, y_2, y_3, z) \mid x_1 = x_2 = x_3 = 0, y_1^2 + y_2^2 + y_3^2 < 1, z = 0\}$.

Take the normal bundle ν of $g(S^3)$ in $\mathbb{R}^6 \times \{0\}$. Let E be a manifold which is the total space of ν . Thus we obtain an immersion $\tilde{g} : E \looparrowright \mathbb{R}^6 \times \{0\}$. Since $\pi_2 SO(3) = 0$, ν is the trivial bundle and ∂E is diffeomorphic to $S^3 \times S^2$.

We can take \tilde{g} to satisfy the following conditions.

- (1) $\tilde{g}|_{V^C}$ is an embedding, where V^C is $\tilde{g}^{-1}(\tilde{g}(E) - \{\tilde{g}(E) \cap V\})$.
- (2) $\tilde{g}(E) \cap V$
 $= \{(x_1, x_2, x_3, y_1, y_2, y_3, z) \mid x_1^2 + x_2^2 + x_3^2 < 1, y_1^2 + y_2^2 + y_3^2 \leq \frac{1}{4}, z = 0\}$
 $\cup \{(x_1, x_2, x_3, y_1, y_2, y_3, z) \mid x_1^2 + x_2^2 + x_3^2 \leq \frac{1}{4}, y_1^2 + y_2^2 + y_3^2 < 1, z = 0\}$.

Figure 1.

You can obtain this figure by clicking ‘PostScript’ in the right side of the cite of the abstract of this paper in arXiv (<https://arxiv.org/abs/the number of this paper>).

You can also obtain it from the author's website, which can be found by typing his name in search engine.

Let $f : E \hookrightarrow \mathbb{R}^6 \times \mathbb{R}^1$ be an embedding with the following properties.

- (1) $\tilde{g}|_{V^c} = f|_{V^c}$
- (2) $f(E) \cap V$
 $= \{(x_1, x_2, x_3, y_1, y_2, y_3, z) \mid x_1^2 + x_2^2 + x_3^2 < 1, y_1^2 + y_2^2 + y_3^2 \leq \frac{1}{4},$
 $z = 1 - (x_1^2 + x_2^2 + x_3^2)\}$
 $\cup \{(x_1, x_2, x_3, y_1, y_2, y_3, z) \mid x_1^2 + x_2^2 + x_3^2 \leq \frac{1}{4}, y_1^2 + y_2^2 + y_3^2 < 1, z = 0\}.$

We can make the corner smooth.

Figure 2.

You can obtain this figure by clicking 'PostScript' in the right side of the cite of the abstract of this paper in arXiv (<https://arxiv.org/abs/the number of this paper>).

You can also obtain it from the author's website, which can be found by typing his name in search engine.

The submanifold $f(\partial E)$ is called K_1 . Then the projection P_1 of K_1 is $\tilde{g}(\partial E)$.

Then we have: The singular point set of the projection P_1 is $\{(x_1, x_2, x_3, y_1, y_2, y_3, z) \mid x_1^2 + x_2^2 + x_3^2 = \frac{1}{4}, y_1^2 + y_2^2 + y_3^2 = \frac{1}{4}, z = 0\}$. It consists of double points. It is diffeomorphic to $S^2 \times S^2$. It is connected. $\mu(P_1) = 1$.

We prove: K_0 is not equivalent to any of $K_1, -K_1, K_1^*$ nor $-K_1^*$.

Proof. Let K be a codimension two submanifold of S^{n+2} . Let X_K denote the infinite cyclic covering space of the complement associated with the natural homomorphism map $\pi_1(S^{n+2} - K) \rightarrow H_1(S^{n+2} - K; \mathbb{Z}) \cong \mathbb{Z}$. We consider $H_*(X_K; \mathbb{Z})$ as a module over $\Lambda = \mathbb{Z}[t, t^{-1}]$. See [M], [L1] etc. for properties of such spaces and those of such modules.

We can regard that K_i is in $S^7 (i = 0, 1)$ naturally. We consider $H_3(X_{K_i}; \mathbb{Z})$. By the construction of K_i , we have:

- (1) $H_3(X_{K_0}; \mathbb{Z}) \cong 0$.
- (2) $H_3(X_{K_1}; \mathbb{Z}) \cong H_3(X_{-K_1}; \mathbb{Z}) \cong H_3(X_{K_1^*}; \mathbb{Z}) \cong H_3(X_{-K_1^*}; \mathbb{Z}) \cong \Lambda/(t-1) \cdot \Lambda$.

Therefore K_0 is equivalent to neither $K_1, -K_1, K_1^*$ nor $-K_1^*$.

We next prove the case of $n > 5$.

We define an n -dimensional submanifold $K_i^{(n)} \subset \mathbb{R}^{n+2}$ as follows. ($n \geq 5, i = 0, 1$.)

Let $K_i^{(5)}$ be K_i .

Put $\mathbb{R}^{n+2} = \{x|x \in \mathbb{R}\} \times \mathbb{R}^n \times \{t|t \in \mathbb{R}\}$. Suppose the projection map is $\mathbb{R}^{n+2} \rightarrow \{x|x \in \mathbb{R}\} \times \mathbb{R}^n \times \{t|t = 0\}$.

We assume $K_i^{(n)} \subset \{x|x \geq 0\} \times \mathbb{R}^n \times \{t|t \in \mathbb{R}\}$ and $K_i^{(n)} \cap \{x|x = 0\} \times \mathbb{R}^n \times \{t|t \in \mathbb{R}\}$ is an n -disc.

We define $K_i^{(n+1)} \subset \mathbb{R}^{(n+3)}$ as follows. We consider $\mathbb{R}^{(n+3)} = \{(x, y)|x, y \in \mathbb{R}\} \times \mathbb{R}^n \times \{t|t \in \mathbb{R}\}$. We regard $\mathbb{R}^{(n+3)}$ as the result of rotating $\{x|x \geq 0\} \times \mathbb{R}^n \times \{t|t \in \mathbb{R}\}$ around $\{x|x = 0\} \times \mathbb{R}^n \times \{t|t \in \mathbb{R}\}$. When rotating it, rotate $K_i^{(n)}$ as well. The result is called $K_i^{(n+1)}$.

By the construction of $K_i^{(n)}$, we have:

(1) For the projection of $P_i^{(n)}$ of $K_i^{(n)}$, $\mu(P_i^{(n)})=1$. The singular point set of P_i consists of double points.

(2) $H_3(X_{K_0^{(n)}}; \mathbb{Z}) \cong 0$.

(3) $H_3(X_{K_1^{(n)}}; \mathbb{Z}) \cong H_3(X_{-K_1^{(n)}}; \mathbb{Z}) \cong H_3(X_{K_1^{(n)*}}; \mathbb{Z}) \cong H_3(X_{-K_1^{(n)*}}; \mathbb{Z}) \cong \Lambda/(t-1) \cdot \Lambda$.

The computation for K_1 follows because the 0-section of E is a generator for $H_3(E)$ (Compare [Kf1], p.43, 190, 229).

Therefore $K_0^{(n)}$ is equivalent to neither $K_1^{(n)}$, $-K_1^{(n)}$, $K_1^{(n)*}$ nor $-K_1^{(n)*}$.

3. THE PROOF OF THEOREM 1

We first prove the case of $n = 5$.

We use $f(E)$ in §2.

We suppose that $f(E) - V \subset \mathbb{R}^6 \times \{0\}$. Take a 6-ball $B^6 \subset \mathbb{R}^6 \times \{0\} \subset \mathbb{R}^6 \times \mathbb{R}$. In $B^6 \times \mathbb{R}$, take a submanifold A_1 which is a parallel displacement of the submanifold $f(E)$. In $(\mathbb{R}^6 - B^6) \times \mathbb{R}$, take a submanifold A_2 which is a parallel displacement of the submanifold $f(E)$ with the opposite orientation.

Recall $E = S^3 \times D^3$. We can put $A_i = S_i^3 \times D^3 = (D_{iS}^3 \cup D_{iN}^3) \times D^3 = (D_{iS}^3 \times D^3) \cup (D_{iN}^3 \times D^3)$ ($i=1,2$). Suppose $(D_{iS}^3 \times D^3)$ is embedded in $\mathbb{R}^6 \times \{0\}$.

Take submanifolds S_1^2 and S_2^2 diffeomorphic to the 2-sphere in ∂B^6 so that the linking number is one.

Figure 3.

You can obtain this figure by clicking ‘PostScript’ in the right side of the cite of the abstract of this paper in arXiv (<https://arxiv.org/abs/the number of this paper>).

You can also obtain it from the author’s website, which can be found by typing his name in search engine.

There are orientation preserving diffeomorphism maps h such that $h(\mathbb{R}^6 \times \{t\}) = \mathbb{R}^6 \times \{t\}$.

By using such a diffeomorphism map, we move A_1 so that:

(1) $(\text{Int}D_{1S}^3) \times D^3 \subset (\mathbb{R}^6 \times \{0\} - B^6)$

(2) $(\text{Int}D_{1N}^3) \times D^3 \subset B^6 \times \mathbb{R}$. The singular point set of the projection of A_1 is in B^6 .

(3) $A_1 \cap \partial B^6 = (\partial D_{1S}^3) \times D^3 = (\partial D_{1N}^3) \times D^3$ and $\partial D_{1s}^3 = \partial D_{1N}^3 = S_1^2$.

By using such a diffeomorphism map, we move A_2 so that:

(1) $(\text{Int}D_{2S}^3) \times D^3 \subset B^6$

(2) $(\text{Int}D_{2N}^3) \times D^3 \subset (\mathbb{R}^6 - B^6) \times \mathbb{R}$. The singular point set of the projection of A_2 is in $\mathbb{R}^6 - B^6$.

(3) $A_2 \cap \partial B^6 = (\partial D_{2S}^3) \times D^3 = (\partial D_{2N}^3) \times D^3$ and $\partial D_{2s}^3 = \partial D_{2N}^3 = S_2^2$.

We define K to be

$$\overline{\partial B^6 - \partial(D_{1N}^3 \times D^3) - \partial(D_{2N}^3 \times D^3) \cup [\partial(D_{1N}^3 \times D^3) \cup \partial(D_{2N}^3 \times D^3)] - \partial B^6}.$$

Figure 4.

You can obtain this figure by clicking ‘PostScript’ in the right side of the cite of the abstract of this paper in arXiv (<https://arxiv.org/abs/the number of this paper>).

You can also obtain it from the author’s website, which can be found by typing his name in search engine.

By the construction, we have:

(1) K is diffeomorphic to the 5-sphere. K is a 5-knot.

(2) For the projection P of K , $\mu(P)=2$. The singular point set of P consists of double points.

(3) A Seifert matrix of K is $\begin{pmatrix} 1 & 1 \\ 0 & -1 \end{pmatrix}$. (See [L1] and [L2] for Seifert matrices.) Hence $H_3(X_K; \mathbb{Z}) \cong \Lambda/(t^2 - 3t + 1) \cdot \Lambda$. Therefore K is truly knotted.

This completes the proof in the case of $n = 5$.

We next prove the case of $n > 5$.

Let $K^{(5)}$ be K . Let $K^{(n+1)}$ be the spun knot of $K^{(n)}$ ($n \geq 5$). (See [Z] for spun knots.)

We take the axis as in the proof of the $n > 5$ case in §2. Then the projection $P^{(n+1)}$ of $K^{(n+1)}$ is the result of rotating $P^{(n)}$ around the axis. Hence $\mu(P)=2$. The singular point set of P consists of double points.

By the construction, we have $H_3(X_{K^{(n+1)}}; \mathbb{Z}) \cong \Lambda/(t^2 - 3t + 1) \cdot \Lambda$. Therefore $K^{(n+1)}$ is truly knotted.

This completes the proof.

4. THE PROOF OF THEOREM 2

We use K , $K^{(n)}$, and $P^{(n)}$ in §3.

We first prove the case of $n = 5$.

Let K' be a 5-knot. Suppose that the projection of K' is the projection P of K .

Then a Seifert matrix of K' is one of the following.

$$\begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}, \begin{pmatrix} 1 & 1 \\ 0 & -1 \end{pmatrix}, \begin{pmatrix} 1 & -1 \\ 0 & 1 \end{pmatrix}, \begin{pmatrix} 1 & -1 \\ 0 & -1 \end{pmatrix}, \begin{pmatrix} -1 & 1 \\ 0 & 1 \end{pmatrix}, \begin{pmatrix} -1 & 1 \\ 0 & -1 \end{pmatrix}, \begin{pmatrix} -1 & -1 \\ 0 & 1 \end{pmatrix}, \begin{pmatrix} -1 & -1 \\ 0 & -1 \end{pmatrix}.$$

Hence $H_3(X_{K'}; \mathbb{Z})$ is not trivial. Therefore K' is truly knotted.

We next prove the case of $n > 5$.

Let $K^{(n)'}$ be an n -knot ($n > 5$). Suppose the projection of $K^{(n)'}$ is the projection $P^{(n)}$ of $K^{(n)}$. Then $K^{(n)'}$ is a spun knot of an $(n - 1)$ -knot whose projection is $P^{(n-1)}$. Hence $H_3(X_{K^{(n)'}}; \mathbb{Z})$ is not trivial. Therefore $K^{(n)'}$ is truly knotted.

REFERENCES

- [A] J. W. Alexander, *Topological invariant of knots and links*, TransAMS **30** (1923), 275-360.
- [BL] J. S. Birman and X. S. Lin, *Knot polynomials and Vassiliev's invariants*, Invent. Math. **111** (1993), 225-270.
- [CS1] J. S. Carter and M. Saito, *Canceling branch points on projections of surfaces in 4-space*, PAMS **116** (1992), 229-237.
- [CS2] J. S. Carter and M. Saito, *Knotted surfaces, braid movies, and beyond*, Knots and quantum gravity, edited by J. C. Baez, Clarendon Press, Oxford (1994), 191-229.
- [CS3] J. S. Carter and M. Saito, *Surfaces in 3-space that do no lift to embeddings in 4-space*, preprint.
- [C] J. Conway, *An enumeration of knots and links, and their algebraic properties*, Computational problems in Abstract Algebra, Pergamon Press, New York (1970).
- [CF] R. H. Crowell and R. H. Fox, *Introduction to knot theory*, Boston:Gin (1963).
- [G] C. Giller, *Towards a classical knot theory for surfaces in \mathbb{R}^4* , Illinois.J **26** (1982), 591-631.
- [J] V. F. R. Jones, *Hecke Algebra representations of braid groups and link*, Ann.Math. **126** (1987), 335-388.
- [Kf1] L. Kauffman, *On knots*, Ann of math studies **115** (1987).
- [Kf2] L. Kauffman, *Knots and Physics*, World Scientific, Series on Knots and Everything **1** (1991).
- [Km] S. Kamada, *A characterization of groups of closed orientable surfaces in 4-space*, Topology **33** (1994), 113-122.
- [KSS] A. Kawauchi, T. Shibuya and S. Suzuki, *Descriptions of surfaces in four-space I; Normal forms*, Math.Sem.Notes.Kobe.Univ. **10** (1982), 75-125.
- [Ko] M. Kontsevich, *Vassiliev's knot invariants*, Adv. in Soviet Math. **16** (1993), 137-150.
- [L1] J. Levine, *Polynomial invariants of knots of codimension two*, Ann. of Math. **84** (1966), 537-554.
- [L2] J. Levine, *Knot cobordism in codimension two*, Comment. Math. Helv. **44** (1969), 229-244.
- [M] J. W. Milnor, *Infinite cyclic coverings*, Topology of Manifolds (Michigan State University 1967)Prindle, Weber and Schmidt, Boston (1968).
- [O] E. Ogasa, *The projections of n -knots which are not the projection of any knot which is unknot* (preprint).
- [Re] K. Reidemeister, *Knotentheorie*, Ergebn.Math.Grenzgeb. Springer Verlag (1932).
- [Ro] D. Roseman, *Projection of codimension two embeddings*, Preprint.
- [SS] O.Saeki and K.Sakuma, *Immersed n -manifolds in \mathbb{R}^{2n} and the double points of their generic projections into \mathbb{R}^{2n-1}* , Trans.Am.Math.Soc. **348** (1996), 2585-2606.
- [Sh] A. Shima, *An unknotting theorem for tori in S^4 (J.Math.Sci.Univ.Tokyo,4,1997,279-339), II (Kobe J.Math.,13,1996,9-25)*.
- [Su] S. Suzuki, *Knotted Problems of 2-spheres in the 4-sphere*, Math.Sem.Notes.Kobe.Univ. **4** (1976), 241-371.
- [V] V. A. Vassiliev, *Complements of Discriminants of smooth maps: Topology and Applications*, Translations of Mathematical Monographs, American Mathematical Society **98** (1994).
- [W] E. Witten, *Quantum field theory and the Jones polynomial*, Commun. Math. Phys. **121** (1989), 351-399.
- [Z] E. Zeeman, *Twisting spun knots*, Trans. Am. Math. Soc. **115** (1965), 471-495.