

# BOUNDS FOR A SPECTRAL EXPONENTIAL SUM

OLGA BALKANOVA AND DMITRY FROLENKOV

ABSTRACT. We prove new upper bounds for a spectral exponential sum by refining the process by which one evaluates mean values of  $L$ -functions multiplied by an oscillating function. In particular, we introduce a method which is capable of taking into consideration the oscillatory behaviour of the function. This gives an improvement of the result of Luo and Sarnak when  $T \geq X^{1/6+2\theta/3}$ . Furthermore, this proves the conjecture of Petridis and Risager in certain ranges.

## CONTENTS

1. Introduction	1
2. Notation	4
3. Initial steps of the proof	9
4. Exact formula	11
5. Analysis of $\Sigma(s)$	12
6. Analysis of $\Sigma_B(s)$	15
7. Special functions	17
8. Estimates for the moment of Maaß sym <sup>2</sup> $L$ -functions	24
9. Estimates for the spectral exponential sum	26
Acknowledgements	30
References	30

## 1. INTRODUCTION

This paper presents a new approach to studying the spectral exponential sum

$$(1.1) \quad S(T, X) = \sum_{t_j \leq T} X^{it_j},$$

---

2010 *Mathematics Subject Classification.* Primary: 11F12.

*Key words and phrases.* spectral exponential sum; symmetric square  $L$ -functions; generalized Dirichlet  $L$ -functions; Prime Geodesic Theorem; Kuznetsov trace formula.

where  $\kappa_j = 1/4 + t_j^2$  are the eigenvalues of the hyperbolic Laplacian for  $PSL_2(\mathbf{Z})$ . The sum (1.1) is attracting considerable interest because it is closely related to some classical problems, including prime geodesic theorem and hyperbolic lattice counting problem. See, for example, [7], [12], [17], [19], [20].

The prime geodesic theorem is concerned with the asymptotic behaviour of

$$\pi_\Gamma(X) = \#\{\{P\} : NP \leq X\},$$

where  $\{P\}$  denotes a primitive hyperbolic class in  $PSL_2(\mathbf{Z})$  and  $NP$  denotes its norm.

By partial summation, the problem can be formulated in terms of

$$\Psi_\Gamma(X) = \sum_{NP \leq X} \Lambda(P),$$

where the sum is over all hyperbolic classes and  $\Lambda(P) = \log NP_0$  if  $\{P\}$  is a power of the primitive hyperbolic class  $\{P_0\}$ .

Iwaniec [12, Lemma 1] proved for  $1 \leq T \leq X^{1/2} \log^{-2} X$  the following explicit formula

$$(1.2) \quad \Psi_\Gamma(X) = X + X^{1/2} \sum_{|t_j| \leq T} \frac{X^{it_j}}{1/2 + it_j} + O\left(\frac{X}{T} \log^2 X\right),$$

which is closely related to the spectral exponential sum (1.1).

The trivial bound on (1.1), namely

$$S(T, X) \ll T^2,$$

follows from Weyl's law and yields

$$(1.3) \quad \Psi_\Gamma(X) = X + O(X^{3/4} \log X).$$

In order to improve (1.3) it is required to exploit cancellation in sum (1.1). There are several non-trivial estimates in the literature. The first estimate

$$(1.4) \quad S(T, X) \ll TX^{11/48+\epsilon}$$

was obtained by Iwaniec in [12]. As a consequence, he showed that

$$\Psi_\Gamma(X) = X + O(X^{35/48+\epsilon}).$$

Luo and Sarnak [17] proved another estimate

$$(1.5) \quad S(T, X) \ll T^{5/4} X^{1/8} \log^2 T$$

and deduced from this that

$$\Psi_\Gamma(X) = X + O(X^{7/10+\epsilon}).$$

In [6, Eq. (7.1)] Cai showed that

$$(1.6) \quad S(T, X) \ll (TX)^\epsilon (T^{2/5} X^{11/30} + T^{3/2}),$$

and consequently,

$$\Psi_\Gamma(X) = X + O(X^{71/102+\epsilon}).$$

Finally, the prime geodesic theorem in the strongest presently known form

$$\Psi_\Gamma(X) = X + O(X^{2/3+\theta/6}) \text{ with } \theta = 1/6 + \epsilon,$$

was proved by Soundararajan and Young in [21] by combining Luo-Sarnak's bound (1.5) and a new approach without recourse to the spectral exponential sum.

Petridis and Risager [20, Conj. 2.2] conjectured square root cancellation in sum (1.1), namely

$$(1.7) \quad S(T, X) \ll T(TX)^\epsilon.$$

Furthermore, they showed that estimate (1.7) yields not only the best possible error term  $O(X^{1/2+\epsilon})$  in the prime geodesic theorem, but also the best error term on average for the hyperbolic lattice problem. See [20] for more details.

In the appendix of [20], Laaksonen proved that the conjecture of Petridis and Risager is true for a fixed  $X$  as  $T \rightarrow \infty$ . Moreover, for

$$(1.8) \quad \kappa(X) := \|X^{1/2} + X^{-1/2}\|,$$

where  $\|x\|$  is the distance from  $x$  to the nearest integer, Laaksonen mentioned in [20, Experimental Observation 2] that  $S(T, X)$  has a peak when  $\kappa(X) = 0$ .

The quantity  $\kappa(X)$  appears also in our result. Furthermore, our estimates depend on the parameter  $\theta$ , which denotes the subconvexity exponent in the conductor aspect for Dirichlet  $L$ -functions of real primitive characters. The current record is  $\theta = 1/6 + \epsilon$  due to Conrey and Iwaniec [8].

**Theorem 1.1.** *The following estimates hold:*

$$(1.9) \quad S(T, X) \ll \max(X^{1/4+\theta/2} T^{1/2}, X^{\theta/2} T) \log^3 T;$$

$$(1.10) \quad S(T, X) \ll T \log^2 T \quad \text{if } T > \frac{X^{1/2+7\theta/6}}{\kappa(X)}.$$

Theorem 1.1 shows that the conjecture of Petridis and Risager is true for  $T > X^{1/2}$  under the Lindelöf conjecture, and for  $T > X^{1/2+7\theta/6}/\kappa(X)$  unconditionally.

Furthermore, estimate (1.9) improves (1.5) when  $T \geq X^{1/6+2\theta/3+\epsilon}$ . Consequently, this result has an application to the prime geodesic theorem, as we now show. Using summation by parts, (1.5) and (1.9) we obtain

$$(1.11) \quad \Psi_{\Gamma}(X) = X + O\left(X^{2/3+\theta/6} \log^3 X\right).$$

The error term in asymptotic formula (1.11) coincides with the one proved by Soundararajan and Young. Note that using (1.6) instead of (1.5) would not lead to further improvement.

The proof of Theorem 1.1 is based on the upper bound for the mean value of Maaß symmetric square  $L$ -functions on the critical line multiplied by  $X^{it_j}$ .

**Theorem 1.2.** *Let  $X \gg 1$  then for  $s = 1/2 + ir$ ,  $|r| \ll T^\epsilon$  one has*

$$(1.12) \quad \sum_j \alpha_j X^{it_j} \exp(-t_j/T) L(\text{sym}^2 u_j, s) \ll (1 + |r|)^A T \log^3 T + \frac{(1 + |r|)^A X^\theta}{X^{1/4}} \left( X^{1/2} \min\left(T, \frac{X^{1/2}}{\kappa(X)}\right)^{1/2} + \min\left(T, \frac{X^{1/2}}{\kappa(X)}\right)^{3/2} \right).$$

The paper is organised as follows. In Section 2 we introduce the required notation. In Section 3 initial steps of the proof are described following the paper of Iwaniec [12]. Sections 4, 5, 6 are devoted to proving an exact formula for the first moment of Maaß symmetric square  $L$ -functions multiplied by an oscillating function. In Section 7 we analyse special functions that appear in the exact formula. Consequently, we prove Theorem 1.2 in Section 8. The main result, namely Theorem 1.1, is proved in Section 9.

## 2. NOTATION

For a complex number  $v$  let

$$\tau_v(n) = \sum_{n_1 n_2 = n} \left(\frac{n_1}{n_2}\right)^v = n^{-v} \sigma_{2v}(n),$$

where

$$\sigma_v(n) = \sum_{d|n} d^v.$$

Note that

$$(2.1) \quad \sum_{n=1}^{\infty} \frac{\tau_{ir}(n^2)}{n^s} = \frac{\zeta(s)\zeta(s+2ir)\zeta(s-2ir)}{\zeta(2s)},$$

where  $\zeta(s)$  is the Riemann zeta function.

Define the Mellin transform of  $f(x)$  as follows

$$\tilde{f}(s) = \int_0^\infty f(x)x^{s-1}dx.$$

The Kloosterman sum is given by

$$S(n, m; c) = \sum_{\substack{a \pmod{c} \\ (a, c)=1}} e\left(\frac{an + a^*m}{c}\right), \quad aa^* \equiv 1 \pmod{c},$$

where  $e(x) = \exp(2\pi ix)$ . According to the Weil bound (see [22])

$$(2.2) \quad |S(m, n; c)| \leq \tau_0(c)\sqrt{(m, n, c)}\sqrt{c}.$$

The Lerch zeta function

$$(2.3) \quad \zeta(\alpha, \beta, s) = \sum_{n+\alpha>0} \frac{e(n\beta)}{(n+\alpha)^s}$$

satisfies the functional equation (see [15])

$$(2.4) \quad \zeta(\alpha, 0, s) = \frac{\Gamma(1-s)}{(2\pi)^{1-s}} \left( ie\left(-\frac{s}{4}\right) \zeta(0, -\alpha, 1-s) - ie\left(\frac{s}{4}\right) \zeta(0, \alpha, 1-s) \right).$$

Introduce the generalized Dirichlet  $L$ -function (see [3, 24] for details)

$$(2.5) \quad \mathcal{L}_n(s) = \frac{\zeta(2s)}{\zeta(s)} \sum_{q=1}^\infty \frac{\rho_q(n)}{q^s} = \sum_{q=1}^\infty \frac{\lambda_q(n)}{q^s},$$

where

$$(2.6) \quad \rho_q(n) := \#\{x \pmod{2q} : x^2 \equiv n \pmod{4q}\},$$

$$(2.7) \quad \lambda_q(n) := \sum_{q_1^2 q_2 q_3 = q} \mu(q_2) \rho_{q_3}(n).$$

For  $n \neq 0$  one has

$$(2.8) \quad \mathcal{L}_n(1/2 + it) \ll n^\theta (1 + |t|)^A,$$

where  $\theta$  and  $A$  are subconvexity exponents for Dirichlet  $L$ -functions of real primitive characters. The best known result in the  $q$  aspect, namely  $\theta = 1/6 + \epsilon$ , was first obtained by Conrey and Iwaniec [8]. Young [23] proved the hybrid subconvexity bound with  $A = \theta = 1/6 + \epsilon$ .

Denote by  $\{u_j\}$  the orthonormal basis of the space of Maaß cusp forms consisting of common eigenfunctions of all Hecke operators and the hyperbolic Laplacian. Let  $\{\lambda_j(n)\}$  be the eigenvalues of Hecke operators acting on  $u_j$ . Let  $\kappa_j = 1/4 + t_j^2$  be the eigenvalues of the

hyperbolic Laplacian acting on  $u_j$ . Elements of the basis have a Fourier expansion of the following form

$$u_j(x + iy) = \sqrt{y} \sum_{n \neq 0} \rho_j(n) K_{it_j}(2\pi|n|y) e(nx),$$

where  $K_\alpha(x)$  is the  $K$ -Bessel function and

$$\rho_j(n) = \rho_j(1) \lambda_j(n).$$

Note that for  $n, m \geq 1$

$$(2.9) \quad \lambda_j(n) \lambda_j(m) = \sum_{d|(m,n)} \lambda_j\left(\frac{nm}{d^2}\right).$$

Introduce the normalizing coefficient

$$(2.10) \quad \alpha_j := \frac{|\rho_j(1)|^2}{\cosh \pi t_j}.$$

For  $\Re s > 1$ , the symmetric square  $L$ -function is given by

$$(2.11) \quad L(\text{sym}^2 u_j, s) := \zeta(2s) \sum_{n=1}^{\infty} \frac{\lambda_j(n^2)}{n^s}.$$

For  $\Re s > 1$  define the Rankin zeta-function

$$(2.12) \quad L(u_j \otimes u_j, s) := \sum_{n=1}^{\infty} \frac{|\rho_j(n)|^2}{n^s}.$$

We will use the following properties (see [16, p. 216] and [12, proof of Lemma 8])

$$(2.13) \quad \text{res}_{s=1} L(u_j \otimes u_j, s) = \frac{2}{\zeta(2)} \cosh(\pi t_j),$$

$$(2.14) \quad L(u_j \otimes u_j, s) = |\rho_j(1)|^2 \frac{\zeta(s)}{\zeta(2s)} L(\text{sym}^2 u_j, s).$$

Let  $\varphi(x)$  be a smooth function on  $[0, \infty)$  such that

$$\varphi(0) = 0, \quad \varphi^{(j)}(x) \ll (1+x)^{-2-\epsilon}, \quad j = 0, 1, 2.$$

Let  $J_\nu(x)$  be the  $J$ -Bessel function. Define the series of integral transforms

$$(2.15) \quad \varphi_0 = \frac{1}{2\pi} \int_0^\infty J_0(y) \varphi(y) dy,$$

$$(2.16) \quad \varphi_B(x) = \int_0^1 \int_0^\infty \xi x J_0(\xi x) J_0(\xi y) \varphi(y) dy d\xi,$$

$$(2.17) \quad \varphi_H(x) = \int_1^\infty \int_0^\infty \xi x J_0(\xi x) J_0(\xi y) \varphi(y) dy d\xi,$$

$$(2.18) \quad \hat{\varphi}(t) = \frac{\pi i}{2 \sinh(\pi t)} \int_0^\infty (J_{2it}(x) - J_{-2it}(x)) \varphi(x) \frac{dx}{x},$$

$$(2.19) \quad \check{\varphi}(l) = \int_0^\infty J_l(y) \varphi(y) \frac{dy}{y}.$$

The following decomposition holds

$$(2.20) \quad \varphi(x) = \varphi_H(x) + \varphi_B(x),$$

where

$$(2.21) \quad \varphi_B(x) = \sum_{l \equiv 1(2)} 2l \check{\varphi}(l) J_l(x).$$

**Lemma 2.1.** (*Kuznetsov trace formula*) For all  $m, n \geq 1$

$$(2.22) \quad \sum_{j=1}^{\infty} \alpha_j \lambda_j(m) \lambda_j(n) \hat{\varphi}(t_j) + \frac{2}{\pi} \int_0^\infty \frac{\tau_{ir}(m) \tau_{ir}(n)}{|\zeta(1+2ir)|^2} \hat{\varphi}(r) dr = \\ \delta(m, n) \varphi_0 + \sum_{q=1}^{\infty} \frac{S(m, n; q)}{q} \varphi_H \left( \frac{4\pi \sqrt{mn}}{q} \right).$$

*Proof.* See [13] or [14]. □

Following [9] and [17]<sup>1</sup> let

$$(2.23) \quad \varphi(x) = \frac{\sinh \beta}{\pi} x \exp(ix \cosh \beta)$$

with

$$(2.24) \quad \beta = \frac{1}{2} \log X + \frac{i}{2T}.$$

It is useful to introduce the following notation

$$(2.25) \quad c := -i \cosh \beta = a - ib,$$

$$(2.26) \quad \begin{cases} a := \sinh(\log \sqrt{X}) \sin((2T)^{-1}); \\ b := \cosh(\log \sqrt{X}) \cos((2T)^{-1}). \end{cases}$$

---

<sup>1</sup>Note that there is a typo (the imaginary unit  $i$  is placed in the denominator instead of the numerator) in [9, p.68 line -1] and [17, p.233 line -3] in the definition of  $\hat{\varphi}(t)$ . See (2.18) for the corrected version. This explains the sign change in the definition of  $\varphi(x)$ .

Note that

$$(2.27) \quad \arg c = -\pi/2 + \gamma, \quad 0 < \gamma \ll T^{-1}.$$

**Lemma 2.2.** *The following holds*

$$(2.28) \quad \hat{\varphi}(t) = \frac{\sinh(\pi t + 2i\beta t)}{\sinh(\pi t)} = X^{it} \exp(-t/T) + O(\exp(-\pi t)),$$

$$(2.29) \quad \varphi_0 = \frac{-\cosh \beta}{2\pi^2 \sinh^2 \beta},$$

$$(2.30) \quad \varphi_B(x) \ll X^{-1/2} \min(x, x^{1/2}),$$

$$(2.31) \quad \varphi_B(x) = \frac{2}{\pi} \sum_{k=1}^{\infty} (-1)^k (2k-1) \exp(-(2k-1)\beta) J_{2k-1}(x).$$

*Proof.* It follows from [11, Eq. 6.6621.1], [18, Eq. 15.4.18] that

$$(2.32) \quad \int_0^{\infty} J_{2it}(x) \exp(ix \cosh \beta) dx = -\frac{\exp(-(\pi + 2i\beta)t)}{i \sinh \beta}.$$

Using (2.18), (2.23) and (2.32) we prove (2.28). Differentiating equation (2.32) with respect to  $\beta$  and taking  $t = 0$  we obtain

$$(2.33) \quad \int_0^{\infty} J_0(x) \exp(ix \cosh \beta) x dx = -\frac{\cosh \beta}{\sinh \beta}.$$

Now (2.29) follows from (2.15), (2.23) and (2.33). Estimate (2.30) can be proved in the same way as [9, Lemma 11]. Namely,

$$\varphi_B(x) \ll |\sinh(2\beta)| \int_0^1 \frac{\min(\xi x, \sqrt{\xi x})}{|\cosh^2 \beta - \xi^2|^{3/2}} d\xi \ll X^{-1/2} \min(x, x^{1/2}).$$

To prove (2.31) it is required to substitute (2.23) to (2.21) and use [11, Eq. 6.6621.1], [18, Eq. 15.4.18].  $\square$

---

<sup>2</sup>Note that there is a typo in [9, Eq. 7.7]: the minus sign is missed. For this reason, the formula (2.33) has an additional minus sign compared to [9, Eq. 7.8]. Consequently, we obtain the same expression (2.29) for  $\varphi_0$  as in [9, Eq. 7.8] regardless of the sign change in the definition of  $\varphi(x)$ .



## 3. INITIAL STEPS OF THE PROOF

To prove Theorem 1.1 we follow the approach of Iwaniec [12]. See also [17, Section 6] for more details. First of all, the problem can be reduced to the analysis of the sum

$$(3.1) \quad \sum_j X^{it_j} \exp(-t_j/T)$$

at the cost of the error  $O(T \log T)$ . To this end, we introduce a smooth function  $g(x)$  such that  $g(x) = 1$  for  $1 \leq x \leq T$  and  $g(x) = 0$  for  $x \leq 1/2$  and  $x \geq T + 1/2$ . Then

$$\sum_{t_j \leq T} X^{it_j} = \sum_j X^{it_j} \exp(-t_j/T) g(t_j) \exp(t_j/T) + O(T).$$

Let

$$\hat{g}(x) = \int_{-\infty}^{\infty} g(\xi) \exp(\xi/T) e(x\xi) d\xi.$$

According to [17, Section 6]

$$(3.2) \quad \hat{g}(x) \ll \min(T, 1/|x|).$$

Finally we have the following lemma (see [17, Section 6]).

**Lemma 3.1.** *One has*

$$(3.3) \quad \sum_{t_j \leq T} X^{it_j} = \int_{-1}^1 \hat{g}(\xi) \left( \sum_j (X \exp(-2\pi\xi))^{it_j} \exp(-t_j/T) \right) d\xi + O(T \log T).$$

Let  $h(x)$  be a smooth function supported in  $[N, 2N]$  such that

$$(3.4) \quad |h^{(j)}(x)| \ll N^{-j}, \text{ for } j = 0, 1, 2, \dots \quad \int_{-\infty}^{\infty} h(x) dx = N.$$

The first idea of Iwaniec was to investigate the following expression

$$(3.5) \quad \sum_n h(n) \sum_j \frac{|\rho_j(n)|^2}{\cosh(\pi t_j)} X^{it_j} \exp(-t_j/T).$$

The sum over  $n$  can be evaluated using the Mellin inversion formula for  $h(n)$  and the fact that Rankin zeta function (2.12) has a pole at the point  $s = 1$  with the residue given by equation (2.13). As in [12, Lemma 8] one has

$$(3.6) \quad \sum_n h(n) \frac{|\rho_j(n)|^2}{\cosh(\pi t_j)} = \frac{2}{\zeta(2)} N + \frac{1}{2\pi i} \int_{(1/2)} \tilde{h}(s) \frac{L(u_j \otimes u_j, s)}{\cosh(\pi t_j)} ds.$$

Substituting (3.6) to (3.5), we obtain

$$(3.7) \quad \sum_j X^{it_j} \exp(-t_j/T) = \frac{\zeta(2)}{2N} \sum_n h(n) \sum_j \frac{|\rho_j(n)|^2}{\cosh(\pi t_j)} X^{it_j} \exp(-t_j/T) \\ - \frac{\zeta(2)}{2N} \frac{1}{2\pi i} \int_{(1/2)} \tilde{h}(s) \sum_j X^{it_j} \exp(-t_j/T) \frac{L(u_j \otimes u_j, s)}{\cosh(\pi t_j)} ds.$$

The standard tool for studying the first sum on the right-hand side of equation (3.7) is the Kuznetsov trace formula.

The second idea of Iwaniec was not to apply Kuznetsov trace formula directly but to find a function  $\varphi(x)$  such that its transform (2.18) approximates  $X^{it} \exp(-t/T)$  with a small error term. The procedure of finding a suitable  $\varphi(x)$  is not straightforward and several examples of such functions are available in the literature, i.e. [12, Eq. 29], [9, Eq. 7.2]. The approximation by [9, Eq. 7.2] produces a smaller error term, and therefore, similarly to [17] we choose  $\varphi(x)$  defined by (2.23). One can compute  $\hat{\varphi}(t)$  explicitly, see equation (2.28). Replacing  $X^{it} \exp(-t/T)$  by  $\hat{\varphi}(t)$  in (3.7), we have

$$(3.8) \quad \sum_j X^{it_j} \exp(-t_j/T) + O(1) = \frac{\zeta(2)}{2N} \sum_n h(n) \sum_j \frac{|\rho_j(n)|^2}{\cosh(\pi t_j)} \hat{\varphi}(t_j) \\ - \frac{\zeta(2)}{2N} \frac{1}{2\pi i} \int_{(1/2)} \tilde{h}(s) \sum_j \hat{\varphi}(t_j) \frac{L(u_j \otimes u_j, s)}{\cosh(\pi t_j)} ds.$$

Applying Kuznetsov's trace formula (2.22) to the first sum over  $j$  on the right-hand side of equation (3.8) and arguing as in the paper of Luo-Sarnak [17, p. 234], we obtain

$$(3.9) \quad \sum_j X^{it_j} \exp(-t_j/T) \ll \frac{NX^{1/2} \log T}{(\max(1, NX^{1/2}/T))^{1/2}} + \sqrt{N/X} \log^2 N \\ + T \log^2 T + \frac{1}{N} \left| \int_{(1/2)} \tilde{h}(s) \sum_j \hat{\varphi}(t_j) \frac{L(u_j \otimes u_j, s)}{\cosh(\pi t_j)} ds \right|.$$

**Remark 3.2.** Estimate (3.9) coincides with the bound of Luo and Sarnak if  $\max(1, NX^{1/2}/T) = NX^{1/2}/T$ . See [17, p. 235, lines 2-4]. This maximum arises naturally when we estimate the sum of Kloosterman sums

$$S_n(\phi) := \sum_{q=1}^{\infty} \frac{S(n, n; q)}{q} \varphi\left(\frac{4\pi n}{q}\right) \ll \sum_{q=1}^{\infty} \frac{|S(n, n; q)|}{q} X^{1/2} \frac{n}{q} \exp(-4\pi an/q)$$

using the Weil bound. Since  $a \asymp X^{1/2}/(4T)$  (see (2.26)), the sum over  $q$  can be viewed as a sum over  $q > \max(1, NX^{1/2}/T)$ .

To analyse the integral in (3.9), Luo and Sarnak used the following "mean Lindelöf" estimate [17, Eq. 5]

$$(3.10) \quad \sum_{t_j \leq T} \frac{L(u_j \otimes u_j, 1/2 + ir)}{\cosh(\pi t_j)} \ll (1 + |r|)^4 T^2 \log^2 T.$$

In order to improve their result we consider the whole sum

$$\sum_j \hat{\varphi}(t_j) \frac{L(u_j \otimes u_j, 1/2 + ir)}{\cosh(\pi t_j)},$$

trying to use the oscillations of the function  $\hat{\varphi}(t_j)$ .

It follows from the properties of  $h(x)$  that  $|\tilde{h}(s)| \ll N^{1/2}(1 + |r|)^{-A}$  for any  $A$  and  $s = 1/2 + ir$ . Thus applying equation (3.10) to handle  $|r| > T^\epsilon := r_0$ , we need to work only with  $|r| \leq r_0$ . Finally, using equations (2.14) and (2.10), we obtain

$$(3.11) \quad \sum_j X^{it_j} \exp(-t_j/T) \ll \frac{NX^{1/2} \log T}{(\max(1, NX^{1/2}/T))^{1/2}} + \sqrt{N/X} \log^2 N + T \log^2 T + \frac{1}{N^{1/2}} \int_{-r_0}^{r_0} \left| \frac{\zeta(1/2 + ir)}{\zeta(1 + 2ir)} \sum_j \alpha_j \hat{\varphi}(t_j) L(\text{sym}^2 u_j, 1/2 + ir) \right| dr.$$

#### 4. EXACT FORMULA

This section is devoted to the analysis of the first moment

$$(4.1) \quad M_1(s) := \sum_j \alpha_j \hat{\varphi}(t_j) L(\text{sym}^2 u_j, s)$$

with  $s = 1/2 + ir$ ,  $|r| \leq T^\epsilon$ . The key ideas are similar to the ones used in [1] for the holomorphic case. The main difference is that instead of the Petersson trace formula we now work with the Kuznetsov trace formula (2.22).

In order to prove an exact formula for  $M_1(1/2 + ir)$ , we apply the technique of analytic continuation. For  $\Re s > 3/2$  define

$$(4.2) \quad \Sigma(s) = \zeta(2s) \sum_{n=1}^{\infty} \frac{1}{n^s} \sum_{q=1}^{\infty} \frac{S(1, n^2; q)}{q} \varphi\left(\frac{4\pi n}{q}\right),$$

$$(4.3) \quad \Sigma_B(s) = \zeta(2s) \sum_{n=1}^{\infty} \frac{1}{n^s} \sum_{q=1}^{\infty} \frac{S(1, n^2; q)}{q} \varphi_B \left( \frac{4\pi n}{q} \right).$$

Convergence of the double series in (4.2) and (4.3) for  $\Re s > 3/2$  follows from (2.2), (2.23) and (2.30).

**Lemma 4.1.** *For  $\Re s > 3/2$  one has*

$$(4.4) \quad M_1(s) = \zeta(2s) \varphi_0 + \Sigma(s) - \Sigma_B(s) - \frac{2\zeta(s)}{\pi} \int_0^{\infty} \frac{\zeta(s+2it)\zeta(s-2it)}{|\zeta(1+2it)|^2} \hat{\varphi}(t) dt.$$

*Proof.* Applying Kuznetsov trace formula (2.22) and using identity (2.1), we obtain

$$\begin{aligned} M_1(s) &= \zeta(2s) \sum_{n=1}^{\infty} \frac{1}{n^s} \sum_j \alpha_j \hat{\varphi}(t_j) \lambda_j(1) \lambda_j(n^2) = \\ &= \zeta(2s) \varphi_0 - \frac{2\zeta(s)}{\pi} \int_0^{\infty} \frac{\zeta(s+2it)\zeta(s-2it)}{|\zeta(1+2it)|^2} \hat{\varphi}(t) dt + \\ & \quad \zeta(2s) \sum_{n=1}^{\infty} \frac{1}{n^s} \sum_{q=1}^{\infty} \frac{S(1, n^2; q)}{q} \varphi_H \left( \frac{4\pi n}{q} \right). \end{aligned}$$

Then equation (2.20) yields the lemma.  $\square$

In order to extend exact formula (4.4) to the critical line  $\Re s = 1/2$ , it is required to continue analytically double sums (4.2) and (4.3). This is the subject of the two subsequent sections.

## 5. ANALYSIS OF $\Sigma(s)$

The strategy of working with  $\Sigma(s)$  is the same as in the proof of [1, Lemma 5.1]. First, we change the order of summation using the fact that  $\Re s > 3/2$ . Second, we apply the Mellin inverse formula for  $\varphi(x)$ , open the Kloosterman sum and obtain the Lerch zeta function. After that we move the line of integration to the region, where the functional equation for the Lerch zeta function can be applied.

However, it turns out that the Mellin transform of  $\varphi(x)$  does not allow moving the line of integration to the desired region. The reason is that the function  $\varphi(x)$  behaves asymptotically like  $x$  when  $x \rightarrow 0$ , and this is insufficient for absolute convergence. To overcome this difficulty we use the so-called "Hecke trick". Accordingly, instead of working directly with  $\varphi(x)$  and  $\Sigma(s)$ , we introduce for a complex variable  $\lambda$  with  $\Re \lambda > 1$  two functions

$$(5.1) \quad \varphi(\lambda, x) = \frac{\sinh \beta}{\pi} x^\lambda \exp(ix \cosh \beta),$$

$$(5.2) \quad \Sigma(\lambda, s) = \sum_{n=1}^{\infty} \frac{1}{n^s} \sum_{q=1}^{\infty} \frac{S(1, n^2; q)}{q} \varphi \left( \lambda, \frac{4\pi n}{q} \right),$$

where  $\beta$  is defined by (2.24). We first prove a formula for  $\Sigma(\lambda, s)$  under the assumption that  $\Re\lambda > \Re s$  and then extend it by analytic continuation to the point  $\lambda = 1$ .

**Lemma 5.1.** *For  $\Re\lambda > \Re s > 3/2$  one has*

$$(5.3) \quad \Sigma(\lambda, s) = (4\pi)^{s-1} \tilde{\varphi}(\lambda, 1-s) \mathcal{L}_{-4}(s) + 2(2\pi)^{s-1} \sum_{n=1}^{\infty} \frac{1}{n^{1-s}} \mathcal{L}_{n^2-4}(s) I \left( \lambda, \frac{n}{2} \right),$$

where

$$(5.4) \quad I(\lambda, x) := \frac{1}{2\pi i} \int_{(\Delta)} \tilde{\varphi}(\lambda, w) \Gamma(1-s-w) \sin \left( \pi \frac{s+w}{2} \right) x^w dw$$

with  $-\Re\lambda < \Delta < -\Re s$ .

*Proof.* To change the order of summation in (5.2), absolute convergence of the both series is required. Applying (2.24), (2.25) and (2.26), we have

$$(5.5) \quad |\varphi(\lambda, x)| \ll X^{1/2} x^{\Re(\lambda)} \exp(-ax).$$

Using (2.2) we obtain that both sums in (5.2) are absolutely convergent for  $\Re(s) > 3/2$ . According to [5, p. 312, Eq. 1], the Mellin transform of  $\varphi(\lambda, x)$  for  $\Re w > -\Re\lambda$  is equal to

$$(5.6) \quad \tilde{\varphi}(\lambda, w) = \frac{\sinh \beta \Gamma(w + \lambda)}{\pi c^{w+\lambda}}.$$

It follows that

$$\Sigma(\lambda, s) = \zeta(2s) \sum_{q=1}^{\infty} \frac{1}{q} \frac{1}{2\pi i} \int_{(\Delta)} \tilde{\varphi}(\lambda, w) \sum_{n=1}^{\infty} \frac{S(1, n^2; q)}{n^{w+s}} \left( \frac{q}{4\pi} \right)^w dw.$$

We assume that  $\max(-\Re\lambda, 1 - \Re s) < \Delta < -1/2$  to guarantee absolute convergence of the integral over  $w$  and the sums over  $q, n$ . Note that due to (2.27) the function  $\tilde{\varphi}(\lambda, w)$  is of exponential decay in terms of  $\Im w$ . By [1, Lemma 5.1] one has

$$\sum_{n=1}^{\infty} \frac{S(1, n^2; q)}{n^{w+s}} = \sum_{d \pmod{q}} \frac{S(1, d^2; q)}{q^{w+s}} \zeta \left( \frac{d}{q}, 0, w+s \right).$$

Note that for all  $d$ , the Lerch zeta function has a simple pole at  $w = 1-s$  with residue one. We move the  $w$ -contour to the left up to  $\Delta_1 := -s - \epsilon$ ,

crossing a simple pole at  $w = 1 - s$ . For the resulting integral we apply functional equation (2.4). Finally,

$$\begin{aligned} \Sigma(\lambda, s) &= \zeta(2s)(4\pi)^{s-1}\tilde{\varphi}(\lambda, 1-s) \sum_{q=1}^{\infty} \frac{1}{q^{1+s}} \sum_{d \pmod{q}} S(1, d^2; q) + \\ &\quad \zeta(2s) \sum_{q=1}^{\infty} \frac{1}{q} \frac{1}{2\pi i} \int_{(\Delta_1)} \tilde{\varphi}(\lambda, w) \frac{2(2\pi)^{s+w-1}}{q^{s+w}} \Gamma(1-s-w) \times \\ &\quad \sin\left(\pi \frac{s+w}{2}\right) \sum_{d \pmod{q}} S(1, d^2; q) \zeta(0, d/q; 1-s-w) \left(\frac{q}{4\pi}\right)^w dw. \end{aligned}$$

Opening the Lerch zeta function and using [1, Lemma 4.1] we prove the lemma.  $\square$

**Lemma 5.2.** *The following relation holds*

$$(5.7) \quad I(\lambda, x) = \frac{\sinh \beta}{\pi c^\lambda} 2^{\lambda-s} \frac{\Gamma((1+\lambda-s)/2)\Gamma(1+(\lambda-s)/2)}{2\Gamma(1/2)} \\ \times \left(\frac{x}{c}\right)^{1-s} \left( \left(1 + \sqrt{-x^2/c^2}\right)^{s-\lambda-1} + \left(1 - \sqrt{-x^2/c^2}\right)^{s-\lambda-1} \right).$$

*Proof.* Moving the contour of integration in (5.4) to the right, we cross simple poles at  $w = 1 - s + j$ ,  $j = 0, 1, 2, \dots$ . Therefore,

$$I(\lambda, x) = \frac{\sinh \beta}{\pi c^\lambda} \left(\frac{x}{c}\right)^{1-s} \sum_{j=0}^{\infty} \frac{(-1)^j}{(2j)!} \Gamma(1+\lambda-s+2j) \left(\frac{x}{c}\right)^{2j}.$$

Using the duplication formula twice

$$\Gamma(1+\lambda-s+2j) = \frac{2^{\lambda-s+2j}}{\sqrt{\pi}} \Gamma((1+\lambda-s)/2+j) \Gamma(1+(\lambda-s)/2+j),$$

$$\Gamma(2j+1) = \frac{2^{2j}}{\sqrt{\pi}} \Gamma(j+1/2) \Gamma(j+1),$$

we obtain

$$(5.8) \quad I(\lambda, x) = \frac{\sinh \beta}{\pi c^\lambda} 2^{\lambda-s} \frac{\Gamma((1+\lambda-s)/2)\Gamma(1+(\lambda-s)/2)}{\Gamma(1/2)} \\ \times \left(\frac{x}{c}\right)^{1-s} {}_2F_1\left(\frac{1+\lambda-s}{2}, 1 + \frac{\lambda-s}{2}, \frac{1}{2}; -\frac{x^2}{c^2}\right).$$

Finally, [18, Eq. 15.4.7] yields the lemma.  $\square$

Analyzing the right-hand side of equation (5.3) we see that  $\Sigma(\lambda, s)$  can be continued analytically to the region of convergence of the series

$$\sum_{n=1}^{\infty} \frac{1}{n^{1-s}} \mathcal{L}_{n^2-4}(s) I\left(\lambda, \frac{n}{2}\right).$$

In view of (5.7) and (2.8), we prove analytic continuation of  $\Sigma(\lambda, s)$  to the region of our interest, namely  $\Re s = 1/2$  and  $\lambda = 1$ .

**Lemma 5.3.** *For  $\Re s = 1/2$  one has*

$$(5.9) \quad \Sigma(s) = (4\pi)^{s-1} \tilde{\varphi}(1, 1-s) \mathcal{L}_{-4}(s) + 2(2\pi)^{s-1} \sum_{n=1}^{\infty} \frac{1}{n^{1-s}} \mathcal{L}_{n^2-4}(s) I\left(\frac{n}{2}\right),$$

where

$$(5.10) \quad I(x) = \frac{\sinh \beta}{\pi c} 2^{1-s} \left(\frac{x}{c}\right)^{1-s} \frac{\Gamma(1-s/2)\Gamma(3/2-s/2)}{\Gamma(1/2)} \times \\ \left( \left(1 + \sqrt{-x^2/c^2}\right)^{s-2} + \left(1 - \sqrt{-x^2/c^2}\right)^{s-2} \right).$$

## 6. ANALYSIS OF $\Sigma_B(s)$

Substituting expression (2.31) to formula (4.3) we obtain

$$(6.1) \quad \Sigma_B(s) = \frac{1}{\pi^2} \sum_{k=1}^{\infty} (2k-1) \exp(-(2k-1)\beta) \Sigma_B(k, s),$$

where

$$(6.2) \quad \Sigma_B(k, s) = 2\pi(-1)^k \zeta(2s) \sum_{n=1}^{\infty} \frac{1}{n^s} \sum_{q=1}^{\infty} \frac{S(1, n^2; q)}{q} J_{2k-1}\left(\frac{4\pi n}{q}\right).$$

The analysis of the function  $\Sigma_B(k, s)$  was performed in [1, Lemma 5.1]. The main difference is that in [1, Lemma 5.1] we assumed that  $k \geq 6$  and now the sum is over all  $k \geq 1$ . The case  $k = 1$  is the most difficult one. In all other cases we can proceed exactly as in the proof of [1, Lemma 5.1]. The major problem for  $k = 1$  is that the line of integration cannot be moved to  $\Delta_1 := -s - \epsilon$  because it is required that  $1 - 2k < -s - \epsilon < -3/2 - \epsilon$ , which is possible only for  $k \geq 2$ . To avoid this problem we again use the "Hecke trick" in the same manner as in [4]. More precisely, we replace  $k$  in the order of the Bessel function by a parameter  $\lambda$  and assume that  $\Re \lambda > 5/4$ . In doing so we can carry on the

analysis of [1, Lemma 5.1] and obtain for  $\Re\lambda > (1 + \Re s)/2$ ,  $\Re s > 3/2$  the following expression

$$(6.3) \quad \Sigma_B(\lambda, s) = (2\pi)^s i^{2k} \frac{\Gamma(\lambda - s/2)}{\Gamma(\lambda + s/2)} \mathcal{L}_{-4}(s) + (2\pi)^s i^{2k} \sum_{n=1}^{\infty} \frac{1}{n^{1-s}} \mathcal{L}_{n^2-4l^2}(s) I_B(\lambda, n),$$

where

$$(6.4) \quad I_B(\lambda, x) := \frac{1}{2\pi i} \int_{(\Delta)} \frac{\Gamma(\lambda - 1/2 + w/2)}{\Gamma(\lambda + 1/2 - w/2)} \Gamma(1 - s - w) \sin\left(\pi \frac{s + w}{2}\right) x^w dw$$

with  $1 - 2\Re\lambda < \Delta < 1 - \Re s$ . As in [1, Lemma 5.1] we prove that for  $x \geq 2$  the following holds

$$(6.5) \quad I_B(\lambda, x) = \frac{2^{2\lambda}}{2^s \sqrt{\pi}} \cos\left(\pi\lambda - \frac{\pi s}{2}\right) \frac{\Gamma(\lambda - s/2) \Gamma(\lambda + 1/2 - s/2)}{\Gamma(2\lambda)} x^{1-2\lambda} \times {}_2F_1\left(\lambda - s/2, \lambda + 1/2 - s/2, 2\lambda; \frac{4}{x^2}\right).$$

The right-hand side of (6.3) yields analytic continuation of  $\Sigma_B(\lambda, s)$  to the region of convergence of the series

$$\sum_{n=1}^{\infty} \frac{1}{n^{1-s}} \mathcal{L}_{n^2-4l^2}(s) I_B(\lambda, n).$$

Using (6.5), (2.8) we obtain analytic continuation of  $\Sigma_B(\lambda, s)$  to  $\Re s = 1/2$ ,  $\lambda = 1$ .

**Lemma 6.1.** *For  $\Re s = 1/2$  the following formula holds*

$$(6.6) \quad \Sigma_B(s) = \frac{1}{\pi^2} \sum_{k=1}^{\infty} (2k-1) \exp(-(2k-1)\beta) \left( \frac{(2\pi)^s i^{2k} \Gamma(k - s/2)}{2 \Gamma(k + s/2)} \mathcal{L}_{-4}(s) \right. \\ + \frac{(2\pi)^s}{\sqrt{\pi}} \zeta(2s-1) \cos\left(\frac{\pi s}{2}\right) \frac{\Gamma(k - s/2) \Gamma(k + 1/2 - s/2)}{\Gamma(k + s/2) \Gamma(k - 1/2 + s/2)} \Gamma(s - 1/2) \\ + \frac{(2\pi)^s \sin(\pi s/2)}{\sqrt{\pi}} \mathcal{L}_{-3}(s) \Phi_k(s, 1/4) \\ \left. + \frac{\pi^s \cos(\pi s/2)}{\sqrt{\pi}} \sum_{n>2} \mathcal{L}_{n^2-4}(s) n^s \Psi_k(s, 4/n^2) \right),$$



where

(6.7)

$$\Phi_k(s, x) := \frac{\Gamma(k - s/2)\Gamma(1 - k - s/2)}{\Gamma(1/2)} {}_2F_1\left(k - \frac{s}{2}, 1 - k - \frac{s}{2}, \frac{1}{2}; \frac{1}{4}\right),$$

(6.8)

$$\Psi_k(s, x) := x^k \frac{\Gamma(k - s/2)\Gamma(k + 1/2 - s/2)}{\Gamma(2k)} {}_2F_1\left(k - \frac{s}{2}, k + \frac{1 - s}{2}, 2k; x\right).$$

## 7. SPECIAL FUNCTIONS

In order to obtain an upper bound for  $M_1(1/2 + ir)$ , it is required to estimate  $I(x)$ ,  $\Phi_k(s, x)$ ,  $\Psi_k(s, x)$  defined by (5.10), (6.7), (6.8) for  $\Re s = 1/2$ .

**Lemma 7.1.** *For  $s = 1/2 + ir$  one has*

$$(7.1) \quad I(x) \ll (1 + |r|)^{3/2} \left(\frac{x}{|c|}\right)^{1/2} \left|1 - \sqrt{-x^2/c^2}\right|^{-3/2}.$$

*Proof.* Let

$$(7.2) \quad \gamma_{\pm} = \arg(1 \pm \sqrt{-x^2/c^2}).$$

To estimate  $I(x)$  given by (5.10) we use the relation

$$(7.3) \quad |z^a| = |z|^{\Re a} \exp(-\Im a \arg z),$$

obtaining

$$\begin{aligned} I(x) \ll (1 + |r|)^{3/2} \left(\frac{x}{|c|}\right)^{1/2} \exp(-\pi|r|/2) \\ \times \left( \frac{\exp(-r(\arg c + \gamma_-))}{\left|1 - \sqrt{-x^2/c^2}\right|^{3/2}} + \frac{\exp(-r(\arg c + \gamma_+))}{\left|1 + \sqrt{-x^2/c^2}\right|^{3/2}} \right). \end{aligned}$$

Thus to prove (7.1) we need to show that

$$(7.4) \quad -\pi|r|/2 - r(-\pi/2 + \gamma + \gamma_{\pm}) \leq 0.$$

It follows from (2.27) that

$$(7.5) \quad 1 \pm \sqrt{-x^2/c^2} = 1 \pm \frac{x}{|c|} \cos \gamma \mp i \frac{x}{|c|} \sin \gamma.$$

In the plus case we have  $-\gamma < \gamma_+ < 0$  and thus (7.4) is satisfied. In the minus case we have  $0 < \gamma_- < \pi - \gamma$  and (7.4) is also satisfied.  $\square$

**Lemma 7.2.** *For  $s = 1/2 + ir$ , one has*

$$(7.6) \quad \Sigma(s) \ll \frac{(1 + |r|)^A X^\theta}{X^{1/4}} (X^{1/2} T^{1/2} + T^{3/2}),$$

where  $A$  is some positive constant.

*Proof.* Estimating (5.9) by absolute value and using (2.8) and (7.1) we obtain

$$(7.7) \quad \Sigma(s) \ll \frac{(1 + |r|)^A}{|c|^{1/2}} \sum_{n=1}^{\infty} n^{2\theta} \left| 1 - \sqrt{-n^2/(4c^2)} \right|^{-3/2}.$$

In view of (7.5) one has

$$\left| 1 - \sqrt{-x^2/c^2} \right|^2 = \left( 1 - \frac{x}{|c|} \right)^2 + 2 \frac{x}{|c|} \sin^2(\gamma/2).$$

Consequently,

$$(7.8) \quad \Sigma(s) \ll \frac{(1 + |r|)^A X^\theta}{X^{1/4}} \sum_{n=1}^{\infty} f\left(\frac{n}{2|c|}\right),$$

where

$$(7.9) \quad f(x) = \frac{x^{2\theta}}{((1-x)^2 + x\delta)^{3/4}}, \quad \delta = 2 \sin^2(\gamma/2) \asymp \frac{1}{8T^2}.$$

Calculating the first derivative of  $f(x)$  and solving the equation  $f'(x) = 0$ , we show that the function  $f(x)$  has only one critical point  $x_0 = 1 + O(\delta)$  belonging to  $(0, \infty)$ . Therefore,

$$(7.10) \quad \sum_{n=1}^{\infty} f\left(\frac{n}{2|c|}\right) = \int_0^{\infty} f\left(\frac{x}{2|c|}\right) dx + O\left(\max_{0 < x < \infty} |f(x)|\right) \ll \\ X^{1/2} \int_0^{1-\delta^{1/2}} \frac{x^{2\theta} dx}{(1-x)^{3/2}} + X^{1/2} \int_{1-\delta^{1/2}}^{1+\delta^{1/2}} \frac{x^{2\theta} dx}{(x\delta)^{3/4}} \\ + X^{1/2} \int_{1+\delta^{1/2}}^{\infty} \frac{x^{2\theta} dx}{(1-x)^{3/2}} + T^{3/2} \ll X^{1/2} T^{1/2} + T^{3/2}.$$

Substituting (7.10) to (7.8) yields the required result.  $\square$

It is not obvious how to improve (7.6) for general  $X$  and  $T$ . However, we can prove a better estimate in the special case when  $T$  is sufficiently large and  $\|X^{1/2} + X^{-1/2}\| \neq 0$ , where  $\|x\|$  denotes the distance between  $x$  and the nearest integer. Let

$$(7.11) \quad \kappa(X) = \|X^{1/2} + X^{-1/2}\|.$$

**Lemma 7.3.** *Assume that  $\kappa(X) \neq 0$  and*

$$(7.12) \quad T \gg \frac{X^{1/4+\epsilon}}{\kappa^{1/2}(X)}.$$

For  $s = 1/2 + ir$  one has

$$\Sigma(s) \ll \frac{(1+|r|)^A X^\theta}{X^{1/4}} \left( X^{1/2} \min \left( T, \frac{X^{1/2}}{\kappa(X)} \right)^{1/2} + \min \left( T, \frac{X^{1/2}}{\kappa(X)} \right)^{3/2} \right),$$

where  $A$  is some positive constant.

*Proof.* We would like to improve the estimate on the sum over  $n$  in (7.8). Calculating the first derivative of  $f(x)$  and solving the equation  $f'(x) = 0$  we obtain that the function  $f(x)$  has only one critical point  $x_0 = 1 + O(\delta)$  belonging to  $(0, \infty)$ . According to (7.10) the largest contribution to the estimate on  $\Sigma(s)$  comes from the sum over

$$|2|c| - n| \ll 2|c|T^{-1}.$$

Note that it may happen that for a sufficiently large  $T$  this region does not contain any integer number.

First of all, we analyse the value of  $\|2|c|\|$ . It follows from (2.25) that

$$(7.13) \quad \begin{aligned} 2|c| &= 2 \left( \cosh^2(\log \sqrt{X}) - \sin^2((2T)^{-1}) \right)^{1/2} \\ &= X^{1/2} + X^{-1/2} + O(X^{-1/2}T^{-2}). \end{aligned}$$

If  $X^{1/2} + X^{-1/2}$  is an integer then  $\|2|c|\| = O(X^{-1/2}T^{-2})$ . Since the sum over  $n$  in (7.8) contains summands

$$f\left(\frac{[2|c|]}{2|c|}\right) + f\left(\frac{[2|c|] + 1}{2|c|}\right) \ll \frac{1}{((\|2|c|\|/(2|c|))^2 + \delta)^{3/4}} \ll T^{3/2},$$

the estimate (7.6) cannot be improved in this case.

If  $X^{1/2} + X^{-1/2}$  is not an integer, then for

$$(7.14) \quad T \gg X^{-1/4+\epsilon} \|X^{1/2} + X^{-1/2}\|^{-1/2}$$

it follows that  $\|2|c|\|$  is very close to  $\|X^{1/2} + X^{-1/2}\|$ . The sum over  $n$  in (7.8) can be decomposed into two sums

$$(7.15) \quad \Sigma(s) \ll \frac{(1+|r|)^A X^\theta}{X^{1/4}} \left( \sum_{n=1}^{[2|c|]} f\left(\frac{n}{2|c|}\right) + \sum_{n=[2|c|]+1}^{\infty} f\left(\frac{n}{2|c|}\right) \right).$$

For  $n \leq [2|c|]$  the function  $f(n/(2|c|))$  is increasing and for  $n \geq [2|c|] + 1$  it is decreasing provided that

$$(7.16) \quad T \gg X^{1/4+\epsilon} \|X^{1/2} + X^{-1/2}\|^{-1/2}.$$

As a result,

$$\sum_{n=1}^{[2|c|]} f\left(\frac{n}{2|c|}\right) = \int_1^{[2|c|]} f\left(\frac{x}{2|c|}\right) dx + O\left(f\left(\frac{[2|c|]}{2|c|}\right)\right),$$

$$\sum_{n=[2|c|]+1}^{\infty} f\left(\frac{n}{2|c|}\right) = \int_{[2|c|]+1}^{\infty} f\left(\frac{x}{2|c|}\right) dx + O\left(f\left(\frac{[2|c|]+1}{2|c|}\right)\right).$$

Evaluating these integrals we obtain that the first sum can be bounded as follows

$$\sum_{n=1}^{[2|c|]} f\left(\frac{n}{2|c|}\right) \ll X^{1/2} \min\left(T^{1/2}, \left(\frac{2|c|}{\|2|c|\|}\right)^{1/2}\right) + \min\left(T^{3/2}, \left(\frac{2|c|}{\|2|c|\|}\right)^{3/2}\right),$$

and the same estimate is also valid for the second sum.  $\square$

**Lemma 7.4.** *For  $s = 1/2 + 2ir$ ,  $0 < x < 1$  the following estimates hold*

$$(7.17) \quad \exp(\pi|r|)\Psi_k(s, x) \ll (1-x)^{1/4} \frac{(1+|r|)^{2k-2a-1}}{2^{2k} k^{a+1/2}} \left(\frac{x}{1-x}\right)^{k-a},$$

$$(7.18) \quad \exp(\pi|r|)\Psi_k(s, x) \ll (1-x)^{1/4} \frac{(1+|r|)^{2b}}{k^{2b-1/2}} \left(\frac{x}{1-x}\right)^b,$$

where  $a, b$  are some absolute constants such that  $0 < a < k - 1/4$ ,  $1/4 < b < k$ .

*Proof.* Using [18, Eq. 15.8.1] we obtain

$$\Psi_k(s, x) = \frac{x^k}{(1-x)^{k-s/2}} \frac{\Gamma(k-s/2)\Gamma(k+1/2-s/2)}{\Gamma(2k)} \times {}_2F_1\left(k - \frac{s}{2}, k - \frac{1-s}{2}, 2k; -\frac{x}{1-x}\right).$$

Writing the Mellin-Barnes integral representation for the hypergeometric function [18, Eq. 15.6.6], we have

$$(7.19) \quad \Psi_k(s, x) = \frac{x^k}{(1-x)^{k-s/2}} \frac{\Gamma(k+1/2-s/2)}{\Gamma(k-1/2+s/2)} \times \frac{1}{2\pi i} \int_{\Delta} \frac{\Gamma(k-s/2+z)\Gamma(k-1/2+s/2+z)\Gamma(-z)}{\Gamma(2k+z)} \left(\frac{x}{1-x}\right)^z dz,$$

where  $1/4 - k < \Delta < 0$ . To prove (7.17) we move the contour of integration to the line  $\Re z = -a$  such that  $0 < a < k - 1/4$ . Arguing in the same way as in [10] we prove

$$(7.20) \quad \Psi_k(s, x) \ll (1-x)^{1/4} (k+|r|)^{1/2} \left( \frac{x}{1-x} \right)^{k-a} \frac{|\Gamma(k-1/4-a+ir)|^2}{\Gamma(2k-a)}.$$

To simplify notation in estimates of the quotient of gamma factors, let us assume that  $a := 1/4 + n$ , where  $n$  is a positive integer. Note that all our arguments are valid for an arbitrary  $a$ . One has

$$|\Gamma(k-1/2-n+ir)| = \prod_{j=1}^{k-n-2} |j+1/2+ir| |\Gamma(3/2+ir)| \ll (1+|r|)^{k-n-1} |\exp(-\pi|r|/2) \Gamma(k-1/2-n)|$$

since

$$|j+1/2+ir| \leq j+1/2+|r| \leq (j+1/2)(1+|r|), \text{ for } j \geq 1.$$

Therefore,

$$(7.21) \quad \frac{|\Gamma(k-1/4-a+ir)|^2}{\Gamma(2k-a)} \ll (1+|r|)^{2k-2a-3/2} |\exp(-\pi|r|)| \frac{\Gamma^2(k-1/4-a)}{\Gamma(2k-a)} \ll \frac{(1+|r|)^{2k-2a-3/2} |\exp(-\pi|r|)|}{2^{2k} k^{1+a}}.$$

Substituting (7.21) to (7.20) we obtain (7.17).

To prove estimate (7.18) we move the contour of integration in (7.19) to the line  $\Re z = b - k$  such that  $1/4 < b < k$ . Consequently,

$$(7.22) \quad \Psi_k(s, x) \ll (1-x)^{1/4} \left( \frac{x}{1-x} \right)^b (k+|r|)^{1/2} \times \int_{-\infty}^{\infty} \left| \frac{\Gamma(b-1/4+i(t-r)) \Gamma(b-1/4+i(t+r)) \Gamma(k-b-it)}{\Gamma(k+b+it)} \right| dt \ll (1-x)^{1/4} \left( \frac{x}{1-x} \right)^b (k+|r|)^{1/2} \times \int_{-\infty}^{\infty} \frac{\exp(-\pi(|t-r|+|t+r|)/2) (1+|t-r|)^{b-3/4} (1+|t+r|)^{b-3/4}}{(k+|t|)^{2b}} dt \ll (1-x)^{1/4} \left( \frac{x}{1-x} \right)^b (k+|r|)^{1/2} \exp(-\pi|r|) \frac{(1+|r|)^{2b-1/2}}{k^{2b}}.$$

This completes the proof of (7.18).  $\square$

**Lemma 7.5.** *For  $s = 1/2 + 2ir$  the following estimate holds*

$$(7.23) \quad \exp(\pi|r|)\Phi_k(s, 1/4) \ll \frac{4^k}{k + |r|}.$$

*Proof.* Using [18, Eq. 15.8.1] and the Mellin-Barnes integral for the hypergeometric function [18, Eq. 15.6.6], we obtain

$$(7.24) \quad \exp(\pi|r|)\Phi_k(s, 1/4) = \exp(\pi|r|) \frac{(4/3)^k \Gamma(1 - k - s/2)}{\Gamma(k - 1/2 + s/2)} \times \\ \frac{1}{2\pi i} \int \frac{\Gamma(k - s/2 + z) \Gamma(k - 1/2 + s/2 + z) \Gamma(-z)}{\Gamma(1/2 + z)} \frac{dz}{3^z}.$$

Applying [18, Eq. 5.5.3] and moving the contour of integration in (7.24) to the line  $\Re z = 3/4 - k$  yields the estimate

$$\exp(\pi|r|)\Phi_k(s, 1/4) \ll \frac{4^k}{|\Gamma(k - 1/4 + ir)\Gamma(k + 1/4 + ir)|} \times \\ \int_{-\infty}^{\infty} \left| \frac{\Gamma(1/2 + i(t - r)) \Gamma(1/2 + i(t + r)) \Gamma(k - 3/4 - it)}{\Gamma(-k + 5/4 + it)} \right| dt.$$

Applying [18, Eq. 5.5.3] once again together with [18, Eq. 5.4.4] gives

$$(7.25) \quad \exp(\pi|r|)\Phi_k(s, 1/4) \ll \frac{4^k}{|\Gamma(k + ir)|^2} \int_0^{\infty} \frac{|\Gamma(k + it)|^2 \exp(\pi t)}{(\cosh(2\pi t) + \cosh(2\pi r))^{1/2} (k + t)} dt.$$

Using the methods of [10, Section 2] one can show that

$$(7.26) \quad |\Gamma(k + it)|^2 \asymp \exp(-\pi|t|) \begin{cases} \Gamma^2(k) \exp(|t|(\pi - g_1(|t|/(k - 1))), & \text{if } 0 \leq |t| \leq k - 1, \\ |t|^{2k-1} \exp(|t|g_2((k - 1)/|t|)), & \text{if } k - 1 < |t|. \end{cases}$$

To estimate (7.25) we consider separately two cases:  $k - 1 \leq |r|$  and  $k - 1 > |r|$ . For  $k - 1 \leq |r|$  using (7.26) we obtain

$$\exp(\pi|r|)\Phi_k(s, 1/4) \ll \frac{4^k}{|\Gamma(k + ir)|^2} \\ \times \left( \Gamma^2(k) \exp(-\pi|r|) k^{-1} \int_0^{k-1} \exp(t(\pi - g_1(t/(k - 1)))) dt + \right. \\ \left. \exp(-\pi|r|) \int_{k-1}^r \exp((2k - 2) \log t + tg_2((k - 1)/t)) dt + \right. \\ \left. \int_r^{\infty} \exp(-\pi t + (2k - 2) \log t + tg_2((k - 1)/t)) dt \right).$$

Estimating the integrals by the means of [10, Lemma 3] we show that

$$\exp(\pi|r|)\Phi_k(s, 1/4) \ll \frac{4^k}{|\Gamma(k+ir)|^2} \exp(-\pi|r|)r^{2k-2} \exp(r g_2((k-1)/r)).$$

Applying (7.26) we finally prove that

$$\exp(\pi|r|)\Phi_k(s, 1/4) \ll 4^k/(1+|r|).$$

The case  $k-1 > |r|$  can be treated in the same way.  $\square$

**Remark 7.6.** *In the case when  $s = 1/2$  one has (see [1, Eq. 2.12]) a much better bound  $\Phi_k(1/2, 1/4) \ll k^{-1/2}$ . We expect that for  $s = 1/2 + 2ir$  the estimate of the following shape*

$$(7.27) \quad \exp(\pi|r|)\Phi_k(s, 1/4) \ll (1+|r|)^a k^{-1/2}$$

*should be correct for some positive constant  $a$ .*

**Lemma 7.7.** *For  $s = 1/2 + 2ir$ ,  $X > 5$  one has*

$$(7.28) \quad \Sigma_B(s) \ll (1+|r|)^A X^{-1/2},$$

*where  $A$  is some positive constant.*

*Proof.* To estimate (6.6) we use (2.24), (2.8), Lemmas 7.4 and 7.5, together with some classical estimates on the Gamma function and the Riemann zeta function. Consequently,

$$(7.29) \quad \sum_{k=1}^{\infty} (2k-1) \exp(-(2k-1)\beta) \frac{(2\pi)^s \sin(\pi s/2)}{\sqrt{\pi}} \mathcal{L}_{-3}(s) \Phi_k(s, 1/4) \\ \ll (1+|r|)^A \sum_{k=1}^{\infty} (2/\sqrt{X})^{2k-1} \ll (1+|r|)^A X^{-1/2}.$$

To estimate the contribution of

$$\sum_{k=1}^{\infty} (2k-1) \exp(-(2k-1)\beta) \frac{\pi^s \cos(\pi s/2)}{\sqrt{\pi}} \sum_{n>2} \mathcal{L}_{n^2-4}(s) n^s \Psi_k(s, 4/n^2)$$

we divide the sum over  $n$  in two parts with the conditions  $n \leq 4(1+|r|)$  and  $n > 4(1+|r|)$ . For the first sum we apply (7.18) and obtain

$$(1+|r|)^A \sum_{k=1}^{\infty} \frac{1}{k^{2b-3/2} X^{k-1/2}} \ll (1+|r|)^A X^{-1/2}.$$

For the second sum over  $n > 4(1 + |r|)$  we use the estimate (7.17) and obtain

$$(1 + |r|)^{A_1} \sum_{k=1}^{\infty} \frac{k}{X^{k-1/2}} \sum_{n > 4(1+|r|)} \frac{n^{2\theta+\epsilon}}{(n^2/4 - 1)^{k-a-1/4}} \frac{(1 + |r|)^{2k-2a-1}}{2^{2k} k^{a+1/2}} \\ \ll (1 + |r|)^A X^{-1/2}.$$

□

**Remark 7.8.** *We only need the condition  $X > 5$  in Lemma 7.7 to estimate the contribution of  $\Phi_k(s, 1/4)$ . In particular, estimate (7.27) would result in a bound of the type*

$$(7.30) \quad \Sigma_B(s) \ll \frac{(1 + |r|)^A}{(X - 1)^B}$$

for any  $X > 1$  with some positive constants  $A$  and  $B$ .

## 8. ESTIMATES FOR THE MOMENT OF MAASS $\text{sym}^2$ $L$ -FUNCTIONS

**Lemma 8.1.** *For  $\Re s = 1/2$  the following formula holds*

$$(8.1) \quad M_1(s) = \zeta(2s)\varphi_0 + \Sigma(s) - \Sigma_B(s) - \\ \frac{2\zeta(s)}{\pi} \int_0^\infty \frac{\zeta(s + 2it)\zeta(s - 2it)}{|\zeta(1 + 2it)|^2} \hat{\varphi}(t) dt - 2 \frac{\zeta(2s - 1)}{\zeta(2 - s)} \hat{\varphi}\left(\frac{1 - s}{2i}\right),$$

where  $\Sigma(s)$  is given by (5.9) and  $\Sigma_B(s)$  by (6.6).

*Proof.* In order to prove that there exist an analytic continuation of (4.4) to the critical line  $\Re s = 1/2$ , it is only left to consider

$$(8.2) \quad \frac{2\zeta(s)}{\pi} \int_0^\infty \frac{\zeta(s + 2it)\zeta(s - 2it)}{|\zeta(1 + 2it)|^2} \hat{\varphi}(t) dt \\ = \frac{\zeta(s)}{2\pi i} \int_{(0)} \frac{\zeta(s + z)\zeta(s - z)}{\zeta(1 + z)\zeta(1 - z)} \hat{\varphi}\left(\frac{z}{2i}\right) dz.$$

The continuation of  $\Sigma(s)$  and  $\Sigma_B(s)$  is given by (5.9) and (6.6), respectively. Arguing in the same manner as in [2, Theorem 7.3] we obtain (8.1). □

The procedure of analytic continuation of (8.1) to the critical point  $s = 1/2$  is not straightforward since there are two summands on the right-hand side of (8.1) with simple poles at  $s = 1/2$ .



**Lemma 8.2.** *The following formula holds at the critical point*

(8.3)

$$\begin{aligned} M_1(1/2) = & -\frac{1}{2\pi^2} \sum_{k=1}^{\infty} (2k-1) \exp(-(2k-1)\beta) (\psi(k-1/4) + \psi(k+1/4)) \\ & - \frac{\cosh \beta}{4\pi^2 \sinh^2 \beta} \left( 3\gamma + \frac{\pi}{2} - 3 \log(2\pi) \right) + \Sigma(1/2) - \Sigma_B(1/2) \\ & - \frac{2\zeta(1/2)}{\pi} \int_0^{\infty} \frac{\zeta(1/2+2it)\zeta(1/2-2it)}{|\zeta(1+2it)|^2} \hat{\varphi}(t) dt - 2 \frac{\zeta(0)}{\zeta(3/2)} \hat{\varphi}\left(\frac{1}{4i}\right), \end{aligned}$$

where  $\Sigma(1/2)$  is given by (5.9) and  $\Sigma_B(1/2)$  by (6.6) with the omitted summand containing  $\zeta(2s-1)$ .

*Proof.* The two summands that have a simple pole at  $s = 1/2$  are

$$(8.4) \quad \zeta(2s)\varphi_0 - \frac{1}{\pi^2} \frac{(2\pi)^s}{\sqrt{\pi}} \zeta(2s-1) \cos\left(\frac{\pi s}{2}\right) \Gamma(s-1/2) \times \\ \sum_{k=1}^{\infty} (2k-1) \exp(-(2k-1)\beta) \frac{\Gamma(k-s/2)\Gamma(k+1/2-s/2)}{\Gamma(k+s/2)\Gamma(k-1/2+s/2)}.$$

For simplicity let  $s = 1/2 + u$  with  $u \rightarrow 0$ . Using (2.29) and the functional equation for the Riemann zeta-function we obtain that (8.4) is equal to

(8.5)

$$\begin{aligned} \zeta(1+2u) \frac{-\cosh \beta}{2\pi^2 \sinh^2 \beta} - \frac{\zeta(1-2u)}{\pi^2} \frac{(2\pi)^{1/2+3u}}{\sqrt{\pi}} \cos\left(\frac{\pi}{4} + \frac{\pi u}{2}\right) \frac{\Gamma(1-2u)}{\Gamma(1-u)} \times \\ \sum_{k=1}^{\infty} (2k-1) \exp(-(2k-1)\beta) \frac{\Gamma(k-1/4-u/2)\Gamma(k+1/4-u/2)}{\Gamma(k+1/4+u/2)\Gamma(k-1/4+u/2)}. \end{aligned}$$

Note that (8.5) is holomorphic at  $u = 0$ . This follows from the relation

$$\sum_{k=1}^{\infty} (2k-1) \exp(-(2k-1)\beta) = \frac{\cosh \beta}{2 \sinh^2 \beta}.$$

Computing the limit of (8.5) as  $u$  tends to 0 by L'hospital's rule we prove the lemma.  $\square$

**Theorem 8.3.** *Let  $X > 5$  then for  $s = 1/2 + ir$  the following holds*

$$(8.6) \quad M_1(s) \ll (1+|r|)^A T \log^3 T + \\ \frac{(1+|r|)^A X^\theta}{X^{1/4}} \left( X^{1/2} \min\left(T, \frac{X^{1/2}}{\kappa(X)}\right)^{1/2} + \min\left(T, \frac{X^{1/2}}{\kappa(X)}\right)^{3/2} \right).$$

*Proof.* This is a consequence of (7.28), (7.3), (7.6) and (8.1). To estimate the integral containing Riemann zeta functions in (8.1) it is sufficient to use the standard bound  $|\zeta^{-1}(1+it)| \ll \log t$ , Cauchy-Schwarz inequality and the estimate on the second moment of the Riemann-zeta function

$$\int_0^T |\zeta(1/2+it)|^2 dt \ll T \log T.$$

As a result,

$$\frac{2\zeta(s)}{\pi} \int_0^\infty \frac{\zeta(s+2it)\zeta(s-2it)}{|\zeta(1+2it)|^2} \hat{\varphi}(t) dt \ll (1+|r|)^A T \log^3 T.$$

From (2.18) it follows that

$$2 \frac{\zeta(2s-1)}{\zeta(2-s)} \hat{\varphi}\left(\frac{1-s}{2i}\right) \ll (1+|r|)^A X^{1/4}.$$

This error term is absorbed by other terms in (8.6).  $\square$

## 9. ESTIMATES FOR THE SPECTRAL EXPONENTIAL SUM

**Theorem 9.1.** *For  $X > 5$  one has*

$$(9.1) \quad \sum_j X^{it_j} \exp(-t_j/T) \ll S(X, T, \kappa(X)) \log^2 T,$$

where

$$(9.2) \quad S(X, T, \kappa(X)) = \begin{cases} X^{1/4+\theta/2} T^{1/2} & \text{if } T \leq X^{1/2}; \\ X^{\theta/2} T & \text{if } X^{1/2} < T \leq \frac{X^{1/2}}{\kappa(X)}; \\ X^{\theta/2} T^{1/4} \left(\frac{X^{1/2}}{\kappa(X)}\right)^{3/4} & \text{if } \frac{X^{1/2}}{\kappa(X)} < T \leq \frac{X^{1/2+2\theta/3}}{\kappa(X)}; \\ T & \text{if } \frac{X^{1/2+2\theta/3}}{\kappa(X)} < T. \end{cases}$$

*Proof.* Substituting (8.6) to (3.11) we obtain

$$\begin{aligned} \sum_j X^{it_j} \exp(-t_j/T) &\ll \frac{NX^{1/2} \log T}{(\max(1, NX^{1/2}/T))^{1/2}} + \sqrt{N/X} \log^2 N + T \log^2 T + \\ &\frac{1}{N^{1/2}} \left( T \log^3 T + \frac{X^\theta}{X^{1/4}} \left( X^{1/2} \min\left(T, \frac{X^{1/2}}{\kappa(X)}\right)^{1/2} + \min\left(T, \frac{X^{1/2}}{\kappa(X)}\right)^{3/2} \right) \right). \end{aligned}$$

Now we consider different cases.

If  $T \leq X^{1/2}$  then taking  $N = X^\theta \log^{-1} T$  we obtain

$$(9.3) \quad \sum_j X^{it_j} \exp(-t_j/T) \ll X^{1/4+\theta/2} T^{1/2} \log^{1/2} T.$$

If  $X^{1/2} < T \leq X^{1/2}/\kappa(X)$  then choosing  $N = TX^{\theta-1/2} \log^{-1} T$  we prove that

$$(9.4) \quad \sum_j X^{it_j} \exp(-t_j/T) \ll X^{\theta/2} T \log^{1/2} T.$$

If  $X^{1/2}/\kappa(X) < T \leq X^{1/2+2\theta/3}/\kappa(X)$  then letting

$$N = \frac{X^\theta}{T^{1/2} X^{1/2} \log T} \left( \frac{X^{1/2}}{\kappa(X)} \right)^{3/2}$$

it follows that

$$(9.5) \quad \sum_j X^{it_j} \exp(-t_j/T) \ll X^{\theta/2} T^{1/4} \left( \frac{X^{1/2}}{\kappa(X)} \right)^{3/4} \log^{1/2} T + T \log^2 T \ll X^{\theta/2} T^{1/4} \left( \frac{X^{1/2}}{\kappa(X)} \right)^{3/4} \log^2 T.$$

If  $X^{1/2+2\theta/3}/\kappa(X) < T$  then taking  $N = X^{2\theta/3}/\kappa(X)$  we have

$$(9.6) \quad \sum_j X^{it_j} \exp(-t_j/T) \ll T \log^2 T + \frac{X^{1/2+2\theta/3}}{\kappa(X)} \log T \ll T \log^2 T.$$

□

Note that it is sufficient to use only the first estimate in (9.2) in order to prove (1.11). Other estimates in (9.2), as well as Lemma 7.3, serve to establish (1.7) unconditionally in some ranges.

**Theorem 9.2.** *For  $X > 5$  the following estimates hold*

$$(9.7) \quad \sum_{t_j \leq T} X^{it_j} \ll \max(X^{1/4+\theta/2} T^{1/2}, X^{\theta/2} T) \log^3 T,$$

$$(9.8) \quad \sum_{t_j \leq T} X^{it_j} \ll T \log^2 T, \quad \text{if } T > \frac{X^{1/2+7\theta/6}}{\kappa(X)}.$$

*Proof.* To prove the theorem we need to substitute the bound (9.1) to equation (3.3). Consequently,

$$(9.9) \quad \sum_{t_j \leq T} X^{it_j} \ll \log^2 T \int_{-1}^1 |\hat{g}(\xi)| S(X \exp(-2\pi\xi), T, \kappa(X \exp(-2\pi\xi))) d\xi.$$

Applying the estimate

$$(9.10) \quad S(X, T, \kappa(X)) \leq \max(X^{1/4+\theta/2} T^{1/2}, X^{\theta/2} T)$$

that follows from (9.2), we immediately obtain (9.7). Estimate (9.8) can be proved using the fact that

$$(9.11) \quad S(X, T, \kappa(X)) = T \quad \text{if} \quad X^{1/2+2\theta/3}/\kappa(X) < T.$$

To this end, we decompose the integral (9.9) into two parts. The first one is over  $\xi$  such that  $\kappa(X \exp(-2\pi\xi)) \leq \delta$ . To estimate this integral we will apply (9.10). To estimate the second integral we will use (9.11). According to these estimates, we will choose an optimal value of parameter  $\delta$ .

First, making the change of variables we have

$$(9.12) \quad \sum_{t_j \leq T} X^{it_j} \ll \frac{\log^2 T}{X} \int_{X \exp(-2\pi)}^{X \exp(2\pi)} \left| \hat{g} \left( \frac{\log(Z/X)}{2\pi} \right) \right| S(Z, T, \kappa(Z)) dZ.$$

Denote  $Z_n = (n + \sqrt{n^2 - 4})^2/4$  and let

$$Z_{n_1-1} < X \exp(-2\pi) \leq Z_{n_1}, \quad Z_{n_2} \leq X \exp(2\pi) < Z_{n_2+1}.$$

Consequently,

$$\begin{aligned} \sum_{t_j \leq T} X^{it_j} &\ll \frac{\log^2 T}{X} \int_I \left| \hat{g} \left( \frac{\log(Z/X)}{2\pi} \right) \right| S(Z, T, \kappa(Z)) dZ \\ &+ \frac{\log^2 T}{X} \sum_{n_1 \leq n \leq n_2} \int_{|Z-Z_n| \leq \delta\sqrt{Z_n}} \left| \hat{g} \left( \frac{\log(Z/X)}{2\pi} \right) \right| S(Z, T, \kappa(Z)) dZ, \end{aligned}$$

where

$$I = [X \exp(-2\pi), X \exp(2\pi)] \setminus \bigcup_{n_1 \leq n \leq n_2} [Z_n - \delta\sqrt{Z_n}, Z_n + \delta\sqrt{Z_n}].$$

For  $Z \in I$  one has  $\kappa(Z) \gg \delta$ . To estimate the integral over  $I$  we use (9.11). Bounds for the remaining integrals rely on (9.10). Finally, we obtain for  $T > X^{1/2+2\theta/3}/\delta$

$$(9.13) \quad \sum_{t_j \leq T} X^{it_j} \ll T \log^3 T + \frac{TX^{\theta/2} \log^2 T}{X} \sum_{n_1 \leq n \leq n_2} \int_{|Z-Z_n| \leq \delta\sqrt{Z_n}} \left| \hat{g} \left( \frac{\log(Z/X)}{2\pi} \right) \right| dZ.$$

It follows from (3.2) that

$$\left| \hat{g} \left( \frac{\log(Z/X)}{2\pi} \right) \right| \ll T \quad \text{when} \quad |Z - X| \ll \frac{X}{T}.$$

Note that if

$$(9.14) \quad \kappa(X) \gg \delta \quad \text{and} \quad \kappa(X)X^{1/2} \gg \frac{X}{T},$$

then the interval  $|Z - X| \ll \frac{X}{T}$  does not intersect with

$$\bigcup_{n_1 \leq n \leq n_2} [Z_n - \delta\sqrt{Z_n}, Z_n + \delta\sqrt{Z_n}].$$

Thus applying (3.2) we obtain

$$(9.15) \quad \int_{|Z-Z_n| \leq \delta\sqrt{Z_n}} \left| \hat{g} \left( \frac{\log(Z/X)}{2\pi} \right) \right| dZ \ll X \int_{|X \exp(y) - Z_n| \leq \delta\sqrt{Z_n}} \frac{dy}{y}.$$

Let  $Z_{n_0}$  be the nearest  $Z_n$  to  $X$  and let  $Y_j = Z_{n_0+j}$ . Then

$$(9.16) \quad \sum_{n_1 \leq n \leq n_2} \int_{|Z-Z_n| \leq \delta\sqrt{Z_n}} \left| \hat{g} \left( \frac{\log(Z/X)}{2\pi} \right) \right| dZ \\ \ll \sum_{|j| \ll X^{1/2}} X \int_{|X \exp(y) - Y_j| \leq \delta\sqrt{Y_j}} \frac{dy}{y}.$$

The interval of integration in (9.16) is equal to

$$(9.17) \quad \log \left( \frac{Y_j}{X} - \frac{\delta\sqrt{Y_j}}{X} \right) \leq y \leq \log \left( \frac{Y_j}{X} + \frac{\delta\sqrt{Y_j}}{X} \right).$$

Since the point  $Y_j$  is very close to the point  $X \pm \kappa(X)\sqrt{X} + j\sqrt{X}$ , the interval has the following form

$$(9.18) \quad \log \left( 1 + \frac{j \pm \kappa(X) - \delta}{X^{1/2}} \right) \leq y \leq \log \left( 1 + \frac{j \pm \kappa(X) + \delta}{X^{1/2}} \right).$$

Using the fact that for a small  $\alpha$

$$\int_{\log a}^{\log(a+\alpha)} \frac{dy}{y} \ll \frac{\alpha}{\log a}$$

and applying estimate (9.14), we prove

$$(9.19) \quad \int_{|X \exp(y) - Y_j| \leq \delta\sqrt{Y_j}} \frac{dy}{y} \ll \frac{\delta}{|j + \kappa(X)|}.$$

Substituting (9.19) to (9.16) yields

$$(9.20) \quad \sum_{n_1 \leq n \leq n_2} \int_{|Z-Z_n| \leq \delta\sqrt{Z_n}} \left| \hat{g} \left( \frac{\log(Z/X)}{2\pi} \right) \right| dZ \ll \frac{X\delta}{\kappa(X)} \log X.$$

Substituting (9.20) to (9.13) we obtain

$$(9.21) \quad \sum_{t_j \leq T} X^{it_j} \ll T \log^3 T + \frac{TX^{\theta/2} \log^3 T}{\kappa(X)} \delta$$

under conditions (9.14) and for  $T > X^{1/2+2\theta/3}/\delta$ . Choosing  $\delta = \kappa(X)X^{-\theta/2}$ , we conclude that (9.14) is satisfied under assumptions of the lemma. This completes the proof of (9.8).  $\square$

#### ACKNOWLEDGEMENTS

We thank the referee for many helpful suggestions.

#### REFERENCES

- [1] O. Balkanova and D. Frolenkov, *On the mean value of symmetric square  $L$ -functions*, Algebra Number Theory 12 (2018), 35–59.
- [2] O. Balkanova and D. Frolenkov, *Convolution formula for the sums of generalized Dirichlet  $L$ -functions*, Revista Math. Iberoamericana, to appear, arXiv:1709.01365 [math.NT].
- [3] V.A. Bykovskii, *Density theorems and the mean value of arithmetic functions on short intervals*. (Russian) Zap. Nauchn. Sem. S.-Peterburg. Otdel. Mat. Inst. Steklov. (POMI) 212 (1994), Anal. Teor. Chisel i Teor. Funktsii. 12, 56–70, 196; translation in J. Math. Sci. (New York) 83 (1997), no. 6, 720–730.
- [4] V. A. Bykovskii and D. A. Frolenkov, *Asymptotic formulas for the second moments of  $L$ -series associated to holomorphic cusp forms on the critical line*, Izvestiya: Mathematics 81:2, 2017, 239–268.
- [5] H. Beitman and A. Erdelyi, *Tables of integral transforms*, Vol. 1, McGraw-Hill, New York, 1954.
- [6] Y. Cai, *Prime geodesic theorem*, J. Theor. Nombres Bordeaux 14:1 (2002), 59–72.
- [7] G. Cherubini and J. Guerreiro, *Mean square in the prime geodesic theorem*, Algebra Number Theory 12 (2018), 571–597.
- [8] J. B. Conrey and H. Iwaniec, *The cubic moment of central values of automorphic  $L$ -functions*, Ann. of Math. (2) 151 (2000), 1175–1216.
- [9] J. M. Deshouillers and H. Iwaniec, *The non-vanishing of Rankin-Selberg zeta-functions at special points*, Selberg trace formula and related topics, Contemp. Math. 53, Amer. Math. Soc., Providence, RI, 1986, 59–95.
- [10] D. A. Frolenkov, *On the uniform bounds on hypergeometric function*, Far Eastern Math. J. 15:2 (2015), 288–298.
- [11] I. S. Gradshteyn and I. M. Ryzhik, *Table of Integrals, Series, and Products*. Edited by A. Jeffrey and D. Zwillinger. Academic Press, New York, 7th edition, 2007.
- [12] H. Iwaniec, *Prime geodesic theorem*, J. Reine. Angew. Math. 349 (1984), 136–159.
- [13] H. Iwaniec, *Introduction to the spectral theory of automorphic forms*, Revista Matemática Iberoamericana, 1995.
- [14] N. V. Kuznetsov, *Petersson’s conjecture for cusp of weight zero and Linnik’s conjecture, Sums of Kloosterman sums*, Mat. Sb. 111 (1980), 334–383.

- [15] M. Lerch, *Note sur la fonction*  $R(w, x, s) = \sum_{k=0}^{\infty} \frac{e^{2\pi i n a}}{(n+c)^s}$ , Acta Math. 11 (1887), 19–24.
- [16] W. Luo, *Values of symmetric square L-functions at 1*, J. Reine Angew. Math. 506 (1999), 215–235.
- [17] W. Luo and P. Sarnak, *Quantum ergodicity of eigenfunctions on  $PSL_2(Z)/H^2$* , Pub. math. de l’I.H.E.S. 81 (1995), 207–237.
- [18] F. W. J. Olver, D. W. Lozier, R. F. Boisvert and C. W. Clarke, *NIST Handbook of Mathematical Functions*, Cambridge University Press, Cambridge (2010).
- [19] Y. N. Petridis and M. S. Risager, *Averaging over Heegner points in the hyperbolic circle problem*, IMRN (2017), <https://doi.org/10.1093/imrn/rnx026>.
- [20] Y. N. Petridis and M. S. Risager, *Local average in hyperbolic lattice point counting, with an appendix by N. Laaksonen*, Math. Z. 285 (2017), no. 3-4, 1319–1344.
- [21] K. Soundararajan and M. P. Young, *The prime geodesic theorem*, J. Reine Angew. Math. 676 (2013), 105–120.
- [22] A. Weil, *On some exponential sums*, Proc. Natl. Acad. Sci. USA 34 (1948), 204–207.
- [23] M.P. Young, *Weyl-type hybrid subconvexity bounds for twisted L-functions and Heegner points on shrinking sets*, J. Eur. Math. Soc. 19 (2017), 1545–1576.
- [24] D. Zagier, *Modular forms whose Fourier coefficients involve zeta-functions of quadratic fields*. Modular functions of one variable, VI (Proc. Second Internat. Conf., Univ. Bonn, Bonn, 1976), pp. 105–169. Lecture Notes in Math., Vol. 627, Springer, Berlin, 1977.

DEPARTMENT OF MATHEMATICAL SCIENCES, CHALMERS UNIVERSITY OF TECHNOLOGY AND UNIVERSITY OF GOTHENBURG, CHALMERS TVÄRGATA 3, 412 58 GOTHENBURG, SWEDEN

*E-mail address:* `olgabalkanova@gmail.com`

NATIONAL RESEARCH UNIVERSITY HIGHER SCHOOL OF ECONOMICS, MOSCOW, RUSSIA AND STEKLOV MATHEMATICAL INSTITUTE OF RUSSIAN ACADEMY OF SCIENCES, 8 GUBKINA ST., MOSCOW, 119991, RUSSIA

*E-mail address:* `frolenkov@mi.ras.ru`