

# ON SOME RANDOM VARIABLES INVOLVING BERNOULLI RANDOM VARIABLE

ROMEO MEŠTROVIĆ

**ABSTRACT.** Motivated by the recent investigations given in [25] and the fact that Bernoulli probability-type models were often used in the study on some problems in theory of compressive sensing, here we define and study the complex-valued discrete random variables  $\tilde{X}_l(m, N)$  ( $0 \leq l \leq N - 1$ ,  $1 \leq m \leq N$ ). Each of these random variables is defined as a linear combination of  $N$  independent identically distributed  $0 - 1$  Bernoulli random variables. We prove that for  $l \neq 0$ ,  $\tilde{X}_l(m, N)$  is the zero-mean random variable, and we also determine the variance of  $\tilde{X}_l(m, N)$  and its real and imaginary parts. Notice that  $\tilde{X}_l(m, N)$  belongs to the class of sub-Gaussian random variables that are significant in some areas of theory of compressive sensing. In particular, we prove some probability estimates for the mentioned random variables. These estimates are used to establish the upper bounds of the sub-Gaussian norm of their real and imaginary parts. We believe that our results should be implemented in certain applications of sub-Gaussian random variables for solving some problems in compressive sensing of sparse signals.

## 1. INTRODUCTION AND PRELIMINARY RESULTS

In the statistical analysis for efficient detection of signal components when missing data samples are present developed by LJ. Stanković, S. Stanković and M. Amin in [25] (cf. [26] and [23]), a crucial role plays a class of complex-valued random variables denoted in [10] as  $X_l(m, N)$  ( $0 \leq l \leq N - 1$ ,  $1 \leq m \leq N$ ). Furthermore, in [12] (also see [13]) it was generalized the random variable  $X_l(m, N)$ . Motivated by these random variables and the known fact that Bernoulli probability-type models, involving Bernoulli random variable, were often used in the study on some compressive sensing-type problems (see, e.g., the famous paper [2] by Candès, Romberg and Tao and [16]), here we define and study complex-valued random variables  $\tilde{X}_l(m, N)$  ( $0 \leq l \leq N - 1$ ,  $1 \leq m \leq N$ ). For more information on the development of compressive sensing (also known as *compressed sensing*, *compressive sampling*, or *sparse recovery*), see [5], [6], [22, Chapter 10] and [24]. For an excellent survey on this topic with applications and related references, see [27] (also see [18]).

**Definition 1.1.** Let  $N$ ,  $l$  and  $m$  be nonnegative integers such that  $0 \leq l \leq N - 1$  and  $1 \leq m \leq N$ . Let  $B_n$  ( $n = 1, \dots, N$ ) be a sequence of independent identically distributed *Bernoulli random variables* (binomial distributions) taking only the values

---

2010 *Mathematics Subject Classification.* 60C05, 94A12, 11A07, 05A10.

*Key words and phrases.* Compressive sensing, Complex-valued discrete random variable, Bernoulli random variable, Sub-Gaussian random variable, Sub-Gaussian norm, Orlicz norm, McDiarmid's inequality.

0 and 1 with probability 0 and  $m/N$ , respectively, i.e.,

$$(1) \quad B_n = \begin{cases} 0 & \text{with probability } 1 - \frac{m}{N} \\ 1 & \text{with probability } \frac{m}{N}. \end{cases}$$

Then the discrete random variable  $\tilde{X}_l(m, N) = \tilde{X}_l(m, N)$  is defined as a sum

$$(2) \quad \tilde{X}_l(m, N) = \sum_{n=1}^N \exp\left(-\frac{2jnl\pi}{N}\right) B_n.$$

We see from (2) that the real and imaginary part of  $\tilde{X}_l(m, N)$  are real-valued random variables  $\tilde{U}_l(m, N)$  and  $\tilde{V}_l(m, N)$ , respectively defined by

$$(3) \quad \tilde{U}_l(m, N) = \sum_{n=1}^N \cos\left(\frac{2jnl\pi}{N}\right) B_n$$

and

$$(4) \quad \tilde{V}_l(m, N) = -\sum_{n=1}^N \sin\left(\frac{2jnl\pi}{N}\right) B_n.$$

From Definition 1.1 it follows that the range of the random variable  $\tilde{X}_l(m, N)$  consists of all possible  $2^N - 1$  sums of elements of (multi)set  $\{e^{-j2nl\pi/N} : n = 1, 2, \dots, N\}$ .

Notice that for  $l = 0$   $\tilde{X}_l(m, N)$  becomes

$$\tilde{X}_0(m, N) = \sum_{k=1}^N B_k \sim B\left(N, \frac{m}{N}\right),$$

where  $B(N, m/N)$  is the binomial distribution with parameters  $N$  and  $p = m/N$  and the probability mass function given by

$$\text{Prob}\left(B\left(N, \frac{m}{N}\right) = k\right) = \binom{N}{k} \left(\frac{m}{N}\right)^k \left(1 - \frac{m}{N}\right)^{N-k}.$$

Notice that by *De Moivre-Laplace central limit theorem*, the distribution of  $B(N, m/N)$  is close to that of normal random variable  $\mathcal{N}\left(m, \sqrt{m(N-m)/N}\right)$  for sufficiently large  $N$ .

If  $N = 2l$ , where  $l \geq 1$  is an integer, then  $\tilde{X}_l(m, N)$  becomes

$$\tilde{X}_l(m, 2l) = \sum_{k=1}^{2l} (-1)^k B_k = \sum_{k=1}^l B_{2k} - \sum_{k=1}^l B_{2k-1} \sim B'\left(l, \frac{m}{2l}\right) - B''\left(l, \frac{m}{2l}\right),$$

where  $B'(2l, m/(2l))$  and  $B''(2l, m/(2l))$  are independent identically distributed binomial distributions with parameters  $l$  and  $p = m/N$  (i.e., two independent copies of  $B(2l, m/(2l))$ ).

**Remark 1.2.** Set  $\Sigma = \{n : 1 \leq n \leq N \text{ and } B_n = 1\}$ , where  $B_n$  is a random variable defined by (1). Then the cardinality  $|\Sigma|$  of  $\Sigma$  is also random, following a binomial distribution whose expected value is equal to

$$\mathbb{E}[|\Sigma|] = N \cdot \frac{m}{N} = m.$$

We point out also that for large values of  $N$  and  $m \ll N$  there holds  $|\Sigma|/m \approx 1$  with high probability. In other words, for such values of  $N$  and  $m$ , the number of observed frequencies is  $\approx m$  with high probability.

## 2. THE RESULTS

We start with the following result.

**Proposition 2.1.** *Let  $N$ ,  $l$  and  $m$  be positive integers such that  $l \leq N - 1$  and  $1 \leq m \leq N$ . Let  $\tilde{X}_l(m, N) = \tilde{U}_l(m, N) + j\tilde{V}_l(m, N)$  be the discrete random variable from Definition 1.1, where the random variables  $\tilde{U}_l(m, N)$  and  $\tilde{V}_l(m, N)$  are the real and imaginary part of  $\tilde{X}_l(m, N)$ , respectively. Then the expected values of  $\tilde{X}_l(m, N)$ ,  $\tilde{U}_l(m, N)$  and  $\tilde{V}_l(m, N)$  are respectively given by*

$$(5) \quad \mathbb{E}[\tilde{X}_l(m, N)] = \mathbb{E}[\tilde{U}_l(m, N)] = \mathbb{E}[\tilde{V}_l(m, N)] = 0.$$

*If in addition we suppose that  $N \neq 2l$ , then for the variance of  $\tilde{X}_l(m, N)$ ,  $\tilde{U}_l(m, N)$  and  $\tilde{V}_l(m, N)$ , respectively there holds*

$$(6) \quad \text{Var}[\tilde{X}_l(m, N)] = \frac{m(N - m)}{N},$$

and

$$(7) \quad \text{Var}[\tilde{U}_l(m, N)] = \text{Var}[\tilde{V}_l(m, N)] = \frac{m(N - m)}{2N}.$$

As an immediate consequence of equalities (5) and (7) of Proposition 2.1 and the general formula  $\text{Var}[X] = \mathbb{E}[X^2] - (\mathbb{E}[X])^2$  for every real-valued random variable  $X$ , we obtain the following result.

**Corollary 2.2.** *Under the assumptions and notations of Proposition 2.1 and the additional assumption that  $N \neq 2l$  there holds*

$$(8) \quad \mathbb{E}[(\tilde{U}_l(m, N))^2] = \mathbb{E}[(\tilde{V}_l(m, N))^2] = \frac{m(N - m)}{2N}.$$

Finally, we will prove the following main result concerning some probability estimates for the random variable  $\tilde{X}_l(m, N)$  and its real and imaginary parts.

**Theorem 2.3.** *Let  $N$ ,  $l$  and  $m$  be positive integers such that  $l \leq N - 1$ ,  $N \neq 2l$  and  $1 \leq m \leq N$ . Then under notations of Proposition 2.1, the following probability estimates are satisfied for each nonnegative real number  $t$  :*

- (i)  $\text{Prob} \left( |\tilde{U}_l(m, N)| \geq t \right) \leq 2 \exp \left( -\frac{4t^2}{N} \right);$
- (ii)  $\text{Prob} \left( |\tilde{V}_l(m, N)| \geq t \right) \leq 2 \exp \left( -\frac{4t^2}{N} \right);$
- (iii)  $\text{Prob} \left( \left| |\tilde{X}_l(m, N)| - \mathbb{E}[|\tilde{X}_l(m, N)|] \right| \geq t \right) \leq 2 \exp \left( -\frac{2t^2}{N} \right).$

**Remark 2.4.** Let us observe that the expressions on the right hand side of inequalities (i), (ii) and (iii) of Theorem 2.3 does not depend on  $m$  (or equivalently, they does not depend on parameter  $p := m/N$  of binomial distributions  $B_1, B_2, \dots, B_N$  which occur in the linear expressions for random variables  $\tilde{U}_l(m, N)$ ,  $\tilde{V}_l(m, N)$  and  $\tilde{X}_l(m, N)$ ). Notice that by using Theorem 3.4.6 of [19], it can be proved the following estimate

for the variables  $\tilde{U}_l(m, N)$  and  $\tilde{V}_l(m, N)$  (denoted below as  $X$  with  $p = m/N$ ) under condition that  $m < N/2$ , i.e.,  $p < 1/2$ :

$$(9) \quad \text{Prob}(|X| \geq t) \leq \exp\left(-\frac{\ln\left(\frac{1-p}{p}\right)t^2}{N(1-2p)}\right) \quad \text{for each } t \geq 0.$$

In the limit as  $p \rightarrow 1/2$ , the right hand side of the above equality becomes  $\exp(-2t^2/N)$ . Therefore, the estimates (i) and (ii) of Theorem 2.3 together with the estimate (9) immediately yield the following improvement of (i) and (ii).

**Theorem 2.5.** *Under the assumptions and notations of Theorem 2.1, if  $m < N/2$  then the following probability estimate holds for the variables  $\tilde{U}_l(m, N)$  and  $\tilde{V}_l(m, N)$  (denoted below as  $X$ ):*

$$(10) \quad \text{Prob}(|X| \geq t) \leq \exp\left(-\max\left\{\frac{4t^2}{N} - \ln 2, \frac{t^2 \ln \frac{N-m}{m}}{N-2m}\right\}\right) \quad \text{for each } t \geq 0.$$

It is easy to see that if  $m = o(N)$  as  $N \rightarrow \infty$ , then the maximum on the right hand side of (10) is attained at the second expression for all sufficiently large values  $N$  and for each  $t \geq 0$ . For example, if  $m = \lfloor \sqrt{N} \rfloor$ , then the mentioned maximum is attained at the second expression for each  $N \geq 2304$ , i.e., for each  $m \geq \lfloor \sqrt{2304} \rfloor = 48$ . A computation shows that the mentioned maximum is attained at the first expression for every pair  $(N, m)$  with  $N = \lfloor cm \rfloor$ , where  $c$  is any real number in the interval  $(2, 47]$ .

Let us recall that a real-valued random variable  $X$  is *sub-Gaussian* if its distribution is dominated by a normal distribution. Precisely, a real-valued random variable  $X$  is sub-Gaussian if there holds

$$\text{Prob}(|X| > t) \leq \exp\left(1 - \frac{t^2}{C^2}\right) \quad \text{for all } t \geq 0,$$

where  $C > 0$  is a real constant that does not depend on  $t$ .

Sub-Gaussian random variables are introduced by Kahane [7]. Notice that normal and all bounded random variables are sub-Gaussian, while exponential random variables are not. A systematic introduction into sub-Gaussian random variables can be found in [30, Lemma 5.5 in Section 5.2.3 and Subsection 5.2.5]; here we briefly mention the basic definitions. Notice that the *Restricted Isometry Property* (RIP) introduced in [3] holds with high probability for any matrix generated by a sub-Gaussian random variable (also see [4] and [21]). In particular, RIP of Bernoulli sensing matrices is studied in [20].

Let us recall that (cf. [31, Definitions 2.5.6 and Example 2.7.13]) the *sub-Gaussian norm*  $\|\cdot\|_{\psi_2}$  which is for the sub-Gaussian real-valued random variable  $X$  is defined as

$$(11) \quad \|X\|_{\psi_2} = \inf\{K > 0 : \mathbb{E}\left[\exp\left(\frac{X^2}{K^2}\right)\right] \leq 2\}.$$

Notice that the sub-Gaussian norm  $\|\cdot\|_{\psi_2}$  given by (11) is a particular case of the *Orlicz norm* with the *Orlicz function*  $\psi_2(x) = \exp(x^2) - 1$  (see [11, Section 2]). For more information on the Orlicz functions and the associated topological vector spaces,

see [17] (also see [14, Chapter 7] and [15]). In view of the above mentioned facts, a random variable  $X$  is sub-Gaussian if and only if

$$\mathbb{E} \left[ \exp \left( \frac{X^2}{\psi} \right) \right] \leq 2$$

for some real constant  $\psi > 0$ . Hence, any bounded real-valued random variable  $X$  is sub-Gaussian, and clearly, there holds

$$(12) \quad \|X\|_{\psi_2} \leq \frac{1}{\sqrt{\ln 2}} \|X\|_{\infty} \approx 1.20112 \|X\|_{\infty},$$

where  $\|\cdot\|_{\infty}$  is the usual *supremum norm*. Moreover, if  $X$  is a centered normal random variable with variance  $\sigma^2$ , then  $X$  is sub-Gaussian with  $\|X\|_{\psi_2} \leq C\sigma$ , where  $C$  is an absolute constant [30, Subsection 5.2.4].

Another definition of the sub-Gaussian norm  $\|X\|'_{\psi_2}$  for the sub-Gaussian random variable  $X$  was given in [30, Definition 5.7] as

$$\|X\|'_{\psi_2} = \sup_{p \geq 1} (p^{-1/2} (E[|X|^p])^{1/p}).$$

Obviously, there holds

$$\|X\|'_{\psi_2} \leq \|X\|_{\infty}.$$

Since  $\tilde{U}_l(m, N)$  and  $\tilde{V}_l(m, N)$  are bounded random variable with  $\mathbb{E}[\tilde{U}_l(m, N)] = \mathbb{E}[\tilde{V}_l(m, N)] = 0$ , the first assertion of the following result is true.

**Proposition 2.6.** *Let  $N, l$  and  $m$  be positive integers such that  $l \leq N - 1$  and  $1 \leq m \leq N$ . Then  $\tilde{U}_l(m, N)$  and  $\tilde{V}_l(m, N)$  are centered sub-gaussian random variables. Moreover, for any positive real number  $K$  such that  $K > \sqrt{N}/2$  there holds*

$$\mathbb{E} \left[ \frac{(\tilde{U}_l(m, N))^2}{K^2} \right] \leq 1 + \frac{8eNK^2}{(4K^2 - N)^2}$$

and

$$\mathbb{E} \left[ \frac{(\tilde{V}_l(m, N))^2}{K^2} \right] \leq 1 + \frac{8eNK^2}{(4K^2 - N)^2}.$$

Taking into account the definition of sub-Gaussian norm  $\|\cdot\|_{\psi_2}$  given by (11), as an immediate consequence of Proposition 2.6, we obtain the following result.

**Corollary 2.7.** *Let  $N, l$  and  $m$  be positive integers such that  $l \leq N - 1$  and  $1 \leq m \leq N$ . Then*

$$\|\tilde{U}_l(m, N)\|_{\psi_2} \leq \frac{\sqrt{N}(\sqrt{2e} + \sqrt{2e+4})}{4} \approx 1.35809\sqrt{N}.$$

and

$$\|\tilde{V}_l(m, N)\|_{\psi_2} \leq \frac{\sqrt{N}(\sqrt{2e} + \sqrt{2e+4})}{4} \approx 1.35809\sqrt{N}$$

Clearly, from (3) and (4) it follows that  $\|\tilde{U}_l(m, N)\|_{\infty} \leq N$  and  $\|\tilde{V}_l(m, N)\|_{\infty} \leq N$  there holds. This together with the inequality (12) yields

$$\|\tilde{U}_l(m, N)\|_{\psi_2} \leq \frac{N}{\sqrt{\ln 2}} \quad \text{and} \quad \|\tilde{V}_l(m, N)\|_{\psi_2} \leq \frac{N}{\sqrt{\ln 2}}.$$

Notice that the estimates from Corollary 2.7 probably are not sharp, but they are much better than these given by the above two inequalities.

**Remark 2.8.** Put

$$Q(x) := \frac{1}{\sqrt{2\pi}} \int_x^\infty \exp\left(-\frac{t^2}{2}\right) dt \quad \text{for all } x \in \mathbb{R}^+ \setminus \{0\},$$

i.e.,  $Q(x)$  is the complementary standard Gaussian cumulative density function (also known as the  $Q$ -function). Then for  $Q(x)$  the following exponential upper and lower bounds holds (see, e.g., [29, Section 3.3]):

$$\frac{1}{\sqrt{2\pi}} \cdot \frac{x}{1+x^2} \cdot \exp\left(-\frac{x^2}{2}\right) < Q(x) < \frac{1}{\sqrt{2\pi}x} \cdot \exp\left(-\frac{x^2}{2}\right) \quad \text{for all } x \in \mathbb{R}.$$

The left hand side of the above double inequality allows us to replace the right hand side of inequalities (i) and (ii) of Theorem 2.3 with the expression involving the function  $Q\left(2t\sqrt{2}/\sqrt{N}\right)$ ; namely, if  $X = |\tilde{U}_l(m, N)|$  or  $Y = |\tilde{V}_l(m, N)|$ , then

$$\text{Prob}(X \geq t) \leq \frac{(N + 8t^2)\sqrt{\pi}}{t\sqrt{N}} Q\left(\frac{2t\sqrt{2}}{\sqrt{N}}\right).$$

**Remark 2.9.** From (6) and [10, (19) of Theorem 2.4] it follows that

$$(13) \quad \frac{\text{Var}[X_l(m, N)]}{\text{Var}[\tilde{X}_l(m, N)]} = \frac{N}{N-1}.$$

For related discussion on the proprtion (13), see [11, Remark 1.2].

Proofs of Proposition 2.1, Theorem 2.3 and Proposition 2.6 are given in Section 3.

### 3. PROOFS OF THE RESULTS

For the proof of Proposition 2.1, we will need the following result.

**Lemma 3.1.** ([10, (38) of Lemma 4.1]). *Let  $N$  and  $l$  be positive integers such that  $l \leq N - 1$  and  $N \neq 2l$ , then*

$$(14) \quad \sum_{k=1}^N \cos \frac{4kl\pi}{N} = \sum_{k=1}^N \sin \frac{4kl\pi}{N} = 0.$$

*Proof of Proposition 2.1.* By the equality (2) of Definition 1.1 and the obvious fact that  $\mathbb{E}[B_n] = m/N$  for each  $n = 1, \dots, N$ , and using the linearity property of the expectation, we find that for each  $l \neq 0$  with  $l \leq N - 1$ ,

$$\begin{aligned} \mathbb{E}\left[\tilde{X}_l(m, N)\right] &= \mathbb{E}\left[\sum_{n=1}^N B_n \exp\left(-\frac{2jnl\pi}{N}\right)\right] = \sum_{n=1}^N \exp\left(-\frac{2jnl\pi}{N}\right) \mathbb{E}[B_n] \\ &= \frac{m}{N} \sum_{n=1}^N \exp\left(-\frac{2jnl\pi}{N}\right) = \frac{m}{N} \cdot \sum_{n=0}^{N-1} \exp\left(-\frac{2jnl\pi}{N}\right) \\ &= \frac{m}{N} \cdot \exp\left(-\frac{2jl\pi}{N}\right) \cdot \frac{\exp(-2jl\pi) - 1}{\exp\left(-\frac{2jl\pi}{N}\right) - 1} = 0. \end{aligned}$$

The above equality and the expression  $\mathbb{E}[\tilde{X}_l(m, N)] = \mathbb{E}[\tilde{U}_l(m, N)] + j\mathbb{E}[\tilde{V}_l(m, N)]$  imply all the equalities in (5).

Further, using the additive property for the variance of a finite sum of *uncorrelated random variables* (and hence, for a finite sum of independent identically distributed random variables), for all  $l = 0, 1, \dots, N - 1$  we obtain

$$\begin{aligned} \text{Var}[\tilde{X}_l(m, N)] &= \sum_{n=1}^N \text{Var} \left[ \exp \left( -\frac{2jnl\pi}{N} \right) B_n \right] \\ &= \sum_{n=1}^N \left( \left| \exp \left( -\frac{2jnl\pi}{N} \right) \right|^2 \text{Var}[B_n] \right) \\ &= \sum_{n=1}^N \text{Var}[B_n] = N \cdot \text{Var}[B_1] = N \cdot (\mathbb{E}[B_1^2] - (\mathbb{E}[B_1])^2) \\ &= N \cdot \left( \frac{m}{N} - \left( \frac{m}{N} \right)^2 \right) = \frac{m(N-m)}{N}, \end{aligned}$$

which implies the first expression in (5). Similarly, by using the expansion

$$\tilde{U}_l(m, N) = \sum_{n=1}^N \cos \frac{2nl\pi}{N} B_n,$$

the trigonometric identity  $\cos^2 \alpha = (1 + \cos 2\alpha)/2$  and the first identity of (14) from Lemma 3.1, we obtain

$$\begin{aligned} \text{Var}[\tilde{U}_l(m, N)] &= \sum_{n=1}^N \text{Var} \left[ \cos \frac{2nl\pi}{N} B_n \right] = \sum_{n=1}^N \cos^2 \frac{2nl\pi}{N} \text{Var}[B_n] \\ &= \sum_{n=1}^N \cos^2 \frac{2nl\pi}{N} \left( \frac{m}{N} - \left( \frac{m}{N} \right)^2 \right) = \frac{m(N-m)}{N^2} \sum_{n=1}^N \cos^2 \frac{2nl\pi}{N} \\ &= \frac{m(N-m)}{N^2} \sum_{n=1}^N \frac{1 + \cos \frac{4nl\pi}{N}}{2} = \frac{m(N-m)}{N^2} \left( \frac{N}{2} + \frac{1}{2} \sum_{n=1}^N \cos \frac{4nl\pi}{N} \right) \\ &= \frac{m(N-m)}{2N}, \end{aligned}$$

whence it follows the equality (6).

Proceeding analogously as above, by using the expansion

$$\tilde{V}_l(m, N) = - \sum_{n=1}^N \sin \left( \frac{2nl\pi}{N} \right) B_n,$$

the trigonometric identity  $\sin^2 \alpha = (1 - \cos 2\alpha)/2$  and the second identity of (14) from Lemma 3.1, we obtain the expression (7). This completes proof of Proposition 2.1.  $\square$

Proof of Theorem 2.3 is based on the following *McDiarmid's inequality*, also known as the *bounded-difference inequality* given in [8, Theorem 3.1] (also see [9]).

**Theorem 3.2.** (*McDiarmid's inequality*). *Let  $X_1, \dots, X_N$  be independent (not necessarily identically distributed) real-valued random variables all taking values in a measurable set  $\chi$ . Further, let  $f : \chi^N \mapsto \mathbb{R}$  be a measurable function of  $X_1, \dots, X_N$  (random variable) that satisfies the inequality (the bounded difference assumption)*

$$(15) \quad \left| f(x_1, \dots, x_k, \dots, x_N) - f(x_1, \dots, x'_k, \dots, x_N) \right| \leq c_k$$

for all  $x_1, \dots, x_N, x'_k \in \chi$  with any fixed  $k \in \{1, \dots, N\}$ , where  $c_k$  ( $k = 1, \dots, N$ ) are arbitrary nonnegative real constants. Consider a random variable  $X = f(X_1, \dots, X_N)$ . Then for every real number  $t \geq 0$ ,

$$(16) \quad \text{Prob}(|X - \mathbb{E}[X]| \geq t) \leq 2 \exp\left(-\frac{2t^2}{\sum_{k=1}^N c_k^2}\right).$$

*Proof of Theorem 2.3.* First notice that by definition,

$$\tilde{U}_l(m, N) = \sum_{k=1}^N \cos \frac{2kl\pi}{N} X_k,$$

where  $X_k = B_k$  ( $k = 1, \dots, N$ ) are independent *Bernoulli 0-1 random variables*. Accordingly, if we define the real-valued function  $f : \{0, 1\}^N \mapsto \mathbb{R}$  defined on two-points set  $\{0, 1\}$  as

$$(17) \quad f(x_1, \dots, x_N) = \sum_{k=1}^N \cos \frac{2kl\pi}{N} x_k, \quad \text{for all } x_1, \dots, x_N \in \{0, 1\},$$

then we can write

$$\tilde{U}_l(m, N) = f(X_1, \dots, X_N).$$

Substituting the expression for the function  $f$  given by (17) into the inequality (15), it becomes

$$(18) \quad \left| \cos \frac{2kl\pi}{N} x_k - \cos \frac{2kl\pi}{N} x'_k \right| = |x_k - x'_k| \cdot \left| \cos \frac{2kl\pi}{N} \right| \leq c_k,$$

for all  $x_k, x'_k \in \{0, 1\}$  with  $k = 1, \dots, N$ .

Clearly, for any fixed  $k \in \{1, \dots, N\}$  the inequality (18) holds with  $c_k = \left| \cos \frac{2kl\pi}{N} \right|$ .

Then by using the identity  $\cos^2 \alpha = (1 + \cos 2\alpha)/2$  and the first identity of (14) from Lemma 3.1, we obtain

$$\begin{aligned} \sum_{k=1}^N c_k^2 &= \sum_{k=1}^N \cos^2 \frac{2kl\pi}{N} = \sum_{k=1}^N \frac{1 + \cos \frac{4kl\pi}{N}}{2} = \\ &= \frac{N}{2} + \frac{1}{2} \sum_{k=1}^N \cos \frac{4kl\pi}{N} = \frac{N}{2}. \end{aligned}$$

Substituting the above expression  $\sum_{k=1}^N c_k^2 = N/2$  into (16), we obtain that for every  $t \geq 0$ ,

$$\text{Prob}\left(|\tilde{U}_l(m, N)| \geq t\right) \leq 2 \exp\left(-\frac{4t^2}{N}\right).$$

This proves the inequality (i) of Theorem 2.3.



As by the identity  $\sin^2 \alpha = (1 - \cos 2\alpha)/2$  and the second identity of (14) from Lemma 3.1,

$$\begin{aligned} \sum_{k=1}^N \sin^2 \frac{2kl\pi}{N} &= \sum_{k=1}^N \frac{1 - \cos \frac{4kl\pi}{N}}{2} = \\ &= \frac{N}{2} - \frac{1}{2} \sum_{k=1}^N \cos \frac{4kl\pi}{N} = \frac{N}{2} \end{aligned}$$

and

$$\tilde{V}_l(m, N) = \sum_{k=1}^N \sin \frac{2kl\pi}{N} B_k,$$

proof of the inequality (ii) of Theorem 2.3 can be deduced in the same manner as that of (i), and hence, may be omitted.

Finally, in order to prove the inequality (iii) of Theorem 2.3, notice that  $|\tilde{X}_l(m, N)|$  is a real-valued random variable such that

$$(19) \quad \left| \tilde{X}_l(m, N) \right| = \left| \sum_{k=1}^N \exp \left( -\frac{2jkl\pi}{N} \right) B_k \right|.$$

Similarly as above, take  $X_k = B_k$  ( $k = 1, \dots, N$ ) and define the real-valued function  $g : \{0, 1\}^N \mapsto \mathbb{R}$  as

$$(20) \quad g(x_1, \dots, x_N) = \left| \sum_{k=1}^N \exp \left( -\frac{2jkl\pi}{N} \right) x_k \right|, \quad \text{for all } x_1, \dots, x_N \in \{0, 1\}.$$

Substituting the expression for the function  $f$  given by (20) into the inequality (15), it becomes

$$(21) \quad \begin{aligned} &\left| \exp \left( -\frac{2jkl\pi}{N} \right) x_k - \exp \left( -\frac{2jkl\pi}{N} \right) x'_k \right| \\ &= |x_k - x'_k| \cdot \left| \exp \left( -\frac{2jkl\pi}{N} \right) \right| = |x_k - x'_k| \leq c_k, \end{aligned}$$

for all  $x_k, x'_k \in \{0, 1\}$  with  $k = 1, \dots, N$ . Obviously, for any fixed  $k \in \{1, \dots, N\}$  the inequality (21) holds with  $c_k = 1$ . Finally, substituting the value  $\sum_{k=1}^N c_k^2 = N$  into (16), we obtain that for every  $t \geq 0$ ,

$$\text{Prob} \left( \left| |\tilde{X}_l(m, N)| - \mathbb{E}[|\tilde{X}_l(m, N)|] \right| \geq t \right) \leq 2 \exp \left( -\frac{2t^2}{N} \right).$$

Therefore, the estimate (iii) holds and proof of the theorem is completed.  $\square$

*Proof of Proposition 2.6.* Since the estimates (i) and (ii) of Theorem 2.3 are the same, it is sufficient to prove the first inequality of Proposition 2.6. We follow the part of proof of Proposition 2.5.2 in [31, Subsection 2.5.1]. It is known (see, e.g., [28, Appendix A, p. 434]) that for any nonnegative real-valued random variable  $X$  with probability distribution function  $F(x) := \text{Prob}(X < x)$  ( $x \in \mathbb{R}$ ),

$$\mathbb{E}[X] = \int_0^\infty (1 - F(v)) dv,$$

which can be written as

$$\mathbb{E}[X] = \int_0^\infty \text{Prob}(X \geq v) dv.$$

Applying the above equality for  $X = \left| \tilde{U}_l(m, N) \right|^n$ , after the change of variables  $v = t^{2n}$ ,  $dv = 2nt^{2n-1}dt$ , using the estimate (i) of Theorem 2.3 and the well known inequality  $\Gamma(x) \leq x^x e^{1-x}$  ( $x \geq 1$ , where  $\Gamma(x) = \int_0^\infty s^{x-1} e^{-s} ds$  for  $x > 0$ , is the *complete gamma function*), for every integer  $n \geq 1$  we obtain

$$\begin{aligned} \mathbb{E} \left[ \tilde{U}_l(m, N)^{2n} \right] &= \int_0^\infty \text{Prob} \left( \left| \tilde{U}_l(m, N) \right| \geq t \right) 2nt^{2n-1} dt \leq 4n \int_0^\infty \exp \left( -\frac{4t^2}{N} \right) t^{2n-1} dt \\ &= \left( \text{the change of variables } \frac{4t^2}{N} = s, dt = \frac{\sqrt{N} ds}{4\sqrt{s}} \right) \\ &= \frac{nN^n}{2^{2n-1}} \int_0^\infty e^{-s} s^{n-1} ds = \frac{nN^n}{2^{2n-1}} \Gamma(n) \leq \frac{nN^n}{2^{2n-1}} \cdot n^n e^{1-n} \\ &= \frac{n^{n+1} N^n}{2^{2n-1} e^{n-1}}. \end{aligned}$$

By using the Taylor expansion of  $\tilde{U}_l(m, N)^2/K^2$  ( $K$  is a positive real constant), the linearity of the expected value, the above inequality and the well known inequality  $n! \geq (n/e)^n$  ( $n \in \mathbb{N}$ ), we find that

$$\begin{aligned} \mathbb{E} \left[ \frac{\tilde{U}_l(m, N)^2}{K^2} \right] &= 1 + \sum_{n=1}^\infty \frac{\mathbb{E} \left[ \tilde{U}_l(m, N)^{2n} \right]}{K^{2n} n!} \\ (22) \quad &\leq 1 + \sum_{n=1}^\infty \frac{n^{n+1} N^n}{2^{2n-1} e^{n-1} K^{2n} n!} \leq 1 + \sum_{n=1}^\infty \frac{n^{n+1} N^n}{2^{2n-1} e^{n-1} K^{2n}} \cdot e^n n^n \\ &= 1 + 2e \sum_{n=1}^\infty n \left( \frac{N}{4K^2} \right)^n. \end{aligned}$$

Since for  $|x| < 1$  we have

$$\begin{aligned} \sum_{n=1}^\infty nx^n &= x \sum_{n=1}^\infty nx^{n-1} = \frac{d}{dx} \sum_{n=1}^\infty x^n = \frac{d}{dx} \frac{x}{1-x} \\ &= \frac{x}{(1-x)^2}, \end{aligned}$$

substituting the above identity with  $N/(4K^2) = x$  (which is by the assumption  $< 1$ ) into (22), we get

$$\mathbb{E} \left[ \frac{\tilde{U}_l(m, N)^2}{K^2} \right] \leq 1 + \frac{8eNK^2}{(4K^2 - N)^2},$$

as asserted. □

## REFERENCES

- [1] E.J. Candès, The restricted isometry property and its implications for compressed sensing, *Comptes Rendus de l'Académie des Sciences-Série I-Mathematics* **346**, No. 9-10 (2008), 589–592.
- [2] E.J. Candès, J. Romberg and T. Tao, Robust uncertainty principles: Exact signal reconstruction from highly incomplete frequency information, *IEEE Transactions on Information Theory* **52**, No. 2 (2006), 489–509.
- [3] E.J. Candès and T. Tao, Decoding by linear programming, *IEEE Transactions on Information Theory* **51**, No. 12 (2005), 4203–4215.
- [4] E.J. Candès and T. Tao, Near-optimal signal recovery from random projections: Universal encoding strategies?, *IEEE Transactions on Information Theory* **52**, No. 12 (2006), 5406–5425.
- [5] D.L. Donoho, Compressed sensing, *IEEE Transactions on Information Theory* **52**, No. 4 (2006), 1289–1306.
- [6] S. Foucart and H. Rauhut, *A Mathematical Introduction to Compressive Sensing*, Springer, 2013.
- [7] J.P. Kahane, Propriétés locales des fonctions à séries de Fourier aléatoires, *Studia Mathematica* **19** (1960), 1–25 (French).
- [8] C. McDiarmid, Centering sequences with bounded differences, *Combinatorics, Probability and Computing* **6**, no. 1 (1997), 79–86.
- [9] C. McDiarmid, On the method of bounded differences. In *Surveys in Combinatorics*, vol. 141, pages 148–188, Cambridge University Press, 1989.
- [10] R. Meštrović, On some discrete random variables arising from recent study on statistical analysis on compressive sensing, available at [arXiv:1803.02260v1](https://arxiv.org/abs/1803.02260v1) [math.ST], 2018, 22 pages; also available in “Preprints” at <https://sites.google.com/site/romeomestrovic/>.
- [11] R. Meštrović, A note on some sub-Gaussian random variables, available at [arXiv:1803.04521v1](https://arxiv.org/abs/1803.04521v1) [math.PR], 2018, 18 pages; also available in “Preprints” at <https://sites.google.com/site/romeomestrovic/>.
- [12] R. Meštrović, Generalization of some random variables involving in certain compressive sensing problems, in preparation.
- [13] R. Meštrović, Three combinatorial identities with applications, in preparation.
- [14] R. Meštrović and Ž. Pavićević, *Privalov spaces of the unit disk* (Research monograph), University of Montenegro, Podgorica, 2009.
- [15] R. Meštrović, Ž. Pavićević and N. Labudović, Remarks on generalized Hardy algebras, *Mathematica Montisnigri* **11** (1999), 25–42, available in “Publications” at <https://sites.google.com/site/romeomestrovic/>.
- [16] D.G. Mixon, W.U. Bajwa and R. Calderbank, Frame Coherence and Sparse Signal Processing, available at [arXiv:1105.4279v1](https://arxiv.org/abs/1105.4279v1) [math.FA], 2011.
- [17] J. Musielak, *Orlicz spaces and modular spaces*, Lecture Notes in Mathematics, **1034**, Springer-Verlag, 1983.
- [18] I. Orović, V. Papić, C. Ioana, X. Liu and S. Stanković, Compressive sensing in Signal processing: algorithms and transform domain formulations, *Mathematical Problems in Engineering* (Special Issue “Algorithms for Compressive Sensing Signal Reconstruction with Applications”) vol. **2016** (2016), Article ID 7616393, 16 pages.
- [19] M. Raginsky and I. Sason, *Concentration of Measure Inequalities in Information Theory, Communications, and Coding*, Foundations and Trends/copyright in Communications and Information Theory, vol. **10**, no. 1–2, pp. 1–246, Oct. 2013. doi: 10.1561/01000000064.
- [20] H. Rauhut, Compressive sensing and structured random matrices, *Theoretical foundations and numerical methods for sparse recovery* **9** (2010), 1–92.
- [21] M. Rudelson and R. Vershynin, On sparse reconstruction from Fourier and Gaussian measurements, *Communications on Pure and Applied Mathematics* **61**, No. 8 (2008), 1025–1045.
- [22] L.J. Stanković, *Digital Signal Processing with Selected Topics*, CreateSpace Independent Publishing Platform, An Amazon.com Company, 2015.

- [23] LJ. Stanković, M. Daković, I. Stanković and S. Vujović, On the errors in randomly sampled non-sparse signals reconstructed with a sparsity assumption, *IEEE Geoscience and Remote Sensing Letters* **14**, No. 12 (2017), 2453–2456.
- [24] LJ. Stanković, M. Daković and T. Thayaparan, *Time-Frequency Signal Analysis with Applications*, Artech House, 2013.
- [25] LJ. Stanković, S. Stanković and M. Amin, Missing samples analysis in signals for applications to L-estimation and compressive sensing, *Signal Processing* **94**, No. 1 (2014), 401–408.
- [26] LJ. Stanković, S. Stanković, I. Orović and M. Amin, Robust time-frequency analysis based on the L-estimation and compressive sensing, *IEEE Signal Processing Letters* **20**, No. 5 (2013), 499–502.\*
- [27] S. Stanković, Compressive sensing, Theory, Algorithms and Applications (invited lecture), *4th Mediterranean Conference on Embedded Computing, MECO-2015*, Budva, Montenegro, 2015; available at [www.tfsa.ac.me/pap/tfsa-001115.pdf](http://www.tfsa.ac.me/pap/tfsa-001115.pdf).
- [28] H.C. Tijms, *A first course in stochastic models*, John Wiley and Sons, 2003.
- [29] S. Verdú, *Multiuser Detection*, Cambridge University Press, 1998.
- [30] R. Vershynin, *Introduction to the non-asymptotic analysis of random matrices*, in *Compressed Sensing: Theory and Applications*, pp. 210–268, edited by Y. Eldar and G. Kutyniok, Eds. Cambridge, Cambridge University Press, 2012.
- [31] R. Vershynin, *High-dimensional probability: An introduction with applications in Data science*, 297 pages, August 24, 2017, available at <https://www.math.uci.edu/~rvershyn/>.

MARITIME FACULTY KOTOR, UNIVERSITY OF MONTENEGRO, 85330 KOTOR, MONTENEGRO  
E-mail address: romeo@ac.me