

(WEAK) INCIDENCE BIALGEBRAS OF MONOIDAL CATEGORIES

ULRICH KRÄHMER AND LUCIA ROTHERAY

ABSTRACT. Incidence coalgebras of categories in the sense of Joni and Rota are studied, specifically cases where a monoidal product on the category turns these into (weak) bialgebras. The overlap with the theory of combinatorial Hopf algebras and that of Hopf quivers is discussed, and examples including trees, skew shapes, Milner's bigraphs and crossed modules are considered.

CONTENTS

1. Introduction	1
2. Definitions and notation	2
3. Bialgebras from monoidal categories	3
3.1. Example: Relations on monoids	4
3.2. Example: Skew shapes and symmetric functions	5
3.3. Example: Rooted forests and the Connes-Kreimer Hopf algebra	6
3.4. Example: A toy model for bigraphs	7
3.5. A notable non-example: Hopf quivers	10
4. Weak Hopf algebras from monoidal categories	12
4.1. Example: Relations on groups	14
4.2. Example: Automorphism 2-groups	14
4.3. Crossed modules	14
References	15

1. INTRODUCTION

Incidence coalgebras of categories, defined in [20], have been studied in several areas, notably Möbius inversion [24, 25] and combinatorial Hopf algebras (see e.g. [11, 16, 17, 26]). The latter notion refers to Hopf algebras with a vector space basis indexed by a family of combinatorial objects (e.g. graphs or integer partitions), but the precise definition varies in the literature. The product and coproduct reflect unions and compositions of these objects, hence the underlying coalgebra is (or is closely related to) an incidence coalgebra of a category \mathcal{C} .

In this case, one might wonder if the multiplication corresponds to a monoidal product on \mathcal{C} . This was explored by several authors, see [14] for a recent account. Here, we characterise two classes of monoidal categories giving rise to pointed respectively weak bialgebras:

Theorem 1. *If a monoidal product on a Möbius category \mathcal{C} has the unique lifting of factorisation property, then its linearisation turns the incidence coalgebra of \mathcal{C} into a pointed bialgebra. This is a Hopf algebra provided that the monoid of objects*

It is our pleasure to thank Gabriella Böhm, Joachim Kock and Michele Sevegnani for discussions and suggestions.

is a group. Similarly, if \mathcal{C} is a locally finite strict 2-group, the monoidal product turns the incidence coalgebra of \mathcal{C} into a weak Hopf algebra.

See the main text for definitions.

A main goal of this paper is to discuss how several well-known examples of Hopf algebras fit into this picture, including the Connes-Kreimer Hopf algebra of rooted trees and symmetric functions. Milner's bigraphs, a combinatorial structure employed in theoretical computer science, are considered as a new example. We discuss how the Hopf algebraic techniques applied to the problem of renormalisation in physics could lead to new approaches to bigraphical systems, e.g. in studying reaction rules, and it is our hope that there are other parallels to be drawn between bigraphical systems and physical ones, for example Dyson-Schwinger type equations for generating sub-bialgebras.

We also show that not all combinatorial Hopf algebras (in the above sense) can be described this way, even when the coalgebra structure is the incidence coalgebra of a category. As an illustrative, and perhaps counterintuitive, example we investigate the Hopf quivers of Cibils and Rosso:

Theorem 2. *Let $Q = (Q_0, Q_1)$ be a Hopf quiver, k a field and $\cdot : kQ \times kQ \rightarrow kQ$ the multiplication in the associated Hopf algebra $k\mathcal{C}$ as defined in [7]. Then \cdot is the linear extension of a monoidal product on Q if and only if $Q_1 = \emptyset$.*

The structure of the paper is as follows:

In Section 2 we recall some basic definitions, including that of an incidence coalgebra of a category, Möbius categories and the unique lifting of factorisations property.

In Section 3 we study monoidal Möbius categories whose monoidal product is an ULF functor. We prove that these define bialgebras, and discuss the examples mentioned above.

In Section 4 we recall the definition of a weak bialgebra, and show that the monoidal product in a 2-group satisfies a weak version of the unique lifting of factorisation property, which leads to the last statement in Theorem 1.

2. DEFINITIONS AND NOTATION

Throughout, all categories are assumed to be small and all monoidal categories to be strict. We use \mathcal{C} to denote both a category and its set of morphisms, and $\mathcal{C}(x, y)$ the subset of morphisms from x to y . We denote the set of objects by $Ob\mathcal{C}$ and the set of identity morphisms in \mathcal{C} by $Id\mathcal{C}$. The identity morphism at $x \in Ob\mathcal{C}$ is written i_x . The monoidal product is denoted $\cdot : \mathcal{C} \times \mathcal{C} \rightarrow \mathcal{C}$.

Definition 1 (Decompositions and length). Given a morphism $f \in \mathcal{C}$ and $n \in \mathbb{N}$, we define the set

$$N_n(f) := \{(a_1, \dots, a_n) \in \mathcal{C}^{\times n} \mid a_1 \circ \dots \circ a_n = f\}$$

of n -decompositions of f , and let $\hat{N}_n(f)$ be the non-degenerate subset, i.e. those decompositions for which no a_i is an identity morphism. Further, we set

$$\hat{N}(f) := \bigcup_{n \in \mathbb{N}} \hat{N}_n(f), \quad \ell(f) := \sup\{n \mid \hat{N}_n(f) \neq \emptyset\}$$

and call $\ell(f)$ the length of f .

Note that by definition, $N_n := \bigcup_{f \in \mathcal{C}} N_n(f)$ is the set of n -simplices in the nerve of the category \mathcal{C} . This leads to the more topological viewpoint of [14]. The first of the following concepts appears under various names in the literature; we follow the terminology of Joni and Rota [20] and Leroux [25], see also [24].

Definition 2 (Locally finite and Möbius categories). A category \mathcal{C} is called locally finite respectively Möbius if $|N_2(f)|$ respectively $|\tilde{N}(f)|$ is finite for every $f \in \mathcal{C}$.

Evidently, a Möbius category does not contain any nontrivial isomorphisms or idempotents.

Lemma 1. \mathcal{C} is Möbius if and only if \mathcal{C} is locally finite and every morphism has finite length.

For the proof see [25] or [24, Proposition 2.8].

Note that the sequence of sets

$$\mathcal{C}_n := \{f \in \mathcal{C} \mid \ell(f) \leq n\}$$

is a filtration of \mathcal{C} which is exhaustive if and only if every morphism has finite length, in particular when \mathcal{C} is Möbius. The set \mathcal{C}_0 contains only identity morphisms, and \mathcal{C} is called strongly one-way if $\text{Id}\mathcal{C} = \mathcal{C}_0$. This also holds if \mathcal{C} is Möbius. The following is due to Joni and Rota [20] – we consider a variation in which the coproduct is multiplied by a nonzero scalar, as this rescaling arises naturally from the compatibility with the associative products discussed below:

Theorem 3 ((Scaled) incidence coalgebra of a category). *Let \mathcal{C} be a category, k a field, $k\mathcal{C} := \text{span}_k \mathcal{C}$ the free k -module on \mathcal{C} and $\lambda \in k$, $\lambda \neq 0$. Then the following defines a coassociative counital coalgebra structure on $k\mathcal{C}$ if and only if \mathcal{C} is locally finite:*

$$\Delta(f) := \frac{1}{\lambda} \sum_{(a,b) \in N_2(f)} a \otimes b, \quad \varepsilon(f) := \begin{cases} \lambda, & f \in \text{Id}\mathcal{C}, \\ 0, & f \notin \text{Id}\mathcal{C} \end{cases}$$

We will combine such coalgebra structures on $k\mathcal{C}$ with the algebra structure defined by a monoidal product on \mathcal{C} :

Lemma 2. *Let $(\mathcal{C}, \cdot, 1)$ be a monoidal category, k a field, and $k\mathcal{C} := \text{span}_k \mathcal{C}$ the free k -module on \mathcal{C} . Then $(k\mathcal{C}, \cdot, i_1)$ defines an associative, unital k -algebra.*

To address compatibility of product and coproduct, we require the following definition:

Definition 3 (Lifting of factorisations property). For $n \in \mathbb{N} \setminus \{0\}$, a functor $F : \mathcal{C} \rightarrow \mathcal{D}$ has the n -to-one lifting of factorisations (nLF) property if the map

$$\begin{aligned} N_2(f) &\rightarrow N_2(Ff) \\ (a, b) &\mapsto (Fa, Fb) \end{aligned}$$

is an n -to-one surjection for all $f \in \mathcal{C}$.

This generalises the unique lifting of factorisations (or ULF) property defined in e.g. [14, 15, 24]. For consistency with the literature we will use the name ULF rather than 1LF.

Lemma 3. *If $F : \mathcal{C} \rightarrow \mathcal{D}$ has the ULF property, then F reflects identities, i.e. $Ff \in \text{Id}\mathcal{D} \Leftrightarrow f \in \text{Id}\mathcal{C}$.*

Proof. $f \in \text{Id}\mathcal{C} \Rightarrow Ff \in \text{Id}\mathcal{D}$ holds as F is a functor. $Ff \in \text{Id}\mathcal{D} \Rightarrow f \in \text{Id}\mathcal{C}$ follows from the injectivity of the map $N_2(f) \rightarrow N_2(Ff)$. \square

3. BIALGEBRAS FROM MONOIDAL CATEGORIES

In this section we consider the following:

Definition 4 (Combinatorial category). A combinatorial category is a monoidal Möbius category $(\mathcal{C}, \cdot, 1)$ whose product $\cdot : \mathcal{C} \times \mathcal{C} \rightarrow \mathcal{C}$ has the ULF property.

Theorem 4. *Let \mathcal{C} be a locally finite category, k a field, and $k\mathcal{C} := \text{span}_k \mathcal{C}$ the free k -module on \mathcal{C} . We understand \otimes to mean the tensor product \otimes_k and $(k\mathcal{C}, \Delta, \varepsilon)$ to be the incidence coalgebra of \mathcal{C} .*

- (1) *If \mathcal{C} is Möbius, the coalgebra $k\mathcal{C}$ is pointed.*
- (2) *If \mathcal{C} is combinatorial, $(k\mathcal{C}, \cdot, i_1, \Delta, \varepsilon)$ defines a k -bialgebra.*
- (3) *If \mathcal{C} is combinatorial, then $k\mathcal{C}$ is a Hopf algebra if and only if the monoid $(\mathcal{C}_0, \cdot, 1)$ formed by the objects is a group.*

Proof. (1) As \mathcal{C} is Möbius, the length filtration is an exhaustive coalgebra filtration, and we apply [29, Lemma 4.1.2].
 (2) The Möbius condition ensures that $\Delta(i_1) = i_1 \otimes i_1$. The ULF property of the monoidal product ensures that Δ and ε are multiplicative (Lemma 3 shows this for ε).
 (3) Application of part (1) and [29, Lemma 7.6.2]. \square

We will now present some examples of combinatorial categories and the bialgebras associated to them as in Theorem 4. These are combinatorial bialgebras in the sense described in the introduction - hence the name combinatorial category.

3.1. Example: Relations on monoids. We begin with a simple class of examples to illustrate the definitions.

Let \preceq be a reflexive and transitive relation on a monoid M for which

$$(1) \quad x \preceq y, z \preceq t \Rightarrow x \cdot z \preceq y \cdot t$$

holds. Then

$$\mathcal{C}_M := \{(x, y) \in M \times M \mid x \preceq y\}$$

is the morphism set of a monoidal category with object set M and (x, y) the (unique) morphism $y \rightarrow x$. The operations \circ, \cdot are given by

$$(x, y) \circ (y, z) := (x, z), \quad (x, y) \cdot (z, t) := (x \cdot z, y \cdot t).$$

The identity morphisms are given by $i_x = (x, x)$. By definition, we have:

- Proposition 1.** (1) *The category \mathcal{C}_M is locally finite if and only if all intervals $[x, y] := \{z \in M \mid x \preceq z \preceq y\}$ are finite.*
 (2) *The category \mathcal{C}_M is Möbius if and only if given $x \preceq y \in M$ there is $\ell(x, y) \in \mathbb{N}$ such that there is no chain $x = z_1 \preceq z_2 \preceq \dots \preceq z_l = y$ with $z_i \neq z_{i+1}$ and $l > \ell(x, y)$.*
 (3) *The monoidal structure is ULF if and only if it defines for all $x \preceq y, z \preceq t \in M$ a bijection $[x, y] \times [z, t] \rightarrow [x \cdot z, y \cdot t]$.*

Note that if \mathcal{C}_M is Möbius, \preceq is a partial ordering.

Remark 1. Let (M, \leq) be a monoid with a partial ordering satisfying (1). On the set \mathcal{C}_M of morphisms define $(x, x) \simeq (1, 1) \forall x \in M$. Taking the quotient by the corresponding bialgebra ideal removes non-unit group-like elements and produces a Hopf algebra which is as close as possible to the original bialgebra.

Various authors, for example [3, 20], consider Hopf algebras of intervals in posets and impose exactly this condition to obtain Hopf algebras rather than bialgebras. Rather than starting with a monoid and imposing a partial order they start with a grading poset and define a product on its intervals (morphisms for us), which induces a monoid structure on the original set, compatible with the grading.

In particular, when \preceq is an equivalence relation, then the interval $[x, y]$ is simply the equivalence class of x (and of y), and (1) demands that the monoid structure descends to the quotient M / \preceq . The category \mathcal{C}_M is locally finite if and only if all

equivalence classes are finite, but it is not Möbius unless \preceq is $=$, as any (x, y) with $x \preceq y$ but $x \neq y$ is a nontrivial isomorphism. As a coalgebra, we have

$$k\mathcal{C}_M \cong \bigoplus_{x \in M/\preceq} M_{|x|}(k),$$

where $M_n(k)$ is the matrix coalgebra of size $n \times n$, so $k\mathcal{C}_M$ is cosemisimple. The monoidal structure is ULF if and only if $M/\preceq \rightarrow (\mathbb{N}, \cdot)$, $x \mapsto |x|$ is a map of monoids.

Example 1. As an example of a combinatorial bialgebra arising in this way, consider $M = \langle S \rangle$, the free monoid on $S = \{x, y\}$, in which two words are equivalent if they have the same length. Then $\mathcal{C}_M = \langle S \times S \rangle$ and $k\langle S \times S \rangle$ is the free algebra on four generators $\alpha = (x, x), \beta = (x, y), \gamma = (y, x), \delta = (y, y)$ whose coproduct is given by

$$\Delta(\alpha) = \alpha \otimes \alpha + \beta \otimes \gamma, \quad \Delta(\beta) = \alpha \otimes \beta + \beta \otimes \delta,$$

$$\Delta(\gamma) = \gamma \otimes \alpha + \delta \otimes \gamma, \quad \Delta(\delta) = \gamma \otimes \beta + \delta \otimes \delta.$$

In particular, it is not a Hopf algebra.

3.2. Example: Skew shapes and symmetric functions. We now elaborate further on a specific case of the previous example and realise the Hopf algebra Λ of symmetric functions (see e.g. [16, Section 2.1]) as a quotient of an incidence bialgebra.

Definition 5 (Path). For $n \in \mathbb{N}$, a path of length n is a word of length n in the alphabet $\{0, 1\}$. For a path $p = p_1 \dots p_n$ we define its height $h(p) = \sum_i p_i$ and width $w(p) = n - h(p)$.

Definition 6 (Skew shape). For two paths q, p of equal height and width we define:

$$q \leq p \Leftrightarrow \sum_{j=1}^i p_j \geq \sum_{j=1}^i q_j \forall i = 1, \dots, h + w$$

$$q < p \Leftrightarrow \sum_{j=1}^i p_j > \sum_{j=1}^i q_j \forall i = 1, \dots, h + w - 1.$$

A skew shape is a pair (q, p) of paths with $q \leq p$. If $q < p$, the shape is called connected.

Definition 7 (Category of skew shapes). We consider a discrete category \mathcal{C}_S with $Ob\mathcal{C}_S$ the set of all paths and $\mathcal{C}_S(q, p) = \{(q, p)\}$ if $q \leq p$ and \emptyset otherwise. The composition and product are defined by

$$(q, p) \circ (r, q) = (r, p)$$

$$p \cdot p' = (p_1, \dots, p_n, p'_1, \dots, p'_m)$$

$$(q, p) \cdot (q', p') = (q \cdot q', p \cdot p')$$


We depict a path in the plane by starting at the origin and drawing each 0 and 1 as a step of unit length in the direction right respectively up. A skew shape $\mu = (q, p)$ is drawn as the area bounded by q and p .

Example 2. Consider the paths

$$r = 00101, \quad q = 01010, \quad p = 10100$$

We will draw trees with the root at the top, and not indicate the total orderings explicitly (we consider leaves and roots as ordered from left to right).

Example 3.



The category \mathcal{C}_{RF} is combinatorial and so by Theorem 4 defines a bialgebra $\mathbb{Q}\mathcal{C}_{RF}$. This bialgebra, also considered explicitly in [23, 27] is made implicit use of in [2, 30] and is closely related to the Hopf algebras of trees defined in [2, 11]. In fact, we can view these Hopf algebras of trees as quotients of $\mathbb{Q}\mathcal{C}_{RF}$.

Definition 10 (Core, core equivalence). For any forest f , let $core(f)$ denote the (non-operadic) rooted forest obtained by deleting $R(f)$, $L(f)$ and all incident edges. The vertices originally adjacent to the roots in f become the roots in $core(f)$. Two forests f, g are core equivalent iff $core(f) = core(g)$.

In particular note that for any $i_n \in Id\mathcal{C}_{RF}$, $core(i_n) = i_0$, the empty tree. Let $I \triangleleft \mathbb{Q}\mathcal{C}_{RF}$ denote the ideal generated by $\{f - g \mid core(f) = core(g)\}$ in $k\mathcal{C}$.

Lemma 5. I is a bialgebra ideal and the quotient $\mathbb{Q}\mathcal{C}_{RF}/I$ is isomorphic to the planar (i.e. non-commutative) Connes-Kreimer Hopf algebra H of rooted trees [9, 11].

Core equivalence and the Connes-Kreimer quotient Hopf algebra are also discussed in [23]. Similarly, it is observed in [22, Section 4.4.2] that in general the quotient of any $k\mathcal{C}$ by the bialgebra ideal generated by all $i_x - i_1$ (which is contained in I) is a Hopf algebra.

3.4. Example: A toy model for bigraphs. Bigraphs [6, 28] are a combinatorial structure originally developed in theoretical computer science to model mobile computation. Bigraphs as defined by Milner form the morphisms of a monoidal pre-category [28, Section 2.2-2.3], i.e. a monoidal category in which not all compositions or monoidal products are defined. Here we consider a restricted definition of bigraphs, which define a combinatorial category. This toy model admits the basic bigraph operations (composition, reactions and reductions), but does not include all features. For example, we only consider private names for bigraphs, and essentially work in the support quotient of Milner's pre-category.

Definition 11 (Place graph). A place graph consists of finite disjoint sets R, S, V , total orders on R and S , and a map $prnt: V \cup S \rightarrow V \cup R$ such that:

- $R \subseteq \text{im } prnt$
- for all $v \in V \cup S$, $prnt^\infty(v) \in R$
- $prnt^\infty(v): S \rightarrow R$ is non-decreasing

The elements of R, S and V are called roots, sites and vertices respectively. Let $f_i = (R_i, S_i, V_i, prnt_i)$, $i = 1, 2$ be place graphs with $R_1 = S_2$ and all other involved sets pairwise disjoint. The composition $f_2 \circ f_1$ is defined as $(R_2, S_1, V_1 \oplus V_2, prnt)$, where $prnt(v) := prnt_i(v)$ for $v \in V_i, S_i$ except when $prnt_1(v) \in R_1 = S_2$, in which case $prnt(v) := prnt_2(prnt_1(v))$.

Definition 12 (Link graph). A link graph consists of finite disjoint sets P, X, Y with X, Y totally ordered and an equivalence relation \sim on $P \cup X \cup Y$. Each equivalence class must contain at least two elements and cannot be entirely contained in X or Y .

The elements of P, X and Y are called ports, inner names and outer names respectively. If $f_i = (P_i, X_i, Y_i, \sim_i)$, $i = 1, 2$, are link graphs with $Y_1 = X_2$, then

$$f_2 \circ f_1 := (P_1 \oplus P_2, X_1, Y_2, \sim)$$

where \sim is generated by \sim_1 and \sim_2 (including $x \sim y$ if $x \in X_1 \cup P_1, y \in P_2 \cup Y_2$, and $x \sim_1 z \sim_2 y$ for some $z \in Y_1 = X_2$).

Definition 13 (Bigraph). A bigraph $g = (g_p, g_l, \rho)$ consists of

- (1) A place graph $g_p = (R, S, V, prnt)$.
- (2) A link graph $g_l = (P, X, Y, \sim)$.
- (3) A map $\rho: P \rightarrow V$ assigning a vertex to each port.


The pairs $(|S|, |X|)$ and $(|R|, |Y|)$ are called the inner and outer interfaces of g .

It is often easier to represent a bigraph by a drawing than to define it formally. To do this, we represent roots and sites by boxes, vertices by circles and ports by dots. The parent map is indicated by nesting, the map ρ by placing a port directly on its associated vertex and the equivalence relation \sim by lines joining equivalent elements.

Example 4. The bigraph with data

$$\begin{aligned} V &= \{v_0\}, \quad R = \{r_0\} \quad S = \{s_0, s_1\} \\ prnt(s_0) &= prnt(v_0) = r_0, \quad prnt(s_1) = v_0 \\ X &= \{x_0, x_1\}, \quad Y = \emptyset \quad P = \{p_0, p_1\} \\ x_0 &\sim p_0, \quad x_1 \sim p_1 \\ \rho(p_0) &= \rho(p_1) = v_0 \end{aligned}$$

will be drawn as:



Definition 14 (Category of bigraphs). We consider a category \mathcal{C}_B with $Ob\mathcal{C}_B = \mathbb{N} \times \mathbb{N}$ and $\mathcal{C}_B((m, x), (n, y))$ the set of all bigraphs with inner and outer interfaces (m, x) and (n, y) respectively. The composition of two bigraphs $g = (g_p, g_l, \rho_g)$ and $f = (f_p, f_l, \rho_f)$ is given componentwise:

$$g \circ f = (g_p \circ f_p, g_l \circ f_l, \rho_g \oplus \rho_f)$$


if the inner interface of g equals the outer interface of f . The product of g and f is given componentwise by ordered unions (as for trees):

$$fg = (f_p \oplus g_p, f_l \oplus g_l, \rho_g \oplus \rho_f).$$


We denote the identity at the interface $\langle m, x \rangle$ by $i_{m,x}$. The monoid $(Id\mathcal{C}_B, \cdot)$ is

freely generated by the set $\{i_{0,1}, i_{1,0}\} = \left\{ \begin{array}{c} 0 \\ \vdots \\ 0 \end{array} ; \begin{array}{c} 0 \\ \square \\ 0 \end{array} \right\}$.

Example 5 (Composition and product). Consider the bigraphs



Then




This category is combinatorial and so by Theorem 4 there exists an incidence bialgebra $k\mathcal{C}_B$.

Example 6 (Coproduct in $k\mathcal{C}_B$). Consider the bigraph f' from the previous example.


$$\Delta(f') = f' \otimes i_{2,2} + i_{1,0} \otimes f' + \begin{array}{c} \boxed{\begin{array}{c} 0 \\ 1 \end{array}} \\ \otimes \end{array} \begin{array}{c} \boxed{0} \\ \otimes \end{array} \begin{array}{c} \boxed{1} \end{array}$$

We feel that the Hopf algebraic techniques used to study \mathcal{C}_{RT} and H_{CK} could provide new approaches to studying bigraphical systems. In particular, one possible application is in the study of reaction rules. Put simply, a reaction rule is a map $r : \mathcal{C}_B \rightarrow \mathcal{C}_B$ which removes a certain subset of bigraphs by mapping them to simpler ones. These rules are not necessarily compatible with the composition, i.e. in general, we have $r(g \circ f) \neq r(g) \circ r(f)$.


Example 7. Consider the reaction rule for a version of \mathcal{C}_B in which vertices can be labelled by $\{A, B, C\}$:



where the grey boxes f, g denote any substructure. This rule demands that the A and B vertices in the initial graph are siblings (i.e. $\text{prnt}^n(A) \in R$ and $\text{prnt}^n(B) \in R$ for the same n). This means that r maps the graphs



to



The second graph shows no reaction even though the bigraph does contain the relevant subgraph.

The coproduct allows us to see such blocked reactions. For our example, define the new map

$$r' := (\text{id} \otimes rM \circ) \Delta : k\mathcal{C}_B \rightarrow k\mathcal{C}_B \otimes k\mathcal{C}_B,$$

where

$$M := \begin{array}{|c|} \hline \square \\ \hline \end{array}^0 + \begin{array}{|c|} \hline \square \\ \hline \square \\ \hline \end{array}^0 + \begin{array}{|c|c|c|} \hline \square & \square & \square \\ \hline \end{array}^0 + \dots$$

Applying this, we have:

$$\begin{aligned} r' \left(\begin{array}{|c|c|} \hline \begin{array}{|c|} \hline \square \\ \hline \end{array} & \begin{array}{|c|} \hline \square \\ \hline \end{array} \\ \hline \end{array}^0 \right) &= \dots + \begin{array}{|c|} \hline \square \\ \hline \square \\ \hline \end{array}^0 \otimes r \left(\begin{array}{|c|c|} \hline \begin{array}{|c|} \hline \square \\ \hline \end{array} & \begin{array}{|c|} \hline \square \\ \hline \end{array} \\ \hline \end{array}^0 \right) + \dots \\ &= \begin{array}{|c|} \hline \square \\ \hline \square \\ \hline \end{array}^0 \otimes \dots + \begin{array}{|c|c|} \hline \begin{array}{|c|} \hline \square \\ \hline \end{array} & \begin{array}{|c|} \hline \square \\ \hline \end{array} \\ \hline \end{array}^0 + \dots \end{aligned}$$

where $i \in \mathbb{N}$ is determined by the interfaces of f and g .

3.5. A notable non-example: Hopf quivers. We now consider an example of an incidence coalgebra which admits a multiplication making it a Hopf algebra, but is not (generally) an example of our construction.

Let $Q = (Q_0, Q_1)$ be a quiver, k be a field, and \mathcal{C}_Q be the category of all paths in Q , which is locally finite. By definition, the coalgebra $k\mathcal{C}_Q$ is the path coalgebra kQ of the quiver.

Cibils and Rosso [7] (see also [8]) showed that a graded Hopf algebra structure on kQ endows Q_0 with a group and kQ_1 with a kQ_0 -Hopf bimodule structure. This in turn gives rise to what Cibils and Rosso call ramification data.

Definition 15 (Ramification data). Let G be a group. By ramification data for G , we mean a sum $r = \sum_{C \in \mathfrak{C}} r_C C$ for \mathfrak{C} the set of conjugacy classes of G and all $r_C \in \mathbb{N}$.

Definition 16 (Hopf quiver). Let G be a group and r some ramification data for G . The quiver with $Q_0 = G$ and r_C arrows from g to cg for each $g, c \in G$, where C is the conjugacy class containing c , is called the Hopf quiver determined by (G, r) .

Theorem 5. *The path coalgebra kQ of a quiver Q admits graded Hopf algebra structures if and only if Q is a Hopf quiver.*

To be more precise, for a Hopf quiver Q , there is a canonical Hopf algebra structure on kQ , with $kQ_1 \cong \bigoplus_{C \in \mathfrak{e}} r_C kC \otimes kQ_0$ as vector space. The kQ_0 -bimodule structure is given by

$$x(c \otimes g)y = xcx^{-1} \otimes xgy$$

and the kQ_0 -bicomodule structure is given by

$$c \otimes g \mapsto cg \otimes (c \otimes g) \otimes g.$$

The group structure on Q_0 and the kQ_0 -bimodule structure on kQ_1 define the algebra structure on kQ in lowest degrees, which is extended universally to all of kQ , see [7, Theorem 3.8]. However, given a Hopf quiver, there are in general also Hopf algebra structures on kQ compatible with r that are different from the canonical one. The choice is in the kQ_0 -bimodule structure of kQ_1 which amounts to a choice of an n_C -dimensional representation of the centraliser of an element $c \in C$. The extension of the product to paths is then unique.

Huang and Torecillas [18] proved that a quiver path coalgebra kQ always admits graded bialgebra structures. The results of Green and Solberg [13] are also closely related, but different in that they study path algebras rather than path coalgebras. Here we focus on the Hopf algebra setting as considered in [7].

Our key aim is to stress that the product in a quiver Hopf algebra kQ is not a linear extension of a monoidal product on \mathcal{C}_Q unless $Q_1 = \emptyset$. To do this, we classify all monoidal structures on path categories of quivers whose vertices form a group under \cdot :

Lemma 6. *Assume that the path category \mathcal{C}_Q of a quiver is monoidal such that (Q_0, \cdot) forms a group. Then:*

- (1) *The monoidal product defines commuting left and right Q_0 -actions on Q_1 .*
- (2) *The path length ℓ is a grading with respect to both \cdot and \circ .*
- (3) *Either Q_1 is empty, or there exists an element $z \in Z(Q_0)$ such that Q_1 contains for each $a \in Q_0$ exactly one arrow $f_a: a \rightarrow z \cdot a$.*

Proof. If $Q_1 = \emptyset$ there is nothing to prove, so we assume $Q_1 \neq \emptyset$.

1. To begin with, we prove that for any identity morphism i_a , $a \in Q_0$, and any arrow $f: b \rightarrow c \in Q_1$, we have $i_a \cdot f, f \cdot i_a \in Q_1$. Indeed, we have

$$i_a \cdot f = g_1 \circ \dots \circ g_n$$

for unique arrows $g_i \in Q_1$, and then

$$f = i_{a^{-1}} \cdot (g_1 \circ \dots \circ g_n) = (i_{a^{-1}} \cdot g_1) \circ \dots \circ (i_{a^{-1}} \cdot g_n)$$

where we used that $i_a \cdot i_{a^{-1}} = i_{a \cdot a^{-1}} = i_1$ and that \cdot is a monoidal product. It is impossible that $i_{a^{-1}} \cdot g_i = i_b$ for some $b \in Q_0$, as we would then have $g_i = i_a \cdot i_b = i_{a \cdot b}$. So the right hand side of the above equation is a path of length at least n while the left hand side has length 1, hence $n = 1$. Analogously, one proves $f \cdot i_a \in Q_1$. That \cdot defines commuting actions is immediate.

2. For any arrow f , let $s(f), t(f)$ denote its source and target vertices respectively. As \cdot is a monoidal product, we have for any two arrows f, g :

$$(2) \quad f \cdot g = (i_{t(f)} \cdot g) \circ (f \cdot i_{s(g)}) = (f \cdot i_{t(g)}) \circ (i_{s(f)} \cdot g).$$

By what has been shown already, this is a path of length 2. Continuing inductively, one proves that $\ell(f \cdot g) = \ell(f) + \ell(g)$ for all paths $f, g \in \mathcal{C}_Q$.

3. We also conclude from (2) that

$$s(f) \cdot t(g) = t(f) \cdot s(g) \Rightarrow t(g) \cdot s(g)^{-1} = s(f)^{-1} \cdot t(f) =: z \in Q_0$$

is constant (independent of f). So

$$t(f) = s(f) \cdot z = z \cdot s(f)$$

and any arrow $f \in Q_1$ with source a has the same target $z \cdot a = a \cdot z$.

Given any arrow $f: a \rightarrow a \cdot z$ and $b \in Q_0$, there exists $i_b \cdot f: b \cdot a \rightarrow z \cdot b \cdot a$. This means that the same number of arrows go out of each vertex and that z is in the centre of Q_0 .

Finally, assume there are two arrows f, g with source 1 (the unit element of Q_0) and consider again (2):

$$f \cdot g = (i_z \cdot g) \circ f = (f \cdot i_z) \circ g.$$

We deduce that $f = g$. □

If $Q_1 = \emptyset$, then $\mathcal{C}_Q = \{i_a : a \in Q_0\}$ is just a group. The monoidal product is the group multiplication, and this is ULF. The Hopf algebra $k\mathcal{C}_Q$ is the group algebra of Q_0 and is combinatorial.

If $Q_1 \neq \emptyset$, there are two sub-cases: $z = 1$ and $z \neq 1$. In the first case, each vertex $a \in Q_0$ has a unique arrow $f_a: a \rightarrow a$. In the second case, each vertex a has one incoming arrow $f_{z^{-1} \cdot a}: z^{-1}a \rightarrow a$ and one outgoing arrow $f_a: a \rightarrow za$.

In both cases, any morphism in \mathcal{C}_Q can be uniquely expressed as

$$f_{z^n \cdot a} \circ \dots \circ f_a, \quad a \in Q_0, n \in \mathbb{N},$$

to be interpreted as i_a when $n = 0$. The monoidal product is given by

$$(3) \quad (f_{z^n \cdot a} \circ \dots \circ f_a) \cdot (f_{z^l \cdot b} \circ \dots \circ f_b) = f_{z^{n+l} \cdot a \cdot b} \circ \dots \circ f_{a \cdot b}.$$

This product is evidently not ULF, so we obtain:

Theorem 6. *Given a Hopf quiver Q , the quiver Hopf algebra kQ is a case of Theorem 4 if and only if $Q_1 = \emptyset$*

Remark 3 (Another, similar, non-example). In [10], Crossley defines several Hopf algebra structures on $k\langle S \rangle$ for a set S . In two of them, the coalgebra structure is an incidence coalgebra (on the free monoidal category with one object and a morphism for each $s \in S$). However, just like the quiver Hopf algebras of Cibils and Rosso, these two Hopf algebras are not examples of our construction, as in both cases, $\langle S \rangle \cdot \langle S \rangle \not\subseteq \langle S \rangle$.

4. WEAK HOPF ALGEBRAS FROM MONOIDAL CATEGORIES

In this section we consider the following:

Definition 17 (2-group). A strict 2-group is a strict monoidal groupoid in which every object is invertible, i.e. in which we have $\forall x \in \text{Ob}\mathcal{C} \exists y \in \text{Ob}\mathcal{C} : x \cdot y = 1 = y \cdot x$.

Lemma 7. *Let \mathcal{C} be a 2-group. Then for any $f \in \mathcal{C}$ there exists $\bar{f} \in \mathcal{C}$ such that $\bar{f} \cdot f = i_1 = f \cdot \bar{f}$.*

Proof. Take $\bar{f} = i_{\overline{t(f)}} \cdot f^{-1} \cdot i_{\overline{s(f)}}$, where $\overline{t(f)}, \overline{s(f)}$ denote the inverses in $(\text{Ob}\mathcal{C}, \cdot)$. □

Strict 2-groups are the objects of a subcategory of **Grpd**, where we keep only the functors of groupoids which preserve the group structure of \mathcal{C} with respect to \cdot .

Definition 18 (Source subgroup). In any 2-group $(\mathcal{C}, \cdot, 1)$ we define the source subgroup

$$S := \bigcup_{x \in \text{Ob}\mathcal{C}} \mathcal{C}(1, x),$$

which contains all morphisms with source 1. (S, \cdot) is a normal subgroup of (\mathcal{C}, \cdot) .

Lemma 8. *A 2-group $(\mathcal{C}, \cdot, 1)$ is locally finite if and only if $|S|$ is finite. In this case, the monoidal product has the $|S|$ -LF property (Definition 3).*

Proof. Multiplication by i_x defines a bijection $\mathcal{C}(1, y) \rightarrow \mathcal{C}(x, xy)$ for all $x, y \in \text{Ob}\mathcal{C}$. As \mathcal{C} is a groupoid, we have

$$\begin{aligned} N_2(f) &= \{(f \circ g^{-1}, g) \mid s(g) = s(f)\} \\ \Rightarrow |N_2(f)| &= |\{g \mid s(g) = s(f)\}| = |S| \end{aligned}$$

for all $f \in \mathcal{C}$, and the map $N_2(f) \times N_2(g) \rightarrow N_2(f \cdot g)$ from Definition 3 has inverse

$$N_2(f \cdot g) \ni (x, y) \mapsto \{(x, y), (\bar{x} \cdot a, \bar{y} \cdot b) \mid (x, y) \in N_2(f)\}. \quad \square$$

Remark 4. The second part of this lemma implies that a 2-group \mathcal{C} is a Möbius category if and only if $\mathcal{C} = \text{Id}\mathcal{C}$.

The $|S|$ -LF property is sufficient to define compatible algebra and coalgebra structures on $k\mathcal{C}$ using the scaled version of the incidence coalgebra (Definition 3). As it is less well-known than the definition of a Hopf algebra, we recall here the definition of a weak Hopf algebra, following [4]:

Definition 19 (Weak Hopf algebra). Let k be a field. A weak k -Hopf algebra is a tuple $(A, \mu, 1, \Delta, \epsilon, S)$ satisfying:

- (A1) $(A, \mu, 1)$ is an associative unital k -algebra;
- (A2) (A, Δ, ϵ) is a coassociative counital k -coalgebra
- (A3) The coproduct is multiplicative: $\Delta(ab) = \Delta(a)\Delta(b)$
- (A4) The counit is weakly multiplicative:

$$\epsilon(ab) = \epsilon(a1_{(1)})\epsilon(1_{(2)}b) = \epsilon(a1_{(2)})\epsilon(1_{(1)}b)$$

- (A5) The unit is weakly comultiplicative:

$$(\Delta(1) \otimes 1)(1 \otimes \Delta(1)) = \Delta^2(1) = (1 \otimes \Delta(1))(\Delta(1) \otimes 1)$$

- (A6) $S : A \rightarrow A$ is a k -linear map (called the antipode) which satisfies

$$\begin{aligned} S(a) &= S(a_{(1)})a_{(2)}S(a_{(3)}) \\ S(a_{(1)})a_{(2)} &= \epsilon(1_{(1)}a)1_{(2)} \\ a_{(1)}S(a_{(2)}) &= 1_{(2)}\epsilon(a1_{(2)}). \end{aligned}$$

Remark 5. Under the assumption of axioms (A1), (A2) and (A3), the version of axiom (A4) given here is equivalent to the formulation in [4]:

$$\epsilon(abc) = \epsilon(ab_{(1)})\epsilon(b_{(2)}c) = \epsilon(ab_{(2)})\epsilon(b_{(1)}c)$$

We can now state the central theorem of this section:

Theorem 7. *Let $(\mathcal{C}, \cdot, 1)$ be a locally finite 2-group, k a field of characteristic zero and $k\mathcal{C}$ the free k -module on \mathcal{C} . We understand \otimes to be the tensor product \otimes_k and $(k\mathcal{C}, \Delta, \epsilon)$ the scaled incidence coalgebra of $k\mathcal{C}$ with $\lambda = |S|$ (Definition 3). Then $(k\mathcal{C}, \cdot, i_1, \Delta, \epsilon, S)$ is a weak Hopf algebra with antipode $S(f) = \bar{f}^{-1}$.*

Proof. (A1) and (A2) are immediate by Theorems 3 and 2. (A3) then follows by Lemma 8, (A4) by Lemma 7. For (A5) we observe that

$$\begin{aligned} N_2(i_1) \times N_2(i_1) &\rightarrow N_3(i_1) \\ ((a, b); (c, d)) &\mapsto (a, b \cdot c, d) \end{aligned}$$

defines a bijection and that $b \circ c = b \cdot c = c \cdot b$ for $s(b) = t(c) = 1$. (A6) can be verified by a short computation. \square

4.1. Example: Relations on groups. Here we present the analogy to Example 1: let \sim be a reflexive transitive relation on a group G which is compatible with the multiplication as in (1) and consider the category \mathcal{C} with $Ob\mathcal{C} = G$, $\mathcal{C}(h, g) = \{(h, g) \mid g \sim h\}$. This category is a locally finite 2-group iff \sim is an equivalence relation and the equivalence classes are finite.

By Theorem 7, $k\mathcal{C}$ admits a weak incidence Hopf algebra structure with a co-product which runs through equivalence classes:

$$\Delta(h, g) = \sum_{g \sim f \sim h} (h, f) \otimes (f, g).$$

In the case that \sim is equality, the weak Hopf algebra obtained is the group algebra kG . This can be alternatively stated in the following way: let G be a group and $N \triangleleft G$ a normal subgroup. Define a 2-group \mathcal{C} by $Ob\mathcal{C} = G$ and

$$|\mathcal{C}(g, h)| = \begin{cases} 1, & \exists z \in N : h = zg \\ 0, & \text{else} \end{cases}.$$

Then $k\mathcal{C}$ admits a weak incidence Hopf algebra if and only if $|N|$ is finite.

4.2. Example: Automorphism 2-groups. Let $\mathcal{C} = (Ob\mathcal{C}, Mor_1\mathcal{C}, Mor_2\mathcal{C})$ be a strict 2-category. Then for each $x \in Ob\mathcal{C}$ there exists a strict 2-group $AUT(x)$ whose objects are the automorphisms of x as an object of \mathcal{C} and whose morphisms are the 2-isomorphisms between these. The product of two objects f, f' in $AUT(x)$ is given by their composition in \mathcal{C} and the unit object is i_x . The composition of morphisms in $AUT(x)$ is given by the vertical composition of 2-morphisms in \mathcal{C} and the product of morphisms in $AUT(x)$ is given by the horizontal composition in \mathcal{C} . According to Theorem 7, $kAUT(x)$ admits a weak incidence Hopf algebra structure iff $\{f \in Mor_1\mathcal{C} \mid f \sim i_x\}$ is finite.

This example overlaps with Example 4.1. Consider the 2-category **Grp**, whose objects are groups, 1-morphisms are group homomorphisms and 2-morphisms are given by inner automorphisms in the target group, i.e. for $f_1, f_2 : G \rightarrow H$, there exists $\phi_h : f_1 \rightarrow f_2$ iff there exists $h \in H$ such that $f_1 = f_h \circ f_2$, where f_h is the morphism in $Inn(H)$ given by conjugation by h . If G is an object in **Grp**, $ObAUT(G) := Aut(G)$, the usual group of automorphisms of G , and $AUT(G)$ (the set of morphisms) contains an arrow between any pair of automorphisms which are related by an inner automorphism of G . Hence $kAUT(G)$ admits a weak Hopf algebra structure iff $Inn(G)$ is finite.

4.3. Crossed modules. Here we will reformulate the result of Theorem 7 in the language of crossed modules.

Definition 20 (Crossed module). A crossed module (G, H, τ, α) consists of:

- Groups G, H
- A group action $\alpha : G \times H \rightarrow H$
- A group homomorphism $\tau : H \rightarrow G$.

such that α and τ satisfy

$$\begin{aligned}\tau(\alpha(g, h)) &= g\tau(h)g^{-1} \\ \alpha(\tau(h), h') &= hh'h^{-1}.\end{aligned}$$

Crossed modules form the objects of a category whose morphisms $(f_1, f_2) : (G, H, \tau, \alpha) \rightarrow (G', H', \tau', \alpha')$ consist of pairs of group homomorphisms $f_1 : G \rightarrow G'$ and $f_2 : H \rightarrow H'$ such that $\tau'f_2 = f_1\tau$.

Theorem 8 (Brown and Spencer). *The categories of strict 2-groups and crossed modules are equivalent.*

For the original statement and proof see [5], for further exploration see e.g. [1, 12].

Given a 2-group $(\mathcal{C}, \cdot, 1)$, the corresponding crossed module (G, H, α, τ) is defined as follows: set $G = \text{Ob}\mathcal{C}$, $H = S$, $\alpha(g, h) = i_g \cdot h \cdot i_{g^{-1}}$ and $\tau(h) = t(h)$.

Conversely, given a crossed module (G, H, α, τ) , the corresponding strict 2-group $(\mathcal{C}, \cdot, 1)$ has $\text{Ob}\mathcal{C} = G$, $\mathcal{C} = H \rtimes_{\alpha} G$, $s(h, g) = g$ and $t(h, g) = \tau(h)g$. Composition of morphisms is given by $(h', \tau(h)g) \circ (h, g) = (h'h, g)$.

This equivalence gives us the following:

Corollary 1 (to Theorem 7). *Let (G, H, τ, α) be a crossed module, k a field and $k(H \times G)$ the free k -module on the set $H \times G$. Then $k(H \times G)$ admits the following weak Hopf algebra structure iff $|H|$ is finite:*

$$\begin{aligned}(h, g) \cdot (h', g') &= (h\alpha(g, h'), gg') \\ \Delta(h, g) &= \sum_{h'h''=h} (h'', \tau(h')g) \otimes (h', g) \\ S(h, g) &= (\alpha(g^{-1}, h^{-1})^{-1}, g^{-1})\end{aligned}$$

If $\tau : H \rightarrow G$ is injective, we are in the case of Example 4.1, i.e. H is isomorphic to a normal subgroup of G .

REFERENCES

- [1] John C. Baez and Aaron D. Lauda, *Higher-Dimensional Algebra V: 2-Groups*, Theory and Applications of Categories 12 (2004), 423-491.
- [2] Christoph Bergbauer and Dirk Kreimer, *Hopf algebras in renormalisation theory: locality and Dyson-Schwinger equations from Hochschild cohomology*, IRMA Lect.Math.Theor.Phys. 10 (2006) 133-164.
- [3] Nantel Bergeron, Stefan Mykytiuk, Frank Sottile, Stephanie van Willigenburg, *Non-commutative Pieri operators on posets*, J. Combin. Th. Ser. A, Vol. 91, No. 1/2, Aug 2000, pp. 84-110.
- [4] Gabriella Böhm, Florian Nill, Kornél Szlachányi *Weak Hopf Algebras I. Integral Theory and C*-structure* arXiv:math/9805116
- [5] R. Brown and C. B. Spencer, *G-groupoids, crossed modules, and the classifying space of a topological group* Proc. Kon. Akad. v. Wet. 79 (1976), 296-302
- [6] Michele Sevegnani and Muffy Calder *Bigraphs with sharing*, TR-2010-310, DCS Technical Report Series, Department of Computing Science, University of Glasgow, 2010
- [7] Claude Cibils and Marc Rosso, *Hopf quivers*, 2000 arXiv:math/0009106.
- [8] Claude Cibils, Aaron Lauve and Sarah Witherspoon, *Hopf quivers and Nichols algebras in positive characteristic* 2009
- [9] Alain Connes and Dirk Kreimer, *Hopf Algebras, Renormalization and Noncommutative Geometry* Commun.Math.Phys. 199 (1998) 203-242.
- [10] M.D. Crossley, *Some Hopf Algebras of Words*, Glasgow Math. J. 48 (2006) 575-582.
- [11] Loïc Foissy, *An introduction to Hopf algebras of trees*, preprint.
- [12] Magnus Forrester-Barker, *Group Objects and Internal Categories*, 2002 arXiv:math/0212065v1.
- [13] E. L. Green and Ø. Solberg, *Basic Hopf algebras and quantum groups*, Mathematische Zeitschrift 229(1):45-76 · September 1998
- [14] Imma Gálvez-Carillo, Joachim Kock and Andrew Tonks, *Decomposition Spaces in Combinatorics* 2016 arXiv:1612.09225v2

- [15] Imma Gálvez-Carillo, Joachim Kock and Andrew Tonks, *Decomposition Spaces, incidence algebras and Möbius inversion III: the decomposition space of Möbius intervals*. 2015 arXiv:1512.07580
- [16] Darij Grinberg and Victor Reiner. *Hopf algebras in combinatorics*. 2014 arXiv:1409.8356
- [17] Ralf Holtkamp, *On Hopf algebra structures over free operads*. Adv. Math. 207 (2006), no. 2, 544–565.
- [18] Hua-Lin Huang and Blas Torrecillas, *Quiver bialgebras and monoidal categories*, Colloq. Math. 131 (2013), 287-300.
- [19] Ole Høgh Jensen and Robin Milner, *Bigraphs and mobile processes (revised)*, University of Cambridge computer laboratory technical report, 2004.
- [20] S. A. Joni and G.-C. Rota, *Coalgebras and bialgebras in combinatorics*, Stud. Appl. Math., 61 (1979), pp. 93–139.
- [21] André Joyal and Ross Street *Braided monoidal categories* Macquarie Mathematics Report No. 860081, November 1986.
- [22] Ralph M. Kaufmann, Benjamin C. Ward *Feynman Categories* arXiv:1312.1269v3
- [23] Joachim Kock, *Polynomial functors and combinatorial Dyson-Schwinger equations*, J. Math. Phys. 58 (2017), 041703.
- [24] F. W. Lawvere and M. Menni, *The Hopf Algebra of Möbius Intervals* Theory and Applications of Categories, Vol. 24, No. 10, 2010, pp. 221–265.
- [25] P. Leroux *Les Catégories de Möbius* Cahiers de Topologie et Géométrie Différentielle Catégoriques, 16(3):280–282, 1975.
- [26] Jean-Louis Loday and María Ronco, *Combinatorial Hopf algebras*, 2008 arXiv:0810.0435.
- [27] Dominique Manchon, *Hopf algebras, from basics to applications to renormalization*, Comptes Rendus des Rencontres Mathématiques de Glanon 2001.
- [28] Robin Milner, *The space and motion of communicating agents*, Cambridge University Press 2008.
- [29] David E. Radford, *Hopf algebras*, World Scientific, 2012.
- [30] Lucia Rotheray, *Hopf Subalgebras from Green's Functions* MSc thesis, Humboldt-Universität zu Berlin, 2015.
- [31] Karen Yeats, *A Hopf algebraic approach to Schur function identities* arXiv:1511.06337v4

INSTITUT FÜR GEOMETRIE, TECHNISCHE UNIVERSITÄT DRESDEN

E-mail address: `ulrich.kraehmer@tu-dresden.de`, `lucia.rotheray@tu-dresden.de`