

# Equivariant Euler characteristics of $\overline{\mathcal{M}}_{g,n}$

Adrian Diaconu

## Abstract

Let  $\overline{\mathcal{M}}_{g,n}$  be the moduli space of  $n$ -pointed stable genus  $g$  curves, and let  $\mathcal{M}_{g,n}$  be the moduli space of  $n$ -pointed smooth curves of genus  $g$ . In this paper, we obtain an asymptotic expansion for the characteristic of the free modular operad  $\mathbb{M}\mathcal{V}$  generated by a stable  $\mathbb{S}$ -module  $\mathcal{V}$ , allowing to effectively compute  $\mathbb{S}_n$ -equivariant Euler characteristics of  $\overline{\mathcal{M}}_{g,n}$  in terms of  $\mathbb{S}_{n'}$ -equivariant Euler characteristics of  $\mathcal{M}_{g',n'}$  with  $0 \leq g' \leq g$ ,  $\max\{0, 3-2g'\} \leq n' \leq 2(g-g') + n$ . This answers a question posed by Getzler and Kapranov by making their integral representation of the characteristic of the modular operad  $\mathbb{M}\mathcal{V}$  effective. To illustrate how the asymptotic expansion is used, we give formulas expressing the generating series of the  $\mathbb{S}_n$ -equivariant Euler characteristics of  $\overline{\mathcal{M}}_{g,n}$ , for  $g = 0, 1$  and  $2$ , in terms of the corresponding generating series associated with  $\mathcal{M}_{g,n}$ .

## Contents

<b>1</b>	<b>Introduction</b>	<b>2</b>
<b>2</b>	<b>Notation and preliminaries</b>	<b>3</b>
<b>3</b>	<b>A semi-classical expansion</b>	<b>6</b>
3.1	The critical points . . . . .	7
3.2	The asymptotic expansion . . . . .	9
<b>4</b>	<b>Equivariant Euler characteristics</b>	<b>11</b>

---

School of Mathematics, University of Minnesota, Minneapolis, MN 55455  
E-mail: cad@umn.edu

# 1 Introduction

Let  $\mathcal{V}$  be a stable  $\mathbb{S}$ -module, i.e., a collection of chain complexes  $\{\mathcal{V}((g, n))\}_{g, n \geq 0}$  with an action of the symmetric group  $\mathbb{S}_n$  on  $\mathcal{V}((g, n))$ , and such that  $\mathcal{V}((g, n)) = 0$  when  $2g + n - 2 \leq 0$ , and let

$$\text{Ch}(\mathcal{V}) = \sum_{2(g-1)+n > 0} \hbar^{g-1} \text{ch}(\mathcal{V}((g, n)))$$

denote the characteristic of  $\mathcal{V}$ , where  $\text{ch}(\mathcal{V}((g, n)))$  is the characteristic of the  $\mathbb{S}_n$ -representation  $\mathcal{V}((g, n))$ . To the stable  $\mathbb{S}$ -module  $\mathcal{V}$ , there is the associated free modular operad  $\mathbb{M}\mathcal{V}$  generated by  $\mathcal{V}$ ; as such, we can also take the characteristic of  $\mathbb{M}\mathcal{V}$ . Throughout, we shall consider only  $\mathbb{S}$ -modules in the category of  $\ell$ -adic Galois representations.

The aim of this paper is to make quite effective a beautiful result of Getzler and Kapranov [15, Theorem 8.13], expressing the relationship between the characteristics of  $\mathcal{V}$  and  $\mathbb{M}\mathcal{V}$  in the form

$$\text{Ch}(\mathbb{M}\mathcal{V}) = \text{Log}(\exp(\Delta)\text{Exp}(\text{Ch}(\mathcal{V}))) \quad (1)$$

where  $\Delta$  is a certain analogue of the Laplacian,  $\text{Exp}(f)$  is the *plethystic* exponential of  $f$ , and  $\text{Log}(f)$  is the inverse of the plethystic exponential; see Section 2 for precise definitions. The above equality is a consequence of how the functor  $\mathbb{M}$  on the category of stable  $\mathbb{S}$ -modules is tailored to the way the boundary strata of  $\overline{\mathcal{M}}_{g,n}$  are obtained by gluing together moduli spaces  $\mathcal{M}_{g',n'}$ .

The formula (1) is a natural generalization of Wick's theorem [5], which gives the integral formula

$$\sum_{2(g-1)+n > 0} \text{M}v_{g,n} \hbar^{g-1} \frac{\xi^n}{n!} = \log \int_{\mathbb{R}} \exp \left( \sum_{2(g-1)+n > 0} v_{g,n} \hbar^{g-1} \frac{x^n}{n!} - \frac{(x-\xi)^2}{2\hbar} \right) \frac{dx}{\sqrt{2\pi\hbar}}. \quad (2)$$

Here  $\{v_{g,n} : 2(g-1) + n > 0\}$  is a set of variables, and

$$\text{M}v_{g,n} := \sum_{G \in \text{Ob } \Gamma_{g,n}} \frac{1}{|\text{Aut}(G)|} \prod_{v \in \text{Vert}(G)} v_{g(v), n(v)} \quad (\text{for } 2(g-1) + n > 0)$$

where  $\Gamma_{g,n}$  is the *finite* category whose objects are isomorphism classes of stable graphs of genus  $g$  with  $n$  ordered legs, and whose morphisms are the automorphisms; see [15] for details. Thus if we define

$$b_g = b_g(\xi) = \sum_{n \geq \max\{0, 3-2g\}} \text{M}v_{g,n} \frac{\xi^n}{n!} \quad \text{and} \quad a_g = a_g(x) = \sum_{n \geq \max\{0, 3-2g\}} v_{g,n} \frac{x^n}{n!}$$

then, by performing an asymptotic expansion of the integral in the right-hand side of (2), one can obtain formulas expressing the coefficients  $b_g$  in terms of  $a_{g'}$  with  $g' \leq g$ .

Our main result, Theorem 3.2, provides an asymptotic expansion of the right-hand side of (1), thus answering a question posed by Getzler and Kapranov in [15], p. 113. Letting  $\mathbf{a}_g$  and  $\mathbf{b}_g$  denote, as above, the coefficients of  $\hbar^{g-1}$  in  $\text{Ch}(\mathcal{V})$  and  $\text{Ch}(\mathbb{M}\mathcal{V})$ , respectively, then as consequences of Theorem 3.2, we shall obtain formulas for  $\mathbf{b}_0$ ,  $\mathbf{b}_1$  and  $\mathbf{b}_2$  in terms of  $\mathbf{a}_0$ ,  $\mathbf{a}_1$  and  $\mathbf{a}_2$ , see Section 4. The formulas for  $\mathbf{b}_0$  and  $\mathbf{b}_1$  (Theorem 4.1 and Theorem 4.3, respectively) are not new, see [15, Theorem 7.17] or [13, Theorem 5.9] for the calculation of  $\mathbf{b}_0$ , and [14] or [24] for that of  $\mathbf{b}_1$ . The proofs of these two results were merely included as examples of how the coefficients  $\mathbf{b}_g$  are calculated.

The argument used to prove Theorem 3.2 represents the natural generalization of the method used by Bini and Harer [6, Section 3] to study the asymptotic expansion of the integral in (2). It is also possible to give an analogous interpretation of the asymptotic expansion discussed in this paper to that in [6, Proposition 3.6], as an expansion over stable graphs, to obtain explicit formulas for the coefficients  $\mathbf{b}_g$  when  $g \geq 3$ . The general formula for  $\mathbf{b}_g$  can then be combined with a result of Gorsky [16], where the author establishes a formula for the generating series of the numerical  $\mathbb{S}_n$ -equivariant Euler characteristics of  $\mathcal{M}_{g,n}$ , to obtain the corresponding numerical  $\mathbb{S}_n$ -equivariant characteristics of  $\overline{\mathcal{M}}_{g,n}$ .

Although we shall work throughout just in the tensor symmetric abelian category  $\mathbf{Rep}_{\mathbb{Q}_\ell}(\mathrm{Gal}(\overline{\mathbb{Q}}/\mathbb{Q}))$  of  $\ell$ -adic Galois representations,<sup>1</sup> the results in this paper are valid for stable  $\mathbb{S}$ -modules in any symmetric monoidal category with finite colimits and additive over a field of characteristic zero.

**Acknowledgements.** I am grateful to Jonas Bergström and Dan Petersen for their comments and suggestions. I would also like to thank them for making available to me the formulas for the  $\mathbb{S}_n$ -equivariant Euler characteristics of  $\overline{\mathcal{M}}_{2,n}$  for small values of  $n$ .

## 2 Notation and preliminaries

**Symmetric functions.** Let  $\mathbb{S}_k$  denote the symmetric group on  $k$  letters. The completed ring of symmetric functions in infinitely many variables (see [15]) is defined by

$$\Lambda = \varprojlim \mathbb{Z}[x_1, \dots, x_k]^{\mathbb{S}_k}.$$

We have the standard functions

$$e_n = \sum_{i_1 < \dots < i_n} x_{i_1} \cdots x_{i_n} \quad h_n = \sum_{i_1 \leq \dots \leq i_n} x_{i_1} \cdots x_{i_n} \quad \text{and} \quad p_n = \sum_{i \geq 1} x_i^n$$

called the elementary symmetric functions, the complete symmetric functions and the power sums, respectively. The ring  $\Lambda$  is also the completion of the graded polynomial ring  $\mathbb{Z}[e_1, e_2, \dots]$ , and thus, by the well-known identities among the standard symmetric functions, we have

$$\Lambda = \mathbb{Z}[e_1, e_2, \dots] = \mathbb{Z}[h_1, h_2, \dots] \quad \text{and} \quad \Lambda \otimes_{\mathbb{Z}} \mathbb{Q} = \mathbb{Q}[p_1, p_2, \dots].$$

In addition, for  $\lambda = (\lambda_1 \geq \dots \geq \lambda_n > 0)$  we have the Schur functions

$$s_\lambda = \det(h_{\lambda_i - i + j})_{1 \leq i, j \leq n}$$

which also generate  $\Lambda$ .

**Plethysm.** One defines (see [22] and [15]) the associative operation “ $\circ$ ” on  $\Lambda$ , called *plethysm* (or *composition*), characterized by the following conditions:

1.  $(f_1 + f_2) \circ g = f_1 \circ g + f_2 \circ g$
2.  $(f_1 f_2) \circ g = (f_1 \circ g)(f_2 \circ g)$
3. If  $f = f(p_1, p_2, \dots)$ , we have  $p_n \circ f = f(p_n, p_{2n}, \dots)$ .

In other words,  $\Lambda$  is a  $\lambda$ -ring, see [20] and [11] for generalities on  $\lambda$ -rings; it is the complete filtered  $\lambda$ -ring obtained by completing the free  $\lambda$ -ring on one generator  $\mathbb{Z}[e_1, e_2, \dots]$ . In this language  $p_n \circ f = \psi_n(f)$ , where  $\psi_n$  denote the Adams operations.

Consider the ring  $\Lambda(\!(\hbar)\!) of Laurent series with the topology induced by the descending filtration$

$$F^n \Lambda(\!(\hbar)\!) = \left\{ \sum_i f_i \hbar^i : f_i \in F^{n-2i} \Lambda \right\}.$$

The plethysm extends to  $\Lambda \times \Lambda(\!(\hbar)\!)$  by keeping conditions 1., 2. above, and replacing the last condition by:

$$p_n \circ f(\hbar, p_1, p_2, \dots) = f(\hbar^n, p_n, p_{2n}, \dots).$$

<sup>1</sup>In fact, we shall only deal with virtual  $\ell$ -adic Galois representations (Euler characteristics); thus it will suffice to work instead with the characters of the representations at Frobenius elements.

For  $f \in F^1\Lambda(\hbar)$ , put  $\Psi(f) := \sum_{n \geq 1} \frac{1}{n} \psi_n(f)$ , where  $\psi_n(f) = p_n \circ f$ , and let  $\text{Exp}(f) = \exp(\Psi(f))$ ; we have

$$\text{Exp}(f + g) = \text{Exp}(f)\text{Exp}(g).$$

The mapping  $\Psi : F^1\Lambda(\hbar) \rightarrow F^1\Lambda(\hbar)$  has an inverse  $\Psi^{-1} : F^1\Lambda(\hbar) \rightarrow F^1\Lambda(\hbar)$  given by

$$\Psi^{-1}(f) = \sum_{m \geq 1} \frac{\mu(m)}{m} \psi_m(f)$$

where  $\mu(m)$  is the usual Möbius function; see [23, Lemma 20]. Thus if we define  $\text{Log} : 1 + F^1\Lambda(\hbar) \rightarrow F^1\Lambda(\hbar)$  by  $\text{Log}(g) = \Psi^{-1}(\log g)$ , then  $\text{Exp}$  and  $\text{Log}$  are inverses of each other.

Finally, one defines ([15, Section 8]) an analogue of the Laplacian on  $\Lambda(\hbar)$  by

$$\Delta = \sum_{n=1}^{\infty} \hbar^n \left( \frac{n}{2} \frac{\partial^2}{\partial p_n^2} + \frac{\partial}{\partial p_{2n}} \right).$$

Note that  $\Delta$  preserves the filtration of  $\Lambda(\hbar)$ .

**The moduli spaces  $\mathcal{M}_{g,n}$  and  $\overline{\mathcal{M}}_{g,n}$ .** For  $g, n \in \mathbb{N}$  with  $2(g-1) + n > 0$ , let  $\overline{\mathcal{M}}_{g,n}$  denote the proper and smooth Deligne-Mumford stack of *stable curves of arithmetic genus  $g$  with  $n$  ordered distinct smooth points*. Let  $\mathcal{M}_{g,n} \subset \overline{\mathcal{M}}_{g,n}$  be the open substack of irreducible and non-singular curves; both  $\mathcal{M}_{g,n}$  and  $\overline{\mathcal{M}}_{g,n}$  are defined over  $\text{Spec}(\mathbb{Z})$ , and the group  $\mathbb{S}_n$  acts on them by permuting the marked points on the curves. Moreover, we know that the boundary  $\overline{\mathcal{M}}_{g,n} \setminus \mathcal{M}_{g,n}$  is a normal crossings divisor.

The stack  $\overline{\mathcal{M}}_{g,n}$  admits a stratification determined by stable graphs of genus  $g$  with  $n$  ordered legs; see [1, Chap. XII.10]. To each stable graph  $G$ , there corresponds the smooth, locally closed stratum  $\mathcal{M}(G) \subset \overline{\mathcal{M}}_{g,n}$  parametrizing curves with dual graph isomorphic to  $G$ . The stratum  $\mathcal{M}(G)$  is canonically isomorphic to the quotient stack

$$\left[ \left( \prod_{v \in \text{Vert}(G)} \mathcal{M}_{g(v), n(v)} \right) / \text{Aut}(G) \right].$$

Here  $\text{Vert}(G)$  denotes the set of vertices of  $G$ ,  $g(v)$  is the geometric genus of the component (of a stable curve) corresponding to  $v$  and  $n(v)$  is the valence of the vertex  $v$ . The automorphism group  $\text{Aut}(G)$  of  $G$  is the set of graph automorphisms preserving the genus function  $g$  and the ordering of the legs.

For generalities on these moduli spaces, see [8], [19], [1] and [17].

**Euler characteristics.** Put  $G_{\mathbb{Q}} = \text{Gal}(\overline{\mathbb{Q}}/\mathbb{Q})$ , and for a prime  $\ell$ , denote by  $\mathbf{Rep}_{\mathbb{Q}_{\ell}}(G_{\mathbb{Q}})$  the abelian category of  $\ell$ -adic Galois representations of  $G_{\mathbb{Q}}$ . Let  $\mathbf{K}_0(\mathbf{Rep}_{\mathbb{Q}_{\ell}}(G_{\mathbb{Q}}))$  denote the Grothendieck ring of this category; it carries a natural  $\lambda$ -ring structure, cf. [27, exposé 5] or [12], the  $\lambda$ -operations enjoying the property:

$$\lambda^m([V]) = [\lambda^m V] \text{ for } m \geq 0.$$

We also recall that a semi-simple  $\ell$ -adic Galois representation  $V$  is determined by the traces  $\text{Tr}(\sigma_p | V)$  of Frobenius elements  $\sigma_p$  on the primes  $p$  at which  $V$  is unramified (see, for instance, [7, Proposition 2.6]).

Let  $\mathcal{M}_{g,n/\overline{\mathbb{Q}}}$  denote the  $\overline{\mathbb{Q}}$ -stack corresponding to  $\mathcal{M}_{g,n}$ , i.e., the generic fiber  $\mathcal{M}_{g,n/\overline{\mathbb{Q}}}$  of  $\mathcal{M}_{g,n} \rightarrow \text{Spec}(\mathbb{Z})$  base changed from  $\mathbb{Q}$  to  $\overline{\mathbb{Q}}$ . The action of the symmetric group  $\mathbb{S}_n$  on  $\mathcal{M}_{g,n}$  induces an isotypic decomposition of the  $\ell$ -adic cohomology  $H_c^i(\mathcal{M}_{g,n/\overline{\mathbb{Q}}}, \mathbb{Q}_{\ell})$  as an  $\mathbb{S}_n$ -module,

$$H_c^i(\mathcal{M}_{g,n/\overline{\mathbb{Q}}}, \mathbb{Q}_{\ell}) \cong \bigoplus_{\lambda \vdash n} H_{c,\lambda}^i(\mathcal{M}_{g,n/\overline{\mathbb{Q}}}, \mathbb{Q}_{\ell})$$

where for an irreducible representation  $V_{\lambda}$  of  $\mathbb{S}_n$  indexed by the partition  $\lambda$  of  $n$ ,

$$H_{c,\lambda}^i(\mathcal{M}_{g,n/\overline{\mathbb{Q}}}, \mathbb{Q}_{\ell}) = V_{\lambda} \otimes \text{Hom}_{\mathbb{S}_n}(V_{\lambda}, H_c^i(\mathcal{M}_{g,n/\overline{\mathbb{Q}}}, \mathbb{Q}_{\ell})).$$

For a partition  $\lambda$  of  $n$ , put

$$\mathbf{e}_{c,\lambda}(\mathcal{M}_{g,n/\overline{\mathbb{Q}}}) = \sum_i (-1)^i [H_{c,\lambda}^i(\mathcal{M}_{g,n/\overline{\mathbb{Q}}}, \mathbb{Q}_{\ell})] \in \mathbf{K}_0(\mathbf{Rep}_{\mathbb{Q}_{\ell}}(G_{\mathbb{Q}}));$$

in addition, we fix throughout a finite field  $\mathbb{F}$  of characteristic different from  $\ell$  and an algebraic closure of it  $\overline{\mathbb{F}}$ , and define similarly the Euler characteristic  $\mathbf{e}_{c,\lambda}(\mathcal{M}_{g,n/\overline{\mathbb{F}}})$ . Let  $\mathcal{V}((g, n)) = H_c^*(\mathcal{M}_{g,n/\overline{\mathbb{Q}}}, \mathbb{Q}_\ell)$ , and set

$$\mathrm{ch}_n(\mathcal{V}((g, n))) := \sum_{\lambda \vdash n} \frac{1}{\dim V_\lambda} \mathbf{e}_{c,\lambda}(\mathcal{M}_{g,n/\overline{\mathbb{Q}}}) s_\lambda$$

see also [4] where this characteristic is computed for small values of  $g$  and  $n$ .

Letting  $F : \mathcal{M}_{g,n/\overline{\mathbb{F}}} \rightarrow \mathcal{M}_{g,n/\overline{\mathbb{F}}}$  denote the Frobenius morphism, we note that by Grothendieck's fixed point formula [2, 3]<sup>2</sup>, the trace of the geometric Frobenius  $F^*$  on the characteristic  $\mathrm{ch}_n$  of the graded  $\mathbb{S}_n$ -module  $H_c^*(\mathcal{M}_{g,n/\overline{\mathbb{F}}}, \mathbb{Q}_\ell)$  is given by

$$t_{g,n}(\mathbf{p}) = t_{g,n}(p_1, p_2, \dots) := \sum_{\lambda \vdash n} \frac{1}{\dim V_\lambda} \mathrm{Tr}(F^* | \mathbf{e}_{c,\lambda}(\mathcal{M}_{g,n/\overline{\mathbb{F}}})) s_\lambda = \frac{1}{n!} \sum_{\sigma \in \mathbb{S}_n} |\mathcal{M}_{g,n/\overline{\mathbb{F}}}^{\sigma F}| p_{c(\sigma)}$$

where  $c(\sigma)$  denotes the cycle type of  $\sigma$ , and if  $c(\sigma) = (1^{\rho(1)}, \dots, n^{\rho(n)})$  then  $p_{c(\sigma)} = p_1^{\rho(1)} \dots p_n^{\rho(n)}$ . Here we have also used Frobenius' formula

$$s_\lambda = \frac{1}{n!} \sum_{\sigma \in \mathbb{S}_n} \chi^\lambda(\sigma) p_{c(\sigma)}$$

$\chi^\lambda$  being the character of  $V_\lambda$ . Since  $|\mathcal{M}_{g,n/\overline{\mathbb{F}}}^{\sigma F}|$  depends only upon the cycle type of  $\sigma$ , we can also write

$$t_{g,n}(\mathbf{p}) = \sum_{\rho \vdash n} |\mathcal{M}_{g,n/\overline{\mathbb{F}}}^{\rho F}| \frac{p_\rho}{z_\rho}$$

where, for convenience, we set  $|\mathcal{M}_{g,n/\overline{\mathbb{F}}}^{\rho F}| := |\mathcal{M}_{g,n/\overline{\mathbb{F}}}^{\sigma F}|$  for  $\sigma \in \mathbb{S}_n$  with  $c(\sigma) = \rho$ , and

$$z_\rho = \prod_{i=1}^n i^{\rho(i)} \rho(i)! \quad \text{if } \rho = (1^{\rho(1)}, \dots, n^{\rho(n)}).$$

Note that, for  $k \geq 1$ , we have

$$\psi_k(t_{g,n})(\mathbf{p}) = \sum_{\rho \vdash n} |\mathcal{M}_{g,n/\overline{\mathbb{F}}}^{\rho F^k}| \frac{p_{k\rho}}{z_\rho}$$

Now the characteristic of the stable  $\mathbb{S}$ -module  $\mathcal{V} = \{\mathcal{V}((g, n))\}$  (that is, the representation  $\mathcal{V}$  of the groupoid  $\mathbb{S} = \coprod_{n \geq 0} \mathbb{S}_n$ ) is defined by the formal Laurent series

$$\mathrm{Ch}(\mathcal{V}) = \sum_{2(g-1)+n > 0} \hbar^{g-1} \mathrm{ch}_n(\mathcal{V}((g, n)))$$

with the coefficients

$$\sum_{n \geq \max\{0, 3-2g\}} \mathrm{ch}_n(\mathcal{V}((g, n))) \in K_0(\mathbf{Rep}_{\mathbb{Q}_\ell}(G_{\mathbb{Q}}))[[h_1, h_2, \dots]].$$

The corresponding generating series of  $t_{g,n}(\mathbf{p})$  will be denoted by  $T(\mathcal{V})$ .

The free modular operad  $\mathbb{M}\mathcal{V} = \{\mathbb{M}\mathcal{V}((g, n))\}$  generated by  $\mathcal{V}$  is obtained by taking  $\mathbb{M}\mathcal{V}((g, n)) = H_c^*(\overline{\mathcal{M}}_{g,n/\overline{\mathbb{Q}}}, \mathbb{Q}_\ell)$ ; the characteristic  $\mathrm{Ch}(\mathbb{M}\mathcal{V})$  and the corresponding generating series  $T(\mathbb{M}\mathcal{V})$  are defined as for  $\mathcal{V}$ .

The connection between the characteristics  $\mathrm{Ch}(\mathcal{V})$  and  $\mathrm{Ch}(\mathbb{M}\mathcal{V})$  is given by the following theorem of Getzler and Kapranov [15, Theorem 8.13]:

**Theorem 2.1.** — *We have*

$$\mathrm{Ch}(\mathbb{M}\mathcal{V}) = \mathrm{Log}(\exp(\Delta)\mathrm{Exp}(\mathrm{Ch}(\mathcal{V}))).$$

Here  $\mathrm{Exp}(-)$  and  $\mathrm{Log}(-)$  are defined as before.

**Integral representation.** Following Getzler and Kapranov, we shall now express  $\mathrm{Ch}(\mathbb{M}\mathcal{V})$  as a formal Fourier transform. The resulting formula is in complete analogy with the formula (2) in Wick's theorem.

<sup>2</sup>The smooth Deligne-Mumford stack  $\mathcal{M}_{g,n}$  is of finite type and has relative dimension  $3(g-1) + n$  over  $\mathrm{Spec}(\mathbb{Z})$ , thus fulfilling the conditions of [2, Theorem 3.1.2].

For a partition  $\rho = (1^{\rho(1)}, 2^{\rho(2)}, \dots)$ , where  $\rho(j) = 0$  for all but finitely many  $j$ , put, as before,  $p_\rho = p_1^{\rho(1)} p_2^{\rho(2)} \dots$ . Let  $\Lambda_{\text{alg}}$  denote the space of *finite* linear combinations of the  $p_\rho$ . On  $\text{Spec}(\Lambda_{\text{alg}} \otimes \mathbb{R}) \cong \mathbb{R}^\infty$  with coordinates  $p_1, p_2, \dots$ , let  $d\mu$  denote the formal Gaussian measure defined by

$$\begin{aligned} d\mu &= \prod_{n-\text{odd}} e^{-p_n^2/2n\hbar^n} \frac{dp_n}{\sqrt{2\pi n\hbar^n}} \prod_{n-\text{even}} e^{-p_n^2/2n\hbar^n + p_n/n\hbar^{n/2}} \frac{dp_n}{e^{1/2n} \sqrt{2\pi n\hbar^n}} \\ &= \text{Exp}(-e_2/\hbar) \prod_{n=1}^{\infty} \frac{dp_n}{e^{\varepsilon_n/2n} \sqrt{2\pi n\hbar^n}} \end{aligned}$$

where  $\varepsilon_n = 0$  or  $1$  according as  $n$  is odd or even. With this measure, for each monomial  $p_\rho = p_1^{\rho(1)} p_2^{\rho(2)} \dots$ , define the integral

$$\begin{aligned} \int_{\mathbb{R}^\infty}^* p_\rho d\mu(\mathbf{p}) &= \prod_{n-\text{odd}} \int_{-\infty}^{\infty} p_n^{\rho(n)} e^{-p_n^2/2n\hbar^n} \frac{dp_n}{\sqrt{2\pi n\hbar^n}} \\ &\cdot \prod_{n-\text{even}} \int_{-\infty}^{\infty} p_n^{\rho(n)} e^{-p_n^2/2n\hbar^n + p_n/n\hbar^{n/2}} \frac{dp_n}{e^{1/2n} \sqrt{2\pi n\hbar^n}}. \end{aligned}$$

Notice that almost all integrals in the products equal 1. We extend  $\int_{\mathbb{R}^\infty}^*$  to a map on  $K_0(\mathbf{Rep}_{\mathbb{Q}_\ell}(G_{\mathbb{Q}}))[[\hbar_1, \hbar_2, \dots]](\hbar)$  by linearity.

With this definition, we have the following interpretation of the formula in Theorem 2.1, see [15, Theorem 8.18]:

**Theorem 2.2.** — *We have*

$$\hbar^{-1} h_2 + \text{Ch}(\mathbb{M}\mathcal{V}) = \text{Log} \left( \int_{\mathbb{R}^\infty}^* \text{Exp}(\hbar^{-1} p_1 q_1 + \text{Ch}(\mathcal{V})) d\mu(\mathbf{p}) \right) \quad (3)$$

where the left-hand side is considered as a function of  $\mathbf{q} = (q_1, q_2, \dots)$  and  $\hbar$ .

For our purposes it will be more convenient to switch the roles of  $\mathbf{p} = (p_1, p_2, \dots)$  and  $\mathbf{q} = (q_1, q_2, \dots)$ , and write (3) in the form

$$\text{Exp}(\text{Ch}(\mathbb{M}\mathcal{V})) = \int_{\mathbb{R}^\infty} e^{\mathcal{K}(\mathbf{p}, \mathbf{q}, \hbar)} d^* \mathbf{q} \quad (4)$$

with  $\mathcal{K}(\mathbf{p}, \mathbf{q}, \hbar)$  defined by

$$\mathcal{K}(\mathbf{p}, \mathbf{q}, \hbar) = - \sum_{m=1}^{\infty} \left\{ (q_m - p_m - \varepsilon_m \hbar^{m/2})^2 / 2m\hbar^m \right\} + \Psi(\text{Ch}(\mathcal{V}))$$

and the measure

$$d^* \mathbf{q} = \prod_{m=1}^{\infty} \frac{dq_m}{\sqrt{2\pi m\hbar^m}}.$$

Formula (4) follows easily by exponentiating (3) and applying the identities  $e_2 = (q_1^2 - q_2)/2$  and  $h_2 = (p_1^2 + p_2)/2$ .

### 3 A semi-classical expansion

To express the coefficients of  $\text{Ch}(\mathbb{M}\mathcal{V})$  in terms of those of  $\text{Ch}(\mathcal{V})$ , both  $\text{Ch}(\mathcal{V})$  and  $\text{Ch}(\mathbb{M}\mathcal{V})$  considered as formal Laurent series in  $\hbar$ , we shall study the integral (4) corresponding to the generating series  $T(\mathcal{V})$  of  $t_{g,n}(\mathbf{p})$  over any finite field  $\mathbb{F}$  of characteristic different from  $\ell$ . Here, we recall that the Euler characteristics  $e_{c,\lambda}$  are elements of the Grothendieck ring  $K_0(\mathbf{Rep}_{\mathbb{Q}_\ell}(G_{\mathbb{Q}}))$ .

In what follows, we shall denote by  $\text{ch}_g(\mathcal{V})$  (resp.  $\text{ch}_g(\mathbb{M}\mathcal{V})$ ) the coefficient of  $\hbar^{g-1}$  in the series  $T(\mathcal{V})$  (resp.  $T(\mathbb{M}\mathcal{V})$ ), i.e.,

$$\text{ch}_g(\mathcal{V})(\mathbf{q}) = \sum_{n \geq \max\{0, 3-2g\}} \sum_{\rho \vdash n} |\mathcal{M}_{g,n/\mathbb{F}}^{\rho F}| \frac{q_\rho}{z_\rho}$$

and

$$\text{ch}_g(\mathbb{M}\mathcal{V})(\mathbf{p}) = \sum_{n \geq \max\{0, 3-2g\}} \sum_{\rho \vdash n} |\overline{\mathcal{M}}_{g,n/\mathbb{F}}^{\rho F}| \frac{p_\rho}{z_\rho}.$$

To make use of Theorem 2.2 of Getzler and Kapranov in the form (4) to obtain formulas expressing the coefficients  $\text{ch}_g(\mathbb{M}\mathcal{V})$  in terms of the coefficients  $\text{ch}_{g'}(\mathcal{V})$  with  $g' \leq g$ , we shall perform a semi-classical expansion of the corresponding integral giving  $T(\mathbb{M}\mathcal{V})$ . The analogous expansion in the context of the usual orbifold Euler characteristics  $\chi(\mathcal{M}_{g,n})$  and  $\chi(\overline{\mathcal{M}}_{g,n})$  is discussed by Bini and Harer in [6, 3.1]. Our arguments are a natural extension of theirs.

### 3.1 The critical points

To apply the principle of stationary phase to the integral giving  $T(\mathbb{M}\mathcal{V})$ , we first need to determine the critical points of the function (still denoted by  $\mathcal{K}(\mathbf{p}, \mathbf{q}, \hbar)$ ) given by

$$\mathcal{K}(\mathbf{p}, \mathbf{q}, \hbar) = - \sum_{m=1}^{\infty} \left\{ (q_m - p_m - \varepsilon_m \hbar^{m/2})^2 / 2m \hbar^m \right\} + \Psi(T(\mathcal{V})).$$

More precisely, for  $m = 1, 2, \dots$ , we have to find the power series

$$\bar{q}_m = \bar{q}_m(\mathbf{p}, \hbar) = \sum_{s=0}^{\infty} \bar{c}_{m,s}(\mathbf{p}) \hbar^s$$

satisfying the system of differential equations

$$\frac{\partial \mathcal{K}}{\partial q_m}(\mathbf{p}, \bar{\mathbf{q}}, \hbar) = 0 \quad (\bar{\mathbf{q}} = (\bar{q}_1, \bar{q}_2, \dots))$$

for  $m = 1, 2, \dots$ , or written explicitly,

$$\bar{q}_m = p_m + \varepsilon_m \hbar^{m/2} + \sum_{k|m} \frac{m}{k} \sum_{g=0}^{\infty} \hbar^{kg+m-k} \frac{\partial \psi_k(\text{ch}_g(\mathcal{V}))}{\partial q_m}(\bar{\mathbf{q}}). \quad (5)$$

Here  $\psi_k(\text{ch}_g(\mathcal{V}))$  is given by

$$\psi_k(\text{ch}_g(\mathcal{V}))(\mathbf{q}) = \sum_{n \geq \max\{0, 3-2g\}} \sum_{\rho \vdash n} |\mathcal{M}_{g,n/\bar{\mathbb{F}}}^{\rho F^k}| \frac{q_{k\rho}}{z_\rho}.$$

For  $m \geq 1$ , define  $c_m(\mathbf{q})$  by

$$c_m(\mathbf{q}) = q_m - \frac{\partial \psi_m(\text{ch}_0(\mathcal{V}))}{\partial q_m}(\mathbf{q}).$$

Note that  $c_m = \psi_m(c_1)$ . If we put  $C(\mathbf{q}) = (c_m(\mathbf{q}))_{m \geq 1}$  and  $\bar{C}(\mathbf{p}) = (\bar{c}_{m,0}(\mathbf{p}))_{m \geq 1}$ , then equating the constant terms (i.e., setting  $\hbar = 0$ ) in both sides of (5), we must have

$$C(\bar{C}(\mathbf{p})) = \mathbf{p}.$$

*Remark.* In [18, Lemma (2.8)], Kisin and Lehrer obtained the formula

$$\frac{|\mathcal{M}_{0,n/\bar{\mathbb{F}}}^{\rho F}|}{z_\rho} = \frac{\prod_{i \geq 1} \binom{s_i(q)/i}{\rho(i)}}{q(q^2-1)} \quad (\text{with } \rho = (1^{\rho(1)}, \dots, n^{\rho(n)}) \text{ and } |\mathbb{F}| = q)$$

where  $s_1(q) = q + 1$ , and  $s_i(q) = \sum_{d|i} \mu\left(\frac{i}{d}\right) q^d$  for  $i \geq 2$ . Accordingly, we can write

$$\text{ch}_0(\mathcal{V})(\mathbf{p}) = \frac{\left\{ \prod_{n \geq 1} (1 + p_n)^{s_n(q)/n} \right\} - 1}{q(q^2-1)} - \frac{p_1^2}{2(q-1)} - \frac{p_1}{q(q-1)} - \frac{p_2}{2(q+1)};$$

see also [13]. Thus we can express

$$c_1(\mathbf{p}) = p_1 - \frac{\partial \text{ch}_0(\mathcal{V})}{\partial p_1}(\mathbf{p}) = \frac{q p_1}{q-1} - \frac{\left\{ \prod_{n \geq 1} (1 + p_n)^{\frac{1}{n} \sum_{d|n} \mu\left(\frac{n}{d}\right) q^d} \right\} - 1}{q(q-1)}. \quad (6)$$

Since  $c_1(\mathbf{q})$  has no constant term, and the linear part of  $C(\mathbf{q})$  is easily seen to be invertible, it follows that  $C(\mathbf{q})$  admits, indeed, a compositional inverse. Note that

$$\psi_m(\bar{c}_{1,0})(C(\mathbf{p})) = p_m \circ \bar{c}_{1,0}(C(\mathbf{p})) = p_m \quad (\text{for all } m \geq 1)$$

and thus  $\bar{c}_{m,0} = \psi_m(\bar{c}_{1,0})$ , that is,  $c_1$  and  $\bar{c}_{1,0}$  are *plethystic* inverses.

**Proposition 3.1.** — *The system (5) has a unique solution of the form*

$$\bar{q}_m(\mathbf{p}, \hbar) = \bar{c}_{m,0}(\mathbf{p}) + O(\hbar^{(m+1-\varepsilon_m)/2})$$

for all  $m \geq 1$ , with  $\varepsilon_m = 0$  or  $1$  according as  $m$  is odd or even. The coefficients  $\bar{c}_{m,m/2}$  of this solution are given by the formula

$$\bar{c}_{m,m/2} = \frac{1 + 2\psi_{\frac{m}{2}}\left(\frac{\partial \text{ch}_0(\mathcal{V})}{\partial p_2}\right)}{1 - \psi_m\left(\frac{\partial^2 \text{ch}_0(\mathcal{V})}{\partial p_1^2}\right)} \circ \bar{c}_{1,0}$$

for all even  $m \geq 2$ .

*Proof.* Splitting the right-hand side of (5) according as  $kg + m - k$  (the exponent of  $\hbar$ ) is zero (i.e.,  $g = 0$  and  $k = m$ ) or not, we rewrite (5) as

$$c_m(\bar{\mathbf{q}}) - p_m = \varepsilon_m \hbar^{m/2} + \sum \left( \frac{m \hbar^{k(g-1)+m}}{k} \right) \frac{\partial \psi_k(\text{ch}_g(\mathcal{V}))}{\partial q_m}(\bar{\mathbf{q}}) \quad (7)$$

the sum being over  $k \mid m$  and  $g \geq 0$  such that  $k(g-1) + m > 0$ . Equating the coefficients of  $\hbar^s$ , for any  $s \geq 1$ , on both sides of (7), we get recursive relations of the form

$$\sum_{r \geq 1} \bar{c}_{mr,s}(\mathbf{p}) \psi_m\left(\frac{\partial c_1}{\partial p_r}\right)(\bar{C}(\mathbf{p})) = R_m(\mathbf{p}) \quad (\text{for all } m \geq 1) \quad (8)$$

where  $R_m(\mathbf{p})$  is an expression involving only coefficients  $\bar{c}_{t,s'}(\mathbf{p})$  ( $t \geq 1$ ) with  $s' < s$ . Notice that the right-hand side of (7) is  $O(\hbar^{(m+1-\varepsilon_m)/2})^3$ ; thus the coefficient of  $\hbar^s$  in  $c_m(\bar{\mathbf{q}})$  must be zero when  $m - \varepsilon_m \geq 2s$ , that is,  $m \geq 2s + 1$ .

We proceed now by induction on  $s$ . Having the coefficients  $\bar{c}_{m,0} = \psi_m(\bar{c}_{1,0})$  already determined, assume that  $\bar{c}_{m,s'}(\mathbf{p})$ , for all  $s' < s$  and  $m \geq 1$ , were also determined, and that, for every  $s' < s$ , we have  $\bar{c}_{m,s'} = 0$  for all  $m \geq 2s' + 1$ . Under these assumptions, we have  $R_m(\mathbf{p}) = 0$  if  $m \geq 2s + 1$ . Indeed, the right-hand side of (7) is certainly  $O(\hbar^{s+1})$ , and the coefficients in  $c_m(\bar{\mathbf{q}})$  getting into  $R_m(\mathbf{p})$  are of the form  $\bar{c}_{mr,s'}$  with  $s' < s$ ; these are all zero, since  $mr > 2s' + 1$ . Let  $\bar{c}_{m,s} := 0$  for  $m \geq 2s + 1$ , hence (8) holds trivially for these values of  $m$ . From (6), it is easy to see that  $\psi_m\left(\frac{\partial c_1}{\partial p_1}\right)(\bar{C}(\mathbf{p}))$  is non-zero for all  $m \geq 1$ , implying that the linear system

$$\sum_{1 \leq r \leq \frac{2s}{m}} \bar{c}_{mr,s}(\mathbf{p}) \psi_m\left(\frac{\partial c_1}{\partial p_r}\right)(\bar{C}(\mathbf{p})) = R_m(\mathbf{p}) \quad (1 \leq m \leq 2s)$$

has a unique solution; thus (8) has a unique solution of the form we asserted. This completes the induction.

To compute the coefficients  $\bar{c}_{m,m/2}(\mathbf{p})$ , we just notice that (8) corresponding to  $s = m/2$  is

$$\bar{c}_{m,m/2}(\mathbf{p}) - \bar{c}_{m,m/2}(\mathbf{p}) \psi_m\left(\frac{\partial^2 \text{ch}_0(\mathcal{V})}{\partial p_1^2}\right)(\bar{C}(\mathbf{p})) = 1 + 2\psi_{\frac{m}{2}}\left(\frac{\partial \text{ch}_0(\mathcal{V})}{\partial p_2}\right)(\bar{C}(\mathbf{p})).$$

Since  $\bar{c}_{r,0} = \psi_r(\bar{c}_{1,0})$ , for all  $r \geq 1$ , it follows that

$$\bar{c}_{m,m/2} = \frac{1 + 2\psi_{\frac{m}{2}}\left(\frac{\partial \text{ch}_0(\mathcal{V})}{\partial p_2}\right)}{1 - \psi_m\left(\frac{\partial^2 \text{ch}_0(\mathcal{V})}{\partial p_1^2}\right)} \circ \bar{c}_{1,0}$$

as claimed. This completes the proof.  $\square$

*Remark.* It is easy to see that the infinite homogeneous linear system associated to (8) (with  $m \geq 2s + 1 \geq 3$ ) has only the trivial solution if we assume that the unknowns  $\bar{c}_{mr,s}(\mathbf{p})$  are subject to a growing condition. Indeed, rewriting (6) as

$$\prod_{n \geq 1} (1 + p_n)^{\frac{1}{n} \sum_{d \mid n} \mu\left(\frac{n}{d}\right) q^d} = -q(q-1)c_1(\mathbf{p}) + q^2 p_1 + 1$$

one finds easily that

$$\psi_m\left(\frac{\partial c_1}{\partial p_1}\right)(\mathbf{p}) = \frac{1 + q^m c_m(\mathbf{p}) - q^m p_m}{1 + p_m} \quad \text{and} \quad \psi_m\left(\frac{\partial c_1}{\partial p_r}\right)(\mathbf{p}) = \frac{s_r(q^m)(q^m(q^m - 1)c_m(\mathbf{p}) - q^{2m} p_m - 1)}{r q^m (q^m - 1)(1 + p_{mr})} \quad (\text{if } r \geq 2).$$

<sup>3</sup>The exponent  $k(g-1) + m$  of  $\hbar$  is at least  $(m+1-\varepsilon_m)/2$ .



Letting  $|p_n| < \delta^n/q^n$  ( $n \geq 1$ ), for a fixed  $0 < \delta < 1$ , we have that

$$\begin{aligned} \left| \psi_m \left( \frac{\partial c_1}{\partial p_r} \right) (\mathbf{p}) \right| &< \frac{q^{mr-m}}{r(q^m-1)(1-q^{-mr})} (1+q^{-m}\delta^m)^{-1} \cdot \prod_{n \geq 1} (1+q^{-mn}\delta^{mn})^{s_n(q^m)/n} \\ &= \frac{q^{mr-m}}{r(q^m-1)(1-q^{-mr})} \cdot \frac{1-q^{-m}\delta^{2m}}{1-\delta^m}. \end{aligned}$$

Similarly, as long as  $\delta$  is not too close to 1 (for instance, one can take  $\delta \leq 1 - \frac{3}{q^{2s+1}-1}$ ), we have the lower bound

$$\left| \psi_m \left( \frac{\partial c_1}{\partial p_1} \right) (\mathbf{p}) \right| > \frac{q^m}{q^m-1} \left( 1 - \frac{1-q^{-m}\delta^{2m}}{q^m(1-\delta^m)(1-q^{-m})} \right) > 0 \quad (\text{for } m \geq 2s+1 \text{ and } r \geq 2).$$

It follows that, for  $|p_n| < \frac{1}{q^n} \left( 1 - \frac{3}{q^{2s+1}-1} \right)^n$ , we have the estimate

$$\left| \psi_m \left( \frac{\partial c_1}{\partial p_r} \right) (\mathbf{p}) / \psi_m \left( \frac{\partial c_1}{\partial p_1} \right) (\mathbf{p}) \right| < \frac{q^{mr-m}}{2r} \quad (m \geq 2s+1 \text{ and } r \geq 2).$$

Our assertion follows now – for instance, from the classical results of von Koch [21]; clearly this gives rise inductively to the solution  $\bar{q}_m(\mathbf{p}, \hbar)$  of (5) in the proposition.

### 3.2 The asymptotic expansion

To obtain the asymptotic expansion of  $T(\mathbb{M}\mathcal{V})$ , we first expand  $\mathcal{K}(\mathbf{p}, \mathbf{q}, \hbar)$  as a power series centered at the unique solution  $\bar{\mathbf{q}} = (\bar{q}_1, \bar{q}_2, \dots)$  of (5). Thus, recalling that

$$\mathcal{K}(\mathbf{p}, \mathbf{q}, \hbar) = - \sum_{m=1}^{\infty} \left\{ (q_m - p_m - \varepsilon_m \hbar^{m/2})^2 / 2m\hbar^m \right\} + \Psi(T(\mathcal{V}))$$

we can express

$$\mathcal{K}(\mathbf{p}, \mathbf{q}, \hbar) = \mathcal{K}(\mathbf{p}, \bar{\mathbf{q}}, \hbar) - \sum_{m=1}^{\infty} \frac{t_m^2}{2m\hbar^m} + \sum_{|\alpha| \geq 2} \partial^\alpha \Psi(T(\mathcal{V}))(\bar{\mathbf{q}}, \hbar) \frac{\mathbf{t}^\alpha}{\alpha!} \quad (\text{with } t_m = q_m - \bar{q}_m \text{ and } \mathbf{t} = \mathbf{q} - \bar{\mathbf{q}}).$$

We set

$$\mathcal{L}(\mathbf{p}, \mathbf{q}, \hbar) = \mathcal{L}(\mathbf{p}, \mathbf{t}, \hbar) := \sum_{|\alpha| \geq 2} \partial^\alpha \Psi(T(\mathcal{V}))(\bar{\mathbf{q}}, \hbar) \frac{\mathbf{t}^\alpha}{\alpha!} - \frac{1}{2} \sum_{m \geq 1} \frac{\partial^2 \Psi(T(\mathcal{V}))}{\partial q_m^2}(\bar{\mathbf{q}}, \hbar) t_m^2$$

so that

$$\mathcal{K}(\mathbf{p}, \mathbf{q}, \hbar) = \mathcal{K}(\mathbf{p}, \bar{\mathbf{q}}, \hbar) - \sum_{m=1}^{\infty} \left\{ \left( 1 - m\hbar^m \frac{\partial^2 \Psi(T(\mathcal{V}))}{\partial q_m^2}(\bar{\mathbf{q}}, \hbar) \right) \frac{t_m^2}{2m\hbar^m} \right\} + \mathcal{L}(\mathbf{p}, \mathbf{q}, \hbar).$$

**Theorem 3.2.** — For coordinates  $t_1, t_2, \dots$  in  $\text{Spec}(\Lambda_{\text{alg}} \otimes \mathbb{R}) \cong \mathbb{R}^\infty$ , set

$$\tilde{t}_m := \left( 1 - m\hbar^m \frac{\partial^2 \Psi(T(\mathcal{V}))}{\partial q_m^2}(\bar{\mathbf{q}}, \hbar) \right)^{-1/2} \sqrt{2m\hbar^m} t_m \quad \text{and} \quad \tilde{\mathbf{t}} = (\tilde{t}_1, \tilde{t}_2, \dots).$$

If  $d\nu(\mathbf{t})$  denotes the formal Gaussian measure

$$d\nu(\mathbf{t}) = \prod_{m=1}^{\infty} e^{-t_m^2} \frac{dt_m}{\sqrt{\pi}}$$

we have

$$\Psi(T(\mathbb{M}\mathcal{V})) = \mathcal{K}(\mathbf{p}, \bar{\mathbf{q}}, \hbar) - \frac{1}{2} \sum_{m=1}^{\infty} \log \left( 1 - m\hbar^m \frac{\partial^2 \Psi(T(\mathcal{V}))}{\partial q_m^2}(\bar{\mathbf{q}}, \hbar) \right) + \log \left( \int_{\mathbb{R}^\infty}^* e^{\mathcal{L}(\mathbf{p}, \tilde{\mathbf{t}}, \hbar)} d\nu(\mathbf{t}) \right). \quad (9)$$

Moreover, as  $\hbar \rightarrow 0^+$ , we have the first-order asymptotic expansion

$$\log \left( \int_{\mathbb{R}^\infty}^* e^{\mathcal{L}(\mathbf{p}, \tilde{\mathbf{t}}, \hbar)} d\nu(\mathbf{t}) \right) = \left( \frac{\left( \frac{\partial^2 \text{ch}_0}{\partial q_1 \partial q_2}(\bar{\mathbf{q}}) \right)^2}{\left( 1 - \frac{\partial^2 \text{ch}_0}{\partial q_1^2}(\bar{\mathbf{q}}) \right) \left( 1 - \frac{\partial^2 \psi_2(\text{ch}_0)}{\partial q_2^2}(\bar{\mathbf{q}}) \right)} + \frac{5}{24} \frac{\left( \frac{\partial^3 \text{ch}_0}{\partial q_1^3}(\bar{\mathbf{q}}) \right)^2}{\left( 1 - \frac{\partial^2 \text{ch}_0}{\partial q_1^2}(\bar{\mathbf{q}}) \right)^3} + \frac{1}{8} \frac{\frac{\partial^4 \text{ch}_0}{\partial q_1^4}(\bar{\mathbf{q}})}{\left( 1 - \frac{\partial^2 \text{ch}_0}{\partial q_1^2}(\bar{\mathbf{q}}) \right)^2} \right) \hbar + O(\hbar^2)$$

where  $\text{ch}_0(\mathbf{q}) = \text{ch}_0(\mathcal{V})(\mathbf{q}) = \sum_{n \geq 3} \sum_{\rho \vdash n} |\mathcal{M}_{0,n/\mathbb{F}}^{\rho F}| \frac{q_\rho}{z_\rho}$ .

*Proof.* By Theorem 2.1, and Theorem 2.2 in the form (4), we have

$$\begin{aligned} \text{Exp}(T(\mathbb{M}\mathcal{V})) &= \exp(\Delta)\text{Exp}(T(\mathcal{V})) = \int_{\mathbb{R}^\infty} e^{\mathcal{K}(\mathbf{p}, \mathbf{q}, \hbar)} d^* \mathbf{q} \\ &= e^{\mathcal{K}(\mathbf{p}, \bar{\mathbf{q}}, \hbar)} \int_{\mathbb{R}^\infty} e^{\mathcal{L}(\mathbf{p}, \mathbf{t}, \hbar)} \exp\left[-\sum_{m=1}^{\infty} \left(1 - m\hbar^m \frac{\partial^2 \Psi(T(\mathcal{V}))}{\partial q_m^2}(\bar{\mathbf{q}}, \hbar)\right) \frac{t_m^2}{2m\hbar^m}\right] d^* \mathbf{t}. \end{aligned}$$

Substituting  $t_m$  by  $(1 - m\hbar^m \frac{\partial^2 \Psi(T(\mathcal{V}))}{\partial q_m^2}(\bar{\mathbf{q}}, \hbar))^{-1/2} \sqrt{2m\hbar^m} t_m$  for all  $m \geq 1$ , and then taking the logarithm, we find that

$$\Psi(T(\mathbb{M}\mathcal{V})) = \mathcal{K}(\mathbf{p}, \bar{\mathbf{q}}, \hbar) - \frac{1}{2} \sum_{m=1}^{\infty} \log\left(1 - m\hbar^m \frac{\partial^2 \Psi(T(\mathcal{V}))}{\partial q_m^2}(\bar{\mathbf{q}}, \hbar)\right) + \log\left(\int_{\mathbb{R}^\infty} e^{\mathcal{L}(\mathbf{p}, \tilde{\mathbf{t}}, \hbar)} d\nu(\mathbf{t})\right).$$

This is the formula for  $\Psi(T(\mathbb{M}\mathcal{V}))$  stated in our theorem.

Now let  $\alpha = (\alpha_1, \dots, \alpha_s)$  be a multi-index with  $|\alpha| \geq 2$ , and put  $\tilde{\alpha} := (1^{\alpha_1}, \dots, s^{\alpha_s})$ . It is clear that when computing the partial derivative  $\partial^\alpha \Psi(T(\mathcal{V}))(\bar{\mathbf{q}}, \hbar)$ , the only terms

$$\frac{\hbar^{k(g-1)}}{k} |\mathcal{M}_{g, n/\bar{g}}^{\rho F^k}| \frac{q_{k\rho}}{z_\rho}$$

in  $\Psi(T(\mathcal{V}))$  contributing nontrivially are among those corresponding to  $k$ 's dividing  $\tilde{\alpha}$  (i.e.,  $k$  divides every part of  $\tilde{\alpha}$ ). Using this, one finds easily that

$$\begin{aligned} \mathcal{L}(\mathbf{p}, \tilde{\mathbf{t}}, \hbar) &= \frac{1}{2\hbar^2} \left\{ \frac{\partial^2 (q_2 \circ \text{ch}_0(\mathcal{V}))}{\partial q_2 \partial q_4}(\bar{\mathbf{q}}) \tilde{t}_2 \tilde{t}_4 + \frac{\partial^3 (q_2 \circ \text{ch}_0(\mathcal{V}))}{\partial q_2^3}(\bar{\mathbf{q}}) \frac{\tilde{t}_2^3}{6} \right\} + \frac{1}{\hbar} \left\{ \frac{\partial^2 \text{ch}_0(\mathcal{V})}{\partial q_1 \partial q_2}(\bar{\mathbf{q}}) \tilde{t}_1 \tilde{t}_2 + \frac{\partial^3 \text{ch}_0(\mathcal{V})}{\partial q_1^3}(\bar{\mathbf{q}}) \frac{\tilde{t}_1^3}{6} \right\} \\ &\quad + \frac{1}{\hbar} \left\{ \frac{\partial^2 \text{ch}_0(\mathcal{V})}{\partial q_1 \partial q_3}(\bar{\mathbf{q}}) \tilde{t}_1 \tilde{t}_3 + \frac{\partial^3 \text{ch}_0(\mathcal{V})}{\partial q_1^2 \partial q_2}(\bar{\mathbf{q}}) \frac{\tilde{t}_1^2 \tilde{t}_2}{2} + \frac{\partial^4 \text{ch}_0(\mathcal{V})}{\partial q_1^4}(\bar{\mathbf{q}}) \frac{\tilde{t}_1^4}{24} \right\} + O(\hbar^{3/2}); \end{aligned}$$

it is also easy to see that

$$\begin{aligned} &\left(1 - m\hbar^m \frac{\partial^2 \Psi(T(\mathcal{V}))}{\partial q_m^2}(\bar{\mathbf{q}}, \hbar)\right)^{-1/2} \\ &= \left(1 - \frac{\partial^2 \psi_m(\text{ch}_0(\mathcal{V}))}{\partial q_m^2}(\bar{\mathbf{q}})\right)^{-1/2} \left(1 + \frac{\delta_{m,1} \frac{\partial^2 \text{ch}_1(\mathcal{V})}{\partial q_1^2}(\bar{\mathbf{q}}) + 2\delta_{m,2} \frac{\partial^2 \text{ch}_0(\mathcal{V})}{\partial q_2^2}(\bar{\mathbf{q}}) \hbar}{1 - \frac{\partial^2 \psi_m(\text{ch}_0(\mathcal{V}))}{\partial q_m^2}(\bar{\mathbf{q}})} \frac{\hbar}{2}\right) + O(\hbar^2). \end{aligned} \quad (10)$$

Here  $\delta_{m,j}$  is the Kronecker delta. Replacing  $\tilde{t}_1, \tilde{t}_2, \tilde{t}_3, \dots$  by their defining expressions and applying (10), we obtain an asymptotic expansion of  $\mathcal{L}(\mathbf{p}, \tilde{\mathbf{t}}, \hbar)$  of the form

$$\begin{aligned} \mathcal{L}(\mathbf{p}, \tilde{\mathbf{t}}, \hbar) &= (A_{1,2} + B_{1,2}\hbar)\sqrt{\hbar} t_1 t_2 + (A_{1,3} + B_{1,3}\hbar)\sqrt{\hbar} t_1^3 + A_{2,4}\hbar t_2 t_4 + A_{2,3}\hbar t_2^3 \\ &\quad + A_{1,3}\hbar t_1 t_3 + A_{1,2,2}\hbar t_1^2 t_2 + A_{1,4}\hbar t_1^4 + O(\hbar^{3/2}) \end{aligned}$$

where, for instance,

$$A_{1,2} = 2\sqrt{2} \cdot \frac{\frac{\partial^2 \text{ch}_0(\mathcal{V})}{\partial q_1 \partial q_2}(\bar{\mathbf{q}})}{\sqrt{\left(1 - \frac{\partial^2 \text{ch}_0(\mathcal{V})}{\partial q_1^2}(\bar{\mathbf{q}})\right)\left(1 - \frac{\partial^2 \psi_2(\text{ch}_0(\mathcal{V}))}{\partial q_2^2}(\bar{\mathbf{q}})\right)}}, \quad A_{1,3} = \frac{\sqrt{2}}{3} \cdot \frac{\frac{\partial^3 \text{ch}_0(\mathcal{V})}{\partial q_1^3}(\bar{\mathbf{q}})}{\left(1 - \frac{\partial^2 \text{ch}_0(\mathcal{V})}{\partial q_1^2}(\bar{\mathbf{q}})\right)^{3/2}}$$

and

$$A_{1,4} = \frac{1}{6} \cdot \frac{\frac{\partial^4 \text{ch}_0(\mathcal{V})}{\partial q_1^4}(\bar{\mathbf{q}})}{\left(1 - \frac{\partial^2 \text{ch}_0(\mathcal{V})}{\partial q_1^2}(\bar{\mathbf{q}})\right)^2}.$$

Finally, by applying the familiar Gaussian integral identity

$$\int_{\mathbb{R}} t^k e^{-t^2} \frac{dt}{\sqrt{\pi}} = \begin{cases} 2^{-k/2} (k-1)!! & \text{if } k \text{ is even} \\ 0 & \text{if } k \text{ is odd} \end{cases}$$

one finds easily that

$$\log \left( \int_{\mathbb{R}^\infty}^* e^{\mathcal{L}(\mathbf{p}, \bar{\mathbf{t}}, \hbar)} d\nu(\mathbf{t}) \right) = \left( \frac{1}{8} A_{1,2}^2 + \frac{15}{16} A_{1,3}^2 + \frac{3}{4} A_{1,4} \right) \hbar + O(\hbar^2)$$

and the asymptotic expansion stated in the theorem follows. This completes the proof.  $\square$

We note that by expressing the exponential  $e^{\mathcal{L}(\mathbf{p}, \bar{\mathbf{t}}, \hbar)}$  as a power series and then integrating, one can write the full asymptotic expansion of  $\Psi(T(\mathbb{M}\mathcal{V}))$ . One should also be able to interpret this asymptotic expansion, as in [6, Proposition 3.6], as an expansion over stable graphs. We shall not pursue this further since the calculations are quite cumbersome. Instead, we shall just use Theorem 3.2 to obtain formulas for the generating series  $\text{ch}_g(\mathbb{M}\mathcal{V})$  when  $g \leq 2$ .

## 4 Equivariant Euler characteristics

In this section we shall apply Theorem 3.2 to express  $\text{ch}_g(\mathbb{M}\mathcal{V})$ , when  $g = 0, 1$  and  $2$ , in terms of  $\text{ch}_{g'}(\mathcal{V})$ ,  $g' \leq g$ . The formulas for  $\text{ch}_0(\mathbb{M}\mathcal{V})$  and  $\text{ch}_1(\mathbb{M}\mathcal{V})$  are not new; they were first obtained by Getzler and Kapranov [15] when  $g = 0$ , and by Getzler [14] when  $g = 1$ .

From now on we adopt the following notation. Put  $\mathbf{a}_g := \text{ch}_g(\mathcal{V})$ ,  $\mathbf{b}_g := \text{ch}_g(\mathbb{M}\mathcal{V})$ , and for  $g, \alpha_1, \dots, \alpha_s \in \mathbb{N}$  ( $\alpha_s \geq 1$ ), write

$$\mathbf{a}_g^{(\alpha_1, \dots, \alpha_s)} = \frac{\partial^{\alpha_1 + \dots + \alpha_s} \text{ch}_g(\mathcal{V})}{\partial q_1^{\alpha_1} \dots \partial q_s^{\alpha_s}} \quad \text{and} \quad \mathbf{b}_g^{(\alpha_1, \dots, \alpha_s)} = \frac{\partial^{\alpha_1 + \dots + \alpha_s} \text{ch}_g(\mathbb{M}\mathcal{V})}{\partial q_1^{\alpha_1} \dots \partial q_s^{\alpha_s}}.$$

In what follows, we shall need the notion of the *Legendre transform* for symmetric functions, see [15, Theorem 7.15]. To define this notion, let  $\mathbb{Q}[x]_*$  denote the set of all formal power series  $f \in \mathbb{Q}[x]$  of the form

$$f(x) = \sum_{s=2}^{\infty} \frac{a_s x^s}{s!} \quad (\text{with } a_2 \neq 0).$$

Let  $\text{rk} : \Lambda \rightarrow \mathbb{Q}[x]$  be the homomorphism defined by  $h_n \mapsto x^n/n!$ , and denote by  $\Lambda_*$  the set of symmetric functions  $f \in \Lambda$  such that  $\text{rk}(f) \in \mathbb{Q}[x]_*$ ; note that  $\text{rk}(f)$  can also be obtained from  $f(p_1, p_2, \dots)$  by setting:  $p_1 = x$ , and  $p_n = 0$  if  $n \geq 2$ . If  $f \in \Lambda_*$ , there is a unique element  $g = \mathcal{L}f \in \Lambda_*$ , called the *Legendre transform* of  $f$ , determined by the formula

$$g \circ \frac{\partial f}{\partial p_1} + f = p_1 \frac{\partial f}{\partial p_1}.$$

With the above notation and terminology, we have (see [15, Theorem 7.17] or [13, Theorem 5.9]):

**Theorem 4.1.** — Put  $f = e_2 - \mathbf{a}_0$  and  $g = h_2 + \mathbf{b}_0$ . Then  $\mathcal{L}f = g$ .

*Proof.* By the definition of the Legendre transform and the identities  $e_2 = (p_1^2 - p_2)/2$  and  $h_2 = (p_1^2 + p_2)/2$ , we have to show that

$$\left( \frac{p_1^2 + p_2}{2} + \mathbf{b}_0 \right) \circ (p_1 - \mathbf{a}_0^{(1)}) = \frac{p_1^2 + p_2}{2} - p_1 \mathbf{a}_0^{(1)} + \mathbf{a}_0.$$

To see this, we first observe that, by (10), the only piece in the right-hand side of (9) contributing negative powers of  $\hbar$  to  $\Psi(T(\mathbb{M}\mathcal{V}))$  is  $\mathcal{K}(\mathbf{p}, \bar{\mathbf{q}}, \hbar)$ . From the definition of  $\mathcal{K}(\mathbf{p}, \mathbf{q}, \hbar)$  it is clear that

$$\mathcal{K}(\mathbf{p}, \bar{\mathbf{q}}, \hbar) = \sum_{m=1}^{\infty} \left\{ - \left( \bar{q}_m - p_m - \varepsilon_m \hbar^{m/2} \right)^2 + 2\psi_m(\mathbf{a}_0)(\bar{\mathbf{q}}) \right\} / 2m\hbar^m + O(1)$$

with  $\psi_m(\mathbf{a}_0)(\bar{\mathbf{q}})$  given explicitly by

$$\psi_m(\mathbf{a}_0)(\bar{\mathbf{q}}) = \sum_{n \geq 3} \sum_{\rho = n} |\mathcal{M}_{0, n/\bar{\rho}}^{\rho F^m}| \frac{\bar{q}_{m\rho}}{z_\rho}.$$

Thus we have:

$$\Psi(T(\mathbb{M}\mathcal{V})) = \sum_{m=1}^{\infty} \left\{ - \left( \bar{q}_m - p_m - \varepsilon_m \hbar^{m/2} \right)^2 + 2\psi_m(\mathbf{a}_0)(\bar{\mathbf{q}}) \right\} / 2m\hbar^m + O(1).$$

Now, since

$$\bar{q}_r(\mathbf{p}, \hbar) = \bar{c}_{r,0}(\mathbf{p}) + O(\hbar^{(r+1-\varepsilon_r)/2}) \quad (\text{for all } r \geq 1)$$

it follows that for every partition  $\rho = (1^{\rho(1)}, \dots, n^{\rho(n)})$ , we have

$$\bar{q}_{m\rho}(\mathbf{p}, \hbar)\hbar^{-m} = \left( \bar{c}_{m,0}^{\rho(1)}(\mathbf{p})\hbar^{-m} + \rho(1)\hbar^{(1-m-\varepsilon_m)/2} \bar{c}_{m,0}^{\rho(1)-1}(\mathbf{p})U_m(\mathbf{p}, \hbar) \right) \prod_{l=2}^n \bar{c}_{m,l,0}^{\rho(l)}(\mathbf{p}) + O(1)$$

where  $U_m(\mathbf{p}, \hbar) = (\bar{q}_m(\mathbf{p}, \hbar) - \bar{c}_{m,0}(\mathbf{p}))\hbar^{-(m+1-\varepsilon_m)/2}$ . Thus we can replace  $\bar{q}_{m\rho}\hbar^{-m}$ , for every  $m$ , by

$$\hbar^{-m} \prod_{l=1}^n \bar{c}_{m,l,0}^{\rho(l)} + U_m \hbar^{(1-m-\varepsilon_m)/2} \rho(1) \bar{c}_{m,0}^{\rho(1)-1} \prod_{l=2}^n \bar{c}_{m,l,0}^{\rho(l)}.$$

This immediately yields

$$\hbar^{-m} \psi_m(\mathbf{a}_0)(\bar{\mathbf{q}}) = \hbar^{-m} \psi_m(\mathbf{a}_0)(\bar{C}(\mathbf{p})) + \hbar^{(1-m-\varepsilon_m)/2} U_m(\mathbf{p}, \hbar) (\bar{c}_{m,0}(\mathbf{p}) - p_m) + O(1).$$

Similarly,

$$\begin{aligned} -\left(\bar{q}_m - p_m - \varepsilon_m \hbar^{m/2}\right)^2 / 2\hbar^m &= -(\bar{c}_{m,0}(\mathbf{p}) - p_m)^2 / 2\hbar^m + \varepsilon_m (\bar{c}_{m,0}(\mathbf{p}) - p_m) / \hbar^{m/2} \\ &\quad - \hbar^{(1-m-\varepsilon_m)/2} U_m(\mathbf{p}, \hbar) (\bar{c}_{m,0}(\mathbf{p}) - p_m) + O(1). \end{aligned}$$

Putting these calculations together, we find that

$$\begin{aligned} \Psi(T(\mathbb{M}\mathcal{V})) &= -\frac{1}{2} \Psi\left(\frac{(\bar{c}_{1,0}(\mathbf{p}) - p_1)^2}{\hbar}\right) + \sum_{m\text{-even}} \frac{1}{m} \psi_m\left(\frac{\bar{c}_{1,0}(\mathbf{p}) - p_1}{\sqrt{\hbar}}\right) \\ &\quad + \left(\Psi\left(\frac{\mathbf{a}_0}{\hbar}\right) \circ \bar{c}_{1,0}\right)(\mathbf{p}) + O(1). \end{aligned}$$

To this asymptotic expansion, we apply the operation  $\Psi^{-1}(-) = \sum_{d \geq 1} \frac{\mu(d)}{d} \psi_d(-)$ . Using the identity

$$\sum_{\substack{d|N \\ \frac{N}{d}\text{-even}}} \mu(d) = \begin{cases} 1 & \text{if } N = 2 \\ 0 & \text{if } N \neq 2 \end{cases}$$

and then equating the coefficients of  $\hbar^{-1}$  in the resulting asymptotic expansion, one obtains the formula:

$$\mathbf{b}_0 = -\frac{1}{2}(\bar{c}_{1,0} - p_1)^2 - \frac{1}{2}p_2 + \frac{1}{2}\psi_2(\bar{c}_{1,0}) + (\mathbf{a}_0 \circ \bar{c}_{1,0}). \quad (11)$$

The theorem follows now by applying  $\circ c_1$  (i.e.,  $\circ (p_1 - \mathbf{a}_0^{(1)})$ ) on the right of (11), and by recalling that  $\bar{c}_{1,0} \circ c_1 = p_1$ .  $\square$

**Corollary 4.2.** — *The symmetric functions*

$$p_1 - \mathbf{a}_0^{(1)} \quad \text{and} \quad p_1 + \mathbf{b}_0^{(1)}$$

are plethystic inverses. Thus  $\bar{c}_{1,0} = p_1 + \mathbf{b}_0^{(1)}$ .

*Proof.* Let  $f$  and  $g$  be as in Theorem 4.1. Then

$$\frac{\partial f}{\partial p_1} = c_1 = p_1 - \mathbf{a}_0^{(1)} \quad \text{and} \quad \frac{\partial g}{\partial p_1} = p_1 + \mathbf{b}_0^{(1)}.$$

Since  $\mathcal{L}f = g$ , the first assertion follows at once from [15, Theorem 7.15 (c)]. The second assertion is an immediate consequence of the fact that  $c_1$  and  $\bar{c}_{1,0}$  are plethystic inverses.  $\square$

The following theorem is the main result of [14]; for a direct combinatorial proof, the reader may consult [24].

**Theorem 4.3.** — *We have*

$$\mathbf{b}_1 = \left\{ \mathbf{a}_1 - \frac{1}{2} \sum_{m=1}^{\infty} \frac{\varphi(m)}{m} \log(1 - \psi_m(\mathbf{a}_0^{(2)})) + \frac{\mathbf{a}_0^{(0,1)}(\mathbf{a}_0^{(0,1)} + 1) + \frac{1}{4}\psi_2(\mathbf{a}_0^{(2)})}{1 - \psi_2(\mathbf{a}_0^{(2)})} \right\} \circ (p_1 + \mathbf{b}_0^{(1)})$$

where  $\varphi(m)$  is Euler's totient function.

*Proof.* The proof is similar to that of Theorem 4.1. By Theorem 3.2 and the definition of  $\mathcal{K}(\mathbf{p}, \mathbf{q}, \hbar)$ , we have that

$$\begin{aligned} \Psi(T(\mathbb{M}\mathcal{V})) &= \Psi(\mathbf{a}_1)(\bar{\mathbf{q}}) + \sum_{m=1}^{\infty} \left\{ -\left(\bar{q}_m - p_m - \varepsilon_m \hbar^{m/2}\right)^2 + 2\psi_m(\mathbf{a}_0)(\bar{\mathbf{q}}) \right\} / 2m\hbar^m \\ &\quad - \frac{1}{2} \sum_{m=1}^{\infty} \log \left( 1 - m\hbar^m \frac{\partial^2 \Psi(T(\mathcal{V}))}{\partial q_m^2}(\bar{\mathbf{q}}, \hbar) \right) + O(\hbar). \end{aligned}$$

By Corollary 4.2, the constant term of  $\Psi(\mathbf{a}_1)(\bar{\mathbf{q}})$  is:

$$\Psi(\mathbf{a}_1)(\bar{\mathbf{q}}(\mathbf{p}, 0)) = \Psi(\mathbf{a}_1)(\bar{C}(\mathbf{p})) = (\Psi(\mathbf{a}_1) \circ \bar{c}_{1,0})(\mathbf{p}) = \Psi(\mathbf{a}_1) \circ (p_1 + \mathbf{b}_0^{(1)}(\mathbf{p})).$$

By applying the linear operation  $\Psi^{-1}$  to this, we obtain the first contribution to  $\mathbf{b}_1$  stated in the theorem.

The second contribution corresponds to

$$-\frac{1}{2} \sum_{m=1}^{\infty} \log \left( 1 - m\hbar^m \frac{\partial^2 \Psi(T(\mathcal{V}))}{\partial q_m^2}(\bar{\mathbf{q}}, \hbar) \right) \Big|_{\hbar=0}. \quad (12)$$

From (10) and Corollary 4.2, we have

$$\log \left( 1 - m\hbar^m \frac{\partial^2 \Psi(T(\mathcal{V}))}{\partial q_m^2}(\bar{\mathbf{q}}, \hbar) \right) \Big|_{\hbar=0} = \log(1 - \psi_m(\mathbf{a}_0^{(2)})) \circ (p_1 + \mathbf{b}_0^{(1)}(\mathbf{p}))$$

for all  $m \geq 1$ . Thus the contribution (12) is given by

$$-\frac{1}{2} \left( \sum_{k=1}^{\infty} \log(1 - \psi_k(\mathbf{a}_0^{(2)})) \right) \circ (p_1 + \mathbf{b}_0^{(1)}(\mathbf{p})).$$

As before, we now apply the operation  $\Psi^{-1}(-) = \sum_{r \geq 1} \frac{\mu(r)}{r} \psi_r(-)$ . Summing first over  $m = kr$ , and using the well-known Möbius inversion formula

$$\sum_{r|m} \frac{\mu(r)}{r} = \frac{\varphi(m)}{m}$$

we find that the contribution to  $\mathbf{b}_1$  corresponding to (12) is

$$-\frac{1}{2} \left\{ \sum_{m=1}^{\infty} \frac{\varphi(m)}{m} \log(1 - \psi_m(\mathbf{a}_0^{(2)})) \right\} \circ (p_1 + \mathbf{b}_0^{(1)}(\mathbf{p})).$$

It remains to compute the constant term of

$$\sum_{m=1}^{\infty} \left\{ -\left(\bar{q}_m - p_m - \varepsilon_m \hbar^{m/2}\right)^2 + 2\psi_m(\mathbf{a}_0)(\bar{\mathbf{q}}) \right\} / 2m\hbar^m.$$

Just as in the proof of Theorem 4.1, for every  $m \geq 1$ , the constant term of the summand is

$$\frac{1}{m} U_{2m}(\mathbf{p}, 0) \psi_m(\mathbf{a}_0^{(0,1)})(\bar{C}(\mathbf{p})) + \frac{\varepsilon_m}{m} \left\{ \frac{1}{2} (-1 + \psi_m(\mathbf{a}_0^{(2)})(\bar{C}(\mathbf{p}))) U_m^2(\mathbf{p}, 0) + U_m(\mathbf{p}, 0) - \frac{1}{2} \right\};$$

by Proposition 3.1, we know that

$$U_m(\cdot, 0) = \bar{c}_{m, m/2} = \frac{1 + 2\psi_{\frac{m}{2}}(\mathbf{a}_0^{(0,1)})}{1 - \psi_m(\mathbf{a}_0^{(2)})} \circ \bar{c}_{1,0} \quad (\text{when } m \text{ is even}).$$

Applying the operation  $\Psi^{-1}$ , it follows that the final contribution to  $\mathbf{b}_1$  is

$$\frac{(\mathbf{a}_0^{(0,1)})^2 + \mathbf{a}_0^{(0,1)} + \frac{1}{4}\psi_2(\mathbf{a}_0^{(2)})}{1 - \psi_2(\mathbf{a}_0^{(2)})} \circ \bar{c}_{1,0}.$$

The formula of  $\mathbf{b}_1$  stated in the theorem follows now from Corollary 4.2.  $\square$

To state the analogous result when  $g = 2$ , let us introduce some notation. For  $m \geq 2$  even, write  $\bar{c}_{m,m/2} = \mathbf{v}_m \circ \bar{c}_{1,0}$ , where, by Proposition 3.1,

$$\mathbf{v}_m = \frac{1 + 2\psi_{\frac{m}{2}}(\mathbf{a}_0^{(0,1)})}{1 - \psi_m(\mathbf{a}_0^{(2)})}.$$

Define

$$\begin{aligned} \mathbf{w}_1 &= \frac{\mathbf{a}_0^{(1,1)}(1 + 2\mathbf{a}_0^{(0,1)})}{(1 - \mathbf{a}_0^{(2)})(1 - \psi_2(\mathbf{a}_0^{(2)}))} + \frac{\mathbf{a}_1^{(1)}}{1 - \mathbf{a}_0^{(2)}} \\ \mathbf{w}_2 &= \frac{\psi_2(\mathbf{a}_1^{(1)}) + 2\mathbf{a}_1^{(0,1)} + 2\mathbf{a}_0^{(1,1)}\mathbf{w}_1}{1 - \psi_2(\mathbf{a}_0^{(2)})} + \frac{2\mathbf{a}_0^{(0,2)}(1 + 2\mathbf{a}_0^{(0,1)})}{(1 - \psi_2(\mathbf{a}_0^{(2)}))^2} + \frac{1}{2} \frac{(1 + 2\mathbf{a}_0^{(0,1)})^2 \psi_2(\mathbf{a}_0^{(3)})}{(1 - \psi_2(\mathbf{a}_0^{(2)}))^3} \\ &\quad + \frac{(1 + 2\psi_2(\mathbf{a}_0^{(0,1)})) \psi_2(\mathbf{a}_0^{(1,1)})}{(1 - \psi_2(\mathbf{a}_0^{(2)}))(1 - \psi_4(\mathbf{a}_0^{(2)}))} \\ \mathbf{w}_3 &= \frac{3\mathbf{a}_0^{(0,0,1)}}{1 - \psi_3(\mathbf{a}_0^{(2)})} \quad \mathbf{w}_4 = \frac{2(1 + 2\mathbf{a}_0^{(0,1)}) \psi_2(\mathbf{a}_0^{(1,1)})}{(1 - \psi_2(\mathbf{a}_0^{(2)}))(1 - \psi_4(\mathbf{a}_0^{(2)}))} + \frac{4\mathbf{a}_0^{(0,0,0,1)}}{1 - \psi_4(\mathbf{a}_0^{(2)})} \quad \mathbf{w}_6 = \frac{3\psi_2(\mathbf{a}_0^{(0,0,1)})}{1 - \psi_6(\mathbf{a}_0^{(2)})} \end{aligned}$$

and  $\mathbf{w}_m = 0$  for  $m \geq 5$ ,  $m \neq 6$ . By solving for the coefficients  $\bar{c}_{m, (m+1+\varepsilon_m)/2}(\mathbf{p})$  in (8), we see that

$$\bar{c}_{m, (m+1+\varepsilon_m)/2} = \mathbf{w}_m \circ \bar{c}_{1,0}.$$

With these quantities we now define

$$\begin{aligned} \bar{\mathbf{b}}_2 &:= \underbrace{\mathbf{a}_2}_{g=2} + \underbrace{\mathbf{w}_1 \mathbf{a}_1^{(1)} + \mathbf{v}_2 \mathbf{a}_1^{(0,1)} + \frac{1}{2} \mathbf{v}_2 \psi_2(\mathbf{a}_1^{(1)}) + \mathbf{v}_2 \mathbf{w}_1 \mathbf{a}_0^{(1,1)} + \frac{1}{2} \mathbf{v}_2^2 \mathbf{a}_0^{(0,2)} + \mathbf{w}_3 \mathbf{a}_0^{(0,0,1)} + \mathbf{v}_4 \mathbf{a}_0^{(0,0,0,1)}}_{g=1} \\ &\quad + \underbrace{\frac{1}{12} \mathbf{v}_2^3 \psi_2(\mathbf{a}_0^{(3)}) + \frac{1}{2} \mathbf{v}_2 \mathbf{v}_4 \psi_2(\mathbf{a}_0^{(1,1)}) + \frac{1}{2} \mathbf{v}_6 \psi_2(\mathbf{a}_0^{(0,0,1)})}_{g=0 \text{ and } m=2} \\ &\quad + \sum_{m\text{-even}} \left\{ \underbrace{\frac{\mathbf{v}_m \mathbf{w}_m \psi_m(\mathbf{a}_0^{(2)}) + \mathbf{w}_{2m} \psi_m(\mathbf{a}_0^{(0,1)})}{m}}_{g=0, m \leq 6} + \underbrace{\frac{\mathbf{w}_m (1 - \mathbf{v}_m)}{m}}_{-(\bar{q}_m - p_m - h^{m/2})^2 / 2m h^m, m \leq 6} \right\} + \sum_{m\text{-odd}} \left\{ \underbrace{\frac{\mathbf{w}_{2m} \psi_m(\mathbf{a}_0^{(0,1)}) + \frac{1}{2} \mathbf{w}_m^2 \psi_m(\mathbf{a}_0^{(2)})}{m}}_{g=0, m \leq 3} - \underbrace{\frac{\mathbf{w}_m^2}{2m}}_{-(\bar{q}_m - p_m)^2 / 2m h^m, m \leq 3} \right\} \\ &\quad + \frac{1}{2} \left( \underbrace{\frac{\mathbf{w}_1 \mathbf{a}_0^{(3)} + \mathbf{v}_2 \mathbf{a}_0^{(2,1)}}{1 - \mathbf{a}_0^{(2)}} + \frac{\mathbf{v}_2 \psi_2(\mathbf{a}_0^{(3)})}{1 - \psi_2(\mathbf{a}_0^{(2)})} + \frac{\mathbf{a}_1^{(2)}}{1 - \mathbf{a}_0^{(2)}}}_{-\frac{1}{2} \sum_{m=1}^{\infty} \log \left( 1 - m h^m \frac{\partial^2 \Psi(T(\mathcal{V}))}{\partial q_m^2}(\bar{\mathbf{q}}, h) \right)} + \frac{\mathbf{a}_0^{(0,2)}}{1 - \psi_2(\mathbf{a}_0^{(2)})} \right) \\ &\quad + \underbrace{\frac{(\mathbf{a}_0^{(1,1)})^2}{(1 - \mathbf{a}_0^{(2)})(1 - \psi_2(\mathbf{a}_0^{(2)}))} + \frac{5}{24} \frac{(\mathbf{a}_0^{(3)})^2}{(1 - \mathbf{a}_0^{(2)})^3} + \frac{1}{8} \frac{\mathbf{a}_0^{(4)}}{(1 - \mathbf{a}_0^{(2)})^2}}_{\log \left( \int_{\mathbb{R}^{\infty}} e^{\mathcal{L}(\mathbf{p}, \bar{\mathbf{t}}, h)} d\nu(\mathbf{t}) \right)}. \end{aligned} \tag{13}$$

**Theorem 4.4.** — *With the above notation, we have  $\mathbf{b}_2 = \bar{\mathbf{b}}_2 \circ (p_1 + \mathbf{b}_0^{(1)})$ .*

*Proof.* The assertion follows, as before, from Theorem 3.2 by computing the coefficient of  $h$  in the right-hand side of (9)<sup>4</sup>. Each contribution to this coefficient corresponds to a piece of  $\bar{\mathbf{b}}_2$ , as indicated in (13). For instance, let us verify the contribution to  $\mathbf{b}_2$  coming from

$$-\frac{1}{2} \sum_{m=1}^{\infty} \log \left( 1 - m h^m \frac{\partial^2 \Psi(T(\mathcal{V}))}{\partial q_m^2}(\bar{\mathbf{q}}, h) \right).$$

<sup>4</sup>The operation  $\Psi^{-1}$  does not have any effect in this case.

By (10), it suffices to compute the coefficient of  $\hbar$  in

$$-\frac{1}{2} \sum_{m=1}^{\infty} \log(1 - \psi_m(\mathbf{a}_0^{(2)})(\bar{\mathbf{q}})) + \frac{\mathbf{a}_1^{(2)}(\bar{\mathbf{q}})}{1 - \mathbf{a}_0^{(2)}(\bar{\mathbf{q}})} \cdot \frac{\hbar}{2} + \frac{\mathbf{a}_0^{(0,2)}(\bar{\mathbf{q}})}{1 - \psi_2(\mathbf{a}_0^{(2)})(\bar{\mathbf{q}})} \cdot \hbar. \quad (14)$$

The logarithmic terms do not contribute to the coefficient when  $m \geq 3$ . If  $m = 1$  we have

$$\mathbf{a}_0^{(2)}(\bar{\mathbf{q}}) = (\mathbf{a}_0^{(2)} \circ \bar{c}_{1,0})(\mathbf{p}) + \hbar \left( (\mathbf{w}_1 \mathbf{a}_0^{(3)} + \mathbf{v}_2 \mathbf{a}_0^{(2,1)}) \circ \bar{c}_{1,0} \right)(\mathbf{p}) + O(\hbar^2)$$

and

$$\psi_2(\mathbf{a}_0^{(2)})(\bar{\mathbf{q}}) = (\psi_2(\mathbf{a}_0^{(2)}) \circ \bar{c}_{1,0})(\mathbf{p}) + \hbar (\mathbf{v}_2 \psi_2(\mathbf{a}_0^{(3)}) \circ \bar{c}_{1,0})(\mathbf{p}) + O(\hbar^2)$$

if  $m = 2$ . Thus the coefficient of  $\hbar$  in (14) is indeed

$$\left( \frac{1}{2} \left( \frac{\mathbf{w}_1 \mathbf{a}_0^{(3)} + \mathbf{v}_2 \mathbf{a}_0^{(2,1)}}{1 - \mathbf{a}_0^{(2)}} + \frac{\mathbf{v}_2 \psi_2(\mathbf{a}_0^{(3)})}{1 - \psi_2(\mathbf{a}_0^{(2)})} + \frac{\mathbf{a}_1^{(2)}}{1 - \mathbf{a}_0^{(2)}} \right) + \frac{\mathbf{a}_0^{(0,2)}}{1 - \psi_2(\mathbf{a}_0^{(2)})} \right) \circ \bar{c}_{1,0}$$

(evaluated at  $\mathbf{p}$ ); we recall that  $\bar{c}_{1,0} = p_1 + \mathbf{b}_0^{(1)}$ .

All the other contributions are computed similarly. □

We note that using the above results, the generating series  $\mathbf{b}_2$  can be effectively computed. Indeed, since  $\mathbf{a}_0 = \text{ch}_0(\mathcal{V})$  is known (see the remark preceding Proposition 3.1), it suffices to compute the generating series  $\mathbf{a}_1$  and  $\mathbf{a}_2$ . An expression in closed form for  $\mathbf{a}_1$  can be read off directly from Getzler's formula [12, Eq. (5.5)] for the generating series of the  $\mathbb{S}_n$ -equivariant Serre characteristic of  $\mathcal{M}_{1,n}$ . To compute  $\mathbf{a}_2$ , let  $\mathcal{A}_2$  denote the moduli space of principally polarized abelian surfaces. Via the Torelli map, we can view  $\mathcal{M}_2$  as an open substack of  $\mathcal{A}_2$ , and set  $\mathcal{A}_{1,1} = \mathcal{A}_2 \setminus \mathcal{M}_2$ . For  $\lambda = (\lambda_1 \geq \lambda_2 \geq 0)$ , we have (see [10]) natural  $\ell$ -adic smooth étale sheaves  $\mathbb{V}_\lambda$  on  $\mathcal{A}_2 \otimes \mathbb{Z}[1/\ell]$  corresponding to irreducible algebraic representations of  $\text{GSp}_4(\mathbb{Q})$ . From the results of [12] (or rather their  $\ell$ -adic realization), we know that computing the  $\mathbb{S}_n$ -equivariant Euler characteristic  $\text{ch}_n(\mathcal{V}(\!(2, n)\!))$  amounts to the same as computing the Euler characteristics

$$\mathbf{e}_c(\mathcal{X}, \mathbb{V}_\lambda) = \sum_i (-1)^i [H_c^i(\mathcal{X}_{/\bar{\mathbb{Q}}}, \mathbb{V}_\lambda)]$$

for  $\mathcal{X} = \mathcal{A}_2$  and  $\mathcal{X} = \mathcal{A}_{1,1}$ , and all  $\lambda$  with  $\lambda_1 + \lambda_2 \leq n$ . A formula for  $\mathbf{e}_c(\mathcal{A}_2, \mathbb{V}_\lambda)$  was conjectured by Faber and van der Geer in [9]. Their conjecture was proved by Weissauer [28] when  $\lambda$  is regular, that is,  $\lambda_1 > \lambda_2 > 0$ , and by Petersen [26] for arbitrary  $\lambda$ . The characteristic  $\mathbf{e}_c(\mathcal{A}_{1,1}, \mathbb{V}_\lambda)$  can be computed, for instance, using the branching formula in [25, Section 3]. Thus taking the trace of Frobenius one obtains a formula expressing  $\mathbf{a}_2$  in terms of traces of Hecke operators on spaces of elliptic and genus 2 vector-valued Siegel modular forms.

## References

- [1] E. Arbarello, M. Cornalba, and P.A. Griffiths: *Geometry of algebraic curves. Volume II*. With a contribution by Joseph Daniel Harris. Grundle Math. Wiss. **268**, Springer, Heidelberg, 2011, xxx+963.
- [2] K.A. Behrend: *The Lefschetz trace formula for algebraic stacks*. Invent. Math. **112** (1993), no. 1, 127–149.
- [3] K.A. Behrend: *Derived  $\ell$ -adic categories for algebraic stacks*. Mem. Amer. Math. Soc. **163** (2003), no. 774, viii+93.
- [4] J. Bergström and O. Tommasi: *The rational cohomology of  $\overline{\mathcal{M}}_4$* . Math. Ann. **338** (2007), no. 1, 207–239.
- [5] D. Bessis, C. Itzykson, and J.B. Zuber: *Quantum field theory techniques in graphical enumeration*. Adv. Appl. Math. **1** (1980) 109–157.
- [6] G. Bini and J. Harer: *Euler characteristics of moduli spaces of curves*. (JEMS) **13** (2011), no. 2, 487–512.
- [7] H. Darmon, F. Diamond, and R. Taylor: *Fermat's last theorem*. In: Elliptic Curves, Modular Forms & Fermat's Last Theorem (Hong Kong, 1993), pp. 2–140, Int. Press, Cambridge, MA, 1997.
- [8] P. Deligne and D. Mumford: *The irreducibility of the space of curves of given genus*. Inst. Hautes Études Sci. Publ. Math. **36** (1969), 75–109.
- [9] C. Faber and G. van der Geer: *Sur la cohomologie des systèmes locaux sur les espaces de modules des courbes de genre 2 et des surfaces abéliennes*, I, II. C. R. Math. Acad. Sci. Paris **338** (2004), no. 5, 381–384 and no. 6, 467–470.

- [10] G. Faltings and C.-L. Chai: *Degeneration of abelian varieties*. With an appendix by David Mumford. *Ergebnisse der Mathematik und ihrer Grenzgebiete (3)* 22, Springer-Verlag, Berlin, 1990, xii+316.
- [11] W. Fulton and S. Lang: *Riemann-Roch algebra*. *Grundlehren der Mathematischen Wissenschaften* 277, Springer-Verlag, New York, 1985, x+203.
- [12] E. Getzler: *Resolving mixed Hodge modules on configuration spaces*. *Duke Math. J.* **96** (1999), no. 1, 175–203.
- [13] E. Getzler: *Operads and moduli spaces of genus 0 Riemann surfaces*. In: *The moduli space of curves* (Texel Island, 1994), *Progr. Math.* 129, pp. 199–230, Birkhäuser Boston, 1995.
- [14] E. Getzler: *The semi-classical approximation for modular operads*. *Comm. Math. Phys.* **194** (1998), no. 2, 481–492.
- [15] E. Getzler and M.M. Kapranov: *Modular operads*. *Compositio Math.* **110** (1998), no. 1, 65–126.
- [16] E. Gorsky: *The equivariant Euler characteristic of moduli spaces of curves*. *Adv. Math.* **250** (2014), 588–595.
- [17] J. Harris and I. Morrison: *Moduli of curves*. *Graduate Texts in Mathematics* 187, Springer-Verlag, New York, 1998, xiv+366.
- [18] M. Kisin and G.I. Lehrer: *Equivariant Poincaré polynomials and counting points over finite fields*. *J. Algebra* **247** (2002), no. 2, 435–451.
- [19] F.F. Knudsen: *The projectivity of the moduli space of stable curves. II. The stacks  $M_{g,n}$* . *Math. Scand.* **52** (1983), no. 2, 161–199.
- [20] D. Knutson:  *$\lambda$ -rings and the representation theory of the symmetric group*. *Lecture Notes in Mathematics*, Vol. 308, Springer-Verlag, Berlin-New York, 1973, iv+203.
- [21] H. von Koch: *On regular and irregular solutions of some infinite systems of linear equations*. In: *Proceedings of the fifth International Congress of Mathematicians, Vol. I* (Cambridge, 22–28 August 1912), pp. 352–365, Cambridge University Press, 1913. Available at: <http://www.mathunion.org/ICM/ICM1912.1/ICM1912.1.ocr.pdf>
- [22] I.G. Macdonald: *Symmetric functions and Hall polynomials*. With contribution by A.V. Zelevinsky and a foreword by Richard Stanley. *Oxford Classic Texts in the Physical Sciences*, Second Edition, Clarendon Press, Oxford University Press, New York, 2015, xii+475.
- [23] S. Mozgovoy: *A computational criterion for the Kac conjecture*. *J. Algebra* **318** (2007), no. 2, 669–679.
- [24] D. Petersen: *A remark on Getzler’s semi-classical approximation*. In: *Geometry and arithmetic*, EMS Ser. Congr. Rep., pp. 309–316, Eur. Math. Soc., Zürich, 2012.
- [25] D. Petersen: *Cohomology of local systems on loci of  $d$ -elliptic abelian surfaces*. *Michigan Math. J.* **62** (2013), no. 4, 705–720.
- [26] D. Petersen: *Cohomology of local systems on the moduli of principally polarized abelian surfaces*. *Pacific J. Math.* **275** (2015), no. 1, 39–61.
- [27] SGA 6: *Théorie des intersections et théorème de Riemann-Roch*. *Lecture Notes in Math.* 225, Séminaire de Géométrie Algébrique du Bois-Marie 1966–1967, dirigé par P. Berthelot, A. Grothendieck et L. Illusie (avec la collaboration de D. Ferrand, J.P. Jouanolou, O. Jussila, S. Kleiman, M. Raynaud et J.P. Serre), Springer-Verlag, Berlin-New York, 1971, xii+700.
- [28] R. Weissauer: *The trace of Hecke operators on the space of classical holomorphic Siegel modular forms of genus two*. Available at [arXiv:0909.1744](https://arxiv.org/abs/0909.1744).