

ON THE DISTRIBUTION OF RANK AND CRANK STATISTICS FOR INTEGER PARTITIONS

NIAN HONG ZHOU

Abstract

Let k be a positive integer and m be an integer. Garvan's k -rank $N_k(m, n)$ is the number of partitions of n into at least $(k - 1)$ successive Durfee squares with k -rank equal to m . In this paper give some asymptotics for $N_k(m, n)$ with $|m| \geq \sqrt{n}$ as $n \rightarrow \infty$. As a corollary, we give a more complete answer for the Dyson's crank distribution conjecture. We also establish some asymptotic formulas for finite differences of $N_k(m, n)$ with respect to m with $m \gg \sqrt{n} \log n$.

1 INTRODUCTION AND STATEMENT OF RESULTS

A partition of an integer n is a sequence of non-increasing positive integers whose sum equals n . As usual, let $p(n)$ be the number of integer partitions of integer n . It is well known that the generating function for $p(n)$ is given by

$$\sum_{n \in \mathbb{Z}} p(n)q^n = \frac{1}{(q; q)_\infty},$$

where $(a; q)_\infty = \prod_{j \geq 0} (1 - aq^j)$ for any $a \in \mathbb{C}$ and $|q| < 1$.

In 1944, F.J. Dyson [1] introduced the rank statistic for integer partitions. The rank of a partition was introduced to explain Ramanujan's famous partition congruences with modulus 5 and 7. If $N(m, n)$ denotes the number of partitions of n with rank m , then it is well known that

$$\sum_{n \geq 0} N(m, n)q^n = \frac{1}{(q; q)_\infty} \sum_{n \geq 1} (-1)^{n-1} q^{n(3n-1)/2 + |m|n} (1 - q^n).$$

However the rank fails to explain Ramanujans congruence modulo 11. Therefore Dyson conjectured the existence of another statistic which he called the crank which would explain Ramanujans partition congruence modulo 11. The crank was found by G.E. Andrews and F.G. Garvan [2, 3]. Let $M(m, n)$ be the number of partitions of n with crank m , then

$$\sum_{n \geq 0} M(m, n)q^n = \frac{1}{(q; q)_\infty} \sum_{n \geq 1} (-1)^{n-1} q^{n(n-1)/2 + |m|n} (1 - q^n).$$

In [4], Dyson gave the following asymptotic formulae conjecture for the crank statistic for integer partitions:

$$M(m, n) \sim \frac{\pi}{4\sqrt{6n}} \operatorname{sech}^2\left(\frac{\pi m}{2\sqrt{6n}}\right) p(n), n \rightarrow +\infty. \quad (1.1)$$

He then asked the question about the precise range of m in (1.1), which holds and about the error term.

The case of fixed m of (1.1) has been investigated by R. Mao [5], K. Bringmann and J. Manschot [6], and B. Kim, E. Kim and J. Seo [7]. In [8], K. Bringmann and J. Dousse proved

Theorem 1.1. ^aFor $|m| \leq \frac{1}{\pi} \sqrt{\frac{n}{6}} \log n$, we have as n tends to infinity,

$$M(m, n) = \frac{\pi}{4\sqrt{6n}} \operatorname{sech}^2\left(\frac{\pi m}{2\sqrt{6n}}\right) p(n) \left(1 + O\left(\frac{|m|^{1/3} + 1}{n^{1/4}}\right)\right). \quad (1.2)$$

Let $N_k(m, n)$ be the number of partitions of n into at least $(k-1)$ successive Durfee squares with k -rank equal to $m \in \mathbb{Z}$ and $k \in \mathbb{Z}_+$. F.G. Garvan [9] proved

$$\sum_{n \geq 0} N_k(m, n) q^n = \frac{1}{(q; q)_\infty} \sum_{n \geq 1} (-1)^{n-1} q^{n((2k-1)n-1)/2 + |m|n} (1 - q^n). \quad (1.3)$$

It is clear that $M(m, n) = N_1(m, n)$ and $N(m, n) = N_2(m, n)$.

In [10], J. Dousse and M.H. Mertens proved that the same asymptotic formula (1.2) holds for $N(m, n)$. D. Parry and R.C. Rhoades [11] proved that

Theorem 1.2. ^b For $\sqrt{n} \ll |m| = o(n^{3/4})$ with $n \rightarrow \infty$, one has

$$N_k(m, n) \sim \frac{\pi}{\sqrt{6n}} \left(e^{\frac{\pi(m+k)}{2\sqrt{6n}}} + e^{-\frac{\pi(m+k)}{2\sqrt{6n}}} \right)^{-2} p(n). \quad (1.4)$$

In this paper, we shall let k be a fixed positive integer and focus on the asymptotics of Garvan k -rank $N_k(m, n)$ with $|m| \geq \sqrt{n}$ and n tends to infinity. Our main result as follows.

Theorem 1.3. Let $F_k(m, n) = p(n - |m| - k + 1) - p(n - |m| - k)$. We have:

1. If $m \geq (n+3)/2 - 2k$ then

$$N_k(m, n) = F_k(m, n).$$

2. If $\sqrt{n} \leq |m| < n/2$ then

$$N_k(m, n) = F_k(m, n) \left(1 + O\left(e^{-\frac{\pi|m|}{\sqrt{6n}}} + \sqrt{n} e^{-n^{1/3}} \right) \right).$$

In particular,

$$N_k(m, n) = \frac{\pi}{\sqrt{6}} \frac{p(n - |m|)}{\sqrt{n - |m|}} \left(1 + O\left(\frac{1}{\sqrt{n - |m|}} + e^{-\frac{\pi|m|}{\sqrt{6n}}} \right) \right)$$

holds for all m such that $|m| \geq \sqrt{n}$ and $n - |m| \rightarrow +\infty$.

From the asymptotics for $M(m, n)$ and $N(m, n)$, that is Theorem 1.1 for $M(m, n)$ and $N(m, n)$, and above Theorem 1.3, we have the following corollary.

Corollary 1.4. Let $k = 1, 2$ and $m, n \in \mathbb{Z}$. We have as $n - |m| \rightarrow +\infty$,

$$\frac{N_k(m, n)}{\widetilde{M}(m, n)} - 1 \ll \frac{1}{\sqrt{n - |m|}} + \min\left(e^{-\frac{\pi|m|}{\sqrt{6n}}}, \frac{|m|^{1/3} + 1}{n^{1/4}} \right), \quad (1.5)$$

^aIn [8], the O -term of (1.1) is $O(|m|^{1/3} n^{-1/4})$, the case of $m = 0$ was missed.

^bIt is need to note that the statement of the main result [11, Theorem 1.2] has some miss. Here is the corrected version.

where

$$\tilde{M}(m, n) = \frac{\pi}{\sqrt{6}} \left(1 + e^{-\frac{\pi|m|}{\sqrt{6n}}}\right)^{-2} \frac{p(n - |m|)}{\sqrt{n - |m|}}.$$

In particular, as $n - |m| \rightarrow +\infty$,

$$N_k(m, n) \sim \tilde{M}(m, n).$$

From Theorem 1.3 and Corollary 1.4, we shall give a more complete answer for the Dyson's crank distribution conjecture.

Theorem 1.5. *Dyson's asymptotic formulae (1.1) is valid if and only if $|m| = o(n^{3/4})$ as n tends to infinity. More precisely,*

$$\frac{M(m, n)}{p(n)} = \frac{\pi}{4\sqrt{6n}} \operatorname{sech}^2\left(\frac{\pi m}{2\sqrt{6n}}\right) \left(1 + O\left(\min\left(e^{-\frac{\pi|m|}{\sqrt{6n}}}, \frac{|m|^{1/3} + 1}{n^{1/4}}\right) + \frac{m^2}{n^{3/2}}\right)\right).$$

Remark 1.1. The error term of Theorem 1.5 smaller than that in Theorem 1.1 when $|m| > \sqrt{n}(\log n - 4 \log \log n) / (\pi \sqrt{6})$ as $n \rightarrow \infty$.

As another application of Theorem 1.3, by a straightforward calculation, we obtain the following asymptotic result for finite differences of $N_k(n, m)$ with respect to m with $m \gg \sqrt{n} \log n$.

Corollary 1.6. *Let $m, n, r, k \in \mathbb{N}$ and r, k be fixed. If $m \geq c \sqrt{n} \log n$ with $c > \frac{(r+1)\sqrt{6}}{2\pi}$ be fixed and $n - m \rightarrow \infty$, then*

$$\sum_{j=0}^r (-1)^j \binom{r}{j} N_k(m + j, n) \sim \left(\frac{\pi}{\sqrt{6(n-m)}}\right)^{r+1} p(n-m). \quad (1.6)$$

In particular,

$$N_k(m, n) - N_k(m + 1, n) \sim \frac{\pi^2}{\sqrt{6(n-m)}} p(n-m)$$

holds for $m \geq c_1 \sqrt{n} \log n$ with $c_1 > \sqrt{6}/\pi$ be fixed and $n - m \rightarrow \infty$.

Remark 1.2. We can improve (1.6) and give the size of the error term by Theorem 1.3. We note that the monotonicity properties of $N(m, n)$ have been investigated by S.H. Chan and R. Mao [12].

The proof of Theorem 1.3 is based on the generating function (1.3) of $N_k(m, n)$ and the Hardy-Ramanujan asymptotic formula for $p(n)$. We present the relation between $N_k(m, n)$ and $p(n)$, and some basic properties of $p(n)$ in Section 2. In Section 3, we prove the main results of this paper.

Acknowledgements. The author would like to thank the anonymous referees for their very helpful comments and suggestions. The author also thank Professor Zhi-Guo Liu for his consistent encouragement.

2 PRELIMINARIES

2.1 BASIC PROPERTIES FOR GARVAN k -RANKS

A key ingredient of our asymptotic results is find the connection $N_k(m, n)$ with $p(n)$. Note that $N_k(m, n) = N_k(-m, n)$, we just need consider the case of $m \geq 0$. We prove the following representation for $N_k(m, n)$.

Proposition 2.1. *Let $N_k(m, n)$ be defined as (1.3), integers $n > 0$ and $m \geq 0$. We have*

$$N_k(m, n) = \sum_{\ell \geq 1} (-1)^{\ell-1} F_k(\ell; m, n),$$

where

$$F_k(\ell; m, n) = p(n - m\ell - Q_k(\ell)) - p(n - m\ell - Q_k(\ell) - \ell)$$

with $Q_k(\ell) = (k - 1/2)\ell^2 - \ell/2$.

Proof. For $m \geq 0$, from

$$\sum_{n \geq 0} N_k(m, n) q^n = \frac{1}{(q; q)_\infty} \sum_{\ell \geq 1} (-1)^{\ell-1} q^{\ell((2k-1)\ell-1)/2+m\ell} (1 - q^\ell)$$

we have

$$\begin{aligned} \sum_{n \geq 0} N_k(m, n) q^n &= \sum_{\ell \geq 1} \sum_{r \in \mathbb{Z}} (-1)^{\ell-1} p(r) q^{r+\ell((2k-1)\ell-1)/2+m\ell} (1 - q^\ell) \\ &= \sum_{n \geq 1} q^n \left(\sum_{\substack{\ell \geq 1, r \in \mathbb{Z} \\ \frac{\ell((2k-1)\ell-1)}{2} + m\ell + r = n}} - \sum_{\substack{\ell \geq 1, \ell \in \mathbb{Z} \\ \frac{\ell((2k-1)\ell+1)}{2} + m\ell + r = n}} \right) (-1)^{\ell-1} p(r). \end{aligned}$$

Then, comparing the coefficients of q on both sides we complete the proof. \square

We need the following Hardy–Ramanujan asymptotic result for $p(n)$, which can be found in [13].

Lemma 2.2. *We have for $n \in \mathbb{Z}_+$,*

$$p(n) = \frac{e^{B\sqrt{n-1/24}}}{4\sqrt{3}(n-1/24)} \left(1 - \frac{1}{B\sqrt{n-1/24}} \right) + O\left(n^{-1}e^{B\sqrt{n}/2}\right),$$

where and throughout this section $B = 2\pi/\sqrt{6}$.

We also need the following approximation for $p(n+r)$ with $r = O(\sqrt{n})$.

Lemma 2.3. *Let integer $r = O(\sqrt{n})$ and n be a positive integer. We have as $n \rightarrow \infty$,*

$$\frac{p(n+r) - p(n)}{p(n)} = \frac{Br}{\sqrt{n}} + O\left(\frac{1 + |r| + |r|^2}{n}\right).$$

In particular, as $n \rightarrow \infty$,

$$\frac{p(n+r) - p(n)}{p(n)} \ll \frac{|r| + 1}{\sqrt{n}}.$$

Proof. It is clear that

$$\begin{aligned} \frac{p(n+r)}{p(n)} &= e^{B(\sqrt{n+r-1/24}-\sqrt{n-1/24})} \left(1 + O\left(\frac{|r|}{n}\right)\right) \\ &= e^{B\sqrt{n}\left(\frac{r-1/24}{2n}-\frac{-1/24}{2n}+O\left(\frac{|r|^2+1}{n^2}\right)\right)} \left(1 + O\left(\frac{|r|}{n}\right)\right) \\ &= \left(1 + \frac{Br}{2\sqrt{n}} + O\left(\frac{|r|^2 + |r| + 1}{n}\right)\right) \left(1 + O\left(\frac{|r|}{n}\right)\right). \end{aligned}$$

by generalized binomial theorem, which completes the proof of the lemma. \square

3 THE PROOF OF THE RESULTS OF THIS PAPER

We prove Theorem 1.3 in Subsection 3.1, Corollary 1.4 and Theorem 1.5 in Subsection 3.2, and Corollary 1.6 in the last subsection of this section.

3.1 THE PROOF OF THEOREM 1.3

Let $m \geq 0$ and $n \geq k - 1$. From Proposition 2.1, it is easy to prove that

$$N_k(m, n) = F_k(1; m, n) = F_k(m, n) \tag{3.1}$$

for $m \geq (n+3)/2 - 2k$. For $n/(4k) \leq m \leq n/2$,

$$\begin{aligned} E_k(m, n) &:= N_k(m, n) - F_k(m, n) \\ &\ll \sum_{2 \leq \ell \leq n/m} p(n - m\ell) \ll p(n - 2m) \ll p(n - m - \lfloor n/(4k) \rfloor) \\ &\ll n^{-1} e^{B\sqrt{n-m-n/(4k)+O(1)}} \ll e^{-n^{1/3}} p(n - m), \end{aligned}$$

by use Lemma 2.2, the asymptotics for $p(n)$. For $n^{1/2} \leq m \leq n/(4k)$, we note that

$$0 \leq m\ell + Q_k(\ell) \leq \frac{n}{2k} + k\left(\frac{n}{2km}\right)^2 + O\left(\frac{n}{m}\right) \leq \frac{3}{4k}n + O(\sqrt{n}),$$

if $0 \leq \ell \leq n/(2km)$. Then by Lemma 2.3 we have

$$\begin{aligned} E_k(m, n) &:= N_k(m, n) - F_k(m, n) \\ &= \sum_{2 \leq \ell \leq n/(2km)} (-1)^{\ell-1} F_k(\ell; m, n) + O\left(\sum_{\ell > n/(2km)} p(n - \ell m)\right) \\ &\ll \sum_{2 \leq \ell \leq n/(2km)} p(n - m\ell - Q_k(\ell)) \frac{\ell + 1}{\sqrt{n - m\ell - Q_k(\ell)}} + \frac{n}{m} p(n - \lfloor n/(2k) \rfloor) \\ &\ll \sum_{2 \leq \ell \leq n/(2km)} \frac{\ell}{\sqrt{n}} p(n - m\ell) + \sqrt{n} p(n - \lfloor n/(2k) \rfloor). \end{aligned}$$

By use the asymptotics for $p(n)$ and $\sqrt{n} \leq m \leq n/(4k)$, we further have

$$\begin{aligned}
 E_k(m, n) &\ll \sum_{2 \leq \ell \leq n/(2km)} \frac{\ell}{n^{3/2}} e^{B\sqrt{n-m\ell}} + n^{-1/2} e^{B\sqrt{n-\lfloor n/(2k) \rfloor}} \\
 &\ll \frac{e^{B\sqrt{n-m}}}{n^{3/2}} \sum_{2 \leq \ell \leq n/(2m)} \ell e^{B(\sqrt{n-m\ell} - \sqrt{n-m})} + \frac{e^{B\sqrt{n-m+n/(2k)+O(1)}}}{\sqrt{n}} \\
 &\ll \frac{p(n-m)}{\sqrt{n}} \sum_{2 \leq \ell \leq n/(2m)} \ell e^{-\frac{B(\ell-1)m}{2\sqrt{n}}} + n^{-1/2} e^{B\sqrt{n-m-n/(4k)+O(1)}} \\
 &\ll \frac{p(n-m)}{\sqrt{n}} + \frac{e^{B\sqrt{n/2}}}{\sqrt{n}} \ll \left(\frac{e^{-\frac{Bm}{2\sqrt{n}}}}{\sqrt{n}} + e^{-n^{1/3}} \right) p(n-m).
 \end{aligned}$$

Further, from Lemma 2.3 we have as $n - |m| \rightarrow +\infty$,

$$\begin{aligned}
 F_k(m, n) &= p(n - (|m| + k) + 1) - p(n - (|m| + k)) \\
 &= \frac{B}{2} \frac{p(n - |m|)}{\sqrt{n - |m|}} \left(1 + O\left(\frac{1}{\sqrt{n - |m|}} \right) \right). \tag{3.2}
 \end{aligned}$$

Therefore, for $\sqrt{n} \ll m < n/2$, from above we find that

$$\frac{N_k(m, n) - F_k(m, n)}{F_k(m, n)} \ll e^{-\frac{B|m|}{2\sqrt{n}}} + \sqrt{n} e^{-n^{1/3}}, \tag{3.3}$$

which completes the proof of Theorem 1.3 by substitute $B = 2\pi/\sqrt{6}$.

3.2 THE PROOF OF COROLLARY 1.4 AND THEOREM 1.5

3.2.1 The proof of Corollary 1.4

For $m = o(n^{3/4})$, by Lemma 2.2 it is easy to find that as $n \rightarrow \infty$,

$$\frac{p(n - |m|)}{\sqrt{n - |m|}} = \frac{p(n)}{\sqrt{n}} e^{-\frac{B|m|}{2\sqrt{n}}} \left(1 + O\left(\frac{n + |m|^2}{n^{3/2}} \right) \right)$$

Hence we have for $m = o(n^{3/4})$,

$$\begin{aligned}
 \tilde{M}(m, n) &= \frac{\pi}{\sqrt{6}} \left(1 + e^{-\frac{\pi|m|}{\sqrt{6n}}} \right)^{-2} \frac{p(n - |m|)}{\sqrt{n - |m|}} \\
 &= \frac{\pi}{4\sqrt{6n}} \operatorname{sech}^2\left(\frac{\pi|m|}{2\sqrt{6n}} \right) p(n) \left(1 + O\left(\frac{n + |m|^2}{n^{3/2}} \right) \right). \tag{3.4}
 \end{aligned}$$

From Theorem 1.1 for $M(m, n)$ and $N(m, n)$ and above we have for $|m| \leq (\sqrt{n} \log n)/(\pi\sqrt{6})$,

$$\tilde{M}(m, n) = N_k(m, n) \left(1 + O\left(\frac{|m|^{1/3} + 1}{n^{1/4}} \right) \right).$$

For $|m| \gg \sqrt{n} \log n$, we have

$$\begin{aligned}\tilde{M}(m, n) &= \left(1 + O\left(e^{-\frac{\pi|m|}{\sqrt{6n}}}\right)\right) \frac{\pi}{\sqrt{6}} \frac{p(n-|m|)}{\sqrt{n-|m|}} \\ &= N_k(m, n) \left(1 + O\left(\frac{1}{\sqrt{n-|m|}} + e^{-\frac{\pi|m|}{\sqrt{6n}}}\right)\right)\end{aligned}$$

by Theorem 1.3. From above we complete the proof of Corollary 1.4.

3.2.2 The proof of Theorem 1.5

It is clear that Theorem 1.5 follows from Theorem 1.3, Corollary 1.4, (3.4) and the following lemma by substitute $B = 2\pi/\sqrt{6}$.

Lemma 3.1. *Let $0 \leq m \leq n$. Then as $n \rightarrow \infty$,*

$$\frac{p(n-m+1) - p(n-m)}{p(n)} \sim \frac{B}{8\sqrt{n}} \operatorname{sech}^2\left(\frac{Bm}{4\sqrt{n}}\right) \quad (3.5)$$

if and only if $1/m = o(n^{-1/2})$ and $m = o(n^{3/4})$.

Proof. For $n \geq m \geq 8n/9$, we have

$$\frac{p(n-m+1) - p(n-m)}{p(n)} \ll \frac{e^{B\sqrt{n}/9}}{n^{-1}e^{B\sqrt{n}}} = ne^{-\frac{2B}{3}\sqrt{n}}$$

by Lemma 2.2 and

$$\frac{B}{8\sqrt{n}} \operatorname{sech}^2\left(\frac{Bm}{4\sqrt{n}}\right) \gg n^{-1}e^{-\frac{Bm}{2\sqrt{n}}} \gg n^{-1}e^{-\frac{B}{2}\sqrt{n}}$$

by the definition of $\operatorname{sech}(x)$. Therefore, if (3.5) is valid then $m < 8n/9$. If $0 \leq m \leq 8n/9$ then

$$\frac{p(n-m+1) - p(n-m)}{p(n)} = \left(1 + O\left(\frac{1}{\sqrt{n}}\right)\right) \frac{Be^{-\frac{Bm}{\sqrt{n-m+\sqrt{n}}}}}{2\sqrt{n}} \left(1 - \frac{m}{n}\right)^{-3/2} \quad (3.6)$$

by Lemma 2.2 and (3.2), and note that

$$\frac{B}{8\sqrt{n}} \operatorname{sech}^2\left(\frac{Bm}{4\sqrt{n}}\right) = \frac{B}{2\sqrt{n}} e^{-\frac{Bm}{2\sqrt{n}}} \frac{1}{\left(1 + e^{-\frac{Bm}{8\sqrt{n}}}\right)^2}. \quad (3.7)$$

Hence clearly, if (3.5) is valid then $m = o(n)$. Let $0 \leq m = o(n)$, from (3.6) we have

$$\frac{p(n-m+1) - p(n-m)}{p(n)} = \left(1 + O\left(\frac{1}{\sqrt{n}}\right)\right) \frac{B}{2\sqrt{n}} e^{-\frac{Bm}{2\sqrt{n}} - \frac{m^2}{8n^{3/2}}(1+o(1))}. \quad (3.8)$$

by generalized binomial theorem. Combining (3.7) and (3.8) we immediately obtain that (3.5) is valid if and only if $1/m = o(n^{-1/2})$ and $m = o(n^{3/4})$. \square

3.3 THE PROOF OF THE COROLLARY 1.6

First of all, let $m \geq \sqrt{n}$ and $n - m$ tends to infinity. We have

$$\begin{aligned} N_k(m, n) &= F_k(m, n) \left(1 + O\left(e^{-\frac{\pi m}{\sqrt{6n}}} + e^{-n^{1/3}} \right) \right) \\ &= p(n - (m + k) + 1) - p(n - (m + k)) + R_k(m, n) \end{aligned} \quad (3.9)$$

with

$$R_k(m, n) \ll \left(e^{-\frac{\pi m}{\sqrt{6n}}} + \sqrt{n} e^{-n^{1/3}} \right) p(n - m)$$

by Theorem 1.3. Now, from [14, Equ. (1.2)] we have for $r \in \mathbb{Z}_{\geq 0}$ be fixed and as $N \rightarrow \infty$

$$\Delta_N^r p(N) = \left(\frac{\pi}{\sqrt{6N}} \right)^r p(N) \left(1 + O\left(\frac{1}{\sqrt{N}} \right) \right), \quad (3.10)$$

where for a function $f(x)$,

$$\Delta_x^0 f(x) := f(x), \quad \Delta_x f(x) := f(x) - f(x - 1)$$

and

$$\Delta_x^\ell p(x) := \Delta_x(\Delta_x^{\ell-1} p(x)) \text{ for } \ell \in \mathbb{N}$$

is the backward difference. Hence clearly,

$$(-1)^r \Delta_m^r N_k(m + r, n) = \sum_{j=0}^r (-1)^j \binom{r}{j} N_k(m + j, n). \quad (3.11)$$

By (3.9) we obtain that

$$\begin{aligned} \Delta_m^r N_k(m + r, n) &= -\Delta_m^{r+1} p(n - ((m + r) + k)) \\ &\quad + O_{k,r} \left(\left(e^{-\frac{\pi m}{\sqrt{6n}}} + e^{-\frac{\pi \sqrt{n/6}}{5}} \right) \sum_{j=0}^{r+1} p(n - m + j) \right). \end{aligned} \quad (3.12)$$

Moreover, we have

$$\Delta_m^{r+1} p(n - ((m + r) + k)) = (-1)^{r+1} \Delta_j^{r+1} p(j) \Big|_{j=n-m-k+1}. \quad (3.13)$$

Using (3.10)–(3.13) and Lemma 2.2, it is clear that

$$\begin{aligned} \left(\frac{\sqrt{6(n-m)}}{\pi} \right)^{r+1} \frac{(-1)^r \Delta_m^r N_k(m + r, n)}{p(n-m)} - 1 &\ll_{k,r} \frac{1}{\sqrt{n-m}} + \frac{(n-m)^{\frac{r+1}{2}}}{e^{\frac{\pi m}{\sqrt{6n}}}} \\ &\ll \frac{1}{\sqrt{n-m}} + n^{\frac{r+1}{2}} e^{-\frac{\pi m}{\sqrt{6n}}} \end{aligned}$$

and the proof of the corollary follows.

REFERENCES

- [1] F. J. Dyson. Some guesses in the theory of partitions. *Eureka*, (8):10–15, 1944.
- [2] George E. Andrews and F. G. Garvan. Dyson’s crank of a partition. *Bull. Amer. Math. Soc. (N.S.)*, 18(2):167–171, 1988.
- [3] F. G. Garvan. New combinatorial interpretations of Ramanujan’s partition congruences mod 5, 7 and 11. *Trans. Amer. Math. Soc.*, 305(1):47–77, 1988.
- [4] Freeman J. Dyson. Mappings and symmetries of partitions. *J. Combin. Theory Ser. A*, 51(2):169–180, 1989.
- [5] Renrong Mao. Asymptotic inequalities for k -ranks and their cumulation functions. *J. Math. Anal. Appl.*, 409(2):729–741, 2014.
- [6] Kathrin Bringmann and Jan Manschot. Asymptotic formulas for coefficients of inverse theta functions. *Commun. Number Theory Phys.*, 7(3):497–513, 2013.
- [7] Byungchan Kim, Eunmi Kim, and Jeehyeon Seo. Asymptotics for q -expansions involving partial theta functions. *Discrete Math.*, 338(2):180–189, 2015.
- [8] Kathrin Bringmann and Jehanne Dousse. On Dyson’s crank conjecture and the uniform asymptotic behavior of certain inverse theta functions. *Trans. Amer. Math. Soc.*, 368(5):3141–3155, 2016.
- [9] Frank G. Garvan. Generalizations of Dyson’s rank and non-Rogers-Ramanujan partitions. *Manuscripta Math.*, 84(3-4):343–359, 1994.
- [10] Jehanne Dousse and Michael H. Mertens. Asymptotic formulae for partition ranks. *Acta Arith.*, 168(1):83–100, 2015.
- [11] Daniel Parry and Robert C. Rhoades. On Dyson’s crank distribution conjecture and its generalizations. *Proc. Amer. Math. Soc.*, 145(1):101–108, 2017.
- [12] Song Heng Chan and Renrong Mao. Inequalities for ranks of partitions and the first moment of ranks and cranks of partitions. *Adv. Math.*, 258:414–437, 2014.
- [13] G. H. Hardy and S. Ramanujan. Asymptotic Formulae in Combinatory Analysis. *Proc. London Math. Soc. (2)*, 17:75–115, 1918.
- [14] A. M. Odlyzko. Differences of the partition function. *Acta Arith.*, 49(3):237–254, 1988.

SCHOOL OF MATHEMATICAL SCIENCES, EAST CHINA NORMAL UNIVERSITY
 500 DONGCHUAN ROAD, SHANGHAI 200241, PR CHINA
nianhongzhou@outlook.com