

# Supercongruences for the $(p - 1)$ th Apéry number

Ji-Cai Liu<sup>1</sup> and Chen Wang<sup>2</sup>

<sup>1</sup>Department of Mathematics, Wenzhou University, Wenzhou 325035, PR China  
jcliu2016@gmail.com

<sup>2</sup>Department of Mathematics, Nanjing University, Nanjing 210093, PR China  
chenwjsnu@163.com

**Abstract.** In this paper, we prove two conjectural supercongruences on the  $(p - 1)$ th Apéry number, which were recently proposed by Z.-H. Sun.

*Keywords:* Apéry numbers; supercongruences; Bernoulli numbers

*MR Subject Classifications:* Primary 11A07; Secondary 11B68, 05A19

## 1 Introduction

In 1979, Apéry [1] introduced the following numbers

$$A_n = \sum_{k=0}^n \binom{n}{k}^2 \binom{n+k}{k}^2 \quad \text{and} \quad A'_n = \sum_{k=0}^n \binom{n}{k}^2 \binom{n+k}{k},$$

in his ingenious proof of the irrationality of  $\zeta(2)$  and  $\zeta(3)$ . These numbers are now known as Apéry numbers. Since the appearance of these numbers, some interesting arithmetic properties have been gradually discovered. For instance, Beukers [2] showed that for primes  $p \geq 5$  and  $m, r \in \mathbb{N}$ ,

$$\begin{aligned} A_{mp^r-1} &\equiv A_{mp^{r-1}-1} \pmod{p^{3r}}, \\ A'_{mp^r-1} &\equiv A'_{mp^{r-1}-1} \pmod{p^{3r}}. \end{aligned}$$

In 2012, Z.-W. Sun [14] proved that for any prime  $p \geq 5$ ,

$$\sum_{k=0}^{p-1} (2k+1)A_k \equiv p + \frac{7}{6}p^4 B_{p-3} \pmod{p^5}.$$

Here the  $n$ th Bernoulli number  $B_n$  is defined as

$$\frac{x}{e^x - 1} = \sum_{n=0}^{\infty} B_n \frac{x^n}{n!}.$$

In the past two decades, some interesting congruence properties for Apéry numbers and similar numbers have been widely studied (see, for example, [2–6, 9, 12, 14–16]).

Our interest concerns the following two conjectural supercongruences on the  $(p - 1)$ th Apéry number, which were recently proposed by Z.-H. Sun [12, Conjecture 2.1 & 2.2].

**Conjecture 1.1** (*Z.-H. Sun*) For any prime  $p \geq 5$ , we have

$$A_{p-1} \equiv 1 + \frac{2}{3}p^3 B_{p-3} \pmod{p^4}, \quad (1.1)$$

$$A'_{p-1} \equiv 1 + \frac{5}{3}p^3 B_{p-3} \pmod{p^4}. \quad (1.2)$$

The aim of this paper is to prove (1.1) and (1.2) by establishing their generalizations.

**Theorem 1.2** Let  $p \geq 7$  be a prime. Then

$$A_{p-1} \equiv 1 + p^3 \left( \frac{4}{3}B_{p-3} - \frac{1}{2}B_{2p-4} \right) + \frac{1}{9}p^4 B_{p-3} \pmod{p^5}. \quad (1.3)$$

**Theorem 1.3** Let  $p \geq 7$  be a prime. Then

$$A'_{p-1} \equiv 1 + p^3 \left( \frac{10}{3}B_{p-3} - \frac{5}{4}B_{2p-4} \right) + \frac{5}{18}p^4 B_{p-3} \pmod{p^5}. \quad (1.4)$$

Letting  $k = 1$  and  $b = p - 3$  in the following Kummer's congruence (see [11, page 193]):

$$\frac{B_{k(p-1)+b}}{k(p-1)+b} \equiv \frac{B_b}{b} \pmod{p},$$

we arrive at

$$B_{2p-4} \equiv \frac{4}{3}B_{p-3} \pmod{p}. \quad (1.5)$$

Substituting (1.5) into (1.3) and (1.4), we get (1.1) and (1.2) for primes  $p \geq 7$ . It is routine to check that (1.1) and (1.2) also hold for  $p = 5$ .

The rest of this paper is organized as follows. In the next section, we prove Theorem 1.2. We show Theorem 1.3 in the last section.

## 2 Proof of Theorem 1.2

Since

$$\binom{p-1+k}{k} = \frac{p}{p+k} \binom{p+k}{k}, \quad (2.1)$$

we have

$$A_{p-1} = \sum_{k=0}^{p-1} \frac{p^2}{(p+k)^2} \binom{p-1}{k}^2 \binom{p+k}{k}^2. \quad (2.2)$$

Note that

$$\begin{aligned} \binom{p-1}{k} \binom{p+k}{k} &= \frac{(p^2-1^2)(p^2-2^2)\cdots(p^2-k^2)}{k!^2} \\ &\equiv (-1)^k \left(1 - p^2 H_k^{(2)}\right) \pmod{p^4}, \end{aligned} \quad (2.3)$$

where  $H_n^{(r)}$  denotes the  $n$ th generalized harmonic number of order  $r$ :

$$H_n^{(r)} = \sum_{k=1}^n \frac{1}{k^r},$$

with the convention that  $H_n = H_n^{(1)}$ . It follows from (2.2) and (2.3) that

$$\begin{aligned} A_{p-1} &= 1 + \sum_{k=1}^{p-1} \frac{p^2}{(p+k)^2} \binom{p-1}{k}^2 \binom{p+k}{k}^2 \\ &\equiv 1 + p^2 \sum_{k=1}^{p-1} \frac{1 - 2p^2 H_k^{(2)}}{(p+k)^2} \pmod{p^6}. \end{aligned} \quad (2.4)$$

Furthermore, we have

$$\frac{1}{(p+k)^2} \equiv \frac{1}{k^2} - \frac{2p}{k^3} + \frac{3p^2}{k^4} \pmod{p^3}. \quad (2.5)$$

Substituting (2.5) into (2.4) gives

$$A_{p-1} \equiv 1 + p^2 H_{p-1}^{(2)} - 2p^3 H_{p-1}^{(3)} + 3p^4 H_{p-1}^{(4)} - 2p^4 \sum_{k=1}^{p-1} \frac{H_k^{(2)}}{k^2} \pmod{p^5}. \quad (2.6)$$

For  $1 \leq k \leq p-1$ , we have

$$\begin{aligned} H_k^{(2)} + H_{p-k}^{(2)} &\equiv H_k^{(2)} + \sum_{i=1}^{p-k} \frac{1}{(p-i)^2} \pmod{p} \\ &= H_{p-1}^{(2)} + \frac{1}{k^2} \\ &\equiv \frac{1}{k^2} \pmod{p}, \end{aligned}$$

and so

$$\frac{H_k^{(2)}}{k^2} + \frac{H_{p-k}^{(2)}}{(p-k)^2} \equiv \frac{1}{k^4} \pmod{p}.$$

It follows that

$$\begin{aligned} \sum_{k=1}^{p-1} \frac{H_k^{(2)}}{k^2} &= \sum_{k=1}^{\frac{p-1}{2}} \left( \frac{H_k^{(2)}}{k^2} + \frac{H_{p-k}^{(2)}}{(p-k)^2} \right) \\ &\equiv \sum_{k=1}^{\frac{p-1}{2}} \frac{1}{k^4} \pmod{p}. \end{aligned} \quad (2.7)$$

By [7, page 353], we have

$$\begin{aligned} \sum_{k=1}^{\frac{p-1}{2}} \frac{1}{k^4} &\equiv \frac{1}{2} \left( \sum_{k=1}^{\frac{p-1}{2}} \frac{1}{k^4} + \sum_{k=1}^{\frac{p-1}{2}} \frac{1}{(p-k)^4} \right) \pmod{p} \\ &= \frac{1}{2} H_{p-1}^{(4)} \\ &\equiv 0 \pmod{p}. \end{aligned} \quad (2.8)$$

It follows from (2.7) and (2.8) that

$$\sum_{k=1}^{p-1} \frac{H_k^{(2)}}{k^2} \equiv 0 \pmod{p}. \quad (2.9)$$

By [7, page 353], we have

$$H_{p-1}^{(3)} \equiv 0 \pmod{p^2}, \quad (2.10)$$

$$H_{p-1}^{(4)} \equiv 0 \pmod{p}. \quad (2.11)$$

Combining (2.6) and (2.9)–(2.11) gives

$$A_{p-1} \equiv 1 + p^2 H_{p-1}^{(2)} \pmod{p^5}. \quad (2.12)$$

Letting  $k = 2$  in [11, Theorem 5.1, (a)] and simplifying gives

$$H_{p-1}^{(2)} \equiv \left( \frac{4}{3} B_{p-3} - \frac{1}{2} B_{2p-4} \right) p + \left( \frac{4}{9} B_{p-3} - \frac{1}{4} B_{2p-4} \right) p^2 \pmod{p^3}. \quad (2.13)$$

Substituting (1.5) into (2.13) yields

$$H_{p-1}^{(2)} \equiv \left( \frac{4}{3} B_{p-3} - \frac{1}{2} B_{2p-4} \right) p + \frac{1}{9} p^2 B_{p-3} \pmod{p^3}. \quad (2.14)$$

The proof of (1.3) follows from (2.12) and (2.14).

### 3 Proof of Theorem 1.3

In order to prove Theorem 1.3, we need the following combinatorial identities.

**Lemma 3.1** *For any non-negative integer  $n$ , we have*

$$\sum_{k=1}^{2n} \frac{(-1)^k}{k} \binom{2n}{k} \binom{2n+1+k}{k} = -2H_{2n}, \quad (3.1)$$

$$\sum_{k=1}^n \frac{(-1)^k}{k} \binom{n}{k} \binom{n+1+k}{k} H_k = 2 \left( \sum_{k=1}^n \frac{(-1)^k}{k^2} + \frac{(-1)^n}{n+1} H_n \right). \quad (3.2)$$

*Proof.* By Schneider's computer algebra package **Sigma** (see [10]), we find that the left-hand side of (3.1) satisfies the following recurrence:

$$\begin{aligned} & (-1 - 2n)\text{SUM}[n] + 2(3 + 2n)\text{SUM}[1 + n] + (-5 - 2n)\text{SUM}[2 + n] \\ &= \frac{17 + 24n + 8n^2}{(1 + n)(2 + n)(3 + 2n)}. \end{aligned}$$

It is easy to verify that the right-hand side of (3.1) also satisfies the recurrence above and both sides of (3.1) are equal for  $n = 0, 1$ .

In a similar way, both sides of (3.2) satisfy the same recurrence:

$$\begin{aligned} & (1 + n)(2 + n)^2(7 + 2n)\text{SUM}[n] + (2 + n)(7 + 2n)(7 + 6n + n^2)\text{SUM}[1 + n] \\ & - (3 + n)(3 + 2n)(2 + 4n + n^2)\text{SUM}[2 + n] - (3 + n)^2(4 + n)(3 + 2n)\text{SUM}[3 + n] = 0. \end{aligned}$$

It is routine to check that both sides of (3.2) are equal for  $n = 0, 1, 2$ . □

*Proof of (1.4).* By (2.1) and (2.3), we have

$$\begin{aligned} A'_{p-1} &= 1 + \sum_{k=1}^{p-1} \frac{p}{p+k} \binom{p-1}{k}^2 \binom{p+k}{k} \\ &\equiv 1 + p \sum_{k=1}^{p-1} \frac{(-1)^k}{p+k} \binom{p-1}{k} \left(1 - p^2 H_k^{(2)}\right) \pmod{p^5}. \end{aligned} \quad (3.3)$$

Letting  $n = p - 1$  and  $x = p$  in the following partial fraction decomposition:

$$\sum_{k=0}^n \frac{(-1)^k}{x+k} \binom{n}{k} = \frac{n!}{(x)_{n+1}},$$

we arrive at

$$\sum_{k=0}^{p-1} \frac{(-1)^k}{p+k} \binom{p-1}{k} = \frac{1}{p \binom{2p-1}{p-1}}.$$

It follows from the above that

$$\begin{aligned} p \sum_{k=1}^{p-1} \frac{(-1)^k}{p+k} \binom{p-1}{k} &= p \sum_{k=0}^{p-1} \frac{(-1)^k}{p+k} \binom{p-1}{k} - 1 \\ &= \frac{1}{\binom{2p-1}{p-1}} - 1. \end{aligned} \quad (3.4)$$

We need the following McIntosh's congruence (see [8, (6)]):

$$\binom{2p-1}{p-1} \equiv 1 - p^2 H_{p-1}^{(2)} \pmod{p^5}. \quad (3.5)$$

Substituting (3.5) into (3.4) and using the fact that  $H_{p-1}^{(2)} \equiv 0 \pmod{p}$ , we arrive at

$$p \sum_{k=1}^{p-1} \frac{(-1)^k}{p+k} \binom{p-1}{k} \equiv p^2 H_{p-1}^{(2)} \pmod{p^5}. \quad (3.6)$$

On the other hand, using  $\binom{p-1}{k} \equiv (-1)^k (1 - pH_k) \pmod{p^2}$  we have

$$\begin{aligned} p^3 \sum_{k=1}^{p-1} \frac{(-1)^k}{p+k} \binom{p-1}{k} H_k^{(2)} \\ \equiv p^3 \sum_{k=1}^{p-1} \frac{H_k^{(2)}}{p+k} - p^4 \sum_{k=1}^{p-1} \frac{H_k H_k^{(2)}}{p+k} \pmod{p^5}. \end{aligned} \quad (3.7)$$

Next, we shall prove that

$$\sum_{k=1}^{p-1} \frac{H_k H_k^{(2)}}{p+k} \equiv \sum_{k=1}^{p-1} \frac{H_k H_k^{(2)}}{k} \equiv 0 \pmod{p}. \quad (3.8)$$

Letting  $n = p - 1$  in (3.2) and noting (2.3), we find that

$$\sum_{k=1}^{p-1} \frac{(1 - p^2 H_k^{(2)}) H_k}{k} \equiv 2 \left( \sum_{k=1}^{p-1} \frac{(-1)^k}{k^2} + \frac{1}{p} H_{p-1} \right) \pmod{p^4}.$$

It follows that

$$p^2 \sum_{k=1}^{p-1} \frac{H_k H_k^{(2)}}{k} \equiv \sum_{k=1}^{p-1} \frac{H_k}{k} - 2 \left( \sum_{k=1}^{p-1} \frac{(-1)^k}{k^2} + \frac{1}{p} H_{p-1} \right) \pmod{p^4}.$$

Since

$$\begin{aligned} \sum_{k=1}^{p-1} \frac{H_k}{k} &= \frac{1}{2} \left( H_{p-1}^2 + H_{p-1}^{(2)} \right), \\ \sum_{k=1}^{p-1} \frac{(-1)^k}{k^2} &= \frac{1}{2} H_{(p-1)/2}^{(2)} - H_{p-1}^{(2)}, \\ H_{p-1} &\equiv 0 \pmod{p^2}, \end{aligned}$$

we have

$$p^2 \sum_{k=1}^{p-1} \frac{H_k H_k^{(2)}}{k} \equiv -\frac{2}{p} H_{p-1} + \frac{5}{2} H_{p-1}^{(2)} - H_{(p-1)/2}^{(2)} \pmod{p^4}.$$

From [8, (6) & (7)], we deduce that

$$\frac{1}{p} H_{p-1} \equiv -\frac{1}{2} H_{p-1}^{(2)} \pmod{p^3}.$$

Thus,

$$p^2 \sum_{k=1}^{p-1} \frac{H_k H_k^{(2)}}{k} \equiv \frac{7}{2} H_{p-1}^{(2)} - H_{(p-1)/2}^{(2)} \pmod{p^3}. \quad (3.9)$$

Letting  $k = 2$  in [11, Theorem 5.2, (a)] and simplifying gives

$$H_{(p-1)/2}^{(2)} \equiv \left( \frac{14}{3} B_{p-3} - \frac{7}{4} B_{2p-4} \right) p + \left( \frac{14}{9} B_{p-3} - \frac{7}{8} B_{2p-4} \right) p^2 \pmod{p^3}. \quad (3.10)$$

Substituting (2.13) and (3.10) into (3.9) gives

$$p^2 \sum_{k=1}^{p-1} \frac{H_k H_k^{(2)}}{k} \equiv 0 \pmod{p^3},$$

which implies (3.8).

Since

$$\frac{1}{p+k} \equiv \frac{1}{k} - \frac{p}{k^2} \pmod{p^2},$$

by (2.9) we have

$$\begin{aligned} \sum_{k=1}^{p-1} \frac{H_k^{(2)}}{p+k} &\equiv \sum_{k=1}^{p-1} \frac{H_k^{(2)}}{k} - p \sum_{k=1}^{p-1} \frac{H_k^{(2)}}{k^2} \\ &\equiv \sum_{k=1}^{p-1} \frac{H_k^{(2)}}{k} \pmod{p^2}. \end{aligned} \quad (3.11)$$

Combining (3.7), (3.8) and (3.11) gives

$$p^3 \sum_{k=1}^{p-1} \frac{(-1)^k}{p+k} \binom{p-1}{k} H_k^{(2)} \equiv p^3 \sum_{k=1}^{p-1} \frac{H_k^{(2)}}{k} \pmod{p^5}. \quad (3.12)$$

Letting  $n = \frac{p-1}{2}$  in (3.1), we get

$$\sum_{k=1}^{p-1} \frac{(-1)^k}{k} \binom{p-1}{k} \binom{p+k}{k} = -2H_{p-1}.$$

By (2.3), we have

$$\sum_{k=1}^{p-1} \frac{1 - p^2 H_k^{(2)}}{k} \equiv -2H_{p-1} \pmod{p^4},$$

and so

$$p^2 \sum_{k=1}^{p-1} \frac{H_k^{(2)}}{k} \equiv 3H_{p-1} \pmod{p^4},$$

which implies that

$$\sum_{k=1}^{p-1} \frac{H_k^{(2)}}{k} \equiv \frac{3}{p^2} H_{p-1} \pmod{p^2}. \quad (3.13)$$

Substituting (3.13) into (3.12) gives

$$p^3 \sum_{k=1}^{p-1} \frac{(-1)^k}{p+k} \binom{p-1}{k} H_k^{(2)} \equiv 3pH_{p-1} \pmod{p^5}. \quad (3.14)$$

From [8, (6) & (7)], we deduce that

$$pH_{p-1} \equiv -\frac{p^2}{2} H_{p-1}^{(2)} \pmod{p^5}. \quad (3.15)$$

Finally combining (3.3), (3.6), (3.14) and (3.15) gives

$$A'_{p-1} \equiv 1 + \frac{5}{2} p^2 H_{p-1}^{(2)} \pmod{p^5}. \quad (3.16)$$

Then the proof of (1.4) follows from (2.14) and (3.16).  $\square$

**Acknowledgments.** The authors are grateful to Professor Zhi-Hong Sun for bringing the reference [11] to their attention and helpful conversations.

## References

- [1] R. Apéry, Irrationalité de  $\zeta(2)$  et  $\zeta(3)$ , Astérisque 61 (1979), 11–13.



- [2] F. Beukers, Some congruences for the Apéry numbers, *J. Number Theory* 21 (1985), 141–155.
- [3] H.H. Chan, S. Cooper and F. Sica, Congruences satisfied by Apéry-like numbers, *Int. J. Number Theory* 6 (2010), 89–97.
- [4] É. Delaygue, Arithmetic properties of Apéry-like numbers, *Compos. Math.* 154 (2018), 249–274.
- [5] I. Gessel, Some congruences for Apéry numbers, *J. Number Theory* 14 (1982), 362–368.
- [6] V.J.W. Guo and J. Zeng, Proof of some conjectures of Z.-W. Sun on congruences for Apéry polynomials, *J. Number Theory* 132 (2012), 1731–1740.
- [7] E. Lehmer, On congruences involving Bernoulli numbers and the quotients of Fermat and Wilson, *Ann. Math.* 39 (1938), 350–360.
- [8] R. Mestrovic, Wolstenholme’s theorem: Its Generalizations and Extensions in the last hundred and fifty years (1862–2012), preprint, 2011, arXiv:1111.3057.
- [9] H. Pan, On divisibility of sums of Apéry polynomials, *J. Number Theory* 143 (2014), 214–223.
- [10] C. Schneider, Symbolic summation assists combinatorics, *Sém. Lothar. Combin.* 56 (2007), B56b, 36 pp.
- [11] Z.-H. Sun, Congruences concerning Bernoulli numbers and Bernoulli polynomials, *Discrete Appl. Math.* 105 (2000), 193–223.
- [12] Z.-H. Sun, Congruences for Apéry-like numbers, preprint, 2018, arXiv:1803.10051.
- [13] Z.-W. Sun, Arithmetic theory of harmonic numbers, *Proc. Amer. Math. Soc.* 140 (2012), 415–428.
- [14] Z.-W. Sun, On sums of Apéry polynomials and related congruences, *J. Number Theory* 132 (2012), 2673–2699.
- [15] Z.-W. Sun, Conjectures and results on  $x^2 \pmod{p^2}$  with  $4p = x^2 + dy^2$ , in: *Number Theory and Related Area* (eds., Y. Ouyang, C. Xing, F. Xu and P. Zhang), *Adv. Lect. Math.* 27, Higher Education Press & International Press, Beijing-Boston, 2013, 149–197, available at <https://arxiv.org/abs/1103.4325>.
- [16] Z.-W. Sun, Congruences involving generalized trinomial coefficients, *Sci. China Math.* 57 (2014), 1375–1400.