

**ASYMPTOTIC GREEDINESS OF THE HAAR SYSTEM
IN THE SPACES $L_p[0, 1]$, $1 < p < \infty$**

FERNANDO ALBIAC, JOSÉ L. ANSORENA, AND PABLO M. BERNÁ

ABSTRACT. Our aim in this paper is to attain a sharp asymptotic estimate for the greedy constant $C_g[\mathcal{H}^{(p)}, L_p]$ of the (normalized) Haar system $\mathcal{H}^{(p)}$ in $L_p[0, 1]$ for $1 < p < \infty$. We will show that the superdemocracy constant of $\mathcal{H}^{(p)}$ in $L_p[0, 1]$ grows as $p^* = \max\{p, p/(p-1)\}$ as p^* goes to ∞ . Thus, since the unconditionality constant of $\mathcal{H}^{(p)}$ in $L_p[0, 1]$ is $p^* - 1$, the well-known general estimates for the greedy constant of a greedy basis obtained from the intrinsic features of greediness (namely, democracy and unconditionality) yield that $p^* \lesssim C_g[\mathcal{H}^{(p)}, L_p] \lesssim (p^*)^2$. Going further, we develop techniques that allow us to close the gap between those two bounds, establishing that, in fact, $C_g[\mathcal{H}^{(p)}, L_p] \approx p^*$.

1. INTRODUCTION

A fundamental and total biorthogonal system for an infinite-dimensional separable Banach space $(\mathbb{X}, \|\cdot\|)$ over the field \mathbb{F} of real or complex scalars, is a family $(\mathbf{x}_j, \mathbf{x}_j^*)_{j \in J}$ in $\mathbb{X} \times \mathbb{X}^*$ verifying

- (i) $\mathbb{X} = \overline{\text{span}\{\mathbf{x}_j : j \in J\}}$,
- (ii) $\mathbb{X}^* = \overline{\text{span}\{\mathbf{x}_j^* : j \in J\}}^{w^*}$, and
- (iii) $\mathbf{x}_j^*(\mathbf{x}_k) = 1$ if $j = k$ and $\mathbf{x}_j^*(\mathbf{x}_k) = 0$ otherwise.

The family $\mathcal{B} = (\mathbf{x}_j)_{j \in J}$ is called a (*Markushevich*) *basis* and the unambiguously determined collection of bounded linear functionals $\mathcal{B}^* = (\mathbf{x}_j^*)_{j \in J}$ is said to be the family of *coordinate functionals* (or *dual basis*) of \mathcal{B} . If the biorthogonal system verifies the condition

- (iv) $\sup_{j \in J} \|\mathbf{x}_j\| \|\mathbf{x}_j^*\| < \infty$

we say that \mathcal{B} is *M-bounded*. Finally, if we have

- (v) $0 < \inf_{j \in J} \|\mathbf{x}_j\| \leq \sup_{j \in J} \|\mathbf{x}_j\| < \infty$

2010 *Mathematics Subject Classification.* 46B15, 41A65.

Key words and phrases. Haar basis, greedy basis, unconditional basis, democratic basis, L_p spaces, Lebesgue-type inequalities.

we say that \mathcal{B} is *semi-normalized* (*normalized* if $\|\mathbf{x}_j\| = 1$ for all $j \in J$). Note that a basis \mathcal{B} verifies simultaneously (iv) and (v) if and only if

$$\sup_{j \in J} \max\{\|\mathbf{x}_j\|, \|\mathbf{x}_j^*\|\} < \infty.$$

Suppose $\mathcal{B} = (\mathbf{x}_j)_{j \in J}$ is a semi-normalized M -bounded basis in a Banach space \mathbb{X} with coordinate functionals $(\mathbf{x}_j^*)_{j \in J}$. Then each $f \in \mathbb{X}$ is uniquely determined by its *coefficient family* $(\mathbf{x}_j^*(f))_{j \in J}$, which belongs to $c_0(J)$. Thus, we can consider its non-increasing rearrangement, which we denote by $(a_m^*[\mathcal{B}, \mathbb{X}](f))_{m=1}^\infty$ (or, simply $(a_m^*(f))_{m=1}^\infty$ if the basis and the space are clear from context).

For each $f \in \mathbb{X}$ there is a 1 – 1 map $\rho: \mathbb{N} \rightarrow J$ such that

$$|\mathbf{x}_{\rho(m)}^*(f)| = a_m^*(f), \quad m \in \mathbb{N}. \quad (1.1)$$

If the family $(\mathbf{x}_j^*(f))_{j \in J}$ contains several terms with the same absolute value then the map ρ for f is not uniquely determined. In order to get uniqueness, we arrange the countable set J by means of a bijection $\sigma: J \rightarrow \mathbb{N}$ and impose the additional condition

$$\sigma(\rho(m)) \leq \sigma(\rho(n)) \text{ whenever } |\mathbf{x}_{\rho(m)}^*(f)| = |\mathbf{x}_{\rho(n)}^*(f)|. \quad (1.2)$$

If f is infinitely supported there is a unique 1 – 1 map $\rho: \mathbb{N} \rightarrow J$ with $\rho(\mathbb{N}) = \text{supp}(f)$ that verifies (1.1) and (1.2). In the case when f is finitely supported, there is a unique bijection $\rho: \mathbb{N} \rightarrow J$ that verifies (1.1) and (1.2). In any case, we will refer to such a map ρ as the *greedy ordering* for f . For each $m \in \mathbb{N}$, the *m th-greedy approximation* to f is the partial sum

$$\mathcal{G}_m[\mathcal{B}, \mathbb{X}](f) := \mathcal{G}_m(f) = \sum_{n=1}^m \mathbf{x}_{\rho(n)}^*(f) \mathbf{x}_{\rho(n)},$$

where ρ is the greedy ordering for f . The sequence $(\mathcal{G}_m(f))_{m=1}^\infty$ is called the greedy algorithm for f with respect to the basis \mathcal{B} .

To quantify the efficiency of the greedy algorithm, Temlyakov [11] introduced the sequence $(\mathbf{L}_m)_{m=1}^\infty$ of *Lebesgue greedy constants*. For each $m \in \mathbb{N}$, $\mathbf{L}_m[\mathcal{B}, \mathbb{X}] := \mathbf{L}_m$ is the smallest constant C such that the Lebesgue-type inequality

$$\|f - \mathcal{G}_m(f)\| \leq C \left\| f - \sum_{j \in A} a_j \mathbf{x}_j \right\|, \quad (1.3)$$

holds for all $f \in \mathbb{X}$, all subsets A of \mathbb{N} with $|A| = m$, and all $a_j \in \mathbb{F}$. Konyagin and Temlyakov [9] then defined a basis \mathcal{B} to be *greedy* if (1.3) holds with a constant $1 \leq C < \infty$ independent of m . The smallest admissible constant C will be denoted by $C_g[\mathcal{B}, \mathbb{X}] = C_g$, and will be

referred to as the *greedy constant* of the basis. In other words, a basis \mathcal{B} is greedy if and only if $\sup_m \mathbf{L}_m = C_g < \infty$, that is, the greedy algorithm provides, up to a multiplicative constant, the best m -term approximation to any vector in the space.

Once we know that a certain basis is greedy, a natural problem in approximation theory is to compute, or at least estimate, its greedy constant. Also of interest is to determine for what values of C a basis is C -greedy under a suitable renorming of the space.

In this paper we focus on the Haar system $\mathcal{H}^{(p)} = (h_I^{(p)})_{I \in \mathcal{D}_0}$ in the spaces $L_p[0, 1]$ (L_p for short from now on). Here, \mathcal{D}_0 denotes the set $\mathcal{D} \cup \{0\}$, where \mathcal{D} is the collection of all dyadic intervals contained in $[0, 1)$, $h_0^{(p)}$ is the constant function 1 on $[0, 1)$, and $h_I^{(p)}$ stands for the L_p -normalized Haar function supported on I , i.e., if $I = [a, b)$ then

$$h_I^{(p)}(t) = \begin{cases} -2^{-(b-a)/p} & \text{if } a \leq t < (a+b)/2, \\ 2^{-(b-a)/p} & \text{if } (a+b)/2 \leq t < b. \end{cases}$$

Thus $\mathcal{H}^{(p)}$ is a normalized M -bounded basis for L_p when $1 \leq p < \infty$ and a normalized M -bounded basis for the space \mathbb{D} of Càglàg functions when $p = \infty$. In either case, the family of coordinate functionals of $\mathcal{H}^{(p)}$ is $\mathcal{H}^{(p')}$ (with the canonical isometric identification of functions in $L_{p'}$ with functionals in $(L_p)^*$, where $p' = p/(p-1)$). In fact, when arranged in the natural way, $\mathcal{H}^{(p)}$ is a Schauder basis (see [2, Proposition 6.1.3]).

Temlyakov [10] showed that $\mathcal{H}^{(p)}$ is a greedy basis in L_p for $1 < p < \infty$. Later on, Dilworth et al. [7] proved that for every $C > 1$ there is a renorming of L_p with respect to which $\mathcal{H}^{(p)}$ is C -greedy. Whether or not the isometric constant $C = 1$ can be achieved up to renorming (see [3]) remains unknown as of today.

Note that neither $\mathcal{H}^{(1)}$ is a greedy basis for L_1 nor $\mathcal{H}^{(\infty)}$ is a greedy basis for \mathbb{D} . Indeed, every greedy (or unconditional) basis for a \mathcal{L}_1 -space is equivalent to the unit vector basis of ℓ_1 and every greedy basis for a \mathcal{L}_∞ -space is equivalent to the unit vector basis of c_0 . Consequently we have

$$\lim_{p \rightarrow 1^+} C_g[\mathcal{H}^{(p)}, L_p] = \infty = \lim_{p \rightarrow \infty} C_g[\mathcal{H}^{(p)}, L_p].$$

The Haar system $\mathcal{H}^{(2)}$ is an orthonormal basis for L_2 , which easily yields $C_g[\mathcal{H}^{(2)}, L_2] = 1$. However, for $p \neq 2$ it seems hopeless to attempt to compute the exact value of $C_g[\mathcal{H}^{(p)}, L_p]$. It is therefore natural to address the problem of obtaining asymptotic estimates for $C_g[\mathcal{H}^{(p)}, L_p]$ as p tends to 1 or to ∞ .

A standard approach to estimate the greedy constant of a greedy basis is to make use of its intrinsic properties instead of relying on the

mere definition. The first movers in this direction were Konyagin and Temlyakov [9], who proved that a basis is greedy if and only if it is *unconditional* and *democratic*. To set the notation, we recall that a basis $\mathcal{B} = (\mathbf{x}_j)_{j \in J}$ for a Banach space \mathbb{X} is said to be unconditional if the series expansion $\sum_{j \in J} \mathbf{x}_j^*(f) \mathbf{x}_j$ converges unconditionally to f for every $f \in \mathbb{X}$. Unconditional bases are characterized as those bases verifying the uniform bound

$$K_{su}[\mathcal{B}, \mathbb{X}] := \sup_{\substack{A \subseteq J \\ |A| < \infty}} \|S_A\| = \sup_{\substack{A \subseteq J \\ |A| < \infty}} \|\text{Id}_{\mathbb{X}} - S_A\| < \infty, \quad (1.4)$$

or, equivalently,

$$K_u[\mathcal{B}, \mathbb{X}] := \sup\{\|M_\varepsilon\| : \varepsilon = (\varepsilon_j)_{j \in J}, |\varepsilon_j| = 1\} < \infty, \quad (1.5)$$

where $S_A = S_A[\mathcal{B}, \mathbb{X}]$ is the coordinate projection on a finite set $A \subseteq J$, i.e.,

$$S_A: \mathbb{X} \rightarrow \mathbb{X}, \quad f \mapsto \sum_{j \in A} \mathbf{x}_j^*(f) \mathbf{x}_j,$$

and $M_\varepsilon = M_\varepsilon[\mathcal{B}, \mathbb{X}]$ is the linear operator from \mathbb{X} into \mathbb{X} given by $\mathbf{x}_j \mapsto \varepsilon_j \mathbf{x}_j$. The *suppression unconditional constant* K_{su} and the *lattice unconditional constant* K_u of a basis are related by the inequalities

$$K_{su}[\mathcal{B}, \mathbb{X}] \leq K_u[\mathcal{B}, \mathbb{X}] \leq \kappa K_{su}[\mathcal{B}, \mathbb{X}],$$

where $\kappa = 2$ if $\mathbb{F} = \mathbb{R}$ and $\kappa = 4$ if $\mathbb{F} = \mathbb{C}$. In turn, a basis \mathcal{B} is said to be *democratic* if there is $1 \leq C < \infty$ such that

$$\left\| \sum_{j \in A} \mathbf{x}_j \right\| \leq C \left\| \sum_{j \in B} \mathbf{x}_j \right\|, \quad |A| = |B| < \infty. \quad (1.6)$$

We will denote by $\Delta[\mathcal{B}, \mathbb{X}]$ the optimal constant C in (1.6). By imposing the extra assumption $A \cap B = \emptyset$ in (1.6) we obtain an equivalent definition of democracy, and $\Delta_d[\mathcal{B}, \mathbb{X}]$ will denote the optimal constant in (1.6) under the extra assumption on disjointness of sets.

Amalgamating some steps in Konyagin–Temlyakov’s proof (see [7, Equation 1]) we get the estimate

$$\max\{K_{su}[\mathcal{B}, \mathbb{X}], \Delta[\mathcal{B}, \mathbb{X}]\} \leq C_g[\mathcal{B}, \mathbb{X}] \leq K_{su}[\mathcal{B}, \mathbb{X}] + \Delta_d[\mathcal{B}, \mathbb{X}] K_u^2[\mathcal{B}, \mathbb{X}]. \quad (1.7)$$

Note that

$$\Delta[\mathcal{B}, \mathbb{X}] \leq \Delta_d[\mathcal{B}, \mathbb{X}](1 + K_{su}[\mathcal{B}, \mathbb{X}]).$$

Hence, when \mathcal{B} runs over a certain family of bases, the left-hand side and the right-hand side terms in inequality (1.7) are of the same order if and only if the constants $K_u[\mathcal{B}, \mathbb{X}]$ are uniformly bounded. This is not the case for $(\mathcal{H}^{(p)})_{p>1}$ as the following theorem of Burkholder exhibits:

Theorem 1.1 ([5]). *If $1 < p < \infty$ then $K_u[\mathcal{H}^{(p)}, L_p] = p^* - 1$, where $p^* = \max\{p, p'\}$.*

Now, we may try to obtain estimates that bring us closer to our goal by using super-democracy instead of democracy. A basis is *super-democratic* if there is a constant $1 \leq C < \infty$ such that

$$\left\| \sum_{j \in A} \varepsilon_j \mathbf{x}_j \right\| \leq C \left\| \sum_{j \in B} \delta_j \mathbf{x}_j \right\|, \quad |A| = |B| < \infty, |\varepsilon_j| = |\delta_j| = 1. \quad (1.8)$$

The smallest admissible constant in (1.8) will be denoted by $\Delta_s[\mathcal{B}, \mathbb{X}]$, and the smallest constant in (1.8) with the extra assumption $A \cap B = \emptyset$ will be denoted by $\Delta_{sd}[\mathcal{B}, \mathbb{X}]$.

Bases that are unconditional and democratic are super-democratic. Hence, greedy bases are characterized as those that are simultaneously unconditional and super-democratic. Quantitatively, a slight improvement of the argument used in the proof of [4, Theorem 1.4] gives

$$\max\{K_{su}[\mathcal{B}, \mathbb{X}], \Delta_{sd}[\mathcal{B}, \mathbb{X}]\} \leq C_g[\mathcal{B}, \mathbb{X}] \leq K_{su}[\mathcal{B}, \mathbb{X}](1 + \Delta_{sd}[\mathcal{B}, \mathbb{X}]).$$

These inequalities allow us to determine the rate of growth of the constants $C_g[\mathcal{B}, \mathbb{X}]$ when \mathcal{B} runs over a family of bases only when $\min\{[K_{su}[\mathcal{B}, \mathbb{X}], \Delta_{sd}[\mathcal{B}, \mathbb{X}]]\}$ is uniformly bounded. But, again, this is not the case for the L_p -normalized Haar system for $1 < p < \infty$. Indeed, on the one hand we have $\Delta_{sd}[\mathcal{B}, \mathbb{X}] \geq \Delta_d[\mathcal{B}, \mathbb{X}]$ for any \mathcal{B} and any \mathbb{X} . On the other hand, the following result (which we shall prove below) yields $\sup_{p>1} \Delta_d[\mathcal{H}^{(p)}, L_p] = \infty$.

Proposition 1.2. *If $1 < p < \infty$ then*

$$\Delta_d[\mathcal{H}^{(p)}, L_p] \geq d_p := \frac{2^{1/p^\#} - 1}{2^{1/p^*} - 1},$$

where $p^\# = \min\{p, p'\}$.

Another important property of bases that comes into play in this scenario is the *symmetry for largest coefficients* (a.k.a. *Property A*). A basis \mathcal{B} is said to be symmetric for largest coefficients if there is a constant $1 \leq C < \infty$ such that

$$\left\| \sum_{j \in A} \varepsilon_j \mathbf{x}_j + f \right\| \leq C \left\| \sum_{j \in B} \delta_j \mathbf{x}_j + f \right\| \quad (1.9)$$

whenever $|A| = |B| < \infty$, $A \cap B = (A \cup B) \cap \text{supp}(f) = \emptyset$, and $|\mathbf{x}_i^*(f)| \leq |\varepsilon_j| = |\delta_k| = 1$ for all $i \in J$, $j \in A$, $k \in B$. We denote by $C_a[\mathcal{B}, \mathbb{X}]$ the optimal constant C in (1.9). A basis is greedy if and only

if it is unconditional and symmetric for largest coefficients. Moreover (see [1, Remark 2.6]),

$$\max\{K_{su}[\mathcal{B}, \mathbb{X}], C_a[\mathcal{B}, \mathbb{X}]\} \leq C_g[\mathcal{B}, \mathbb{X}] \leq C_a[\mathcal{B}, \mathbb{X}]K_{su}[\mathcal{B}, \mathbb{X}]. \quad (1.10)$$

These estimates are useful when one wants to show that the greedy constant of a certain basis is close to 1. However, since $C_a[\mathcal{B}, \mathbb{X}] \geq \Delta_{sd}[\mathcal{B}, \mathbb{X}]$, and the side terms of (1.10) are of different order, they do not provide a tight information about the asymptotic growth of the greedy constants of a family of bases.

Despite the fact that the methods described above are not strong enough to be applied to our problem, in this note we shall reach our goal and prove the following theorem.

Theorem 1.3. $C_g[\mathcal{H}^{(p)}, L_p] \approx p^*$ for $1 < p < \infty$.

The key idea in the proof of Theorem 1.3 will consist of taking advantage of the fact that the Haar system in L_p belongs to a more demanding class of bases than that of greedy bases, namely, the class of *bi-greedy* bases. A basis is said to be bi-greedy if both the basis itself and its dual basis are greedy. Bi-greedy bases were characterized in [6] as those bases that are unconditional and *bi-democratic*. Recall that a basis $\mathcal{B} = (\mathbf{x}_j)_{j=1}^\infty$ is said to be bi-democratic if there is a constant $1 \leq C < \infty$ such that

$$\left\| \sum_{j \in A} \mathbf{x}_j \right\| \left\| \sum_{k \in B} \mathbf{x}_k^* \right\| \leq Cm, \quad |A| = |B| = m. \quad (1.11)$$

We will denote by $\Delta_b[\mathcal{B}, \mathbb{X}]$ the smallest constant C such that (1.11) holds, and we will refer to it as the bi-democratic constant of the basis.

The following new estimate for the greedy constant will also be crucial in the proof of Theorem 1.3.

Theorem 1.4. *Let \mathcal{B} be a bi-democratic and unconditional basis for a Banach space \mathbb{X} . Then*

$$C_g[\mathcal{B}, \mathbb{X}] \leq K_{su}[\mathcal{B}, \mathbb{X}] + \kappa^2 \Delta_b[\mathcal{B}, \mathbb{X}].$$

Section 2 is devoted to proving Theorem 1.4, while in Section 3 we obtain the remaining estimates that in combination with Theorem 1.4 will yield Theorem 1.3.

Throughout this article we use standard facts and notation from Banach spaces and approximation theory. Here, and throughout this paper, the symbol $\alpha_i \lesssim \beta_i$ for $i \in I$ means that the families of positive real numbers $(\alpha_i)_{i \in I}$ and $(\beta_i)_{i \in I}$ verify $\sup_{i \in I} \alpha_i / \beta_i < \infty$. If $\alpha_i \lesssim \beta_i$ and $\beta_i \lesssim \alpha_i$ for $i \in I$ we say $(\alpha_i)_{i \in I}$ are $(\beta_i)_{i \in I}$ are equivalent, and we

write $\alpha_i \approx \beta_i$ for $i \in I$. We refer the reader to [2] for the necessary background.

2. A NEW ESTIMATE FOR THE LEBESGUE TYPE CONSTANTS FOR
THE GREEDY ALGORITHM USING BI-DEMOCRACY

The *fundamental function* of a basis $\mathcal{B} = (\mathbf{x}_j)_{j=1}^\infty$ in a Banach space \mathbb{X} is the sequence given by

$$\varphi_m[\mathcal{B}, \mathbb{X}] = \sup_{|A| \leq m} \left\| \sum_{j \in A} \mathbf{x}_j \right\|, \quad m \in \mathbb{N}.$$

We shall also consider the *super-fundamental function* of the basis, given by

$$\varphi_m^\epsilon[\mathcal{B}, \mathbb{X}] = \varphi_m^\epsilon := \sup_{\substack{|A|=m \\ |\varepsilon_j|=1}} \left\| \sum_{j \in A} \varepsilon_j \mathbf{x}_j \right\|, \quad m \in \mathbb{N}.$$

A standard convexity argument yields

$$\varphi_m[\mathcal{B}, \mathbb{X}] \leq \sup_{\substack{|A|=m \\ |a_j| \leq 1}} \left\| \sum_{j \in A} a_j \mathbf{x}_j \right\| = \varphi_m^\epsilon[\mathcal{B}, \mathbb{X}] \leq \kappa \varphi_m[\mathcal{B}, \mathbb{X}]. \quad (2.1)$$

If \mathbb{Y} is the subspace of \mathbb{X}^* spanned by \mathcal{B}^* , we set

$$\varphi_m^{\epsilon,*}[\mathcal{B}, \mathbb{X}] = \varphi_m^{\epsilon,*} := \varphi_m^\epsilon[\mathcal{B}^*, \mathbb{Y}], \quad m \in \mathbb{N},$$

and define the sequence

$$\mathbf{B}_m[\mathcal{B}, \mathbb{X}] = \mathbf{B}_m := \sup_{r \leq m} \frac{\varphi_r^\epsilon[\mathcal{B}, \mathbb{X}] \varphi_r^{\epsilon,*}[\mathcal{B}, \mathbb{X}]}{r}.$$

Then a basis \mathcal{B} is bi-democratic if and only

$$\Delta_{sb}[\mathcal{B}, \mathbb{X}] := \sup_m \mathbf{B}_m < \infty. \quad (2.2)$$

Quantitatively we have

$$\Delta_b[\mathcal{B}, \mathbb{X}] \leq \Delta_{sb}[\mathcal{B}, \mathbb{X}] \leq \kappa^2 \Delta_b[\mathcal{B}, \mathbb{X}]. \quad (2.3)$$

Remark 2.1. *The identity*

$$\left(\sum_{j \in A} \mathbf{x}_j^* \right) \left(\sum_{j \in A} \mathbf{x}_j \right) = |A|, \quad A \subseteq J, |A| < \infty$$

yields (see [6])

$$\Delta[\mathcal{B}, \mathbb{X}] \leq \Delta_b[\mathcal{B}, \mathbb{X}]$$

Similarly, the identity

$$\left(\sum_{j \in A} \varepsilon_n \mathbf{x}_j^* \right) \left(\sum_{j \in A} \overline{\varepsilon_n} \mathbf{x}_j \right) = |A|, \quad A \subseteq J, |A| < \infty, |\varepsilon_j| = 1$$

gives

$$\Delta_s[\mathcal{B}, \mathbb{X}] \leq \Delta_{sb}[\mathcal{B}, \mathbb{X}].$$

Garrigós et al. [8] gave several estimates for the constants $\mathbf{L}_m[\mathcal{B}, \mathbb{X}]$ involving the sequences of democracy constants and of conditionality constants of the basis. Subsequently, Berná et al. [4] obtained estimates for $\mathbf{L}_m[\mathcal{B}, \mathbb{X}]$ that also involved the sequence of quasi-greedy constants of the basis. The estimate we shall provide involves the sequence of bi-democracy constants and the sequence of conditionality constants $(\mathbf{k}_m^c)_{m=1}^\infty$ given by

$$\mathbf{k}_m^c[\mathcal{B}, \mathbb{X}] = \mathbf{k}_m^c := \sup_{|A| \leq m} \|\text{Id}_{\mathbb{X}} - S_A[\mathcal{B}, \mathbb{X}]\|.$$

Note that the basis \mathcal{B} is unconditional if and only if

$$\sup_m \mathbf{k}_m^c = K_{su}[\mathcal{B}, \mathbb{X}] < \infty.$$

Lemma 2.2. *Let $\mathcal{B} = (\mathbf{x}_j)_{j \in J}$ be a semi-normalized M -bounded basis in a Banach space \mathbb{X} . For every $f \in \mathbb{X}$ and $m \in \mathbb{N}$,*

$$a_m^*[\mathcal{B}, \mathbb{X}](f) \varphi_m^\epsilon[\mathcal{B}, \mathbb{X}] \leq \mathbf{B}_m[\mathcal{B}, \mathbb{X}] \|f\|.$$

Proof. Let $G \subseteq J$ be such that $|G| = m$ and $a_m^* \leq |\mathbf{x}_j^*(f)|$ for all $j \in G$. Define $f^* \in \mathbb{X}^*$ by $f^* = \sum_{j \in G} \overline{\text{sign } \mathbf{x}_j^*(f)} \mathbf{x}_j^*$. Then

$$a_m^*(f) \varphi_m^\epsilon \leq \mathbf{B}_m \frac{m a_m^*(f)}{\varphi_m^{\epsilon, *}} \leq \mathbf{B}_m \frac{\sum_{j \in G} |\mathbf{x}_j^*(f)|}{\|f^*\|} = \mathbf{B}_m \frac{f^*(f)}{\|f^*\|} \leq \mathbf{B}_m \|f\|.$$

□

Theorem 2.3. *Let $\mathcal{B} = (\mathbf{x}_j)_{j \in J}$ be a semi-normalized M -bounded basis in a Banach space \mathbb{X} . For all $m \in \mathbb{N}$ we have*

$$\mathbf{L}_m[\mathcal{B}, \mathbb{X}] \leq \mathbf{k}_{2m}^c[\mathcal{B}, \mathbb{X}] + \mathbf{B}_m[\mathcal{B}, \mathbb{X}].$$

Proof. Let $f \in \mathbb{X}$, $m \in \mathbb{N}$ and $G \subseteq J$ of cardinality m such that $\mathcal{G}_m(f) = S_G(f)$. Let $A \subseteq J$ with $|A| = m$ and $(a_j)_{j \in A} \in \mathbb{F}^A$. Put $g = f - \sum_{j \in A} a_j \mathbf{x}_j$. We have

$$\|f - \mathcal{G}_m(f)\| = \|g - S_{A \cup G}(g) + S_{A \setminus G}(f)\| \leq \|g - S_{A \cup G}(g)\| + \|S_{A \setminus G}(f)\|.$$

Since $|A \cup B| \leq 2m$,

$$\|g - S_{A \cup G}(g)\| \leq \mathbf{k}_{2m}^c \|g\|.$$

Let $r = |A \setminus G| = |G \setminus A|$. Invoking (2.1) and Lemma 2.2,

$$\begin{aligned} \|S_{A \setminus G}(f)\| &\leq \max_{j \in A \setminus G} |\mathbf{x}_j^*(f)| \varphi_r^\varepsilon \\ &\leq \min_{j \in G \setminus A} |\mathbf{x}_j^*(f)| \varphi_r^\varepsilon \\ &= \min_{j \in G \setminus A} |\mathbf{x}_j^*(g)| \varphi_r^\varepsilon \\ &\leq a_r^*(g) \varphi_r^\varepsilon \\ &\leq \mathbf{B}_r \|g\|. \end{aligned}$$

Taking into account that $r \leq m$ and that $(\mathbf{B}_m)_{m=1}^\infty$ is non-decreasing, we get

$$\|S_{A \setminus G}(f)\| \leq \mathbf{B}_m \|g\|.$$

Combining, we obtain the desired result. \square

Proof of Theorem 1.4. Taking the supremum over m in Theorem 2.3 and appealing to (2.3) gives

$$C_g[\mathcal{B}, \mathbb{X}] \leq K_{su}[\mathcal{B}, \mathbb{X}] + \Delta_{sb}[\mathcal{B}, \mathbb{X}] \leq K_{su}[\mathcal{B}, \mathbb{X}] + \kappa^2 \Delta_b[\mathcal{B}, \mathbb{X}].$$

\square

3. ESTIMATES FOR THE HAAR BASIS IN L_p

We start this section with the proof advertised in Section 1 of the lower estimate for the democracy constant.

Proof of Proposition 1.2. If $(J_j)_{j=1}^m$ are disjointly supported intervals in \mathcal{D} we have

$$\left\| \sum_{j=1}^m h_{J_j}^{(p)} \right\|_p = m^{1/p}. \quad (3.1)$$

Let $(I_j)_{j=1}^\infty$ be the sequence in \mathcal{D} defined recursively as follows: $I_1 = [0, 1)$ and I_{j+1} is the left half of I_j . Set $q = p'$. Then

$$\begin{aligned} \left\| \sum_{j=1}^m h_{I_j}^{(p)} \right\|_p^p &= 2^{-m-1} \left| \sum_{k=0}^m 2^{k/p} \right|^p + \sum_{j=0}^{m-1} 2^{-j-1} \left| 2^{j/p} - \sum_{k=0}^{j-1} 2^{k/p} \right|^p \\ &= \frac{(1 - 2^{-(m+1)/p})^p + \sum_{j=0}^{m-1} | - 2^{-(j+1)/p} + (2^{1/q} - 1) |^p}{(2^{1/p} - 1)^p} \\ &= \frac{\|f - g\|_p^p}{(2^{1/p} - 1)^p}, \end{aligned}$$

where

$$g = (2^{-(j+1)/p})_{j=0}^m, \quad f = \underbrace{(2^{1/q} - 1, \dots, 2^{1/q} - 1)}_{m \text{ times}}, 1).$$

We have $\|g\|_p \leq 1$ and $\|f\|_p = (1 + m(2^{1/q} - 1)^p)^{1/p}$. Hence, by Minkowski's inequality,

$$\frac{(1 + m(2^{1/q} - 1)^p)^{1/p} - 1}{2^{1/p} - 1} \leq \left\| \sum_{j=1}^m h_{I_j}^{(p)} \right\|_p \leq \frac{(1 + m(2^{1/q} - 1)^p)^{1/p} + 1}{2^{1/p} - 1}. \quad (3.2)$$

Comparing (3.1) with (3.2), and letting m tend to ∞ , we get

$$\Delta_d[\mathcal{H}^{(p)}, L_p] \geq \max \left\{ \frac{2^{1/q} - 1}{2^{1/p} - 1}, \frac{2^{1/p} - 1}{2^{1/q} - 1} \right\},$$

as desired. \square

Next we establish the upper estimate for the super-bi-democracy constants that we will need.

Proposition 3.1. *If $1 < p < \infty$ then*

$$\Delta_{sb}[\mathcal{H}^{(p)}, L_p] \leq D_p := \frac{8}{(2^{1/p} - 1)(2^{1/p'} - 1)}. \quad (3.3)$$

Proof. For $I \in \mathcal{D}$ let $n(I)$ be such that $|I| = 2^{-n(I)}$. Let $A \subseteq \mathcal{D}$ finite and $\varepsilon = (\varepsilon_I)_{I \in A}$ be such that $|\varepsilon_I| = 1$ for all $I \in A$. For $J \in A$ set

$$R_J = J \setminus \cup \{I : I \in A, n(I) > n(J)\}.$$

Taking into account that, for $n \in \mathbb{N}$, the collection of dyadic intervals $\{I \in \mathcal{D} : n(I) = n\}$ is a partition on $[0, 1)$ we infer that

- $(R_I)_{I \in A}$ is a partition of $K = \cup_{I \in A} I$,
- $R_I \subseteq I$ for every $I \in A$, and
- given $t \in K$ and $k \in \mathbb{N}$ there is at most one interval $I_{t,k} \in A \cap \mathcal{D}_k$ such that $t \in I_{t,k}$; moreover $n(I_{t,k}) \leq n(J)$.

Consequently, for any $t \in K$ we have

$$\begin{aligned}
\left| \sum_{I \in A} \varepsilon_I h_I^{(p)}(t) \right| &\leq \sum_{I \in A} |h_I^{(p)}(t)| \\
&= \sum_{I \in A} 2^{n(I)/p} \chi_I(t) \\
&\leq \sum_{J \in A} \left(\sum_{n=-\infty}^{n(J)} 2^{n/p} \right) \chi_{R_J}(t) \\
&= \frac{1}{1 - 2^{-1/p}} \sum_{J \in A} 2^{n(J)/p} \chi_{R_J}(t).
\end{aligned}$$

Hence, if we set $a_p = 1/(1 - 2^{-1/p})$ we obtain

$$\begin{aligned}
\left\| \sum_{I \in A} \varepsilon_I h_I^{(p)} \right\|_p &\leq a_p \left(\sum_{J \in A} 2^{n(J)} |R_J| \right)^{1/p} \\
&\leq a_p \left(\sum_{J \in A} 2^{n(J)} |J| \right)^{1/p} \\
&= a_p |A|^{1/p}.
\end{aligned}$$

Therefore,

$$\left\| h_0^{(p)} + \sum_{I \in A} \varepsilon_I h_I^{(p)} \right\|_p \leq 1 + a_p |A|^{1/p}.$$

We infer that, for $m \in \mathbb{N}$,

$$\varphi_m^\varepsilon[\mathcal{H}^{(p)}, L_p] \leq \max\{a_p m^{1/p}, 1 + a_p(m-1)^{1/p}\} \leq 2a_p m^{1/p}.$$

The fact that $\varphi_m^{\varepsilon,*}[\mathcal{H}^{(p)}, L_p] = \varphi_m^\varepsilon[\mathcal{H}^{(p')}, L_{p'}]$ for $m \in \mathbb{N}$ yields

$$\frac{\varphi_m^\varepsilon[\mathcal{H}^{(p)}, L_p] \varphi_m^{\varepsilon,*}[\mathcal{H}^{(p)}, L_p]}{m} \leq 4a_p a_{p'} \frac{m^{1/p} m^{1/p'}}{m} = 4a_p a_{p'} = D_p.$$

Thus $\Delta_{sb}[\mathcal{H}^{(p)}, L_p] \leq D_p$. □

Corollary 3.2. *We have*

$$\begin{aligned}
\Delta_b[\mathcal{H}^{(p)}, L_p] &\approx \Delta_s[\mathcal{H}^{(p)}, L_p] \approx \Delta_{sd}[\mathcal{H}^{(p)}, L_p] \approx \Delta[\mathcal{H}^{(p)}, L_p] \approx \Delta_d[\mathcal{H}^{(p)}, L_p] \\
&\approx p^*
\end{aligned}$$

for $1 < p < \infty$.

Proof. Let D_p and d_p be as in Proposition 1.2 and Propostion 3.1. We have $d_p \approx D_p \approx p^*$ for $1 < p < \infty$. Then the result follows by combining Proposition 1.2, Propostion 3.1, and Remark 2.1. \square

We close by providing the conclusion of the proof of our main theorem and enunciating a corollary.

Conclusion of the Proof of Theorem 1.3. Combine Theorem 1.4 with the left-hand side of inequality (1.7), Theorem 3.1 and Theorem 1.1. \square

Corollary 3.3. $C_a[\mathcal{H}^{(p)}, L_p] \approx p^*$ for $1 < p < \infty$.

Proof. Just combine the left-hand side inequality in (1.10), with Corollary 3.2 and Theorem 1.3. \square

ANNEX: SUMMARY OF THE MOST COMMONLY EMPLOYED
CONSTANTS

Symbol	Name of constant	Ref. equation
C_a	Symmetry for largest coeffs. constant	(1.9)
C_g	Greedy constant	(1.3)
Δ	Democracy constant	(1.6)
Δ_b	Bi-democracy constant	(1.11)
Δ_d	Disjoint-democracy constant	(1.6))
Δ_s	Superdemocracy constant	(1.8)
Δ_{sb}	Super bi-democratic constant	(2.2)
Δ_{sd}	Disjoint-superdemocracy constant	(1.8)
K_{su}	Suppression unconditional constant	(1.4)
K_u	Lattice unconditional constant	(1.5)

Acknowledgments. F. Albiac and J. L. Ansorena acknowledge the support of the grant MTM2014-53009-P (MINECO, Spain). F. Albiac was also supported by the grant MTM2016-76808-P (MINECO, Spain). P. Berná was supported by a Ph.D. fellowship from the program FPI-UAM, as well as the grants MTM-2016-76566-P (MINECO, Spain) and 19368/PI/14 (*Fundación Séneca*, Región de Murcia, Spain).

REFERENCES

- [1] F. Albiac and J. L. Ansorena, *Characterization of 1-almost greedy bases*, Rev. Mat. Complut. **30** (2017), no. 1, 13–24.
- [2] F. Albiac and N. J. Kalton, *Topics in Banach space theory, 2nd revised and updated edition*, Graduate Texts in Mathematics, vol. 233, Springer International Publishing, 2016.
- [3] F. Albiac and P. Wojtaszczyk, *Characterization of 1-greedy bases*, J. Approx. Theory **138** (2006), no. 1, 65–86.
- [4] P. M. Berná, Ó. Blasco, and G. Garrigós, *Lebesgue inequalities for the greedy algorithm in general bases*, Rev. Mat. Complut. **30** (2017), no. 1, 369–392.
- [5] D. Burkholder, *A proof of Pelczyński’s conjecture for the Haar system*, Studia Mathematica **91** (1988), no. 1, 79–83.
- [6] S. J. Dilworth, N. J. Kalton, D. Kutzarova, and V. N. Temlyakov, *The thresholding greedy algorithm, greedy bases, and duality*, Constr. Approx. **19** (2003), no. 4, 575–597.
- [7] S. J. Dilworth, D. Kutzarova, E. Odell, Th. Schlumprecht, and A. Zsák, *Renorming spaces with greedy bases*, J. Approx. Theory **188** (2014), 39–56.
- [8] G. Garrigós, E. Hernández, and T. Oikhberg, *Lebesgue-type inequalities for quasi-greedy bases*, Constr. Approx. **38** (2013), no. 3, 447–470.
- [9] S. V. Konyagin and V. N. Temlyakov, *A remark on greedy approximation in Banach spaces*, East J. Approx. **5** (1999), no. 3, 365–379.
- [10] V. N. Temlyakov, *The best m -term approximation and greedy algorithms*, Adv. Comput. Math. **8** (1998), no. 3, 249–265.
- [11] ———, *Greedy approximation*, Acta Numer. **17** (2008), 235–409.

MATHEMATICS DEPARTMENT, UNIVERSIDAD PÚBLICA DE NAVARRA, CAMPUS DE ARROSADÍA, PAMPLONA, 31006 SPAIN
E-mail address: fernando.albiac@unavarra.es

DEPARTMENT OF MATHEMATICS AND COMPUTER SCIENCES, UNIVERSIDAD DE LA RIOJA, LOGROÑO, 26004 SPAIN
E-mail address: joseluis.ansorena@unirioja.es

PABLO M. BERNÁ, MATHEMATICS DEPARTMENT, UNIVERSIDAD AUTÓNOMA DE MADRID, MADRID, 28040 SPAIN.
E-mail address: pablo.berna@uam.es