

A NEW SURVEY: CONE METRIC SPACES

SUZANA ALEKSIĆ¹, ZORAN KADELBERG², ZORAN D. MITROVIĆ³ AND STOJAN
RADENOVIĆ⁴

¹ University of Kragujevac, Faculty of Sciences, Department of Mathematics and Informatics, Radoja Domanovića 12, 34000 Kragujevac, Serbia

² University of Belgrade, Faculty of Mathematics, Studentski trg 16, 11000 Beograd, Serbia

³ University of Banja Luka, Faculty of Electrical Engineering, Patre 5, 78000 Banja Luka, Bosnia and Herzegovina

⁴ University of Belgrade, Faculty of Mechanical Engineering, Department of Mathematics, Kraljice Marije 16, 11120 Belgrade 35, Serbia

E-mail: suzanasimic@kg.ac.rs (S. Aleksić), kadelbur@matf.bg.ac.rs (Z. Kadelburg), zmitrovic@etfbl.net (Z. D. Mitrović), radens@beotel.rs (S. Radenović)

Abstract. The purpose of this new survey paper is, among other things, to collect in one place most of the articles on cone (abstract, K-metric) spaces, published after 2007. This list can be useful to young researchers trying to work in this part of functional and nonlinear analysis. On the other hand, the existing review papers on cone metric spaces are updated.

The main contribution is the observation that it is usually redundant to treat the case when the underlying cone is solid and non-normal. Namely, using simple properties of cones and Minkowski functionals, it is shown that the problems can be usually reduced to the case when the cone is normal, even with the respective norm being monotone. Thus, we offer a synthesis of the respective fixed point problems arriving at the conclusion that they can be reduced to their standard metric counterparts. However, this does not mean that the whole theory of cone metric spaces is redundant, since some of the problems remain which cannot be treated in this way, which is also shown in the present article.

Key Words and Phrases: normal cone; non-normal cone; solid cone; non-solid cone; ordered topological vector space; cone metric space; Minkowski functional.

2010 Mathematics Subject Classification: 47H10, 54H25.

1. INTRODUCTION

Since 1922, when S. Banach proved in his PhD thesis (see [62]) the celebrated Contraction Principle for self-mappings on a complete metric space, several hundreds of researchers have tried to generalize or improve it. Basically, these generalizations were done in two directions—either the contractive condition was replaced by some more general one, or the environment of metric spaces was widened. In the first direction, a lot of improved results appeared, like Kannan’s, Chatterjea’s, Zamfirescu’s, Hardy-Rogers’s, Ćirić’s, Meir-Keeler’s, Boyd-Wong’s, to mention just a few. The other direction of investigations included introduction of semimetric, quasimetric, symmetric, partial metric, b -metric, and many other classes of spaces.

Of course, not all of these attempts were useful in applications, which should be the main motive for such investigations. Some of them were even not real generalizations, since the obtained results appeared to be equivalent to the already known ones.

It seems that Đ. Kurepa was the first to replace the set \mathbb{R} of real numbers by an arbitrary partially ordered set as a codomain of a metric (see [218]). This approach was used by several authors in mid-20th century, who used various names (abstract metric spaces, K -metric spaces, ...) for spaces thus obtained (see, e.g., [199, 210, 345, 363]). The applications were mostly in numerical analysis.

The interest in such spaces increased after the 2007 paper [153] of L. G. Huang and X. Zhang, who re-introduced the mentioned type of spaces under the name of *cone metric spaces*. Their approach included the use of interior points of the cone which defined the partial order of the space. The original paper [153] used also the assumption of normality of the cone, but later most of the results were obtained for non-normal cones, however with more complicated proofs.

It was spotted very early that (topological vector) cone metric spaces were metrizable (see, e.g., [114, 136, 190, 204]). However, this does not necessarily mean that all fixed point results in cone metric spaces reduce to their standard metric counterparts, since at least some of them still depend on the particular cone that is used. Hence, the research in these spaces has continued and more than 300 papers has appeared—most of them are cited in the list of references of the present paper (we probably skipped some of them, but it was not on purpose).

The intention of this article is, first, to recollect the basic properties of cones in (topological) vector spaces which are important for research in cone metric spaces. In particular (non)-normal and (non)-solid cones are investigated in some details. Then, it is shown that each solid cone in a topological vector space can be essentially replaced by a solid and normal cone in a normed space (even with normal constant equal to 1). Thus, most of the theory of (TVS) cone metric spaces can be easily reduced to the respective problems in standard metric spaces.

It has to be noted that some problems still remain to which the previous assertion cannot be applied—we will mention some of them in Section 4.

The present paper can also be treated as a continuation and refinement of the existing survey articles on this theme, like [170, 201, 273].

2. TOPOLOGICAL AND ORDERED VECTOR SPACES

We start our paper with some basic facts about ordered topological vector spaces. For more details, the reader may consult any book on this topic, e.g., [25, 101, 182, 314, 357].

2.1. Topological vector spaces. Throughout the paper, E will denote a vector space over the field \mathbb{R} of real numbers, with zero-vector denoted by θ . A topology τ on E is called a *vector topology* if linear operations $(x, y) \mapsto x + y$ and $(\lambda, x) \mapsto \lambda x$ are continuous on $E \times E$, resp. $\mathbb{R} \times E$. In this case, (E, τ) is called a (*real*) *topological vector space* (TVS). If a TVS has a base of neighbourhoods of θ whose elements are (closed and absolutely) convex sets in E , then it is called *locally convex* (LCS).

The topology of an LCS can be characterized using seminorms. Namely, a non-negative real function p on a vector space E is called a *seminorm* if: 1° $p(x + y) \leq p(x) + p(y)$ for $x, y \in E$; 2° $p(\lambda x) = |\lambda|p(x)$ for $x \in E, \lambda \in \mathbb{R}$. It is easy to prove that a function $p : E \rightarrow \mathbb{R}$ is a seminorm if and only if p is the *Minkowski functional* of an absolutely convex and absorbing set $M \subset E$, i.e., $p = p_M$ is given by

$$x \mapsto p_M(x) = \inf\{\lambda > 0 \mid x \in \lambda M\}.$$

Here, one can take $M = \{x \in E \mid p(x) < 1\}$. The following proposition provides the mentioned characterization.

Proposition 2.1. *Let $\{p_i \mid i \in I\}$ be an arbitrary family of seminorms on a vector space E . If $U_i = \{x \in E \mid p_i(x) < 1\}$, $i \in I$, then the family of all scalar multiples λU , $\lambda > 0$, where U runs through finite intersections $U = \bigcap_{j=1}^n U_j$, forms a base of the neighbourhoods of θ for a locally convex topology τ on E in which all the seminorms p_i are continuous. Each LCS (E, τ) can be described in such a way. If we put in the previous construction $U_i = \{x \in E \mid p_i(x) \leq 1\}$, we obtain a base of neighbourhoods of θ constituted of closed absolutely convex sets.*

The easiest (and well known) examples of locally convex spaces are normed and, in particular, Banach spaces. The topology is then generated by a single norm which is the Minkowski functional of the unit ball.

2.2. Cones in ordered topological vector spaces. Let (E, τ) be a real TVS. A subset C of E is called a *cone* if: 1° C is closed, nonempty and $C \neq \{\theta\}$; 2° $x, y \in C$ and $\lambda, \mu \geq 0$ imply that $\lambda x + \mu y \in C$; 3° $C \cap (-C) = \{\theta\}$. Obviously, each cone generates a partial order \preceq_C in E by

$$x \preceq_C y \iff y - x \in C$$

(we will write just \preceq if it is obvious which is the respective cone). And conversely, a partial order \preceq on E generates a cone $C = \{x \in E \mid x \succeq \theta\}$. If \preceq is a partial order on a TVS E , then (E, τ, \preceq) is called an *ordered topological vector space* (OTVS). If $x \preceq y$ and $x \neq y$, we will write $x \prec y$.

The most important examples of OTVS's are special cases of the following one.

Example 2.1. Let X be a nonempty set and E be the set of all real-valued functions f defined on X . Let a relation \preceq be defined on E by $f \preceq g$ if and only if $f(x) \leq g(x)$ for all $x \in X$. If τ is a vector topology on E such that the respective cone $C = \{f \in E \mid f(x) \geq 0, x \in X\}$ is τ -closed, then (E, τ, \preceq) is an OTVS.

If (E, \preceq) is an ordered vector space (with the respective cone C) and $x, y \in E$, $x \preceq y$, then the set

$$[x, y] = \{z \in E \mid x \preceq z \preceq y\} = (x + C) \cap (y - C)$$

is called an *order-interval*. A subset A of E is *order-bounded* if it is contained in some order-interval. It is *order-convex* if $[x, y] \subset A$ whenever $x, y \in A$ and $x \preceq y$. It has to be remarked that the families of *convex* and *order-convex* subsets of an ordered vector space are incomparable in general.

An OTVS (E, τ, \preceq) is said to be *order-convex* if it has a base of neighborhoods of θ consisting of order-convex subsets. In this case the order cone C is said to be *normal*. In the case of a normed space, this condition means that the unit ball is order-convex, which is equivalent to the condition that there is a number K such that $x, y \in E$ and $\theta \preceq x \preceq y$ imply that $\|x\| \leq K\|y\|$. The smallest constant K satisfying the last inequality is called the *normal constant* of C .

Equivalently, the cone C in $(E, \|\cdot\|, \preceq_C)$ is normal if the Sandwich Theorem holds, i.e., if

$$(2.1) \quad (\forall n) x_n \preceq y_n \preceq z_n \text{ and } \lim_{n \rightarrow \infty} x_n = \lim_{n \rightarrow \infty} z_n = x \text{ imply } \lim_{n \rightarrow \infty} y_n = x.$$

In particular, if a cone is normal with the constant equal to 1, i.e., if $\theta \preceq x \preceq y$ implies that $\|x\| \leq \|y\|$, the cone is called *monotone* w.r.t. the given norm.

The following is an example of a non-normal cone.

Example 2.2. [345] Let $E = C_{\mathbb{R}}^1[0, 1]$ with $\|x\| = \|x\|_{\infty} + \|x'\|_{\infty}$ and $C = \{x \in E \mid x(t) \geq 0, t \in [0, 1]\}$. Consider, for example, $x_n(t) = \frac{t^n}{n}$ and $y_n(t) = \frac{1}{n}$, $n \in \mathbb{N}$. Then $\theta \preceq x_n \preceq y_n$, and $\lim_{n \rightarrow \infty} y_n = \theta$, but $\|x_n\| = \max_{t \in [0, 1]} |\frac{t^n}{n}| + \max_{t \in [0, 1]} |t^{n-1}| = \frac{1}{n} + 1 > 1$; hence $\{x_n\}$ does not converge to zero. It follows by (2.1) that C is a non-normal cone.

The cone C in an OTVS (E, τ) is called *solid* if it has a nonempty interior $\text{int } C$.

There are numerous examples of cones which are solid and normal. Perhaps the easiest is the following one.

Example 2.3. Let $E = \mathbb{R}^n$ with the Euclidean topology τ . Let the cone C be given as in Example 2.1, i.e., $C = \{(x_i)_{i=1}^n \mid x_i \geq 0, i = 1, 2, \dots, n\}$. Then it is easy to see that C is a solid and normal cone in (E, τ) .

Example 2.2 shows that there exists a cone which is solid and non-normal. A lot of spaces which are important in functional analysis have cones which are normal and non-solid. We present just two of them.

Example 2.4. (see, e.g., [176]) Consider $E = c_0$ (the standard Banach space of all real 0-sequences) and let $C = \{x = \{x_n\} \mid x_n \geq 0, n \in \mathbb{N}\}$. It follows easily that C is a normal cone in E . Let us prove that it has an empty interior.

Take any $x = \{x_n\} \in C$. If $x_n = 0$ for $n \geq n_0$, then obviously $x \notin \text{int } C$. Otherwise, for arbitrary $\varepsilon > 0$, there exists $n_0 \in \mathbb{N}$ such that $x_n < \frac{\varepsilon}{2}$ for all $n > n_0$. Construct a sequence $y = \{y_n\}$ by

$$y_n = \begin{cases} x_n, & \text{if } n \leq n_0, \\ -x_n, & \text{for } n > n_0. \end{cases}$$

Then, at least one of y_n is strictly negative, hence $y \notin C$. However, $\|x - y\| = \sup_{n > n_0} |2x_n| < \varepsilon$, therefore, the ball $B(x, \varepsilon)$ is not a subset of C . Thus, $\text{int } C = \emptyset$.

Example 2.5. [242] Let $E = L^2(\mathbb{R}, \mu) \cap C(\mathbb{R})$ be equipped with the norm $\|\cdot\|_E = \|\cdot\|_{L^2} + \|\cdot\|_\infty$, where μ is the Lebesgue measure. Let

$$C = \{h \in E \mid h(t) \geq 0, t \in \mathbb{R}\}$$

be the cone of all positive elements in E . We will show that C has an empty interior. For any $f \in C$ and each $\delta > 0$, define $N_f(\delta) = \{t \in \mathbb{R} \mid f(t) \geq \delta\}$. Then $\mu(N_f(\delta)) < \infty$, and therefore $\mu(\mathbb{R} \setminus N_f(\delta)) = \infty$. Note that

$$\begin{aligned} \mu(\mathbb{R} \setminus N_f(\delta)) &= \mu\left(\bigcup_{r \in \mathbb{Q}} (B(r, \delta) \cap (\mathbb{R} \setminus N_f(\delta)))\right) \\ &\leq \sum_{r \in \mathbb{Q}} \mu(B(r, \delta) \cap (\mathbb{R} \setminus N_f(\delta))) = \infty, \end{aligned}$$

which means that $\mu(A) > 0$, where $A = B(s, \delta) \cap (\mathbb{R} \setminus N_f(\delta))$, for some $s \in \mathbb{Q}$. Define

$$g(x) = \begin{cases} f(x), & x \notin A, \\ f(x) - \delta, & x \in A. \end{cases}$$

Then g is negative on the set A of positive measure and

$$\|f - g\|_{L_2} + \|f - g\|_\infty = \left(\int_A \delta^2 d\mu\right)^{1/2} + \sup_A \delta \leq (2\delta^3)^{1/2} + \delta.$$

Hence, for any $f \in C$ and arbitrary $\varepsilon > 0$, we can find $g \notin C$ with $\|f - g\|_E < \varepsilon$, i.e., there is no ball with centre f that lies inside C . Therefore, C has an empty interior.

We state now the following properties of solid cones.

- [357, Proposition (2.2), page 20] $e \in \text{int } C$ if and only if $[-e, e] = (C - e) \cap (e - C)$ is a τ -neighbourhood of θ .

- If $e \in \text{int } C$, then the Minkowski functional $\|\cdot\|_e = p_{[-e, e]}$ is a norm on the vector space E . Indeed, one has just to prove that $\|x\|_e = 0$ implies that $x = \theta$. Suppose, to the contrary, that $x \neq \theta$. Then, by the definition of infimum, it follows that there is a sequence $\{\lambda_n\}$ of positive reals tending to 0, and such that $x \in \lambda_n[-e, e]$, i.e., $-\lambda_n e \preceq x \preceq \lambda_n e$. It follows that the sequences $\{\lambda_n e - x\}$ and $\{x + \lambda_n e\}$ both belong to C . Since they converge to $-x$ and x , respectively, the closedness of the cone C implies that $-x, x \in C$. Hence, $x = \theta$, which is a contradiction. See also [25, 167].

Note that this conclusion was stated in [200] [the paragraph before Theorem 3.1] under the additional assumption that the cone C is normal. The previous proof shows that this assumption is in fact redundant.

- If $e_1, e_2 \in \text{int } C$, then the respective norms $\|\cdot\|_{e_1}$ and $\|\cdot\|_{e_2}$ are equivalent. In particular, the interiors of the cone C w.r.t. the norms $\|\cdot\|_{e_1}$ and $\|\cdot\|_{e_2}$ coincide; in fact, they are equal to $\text{int}_\tau C$. This follows by the characterization of interior points of a cone in an OTVS.

- If $\text{int}_\tau C \neq \emptyset$, then the topology τ is Hausdorff. Indeed, if $e \in \text{int } C$, then the unit ball $[-e, e]$ is a neighbourhood of θ in the topology generated by the norm

$\|\cdot\|_e$. Since this norm topology is Hausdorff and not stronger than τ , it follows that τ is Hausdorff, too.

We will show now that C is a cone, i.e., it is closed, in the topology generated by $\|\cdot\|_e$. Let $x_n \in C$ and $x_n \rightarrow x$ as $n \rightarrow \infty$, in the norm $\|\cdot\|_e$. This means that $-x \preceq -(x - x_n) \preceq \frac{1}{n}e$ for $n \geq n_0$. Passing to the limit as $n \rightarrow \infty$, we get that $-x \preceq \theta$, i.e., $x \succeq \theta$, and so $x \in C$.

Thus, we have proved the next assertion (see also [22, 101, 180, 284, 285, 346]).

Proposition 2.2. *If C is a solid cone in an OTVS (E, τ, \preceq_C) , then there exists a norm $\|\cdot\|_e$ on E which is monotone and such that the cone C is normal and solid w.r.t. this norm. In particular, the cone C has the same set of interior points, both in topology τ and in the norm generated by $\|\cdot\|_e$.*

Further, following [153], denote

$$(2.2) \quad x \ll y \quad \text{if and only if} \quad y - x \in \text{int } C.$$

Let $\{x_n\}$ be a sequence in (E, τ, \preceq_C) and $c \in \text{int } C$. We will say that $\{x_n\}$ is a c -sequence if there exists $n_0 \in \mathbb{N}$ such that $x_n \ll c$ whenever $n \geq n_0$. The connection between c -sequences and sequences tending to θ is the following.

- If $x_n \rightarrow \theta$ as $n \rightarrow \infty$, then it is a c -sequence for each $c \in \text{int } C$. Indeed, if $c \in \text{int } C$, then $[-c, c]$ is a τ -neighbourhood of θ and it follows that $\text{int}[-c, c] = (\text{int } C - c) \cap (c - \text{int } C)$ is also a τ -neighbourhood of θ . Hence, there exists $n_0 \in \mathbb{N}$ such that, for all $n \geq n_0$, $x_n \in (\text{int } C - c) \cap (c - \text{int } C)$, i.e., $c - x_n \in \text{int } C$, and so $x_n \ll c$.

- The converse of the previous assertion is not true, i.e., $x_n \ll c$ for some $c \in \text{int } C$ and all $n \geq n_0$ does not necessarily imply that $x_n \rightarrow \theta$ (see Example 2.2). However, if the cone C is normal, then these two properties of a sequence $\{x_n\}$ are equivalent.

Note also the following properties of bounded sets.

- If the cone C is solid, then each topologically bounded subset of (E, τ, \preceq_C) is also order-bounded, i.e., it is contained in a set of the form $[-c, c]$ for some $c \in \text{int } C$.

- If the cone C is normal, then each order-bounded subset of (E, τ, \preceq_C) is topologically bounded. Hence, if the cone is both solid and normal, these two properties of subsets of E coincide.

In other words, we have the following property (see, e.g., [345]).

Proposition 2.3. *If the underlying cone of an OTVS is solid and normal, then such TVS must be an ordered normed space.*

The following old result of M. Krein can also be useful when dealing with cones in normed spaces.

Proposition 2.4. [211] *A cone C in a normed space $(E, \|\cdot\|)$ is normal if and only if there exists a norm $\|\cdot\|_1$ on E , equivalent to the given norm $\|\cdot\|$, such that the cone C is monotone w.r.t. $\|\cdot\|_1$.*

3. CONE METRIC SPACES

3.1. Definition and basic properties. As was already said in Introduction, spaces with “metrics” having values in ordered spaces, more general than the set \mathbb{R} of real numbers, appeared under various names (abstract metric spaces, K -metric spaces, ...) since the mid-20th century (see, e.g., [199, 210, 218, 345, 363]). Starting from 2007 and the paper [153], some version of the following definition has been usually used.

Definition 3.1. *Let X be a non-empty set and (E, τ, \preceq) be an OTVS. If a mapping $d : X \times X \rightarrow E$ satisfies the conditions*

- (i) $d(x, y) \succeq \theta$ for all $x, y \in X$ and $d(x, y) = \theta$ if and only if $x = y$;
- (ii) $d(x, y) = d(y, x)$ for all $x, y \in X$;
- (iii) $d(x, z) \preceq d(x, y) + d(y, z)$ for all $x, y, z \in X$,

then d is called a cone metric on X and (X, d) is called a cone metric space.

Convergent and Cauchy sequences can be introduced in the following way.

Definition 3.2. *Let (X, d) be a cone metric spaces with the cone metric having values in (E, τ, \preceq) , where the underlying cone C is solid. Let $\{x_n\}$ be a sequence in X . Then we say that:*

- (1) *the sequence $\{x_n\}$ converges to $x \in X$ if for each $c \in \text{int } C$ there exists $n_0 \in \mathbb{N}$ such that $d(x_n, x) \ll c$ holds for all $n \geq n_0$;*
- (2) *$\{x_n\}$ is a Cauchy sequence if for each $c \in \text{int } C$ there exists $n_0 \in \mathbb{N}$ such that $d(x_m, x_n) \ll c$ holds for all $m, n \geq n_0$;*
- (3) *the space (X, d) is complete if each Cauchy sequence $\{x_n\}$ in it converges to some $x \in X$.*

Note that, according to [180], in the previous definition, equivalent notions are obtained if one replace \ll by \prec or \preceq .

The following crucial observation (which is a consequence of Proposition 2.2) shows that most of the problems in cone metric spaces can be reduced to their standard metric counterparts.

Proposition 3.1. *Let X be a non-empty set and (E, τ, \preceq_C) be an OTVS with a solid cone C . If $e \in \text{int } C$ is arbitrary, then $D(x, y) = \|d(x, y)\|_e$ is a (standard) metric on X . Moreover, a sequence $\{x_n\}$ is Cauchy (convergent) in (X, d) if and only if it is Cauchy (convergent) in (X, D) .*

We will show in Section 4 how this approach simplifies most of the proofs concerning fixed points of mappings.

Note that it is clear that the cone metric with values in $(E, \|\cdot\|_e, \preceq)$ is continuous (as a function of two variables). Also, Sandwich Theorem holds (since the cone C is normal in the new norm).

3.2. Completion and Cantor’s intersection theorem. Most of standard notions from the setting of metric spaces, like accumulation points of sequences, open and closed balls, open and closed subsets, etc., are introduced in the usual way. Also, standard properties (e.g., that closed balls are closed subsets) can be easily deduced, based on Proposition 2.2.

As a sample, we state the following theorem on the completion of a cone metric space and we note that, with the mentioned approach, its proof is much easier than in [11].

Theorem 3.1. *Let (X, d) be a cone metric space over an OTVS (E, τ, \preceq_C) with a solid cone. Then there exists a complete cone metric space (\tilde{X}, \tilde{d}) over an ordered normed space $(E, \|\cdot\|, \preceq_C)$ with a solid and normal cone, containing a dense subset $(\tilde{X}^*, \tilde{d}^*)$, isometric with (X, d) .*

Now, we present a proof of the Cantor's intersection theorem in the setting of cone metric spaces.

Theorem 3.2. *Let (X, d) be a cone metric space over an OTVS (E, τ, \preceq_C) with a solid cone. Then (X, d) is complete if and only if every decreasing sequence $\{B_n\}$ of non-empty closed balls in X for which the sequence of diameters tends to θ as $n \rightarrow \infty$, has a non-empty intersection, and more precisely, there exists a point $x \in X$ such that $\bigcap_{n=1}^{\infty} B_n = \{x\}$.*

Proof. According to Propositions 2.2 and 2.4, without loss of generality, we can assume that (X, d) is a cone metric space over an ordered normed space $(E, \|\cdot\|, \preceq_C)$ with the cone C being solid and normal. Moreover, we can assume that the normal constant of C is equal to 1 (i.e., the norm is monotone w.r.t. C).

Assume that (X, d) is a complete cone metric space, and let $B_n = B[x_n, c_n]$, $n \in \mathbb{N}$, be the sequence of non-empty, closed balls in X with the properties: $B_{n+1} \subset B_n$, $n \in \mathbb{N}$, and $c_n \rightarrow \theta$ as $n \rightarrow \infty$. We first show that $\{x_n\}$ is a Cauchy sequence.

If $m > n$, we have $x_m \in B_m \subset B_n$ which implies that $d(x_m, x_n) \preceq c_n$. Since $c_n \rightarrow \theta$ as $n \rightarrow \infty$, it follows that $d(x_m, x_n) \rightarrow \theta$ as $n \rightarrow \infty$, and consequently $\{x_n\}$ is a Cauchy sequence.

The completeness of (X, d) implies the existence of $x \in X$ such that $x_n \rightarrow x$ as $n \rightarrow \infty$. Since $x_m \in B_n$ for all $m \geq n$, and B_n is a closed subset of X , it follows that $x \in B_n$ for all $n \in \mathbb{N}$, i.e., $x \in \bigcap_{n=1}^{\infty} B_n$.

Suppose that there exists $y \in \bigcap_{n=1}^{\infty} B_n$ such that $y \neq x$. We thus get

$$d(x, y) \preceq d(x, x_n) + d(x_n, y) \preceq 2c_n,$$

which gives $d(x, y) \preceq \theta$ and, as a consequence, we obtain $x = y$, a contradiction. Hence, $\bigcap_{n=1}^{\infty} B_n = \{x\}$.

Conversely, to obtain a contradiction, suppose that (X, d) is not a complete cone metric space. Then there exists a Cauchy sequence $\{x_n\}$ which does not converge in X .

We construct a strictly increasing sequence of positive integers $\{n_i\}$ in the following way. If $c_1 \gg \theta$, then there exists $n_1 \in \mathbb{N}$ such that $m > n_1$ implies $d(x_m, x_{n_1}) \ll c_1$. Now, for $c_2 = \frac{c_1}{2}$ there exists $n_2 > n_1$ such that $d(x_m, x_{n_2}) \ll c_2$ holds for all $m > n_2$. We continue in this manner. For $c_i = \frac{c_{i-1}}{2}$, there exists $n_i > n_{i-1}$ such that $m > n_i$ implies $d(x_m, x_{n_i}) \ll c_i$.

Let us consider the sequence $B_i = B[x_{n_i}, 2c_i]$, $i \in \mathbb{N}$, such that $c_i = \frac{c_1}{2^{i-1}} \rightarrow \theta$ as $i \rightarrow \infty$. We first show that $B_{i+1} \subset B_i$ for all $i \in \mathbb{N}$. From $y \in B_{i+1}$, it follows that $d(y, x_{n_{i+1}}) \preceq 2c_{i+1}$, hence that

$$d(y, x_{n_i}) \preceq d(y, x_{n_{i+1}}) + d(x_{n_{i+1}}, x_{n_i}) \preceq 2c_{i+1} + c_i = 2\frac{c_i}{2} + c_i = 2c_i,$$

and finally that $y \in B[x_{n_i}, 2c_i]$. We thus get $y \in B_i$, which is the desired conclusion.

Let us prove that $\bigcap_{i=1}^{\infty} B_i = \emptyset$. Indeed, if there exists an $x \in \bigcap_{i=1}^{\infty} B_i$, then $x \in B_i$ for all $i \in \mathbb{N}$ and thus $d(x, x_i) \preceq 2c_i$, $i \in \mathbb{N}$. For $m > n_i$, we have

$$d(x, x_m) \preceq d(x, x_{n_i}) + d(x_{n_i}, x_m) \preceq 2c_i + c_i = 3c_i.$$

The normality of the cone C gives $d(x, x_m) \rightarrow \theta$ as $m \rightarrow \infty$ (since $m > n_i$). This contradicts the fact that the Cauchy sequence $\{x_n\}$ does not converge in X . \square

Comparing the previous proof with the one of [167, Theorem 3.10]), we see that our result is more general and with a shorter proof.

4. FIXED POINT RESULTS IN CONE METRIC SPACES

4.1. Some fixed point results. Most of the papers from the present list of references dealt with some (common) fixed point problems in cone metric spaces. However, the majority of the obtained results are in fact direct consequences of the corresponding results from the standard metric spaces. We will show this just on certain examples, but it will be clear that the same procedure can be done in most of the other cases.

Thus, let (X, d) be a cone metric space over an OTVS (E, τ, \preceq) with a solid cone C . Let $f, g : X \rightarrow X$ be two self-mappings satisfying $fX \subset gX$, and assume that one of these two subsets of X is complete. Consider the following contractive conditions:

- (1) (Banach) $d(fx, fy) \preceq \lambda d(gx, gy)$, $\lambda \in (0, 1)$;
- (2) (Kannan) $d(fx, fy) \preceq \lambda d(gx, fx) + \mu d(gy, fy)$, $\lambda, \mu \geq 0$, $\lambda + \mu < 1$;
- (3) (Chatterjea) $d(fx, fy) \preceq \lambda d(gx, fy) + \mu d(gy, fx)$, $\lambda, \mu \geq 0$, $\lambda + \mu < 1$;
- (4) (Reich) $d(fx, fy) \preceq \lambda d(gx, gy) + \mu d(gx, fx) + \nu d(gy, fy)$, $\lambda, \mu, \nu \geq 0$, $\lambda + \mu + \nu < 1$;
- (5) (Zamfirescu) one of the conditions

$$\begin{cases} d(fx, fy) \preceq \lambda d(gx, gy), & \lambda \in (0, 1) \\ d(fx, fy) \preceq \lambda d(gx, fx) + \mu d(gy, fy), & \lambda, \mu \geq 0, \lambda + \mu < 1 \\ d(fx, fy) \preceq \lambda d(gx, fy) + \mu d(gy, fx), & \lambda, \mu \geq 0, \lambda + \mu < 1; \end{cases}$$

- (6) (Hardy-Rogers) $d(fx, fy) \preceq \lambda d(gx, gy) + \mu d(gx, fx) + \nu d(gy, fy) + \xi d(gx, fy) + \pi d(gy, fx)$, $\lambda, \mu, \nu, \xi, \pi \geq 0$, $\lambda + \mu + \nu + \xi + \pi < 1$;
- (7) (Ćirić) $d(fx, fy) \preceq \lambda u(x, y)$, for some $u(x, y)$ belonging to one of the following sets

$$\begin{cases} \left\{ d(gx, gy), \frac{d(gx, fx) + d(gy, fy)}{2}, \frac{d(gx, fy) + d(gy, fx)}{2} \right\} \\ \left\{ d(gx, gy), d(gx, fx), d(gy, fy), \frac{d(gx, fy) + d(gy, fx)}{2} \right\} \end{cases}$$

where $\lambda \in (0, 1)$.

Theorem 4.1. *If one of the conditions (1)–(7) is satisfied for all $x, y \in X$, then the mappings f and g have a unique point of coincidence. Moreover, if f and g are weakly compatible, then they have a unique common fixed point in X .*

Proof. Let $e \in \text{int } C$ be arbitrary. Based on Propositions 2.2, 2.4 and 3.1, consider the metric space (X, D) , where $D(x, y) = \|d(x, y)\|_e$. Then, the self-mappings $f, g : X \rightarrow X$ satisfy one of the following contractive conditions:

- (1') $D(fx, fy) \leq \lambda D(gx, gy)$, $\lambda \in (0, 1)$;
- (2') $D(fx, fy) \leq \lambda D(gx, fx) + \mu D(gy, fy)$, $\lambda, \mu \geq 0$, $\lambda + \mu < 1$;
- (3') $D(fx, fy) \leq \lambda D(gx, fy) + \mu D(gy, fx)$, $\lambda, \mu \geq 0$, $\lambda + \mu < 1$;
- (4') $D(fx, fy) \leq \lambda D(gx, gy) + \mu D(gx, fx) + \nu D(gy, fy)$, $\lambda, \mu, \nu \geq 0$, $\lambda + \mu + \nu < 1$;

(5') one of the conditions

$$\begin{cases} D(fx, fy) \leq \lambda D(gx, gy), & \lambda \in (0, 1) \\ D(fx, fy) \leq \lambda D(gx, fx) + \mu D(gy, fy), & \lambda, \mu \geq 0, \lambda + \mu < 1 \\ D(fx, fy) \leq \lambda D(gx, fy) + \mu D(gy, fx), & \lambda, \mu \geq 0, \lambda + \mu < 1; \end{cases}$$

(6') $D(fx, fy) \leq \lambda D(gx, gy) + \mu D(gx, fx) + \nu D(gy, fy) + \xi D(gx, fy) + \pi D(gy, fx)$,
 $\lambda, \mu, \nu, \xi, \pi \geq 0$, $\lambda + \mu + \nu + \xi + \pi < 1$;

(7') one of the conditions

$$D(fx, fy) \leq \lambda \left\{ D(gx, gy), \frac{D(gx, fx) + D(gy, fy)}{2}, \frac{D(gx, fy) + D(gy, fx)}{2} \right\}$$

$$D(fx, fy) \leq \lambda \left\{ D(gx, gy), D(gx, fx), D(gy, fy), \frac{D(gx, fy) + D(gy, fx)}{2} \right\}$$

where $\lambda \in (0, 1)$.

It is well known (see, e.g., [304] for the case $g = id_X$; the general cases are similar) that, in each of these cases, it follows that the mappings f and g have a unique point of coincidence. The conclusion in the case of weak compatibility is also standard (see, e.g., [177]). \square

For Ćirić's quasicontraction (see also [304]), but for a single mapping, the following cone metric version can be proved in a similar way.

Theorem 4.2. *Let (X, d) be a complete cone metric space over an OTVS (E, τ, \preceq) with a solid cone C . Let $f : X \rightarrow X$ be a self-mapping such that there exists $\lambda \in (0, 1)$ and, for all $x, y \in X$ there exists*

$$u(x, y) \in \{d(x, y), d(x, fx), d(y, fy), d(x, fy), d(y, fx)\}$$

satisfying $d(fx, fy) \preceq \lambda u(x, y)$. Then f has a unique fixed point in X .

4.2. Results that cannot be obtained in the mentioned way. Certain fixed point problems in cone metric spaces cannot be treated as in the previous subsection. This is always the case when the given contractive condition cannot be transformed to an appropriate condition in the corresponding metric space (X, D) . As an example of this kind we recall a Boyd-Wong type result.

Let (E, τ, \preceq_C) be an OTVS with a solid cone. A mapping $\varphi : C \rightarrow C$ is called a *comparison function* if: (i) φ is nondecreasing w.r.t. \preceq ; (ii) $\varphi(\theta) = \theta$ and $\theta \prec \varphi(c) \prec c$ for $c \in C \setminus \{\theta\}$; (iii) $c \in \text{int } C$ implies $c - \varphi(c) \in \text{int } C$; (iv) $c \in C \setminus \{\theta\}$ and $e \in \text{int } C$, then there exists $n_0 \in \mathbb{N}$ such that $\varphi^n(c) \ll e$ for each $n \geq n_0$.

Theorem 4.3. [31, Theorem 2.1] *Let (X, d) be a complete cone metric space over (E, τ, \preceq_C) with a solid cone. Let the mappings $f, g : X \rightarrow X$ be weakly compatible and suppose that for some comparison function $\varphi : C \rightarrow C$ and for all $x, y \in X$ there exists $u \in \{d(gx, gy), d(gx, fx), d(gy, fy)\}$, such that $d(fx, fy) \preceq \varphi(u)$. Then f and g have a unique common fixed point in X .*

An additional example is provided by the following result.

Theorem 4.4. [99] *Let (X, d) be a complete cone metric space over $(E, \|\cdot\|, \preceq_C)$ with a solid cone. Let $f : X \rightarrow X$ and suppose that there exists a positive bounded operator on E with the spectral radius $r(A) < 1$, such that $d(fx, fy) \preceq A(d(x, y))$ for all $x, y \in X$. Then f has a unique fixed point in X .*

In this case, passing to the space $(E, \|\cdot\|_e, \preceq_C)$ cannot help, since the two norms need not be equivalent and we have no information about the boundedness of A and (if it is bounded) how $r(A)$ has changed.

A very important special class of fixed point results are the Caristi-type ones, and the huge literature is devoted to them. When cone metric spaces are concerned, they are treated in details, e.g., in the papers [19, 85, 193, 200–202, 353], so we will give here only some basic information.

The following result was proved in [19].

Theorem 4.5. [19, Theorem 2] *Let (X, d) be a complete cone metric space over an OTVP $(E, \|\cdot\|, \preceq_C)$ with a solid cone. Let $F : X \rightarrow C$ be a lower-semi-continuous map and let $f : X \rightarrow X$ satisfy the condition*

$$d(x, fx) \preceq F(x) - F(f(x))$$

for any $x \in X$. Then f has a fixed point.

It is not clear whether, by passing to the space $(E, \|\cdot\|_e, \preceq)$ using the Minkowski functional, this theorem reduces to the classical Caristi result. However, it is proved in [202] that this is still the case if $e \in \text{int } C$ can be chosen so that the corresponding Minkowski functional $p_{[-e, e]}$ is additive. It is an open question whether this can be done without this assumption.

5. SOME “HYBRID” SPACES

As has been already mentioned, besides cone metric spaces, a lot of other generalizations of standard metric spaces have been introduced and fixed point problems in these settings have been investigated by several authors. There are among them some “hybrid” spaces, i.e., spaces where axioms of several types are used simultaneously. Such are, e.g., cone b -metric spaces, cone- G -metric spaces, cone rectangular metric spaces, partial cone metric spaces, cone spaces with c -distance and some others. We will recall here definitions of two of these classes and how the work in these classes can be easily reduced to the already known ones. Similar conclusions hold for the other mentioned classes, too.

5.1. Cone b -metric spaces. Cone b -metric spaces were introduced (under a different name) in [97] and some fixed point results were presented in this setting.

Definition 5.1. Let X be a non-empty set, (E, τ, \preceq) be an OTVS and $s \geq 1$ be a real number. If a mapping $d : X \times X \rightarrow E$ satisfies the conditions

- (i) $d(x, y) \succeq \theta$ for all $x, y \in X$ and $d(x, y) = \theta$ if and only if $x = y$;
- (ii) $d(x, y) = d(y, x)$ for all $x, y \in X$;
- (iii) $d(x, z) \preceq s(d(x, y) + d(y, z))$ for all $x, y, z \in X$,

then d is called a cone b -metric on X and (X, d) is called a cone b -metric space with parameter s .

Later, these and other researchers obtained a lot of (common) fixed point results (see, [57, 63, 131, 137, 146, 234, 263, 323, 334, 360, 366]). However, similarly as in the case of cone metric spaces, most of these results can be reduced to the case of classical b -metric spaces of Bakhtin [61] and Czerwik [100]. Namely, similarly as Proposition 3.1, the following can be deduced from Proposition 2.2.

Proposition 5.1. Let X be a non-empty set, (E, τ, \preceq_C) be an OTVS with a solid cone C and $s \geq 1$ be a real number. If $e \in \text{int } C$ is arbitrary, then $D(x, y) = \|d(x, y)\|_e$ is a b -metric on X . Moreover, a sequence $\{x_n\}$ is Cauchy (convergent) in (X, d) if and only if it is Cauchy (convergent) in (X, D) .

5.2. Cone rectangular spaces. The following definition was given in [50].

Definition 5.2. Let X be a non-empty set and (E, τ, \preceq) be an OTVS. If a mapping $d : X \times X \rightarrow E$ satisfies the conditions

- (i) $d(x, y) \succeq \theta$ for all $x, y \in X$ and $d(x, y) = \theta$ if and only if $x = y$;
- (ii) $d(x, y) = d(y, x)$ for all $x, y \in X$;
- (iii) $d(x, y) \preceq d(x, w) + d(w, z) + d(z, y)$ for all $x, y \in X$ and all distinct points $w, z \in X \setminus \{x, y\}$,

then d is called a cone rectangular metric on X and (X, d) is called a cone rectangular metric space.

In the papers [50, 174, 235, 238, 298, 339, 346], some fixed point results in cone rectangular spaces were obtained. The following proposition shows how most of these results can be easily reduced to the corresponding known ones in the environment of rectangular spaces defined by Branciari in [73].

Proposition 5.2. Let X be a non-empty set and (E, τ, \preceq_C) be an OTVS with a solid cone C . If $e \in \text{int } C$ is arbitrary, then $D(x, y) = \|d(x, y)\|_e$ is a rectangular metric on X . Moreover, a sequence $\{x_n\}$ is Cauchy (convergent) in (X, d) if and only if it is Cauchy (convergent) in (X, D) .

6. THE CASE OF A NON-SOLID CONE

In this section, we will briefly consider the case when the underlying cone is non-solid, but normal.

Let $(E, \|\cdot\|, \preceq_C)$ be an ordered normed space and let the cone C be normal, but non-solid (see, e.g., Examples 2.4 and 2.5). The definition of a cone metric d on a nonempty set X , over E , is the same as Definition 3.1. However, definition of convergent and Cauchy sequences, as well as of the completeness, must be modified.

Definition 6.1. Let (X, d) be a cone metric space over $(E, \|\cdot\|, \preceq_C)$, with the cone C being normal (possibly non-solid). A sequence $\{x_n\}$ in X is said to converge to $x \in X$ if $d(x_n, x) \rightarrow \theta$ as $n \rightarrow \infty$ (the convergence in the sense of norm in E). Similarly, Cauchy sequences are introduced, as well as the completeness of the space.

In the view of Proposition 2.4, we can assume that the normal constant of C is equal to 1, and further consider just the standard metric space (X, D) , where $D(x, y) = \|d(x, y)\|$.

Open (resp. closed) balls in (X, d) can be introduced as follows: let $x \in X$ and $c \in C \setminus \{\theta\}$. The open (resp. closed) ball with centre x and radius c are given by $B(x, c) = \{y \in X \mid d(y, x) \prec c\}$ (resp. $B(x, c) = \{y \in X \mid d(y, x) \preceq c\}$). It can be easily shown that an open (closed) ball is an open (closed) subset of (X, d) . We will show this, e.g., in the case of a closed ball.

Thus, let $y_n \in B[x, c]$ and $y_n \rightarrow y$ as $n \rightarrow \infty$ in (X, d) . Then we have $d(y, x) \preceq d(y, y_n) + d(y_n, x) \preceq d(y, y_n) + c$. By the Sandwich Theorem (which holds since the cone C is normal), we get that $d(y, x) \preceq c$, which means that $y \in B(x, c)$.

However, note that the collection of open balls $\{B(x, c) \mid x \in X, c \in C \setminus \{\theta\}\}$ does not necessarily form a base of topology on X . Indeed, it is not sure that the intersection of two open balls contains a ball. For example, for $X = \mathbb{R}^2$, $C = \{(x, y) \mid x, y \geq 0\}$ and the points $c_1 = (1, 0)$, $c_2 = (0, 1)$ from $C \setminus \{\theta\}$, there is no $c \in C \setminus \{\theta\}$ such that $c \prec c_1$ and $c \prec c_2$.

Nevertheless, the theorem on completion holds in this case, too.

We conclude with the following open question.

Question 6.1. Construct a cone in a real TVS which is both non-normal and non-solid.

Acknowledgement. The first and second author are grateful to the Ministry of Education, Science and Technological Development of Serbia, Grants No. 174024 and 174002, respectively.

REFERENCES

- [1] Ch. T. Aage, J. N. Salunke, *Some fixed point theorems for expansion onto mappings on cone metric spaces*, Acta Math. Sinica, English Ser., 27 (6) (2011), 1101–1106.
- [2] M. Abbas, Y. J. Cho, T. Nazir, *Common fixed point theorems for four mappings in TVS-valued cone metric spaces*, J. Math. Inequal. 5 (2) (2011), 287–299.
- [3] M. Abbas, M. Ali Khan, S. Radenović, *Common coupled fixed point theorems in cone metric spaces for w -compatible mappings*, Appl. Math. Comput. 217 (2010), 195–202.
- [4] M. Abbas, I. Altun, *Common fixed point results in ordered cone metric spaces*, Hacettepe J. Math. Stat, 42 (2013), 525–531.
- [5] M. Abbas, M. Jovanović, S. Radenović, A. Sretenović, and S. Simić, *Abstract metric spaces and approximating fixed points of a pair of contractive type mappings*, J. Comput. Anal. Appl. 13, (2) (2011), 243–253.
- [6] M. Abbas, G. Jungck, *Common fixed point results for noncommuting mappings without continuity in cone metric spaces*, J. Math. Anal. Appl. 341 (1) (2008), 416–420.
- [7] M. Abbas, B. E. Rhoades, *Fixed and periodic point results in cone metric spaces*, Appl. Math. Lett. 22 (2009), 511–515.
- [8] M. Abbas, B. E. Rhoades, T. Nazir, *Common fixed points for four maps in cone metric spaces*, Appl. Math. Comput. 216 (1) (2010), 80–86.
- [9] M. Abbas, B. E. Rhoades, T. Nazir, *Common fixed points of generalized contractive multi-valued mappings in cone metric spaces*, Math. Commun. 14 (2) (2009), 365–378.

- [10] M. Abbas, P. Vetro, S. F. Khan, *On fixed points of Berinde's contractive mappings in cone metric spaces*, Carpathian J. Math. 26 (2) (2010), 121–133.
- [11] Th. Abdeljawad, *Completion of cone metric spaces*, Hacet. J. Math. Stat. 39 (1) (2010), 67–74.
- [12] Th. Abdeljawad, *Order norm completions of cone metric spaces*, Numer. Funct. Anal. Optim. 32 (5) (2011), 477–495.
- [13] Th. Abdeljawad, *Edelstein-type fixed point theorems in compact TVS-cone metric spaces*, Hacet. J. Math. Stat. 43 (4) (2014), 561–569.
- [14] T. Abdeljawad, E. Karapinar, *Quasiconic metric spaces and generalizations of Caristi-Kirk's theorem*, Fixed Point Theory Appl. Vol. 2009, Article ID 574387, 9 pages.
- [15] T. Abdeljawad, D. Turkoglu, M. Abuloha, *Some theorems and examples of cone Banach spaces*, J. Comput. Anal. Appl. 12 (4) (2010), 739–753.
- [16] T. Abdeljawad, Sh. Rezapour, *Some fixed point results in TVS-cone metric spaces*, Fixed Point Theory, 14 (2) (2013), 263–268.
- [17] S. M. Abusalim, M. S. M. Noorani, *Fixed point and common fixed point theorems on ordered cone b-metric spaces*, Abstract Appl. Math., Vol. 2013, Article ID 815289, 7 pages.
- [18] S. M. Abusalim, M. S. M. Noorani, *Tripled fixed point theorems in cone metric spaces under F-invariant set and c-distance*, J. Nonlinear Sci. Appl. 8 (5) (2015), 750–762.
- [19] R. P. Agarwal, M. A. Khamsi, *Extension of Caristi's fixed point theorem to vector valued metric spaces*, Nonlinear Anal. 74 (2011), 141–145.
- [20] R.P. Agarwal, W. Sintunavarat, P. Kumam, *Coupled coincidence point and common coupled fixed point theorems lacking the mixed monotone property*, Fixed Point Theory Appl., 2013:22 (2013).
- [21] A. G. B. Ahmed, Z. M.Fadail, M. Abbas, Z. Kadelburg, S. Radenović, *Some fixed and periodic point results in abstract metric spaces*, Abstr. Appl. Anal. Vol. 2012, Article ID 908423.
- [22] S. Aleksić, Lj. Paunović, S. Radenović, F. Vetro, *Some critical remarks on the paper "A note on the metrizable of TVS-cone metric spaces"*, Vojno-tehnički glasnik/ Military Technical Courier 65 (3) (2017), 1–8.
- [23] M. A. Alghamdi, S. H. Alnafe'i, S. Radenović, N. Shahzad, *Fixed point theorems for convex contraction mappings on cone metric spaces*, Math. Comput. Modell. 54 (2011), 2020–2026.
- [24] M. Alimohammady, J. Balooee, S. Radojević, V. Rakočević, M. Roohi, *Conditions of regularity in cone metric spaces*, Appl. Math. Comput. 217 (2011), 6359–6363.
- [25] C. D. Aliprantis, R. Tourky, *Cones and Duality*, Grad. Stud. Math. 84, Amer. Math. Soc., Providence, RI, 2007.
- [26] S. H. Alnafe'i, S. Radenović, N. Shahzad, *Fixed point theorems for mappings with convex diminishing diameters on cone metric spaces*, Appl. Math. Lett. 24 (2011), 2162–2166.
- [27] I. Altun, B. Damjanović, D. Đorić, *Fixed point and common fixed point theorems on ordered cone metric spaces*, Appl. Math. Lett. 23 (2010), 310–316.
- [28] I. Altun, G. Durmaz, *Some fixed point theorems in ordered cone metric spaces*, Rend. Circ. Mat. Palermo 58 (2009), 319–325.
- [29] I. Altun, V. Rakočević, *Ordered cone metric spaces and fixed point results*, Comput. Math. Appl. 60 (2010), 1145–1151.
- [30] A. Amini-Harandi, M. Fakhar, *Fixed point theory in cone metric spaces obtained via the scalarization method*, Comput. Math. Appl. 59 (11) (2010), 3529–3534.
- [31] I. Arandjelović, Z. Kadelburg, S. Radenović, *Boyd-Wong-type common fixed point results in cone metric spaces*, Appl. Math. Comput. 217 (2011), 7167–7171.
- [32] I. Arandjelović, D. Kečkić, *A counterexample on a theorem of Khojasteh, Goadarzi and Razani*, Fixed Point Theory Appl., Vol. 2010, Article ID 470141, 6 pages.
- [33] I. Arandjelović, D. Kečkić, *TVS-cone metric spaces as a special case of metric spaces*, arXiv:1202.5930v1 [math. FA] 27 Feb 2012.
- [34] I. Arandjelović, D. Kečkić, *On nonlinear quasicontractions on TVS-cone metric spaces*, Appl. Math. Lett. 24 (2011), 1209–1213.
- [35] M. Arshad, *Fixed points in topological vector space (tvs) valued cone metric spaces*, Adv. Fixed Point Theory 5 (2) (2015), 135–146.
- [36] M. Arshad, J. Ahmad, *On multivalued contractions in cone metric spaces without normality*, The Scientific World J., Vol. 2013, Art. ID 481601, 3 pages.

- [37] M. Arshad, A. Azam, P. Vetro, *Common fixed point of generalized contractive type mappings in cone metric spaces*, IAENG Int. J. Appl. Math. 41 (3) (2012).
- [38] M. Asadi, *Possibility or impossibility of the metrization of cone metric spaces via renorming the Banach spaces*, Nonlinear Funct. Anal. Appl. 19 (3) (2014), 409–413.
- [39] M. Asadi, B. E. Rhoades, H. Soleimani, *Some notes on the paper “The equivalence of cone metric spaces and metric spaces”*, Fixed Point Theory Appl. 2012, 2012:87.
- [40] M. Asadi, H. Soleimani, M. Vaezpour, *An order on subsets of cone metric spaces and fixed points of set-valued contractions*, Fixed Point Theory Appl., Vol. 2009, Article ID 723203.
- [41] M. Asadi, H. Soleimani, S. M. Vaezpour, B. E. Rhoades, *On T-stability of Picard iteration in cone metric spaces*, Fixed Point Theory Appl., Vol. 2009, Article ID 751090, 6 pages.
- [42] M. Asadi, S. M. Vaezpour, V. Rakočević, B. E. Rhoades, *Fixed point theorems for contractive mappings in cone metric spaces*, Math. Commun. 16 (1) (2011), 147–155.
- [43] M. Asadi, M. Vaezpour, B. E. Rhoades, H. Soleimani, *Metrizability of cone metric spaces via renorming the Banach spaces*, Nonlinear Anal. Appl., Vol. 2012, Article ID jnaa-00160, 5 pages.
- [44] H. Aydi, *A common fixed point for a family of compatible mappings involving a quasi-contraction on cone metric spaces*, Thai J. Math. 11 (2013), 157–171.
- [45] H. Aydi, E. Karapinar, S. Moradi, *Coincidence points for expansive mappings under c-distance in cone metric spaces*, Intern. J. Math. Math. Sci., Vol. 2012, Article ID 308921, 11 pages.
- [46] H. Aydi, E. Karapinar, Z. Mustafa, *Coupled coincidence point results on generalized distance in ordered cone metric spaces*, Positivity, Vol. 2013, Art. ID 17, 979–993.
- [47] H. Aydi, B. Nashine, B. Samet, H. Yazidi, *Coincidence and common fixed point results in partially ordered cone metric spaces and applications to integral equations*, Nonlinear Anal. TMA 74 (2011), 6814–6825.
- [48] H. Aydi, B. Samet, C. Vetro, *Coupled fixed point results in cone metric spaces for w-compatible mappings*, Fixed Point Theory Appl. 2011:27 (2011).
- [49] A. Azam, M. Arshad, *Kannan fixed point theorem on generalized metric spaces*, J. Nonlinear Sci. Appl., 1 (1) (2008), 45–48.
- [50] A. Azam, M. Arshad, I. Beg, *Banach contraction principle on cone rectangular metric spaces*, Appl. Anal. Discrete Math. 3 (2009), 236–241.
- [51] A. Azam, M. Arshad, I. Beg, *Existence of fixed points in complete cone metric spaces*, Int. J. Mod. Math. 5 (1) (2010), 91–99.
- [52] A. Azam, I. Beg, *Kannan type mapping in TVS-valued cone metric spaces and their application to Urysohn integral equations*, Sarajevo J. Math. 9 (22) (2013), 243–255.
- [53] A. Azam, I. Beg, M. Arshad, *Fixed point in topological vector space valued cone metric spaces*, Fixed Point Theory Appl., Vol. 2010, Article ID 604084, 9 pages.
- [54] A. Azam, B. Fisher, M. Khan, *Common fixed point theorems in complex valued metric spaces*, Num. Funct. Anal. Optim. 32 (3) (2011), 243–253.
- [55] A. Azam, N. Mehmood, *Multivalued fixed point theorems in tvs-cone metric spaces*, Fixed Point Theory Appl. 2013:184 (2013).
- [56] A. Azam, N. Mehmood, *Fixed point theorems for multivalued mappings in G-cone metric spaces*, J. Inequal. Appl. 2013:354 (2013).
- [57] A. Azam, N. Mehmood, J. Arshad, S. Radenović, *Multivalued fixed point theorems in cone b-metric spaces*, J. Inequal. Appl. 2013:582 (2013).
- [58] A. Azam, B. E. Rhoades, *Some fixed point theorems for a pair of mappings on a cone metric space*, IAENG Int. J. Appl. Math. 42:3 (2012).
- [59] G. V. R. Babu, P. D. Sailaja, *Common fixed point theorems of Ćirić type weak contractions in cone metric spaces*, Thai J. Math. 10 (2012), 517–533.
- [60] J-S. Bae, S-H. Cho, *Fixed points and variational principle with application to equilibrium problems on cone metric spaces*, J. Korean Math. Soc. 50 (2013), 95–109.
- [61] I. A. Bakhtin, *The contraction principle in quasimetric spaces*, Func. An., Ulianowsk, Gos. Ped. Inst. 30 (1989), 26–37.
- [62] S. Banach, *Sur les opérations dans les ensembles abstraits et leur applications aux équations intégrales*, Fund. Math. 3 (1922), 133–181.
- [63] B. Bao, Sh. Xu, L. Shi, V. Ćojbašić Rajić, *Fixed point theorems on generalized c-distance in ordered cone b-metric spaces*, Int. J. Nonlinear Anal. Appl. 6 (1) (2015), 9–22.

- [64] R. Batra, D. Vashistha, *Coupled coincidence point theorems for nonlinear contractions under c -distance in cone metric spaces*, Ann. Funct. Anal. 4 (2013), 138–148.
- [65] R. Batra, D. Vashistha, *Coupled coincidence point theorems for nonlinear contractions under (F, g) -invariant set in cone metric spaces*, J. Nonlinear Sci. Appl. 6 (2013), 86–96.
- [66] R. Batra, D. Vashistha, *Some coupled coincidence point results c -distance in cone metric spaces*, Engin. Math. Letters 2 (2013), 90–114.
- [67] R. Batra, S. Vashistha, R. Kumar, *Coupled coincidence point theorems for mappings without mixed monotone property under c -distance in cone metric spaces*, J. Nonlinear Sci. Appl. 7 (2014), 345–358.
- [68] I. Beg, M. Abbas, T. Nazir, *Generalized cone metric spaces*, J. Nonlinear Sci. Appl., 3 (1) (2010), 21–31.
- [69] I. Beg, M. Abbas, T. Nazir, *Common fixed point results in G -cone metric spaces*, J. Adv. Res. Pure Math. 2 (2010), 94–109.
- [70] I. Beg, A. Azam, M. Arshad, *Common fixed points for maps on topological vector space valued cone metric spaces*, Int. J. Math. Sci., Vol. 2009, Article ID 560264, 8 pages.
- [71] T. Bekeshie, G. A. Naidu, *Recent fixed point theorems for T -contractive mappings and T -weak (almost) contractions in metric and cone metric spaces are not real generalizations*, J. Nonlinear Anal. Optim. 4 (2013), 219–225.
- [72] A. Boccuto, X. Dimitriou, *Ascoli type theorems in the cone metric space setting*, J. Inequal. Appl. 2014:420 (2014).
- [73] A. Branciari, *A fixed point theorem of Banach-Caccioppoli type on a class of generalized metric spaces*, Publ. Math. Debrecen, 57 (2000), 31–37.
- [74] H. Cakalli, A. Sonmez, C. Genc, *On an equivalence of topological vector space valued cone metric spaces and metric spaces*, Appl. Math. Lett. 25 (2012), 429–433.
- [75] N. Cakić, Z. Kadelburg, S. Radenović, A. Razani, *Common fixed point results in cone metric spaces for a family of weakly compatible maps*, Adv. Appl. Math. Sci. 1 (1) (2009), 183–207.
- [76] W. Chaker, A. Ghribi, A. Jeribi, B. Krichen, *Fixed point theorem for quasi-contractive mapping in partial cone metric space*, Afrika Mat. 26 (5-6) (2015), 1153–1159.
- [77] P. Che, L-P. Yang, *Common coupled fixed point theorems for φ -maps in cone metric spaces on w -compatible mappings*, J. Adv. Math. Stud. 7 (2) (2014), 90–103.
- [78] Ch. M. Chen, *On set-valued contractions of Nadler type in tvS - G -cone metric spaces*, Fixed Point Theory Appl. 2012:52 (2012).
- [79] Ch. M. Chen, T. H. Chang, *Common fixed point theorems for a weaker Meir-Keeler type function in cone metric spaces*, Appl. Math. Lett. 23 (11) (2010), 1336–1341.
- [80] Ch.M. Chen, T.H. Chang, K.S. Juang *Common fixed point theorems for the stronger Meir-Keeler type function in cone ball-metric spaces*, Appl. Math. Lett. 25 (2012), 692–697.
- [81] K. Ph. Chi, T.V. An, *Dugundji's theorem for cone metric spaces*, Appl. Math. Lett. 24 (3) (2011), 387–390.
- [82] S. H. Cho, *Fixed point theorems for set-valued generalized contractive maps in cone metric spaces*, Int. J. Math. Anal. 7 (31) (2013), 1543–1555.
- [83] S. H. Cho, J. S. Bae, *Fixed point theorems for multivalued maps in cone metric spaces*, Fixed Point Theory Appl. 2011:87 (2011).
- [84] S. H. Cho, J. S. Bae, *Common fixed point theorems for mappings satisfying property $(E.A)$ on cone metric spaces*, Math. Comput. Modelling 53 (2011), 945–951.
- [85] S. H. Cho, J. S. Bae, K. S. Na *Fixed point theorems for multivalued contractive mappings and multivalued Caristi type mappings in cone metric spaces*, Fixed Point Theory Appl. 2012:133 (2012).
- [86] Y. J. Cho, R. Saadati, Sh. Wang, *Common fixed point theorems on generalized distance in ordered cone metric spaces*, Comput. Math. Appl. 61 (2011), 1254–1260.
- [87] B. S. Choudhury, L. Kumar, T. Som, N. Metiya, *The point of coincidence and common fixed point results in cone metric space*, Adv. Fixed Point Theory 3 (2013), 77–92.
- [88] B. S. Choudhury, L. Kumar, T. Som, N. Metiya, *A weak contraction principle in partially ordered cone metric space with three control functions*, Intern. J. Anal. Appl. 6 (1) (2014), 18–27.
- [89] B. S. Choudhry, N. Metiya, *Fixed points of weak contractions in cone metric spaces*, Nonlinear Anal. TMA 72 (2010), 1589–1593.
- [90] B. S. Choudhry, N. Metiya, *The point of coincidence and common fixed point for a pair of mappings in cone metric spaces*, Comput. Math. Appl. 60 (6) (2010), 1686–1695.

- [91] B.S. Choudhury, N. Metiya, *Coincidence point and fixed point theorems in ordered cone metric spaces*, J. Advan. Math. Stud. 5 (2012), 20–31.
- [92] B.S. Choudhury, N. Metiya, *Fixed point and common fixed point results in ordered cone metric spaces*, An. Şt. Univ. Ovidius Constanţa 20, 1 (2012), 55–72.
- [93] K. J. Chung, *Nonlinear contractions in abstract spaces*, Kodai Math. J. 4 (1981), 288–292.
- [94] Lj. Ćirić, H. Lakzian, V. Rakočević, *Fixed point theorems for w -cone distance contraction mappings in tvs -cone metric spaces*, Fixed Point Theory Appl. 2012:3 (2012).
- [95] Lj. Ćirić, B. Samet, C. Vetro, M. Abbas, *Fixed points results for weak contractive mappings in ordered K -metric spaces*, Fixed Point Theory 13 (1) (2012), 59–72.
- [96] A. S. Cvetković, S. Radenović, S. Simić, M. Stanić, *Generalized φ -contraction for four mappings on cone metric spaces*, Indian J. Math. (IJM), Special Volume on Professor Billy Rhoades, 56 (2) (2014), 209–228.
- [97] A. S. Cvetković, M. Stanić, S. Dimitrijević, S. Simić, *Common fixed point theorems for four mappings on cone metric type space*, Fixed Point Theory Appl., Vol. 2011, Article ID 589725, 15 pages.
- [98] M. Cvetković, *Operatorial contractions on solid cone metric spaces*, J. Nonlinear Convex Anal. 17 (7) (2016), 1399–1408.
- [99] M. Cvetković, V. Rakočević, *Quasi-contraction of Perov type*, Appl. Math. Comput. 235 (2014), 712–722.
- [100] S. Czerwik, *Contraction mappings in b -metric spaces*, Acta Math. Inform. Univ. Ostraviensis 1 (1993), 5–11.
- [101] K. Deimling, *Nonlinear Functional Analysis*, Springer-Verlag, 1985.
- [102] G. Deng, H. Huang, M. Cvetković, S. Radenović, *Cone valued measure of noncompactness and related fixed point theorems*, Bull. Intern. Math. Virtual Institute, 8 (2018), 233–243.
- [103] C. Di Bari, R. Saadati, P. Vetro, *Common fixed point in cone metric spaces for CJM pairs*, Math. Comput. Modell. 54 (9-10) (2012), 2348–2354.
- [104] C. Di Bari, P. Vetro, *Common fixed points in cone metric spaces for MK -pairs and L -pairs*, Ars Combinatoria 99 (2011), 429–437.
- [105] H. S. Ding, M. Jovanović, Z. Kadelburg, S. Radenović, *Common fixed point results for generalized quasicontractions in tvs -cone metric spaces*, J. Comput. Anal. Appl. 15 (3) (2013), 463–470.
- [106] H. S. Ding, Z. Kadelburg, E. Karapinar, S. Radenović, *Common fixed points of weak contractions in cone metric spaces*, Abstr. Appl. Anal., Vol. 2012, Article ID 793862, 18 pages.
- [107] H. S. Ding, L. Li, *Coupled fixed point theorems in partially ordered cone metric spaces*, Filomat 25 (2011), 137–149.
- [108] T. Dominguez, J. Lorenzo, I. Gatica, *Some generalizations of Kannan's fixed point theorem in K -metric spaces*, Fixed Point Theory, 13 (1) (2012), 73–83.
- [109] M. Đorđević, D. Đorić, Z. Kadelburg, S. Radenović, D. Spasić, *Fixed point results under C -distance in TVS -cone metric spaces*, Fixed Point Theory Appl. 2011, 2011:29.
- [110] D. Đorić, *Common fixed point theorems for generalized multivalued contractions on cone metric spaces over a non-normal solid cone*, Fixed Point Theory Appl. 2014:159 (2014).
- [111] D. Đorić, Z. Kadelburg, S. Radenović, *Edelstein-Suzuki-type fixed point results in metric and abstract metric spaces*, Nonlinear Anal. TM&A 75 (2012), 1927–1932.
- [112] Z. Drenkowski, S. Migorski, N. S. Papagegiou, *An Introduction to Nonlinear Analysis: Applications*, Kluwer Academic/ Plenum Publisher, New York 2003.
- [113] D. Đukić, Lj. Paunović, S. Radenović, *Convergence of iterates with errors of uniformly quasi-Lipschitzian mappings in cone metric spaces*, Kragujevac. J. Math. 35 (3) (2011), 399–410.
- [114] Wei-Shih Du, *A note on cone metric fixed point theory and its equivalence*, Nonlinear Anal. TMA 72 (2010), 2259–2261.
- [115] Wei-Shih Du, *Nonlinear contractive conditions for coupled cone fixed point theorems*, Fixed Point Theory Appl. Vol. 2010, Article ID 190606, 20 pages.
- [116] Wei-Shih Du, *New cone fixed point theorems for nonlinear multivalued maps with their applications*, Appl. Math. Lett., 24 (2011), 172–178.
- [117] Wei-Shih Du, *A simple proof of Caristi's fixed point theorem without using Zorn's lemma and transfinite induction*, Thai J. Math. 14 (2) (2016), 259–264.
- [118] A. K. Dubey, U. Mishra, *Some fixed point results of single-valued mapping for c -distance in tvs -cone metric spaces*, Filomat 30 (11) (2016), 2925–2934.

- [119] A. K. Dubey, R. Shukla, R. P. Dubey, *Common fixed point theorems for T-Reich contraction mapping in cone metric spaces*, Adv. Fixed Point Theory 3 (2013), 315–326.
- [120] A. K. Dubey, R. Verma, R. P. Dubey, *Cone metric spaces and fixed point theorems of contractive mappings for c-distance*, Inter. J. Math. Appl. 3 (1) (2015), 83–88.
- [121] E. Durmishi, K. Rasimi, A. Ibraimi, *Finding a metric for a family of contractive functions*, Int. J. Math. Sci. Engg. Appls. (IJMSEA) 8 (6) (2014), 181–188.
- [122] J. Eisenfeld, V. Lakshmikantham, *Remarks on nonlinear contraction and comparison principle in abstract cones*, J. Math. Anal. Appl. 61 (1977), 116–121.
- [123] Z. Ercan, *On the end of the cone metric spaces*, Topology Appl. 166 (2014), 10–14.
- [124] Z. M. Fadail, A. G. B. Ahmad, *Coupled fixed point theorems of single-valued mappings for c-distance in cone metric spaces*, J. Appl. Math., Vol. 2012, Article ID 246516, 20 pages.
- [125] Z. M. Fadail, A. G. B. Ahmad, *Common coupled fixed point theorems of single-valued mapping for c-distance in cone metric space*, Abstract Appl. Anal., Vol. 2012, Article ID 901792, 24 pages.
- [126] Z. M. Fadail, A. G. B. Ahmad, *Coupled coincidence point and common coupled fixed point results in cone b-metric spaces*, Fixed Point Theory Appl., 2013, 2013:177.
- [127] Z. M. Fadail, A. G. B. Ahmad, Z. Golubović, *Fixed point theorems of single-valued mapping for c-distance in cone metric space*, Abstract Appl. Anal., Vol. 2012, Article ID 826815, 11 pages.
- [128] Z. M. Fadail, A. G. B. Ahmad, Lj. Paunović, *New fixed point results of single-valued mapping for c-distance in cone metric space*, Abstract Appl. Anal., Vol. 2012, Article ID 639713, 12 pages.
- [129] Z. M. Fadail, A. G. B. Ahmad, S. Radenović, *Common fixed point and fixed point results under c-distance in cone metric spaces*, Appl. Math. Inform. Sci. Lett. 1 (2) (2012), 47–52.
- [130] Z. M. Fadail, G. S. Rad, V. Öztürk, S. Radenović, *Some remarks on coupled, tripled and n-tupled fixed points theorems in ordered abstract metric spaces*, Far East J. Math. Sci. 97 (7) (2015), 809–839.
- [131] K. Fallahi, M. Abbas, Gh. S. Rad, *Generalized c-distance on cone b-metric spaces endowed with a graph and fixed point results*, Appl. Gen. Topology 18 (2) (2017), 391–400.
- [132] K. Fallahi, A. Petrusel, G. S. Rad, *Fixed point results for pointwise Chatterjea type mappings with respect to a c-distance in cone metric spaces endowed with a graph*, U.P.B. Sci. Bull., Series A, 80 (1) (2018), 47–54.
- [133] X. Fan, Y. Liu, M. Song, Y. Ma, *Some common fixed point theorems in cone metric spaces*, Adv. Fixed Point Theory 4 (3) (2014), 365–377.
- [134] A. P. Farajzadeh, *On the scalarization method in cone metric spaces*, Positivity 18 (2014), 703–708.
- [135] A. P. Farajzadeh, A. Amini-Harandi, D. Baleanu, *Fixed point theory for generalized contractions in cone metric spaces*, Commun. Nonlinear Sci. Numer. Simulat. 17 (2012), 708–712.
- [136] Y. Feng, W. Mao, *The equivalence of cone metric spaces and metric spaces*, Fixed Point Theory 11 (2) (2010), 259–264.
- [137] J. Fernandez, N. Malviya, B. Fisher, *The asymptotically reguylarity and sequences in partial cone b-metric spaces with application*, Filomat 30 (10) (2016), 2749–2760.
- [138] R. Fierro, *Fixed point theorems for set-valued mappings on TVS-cone metric spaces*, Fixed Point Theory Appl. (2015) 2015:221.
- [139] M. Filipović, Lj. Paunović, S. Radenović, M. Rajović, *Remarks on “Cone metric spaces and fixed point theortems of T-Kannan and T-Chatterjea contractive mappings*, Mathematical and Computer Modelling 54 (2011), 1467–1472.
- [140] Lj. Gajić, D. Ilić, V. Rakočević, *On Ćirić maps with generalized contractive iterate at a point and Fisher’s quasicontractions in cone metric spaces*, Appl. Math. Comput. 216 (2010), 2240–2247.
- [141] Lj. Gajić, V. Rakočević, *Quasi-contraction on a nonnonrml cone metric space*, Funct. Anal. Appl. 46 (1) (2012), 62–65.
- [142] Lj. Gajić, V. Rakočević, *On Ćirić maps with a generalized contractive iterate at a point and Fisher’s quasi-contractions in cone metric spaces*, Appl. Math. Comput. 216 (8) (2010), 2240–2247.
- [143] Xun Ge, *A fixed point theorem for correspondences in cone metric spaces*, Fixed Point Theory 15 (2014), 79–86.

- [144] R. George, *Prešić type hybrid contraction and fixed points in cone metric spaces*, Adv. Fixed Point Theory 3 (2013), 93–104.
- [145] R. George, B. Fisher, *Some generalized results of fixed points in cone b-metric spaces*, Math. Moravica 17 (2) (2013), 39–50.
- [146] R. George, H. A. Nabwey, K. P. Reshma, R. Rajagoplan, *Generalized cone b-metric spaces and contraction principles*, Mat. Vesnik 67 (4) (2015), 246–257.
- [147] R. George, K. P. Reshma, R. Rajagopalan, *A generalized fixed point theorem of Prešić type in cone metric spaces and application to Markov process*, Fixed Point Theory Appl. 2011:85 (2011).
- [148] Y. Guo, *A generalization of Banach's contraction principle for some non-obviously contractive operators in a cone metric space*, Turk. J. Math. 35 (2011), 1–8.
- [149] R. H. Haghi, V. Rakočević, S. Rezapour, N. Shahzad, *Best proximity results in regular cone metric spaces*, Rend. Circolo Mat. Palermo 60 (3) (2011), 323–327.
- [150] R. H. Haghi, Sh. Rezapour, *Fixed points of multifunctions on regular cone metric spaces*, Exposit. Math. 28 (2010), 71–77.
- [151] Y. Han, R. Wang, *Coincidence point and common fixed point theorems in cone metric type spaces*, Adv. Fixed Point Theory 5 (3) (2015), 342–358.
- [152] Y. Han, Sh. Xu, *Some new theorems of expanding mappings without continuity in cone metric spaces*, Fixed Point Theory Appl. 2013:3 (2013).
- [153] L. G. Huang, X. Zhang, *Cone metric spaces and fixed point theorems of contractive mappings*, J. Math. Anal. Appl. 332 (2) (2007), 1468–1476.
- [154] H. Huang, *Conditions of non-normality in cone metric spaces*, Mathematica Applicata 25 (4) (2012), 894–898.
- [155] H. Huang, Z. Kadelburg, S. Radenović, *A note on some recent results about multivalued mappings in TVS-cone metric spaces*, J. Adv. Math. Stud. 9 (2) (2016), 330–337.
- [156] H. Huang, S. Radenović, D. Đorić, *A note on the equivalence of some metric and H-cone metric fixed point theorems for multivalued contractions*, Fixed Point Theory Appl., 2014, 2014:43.
- [157] H. Huang, X. Shaoyuan, *Some new topological properties in cone metric spaces*, Research Gate.
- [158] H. Huang, S. Xu, *Fixed point theorems of contractive mappings in cone b-metric spaces and applications*, Fixed Point Theory Appl. 2013, 2013:120.
- [159] X. Huang, J. Luo, Ch. Zhu, X. Wen, *Common fixed point theorem for two pairs of non-self-mappings satisfying generalized Ćirić type contraction condition in cone metric spaces*, Fixed Point Theory Appl. 2014:157 (2014).
- [160] X. Huang, C. Zhu, X. Wen, *Common fixed point theorem for four non-self mappings in cone metric spaces*, Fixed Point Theory Appl., Vol. 2010, Article ID 983802.
- [161] X. Huang, Ch. Zhu, X. Wen, Lj. Lalović *Some common fixed point theorems for a family of non-self mappings in cone metric spaces*, Fixed Point Theory Appl. 2013:144 (2013).
- [162] N. Hussain, M. H. Shah, *KKM mappings in cone b-metric spaces*, Comput. Math. Appl. 62 (2011), 1677–1681.
- [163] D. Ilić, V. Rakočević, *Common fixed points for maps on cone metric space*, J. Math. Anal. Appl. 341 (2) (2008), 876–882.
- [164] D. Ilić, V. Rakočević, *Quasi-contraction on a cone metric space*, Appl. Math. Lett. 22 (2009), 728–731.
- [165] D. Ilić, V. Rakočević, *Kontraktivna preslikavanja na metričkim prostorima i uopštenja*, Univerzitet u Nišu, Prirodno-matematički fakultet, Niš, 2014 [in Serbian].
- [166] Z. Izadi, K. Nourouzi, *Fixed points of correspondences defined on cone metric spaces*, Comput. Math. Appl. 60 (3) (2010), 653–659.
- [167] J. Jachymski, J. Klima, *Cantor's intersection theorem for K-metric spaces with a solid cone and a contraction principle*, J. Fixed Point Theory Appl. 18 (2016), 445–463.
- [168] J. Jachymski, J. Klima, *Around Perov's fixed point theorem for mappings on generalized metric spaces*, Fixed Point Theory 17 (2) (2016), 367–380.
- [169] S. Janković, Z. Golubović, S. Radenović, *Compatible and weakly compatible mappings in cone metric spaces*, Math. Comput. Modeling 52 (2010), 1728–1738.
- [170] S. Janković, Z. Kadelburg, S. Radenović, *On cone metric spaces: a survey*, Nonlinear Anal. 74 (2011), 2591–2601.

- [171] S. Janković, Z. Kadelburg, S. Radenović, B. E. Rhoades, *Assad-Kirk-type fixed point theorems for a pair of nonself mappings on cone metric spaces*, Fixed Point Theory Appl. Vol. 2009, Article ID 761086, 16 pages.
- [172] S. Jiang, Zh. Li, *Extension of Banach contraction principle to partial cone metric spaces over a non-normal solid cone*, Fixed Point Theory Appl. 2013, 2013:250.
- [173] Sh. Jiang, Zh. Li, *Nonlinear quasi-contractions in non-normal cone metric spaces*, Fixed Point Theory Appl. 2014:165 (2014).
- [174] M. Jleli, B. Samet, *The Kannan fixed point theorem in a cone rectangular metric space*, J. Nonlinear Sci. Appl. 2 (3) (2009), 161–167.
- [175] M. Jleli, M. Mursaleen, K. Sadarangani, B. Samet, *A cone measure of noncompactness and some generalizations of Darbo's theorem with applications to functional integral equations*, J. Funct. Spaces, Vol. 2016, Article ID 9896502, 11 pages.
- [176] M. Jovanović, *Contribution to the theory of abstract metric spaces*, PhD thesis, University of Belgrade, Faculty of Mathematics, Belgrade, 2016.
- [177] G. Jungck, *Compatible mappings and common fixed points*, Internat. J. Math. Math. Sci. 9 (1986), 771–779.
- [178] G. Jungck, S. Radenović, S. Radojević, V. Rakočević, *Common fixed point theorems for weakly compatible pairs on cone metric spaces*, Fixed Point Theory Appl., Vol. 2009, Article ID 643840, 13 pages.
- [179] Z. Kadelburg, P. P. Murthy, S. Radenović, *Common fixed points for expansive mappings in cone metric spaces*, Int. J. Math. Analysis, 5 (27) (2011), 1309–1319.
- [180] Z. Kadelburg, Lj. Paunović, S. Radenović, G. Soleimani Rad, *Non-normal cone metric and cone b-metric spaces and fixed point results*, Scientific Publications of the State University of Novi Pazar, Ser. A: Appl. Math. Inform. and Mech. vol. 8 (2) (2016), 177–186.
- [181] Z. Kadelburg, M. Pavlović, S. Radenović, *Common fixed point theorems for ordered contractions and quasicontractions in ordered cone metric spaces*, Comput. Math. Appl. 59 (2010), 3148–3159.
- [182] Z. Kadelburg, S. Radenović, *Subspaces and Quotients of Topological and Ordered Vector Spaces*, University of Novi Sad, Institute of Mathematics, Novi Sad 1997.
- [183] Z. Kadelburg, S. Radenović, *Some common fixed point results in non-normal cone metric spaces*, J. Nonlinear Sci. Appl. 3 (3) (2010), 193–202.
- [184] Z. Kadelburg, S. Radenović, *Meir-Keeler-type conditions in abstract metric spaces*, Appl. Math. Lett. 24 (2011), 1411–1414.
- [185] Z. Kadelburg, S. Radenović, *Some results on set-valued contractions in abstract metric spaces*, Comput. Math. Appl. 62 (2011), 342–350.
- [186] Z. Kadelburg, S. Radenović, *Generalized quasicontractions in orbitally complete abstract metric spaces*, Fixed Point Theory 13 (2) (2012), 527–536.
- [187] Z. Kadelburg, S. Radenović, *Coupled fixed point results under tvs-cone metric spaces and w-cone-distance*, Advances Fixed Point Theory Appl. 2 (1) (2012), 29–46.
- [188] Z. Kadelburg, S. Radenović, *A note on various types of cones and fixed point results in cone metric spaces*, Asian J. Math. Appl. Vol. 2013, Article ID ama0104, 7 pages.
- [189] Z. Kadelburg, S. Radenović, V. Rakočević, *Remarks on "Quasi-contraction on a cone metric space"*, Appl. Math. Lett. 22 (2009), 1674–1679.
- [190] Z. Kadelburg, S. Radenović, V. Rakočević, *A note on equivalence of some metric and cone metric fixed point results*, Appl. Math. Lett. 24 (2011), 370–374.
- [191] Z. Kadelburg, S. Radenović, V. Rakočević, *Topological vector spaces-valued cone metric spaces and fixed point theorems*, Fixed Point Theory Appl., Vol. 2010, Article ID 170253, 17 pages.
- [192] Z. Kadelburg, S. Radenović, B. Rosić, *Strict contractive conditions and common fixed point theorems in cone metric spaces*, Fixed Point Theory Appl., Vol. 2009, Article ID 173838, 14 pages.
- [193] Z. Kadelburg, S. Radenović, S. Simić, *Abstract metric spaces and Caristi-Nguyen-type theorems*, Filomat 25 (3) (2011), 111–124.
- [194] E. Karapinar, *Fixed point theorems in cone Banach spaces*, Fixed Point Theory Appl., Vol. 2009, Article ID 09281, 9 pages.
- [195] E. Karapinar, *Couple fixed point theorems for nonlinear contractions in cone metric spaces*, Comput. Math. Appl. 59 (2010), 3656–3668.

- [196] E. Karapınar, *Some nonunique fixed point theorems of Ćirić type on cone metric spaces*, Fixed Point Theory Appl., Vol. 2010, Article ID 123094, 14 pages.
- [197] E. Karapınar, Wei-Shih Du, *A note on b-cone metric and its related results: Generalizations or equivalence?*, Fixed Point Theory Appl. 2013, 2013:210.
- [198] E. Karapınar, P. Kumam, W. Sintunavarat, *Coupled fixed point theorems in cone metric spaces with a c-distance and applications*, Fixed Point Theory Appl. 2012, 2012:194.
- [199] L. V. Kantorovich, *The majorant principle and Newton's method*, Dokl. Akad. Nauk sssr (n.s.), 76 (1951), 17–20.
- [200] M. A. Khamsi, *Remarks on cone metric spaces and fixed point theorems of contractive mappings*, Fixed Point Theory Appl., Vol. 2010, Article ID 315398, 7 pages.
- [201] M. A. Khamsi, *Generalized metric spaces: A survey*, J. Fixed Point Theory Appl. 17 (3) (2015), 455–475.
- [202] M. A. Khamsi, P. J. Wojciechowski, *On the additivity of the Minkowski functionals*, Nonlinear Funct. Anal. Optim. 34 (6) (2013), 635–647.
- [203] M. S. Khan, M. Samanipour, *Fixed point theorems for some discontinuous operators in cone metric space*, Math. Morav. 12 (2) (2008), 29–34.
- [204] M. Khani, M. Pourmahdian, *On the metrizability of cone metric spaces*, Topology Appl. 158 (2) (2011), 190–193.
- [205] F. Khojasteh, Z. Goodarzi, A. Razani, *Some fixed point theorems of integral type contraction in cone metric spaces*, Fixed Point Theory Appl. Volume 2010, Article ID 189684, 13 pages
- [206] F. Khojasteh, A. Razani, S. Moradi, *A fixed point of generalized T_F -contraction mappings in cone metric spaces*, Fixed Point Theory Appl. 2011:14 (2011).
- [207] D. Klim, D. Wardowski, *Fixed point theorems for set-valued contractions in cone metric spaces*, J. Math. Anal. Appl. 334 (1) (2007), 132–139.
- [208] D. Klim, D. Wardowski, *Dynamic processes and fixed points of set-valued nonlinear contractions in cone metric spaces*, Nonlinear Anal. TMA 71 (2009), 5170–5175.
- [209] W-M. Kong, L-P. Yang, *Quadruple common fixed point theorems for w-compatible mappings in ordered cone metric spaces*, J. Adv. Math. Stud. 7 (2) (2014), 49–65.
- [210] M. A. Krasnosel'ski, P. P. Zabreiko, *Geometrical Methods in Nonlinear Analysis*, Springer, 1984.
- [211] M. Krein, *Propriétés fondamentales des ensembles coniques normaux dans l'espace de Banach*, C.R. (Doklady) Acad. Sci. URSS (N.S) 28 (1940), 13–17.
- [212] M. G. Krein, M. A. Rutman, *Linear operators leaving invariant a cone in a Banach spaces*, Uspekhi Math. Nauk (N.S), 3 (1) (1948), 3–95.
- [213] P. Kumam, N. V. Dung, V. T. L. Hang, *Some equivalences between cone b-metric spaces and b-metric spaces*, Abstr. Appl. Anal. Vol. 2013, Article ID 573740, 8 pages.
- [214] P. Kumam, H. Rahimi, G. Soleimani Rad, *The existence of fixed and periodic point theorems in cone metric type spaces*, J. Nonlinear Sci. Appl. 7 (2014), 255–263.
- [215] A. Kumar, S. Rathee, *Fixed point and common fixed point results in cone metric space and application to invariant approximation*, Fixed Point Theory App. 2015:45 (2015).
- [216] M. Kumar, P. Kumar, S. Kumar, *Some common fixed point theorems using (CLR_g) property in cone metric spaces*, Adv. Fixed Point Theory 2 (2012), 340–356.
- [217] H. Kunze, D. La Torre, F. Mendivil, E. R. Vrscay, *Generalized fractal transforms and self-similar objects in cone metric spaces*, Comut. Math. Appl. 64 (6) (2012), 1761-1769.
- [218] D. R. Kurepa, *Tableaux ramifiés d'ensembles*, C. R.Acad. Sci. Paris 198 (1934), 1563–1565.
- [219] M. A. Kutbi, J. Ahmad, A. Azam, *On fixed point of α - ψ -contractive multi-valued mappings in cone metric spaces*, Abstract Appl. Anal., Vol. 2013, Art. ID 313782.
- [220] H. Lakzian, F. Arabyani, *Some fixed point theorems in cone metric spaces with w-distance*, Int. J. Math. Anal. 3 (2009), 1081–1086.
- [221] A. Latif, N. Hussain, J. Ahmad, *Coincidence points for hybrid contractions in cone metric spaces*, J. Nonlinear Convex Anal. 17 (5) (2016), 899–906.
- [222] A. Latif, F. Y. Shaddad, *Fixed point results for multivalued maps in cone metric spaces*, Fixed Point Theory Appl., Vol. 2010, Article ID 941371, 11 pages.
- [223] K. Li, S. Lin, Y. Ge, *On statistical convergence in cone metric spaces*, Topology Appl. 196 (2015) 641–651.
- [224] Zh. Li, Sh. Jiang, *On fixed point theory of monotone mappings with respect to a partial order introduced by a vector functional in cone metric spaces*, Abstract Appl. Anal., Vol. 2013, Article ID 349305, 6 pages.

- [225] Zh. Li, Sh. Jiang, *Common fixed point theorems of contractions in partial cone metric spaces over non-normal cones*, Abstract Appl. Anal., Vol. 2014, Article ID 653841, pages.
- [226] Zh. Li, Sh. Jiang, *Quasi-contractions restricted with linear bounded mappings in cone metric spaces*, Fixed Point Theory Appl. 2014:87 (2014).
- [227] I-J. Lin, Ch-M. Chen, Ch-H. Chen, T-Y. Cheng, *A note on tvs-G-cone metric fixed point theory*, J. Appl. Math., Vol. 2012, Article ID 407071, 10 pages.
- [228] I-J. Lin, Ch-M. Chen, M. Jovanović, Tz-H. Wu, *Fixed points and endpoints of set-valued contractions in cone metric spaces*, Abstract Appl. Math., Vol. 2012, Article ID 632638, 14 pages.
- [229] S. Lin, Y. Ge, *Compact-valued continuous relations on TVS-cone metric spaces*, Filomat 27 (2) (2013), 327–332.
- [230] S. Lin, K. Li, Y. Ge, *On the metrizability of TVS-cone metric spaces*, Publ. Inst. Math., Nouvelle série, 98 (112) (2015), 271–279.
- [231] W. Long, B. E. Rhoades, M. Rajović, *Coupled coincidence points for two mappings in metric spaces and cone metric spaces*, Fixed Point Theory Appl. 2012:66 (2012).
- [232] N. V. Luong, N. X. Thuan, K. P. R. Rao, *Remarks on coupled fixed point theorems in cone metric spaces*, Mat. Vesnik 65 (1) (2013), 122–136.
- [233] Y. Ma, Y. Liu, M. Song, *Common fixed point theorems for a family of mappings in cone metric spaces*, Adv. Fixed Point Theory 4 (2014), 160–168.
- [234] R. Maitra, *Caccioppoli type fixed point result in cone-b-metric space*, IJISSET 1 (4) (2014), 3 pages.
- [235] S. K. Malhotra, J.B. Sharma, S. Shukla, *g-weak contraction in ordered cone rectangular metric spaces*, The Scientific World J., Vol. 2013, Art. ID 810732.
- [236] S. K. Malhotra, S. Shukla, R. Sen, *A generalization of Banach contraction principle in ordered cone metric spaces*, J. Advan. Math. Stud. 5 (2012), 59–67.
- [237] S. K. Malhotra, S. Shukla, R. Sen, *Some coincidence and common fixed points theorems on cone metric spaces*, Bul. Math. Anal. Appl. 4 (2012), 64–71.
- [238] S. K. Malhotra, S. Shukla, R. Sen, *Some fixed point theorems for ordered Reich type contractions in cone rectangular metric spaces*, Acta Math. Univ. Comenianae 82 (2) (2013), 165–175.
- [239] S. K. Malhotra, S. Shukla, R. Sen, *Some fixed point results in θ -complete partial cone metric spaces*, J. Adv. Math. Stud. 6 (2) (2013), 97–108.
- [240] S. K. Malhotra, S. Shukla, R. Sen, N. Verma, *Generalization of fixed point theorems in partial cone metric spaces*, Int. J. Math. Archive 2 (4) (2011), 610–616.
- [241] S. K. Malhotra, S. Shukla, J. B. Sharma, *Cyclic contractions in θ -complete partial cone metric spaces and fixed point theorems*, Jordan J. Math. Stat. 7 (3) (2014), 233–246.
- [242] N. Mehmood, Personal communication.
- [243] N. Mehmood, A. Azam, S. Aleksić, *Topological vector-space valued cone Banach spaces*, Intern. J. Anal. Appl. 6 (2) (2014), 205–219.
- [244] N. Mehmood, A. Azam, I. Beg, *Fixed points of Edelstein-type multivalued maps*, Rend. Circolo Mat. Palermo 63 (3) (2014), 399–407.
- [245] N. Mehmood, A. Azam, Lj. Kočinac, *Multivalued fixed point results in cone metric spaces*, Topology Appl. 179 (2015) 156–170.
- [246] N. Mehmood, A. Azam, Lj. Kočinac, *Multivalued $\mathcal{R}_{\psi, \phi}$ -weakly contractive mappings in ordered cone metric spaces with applications*, Fixed Point Theory 18 (2) (2017), 673–688.
- [247] S. H. Mohanta, *Common fixed points for mappings in G-cone metric spaces*, J. Nonlinear Anal. Appl., Vol. 2012, Article ID jnaa-00120, 13 pages.
- [248] S. K. Mohanta, R. Maitra, *A characterization of completeness in cone metric spaces*, J. Nonlinear Sci. Appl. 6 (2013), 227–233.
- [249] H. Mohebi, *Topical functions and their properties in a class of ordered Banach spaces*, Part II, in: Continuous Optimization, Current Trends and Modern Applications, Springer, 2005, pp. 343–361.
- [250] S. Moradi, *A simple proof of Olatinwo's theorem*, 58 (1) (2012), 119–125.
- [251] F. Moradlou, P. Salimi, *Edelstein-Suzuki fixed point theorem in metric and cone metric spaces; an implicit relation approach*, Romai J. 9 (1) (2013), 70–89.
- [252] F. Moradlou, P. Salimi, S. Radenović, *Implicit relation and Edelstein-Suzuki type fixed point results in cone metric spaces*, Appl. Math. E-Notes, 14 (2014), 1–12.

- [253] F. Moradlou, P. Salimi, P. Vetro, *Some new extensions of Edelstein-Suzuki-type fixed point theorems to G -metric and G -cone metric spaces*, Acta Math. Sci. 33B (4) (2013), 1049–1058.
- [254] J. Morales, E. Rojas, *Cone metric spaces and fixed point theorems of T -Kannan contractive mappings*, Int. J. Math. Anal. 4 (4) (2010) 175–184.
- [255] P. P. Murthy, K. T. Rashmi, *Tripled common fixed point theorems for w -compatible mappings in ordered cone metric spaces*, Adv. Fixed Point Theory 2 (2012), 157–175.
- [256] P. P. Murthy, K. T. Rashmi, L. N. Mishra, *Sub-compatible maps, weakly commuting maps and common fixed points in cone metric spaces*, Scientific Publications of the State University of Novi Pazar, Ser. A: Appl. Math. Inform. and Mech.vol. 8, 2 (2016), 139–148.
- [257] H. K. Nashine, M. Abbas, *Common fixed point of mappings satisfying implicit contractive conditions in TVS-valued ordered cone metric spaces*, J. Nonlinear Sci. Appl. 6 (2013), 205–215.
- [258] H. K. Nashine, Z. Kadelburg, R. P. Pathak, S. Radenović, *Coincidence and fixed point results in ordered G -cone metric spaces*, Math. Comput. Model. 57 (2013), 701–709.
- [259] H. K. Nashine, Z. Kadelburg, S. Radenović, *Coupled common fixed point theorems for w^* -compatible mappings in ordered cone metric spaces*, Appl. Math. Comput. 218 (2012), 5422–5432.
- [260] J. O. Olaleru, *Some generalizations of fixed point theorems in cone metric spaces*, Fixed Point Theory Appl., Vol. 2009, Article ID 657914, 10 pages.
- [261] J. O. Olaleru, G. A. Okeke, H. Akewe, *Coupled fixed point theorems of integral type mappings in cone metric spaces*, Kragujevac J. Math. 36 (2012), 215–224.
- [262] H. Olaoluwa, J. Olaleru, *Multipled fixed point theorems in cone metric spaces*, Fixed Point Theory Appl. 2014:43 (2014).
- [263] I. M. Olaru, A. Branga, A. Oprea, *Common fixed point results in b -cone metric spaces over topological vector spaces*, General Math. 20 (2012), 57–67.
- [264] I. M. Olaru, N. A. Secelean, *Vector comparison operators in cone metric spaces*, Math. Reports 16(66) (3) (2014), 431–442.
- [265] M. O. Olatinwo, *Coupled fixed point theorems in cone metric spaces*, Ann. Univ. Ferrara 57 (2011), 173–180.
- [266] M. Öztürk, M. Basarir, *On some common fixed point theorems with φ -maps on G -cone metric spaces*, Bull. Math. Anal. Appl. 3 (1) (2011), 121–133.
- [267] M. Öztürk, M. Basarir, *On some coincidence and common fixed point theorems in G -cone metric spaces*, Thai J. Math. 9 (3) (2011), 647–657.
- [268] E. de Pascale, G. Marino, P. Pietramala, *The use of E -metric spaces in the search for fixed points*, Le Matematiche 48 (1993), 367–376.
- [269] H. K. Pathak, N. Shahzad, *Fixed point results for generalized quasi-contraction mappings in abstract metric spaces*, Nonlinear Anal. TMA 71 (2009) 6068–6076.
- [270] Lj. Paunović, *Teorija apstraktnih metričkih prostora*, Leposavić 2017 [in Serbian].
- [271] M. Pavlović, S. Radenović, S. Radojević, *Abstract metric spaces and Sehgal-Guseman-type theorems*, Comput. Math. Appl. 60 (2010), 865–872..
- [272] B. Popović, S. Radenović, S. Shukla, *Fixed point results to tvs-cone b -metric spaces*, Gulf J. Math. 1 (2013) 51–64.
- [273] P. D. Proinov, *A unified theory of cone metric spaces and its applications to the fixed point theory*, Fixed Point Theory Appl. 2013:103 (2013).
- [274] P. D. Proinov, I. A. Nikolova, *Iterative approximation of fixed points of quasi-contraction mappings in cone metric spaces*, J. Inequal. Appl. 2014:226 (2014).
- [275] K. Prudhvi, *Common fixed points of four mappings in cone metric spaces*, Adv. Inequal. Appl. 2014:27 (2014).
- [276] S. Radenović, *A pair of non-self mappings in cone metric spaces*, Kragujevac J. Math. 36 (2) (2012), 189–198.
- [277] S. Radenović, *Common fixed points under contractive conditions in cone metric spaces*, Comput. Math. Appl. 58 (2009) 123–1278.
- [278] S. Radenović, Z. Kadelburg, *Quasi-contractions on symmetric and cone symmetric spaces*, Banach J. Math. Anal. 5 (1) (2011), 38–50.
- [279] S. Radenović, Z. Kadelburg, *Some results on fixed points of multifunctions on abstract metric spaces*, Math. Comput. Model. 53 (2011) 746–754.

- [280] S. Radenović, M. Pavlović, Lj. Paunović, Z. Kadelburg, *A note on generalized operator quasicontractions in cone metric spaces*, Scientific Publications of the state University of Novi Pazar, Ser. A: Appl. Math. Inform. and Mech. 9 (2) (2017), 127–132.
- [281] S. Radenović, V. Rakočević, Sh Rezapour, *Common fixed points for (g, f) -type maps in cone metric spaces*, Appl. Math. Comput. 218 (2011), 480–491.
- [282] S. Radenović, B. E. Rhoades, *Fixed point theorem for two non-self mappings in cone metric spaces*, Comput. Math. Appl. 57 (2009), 1701–1707.
- [283] S. Radenović, S. Simić, N. Cakić, Z. Golubović, *A note on tvs-cone metric fixed point theory*, Math. Comput. Model. 54 (2011), 2418–2422.
- [284] S. Radenović, F. Vetro, *Some remarks on Perov type mappings in cone metric spaces*, Mediterr. J. Math. 14:240 (2017).
- [285] S. Radenović, F. Vetro, S. Xu, *Some results of Perov type mappings*, J. Adv. Math. Stud. 10 (3) (2017), 396–409.
- [286] S. Radojević, Lj. Paunović, S. Radenović, *Abstract metric spaces and Hardy-Rogers-type theorems*, Appl. Math. Lett. 24 (2011), 553–558.
- [287] H. Rahimi, G. S. Rad, *Note on “Common fixed point results for noncommuting mappings without continuity in cone metric spaces”*, Thai J. Math. 11 (3) (2013), 589–599.
- [288] H. Rahimi, G. S. Rad, *Common fixed point theorems and c -distance in ordered cone metric spaces*, Ukrainian Math. 12 (2014), 1845–1861.
- [289] H. Rahimi, G. S. Rad, *Fixed point theorems under c -distance in ordered cone metric spaces*, Int. J. Industrial Math. 6 (2) (2014), Art. ID IJIM-00253, 9 p.
- [290] H. Rahimi, G. S. Rad, P. Kumam, *Coupled common fixed point theorems under weak contractions in cone metric type spaces*, Thai J. Math. 12 (1) (2014), 1–14.
- [291] H. Rahimi, S. Radenović, G. S. Rad, P. Kumam, *Quadrupled fixed point results in abstract metric spaces*, Comput. Appl. Math. 33 (3) (2014), 671–685.
- [292] H. Rahimi, B. E. Rhoades, S. Radenović, G. S. Rad, *Fixed and periodic point theorems for T -contractions on cone metric spaces*, Filomat 27 (5) (2013), 881–888.
- [293] H. Rahimi, P. Vetro, G. S. Rad, *Some common fixed point results for weakly compatible mappings in cone metric type spaces*, Miskolc Math. Notes 14 (2013), 233–243.
- [294] P. Raja, S. M. Vaezpour, *Some extensions of Banach’s contraction principles in complete cone metric spaces*, Fixed Point Theory Appl., Vol. 2008, Article ID 768294, 11 pages.
- [295] V. Č. Rajić, S. Radenović, *A note on tripled fixed point of w -compatible mappings in tvs-cone metric spaces*, Thai J. Math. 12 (3) (2014), 717–728.
- [296] K. P. R. Rao, G. N. V. Kishore, *A unique common triple fixed point theorem in partially ordered cone metric spaces*, Bull. Math. Anal. Appl. 3 (2011), 213–222.
- [297] M. Rashid, N. Mehmood, A. Azam, S. Radenović, *Fuzzy fixed point theorems in ordered cone metric spaces*, Filomat 29 (4) (2015) 887–896.
- [298] R. A. Rashwan, S. M. Saleh, *Some fixed point theorems in cone rectangular metric spaces*, Math. Aeterna 2 (60) (2012), 573–587.
- [299] A. Razani, V. Rakočević, Z. Goodarzi, *Generalized φ -contraction for a pair of mappings on cone metric spaces*, Appl. Math. Comput. 217 (2) (2011), 8889–8906.
- [300] Sh. Rezapour, *Best approximations in cone metric spaces*, Math. Moravica 11 (2007), 85–88.
- [301] Sh. Rezapour, R. H. Haghi, *Fixed points of multifunctions on cone metric spaces*, Numer. Funct. Anal. Optim. 30 (7-8) (2009), 1–8.
- [302] Sh. Rezapour, R. H. Haghi, N. Shahzad, *Some notes on fixed points of quasi-contraction maps*, Appl. Math. Lett. 23 (2010), 498–502.
- [303] Sh. Rezapour, R. Hamlbarani, *Some notes on the paper “Cone metric spaces and fixed point theorems of contractive mappings”* J. Math. Anal. Appl. 345 (2008), 719–724.
- [304] B.E. Rhoades, *A comparison of various definitions of contractive mappings*, Trans. Amer. Math. Soc., 226 (1977), 257–290.
- [305] B. Rzepecki, *On fixed point theorems of Maia type*, Publ. Inst. Math. 28 (42) (1980), 179–186.
- [306] F. Sabetghadam, H. P. Masiha, *Common fixed points for generalized φ -pairs of mappings on cone metric spaces*, Fixed Point Theory Appl., Vol. 2010, Article ID 718340, 8 pages.
- [307] F. Sabetghadam, H. P. Masiha, A. H. Sanatpour, *Some coupled fixed point theorems in cone metric spaces*, Fixed Point Theory Appl., Vol. 2009, Article ID 125426.
- [308] I. Sahin, M. Telci, *Fixed points of contractive mappings on complete cone metric spaces*, Hacet. J. Math. Stat. 38 (1) (2009) 59–67.

- [309] I. Sahin, M. Telci, *A theorem on common fixed points of expansion type mappings in cone metric spaces*, An. Şt. Univ. Ovidius Constanţa 18 (2010), 329–336.
- [310] G. S. Saluja, *Convergence theorems for generalized asymptotically quasi-nonexpansive mappings in cone metric spaces*, CUBO A Math. J. 15 (3) (2013), 71–87.
- [311] B. Samet, *Common fixed point under contractive condition of Ćirić type in cone metric spaces*, Appl. Anal. Discrete Math. 5 (2011), 159–164.
- [312] K. P. R. Sastry, Ch. S. Rao, A. Ch. Sekhar, M. Balaiah, N. A. Aprarao *A coincidence point theorem for expansive maps under c -distance in cone metric spaces*, Int. J. Math. Archive 4 (1) (2013), 204–211.
- [313] F. Shaddad, M. S. M. Noorani, *Fixed point results in quasi-cone metric space*, Abstract Appl. Anal., Vol. 2013, Article ID 303662.
- [314] H. H. Schaefer, *Topological Vector Spaces*, 3rd ed., Berlin-Heidelberg-New York: Springer-Verlag, 1971.
- [315] M. H. Shah, S. Simić, N. Hussain, A. Sretenović, S. Radenović, *Common fixed points theorems for occasionally weakly compatible pairs on cone metric type spaces*, J Comput. Anal. Appl. 14 (2) (2012), 290–297, 2012.
- [316] W. Shatanawi, *Partially ordered cone metric spaces and coupled fixed point results*, Comput. Math. Appl. 60 (2010), 2508–2515.
- [317] W. Shatanawi, *Some common coupled fixed point results in cone metric spaces*, Int. J. Math. Anal. 4 (48) (2010), 2381–2388.
- [318] W. Shatanawi, *On w -compatible mappings and common coupled coincidence point in cone metric spaces*, Appl. Math. Lett. 25 (2012), 925–931.
- [319] W. Shatanawi, *Some coincidence point results in cone metric spaces*, Math. Comput. Modelling 55 (2012), 2023–2028.
- [320] W. Shatanawi, F. Awawdeh, *Some fixed and coincidence point theorems for expansive maps in cone metric spaces*, Fixed Point Theory Appl. 2012:19 (2012).
- [321] W. Shatanawi, E. Karapinar, H. Aydi, *Coupled coincidence points in partially ordered cone metric spaces with a c -distance*, J. Appl. Math., Vol. 2012, Article ID 312078, 15 pages.
- [322] W. Shatanawi, V. Ć. Rajić, S. Radenović, A. Al-Rawashdeh, *Mizoguchi-Takahashi-type theorems in tvs -cone metric spaces*, Fixed Point Theory Appl. 2012:106 (2012).
- [323] L. Shi, Sh. Xu, *Common fixed point theorems for two weakly compatible self-mappings in cone b -metric spaces*, Fixed Point Theory Appl. 2013:120 (2013).
- [324] R. Shrivastava, Z. K. Ansari, M. Sharma, *On generalization of some common fixed point theorems for T -contraction in cone metric space*, Int. J. Phys. Math. Sci. 2011, 83–87.
- [325] S. Simić, *A note on Stone's, Baire's, Ky Fan's and Dugundji's theorem in tvs -cone metric spaces*, Appl. Math. Lett. 24 (6) (2011), 999–1002.
- [326] A. Singh, R. C. Dimri, S. Bhatt, *A unique common fixed point theorem for four maps in cone metric spaces*, Int. J. Math. Anal. 4 (2010), 1511–1517.
- [327] T. Ch. Singh, Y. R. Singh, *Some remarks on D -quasi contraction on cone symmetric space*, Intern. J. Math. Archive 5 (7) (2014), 75–78.
- [328] W. Sintunavarat, Y. J. Cho, P. Kumam, *Common fixed point theorems for c -distance in ordered cone metric spaces*, Comput. Math. Appl. 62 (2011), 1969–1978.
- [329] W. Sintunavarat, Y.J. Cho, P. Kumam, *Coupled fixed point theorems for weak contraction mappings under F -invariant set*, Abstract Appl. Anal. 2012, Article ID 324874, 15 pages.
- [330] G. Soleimani Rad, K. Fallahi, Z. Kadelburg, *New fixed point results under generalized c -distance in tvs -cone b -metric spaces with an application to systems of Fredholm integral equations*, to appear in J. Math. Extensions.
- [331] G. Song, X. Sun, Y. Zhao, G. Wang, *New common fixed point theorems for maps on cone metric spaces*, Appl. Math. Lett. 23 (9) (2010), 1033–1037.
- [332] A. Sönmez, *On paracompactness in cone metric spaces*, Appl. Mathem. Lett. 23 (2010), 494–497.
- [333] A. Sönmez, H. Cakalli, *Cone normed spaces and weighted means*, Mathem. Comput. Modell. 52 (2010), 1660–1666.
- [334] M. P. Stanić, A. S. Cvetković, S. Simić, S. Dimitrijević, *Common fixed point under contractive condition of Ćirić's type on cone metric type spaces*, Fixed Point Theory Appl. 2012:35 (2012).

- [335] R. Sumitra, V. R. Uthariaraj, R. Hemavathy, P. Vijayaraju, *Common fixed point theorem for non-self mappings satisfying generalized Ćirić type contraction condition in cone metric spaces*, Fixed Point Theory Appl., Vol. 2010, Article ID 408086, 17 pages.
- [336] R. Sumitra, V. R. Uthariaraj, R. Hemavathy, *Common fixed point theorem for T-Hardy-Rogers contraction mapping in a cone metric space*, Int. Math. Forum 5 (30) (2010), 1495–1506.
- [337] M. R. H. Tavani, *Some fixed point and common fixed point theorems in G-cone metric spaces*, Adv. Environ. Biol. 8 (7) (2014), 3387–3392.
- [338] M. R. H. Tavani, *Some fixed point theorems in quasi-cone metric spaces*, Adv. Environ. Biol. 8 (7) (2014), 3382–3386.
- [339] F. Tchier, C. Vetro, F. Vetro, *A coincidence-point problem of Perov type on rectangular cone metric spaces*, J. Nonlinear Sci. Appl. 10 (2017), 4307–4317.
- [340] H. L. Tidke, M. B. Dhakne, *Existence and uniqueness of solution of differential equation of second order in cone metric spaces*, Fasc. Math. 45 (2010), 121–131.
- [341] S. K. Tiwari, R. Pal, R. P. Dubey, *Generalization of fixed point theorems in cone metric spaces*, Intern. J. Math. Archive 5 (5) (2014), 115–122.
- [342] M. A. Tootkaboni, A. B. Salec, *Algebraic cone metric spaces and fixed point theorems of contractive mappings*, Fixed Point Theory Appl. 2014:160 (2014).
- [343] D. Turkoglu, M. Abuloha, *Cone metric spaces and fixed point theorems in diametrically contractive mappings*, Acta Math. Sin. (Engl. Ser.) 26 (3) (2010), 489–496.
- [344] D. Turkoglu, M. Abuloha, T. Abdeljavad, *KKM mappings in cone metric spaces and some fixed point theorems*, Nonlinear Anal. TMA 72 (2010), 348–353.
- [345] J. S. Vandergraft, *Newton method for convex operators in partially ordered spaces*, SIAM J. Numer. Anal. 4 (3) (1967), 406–432.
- [346] F. Vetro, S. Radenović, *Some results of Perov type on rectangular cone metric spaces*, J. Fixed Point Theory Appl., doi.org/10.1007/s11784-018-0520-y
- [347] P. Vetro, *Common fixed points in cone metric spaces*, Rend. Circ. Mat. Palermo, Ser. II, 56 (2007), 464–468.
- [348] N. Wairoijana, W. Sintunavarat, P. Kumam, *Common tripled fixed point theorems for W-compatibility mappings along with the CLR_g property in abstract metric spaces*, J. Inequal. Appl. 2014:133 (2014).
- [349] F. Wang, Sh. M. Kang, Sh. Wang, *Fixed point theorems for generalized multivalued mappings in cone metric spaces*, Int. J. Math. Anal. 7 (6) (2013), 265–274.
- [350] Sh. Wang, B. Guo, *Distance in cone metric spaces and common fixed point theorems*, Appl. Math. Lett. 24 (2011), 1735–1739.
- [351] Sh. Wang, M. S. Park, *Common fixed point theorems in generalized ordered cone metric spaces*, Nonlinear Funct. Anal. Appl. 16 (3) (2011), 387–399.
- [352] D. Wardowski, *Endpoints and fixed points of set-valued contractions in cone metric spaces*, Nonlinear Anal. TMA 71 (2009), 512–516.
- [353] K. Włodarczyk, R. Plebaniak, *Maximality principle and general results of Ekeland and Caristi types without lower semicontinuity assumptions in cone uniform spaces with generalized pseudodistances*, Fixed Point Theory Appl. Vol. 2010, Article ID 175453, 37 pages.
- [354] K. Włodarczyk, R. Plebaniak, *Periodic point, endpoint, and convergence theorems for dissipative set-valued dynamic systems with generalized pseudodistances in cone uniform and uniform spaces*, Fixed Point Theory Appl. Vol. 2010, Article ID 864536, 32 pages
- [355] K. Włodarczyk, R. Plebaniak, C. Obczynski, *Convergence theorems, best approximation and best proximity for set-valued dynamic system of relatively quasi-asymptotic contractions in cone uniform spaces*, Nonlinear Anal. TMA 72 (2010), 794–805.
- [356] K. Włodarczyk, R. Plebaniak, M. Dolinski, *Cone uniform, cone locally convex and cone metric spaces, endpoints, set-valued dynamic systems and quasi-asymptotic contractions*, Nonlinear Anal. TMA 71 (2009), 5022–5031.
- [357] Yau-Chuen Wong, Kung-Fu Ng, *Partially Ordered Topological Vector Spaces*, Clarendon Press, Oxford, 1973.
- [358] S. Xiang, S. Xia, *Notes on the metrizablety of cone metric spaces*, College of Science, Guizhon University, 550025.
- [359] S. Xu, Ć. Dolićanin, S. Radenović, *Some remarks on results of Perov type*, J. Adv. Math. Stud. 9 (3) (2016), 361–369.

- [360] Sh. Xu, H. Huang, *Correction: Fixed point theorems of contractive mappings in cone b-metric spaces and applications*, Fixed Point Theory Appl. 2014:55 (2014).
- [361] Sh. Xu, A. Cheng, Y. Han, *Correction: Some new theorems of expanding mappings without continuity in cone metric spaces*, Fixed Point Theory Appl. 2014:178 (2014).
- [362] W. Xu, Ch. Zhu, Ch. Chen, *Suzuki's fixed point theorems for generalized mappings in partial cone metric spaces over a solid cone*, Fixed Point Theory Appl. 2016:47 (2016), 1–18.
- [363] P. P. Zabrejko, *K-metric and K-normed linear spaces: survey*, Collect. Math. 48 (1997), 825–859.
- [364] X. Zhang, *Fixed point theorem of generalized quasi-contractive mapping in cone metric spaces*, Comput. Math. Appl. 62 (2011), 1627–1633.
- [365] X. Zhang, *Some new versions of Hahn-Banach theorem*, Afrika Mat. 25 (3) (2014), 823–831.
- [366] Ch. Zhu, W. Xu, T. Došenović, Z. Golubović, *Common fixed point theorems for cyclic contractive mappings in partial cone b-metric spaces and applications to integral equations*, Nonlinear Anal. Model. Control 21 (6) (2016), 807–827.