

FURTHER REFINEMENTS OF GENERALIZED NUMERICAL RADIUS INEQUALITIES FOR HILBERT SPACE OPERATORS

MONIRE HAJMOHAMADI¹, RAHMATOLLAH LASHKARIPOUR², MOJTABA BAKHERAD³

ABSTRACT. In this paper, we show some refinements of generalized numerical radius inequalities involving the Young and Heinz inequalities. In particular, we present

$$\begin{aligned} & w_p^p(A_1^*T_1B_1, \dots, A_n^*T_nB_n) \\ & \leq \frac{n^{1-\frac{1}{r}}}{2^{\frac{1}{r}}} \left\| \sum_{i=1}^n [B_i^* f^2(|T_i|) B_i]^{rp} + [A_i^* g^2(|T_i^*|) A_i]^{rp} \right\|^{\frac{1}{r}} - \inf_{\|x\|=1} \eta(x), \end{aligned}$$

where $T_i, A_i, B_i \in \mathbb{B}(\mathcal{H})$ ($1 \leq i \leq n$), f and g are nonnegative continuous functions on $[0, \infty)$ satisfying $f(t)g(t) = t$ for all $t \in [0, \infty)$, $p, r \geq 1$, $N \in \mathbb{N}$ and

$$\begin{aligned} \eta(x) = & \frac{1}{2} \sum_{i=1}^n \sum_{j=1}^N \left(\sqrt[2^j]{\langle (A_i^* g^2(|T_i^*|) A_i)^{p x}, x \rangle^{2^{j-1}-k_j} \langle (B_i^* f^2(|T_i|) B_i)^{p x}, x \rangle^{k_j}} \right. \\ & \left. - \sqrt[2^j]{\langle (B_i^* f^2(|T_i|) B_i)^{p x}, x \rangle^{k_j+1} \langle (A_i^* g^2(|T_i^*|) A_i)^{p x}, x \rangle^{2^{j-1}-k_j-1}} \right)^2. \end{aligned}$$

1. INTRODUCTION

Let $\mathbb{B}(\mathcal{H})$ denote the C^* -algebra of all bounded linear operators on a complex Hilbert space \mathcal{H} with an inner product $\langle \cdot, \cdot \rangle$ and the corresponding norm $\| \cdot \|$. In the case when $\dim \mathcal{H} = n$, we identify $\mathbb{B}(\mathcal{H})$ with the matrix algebra \mathbb{M}_n of all $n \times n$ matrices with entries in the complex field. The numerical radius of $T \in \mathbb{B}(\mathcal{H})$ is defined by

$$w(T) := \sup\{ | \langle Tx, x \rangle | : x \in \mathcal{H}, \|x\| = 1 \}.$$

It is well known that $w(\cdot)$ defines a norm on $\mathbb{B}(\mathcal{H})$, which is equivalent to the usual operator norm $\| \cdot \|$. In fact, for any $T \in \mathbb{B}(\mathcal{H})$, $\frac{1}{2}\|T\| \leq w(T) \leq \|T\|$; see [6]. The quantity $w(T)$ is useful in studying perturbation, convergence and approximation problems as well as interactive method, etc. For more information see [1, 2, 4, 5, 7, 8, 13, 19] and references therein.

The classical Young inequality says that if $0 \leq \nu \leq 1$, then $a^\nu b^{1-\nu} \leq \nu a + (1 -$

2010 *Mathematics Subject Classification.* Primary 47A12, Secondary 47A63, 47A30 .

Key words and phrases. Euclidean operator radius; Heinz means; Numerical radius; positive operator; Young inequality.

$\nu)b$ ($a, b > 0$). During the last decades several generalizations, reverses, refinements and applications of the Young inequality in various settings have been given (see [3, 12] and references therein). A refinement of the scalar Young inequality is presented in [12] as follows:

$$a^\nu b^{1-\nu} \leq \nu a + (1-\nu)b - r_0(a^{\frac{1}{2}} - b^{\frac{1}{2}})^2, \quad (1.1)$$

where $r_0 = \min\{\nu, 1-\nu\}$.

Recently, Sababheh and Choi in [15] obtained a refinement of the Young inequality

$$a^\nu b^{1-\nu} \leq \nu a + (1-\nu)b - S_N(\nu), \quad (1.2)$$

in which

$$S_N(\nu) := \sum_{j=1}^N \left((-1)^{r_j} 2^{j-1} \nu + (-1)^{r_j+1} \left[\frac{r_j+1}{2} \right] \right) \left(\sqrt[2^j]{b^{2^{j-1}-k_j} a^{k_j}} - \sqrt[2^j]{a^{k_j+1} b^{2^{j-1}-k_j-1}} \right)^2,$$

where $N \in \mathbb{N}$, $r_j = [2^j \nu]$ and $k_j = [2^{j-1} \nu]$. Here $[x]$ is the greatest integer less than or equal to x . When $N = 1$, inequality (1.2) reduces to (1.1).

It follows from $\nu a + (1-\nu)b \leq (\nu a^r + (1-\nu)b^r)^{\frac{1}{r}}$ ($r \geq 1$) and inequality (1.1) that

$$a^\nu b^{1-\nu} \leq (\nu a^r + (1-\nu)b^r)^{\frac{1}{r}} - S_N(\nu).$$

In particular, for $\nu = \frac{1}{2}$ we get

$$a^{\frac{1}{2}} b^{\frac{1}{2}} \leq \left(\frac{1}{2} \right)^{\frac{1}{r}} (a^r + b^r)^{\frac{1}{r}} - \frac{1}{2} \sum_{j=1}^N \left(\sqrt[2^j]{b^{2^{j-1}-k_j} a^{k_j}} - \sqrt[2^j]{a^{k_j+1} b^{2^{j-1}-k_j-1}} \right)^2.$$

If $N = 1$, then we reach to inequality (2.1) in [12] as follows:

$$a^{\frac{1}{2}} b^{\frac{1}{2}} \leq \left(\frac{1}{2} \right)^{\frac{1}{r}} (a^r + b^r)^{\frac{1}{r}} - \frac{1}{2} (a^{\frac{1}{2}} - b^{\frac{1}{2}})^2.$$

Let $T_i \in \mathbb{B}(\mathcal{H})$ ($1 \leq i \leq n$). The Euclidean operator radius of T_1, \dots, T_n is defined in [14] by

$$w_e(T_1, \dots, T_n) := \sup_{\|x\|=1} \left(\sum_{i=1}^n |\langle T_i x, x \rangle|^2 \right)^{\frac{1}{2}}.$$

In [16], the functional w_p of operators T_1, \dots, T_n for $p \geq 1$ is defined by

$$w_p(T_1, \dots, T_n) := \sup_{\|x\|=1} \left(\sum_{i=1}^n |\langle T_i x, x \rangle|^p \right)^{\frac{1}{p}}.$$

Let $T_1, \dots, T_n \in \mathbb{B}(\mathcal{H})$. Recently, Sheikhsosseini et al. in [18] showed

$$w_p^p(A_1^* T_1 B_1, \dots, A_n^* T_n B_n) \leq \frac{n^{1-\frac{1}{r}}}{2^{\frac{1}{r}}} \left\| \sum_{i=1}^n [B_i^* f^2(|T_i|) B_i]^{rp} + [A_i^* g^2(|T_i^*|) A_i]^{rp} \right\|^{\frac{1}{r}} - \inf_{\|x\|=1} \zeta(x), \quad (1.3)$$

where $\zeta(x) = \frac{1}{2} \sum_{i=1}^n \left(\langle [B_i^* f^2(|T_i|) B_i]^p x, x \rangle^{\frac{1}{2}} - \langle [A_i^* f^2(|T_i^*|) A_i]^p x, x \rangle^{\frac{1}{2}} \right)^2$. They also presented the following inequality

$$w_p(T_1, \dots, T_n) \leq \frac{1}{2} \left[\sum_{i=1}^n \left(\| |T_i|^{2\alpha} + |T_i^*|^{2(1-\alpha)} \| - 2 \inf_{\|x\|=1} \zeta_i(x) \right)^p \right]^{\frac{1}{p}}, \quad (1.4)$$

in which $0 \leq \alpha \leq 1$, $p \geq 1$ and $\zeta_i(x) = \frac{1}{2} (\langle |T_i|^{2\alpha} x, x \rangle^{\frac{1}{2}} - \langle |T_i^*|^{2(1-\alpha)} x, x \rangle^{\frac{1}{2}})^2$.

In the same paper, they showed

$$w_p^p(T_1, \dots, T_n) \leq \frac{1}{2} \left\| \sum_{i=1}^n (|T_i|^{2\alpha p} + |T_i^*|^{2(1-\alpha)p}) \right\| - \inf_{\|x\|=1} \zeta(x), \quad (1.5)$$

where $\zeta(x) = \frac{1}{2} \sum_{i=1}^n (\langle |T_i|^{2\alpha p} x, x \rangle^{\frac{1}{2}} - \langle |T_i^*|^{2(1-\alpha)p} x, x \rangle^{\frac{1}{2}})^2$.

Moreover, they established the inequalities

$$w_p^p(T_1, \dots, T_n) \leq \left\| \sum_{i=1}^n \alpha |T_i|^p + (1-\alpha) |T_i^*|^p \right\| - \inf_{\|x\|=1} \zeta(x), \quad (1.6)$$

and

$$w_p^r(|T_1|, \dots, |T_n|) w_q^r(|T_1^*|, \dots, |T_n^*|) \leq \frac{r}{p} \left\| \sum_{i=1}^n |T_i|^p \right\| + \frac{r}{q} \left\| \sum_{i=1}^n |T_i^*|^q \right\| - \inf_{\|x\|=\|y\|=1} \delta(x, y), \quad (1.7)$$

where $\zeta(x) = \min\{\alpha, 1-\alpha\} \sum_{i=1}^n (\langle |T_i|^p x, x \rangle^{\frac{1}{2}} - \langle |T_i^*|^p x, x \rangle^{\frac{1}{2}})^2$ and

$$\delta(x, y) = \frac{r}{p} \left(\sqrt[2]{\sum_{i=1}^n \langle |T_i|^p x, y \rangle^p} - \sqrt[2]{\sum_{i=1}^n \langle |T_i^*|^q x, y \rangle^q} \right)^2.$$

Assume that $X \in \mathbb{B}(\mathcal{H})$. The mixed Heinz means are defined by

$$H_\alpha(A, B) = \frac{A^\alpha X B^{1-\alpha} + A^{1-\alpha} X B^\alpha}{2},$$

in which $0 \leq \alpha \leq 1$ and $A, B \geq 0$, see [10]. In [17], the authors showed that

$$w^r(A^\alpha X B^{1-\alpha}) \leq \|X\|^r \|\alpha A^r + (1-\alpha) B^r\|, \quad (1.8)$$

where $A, B, X \in \mathbb{B}(\mathcal{H})$ such that A, B are positive, $r \geq 2$ and $0 \leq \alpha \leq 1$.

Using inequality (1.8), they presented an upper bound for Heinz means of matrices as follows:

$$w^r(H_\alpha(A, B)) \leq \|X\|^r \left\| \frac{A^r + B^r}{2} \right\|. \quad (1.9)$$

In this present paper, we refine inequalities (1.3)-(1.9). We also find an upper bound for the functional w_p .

2. MAIN RESULTS

To prove our numerical radius inequalities, we need several known lemmas. The first lemma is a simple result of the classical Jensen, Young and a generalized mixed Cauchy-Schwarz inequalities [9, 11].

Lemma 2.1. *Let $a, b \geq 0$, $0 \leq \nu \leq 1$ and $r \neq 0$. Then*

(a) $a^\nu b^{1-\nu} \leq \nu a + (1 - \nu)b \leq (\nu a^r + (1 - \nu)b^r)^{\frac{1}{r}}$ for $r \geq 1$.

(b) *If $T \in \mathbb{B}(\mathcal{H})$ and $x, y \in \mathcal{H}$ be any vectors, then*

$$|\langle Tx, y \rangle|^2 \leq \langle |T|^{2\nu} x, x \rangle \langle |T^*|^{2(1-\nu)} y, y \rangle.$$

(c) *If f, g are nonnegative continuous functions on $[0, \infty)$ which are satisfying the relation $f(t)g(t) = t$ ($t \in [0, \infty)$), then*

$$|\langle Tx, y \rangle| \leq \|f(|T|)x\| \|g(|T^*|)x\|$$

for all $x, y \in \mathcal{H}$.

Lemma 2.2. (McCarty inequality [11]). *Let $T \in \mathbb{B}(\mathcal{H})$, $T \geq 0$ and $x \in \mathcal{H}$ be a unit vector. Then*

(a) $\langle Tx, x \rangle^r \leq \langle T^r x, x \rangle$ for $r \geq 1$;

(b) $\langle T^r x, x \rangle \leq \langle Tx, x \rangle^r$ for $0 < r \leq 1$.

Now, by using inequality (1.2) we get the first result.

Theorem 2.3. *Let $A, B, X \in \mathbb{B}(\mathcal{H})$ such that A, B are positive, $r \geq 2$ and $0 \leq \nu \leq 1$. Then*

$$w^r(A^\nu X B^{1-\nu}) \leq \|X\|^r \left[\|\nu A^r + (1 - \nu)B^r\| - \inf_{\|x\|=1} \eta(x) \right], \quad (2.1)$$

where

$$\begin{aligned} \eta(x) &= \sum_{j=1}^N \left((-1)^{r_j} 2^{j-1} \nu + (-1)^{r_j+1} \left[\frac{r_j+1}{2} \right] \right) \\ &\quad \times \left(\sqrt[2^j]{\langle B^r x, x \rangle^{2^{j-1}-k_j} \langle A^r x, x \rangle^{k_j}} - \sqrt[2^j]{\langle A^r x, x \rangle^{k_j+1} \langle B^r x, x \rangle^{2^{j-1}-k_j-1}} \right)^2. \end{aligned}$$

Proof. Let $x \in \mathcal{H}$ be unit vector. Then

$$\begin{aligned}
 |\langle A^\nu X B^{1-\nu} x, x \rangle|^r &= |\langle X B^{1-\nu} x, A^\nu x \rangle|^r \\
 &\leq \|X\|^r \|B^{1-\nu} x\|^r \|A^\nu x\|^r \\
 &= \|X\|^r \langle B^{2(1-\nu)} x, x \rangle^{\frac{r}{2}} \langle A^{2\nu} x, x \rangle^{\frac{r}{2}} \\
 &\leq \|X\|^r \langle A^r x, x \rangle^\nu \langle B^r x, x \rangle^{1-\nu} \quad (\text{by Lemma 2.2}) \\
 &\leq \|X\|^r [\nu \langle A^r x, x \rangle + (1-\nu) \langle B^r x, x \rangle] \\
 &\quad - \|X\|^r \sum_{j=1}^N \left((-1)^{r_j} 2^{j-1} \nu + (-1)^{r_j+1} \left[\frac{r_j+1}{2} \right] \right) \times \\
 &\quad \times \left(\sqrt[2^j]{\langle B^r x, x \rangle^{2^{j-1}-k_j} \langle A^r x, x \rangle^{k_j}} - \sqrt[2^j]{\langle A^r x, x \rangle^{k_j+1} \langle B^r x, x \rangle^{2^{j-1}-k_j-1}} \right)^2 \\
 &\quad (\text{by inequality (1.2)}).
 \end{aligned}$$

Taking the supremum over $x \in \mathcal{H}$ with $\|x\| = 1$ in the above inequality we deduce the desired inequality. \square

Remark 2.4. Let $N = 1$ in inequality (2.1). Then

$$w^r(A^\nu X B^{1-\nu}) \leq \|X\|^r \left[\|\nu A^r + (1-\nu) B^r\| - \inf_{\|x\|=1} \eta(x) \right], \quad (2.2)$$

in which $\eta(x) = r_0 \left(\langle A^r x, x \rangle^{\frac{1}{2}} - \langle B^r x, x \rangle^{\frac{1}{2}} \right)^2$ and $r_0 = \min\{\nu, 1-\nu\}$. Hence inequality (2.2) is a refinement of inequality (1.8).

Using Theorem 2.3 we can find an upper bound for Heinz means of matrices that it is a refinement of (1.9).

Theorem 2.5. *Suppose $A, B, X \in \mathbb{B}(\mathcal{H})$ such that A, B are positive. Then*

$$w^r(H_\nu(A, B)) \leq \|X\|^r \left[\left\| \frac{A^r + B^r}{2} \right\| - \frac{1}{2} \inf \zeta(x) \right],$$

where $r \geq 2$, $0 \leq \nu \leq 1$, $n \in \mathbb{N}$ and

$$\begin{aligned}
 \zeta(x) &= \sum_{j=1}^N \left((-1)^{r_j} 2^{j-1} + (-1)^{r_j+1} \left[\frac{r_j+1}{2} \right] \right) \\
 &\quad \times \left(\sqrt[2^j]{\langle B^r x, x \rangle^{2^{j-1}-k_j} \langle A^r x, x \rangle^{k_j}} - \sqrt[2^j]{\langle A^r x, x \rangle^{k_j+1} \langle B^r x, x \rangle^{2^{j-1}-k_j-1}} \right)^2.
 \end{aligned}$$

Proof. For unit vector $x \in \mathcal{H}$, we have

$$\begin{aligned}
& \left| \left\langle \frac{A^\nu X B^{1-\nu} + A^{1-\nu} X B^\nu}{2} x, x \right\rangle \right|^r \\
& \leq \left(\frac{|\langle A^\nu X B^{1-\nu} x, x \rangle| + |\langle A^{1-\nu} X B^\nu x, x \rangle|}{2} \right)^r \\
& \leq \frac{|\langle A^\nu X B^{1-\nu} x, x \rangle|^r + |\langle A^{1-\nu} X B^\nu x, x \rangle|^r}{2} \\
& \leq \frac{\|X\|^r}{2} [\langle \nu A^r + (1-\nu) B^r x, x \rangle] - \frac{\|X\|^r}{2} \sum_{j=1}^N \left((-1)^{r_j} 2^{j-1} \nu + (-1)^{r_j+1} \left\lfloor \frac{r_j+1}{2} \right\rfloor \right) \\
& \quad \times \left(\sqrt[2^j]{b^{2^{j-1}-k_j} a^{k_j}} - \sqrt[2^j]{a^{k_j+1} b^{2^{j-1}-k_j-1}} \right)^2 \\
& \quad + \frac{\|X\|^r}{2} [\langle (1-\nu) A^r + \nu B^r x, x \rangle] - \frac{\|X\|^r}{2} \sum_{j=1}^N \left((-1)^{r_j} 2^{j-1} \nu + (-1)^{r_j+1} \left\lfloor \frac{r_j+1}{2} \right\rfloor \right) \\
& \quad \times \left(\sqrt[2^j]{\langle B^r x, x \rangle^{2^{j-1}-k_j} \langle A^r x, x \rangle^{k_j}} - \sqrt[2^j]{\langle A^r x, x \rangle^{k_j+1} \langle B^r x, x \rangle^{2^{j-1}-k_j-1}} \right)^2 \\
& = \|X\|^r \left[\left\langle \frac{A^r + B^r}{2} x, x \right\rangle \right] - \frac{\|X\|^r}{2} \sum_{j=1}^N \left((-1)^{r_j} 2^{j-1} + (-1)^{r_j+1} \left\lfloor \frac{r_j+1}{2} \right\rfloor \right) \\
& \quad \times \left(\sqrt[2^j]{\langle B^r x, x \rangle^{2^{j-1}-k_j} \langle A^r x, x \rangle^{k_j}} - \sqrt[2^j]{\langle A^r x, x \rangle^{k_j+1} \langle B^r x, x \rangle^{2^{j-1}-k_j-1}} \right)^2.
\end{aligned}$$

If we take the supremum over $x \in \mathcal{H}$ with $\|x\| = 1$, then we deduce the desired inequality. \square

In the next theorem we show a refinement of inequality (1.3).

Theorem 2.6. *Let $T_i, A_i, B_i \in \mathbb{B}(\mathcal{H})$ ($1 \leq i \leq n$) and let f and g be nonnegative continuous functions on $[0, \infty)$ satisfying $f(t)g(t) = t$ for all $t \in [0, \infty)$. Then*

$$w_p^p(A_1^* T_1 B_1, \dots, A_n^* T_n B_n) \leq \frac{n^{1-\frac{1}{r}}}{2^{\frac{1}{r}}} \left\| \sum_{i=1}^n [B_i^* f^2(|T_i|) B_i]^{rp} + [A_i^* g^2(|T_i^*|) A_i]^{rp} \right\|^{\frac{1}{r}} - \inf_{\|x\|=1} \eta(x), \quad (2.3)$$

where $p, r \geq 1$, $N \in \mathbb{N}$ and

$$\begin{aligned}
\eta(x) &= \frac{1}{2} \sum_{i=1}^n \sum_{j=1}^N \left(\sqrt[2^j]{\langle (A_i^* g^2(|T_i^*|) A_i)^{p x}, x \rangle^{2^{j-1}-k_j} \langle (B_i^* f^2(|T_i|) B_i)^{p x}, x \rangle^{k_j}} \right. \\
& \quad \left. - \sqrt[2^j]{\langle (B_i^* f^2(|T_i|) B_i)^{p x}, x \rangle^{k_j+1} \langle (A_i^* g^2(|T_i^*|) A_i)^{p x}, x \rangle^{2^{j-1}-k_j-1}} \right)^2.
\end{aligned}$$

Proof. Let $x \in \mathcal{H}$ be any unit vector. Then

$$\begin{aligned}
 \sum_{i=1}^n |\langle A_i^* T_i B_i x, x \rangle|^p &= \sum_{i=1}^n |\langle T_i B_i x, A_i x \rangle|^p \\
 &\leq \sum_{i=1}^n \|f(|T_i|) B_i x\|^p \|g(|T_i^*|) A_i x\|^p \\
 &\quad \text{(by Lemma (2.1), (c))} \\
 &= \sum_{i=1}^n \langle f(|T_i|) B_i x, f(|T_i|) B_i x \rangle^{\frac{p}{2}} \langle g(|T_i^*|) A_i x, g(|T_i^*|) A_i x \rangle^{\frac{p}{2}} \\
 &= \sum_{i=1}^n \langle B_i^* f^2(|T_i|) B_i x, x \rangle^{\frac{p}{2}} \langle A_i^* g^2(|T_i^*|) A_i x, x \rangle^{\frac{p}{2}} \\
 &\leq \sum_{i=1}^n \langle (B_i^* f^2(|T_i|) B_i)^p x, x \rangle^{\frac{1}{2}} \langle (A_i^* g^2(|T_i^*|) A_i)^p x, x \rangle^{\frac{1}{2}} \\
 &\quad \text{(by Lemma (2.2), (a))} \\
 &\leq \sum_{i=1}^n \left[\left(\frac{1}{2} \langle (B_i^* f^2(|T_i|) B_i)^{pr} x, x \rangle + \frac{1}{2} \langle (A_i^* g^2(|T_i^*|) A_i)^{pr} x, x \rangle \right)^{\frac{1}{r}} \right] \\
 &\quad \text{(by (1.2))} \\
 &\leq \frac{1}{2} \sum_{i=1}^n \sum_{j=1}^N \left(\sqrt[2^j]{\langle (A_i^* g^2(|T_i^*|) A_i)^p x, x \rangle^{2^{j-1}-k_j} \langle (B_i^* f^2(|T_i|) B_i)^p x, x \rangle^{k_j}} \right. \\
 &\quad \left. - \sqrt[2^j]{\langle (B_i^* f^2(|T_i|) B_i)^p x, x \rangle^{k_j+1} \langle (A_i^* g^2(|T_i^*|) A_i)^p x, x \rangle^{2^{j-1}-k_j-1}} \right)^2 \\
 &\leq \frac{n^{1-\frac{1}{r}}}{2^{\frac{1}{r}}} \left\langle \left(\sum_{i=1}^n \left([B_i^* f^2(|T_i|) B_i]^{rp} + [A_i^* g^2(|T_i^*|) A_i]^{rp} \right) \right) x, x \right\rangle^{\frac{1}{r}} \\
 &\leq \frac{1}{2} \sum_{i=1}^n \sum_{j=1}^N \left(\sqrt[2^j]{\langle (A_i^* g^2(|T_i^*|) A_i)^p x, x \rangle^{2^{j-1}-k_j} \langle (B_i^* f^2(|T_i|) B_i)^p x, x \rangle^{k_j}} \right. \\
 &\quad \left. - \sqrt[2^j]{\langle (B_i^* f^2(|T_i|) B_i)^p x, x \rangle^{k_j+1} \langle (A_i^* g^2(|T_i^*|) A_i)^p x, x \rangle^{2^{j-1}-k_j-1}} \right)^2
 \end{aligned}$$

By taking supremum on unit vector x in \mathcal{H} we reach the desired inequality. \square

Corollary 2.7. Let $A_i, B_i \in \mathbb{B}(\mathcal{H})$ ($1 \leq i \leq n$). Then for $r, p \geq 1$ we have

$$w_p^p(A_1^* B_1, \dots, A_n^* B_n) \leq \frac{n^{1-\frac{1}{r}}}{2^{\frac{1}{r}}} \left\| \sum_{i=1}^n (|B_i|^{2rp} + |A_i|^{2rp}) \right\|^{\frac{1}{r}} - \inf_{\|x\|=1} \eta(x),$$

where

$$\begin{aligned}
 \eta(x) &= \frac{1}{2} \sum_{i=1}^n \sum_{j=1}^N \left(\sqrt[2^j]{\langle (|A_i|^{2p} x, x)^{2^{j-1}-k_j} \langle (|B_i|^{2p} x, x)^{k_j} \right.} \\
 &\quad \left. - \sqrt[2^j]{\langle (|B_i|^{2p} x, x)^{k_j+1} \langle (|A_i|^{2p} x, x)^{2^{j-1}-k_j-1} \right)^2.
 \end{aligned}$$

Proof. Choosing $f(t) = g(t) = t^{\frac{1}{2}}$ and $T_i = I$ for $i = 1, 2, \dots, n$ in Theorem 2.6, we get the desired result. \square

Corollary 2.8. *Let $T_i \in \mathbb{B}(\mathcal{H})$ ($1 \leq i \leq n$), let f and g be nonnegative continuous functions on $[0, \infty)$ such that $f(t)g(t) = t$ for all $t \in [0, \infty)$ and $r, p \geq 1$. Then*

$$w_p^p(T_1, \dots, T_n) \leq \frac{n^{1-\frac{1}{r}}}{2^{\frac{1}{r}}} \left\| \sum_{i=1}^n (|f^{2rp}(|T_i|) + g^{2rp}(|T_i^*|)) \right\|_{\frac{1}{r}} - \inf_{\|x\|=1} \eta(x), \quad (2.4)$$

where

$$\begin{aligned} \eta(x) &= \frac{1}{2} \sum_{i=1}^n \sum_{j=1}^N \left(\sqrt[2^j]{\langle g^{2p}(|T_i^*|)x, x \rangle^{2^{j-1}-k_j} \langle f^{2p}(|T_i|)x, x \rangle^{k_j}} \right. \\ &\quad \left. - \sqrt[2^j]{\langle f^{2p}(|T_i|)x, x \rangle^{k_j+1} \langle g^{2p}(|T_i^*|)x, x \rangle^{2^{j-1}-k_j-1}} \right)^2. \end{aligned}$$

In particular,

$$w_p^p(T_1, \dots, T_n) \leq \frac{1}{2} \left\| \sum_{i=1}^n (|T_i|^{2\alpha p} + |T_i^*|^{2(1-\alpha)p}) \right\| - \inf_{\|x\|=1} \eta(x), \quad (2.5)$$

where $0 \leq \alpha \leq 1$ and

$$\begin{aligned} \eta(x) &= \frac{1}{2} \sum_{i=1}^n \sum_{j=1}^N \left(\sqrt[2^j]{\langle |T_i^*|^{2(1-\alpha)p}x, x \rangle^{2^{j-1}-k_j} \langle |T_i|^{2\alpha p}x, x \rangle^{k_j}} \right. \\ &\quad \left. - \sqrt[2^j]{\langle |T_i|^{2\alpha p}x, x \rangle^{k_j+1} \langle |T_i^*|^{2(1-\alpha)p}x, x \rangle^{2^{j-1}-k_j-1}} \right)^2. \end{aligned}$$

Proof. Selecting $A_i = B_i = I$ for $i = 1, 2, \dots, n$ in Theorem 2.6, we get the first result. Letting $f(t) = t^\alpha$, $g(t) = t^{1-\alpha}$, $r = 1$ and $B_i = A_i = I$ for $i = 1, 2, \dots, n$ in inequality (2.4), we reach the second inequality. \square

Remark 2.9. Note that inequality (2.5) is a refinement of inequality (1.5), since

$$\begin{aligned} \frac{1}{2} \sum_{i=1}^n (\langle |T_i|^{2\alpha p}x, x \rangle^{\frac{1}{2}} - \langle |T_i^*|^{2(1-\alpha)p}x, x \rangle^{\frac{1}{2}})^2 &\leq \frac{1}{2} \sum_{i=1}^n \sum_{j=1}^N \left(\sqrt[2^j]{\langle |T_i^*|^{2(1-\alpha)p}x, x \rangle^{2^{j-1}-k_j} \langle |T_i|^{2\alpha p}x, x \rangle^{k_j}} \right. \\ &\quad \left. - \sqrt[2^j]{\langle |T_i|^{2\alpha p}x, x \rangle^{k_j+1} \langle |T_i^*|^{2(1-\alpha)p}x, x \rangle^{2^{j-1}-k_j-1}} \right)^2. \end{aligned}$$

Now by letting $n = 2$, $N = 1$, $T_1 = B$ and $T_2 = C$ in Theorem 2.6, we obtain the following consequence.

Corollary 2.10. *Let $B, C \in \mathbb{B}(\mathcal{H})$. Then for all $p \geq 1$ and $0 \leq \alpha \leq 1$*

$$w_p^p(B, C) \leq \frac{1}{2} \left\| |B|^{2\alpha p} + |B^*|^{2(1-\alpha)p} + |C|^{2\alpha p} + |C^*|^{2(1-\alpha)p} \right\| - \inf_{\|x\|=1} \eta(x),$$

where

$$\eta(x) = \frac{1}{2} \left[(\langle |B|^{2\alpha p}x, x \rangle^{\frac{1}{2}} - \langle |B^*|^{2(1-\alpha)p}x, x \rangle^{\frac{1}{2}})^2 + (\langle |C|^{2\alpha p}x, x \rangle^{\frac{1}{2}} - \langle |C^*|^{2(1-\alpha)p}x, x \rangle^{\frac{1}{2}})^2 \right].$$

Theorem 2.11. *Let $T_i \in \mathbb{B}(\mathcal{H})$ ($1 \leq i \leq n$). Then*

$$w_p(T_1, \dots, T_n) \leq \frac{1}{2} \left[\sum_{i=1}^n \left(\left\| |T_i|^{2\alpha} + |T_i^*|^{2(1-\alpha)} \right\| - 2 \inf_{\|x\|=1} \eta_i(x) \right)^p \right]^{\frac{1}{p}}, \quad (2.6)$$

where $p \geq 1$, $0 \leq \alpha \leq 1$ and

$$\begin{aligned} \eta_i(x) = & \frac{1}{2} \sum_{j=1}^N \left(\sqrt[2^j]{\langle |T_i^*|^{2(1-\alpha)p} x, x \rangle^{2^{j-1}-k_j} \langle |T_i|^{2\alpha p} x, x \rangle^{k_j}} \right. \\ & \left. - \sqrt[2^j]{\langle |T_i|^{2\alpha p} x, x \rangle^{k_j+1} \langle |T_i^*|^{2(1-\alpha)p} x, x \rangle^{2^{j-1}-k_j-1}} \right)^2. \end{aligned}$$

Proof. By using of Lemma 2.1 and inequality (1.2), for any unit vector $x \in \mathcal{H}$ we have

$$\begin{aligned} \sum_{i=1}^n |\langle T_i x, x \rangle|^p & \leq \sum_{i=1}^n \left(\langle |T_i|^{2\alpha} x, x \rangle^{\frac{1}{2}} \langle |T_i^*|^{2(1-\alpha)} x, x \rangle^{\frac{1}{2}} \right)^p \\ & \quad \text{(by Lemma (2.1), (b))} \\ & \leq \frac{1}{2^p} \sum_{i=1}^n \left[\langle |T_i|^{2\alpha} x, x \rangle + \langle |T_i^*|^{2(1-\alpha)} x, x \rangle - \right. \\ & \quad \left. - \sum_{j=1}^N \left(\sqrt[2^j]{\langle |T_i^*|^{2(1-\alpha)p} x, x \rangle^{2^{j-1}-k_j} \langle |T_i|^{2\alpha p} x, x \rangle^{k_j}} \right. \right. \\ & \quad \left. \left. - \sqrt[2^j]{\langle |T_i|^{2\alpha p} x, x \rangle^{k_j+1} \langle |T_i^*|^{2(1-\alpha)p} x, x \rangle^{2^{j-1}-k_j-1}} \right)^2 \right]^p \\ & \quad \text{(by (1.2))} \\ & = \frac{1}{2^p} \sum_{i=1}^n \left[\langle |T_i|^{2\alpha} + |T_i^*|^{2(1-\alpha)} x, x \rangle \right. \\ & \quad \left. - \sum_{j=1}^N \left(\sqrt[2^j]{\langle |T_i^*|^{2(1-\alpha)p} x, x \rangle^{2^{j-1}-k_j} \langle |T_i|^{2\alpha p} x, x \rangle^{k_j}} \right. \right. \\ & \quad \left. \left. - \sqrt[2^j]{\langle |T_i|^{2\alpha p} x, x \rangle^{k_j+1} \langle |T_i^*|^{2(1-\alpha)p} x, x \rangle^{2^{j-1}-k_j-1}} \right)^2 \right]^p. \end{aligned}$$

Thus

$$\begin{aligned}
\left(\sum_{i=1}^n |\langle T_i x, x \rangle|^p \right)^{\frac{1}{p}} &\leq \frac{1}{2} \left[\sum_{i=1}^n \left(\langle |T_i|^{2\alpha} + |T_i^*|^{2(1-\alpha)} x, x \rangle - \right. \right. \\
&\quad \left. \left. - \sum_{j=1}^N \left(\sqrt[2^j]{\langle |T_i^*|^{2(1-\alpha)p} x, x \rangle^{2^{j-1}-k_j} \langle |T_i|^{2\alpha p} x, x \rangle^{k_j}} \right. \right. \right. \\
&\quad \left. \left. \left. - \sqrt[2^j]{\langle |T_i|^{2\alpha p} x, x \rangle^{k_j+1} \langle |T_i^*|^{2(1-\alpha)p} x, x \rangle^{2^{j-1}-k_j-1}} \right)^2 \right)^p \right]^{\frac{1}{p}} \\
&= \frac{1}{2} \left[\sum_{i=1}^n \left(\langle |T_i|^{2\alpha} + |T_i^*|^{2(1-\alpha)} x, x \rangle - 2\eta_i(x) \right)^p \right]^{\frac{1}{p}}.
\end{aligned}$$

Now, by taking the supremum over all unit vector $x \in \mathcal{H}$ we get the desired result. \square

Remark 2.12. If $N = 1$ in inequality (2.6), then we reach to inequality (1.4), it follows from

$$\begin{aligned}
\frac{1}{2} \left(\langle |T_i|^{2\alpha} x, x \rangle^{\frac{1}{2}} - \langle |T_i^*|^{2(1-\alpha)} x, x \rangle^{\frac{1}{2}} \right)^2 &\leq \frac{1}{2} \sum_{j=1}^N \left(\sqrt[2^j]{\langle |T_i^*|^{2(1-\alpha)p} x, x \rangle^{2^{j-1}-k_j} \langle |T_i|^{2\alpha p} x, x \rangle^{k_j}} \right. \\
&\quad \left. - \sqrt[2^j]{\langle |T_i|^{2\alpha p} x, x \rangle^{k_j+1} \langle |T_i^*|^{2(1-\alpha)p} x, x \rangle^{2^{j-1}-k_j-1}} \right)^2,
\end{aligned}$$

that inequality (2.6) is a refinement of inequality (1.4).

Theorem 2.13. *Let $T_i \in \mathbb{B}(\mathcal{H})$ ($1 \leq i \leq n$). Then for $0 \leq \alpha \leq 1$ and $p \geq 2$*

$$w_p^p(T_1, \dots, T_n) \leq \left\| \sum_{i=1}^n (\alpha |T_i|^p + (1-\alpha) |T_i^*|^p) \right\| - \inf_{\|x\|=1} \eta(x), \quad (2.7)$$

where

$$\begin{aligned}
\eta(x) &= \sum_{i=1}^n \left(\sum_{j=1}^N \left((-1)^{r_j} 2^{j-1} \alpha + (-1)^{r_j+1} \left[\frac{r_j+1}{2} \right] \right) \right. \\
&\quad \left. \times \left(\sqrt[2^j]{\langle |T_i^*|^p x, x \rangle^{2^{j-1}-k_j} \langle |T_i|^p x, x \rangle^{k_j}} - \sqrt[2^j]{\langle |T_i|^p x, x \rangle^{k_j+1} \langle |T_i^*|^p x, x \rangle^{2^{j-1}-k_j-1}} \right)^2 \right)
\end{aligned}$$

Proof. For every unit vector $x \in \mathcal{H}$ we have

$$\begin{aligned}
 & \sum_{i=1}^n |\langle T_i x, x \rangle|^p \\
 &= \sum_{i=1}^n (|\langle T_i x, x \rangle|^2)^{\frac{p}{2}} \\
 &\leq \sum_{i=1}^n (\langle |T_i|^{2\alpha} x, x \rangle \langle |T_i^*|^{2(1-\alpha)} x, x \rangle)^{\frac{p}{2}} && \text{(by Lemma 2.1, (b))} \\
 &\leq \sum_{i=1}^n (\langle |T_i|^p x, x \rangle^\alpha \langle |T_i^*|^p x, x \rangle^{1-\alpha}) && \text{(by Lemma 2.2, (b))} \\
 &\leq \sum_{i=1}^n (\alpha \langle |T_i|^p x, x \rangle + (1-\alpha) \langle |T_i^*|^p x, x \rangle) - \\
 &\quad - \sum_{i=1}^n \left(\sum_{j=1}^N \left((-1)^{r_j} 2^{j-1} \alpha + (-1)^{r_j+1} \left[\frac{r_j+1}{2} \right] \right) \right) \\
 &\quad \times \left(\sqrt[2^j]{\langle |T_i^*|^p x, x \rangle^{2^{j-1}-k_j} \langle |T_i|^p x, x \rangle^{k_j}} - \sqrt[2^j]{\langle |T_i|^p x, x \rangle^{k_j+1} \langle |T_i^*|^p x, x \rangle^{2^{j-1}-k_j-1}} \right)^2 \\
 & && \text{(by (1.2))} \\
 &\leq \sum_{i=1}^n \langle (\alpha |T_i|^p + (1-\alpha) |T_i^*|^p) x, x \rangle - \inf_{\|x\|=1} \eta(x) \\
 &= \left\langle \sum_{i=1}^n (\alpha |T_i|^p + (1-\alpha) |T_i^*|^p) x, x \right\rangle - \inf_{\|x\|=1} \eta(x).
 \end{aligned}$$

Now by taking supremum over unit vector $x \in \mathcal{H}$ we get. \square

Remark 2.14. If we put $N = 1$ in inequality (2.7), then we get inequality (1.6). Hence inequality (2.7) is refinement of (1.6).

In [16, Remark 3.10], the author showed

$$w_p^p(B, C) \leq \frac{1}{2} \| |B|^p + |B^*|^p + |C|^p + |C^*|^p \|, \quad (2.8)$$

in which $B, C \in \mathbb{B}(\mathcal{H})$ and $p \geq 2$. In the following result we show a refinement of (2.8).

Corollary 2.15. *Let $B, C \in \mathbb{B}(\mathcal{H})$. Then for $p \geq 2$,*

$$w_p^p(B, C) \leq \frac{1}{2} \| |B|^p + |B^*|^p + |C|^p + |C^*|^p \| - \inf_{\|x\|=1} \eta(x), \quad (2.9)$$

where $\eta(x) = \frac{1}{2} \left((\langle |B|^p x, x \rangle^{\frac{1}{2}} - \langle |B^*|^p x, x \rangle^{\frac{1}{2}})^2 - (\langle |C|^p x, x \rangle^{\frac{1}{2}} - \langle |C^*|^p x, x \rangle^{\frac{1}{2}})^2 \right)$.
 In particular, if $A \in \mathbb{B}(\mathcal{H})$, then

$$w^2(A) \leq \frac{1}{2} \|A^*A + AA^*\|.$$

Proof. If we take $N = 1$, $n = 2$, $T_1 = B$, $T_2 = C$, and $\alpha = \frac{1}{2}$ in Theorem 2.13, we get the first inequality.

In particular case, let $A = B + iC$ be the Cartesian decomposition of A . Then $A^*A + AA^* = 2(B^2 + C^2)$, and $\inf_{\|x\|=1} \eta(x) = 0$. Thus, for $p = 2$, inequality (2.9) can be written as

$$w_2^2(B, C) \leq \|B^2 + C^2\| = \frac{1}{2} \|A^*A + AA^*\|.$$

The desired inequality follows by noting that

$$w_2^2(B, C) = \sup_{\|x\|=1} \{ |\langle Bx, x \rangle|^2 + |\langle Cx, x \rangle|^2 \} = \sup_{\|x\|=1} |\langle Ax, x \rangle|^2 = w^2(A).$$

□

Theorem 2.16. Let $T_i \in \mathbb{B}(\mathcal{H})$ ($1 \leq i \leq n$), $r \geq 1$, and $p \geq q \geq 1$ with $\frac{1}{p} + \frac{1}{q} = \frac{1}{r}$. Then

$$w_p^r(|T_1|, \dots, |T_n|) w_q^r(|T_1^*|, \dots, |T_n^*|) \leq \frac{r}{p} \left\| \sum_{i=1}^n |T_i|^p \right\| + \frac{r}{q} \left\| \sum_{i=1}^n |T_i^*|^q \right\| - \inf_{\|x\|=\|y\|=1} \lambda(x, y), \quad (2.10)$$

where

$$\begin{aligned} \lambda(x, y) = & \sum_{j=1}^N \left((-1)^{r_j} 2^{j-1} \left(\frac{r}{p} + (-1)^{r_j+1} \left[\frac{r_j+1}{2} \right] \right) \left(\sqrt[2^j]{\left(\sum_{i=1}^n \langle |T_i^*|^q |x, y \rangle^q \right)^{2^{j-1}-k_j} \left(\sum_{i=1}^n \langle |T_i|^p |x, y \rangle^p \right)^{k_j}} \right. \right. \\ & \left. \left. - \sqrt[2^j]{\left(\sum_{i=1}^n \langle |T_i|^p |x, y \rangle^p \right)^{k_j+1} \left(\sum_{i=1}^n \langle |T_i^*|^q |x, y \rangle^q \right)^{2^{j-1}-k_j-1}} \right)^2. \end{aligned}$$

Proof. Let $x, y \in \mathbb{B}(\mathcal{H})$ be unit vectors. Applying inequality (1.2), we get

$$\begin{aligned}
 & \left(\left(\sum_{i=1}^n \langle |T_i|x, y\rangle^p \right)^{\frac{1}{p}} \left(\sum_{i=1}^n \langle |T_i^*|x, y\rangle^q \right)^{\frac{1}{q}} \right)^r \leq \frac{r}{p} \sum_{i=1}^n \langle |T_i|x, y\rangle^p + \frac{r}{q} \sum_{i=1}^n \langle |T_i^*|x, y\rangle^q \\
 & \quad - \sum_{j=1}^N \left((-1)^{r_j} 2^{j-1} \left(\frac{r}{p} \right) + (-1)^{r_j+1} \left[\frac{r_j+1}{2} \right] \right) \\
 & \quad \times \left(\sqrt[2^j]{ \left(\sum_{i=1}^n \langle |T_i^*|x, y\rangle^q \right)^{2^{j-1}-k_j} \left(\sum_{i=1}^n \langle |T_i|x, y\rangle^p \right)^{k_j} } \right. \\
 & \quad \left. - \sqrt[2^j]{ \left(\sum_{i=1}^n \langle |T_i|x, y\rangle^p \right)^{k_j+1} \left(\sum_{i=1}^n \langle |T_i^*|x, y\rangle^q \right)^{2^{j-1}-k_j-1} } \right)^2 \\
 & \hspace{15em} \text{(by (1.2))} \\
 & \leq \frac{r}{p} \sum_{i=1}^n \langle |T_i|^p x, y \rangle + \frac{r}{q} \sum_{i=1}^n \langle |T_i^*|^q x, y \rangle - \\
 & \quad - \sum_{j=1}^N \left((-1)^{r_j} 2^{j-1} \left(\frac{r}{p} \right) + (-1)^{r_j+1} \left[\frac{r_j+1}{2} \right] \right) \\
 & \quad \times \left(\sqrt[2^j]{ \left(\sum_{i=1}^n \langle |T_i^*|x, y\rangle^q \right)^{2^{j-1}-k_j} \left(\sum_{i=1}^n \langle |T_i|x, y\rangle^p \right)^{k_j} } \right. \\
 & \quad \left. - \sqrt[2^j]{ \left(\sum_{i=1}^n \langle |T_i|x, y\rangle^p \right)^{k_j+1} \left(\sum_{i=1}^n \langle |T_i^*|x, y\rangle^q \right)^{2^{j-1}-k_j-1} } \right)^2 \\
 & \hspace{15em} \text{(by Lemma (2.2), (a))} \\
 & = \frac{r}{p} \left\langle \left(\sum_{i=1}^n |T_i|^p \right) x, y \right\rangle + \frac{r}{q} \left\langle \left(\sum_{i=1}^n |T_i^*|^q \right) x, y \right\rangle \\
 & \quad - \sum_{j=1}^N \left((-1)^{r_j} 2^{j-1} \left(\frac{r}{p} \right) + (-1)^{r_j+1} \left[\frac{r_j+1}{2} \right] \right) \\
 & \quad \times \left(\sqrt[2^j]{ \left(\sum_{i=1}^n \langle |T_i^*|x, y\rangle^q \right)^{2^{j-1}-k_j} \left(\sum_{i=1}^n \langle |T_i|x, y\rangle^p \right)^{k_j} } \right. \\
 & \quad \left. - \sqrt[2^j]{ \left(\sum_{i=1}^n \langle |T_i|x, y\rangle^p \right)^{k_j+1} \left(\sum_{i=1}^n \langle |T_i^*|x, y\rangle^q \right)^{2^{j-1}-k_j-1} } \right)^2
 \end{aligned}$$

By taking supremum on $x, y \in \mathcal{H}$ with $\|x\| = \|y\| = 1$, we get desired inequality. \square

Remark 2.17. If we put $N = 1$ in inequality (2.10), then we reach to inequality (1.7).

It follows from

$$\begin{aligned} & \frac{r}{p} \left(\sqrt[p]{\sum_{i=1}^n \langle |T_i|x, y \rangle^p} - \sqrt[p]{\sum_{i=1}^n \langle |T_i^*|x, y \rangle^q} \right)^2 \\ & \leq \sum_{j=1}^N \left((-1)^{r_j} 2^{j-1} \left(\frac{r}{p} \right) + (-1)^{r_j+1} \left[\frac{r_j+1}{2} \right] \right) \left(\sqrt[p]{\sum_{i=1}^n \langle |T_i^*|x, y \rangle^q} \right)^{2^{j-1}-k_j} \left(\sum_{i=1}^n \langle |T_i|x, y \rangle^p \right)^{k_j} \\ & \quad - \sqrt[p]{\left(\sum_{i=1}^n \langle |T_i|x, y \rangle^p \right)^{k_j+1} \left(\sum_{i=1}^n \langle |T_i^*|x, y \rangle^q \right)^{2^{j-1}-k_j-1}} \right)^2, \end{aligned}$$

that inequality (2.10) is refinement of (1.7).

Corollary 2.18. *Let $T_i \in \mathbb{B}(\mathcal{H})$ ($1 \leq i \leq n$). Then*

$$w_e(|T_1|, \dots, |T_n|) w_e(|T_1^*|, \dots, |T_n^*|) \leq \frac{1}{2} \left(\left\| \sum_{i=1}^n T_i^* T_i \right\| + \left\| \sum_{i=1}^n T_i T_i^* \right\| \right) - \inf_{\|x\|=\|y\|=1} \lambda(x, y),$$

where

$$\begin{aligned} \lambda(x, y) &= \sum_{j=1}^N \left((-1)^{r_j} 2^{j-1} \left(\frac{1}{2} \right) + (-1)^{r_j+1} \left[\frac{r_j+1}{2} \right] \right) \left(\sqrt[p]{\sum_{i=1}^n \langle |T_i^*|x, y \rangle^q} \right)^{2^{j-1}-k_j} \left(\sum_{i=1}^n \langle |T_i|x, y \rangle^p \right)^{k_j} \\ & \quad - \sqrt[p]{\left(\sum_{i=1}^n \langle |T_i|x, y \rangle^p \right)^{2(k_j+1)} \left(\sum_{i=1}^n \langle |T_i^*|x, y \rangle^q \right)^{2(2^{j-1}-k_j-1)}} \right)^2. \end{aligned}$$

Proof. The result obtained by letting $p = q = 2$ and $r = 1$ in inequality (2.10). \square

Corollary 2.19. *Let $T_1, \dots, T_n \in \mathbb{B}(\mathcal{H})$ be positive operators. Then*

$$w_e(T_1, \dots, T_n) \leq \left\| \sum_{i=1}^n T_i^2 \right\|^{\frac{1}{2}}.$$

REFERENCES

1. O. Axelsson, H. Lu and B. Polman, *On the numerical radius of matrices and its application to iterative solution methods*, Linear Multilinear Algebra. **37** (1994), 225–238.
2. M. Bakherad and F. Kittaneh, *Numerical radius inequalities involving commutators of $G1$ operators*, Complex Anal. Oper. Theory (2017). <https://doi.org/10.1007/s11785-017-0726-9>.
3. M. Bakherad, M. Krnic and M.S. Moslehian, *Reverses of the Young inequality for matrices and operators*, Rocky Mountain J. Math. **46**(4) (2016), 1089–1105.

4. M. Bakherad and Kh. Shebrawi, *Upper bounds for numerical radius inequalities involving off-diagonal operator matrices*, Ann. Funct. Anal. (2018) (to appear).
5. M. Boumazgour, H. Nabwey, *A note concerning the numerical range of a basic elementary operator*, Ann. Funct. Anal. **7** (2016), no. 3, 434–441.
6. K.E. Gustafson and D.K.M. Rao, *Numerical Range, The Field of Values of Linear Operators and Matrices*, Springer, New York, 1997.
7. M. Hajmohamadi, R. Lashkaripour, M. Bakherad, *Some generalizations of numerical radius on off-diagonal part of 2×2 operator matrices.*, J. Math. Inequal. **12**(2) (2018), 447457.
8. M. Hajmohamadi, R. Lashkaripour, M. Bakherad, *Some extensions of the Young and Heinz inequalities for matrices*, Bull. Iranian Math. Soc. (2017) (to appear).
9. G.H. Hardy, J.E. Littlewood and G. Polya, *inequalities*, 2nd ed., Cambridge Univ. Press, Cambridge, (1988).
10. R. Kaur, M.S. Moslehian, M. Singh and C. Conde, *Further refinements of the Heinz inequality*, Linear Algebra Appl. **447** (2014), 26–37.
11. F. Kittaneh, *Notes on some inequalities for Hilbert space operators*, Publ. Res. Inst. Math. Sci. **24**(2) (1988), 283–293.
12. F. Kittaneh and Y. Manasrah, *Improved Young and Heinz inequalities for matrices*, J. Math. Anal. Appl. **361** (2010), 262–269.
13. M.S. Moslehian and M. Sattari, *Inequalities for operator space numerical radius of 2×2 block matrices*, J. Math. Phys. **57**(1) (2016), 015201, 15 pp.
14. G. Popescu, *Unitary invariants in multivariable operator theory*, Mem. Amer. Math. Soc. **200**(941) (2009).
15. M. Sababheh and D. Choi, *A complete refinement of Yongs inequality*, J. Math. Anal. Appl. **440** (2016), 379–393.
16. M. Sattari, M.S. Moslehian and K. Shebrawi, *Extension of Euclidean operator radius inequalities*, Math. Scand. **20** (2017), 129–144.
17. M. Sattari, M.S. Moslehian and T. Yamazaki, *Some generalised numerical radius inequalities for Hilbert space operators*, Linear Algebra Appl. **470** (2014), 1–12.
18. A. Sheikhsosseini, M.S. Moslehian and K. Shebrawi, *Inequalities for generalized Euclidean operator radius via Young's inequality*, J. Math. Anal. Appl. **445**(2) (2016), 1516–1529.
19. A. Zamani, *Some lower bounds for the numerical radius of Hilbert space operators*, Adv. Oper. Theory **2** (2) (2017), 98–107.

^{1,2,3} DEPARTMENT OF MATHEMATICS, FACULTY OF MATHEMATICS, UNIVERSITY OF SISTAN AND BALUCHESTAN, ZAHEDAN, I.R.IRAN.

E-mail address: ¹monire.hajmohamadi@yahoo.com

E-mail address: ²lashkari@hamoon.usb.ac.ir

E-mail address: ³mojtaba.bakherad@yahoo.com; bakherad@member.ams.org