

Non-conforming Crouzeix-Raviart element approximation for Stekloff eigenvalues in inverse scattering

Yidu Yang, Yu Zhang, Hai Bi

School of Mathematical Sciences, Guizhou Normal University,
Guiyang, 550001, China

Abstract

In this paper, we use the non-conforming Crouzeix-Raviart element method to solve a Stekloff eigenvalue problem arising in inverse scattering. The weak formulation corresponding to this problem is non-selfadjoint and does not satisfy H^1 -elliptic condition, and its Crouzeix-Raviart element discretization does not meet the Strang lemma condition. We use the standard duality techniques to prove an extension of Strang lemma. And we prove the convergence and error estimate of discrete eigenvalues and eigenfunctions using the spectral perturbation theory for compact operators. Finally, we present some numerical examples not only on uniform meshes but also in an adaptive refined meshes to show that the Crouzeix-Raviart method is efficient for computing real and complex eigenvalues as expected.

Key words Stekloff eigenvalue, Nonconforming Crouzeix-Raviart element, Strang lemma, Error estimates.

1 Introduction

Steklov eigenvalue problems have important physical background and many applications. For instance, they appear in the analysis of stability of mechanical oscillators immersed in a viscous fluid (see [29] and the references therein), in the study of surface waves [10], in the study of the vibration modes of a structure in contact with an incompressible fluid [11] and in the analysis of the antiplane shearing on a system of collinear faults under slip-dependent friction law [19]. Hence, the finite element methods for solving these problems have attracted more and more scholars' attention. Till now, systematical and profound studies on the finite element approximation mainly focus on Steklov eigenvalue problems which satisfy H^1 -elliptic condition (see, e.g., [3, 4, 5, 6, 11, 13, 15, 22, 36, 41,

42, 46, 51, 52] and the references therein).

Recently Cakoni et al. [21] study a new Stekloff eigenvalue problem arising from the inverse scattering theory:

$$\Delta u + k^2 n(x)u = 0 \text{ in } \Omega, \quad \frac{\partial u}{\partial \gamma} = -\lambda u \text{ on } \partial\Omega, \quad (1.1)$$

where Ω is a bounded domain in \mathbb{R}^d ($d = 2, 3$), $\frac{\partial u}{\partial \gamma}$ is the outward normal derivative, k is the wavenumber and $n(x) = n_1(x) + i\frac{n_2(x)}{k}$ is the index of refraction that is a bounded complex valued function with $n_1(x) > 0$ and $n_2(x) \geq 0$.

Note that the weak formulation of (1.1) (see (2.1)) does not satisfy H^1 -elliptic condition. Cakoni et al. [21] analyze the mathematics properties of (1.1) and use conforming finite element method to solve it. Liu et al. [43] then study error estimates of conforming finite element eigenvalues for (1.1).

In this paper, we will study the non-conforming Crouzeix-Raviart element (C-R element) approximation for the problem (1.1). The C-R element was first introduced by Crouzeix and Raviart in [30] in 1973 to solve the stationary Stokes equation. It was also used to solve linear elasticity equations (see [35, 16]), the Laplace equation/eigenvalues (see [7, 14, 17, 23, 24, 25, 33]), Darcy's equation [2], Steklov eigenvalue (see [3, 13, 41, 46, 52]) and so on. The features of our work are as follows:

1. As we know, the convergence and error estimates of the non-conforming finite element method for an eigenvalue problem is based on the convergence and error estimates of the non-conforming finite element method for the corresponding source problem, and Strang lemma (see [49]) is a fundamental analysis tool. However, the sesquilinear form in the C-R element discretization here does not meet the Strang lemma condition. To overcome this difficulty, referring to §5.7 in [17], we use the standard duality techniques to prove an extension of Strang lemma (see Theorem 2). Based on the theorem, we prove the convergence and error estimates of the C-R method for the corresponding source problem. The current paper, to our knowledge, is the first investigation of applying and extending Strang lemma to elliptic boundary value problem that the corresponding sesquilinear form is non-selfadjoint and not coercive.
2. Cakoni et al. [21] write (1.1) as an equivalent eigenvalue problem of the Neumann-to-Dirichlet operator T . In this paper, we write the C-R element approximation of (1.1) as an equivalent eigenvalue problem of the discrete operator T_h , and prove T_h converges T in the sense of norm in $L^2(\partial\Omega)$, thus using Babuska-Osborn spectral approximation theory [8] we prove first the convergence and error estimates of C-R finite element eigenvalues and eigenfunctions for the problem (1.1).
3. We implement some numerical experiments not only on uniform meshes but also in adaptive refined meshes. It can be seen that the C-R method is efficient for computing real and complex eigenvalues as expected. In

addition, we discover, when the index of refraction $n(x)$ is real and Ω is the L-shaped domain or the square with a slit, the C-R element eigenvalues approximate the exact ones from above, and numerical results in [21, 43] show conforming finite element eigenvalues approximate the exact ones from below, thus we get the upper and lower bounds of eigenvalues.

It should be pointed out that the theoretical analysis and conclusions in this paper are also valid for the extension Crouzeix-Raviart element [38].

In this paper, regarding the basic theory of finite element methods, we refer to [8, 17, 26, 28, 45, 48].

Throughout this paper, C denotes a positive constant independent of h , which may not be the same constant in different places. For simplicity, we use the symbol $a \lesssim b$ to mean that $a \leq Cb$.

2 Preliminary

In this paper, we assume $\Omega \subset \mathbb{R}^d$ ($d = 2, 3$) is a polygonal ($d = 2$) or polyhedral ($d = 3$) domain, and assume $n \in L^\infty(\Omega)$. Let $H^\rho(\Omega)$ denote the Sobolev space on Ω with the norm $\|\cdot\|_{\rho,\Omega}$ (denoted by $\|\cdot\|_\rho$ for simplicity) and the seminorm $|\cdot|_{\rho,\Omega}$ (denoted by $|\cdot|_\rho$ for simplicity) and $H^0(\Omega) = L^2(\Omega)$, and let $H^\rho(\partial\Omega)$ denotes the Sobolev space on $\partial\Omega$ with the norm $\|\cdot\|_{\rho,\partial\Omega}$ and the seminorm $|\cdot|_{\rho,\partial\Omega}$.

Cakoni et al. [21] give the weak form of (1.1): Find $\lambda \in \mathbb{C}$, $u \in H^1(\Omega) \setminus \{0\}$, such that

$$a(u, v) = -\lambda \langle u, v \rangle, \quad \forall v \in H^1(\Omega), \quad (2.1)$$

where

$$a(u, v) = (\nabla u, \nabla v) - (k^2 n u, v), \quad (u, v) = \int_{\Omega} u \bar{v} dx, \quad \langle u, v \rangle = \int_{\partial\Omega} u \bar{v} ds.$$

The source problem associated with (2.1) is as follows: Find $\varphi \in H^1(\Omega)$ such that

$$a(\varphi, v) = \langle f, v \rangle, \quad \forall v \in H^1(\Omega). \quad (2.2)$$

The Neumann eigenvalue problem associated with $n(x)$ is to find $k^2 \in \mathbb{C}$ and a nontrivial u such that

$$\Delta u + k^2 n(x) u = 0 \quad \text{in } \Omega, \quad \frac{\partial u}{\partial \gamma} = 0 \quad \text{on } \partial\Omega. \quad (2.3)$$

In this paper, we always assume k^2 is not an interior Neumann eigenvalue of (2.3). Under this assumption, according to [21] the Neumann-to-Dirichlet map $T : L^2(\partial\Omega) \rightarrow L^2(\partial\Omega)$ can be defined as follows. Let $f \in L^2(\partial\Omega)$, define $A : L^2(\partial\Omega) \rightarrow H^1(\Omega)$ by

$$a(Af, v) = \langle f, v \rangle, \quad \forall v \in H^1(\Omega), \quad (2.4)$$

and $Tf = (Af)'$, where $'$ denotes the restriction to $\partial\Omega$. And (2.1) can be stated as the operator form:

$$Tw = \mu w. \quad (2.5)$$

(2.1) and (2.5) are equivalent, namely, if $(\mu, w) \in \mathbb{C} \times L^2(\partial\Omega)$ is an eigenpair of (2.5), then (λ, Aw) is an eigenpair of (2.1), $\lambda = -\mu^{-1}$; conversely, if (λ, u) is an eigenpair of (2.1), then (μ, u') is an eigenpair of (2.5), $\mu = -\lambda^{-1}$.

From [21] we know $T : L^2(\partial\Omega) \rightarrow L^2(\partial\Omega)$ is compact. If $n(x)$ is real, then T is also self-adjoint.

Consider the dual problem of (2.1): Find $\lambda^* \in \mathbb{C}$, $u^* \in H^1(\Omega) \setminus \{0\}$ such that

$$a(v, u^*) = -\overline{\lambda^*} \langle v, u^* \rangle, \quad \forall v \in H^1(\Omega). \quad (2.6)$$

The source problem associated with (2.6) is as follows: Find $\varphi^* \in H^1(\Omega)$ such that

$$a(v, \varphi^*) = \langle v, g \rangle, \quad \forall v \in H^1(\Omega), \quad (2.7)$$

Define the corresponding Neumann-to-Dirichlet operator operator $T^* : L^2(\partial\Omega) \rightarrow L^2(\partial\Omega)$ by

$$a(v, A^*g) = \langle v, g \rangle, \quad \forall v \in H^1(\Omega), \quad (2.8)$$

and $T^*g = (A^*g)'$. Then (2.6) has the equivalent operator form:

$$T^*u^* = -\lambda^{*-1}u^*. \quad (2.9)$$

It can be proved that T^* is the adjoint operator of T in the sense of inner product $\langle \cdot, \cdot \rangle$. In fact, from (2.4) and (2.8) we have

$$\langle Tf, g \rangle = a(Af, A^*g) = \langle f, A^*g \rangle = \langle f, T^*g \rangle, \quad \forall f, g \in L^2(\partial\Omega).$$

Note that since T^* is the adjoint operator of T , the primal and dual eigenvalues are connected via $\lambda = \overline{\lambda^*}$.

Let $\pi_h = \{\kappa\}$ be a regular d -simplex partition of Ω (see [28], pp. 131). We denote $h = \max_{\kappa \in \pi_h} h_\kappa$ where h_κ is the diameter of element κ . Let \mathcal{E}_h denote the set of all $(d-1)$ -faces of elements $\kappa \in \pi_h$. We split this set as follows: $\mathcal{E}_h = \mathcal{E}_h^i \cup \mathcal{E}_h^b$, with \mathcal{E}_h^i and \mathcal{E}_h^b being the sets of inner and boundary edges, respectively. Let S^h be the C-R element space defined on π_h :

$$S^h = \{v \in L^2(\Omega) : v|_\kappa \in P_1(\kappa), v \text{ is continuous at the barycenters of the } (d-1)\text{-faces of element } \kappa, \forall \kappa \in \pi_h\}.$$

The C-R element approximation of (2.1) is: Find $\lambda_h \in \mathbb{C}$, $u_h \in S^h \setminus \{0\}$, such that

$$a_h(u_h, v) = -\lambda_h \langle u_h, v \rangle, \quad \forall v \in S^h, \quad (2.10)$$

where $a_h(u_h, v) = \sum_{\kappa \in \pi_h} \int_{\kappa} (\nabla u_h \cdot \nabla \bar{v} - k^2 n(x) u_h \bar{v}) dx$.

Define $\|v\|_h = (\sum_{\kappa \in \pi_h} \|v\|_{1,\kappa}^2)^{\frac{1}{2}}$, $\|v\|_{1,\kappa}^2 = \int_{\kappa} (\sum_{i=1}^d |\frac{\partial v}{\partial x_i}|^2 + |v|^2) dx$. Evidently, $\|\cdot\|_h$ is the norm on S^h and it is easy to know that $a_h(\cdot, \cdot)$ is not uniformly S^h -elliptic.

The C-R element approximation of (2.2) is: Find $\varphi_h \in S^h$, such that

$$a_h(\varphi_h, v) = \langle f, v \rangle, \quad \forall v \in S^h. \quad (2.11)$$

Since k^2 is not an interior Neumann eigenvalue of (2.3), from spectral approximation theory [26] we know that when h is properly small k^2 is also not a C-R element eigenvalue for (2.3). So the discrete source problem (2.11) is uniquely solvable. Thus, we can define the discrete operator $A_h : L^2(\partial\Omega) \rightarrow S^h$, satisfying

$$a_h(A_h f, v) = \langle f, v \rangle, \quad \forall v \in S^h. \quad (2.12)$$

Let us denote by δS^h the function space defined on $\partial\Omega$, which are restriction of functions in S^h to $\partial\Omega$. Define the discrete operator $T_h : L^2(\partial\Omega) \rightarrow \delta S^h \subset L^2(\partial\Omega)$, satisfying $T_h f = (A_h f)'$. Then (2.10) has the equivalent operator form:

$$T_h w_h = \mu_h w_h, \quad (2.13)$$

namely, if $(\mu_h, w_h) \in \mathbb{C} \times L^2(\partial\Omega)$ is an eigenpair of (2.13), then $(\lambda_h, A_h w_h)$ is an eigenpair of (2.10), $\lambda_h = -\mu_h^{-1}$; conversely, if (λ_h, u_h) is an eigenpair of (2.10), then (μ_h, u_h') is an eigenpair of (2.13), $\mu_h = -\lambda_h^{-1}$.

The non-conforming finite element approximation of (2.6) is given by: Find $\lambda_h^* \in \mathbb{C}$, $u_h^* \in S^h \setminus \{0\}$ such that

$$a_h(v, u_h^*) = -\overline{\lambda_h^*} \langle v, u_h^* \rangle, \quad \forall v \in S^h. \quad (2.14)$$

The C-R element approximation of (2.7) is: Find $\varphi_h^* \in S^h$, such that

$$a_h(v, \varphi_h^*) = \langle v, g \rangle, \quad \forall v \in S^h. \quad (2.15)$$

Define the discrete operator $A_h^* : L^2(\partial\Omega) \rightarrow S^h$ satisfying

$$a_h(v, A_h^* g) = \langle v, g \rangle, \quad \forall v \in S^h, \quad (2.16)$$

and denote $T_h^* g = (A_h^* g)'$, then (2.14) has the following equivalent operator form

$$T_h^* u_h^* = -\lambda_h^{*-1} u_h^*. \quad (2.17)$$

It can be proved that T_h^* is the adjoint operator of T_h in the sense of inner product $\langle \cdot, \cdot \rangle$. Hence, the primal and dual eigenvalues are connected via $\lambda_h = \overline{\lambda_h^*}$.

We need the following regularity estimates which play an important role in

our theoretical analysis. Note that for $v \in H^{\frac{1}{2}}(\partial\Omega)$, $\langle f, v \rangle$ has a continuous extension, still denoted by $\langle f, v \rangle$, to $f \in H^{-\frac{1}{2}}(\partial\Omega)$.

Lemma 1 For any $f \in H^{-\frac{1}{2}}(\partial\Omega)$, let $\langle f, v \rangle$ be the dual product on $H^{-\frac{1}{2}}(\partial\Omega) \times H^{\frac{1}{2}}(\partial\Omega)$ in (2.2), then there exists a unique solution $\varphi \in H^1(\Omega)$ to (2.2) such that

$$\|\varphi\|_1 \lesssim \|f\|_{-\frac{1}{2}, \partial\Omega}. \quad (2.18)$$

Proof Since k^2 is not an interior Neumann eigenvalue of (2.3), there exists a unique solution $\varphi \in H^1(\Omega)$ to (2.2). Denote

$$b(u, v) = \int_{\Omega} \nabla u \cdot \overline{\nabla v} + n(x)u\bar{v}dx.$$

Referring to the proof of (14.11) in [28], we will show that, for any $v \in H^1(\Omega)$, there exists a constant $C(\Omega)$ such that

$$\|v\|_1 \leq C(\Omega)\sqrt{\text{Re}b(v, v)} \equiv C(\Omega)\left(\int_{\Omega} |\nabla v|^2 + n_1(x)|v|^2 dx\right)^{\frac{1}{2}}. \quad (2.19)$$

From (2.19), it is easy to verify that $\sqrt{\text{Re}b(v, v)}$ is a norm on $H^1(\Omega)$ that is equivalent to the norm $\|\cdot\|_1$.

If (2.19) is false, there exists a sequence $\{v_i\} \subset H^1(\Omega)$ such that

$$\|v_i\|_1 = 1 \text{ for all } i \geq 1, \quad \lim_{i \rightarrow \infty} \left(\int_{\Omega} |\nabla v_i|^2 + n_1(x)|v_i|^2 dx\right)^{\frac{1}{2}} = 0. \quad (2.20)$$

Since the $\{v_i\}$ is bounded in $H^1(\Omega)$, there exists a subsequence, again denoted $\{v_i\}$ for notational convenience, that converges in $L^2(\Omega)$. Since $\lim_{i \rightarrow \infty} \|v_i\|_1 = 0$, by (2.20), and since $H^1(\Omega)$ is complete, the sequence $\{v_i\}$ converges in $H^1(\Omega)$. The limit v of this sequence is such that

$$\|v\|_1 = \lim_{i \rightarrow \infty} \|v_i\|_1 = 0,$$

and thus, v is a constant. Using (2.20), we have

$$\left(\int_{\Omega} n_1(x)|v|^2 dx\right)^{\frac{1}{2}} = \lim_{i \rightarrow \infty} \left(\int_{\Omega} n_1(x)|v_i|^2 dx\right)^{\frac{1}{2}} = 0,$$

hence we conclude that $n_1(x) = 0$. But this contradicts the hypothesis $n_1(x) > 0$. It implies that (2.19) holds.

(2.3) can be rewritten as: Find $\tilde{\lambda} \in \mathbb{C}$, $u \in H^1(\Omega) \setminus \{0\}$ such that

$$b(u, v) = \tilde{\lambda}(nu, v), \quad \forall v \in H^1(\Omega), \quad (2.21)$$

Since k^2 is not an interior Neumann eigenvalue for (2.3), $k^2 + 1$ is not an eigenvalue of (2.21). Define the map $B : H^1(\Omega) \rightarrow H^1(\Omega)$ by

$$b(Bg, v) = (ng, v), \quad \forall v \in H^1(\Omega). \quad (2.22)$$

Then (2.21) has the operator form:

$$Bu = \tilde{\lambda}^{-1}u.$$

And B is compact, $\frac{1}{k^2+1}$ is not an eigenvalue of B . So $(B - \frac{1}{k^2+1}I)^{-1} : H^1(\Omega) \rightarrow H^1(\Omega)$ is bounded. Let $\psi \in H^1(\Omega)$ be the solution of the following equation:

$$b(\psi, v) = \langle f, v \rangle, \quad \forall v \in H^1(\Omega), \quad (2.23)$$

then we have $\|\psi\|_1 \lesssim \|f\|_{-\frac{1}{2}, \partial\Omega}$. From (2.22) we obtain

$$\begin{aligned} a(\varphi, v) &= b(\varphi, v) - (k^2 + 1)(n\varphi, v) \\ &= b(\varphi, v) - (k^2 + 1)b(B\varphi, v) = b((I - (k^2 + 1)B)\varphi, v), \end{aligned}$$

which together with (2.2) and (2.23) yields

$$b((I - (k^2 + 1)B)\varphi, v) = b(\psi, v), \quad \forall v \in H^1(\Omega).$$

Thus we have

$$\begin{aligned} \|\varphi\|_1 &= \left\| \left(\frac{1}{k^2+1}I - B \right)^{-1} \frac{1}{k^2+1} \psi \right\|_1 \\ &\leq \left\| \left(\frac{1}{k^2+1}I - B \right)^{-1} \right\|_1 \left\| \frac{1}{k^2+1} \psi \right\|_1 \lesssim \|f\|_{-\frac{1}{2}, \partial\Omega}, \end{aligned}$$

and the proof is completed. \square

Lemma 2 Assume that $\Omega \subset \mathbb{R}^2$ is a polygonal with ω being the largest interior angle, and φ is the solution of (2.2).

i) If $f \in L^2(\partial\Omega)$, then $\varphi \in H^{1+\frac{\pi}{2}}(\Omega)$ and

$$\|\varphi\|_{1+\frac{\pi}{2}} \leq C_\Omega \|f\|_{0, \partial\Omega}. \quad (2.24)$$

ii) If $f \in H^{\frac{1}{2}}(\partial\Omega)$, then $\varphi \in H^{1+r}(\Omega)$ satisfying

$$\|\varphi\|_{1+r} \leq C_\Omega \|f\|_{\frac{1}{2}, \partial\Omega}. \quad (2.25)$$

Here $r = 1$ when $\omega < \pi$, and $r < \frac{\pi}{\omega}$ when $\omega > \pi$, and C_Ω is a priori constant.

Proof Consider the auxiliary boundary value problem:

$$\Delta\varphi_1 + \varphi_1 = 0, \quad \frac{\partial\varphi_1}{\partial\gamma} = f, \quad (2.26)$$

$$\Delta\varphi_2 + \varphi_2 = -k^2n(x)(\varphi_1 + \varphi_2) + \varphi_1 + \varphi_2, \quad \frac{\partial\varphi_2}{\partial\gamma} = 0. \quad (2.27)$$

Let φ_1 and φ_2 be the solution of (2.26) and (2.27), respectively, then it is easy to see that $\varphi = \varphi_1 + \varphi_2$. Since $\Omega \subset \mathbb{R}^2$, from classical regularity results (see [32], or Proposition 4.1 in [3] and Proposition 4.4 in [11]) we have

$$\|\varphi_1\|_{1+(\frac{1}{2}+s)r} \lesssim \|f\|_{s,\partial\Omega}, \quad s = 0, \frac{1}{2},$$

and from classical regularity result for the Laplace problem with homogeneous Neumann boundary condition we have

$$\|\varphi_2\|_{1+r} \lesssim \| -k^2 n(x)(\varphi_1 + \varphi_2) + \varphi_1 + \varphi_2 \|_{0,\Omega}.$$

Thus we get

$$\|\varphi\|_{1+(\frac{1}{2}+s)r} \lesssim \|\varphi_1\|_{1+(\frac{1}{2}+s)r} + \|\varphi_2\|_{1+(\frac{1}{2}+s)r} \lesssim \|f\|_{s,\partial\Omega} + \|\varphi\|_0, \quad s = 0, \frac{1}{2}.$$

Substituting (2.18) into the above inequality we get (2.24) and (2.25). \square

Remark 1 (Regularity in \mathbb{R}^3) When $\Omega \subset \mathbb{R}^3$ is a polyhedral domain, regularity of the solution of the Neumann problem (2.26) has been discussed by many scholars. Referring Theorem 4 in [47], Remark 2.1 in [36], [39] and [32], and using the argument of Lemma 2 in this paper, we think the following regularity hypothesis $R(\Omega)$ is reasonable:

Hypothesis $R(\Omega)$ Let φ be the solution of (2.2) with $f \in L^2(\partial\Omega)$. When $\Omega \subset \mathbb{R}^3$, we have $\varphi \in H^{1+r3}(\Omega)$ for all $r3 \in (0, \frac{1}{2})$ and

$$\|\varphi\|_{1+r3} \leq C_\Omega \|f\|_{0,\partial\Omega}. \quad (2.28)$$

It is easy to know that Lemmas 1-2 and Remark 1 are also valid for the dual problem (2.7).

3 The consistency term and the extension of Strang lemma

Define $S^h + H^1(\Omega) = \{w_h + w : w_h \in S^h, w \in H^1(\Omega)\}$.

Let φ and φ^* be the solution of (2.2) and (2.7), respectively. Define the consistency terms: For any $v \in S^h + H^1(\Omega)$,

$$D_h(\varphi, v) = a_h(\varphi, v) - \langle f, v \rangle, \quad (3.1)$$

$$D_h^*(v, \varphi^*) = a_h(v, \varphi^*) - \langle v, g \rangle. \quad (3.2)$$

In order to analyze error estimates of the consistency terms, we need the following trace inequalities.

Lemma 3 For any $\kappa \in \pi_h$, the following trace inequalities hold:

$$\begin{aligned} \|w\|_{0,\partial\kappa} &\lesssim h_\kappa^{-\frac{1}{2}} \|w\|_{0,\kappa} + h_\kappa^{\frac{1}{2}} |w|_{1,\kappa}, \\ \|\nabla w\|_{0,\partial\kappa} &\lesssim h_\kappa^{-\frac{3}{2}} \|w\|_{0,\kappa} + h_\kappa^{-\frac{1}{2}} |w|_{1,\kappa} + h_\kappa^{s-\frac{1}{2}} |w|_{1+s,\kappa} \quad \left(\frac{1}{2} \leq s \leq 1\right). \end{aligned} \quad (3.3)$$

Proof The conclusion is followed by using the trace theorem on the reference element and the scaling argument (see, e.g., Lemma 2.2 in [52]). \square

The following Green's formula (see (2.7) in [20], (3.11) in [12] and Corollary 2.2 in [37]) will play a crucial role in our analysis:

$$\int_{\partial\kappa} (\nabla w \cdot \gamma) v ds = \int_{\kappa} \Delta w v dx + \int_{\kappa} \nabla w \cdot \nabla v dx \quad \forall \kappa \in \pi_h, \quad (3.4)$$

where $w \in H^{1+\epsilon}(\kappa)$ with $\Delta w \in L^2(\kappa)$ and $v \in H^{1-\epsilon}(\kappa)$ with $0 \leq \epsilon < \frac{1}{2}$.

Lemma 4 Suppose that $\varphi \in H^{1+t}(\Omega)$ ($0 < t < \frac{1}{2}$) is the solution of (2.2) and **Hypothesis** $R(\Omega)$ holds, then

$$\|\nabla \varphi \cdot \gamma\|_{t-\frac{1}{2}, \ell} \lesssim \|\varphi\|_{1+t, \kappa} \quad \forall \kappa \in \pi_h, \ell \in \partial\kappa. \quad (3.5)$$

Proof Inequality (3.5) is contained in the proof of Corollary 3.3 on page 1384 of [12], see also Lemma 2.1 in [20]. For the convenience of readers, we write the proof here.

For any $g \in H^{\frac{1}{2}-t}(\ell)$, it is proven by going to a reference element and using the inverse trace theorem (see page 387 in [40], page 1767 in [20]) that there exists a lifting w_g of g such that $w_g \in H^{1-t}(\kappa)$, $w_g|_{\ell} = g$, $w_g|_{\partial\kappa \setminus \ell} = 0$, and

$$\|\nabla w_g\|_{-t, \kappa} + h_{\kappa}^{t-1} \|w_g\|_{0, \kappa} \leq C \|g\|_{\frac{1}{2}-t, \ell}. \quad (3.6)$$

From Green's formula (3.4), (2.2), Cauchy-Schwarz inequality, the definition of the dual norm and (3.6) we deduce

$$\begin{aligned} \int_{\ell} \nabla \varphi \cdot \gamma \bar{g} ds &= \int_{\ell} \nabla \varphi \cdot \gamma \overline{w_g} ds = \int_{\kappa} \Delta \varphi \overline{w_g} dx + \int_{\kappa} \nabla \varphi \cdot \overline{\nabla w_g} dx \\ &= \int_{\kappa} -k^2 n \varphi \overline{w_g} dx + \int_{\kappa} \nabla \varphi \cdot \overline{\nabla w_g} dx \lesssim \| -k^2 n \varphi \|_{0, \kappa} \| \overline{w_g} \|_{0, \kappa} + \| \nabla \varphi \|_{t, \kappa} \| \overline{\nabla w_g} \|_{-t, \kappa} \\ &\lesssim \| -k^2 n \varphi \|_{0, \kappa} h_{\kappa}^{1-t} \| g \|_{\frac{1}{2}-t, \ell} + \| \nabla \varphi \|_{t, \kappa} \| g \|_{\frac{1}{2}-t, \ell} \lesssim \| \varphi \|_{1+t, \kappa} \| g \|_{\frac{1}{2}-t, \ell}, \end{aligned}$$

thus by the definition of the dual norm we obtain

$$\|\nabla \varphi \cdot \gamma\|_{t-\frac{1}{2}, \ell} = \sup_{g \in H^{\frac{1}{2}-t}(\ell)} \frac{|\int \nabla \varphi \cdot \gamma \bar{g} ds|}{\|g\|_{\frac{1}{2}-t, \ell}} \lesssim \|\varphi\|_{1+t, \kappa}.$$

This completes the proof of the lemma. \square

Based on the standard argument (see, e.g., [3, 41, 52]), the following consistency error estimates will be proved.

Theorem 1 Let φ and φ^* be the solution of (2.2) and (2.7), respectively, and suppose that $\varphi, \varphi^* \in H^{1+t}(\Omega)$ with $t \in [s, 1]$ and **Hypothesis** $R(\Omega)$ holds, then

$$|D_h(\varphi, v)| \lesssim h^t \|\varphi\|_{1+t} \|v\|_h, \quad \forall v \in S^h + H^1(\Omega), \quad (3.7)$$

$$|D_h^*(v, \varphi^*)| \lesssim h^t \|\varphi^*\|_{1+t} \|v\|_h, \quad \forall v \in S^h + H^1(\Omega), \quad (3.8)$$

where $s = \frac{r}{2}$ when $\Omega \subset \mathbb{R}^2$, $s = r/3$ when $\Omega \subset \mathbb{R}^3$.

Proof Let $[[\cdot]]$ denote the jump across an inner face $\ell \in \mathcal{E}_h^i$. By Green's formula (3.4) we deduce

$$\begin{aligned} D_h(\varphi, v) &= a_h(\varphi, v) - \langle f, v \rangle = \sum_{\kappa \in \pi_h} \int_{\kappa} (-\Delta \varphi - k^2 n(x) \varphi) \bar{v} dx \\ &+ \sum_{\kappa \in \pi_h} \int_{\partial \kappa} \frac{\partial \varphi}{\partial \gamma} \bar{v} ds - \int_{\partial \Omega} \frac{\partial \varphi}{\partial \gamma} \bar{v} ds = \sum_{\ell \in \mathcal{E}_h^i} \int_{\ell} \frac{\partial \varphi}{\partial \gamma} [[\bar{v}]] ds. \end{aligned} \quad (3.9)$$

Let ℓ be a $(d-1)$ -face of κ , define

$$P_\ell f = \frac{1}{|\ell|} \int_{\ell} f ds, \quad P_\kappa f = \frac{1}{|\kappa|} \int_{\kappa} f dx.$$

For $\ell \in \mathcal{E}_h^i$, suppose that $\kappa_1, \kappa_2 \in \pi_h$ such that $\kappa_1 \cap \kappa_2 = \ell$. Since $[[v]]$ is a linear function vanishing at the barycenters of ℓ , we have

$$\begin{aligned} \left| \int_{\ell} \frac{\partial \varphi}{\partial \gamma} [[v]] ds \right| &= \left| \int_{\ell} \left(\frac{\partial \varphi}{\partial \gamma} - P_\ell \left(\frac{\partial \varphi}{\partial \gamma} \right) \right) [[v]] ds \right| \\ &= \left| \int_{\ell} \left(\frac{\partial \varphi}{\partial \gamma} - P_\ell \left(\frac{\partial \varphi}{\partial \gamma} \right) \right) ([v] - P_\ell [[v]]) ds \right| \\ &= \left| \int_{\ell} \frac{\partial \varphi}{\partial \gamma} ([v] - P_\ell [[v]]) ds \right|. \end{aligned} \quad (3.10)$$

Then, when $t \in [\frac{1}{2}, 1]$, using Schwarz inequality we deduce

$$\begin{aligned} \left| \int_{\ell} \frac{\partial \varphi}{\partial \gamma} [[v]] ds \right| &\leq \sum_{i=1,2} \|\nabla \varphi \cdot \gamma - P_\ell(\nabla \varphi \cdot \gamma)\|_{0,\ell} \|v|_{\kappa_i} - P_\ell(v|_{\kappa_i})\|_{0,\ell} \\ &\leq \sum_{i=1,2} \|\nabla(\varphi - \varphi_I) \cdot \gamma\|_{0,\ell} \|v|_{\kappa_i} - P_{\kappa_i}(v|_{\kappa_i})\|_{0,\ell}, \end{aligned} \quad (3.11)$$

by Lemma 3 and the standard error estimates for L^2 -projection, we deduce

$$\|\nabla(\varphi - \varphi_I) \cdot \gamma\|_{0,\ell} \lesssim h^{t-\frac{1}{2}} \|\varphi\|_{1+t, \kappa_i}, \quad \|v|_{\kappa_i} - P_{\kappa_i}(v|_{\kappa_i})\|_{0,\ell} \lesssim h^{\frac{1}{2}} \|v\|_{1, \kappa_i}.$$

Substituting the above two estimates into (3.11), we obtain

$$\left| \int_{\ell} \frac{\partial \varphi}{\partial \gamma} [[v]] ds \right| \lesssim \sum_{i=1,2} h^t \|\varphi\|_{1+t, \kappa_i} \|v\|_{1, \kappa_i}, \quad (3.12)$$

and substituting (3.12) into (3.9) we conclude that (3.7) holds. When $t < \frac{1}{2}$, from (3.10) we deduce that

$$\left| \int_{\ell} \frac{\partial \varphi}{\partial \gamma} \overline{[[v]]} ds \right| \leq \|\nabla \varphi \cdot \gamma\|_{t-\frac{1}{2}, \ell} \|[[v]] - P_{\ell}[[v]]\|_{\frac{1}{2}-t, \ell}. \quad (3.13)$$

By using inverse estimate, Lemma 3 and the error estimate of L^2 -projection, we derive

$$\|[[v]] - P_{\ell}[[v]]\|_{\frac{1}{2}-t, \ell} \lesssim h_{\ell}^{t-\frac{1}{2}} \|[[v]] - P_{\ell}[[v]]\|_{0, \ell} \lesssim \sum_{i=1,2} h_{\kappa_i}^t \|v\|_{1, \kappa_i}.$$

Substituting the above estimate and (3.5) into (3.13), we obtain

$$\left| \int_{\ell} \frac{\partial \varphi}{\partial \gamma} \overline{[[v]]} ds \right| \lesssim \sum_{i=1,2} \|\varphi\|_{1+t, \kappa_i} h_{\kappa_i}^t \|v\|_{1, \kappa_i},$$

plugging the above inequality into (3.9) we also get (3.7).

Using the same argument as above, we can prove (3.8). \square

The C-R element approximation (2.11) of (2.2) does not satisfy the condition of Strang lemma, that is $a_h(\cdot, \cdot)$ is not uniformly S^h -elliptic. To overcome this difficulty, Inspired by the work in §5.7 in [17], next we use the standard duality technique to prove an extension version of the well-known Strang lemma.

First, we will use the standard duality argument to prove that $\|\varphi - \varphi_h\|_0$ is a quantity of higher order than $\|\varphi - \varphi_h\|_h$.

Introduce the auxiliary problem: Find $\psi \in H^1(\Omega)$, such that

$$a(v, \psi) = (v, g), \quad \forall v \in H^1(\Omega). \quad (3.14)$$

Let ψ be the solution of (3.14), then from elliptic regularity estimates for homogeneous Neumann boundary value problem we know that there exists $r_N > 0$, such that

$$\|\psi\|_{1+r_N} \lesssim \|g\|_0. \quad (3.15)$$

Let $\hat{E}_h(v, \psi) = a_h(v, \psi) - (v, g)$, then

$$|\hat{E}_h(v, \psi)| \lesssim h^{r_N} \|\psi\|_{1+r_N} \|v\|_h, \quad \forall v \in H^1(\Omega) + S^h. \quad (3.16)$$

Lemma 5 Let φ and φ_h be the solution of (2.2) and (2.11), respectively, and let φ^* and φ_h^* be the solution of (2.7) and (2.15), respectively, then

$$\|\varphi - \varphi_h\|_0 \lesssim h^{r_N} \|\varphi - \varphi_h\|_h, \quad (3.17)$$

$$\|\varphi^* - \varphi_h^*\|_0 \lesssim h^{r_N} \|\varphi^* - \varphi_h^*\|_h. \quad (3.18)$$

Proof By Riesz representation theorem we have

$$\|\varphi - \varphi_h\|_0 = \sup_{g \in L^2(\Omega), g \neq 0} \frac{|(\varphi - \varphi_h, g)|}{\|g\|_0}. \quad (3.19)$$

Let $\psi_I \in S^h$ be the C-R element interpolation function of ψ , the solution of (3.13), then according to the interpolation theory (see [28]) we have

$$\|\psi - \psi_I\|_h \lesssim h^{r_N} \|\psi\|_{1+r_N}. \quad (3.20)$$

By computing, we deduce

$$\begin{aligned} \langle f, \psi - \psi_I \rangle &= \langle f, \psi \rangle - \langle f, \psi_I \rangle = a(\varphi, \psi) - a_h(\varphi_h, \psi_I) \\ &= a(\varphi, \psi) - a_h(\varphi, \psi_I) + a_h(\varphi, \psi_I) - a_h(\varphi_h, \psi_I) \\ &= a_h(\varphi, \psi - \psi_I) + a_h(\varphi - \varphi_h, \psi_I) \\ &= a_h(\varphi, \psi - \psi_I) + a_h(\varphi - \varphi_h, \psi_I - \psi) + a_h(\varphi - \varphi_h, \psi), \end{aligned}$$

and

$$\begin{aligned} a_h(\varphi, \psi - \psi_I) + a_h(\varphi - \varphi_h, \psi) &= a_h(\varphi, \psi - \psi_I) - \langle f, \psi - \psi_I \rangle \\ &\quad + \langle f, \psi - \psi_I \rangle + a_h(\varphi - \varphi_h, \psi) - (\varphi - \varphi_h, g) + (\varphi - \varphi_h, g) \\ &= D_h(\varphi, \psi - \psi_I) + \hat{E}_h(\varphi - \varphi_h, \psi) + \langle f, \psi - \psi_I \rangle + (\varphi - \varphi_h, g). \end{aligned}$$

Combining the above two inequalities we get

$$(\varphi - \varphi_h, g) = a_h(\varphi - \varphi_h, \psi - \psi_I) - D_h(\varphi, \psi - \psi_I) - \hat{E}_h(\varphi - \varphi_h, \psi).$$

Substituting the above equality into (3.19) we get

$$\begin{aligned} \|\varphi - \varphi_h\|_0 &\leq \sup_{g \in L^2(\Omega), g \neq 0} \frac{|a_h(\varphi - \varphi_h, \psi - \psi_I)|}{\|g\|_0} \\ &\quad + \sup_{g \in L^2(\Omega), g \neq 0} \frac{|D_h(\varphi, \psi - \psi_I) + \hat{E}_h(\varphi - \varphi_h, \psi)|}{\|g\|_0}. \end{aligned} \quad (3.21)$$

Let I_h^C be the Lagrange interpolation operator, then $I_h^C \psi \in H^1(\Omega) \cap S^h$. According to (3.1) and (2.2) we have $D_h(\varphi, v) = 0, \forall v \in H^1(\Omega)$, and we deduce

$$\begin{aligned} |D_h(\varphi, \psi - \psi_I)| &= |D_h(\varphi, I_h^C \psi - \psi_I)| = |a_h(\varphi - \varphi_h, I_h^C \psi - \psi_I)| \\ &\lesssim \|\varphi - \varphi_h\|_h \|I_h^C \psi - \psi_I\|_h \lesssim h^{r_N} \|\psi\|_{1+r_N} \|\varphi - \varphi_h\|_h. \end{aligned} \quad (3.22)$$

Substituting (3.16), (3.20) and (3.22) into (3.21) we obtain the desired result (3.17).

Using the same argument as (3.17) we can prove (3.18). \square

Now we are ready to prove the following extension of Strang lemma.

Theorem 2 Let φ and φ_h be the solution of (2.2) and (2.11), respectively, then

$$\begin{aligned} \inf_{v \in S^h} \|\varphi - v\|_h + \sup_{v \in S^h \setminus \{0\}} \frac{|D_h(\varphi, v)|}{\|v\|_h} &\lesssim \|\varphi - \varphi_h\|_h \\ &\lesssim \inf_{v \in S^h} \|\varphi - v\|_h + \sup_{v \in S^h \setminus \{0\}} \frac{|D_h(\varphi, v)|}{\|v\|_h}. \end{aligned} \quad (3.23)$$

Let φ^* and φ_h^* be the solution of (2.7) and (2.15), respectively, then

$$\begin{aligned} & \inf_{v \in S^h} \|\varphi^* - v\|_h + \sup_{v \in S^h \setminus \{0\}} \frac{|D_h^*(v, \varphi^*)|}{\|v\|_h} \lesssim \|\varphi^* - \varphi_h^*\|_h \\ & \lesssim \inf_{v \in S^h} \|\varphi^* - v\|_h + \sup_{v \in S^h \setminus \{0\}} \frac{|D_h^*(v, \varphi^*)|}{\|v\|_h}. \end{aligned} \quad (3.24)$$

Proof Denote

$$\mathcal{A}_h(u, v) = a_h(u, v) + K(u, v), \quad \forall u, v \in S^h + H^1(\Omega),$$

where $K > \|k^2 n\|_{0, \infty}$. Then we know that \mathcal{A}_h satisfies the uniform S^h -ellipticity:

$$|\mathcal{A}_h(v, v)| \geq \min\{1, K - \|k^2 n\|_{0, \infty}\} \|v\|_h^2, \quad \forall v \in S^h.$$

And thus, for any $v \in S^h$,

$$\begin{aligned} \|\varphi_h - v\|_h^2 & \lesssim |\mathcal{A}_h(\varphi_h - v, \varphi_h - v)| \\ & = C|a_h(\varphi - v, \varphi_h - v) + \langle f, \varphi_h - v \rangle - a_h(\varphi, \varphi_h - v) + K\|\varphi_h - v\|_0^2|. \end{aligned}$$

When $\|\varphi_h - v\|_h \neq 0$, dividing both sides of the above by $\|\varphi_h - v\|_h$ we obtain

$$\begin{aligned} \|\varphi_h - v\|_h & \lesssim \|\varphi - v\|_h + \frac{|a_h(\varphi, \varphi_h - v) - \langle f, \varphi_h - v \rangle|}{\|\varphi_h - v\|_h} + K\|\varphi_h - v\|_0 \\ & \lesssim \|\varphi - v\|_h + \sup_{v \in S^h \setminus \{0\}} \frac{|D_h(\varphi, v)|}{\|v\|_h} + \|\varphi - \varphi_h\|_0. \end{aligned}$$

From the triangular inequality and (3.17) we get

$$\begin{aligned} \|\varphi - \varphi_h\|_h & \leq \|\varphi - v\|_h + \|v - \varphi_h\|_h \\ & \lesssim \|\varphi - v\|_h + \sup_{v \in S^h \setminus \{0\}} \frac{|D_h(\varphi, v)|}{\|v\|_h} + h^{r_N} \|\varphi - \varphi_h\|_h. \end{aligned}$$

The second inequality of (3.23) is proved.

From

$$|a_h(\varphi - \varphi_h, v)| \leq \|\varphi - \varphi_h\|_h \|v\|_h, \quad \forall v \in S^h,$$

we get

$$\|\varphi - \varphi_h\|_h \geq \frac{|a_h(\varphi, v) - a_h(\varphi_h, v)|}{\|v\|_h} = \frac{|D_h(\varphi, v)|}{\|v\|_h},$$

which together with $\|\varphi - \varphi_h\|_h \geq \inf_{v \in S^h} \|\varphi - v\|_h$ we obtain the first inequality of (3.23).

Similarly we can prove (3.24). The proof is completed. \square

Now we can state the error estimates of C-R element approximation for (2.2) and (2.7).

Theorem 3 Under the conditions of Theorems 1 and 2, we have

$$\|\varphi - \varphi_h\|_h \leq Ch^t \|\varphi\|_{1+t}, \quad (3.25)$$

$$\|\varphi - \varphi_h\|_{0,\partial\Omega} \leq Ch^{t+s} \|\varphi\|_{1+t} \quad (3.26)$$

$$\|\varphi^* - \varphi_h^*\|_h \leq Ch^t \|\varphi^*\|_{1+t}, \quad (3.27)$$

$$\|\varphi^* - \varphi_h^*\|_{0,\partial\Omega} \leq Ch^{t+s} \|\varphi^*\|_{1+t}. \quad (3.28)$$

Proof From Theorem 2, the interpolation error estimate and Theorem 1 we can obtain (3.25) and (3.27). From (2.2) and (2.11) we deduce

$$\begin{aligned} \langle f, \varphi^* - \varphi_h^* \rangle &= \langle f, \varphi^* \rangle - \langle f, \varphi_h^* \rangle \\ &= a(\varphi, \varphi^*) - a_h(\varphi_h, \varphi_h^*) = a(\varphi, \varphi^*) - a_h(\varphi_h, \varphi^*) + a_h(\varphi_h, \varphi^*) - a_h(\varphi_h, \varphi_h^*) \\ &= a_h(\varphi - \varphi_h, \varphi^*) + a_h(\varphi_h, \varphi^* - \varphi_h^*) \\ &= a_h(\varphi - \varphi_h, \varphi^*) + a_h(\varphi, \varphi^* - \varphi_h^*) - a_h(\varphi - \varphi_h, \varphi^* - \varphi_h^*), \end{aligned}$$

and from (3.1) and (3.2) we get

$$\begin{aligned} a_h(\varphi - \varphi_h, \varphi^*) + a_h(\varphi, \varphi^* - \varphi_h^*) &= a_h(\varphi - \varphi_h, \varphi^*) - \langle \varphi - \varphi_h, g \rangle \\ &\quad + \langle \varphi - \varphi_h, g \rangle + a_h(\varphi, \varphi^* - \varphi_h^*) - \langle f, \varphi^* - \varphi_h^* \rangle + \langle f, \varphi^* - \varphi_h^* \rangle \\ &= D_h^*(\varphi - \varphi_h, \varphi^*) + D_h(\varphi, \varphi^* - \varphi_h^*) + \langle f, \varphi^* - \varphi_h^* \rangle + \langle \varphi - \varphi_h, g \rangle. \end{aligned}$$

Combining the above two equalities we obtain

$$\langle \varphi - \varphi_h, g \rangle = a_h(\varphi - \varphi_h, \varphi^* - \varphi_h^*) - D_h^*(\varphi - \varphi_h, \varphi^*) - D_h(\varphi, \varphi^* - \varphi_h^*),$$

and we have by duality,

$$\begin{aligned} \|\varphi - \varphi_h\|_{0,\partial\Omega} &= \sup_{g \in L^2(\partial\Omega) \setminus \{0\}} \frac{\langle \varphi - \varphi_h, g \rangle}{\|g\|_{0,\partial\Omega}} \lesssim \|\varphi - \varphi_h\|_h \sup_{g \in L^2(\partial\Omega) \setminus \{0\}} \frac{\|\varphi^* - \varphi_h^*\|_h}{\|g\|_{0,\partial\Omega}} \\ &\quad + \sup_{g \in L^2(\partial\Omega) \setminus \{0\}} \frac{|D_h(\varphi, \varphi^* - \varphi_h^*)| + |D_h^*(\varphi - \varphi_h, \varphi^*)|}{\|g\|_{0,\partial\Omega}}. \end{aligned}$$

Substituting (3.7), (3.8), (3.25) and (3.27) into the above inequality, and using the regularity estimates (2.24) and (2.28), we get (3.26). Similarly we can prove (3.28). The proof is completed. \square

Remark 2 Consider the Neumann boundary problem: find $\varphi \in H^1(\Omega)$ such that

$$a(\varphi, v) = \langle f, v \rangle + (\zeta, v), \quad \forall v \in H^1(\Omega). \quad (3.29)$$

Let φ and φ_h be the exact solution and the C-R element solution of (3.29), respectively,

$$D_h(\varphi, v) = a_h(\varphi, v) - \langle f, v \rangle - (\zeta, v),$$

and let φ^* and φ_h^* be the exact solution and the C-R element solution of the dual problem of (3.29), respectively. Then the analysis and conclusions in this section are also valid for (3.29).

4 Error estimates of discrete Stekloff eigenvalues

In this paper we suppose that $\{\lambda_j\}$ and $\{\lambda_{j,h}\}$ are enumerations of the eigenvalues of (2.1) and (2.10) respectively according to the same sort rule, and let $\lambda = \lambda_m$ be the m th eigenvalue with the algebraic multiplicity q and the ascent α , $\lambda_m = \lambda_{m+1} = \dots, \lambda_{m+q-1}$. When $\|T_h - T\|_{0,\partial\Omega} \rightarrow 0$, q eigenvalues $\lambda_{m,h}, \dots, \lambda_{m+q-1,h}$ of (2.10) will converge to λ (see Lemma 5 on page 1091 of [34]).

Let $M(\lambda)$ be the space of generalized eigenvectors associated with λ and T , let $M_h(\lambda_{j,h})$ be the space of generalized eigenvectors associated with $\lambda_{j,h}$ and T_h , and let $M_h(\lambda) = \sum_{j=m}^{m+q-1} M_h(\lambda_{j,h})$. In view of the dual problem (2.6) and (2.14), the definitions of $M(\lambda^*)$, $M_h(\lambda_{j,h}^*)$ and $M_h(\lambda^*)$ are analogous to $M(\lambda)$, $M_h(\lambda_{j,h})$ and $M_h(\lambda)$.

Given two closed subspaces V and U , denote

$$\delta(V, U) = \sup_{\substack{u \in V \\ \|u\|_{0,\partial\Omega}=1}} \inf_{v \in U} \|u - v\|_{0,\partial\Omega}, \quad \hat{\delta}(V, U) = \max\{\delta(V, U), \delta(U, V)\}.$$

And denote $\hat{\lambda}_h = \frac{1}{q} \sum_{j=m}^{m+q-1} \lambda_{j,h}$. Thanks to [8], we get the following Theorem 4.

Theorem 4 Suppose $M(\lambda), M(\lambda^*) \subset H^{1+t}(\Omega)$ ($t \in [r, 1]$ for $\Omega \subset \mathbb{R}^2$, and $t \in [r3, 1]$ for $\Omega \subset \mathbb{R}^3$), and **Hypothesis** $R(\Omega)$ holds. Then

$$\hat{\delta}(M(\lambda), M_h(\lambda)) \lesssim h^{s+t}, \quad (4.1)$$

$$|\hat{\lambda}_h - \lambda| \lesssim h^{2t}, \quad (4.2)$$

$$|\lambda - \lambda_{j,h}| \lesssim h^{\frac{2t}{\alpha}}, \quad j = m, m+1, \dots, m+q-1; \quad (4.3)$$

suppose u_h is an eigenfunction corresponding to $\lambda_{j,h}$ ($j = m, m+1, \dots, m+q-1$), $\|u_h\|_{0,\partial\Omega} = 1$, then there exists an eigenfunction u corresponding to λ , such that

$$\|u_h - u\|_{0,\partial\Omega} \lesssim h^{(s+t)\frac{1}{\alpha}}, \quad (4.4)$$

$$\|u_h - u\|_h \lesssim h^t + h^{(s+t)\frac{1}{\alpha}}; \quad (4.5)$$

where $s = \frac{r}{2}$ when $\Omega \subset \mathbb{R}^2$, $s = r3$ when $\Omega \subset \mathbb{R}^3$.

Proof Note that $\|Tf - T_h f\|_{0,\partial\Omega} = \|Af - A_h f\|_{0,\partial\Omega} = \|\varphi - \varphi_h\|_{0,\partial\Omega}$, from (3.26) with $t = s$ we deduce

$$\begin{aligned} \|T - T_h\|_{0,\partial\Omega} &= \sup_{f \in L^2(\partial\Omega), \|f\|_{0,\partial\Omega}=1} \|Tf - T_h f\|_{0,\partial\Omega} \\ &\lesssim \sup_{f \in L^2(\partial\Omega), \|f\|_{0,\partial\Omega}=1} h^{2s} \|Af\|_{1+s} \lesssim h^{2s} \|f\|_{0,\partial\Omega} \lesssim h^{2s} \rightarrow 0 \quad (h \rightarrow 0) \end{aligned} \quad (4.6)$$

Thus from Theorem 7.1, Theorem 7.2 (inequality (7.12)), Theorem 7.3 and Theorem 7.4 in [8] we get

$$\hat{\delta}(M(\lambda), M_h(\lambda)) \lesssim \|(T - T_h)|_{M(\lambda)}\|_{0, \partial\Omega}, \quad (4.7)$$

$$\begin{aligned} |\lambda - \hat{\lambda}_h| &\lesssim \sum_{i,j=m}^{m+q-1} |\langle (T - T_h)\varphi_i, \varphi_j^* \rangle| \\ &\quad + \|(T - T_h)|_{M(\lambda)}\|_{0, \partial\Omega} \|(T^* - T_h^*)|_{M(\lambda^*)}\|_{0, \partial\Omega}, \end{aligned} \quad (4.8)$$

$$\begin{aligned} |\lambda - \lambda_h| &\lesssim \left\{ \sum_{i,j=m}^{m+q-1} |\langle (T - T_h)\varphi_i, \varphi_j^* \rangle| \right. \\ &\quad \left. + \|(T - T_h)|_{M(\lambda)}\|_{0, \partial\Omega} \|(T^* - T_h^*)|_{M(\lambda^*)}\|_{0, \partial\Omega} \right\}^{1/\alpha}, \end{aligned} \quad (4.9)$$

$$\|u_h - u\|_{0, \partial\Omega} \leq C \|(T_h - T)|_{M(\lambda)}\|_{0, \partial\Omega}^{\frac{1}{\alpha}}, \quad (4.10)$$

where $\varphi_m, \dots, \varphi_{m+q-1}$ are any basis for $M(\lambda)$ and $\varphi_m^*, \dots, \varphi_{m+q-1}^*$ are the dual basis in $M(\lambda^*)$.

From (3.26) with $t \in [r, 1]$ we obtain

$$\begin{aligned} \|(T - T_h)|_{M(\lambda)}\|_{0, \partial\Omega} &= \sup_{f \in M(\lambda), \|f\|_{0, \partial\Omega}=1} \|Tf - T_h f\|_{0, \partial\Omega} \\ &\lesssim h^{s+t} \sup_{f \in M(\lambda), \|f\|_{0, \partial\Omega}=1} \|Af\|_{1+t}. \end{aligned} \quad (4.11)$$

Similarly we have

$$\|(T^* - T_h^*)|_{M(\lambda^*)}\|_{0, \partial\Omega} \lesssim h^{s+t} \sup_{f \in M(\lambda^*), \|f\|_{0, \partial\Omega}=1} \|A^* f\|_{1+t}. \quad (4.12)$$

Substituting (4.11) into (4.7) and (4.10) we get (4.1) and (4.4), respectively.

The remainder is to prove (4.2), (4.3) and (4.5). An easy calculation show that

$$\begin{aligned} \langle (T - T_h)\varphi_i, \varphi_j^* \rangle &= \langle T\varphi_i, \varphi_j^* \rangle - \langle T_h\varphi_i, \varphi_j^* \rangle \\ &= a_h(A\varphi_i, A^*\varphi_j^*) - a_h(A_h\varphi_i, A_h^*\varphi_j^*) \\ &= a_h(A\varphi_i - A_h\varphi_i, A^*\varphi_j^*) + a_h(A_h\varphi_i, A^*\varphi_j^* - A_h^*\varphi_j^*) \\ &= a_h(A\varphi_i - A_h\varphi_i, A^*\varphi_j^*) + a_h(A\varphi_i, A^*\varphi_j^* - A_h^*\varphi_j^*) \\ &\quad - a_h(A\varphi_i - A_h\varphi_i, A^*\varphi_j^* - A_h^*\varphi_j^*). \end{aligned} \quad (4.13)$$

By (3.1) and (3.2) with $f = \varphi_i$, $\varphi = A\varphi_i$, $g = \varphi_j^*$ and $\varphi^* = A^*\varphi_j^*$, we obtain

$$\begin{aligned} a_h(A\varphi_i - A_h\varphi_i, A^*\varphi_j^*) &= D_h^*(A\varphi_i - A_h\varphi_i, A^*\varphi_j^*) + \langle T\varphi_i - T_h\varphi_i, \varphi_j^* \rangle, \\ a_h(A\varphi_i, A^*\varphi_j^* - A_h^*\varphi_j^*) &= D_h(A\varphi_i, A^*\varphi_j^* - A_h^*\varphi_j^*) + \langle \varphi_i, T^*\varphi_j^* - T_h^*\varphi_j^* \rangle \\ &= D_h(A\varphi_i, A^*\varphi_j^* - A_h^*\varphi_j^*) + \langle T\varphi_i - T_h\varphi_i, \varphi_j^* \rangle. \end{aligned}$$

Substituting the above two relations into (4.13) we get

$$\begin{aligned} \langle (T - T_h)\varphi_i, \varphi_j^* \rangle &= -D_h^*(A\varphi_i - A_h\varphi_i, A^*\varphi_j^*) - D_h(A\varphi_i, A^*\varphi_j^* - A_h^*\varphi_j^*) \\ &\quad + a_h(A\varphi_i - A_h\varphi_i, A^*\varphi_j^* - A_h^*\varphi_j^*), \end{aligned} \quad (4.14)$$

which together with (3.7), (3.8), (3.25) and (3.27) yields

$$| \langle (T - T_h)\varphi_i, \varphi_j^* \rangle | \lesssim h^{2t}. \quad (4.15)$$

Substituting (4.15), (4.11) and (4.12) into (4.8) and (4.9) we get (4.2) and (4.3), respectively.

From (2.1) and (2.4) we get

$$a(u, v) = a(A(-\lambda u), v), \quad \forall v \in H^1(\Omega),$$

noting that k^2 is not an eigenvalue of (2.3), we have $u = -\lambda Au$. Similarly, using (2.10) and (2.12) we can get $u_h = -\lambda_h A_h u_h$. Thus from (3.25), (2.18), (2.24), (2.28), (4.3) and (4.4) we deduce

$$\begin{aligned} \|u_h + \lambda A_h u\|_h &= \| -\lambda_h A_h u_h + \lambda A_h u\|_h = \|A_h(\lambda_h u_h - \lambda u)\|_h \\ &\leq \|(A - A_h)(\lambda u - \lambda_h u_h)\|_h + \|A(\lambda_h u_h - \lambda u)\|_h \\ &\lesssim h^s \|A(\lambda u - \lambda_h u_h)\|_{1+s} + \|\lambda_h u_h - \lambda u\|_{0,\partial\Omega} \lesssim \|\lambda_h u_h - \lambda u\|_{0,\partial\Omega} \lesssim h^{(t+s)\frac{1}{\alpha}}, \end{aligned}$$

and by the triangular inequality

$$\|u_h - u\|_h \leq \|u_h + \lambda A_h u\|_h + \|\lambda Au - \lambda A_h u\|_h \lesssim h^{(t+s)\frac{1}{\alpha}} + h^t,$$

i.e., (4.5) holds. The proof is completed. \square

5 Numerical experiments

Consider the problem (1.1) on the test domain $\Omega \subset \mathbb{R}^2$, where $\Omega = (-\frac{\sqrt{2}}{2}, \frac{\sqrt{2}}{2})^2$ is the square, or $\Omega = (-1, 1)^2 \setminus ([0, 1] \times (-1, 0])$ is an L-shaped domain with the largest inner angle $\omega = \frac{3}{2}\pi$, or $\Omega = (-\frac{\sqrt{2}}{2}, \frac{\sqrt{2}}{2})^2 \setminus \{0 \leq x \leq \frac{\sqrt{2}}{2}, y = 0\}$ is the square with a slit which the largest inner angle $\omega = 2\pi$, or Ω is the unit disk, and $k = 1$, $n(x) = 4$ or $n(x) = 4 + 4i$.

We use Matlab 2012a to solve (1.1) on a Lenovo ideaPad PC with 1.8GHZ CPU and 8GB RAM. Our program is compiled under the package of iFEM [27].

Referring to [21, 43], when $n = 4$ we sort eigenvalues in descending order, and when $n = 4 + 4i$ we arrange complex eigenvalues by their imaginary parts from large to small.

For the unit disk, the exact Stekloff eigenvalues are given by (5.3) in [43], and when $n = 4$ the largest six eigenvalues are

$$\lambda_1 = 5.151841, \quad \lambda_{2,3} = 0.223578, \quad \lambda_{4,5} = -1.269100, \quad \lambda_6 = -2.472703,$$

and when $n = 4 + 4i$ the four complex eigenvalues with the largest imaginary parts are

$$\lambda_1 = -0.320506 + 3.121689i, \quad \lambda_{2,3} = -0.136861 + 1.396737i, \quad \lambda_4 = -1.353076 + 0.791723i.$$

For the L-shaped and the slit domain, the reference eigenvalues of the exact eigenvalues are listed in Tables 11-12.

Table 1: The eigenvalues on the square: $n = 4$.

dof	$\lambda_{1,h}$	$\lambda_{2,h}$	$\lambda_{3,h}$	$\lambda_{4,h}$	$\lambda_{5,h}$	$\lambda_{6,h}$
3136	2.2018805	-0.2116751	-0.2116708	-0.9069429	-2.7589883	-2.7522381
12416	2.2023533	-0.2121076	-0.2121070	-0.9077740	-2.7664177	-2.7646187
49408	2.2024690	-0.2122160	-0.2122159	-0.9079851	-2.7683097	-2.7678463
197120	2.2024977	-0.2122431	-0.2122431	-0.9080383	-2.7687870	-2.7686695

Table 2: The eigenvalues on the L-shaped domain: $n = 4$.

dof	$\lambda_{1,h}$	$\lambda_{2,h}$	$\lambda_{3,h}$	$\lambda_{4,h}$	$\lambda_{5,h}$	$\lambda_{6,h}$
9344	2.5335485	0.8592520	0.1246281	-1.0845725	-1.0901869	-1.4147102
37120	2.5333019	0.8583814	0.1245509	-1.0851154	-1.0909141	-1.4163502
147968	2.5332364	0.8580275	0.1245311	-1.0852527	-1.0911151	-1.4167642
590848	2.5332194	0.8578847	0.1245261	-1.0852873	-1.0911726	-1.4168682

5.1 Numerical experiments on uniform meshes

We adopt a uniform mesh π_h for each domain. The numerical results are listed in Tables 1-8. The error curves of the C-R eigenvalues are showed in Figs. 1-4.

From Lemma 2, the regularity results, we know that for the square domain $2r = 2$, for the L-shaped domain $2r \approx \frac{4}{3}$, for the unit square with a slit $2r \approx 1$. From Fig. 1 we can see that the convergence order of $\lambda_{1,h}, \lambda_{2,h}, \dots, \lambda_{6,h}$ are approximately equal to 2 on the square domain; from Fig. 2 we can see that the convergence order of $\lambda_{2,h}$ is approximately equal to $\frac{4}{3} \approx 1.333333$ on the L-shaped domain, and the eigenfunction corresponding to λ_2 has lower smoothness than others; from Fig. 3 we can see that the convergence order of $\lambda_{2,h}$ is approximately equal to 1 on the slit domain, and the eigenfunction corresponding to λ_2 is also less smoother than others, which are coincide with the theoretical results. Although there is an effect of regional approximation for the computation on a disk, namely, replacing the disk Ω with a similar polygonal Ω^h , from Tables 7-8 and Fig. 4 we see that C-R element eigenvalues can approximate the exact ones.

For the square and the L-shaped domain, we also compare the numerical results in Tables 1-2, 5-6 with Tables 5.2-5.3, 5.5-5.6 in [43] and find that, with the increase of dof (or the decrease of mesh size h), the eigenvalues obtained by C-R

Table 3: The eigenvalues on the square with a slit: $n = 4$.

dof	$\lambda_{1,h}$	$\lambda_{2,h}$	$\lambda_{3,h}$	$\lambda_{4,h}$	$\lambda_{5,h}$	$\lambda_{6,h}$
12448	1.4848728	0.4698829	-0.1840366	-0.6898362	-1.8987837	-1.9264514
49472	1.4847611	0.4658257	-0.1841411	-0.6900139	-1.8995947	-1.9278655
197248	1.4847266	0.4637839	-0.1841672	-0.6900592	-1.8998016	-1.9283610
787712	1.4847163	0.4627589	-0.1841737	-0.6900708	-1.8998539	-1.9285538

Table 4: The eigenvalues on the unit disk: $n = 4$.

dof	$\lambda_{1,h}$	$\lambda_{2,h}$	$\lambda_{3,h}$	$\lambda_{4,h}$	$\lambda_{5,h}$	$\lambda_{6,h}$
128628	5.1514757	0.2235716	0.2235710	-1.2689792	-1.2689809	-2.4724133
201444	5.1516049	0.2235739	0.2235738	-1.2690218	-1.2690228	-2.4725174
359676	5.1517065	0.2235760	0.2235759	-1.2690553	-1.2690557	-2.4725980
809421	5.1517811	0.2235773	0.2235773	-1.2690803	-1.2690804	-2.4726559

Table 5: The eigenvalues on the square: $n = 4 + 4i$.

dof	$\lambda_{1,h}$	$\lambda_{2,h}$	$\lambda_{3,h}$	$\lambda_{4,h}$	$\lambda_{5,h}$	$\lambda_{6,h}$
3136	0.687353	-0.342514	-0.342525	-0.948908	-2.779702	-2.786716
	+2.494448i	+0.85089i	+0.850899i	+0.539844i	+0.53745i	+0.539647i
12416	0.686749	-0.342915	-0.342916	-0.949807	-2.792169	-2.794033
	+2.495075i	+0.850782i	+0.850784i	+0.540029i	+0.539839i	+0.540444i
49408	0.686601	-0.343014	-0.343014	-0.950034	-2.795417	-2.795897
	+2.495238i	+0.850755i	+0.850756i	+0.540079i	+0.540498i	+0.540656i
197120	0.686564	-0.343038	-0.343038	-0.950091	-2.796245	-2.796367
	+2.495280i	+0.850749i	+0.850749i	+0.540092i	+0.540671i	+0.540711i

Table 6: The eigenvalues on the L-shaped domain: $n = 4 + 4i$.

dof	$\lambda_{1,h}$	$\lambda_{2,h}$	$\lambda_{3,h}$	$\lambda_{4,h}$	$\lambda_{5,h}$	$\lambda_{6,h}$
9344	0.513857	0.398298	-0.076964	-1.438567	-1.654555	-2.513849
	+2.881404i	+1.459758i	+1.042587i	+0.803689i	+0.766423i	+0.570528i
37120	0.514176	0.397512	-0.077125	-1.440022	-1.656531	-2.516699
	+2.882086i	+1.459328i	+1.042656i	+0.804437i	+0.766548i	+0.571289i
147968	0.514259	0.397218	-0.077165	-1.440388	-1.657092	-2.517426
	+2.882263i	+1.459129i	+1.042672i	+0.80463i	+0.766548i	+0.571486i
590848	0.514280	0.397106	-0.077175	-1.440479	-1.657258	-2.517610
	+2.882308i	+1.459043i	+1.042677i	+0.804678i	+0.766534i	+0.571536i

Table 7: The eigenvalues on the square with a slit: $n = 4 + 4i$.

dof	$\lambda_{1,h}$	$\lambda_{2,h}$	$\lambda_{3,h}$	$\lambda_{4,h}$	$\lambda_{5,h}$	$\lambda_{6,h}$
12448	0.918974	0.299813	-0.262446	-0.741837	-2.615356	-2.840331
	+1.770802i	+1.003519i	+0.757437i	+0.608741i	+0.561764i	+0.493956i
49472	0.919206	0.296211	-0.262573	-0.742028	-2.618344	-2.845935
	+1.770795i	+1.001745i	+0.757447i	+0.608765i	+0.562409i	+0.493673i
197248	0.919276	0.294417	-0.262604	-0.742076	-2.619113	-2.847993
	+1.770791i	+1.000826i	+0.757449i	+0.608772i	+0.562579i	+0.493444i
787712	0.919297	0.293522	-0.262612	-0.742088	-2.619310	-2.848830
	+1.770789i	+1.000356i	+0.75745i	+0.608774i	+0.562623i	+0.493306i




Figure 1: *The error curves of the first six eigenvalues on the square (left: $n = 4$, right: $n = 4 + 4i$)*




Figure 2: *The error curves of the first six eigenvalues on the L-shaped domain (left: $n = 4$, right: $n = 4 + 4i$)*




Figure 3: *The error curves of the first six eigenvalues on the square with a slit (left: $n = 4$, right: $n = 4 + 4i$)*

Table 8: The eigenvalues on the unit disk: $n = 4 + 4i$.

dof	$\lambda_{1,h}$	$\lambda_{2,h}$	$\lambda_{3,h}$	$\lambda_{4,h}$
128628	-0.320420	-0.136864	-0.136865	-1.352964
	+3.124755i	+1.39673i	+1.39673i	+0.79174i
201444	-0.320451	-0.136863	-0.136863	-1.353004
	+3.124732i	+1.396733i	+1.396733i	+0.791734i
359676	-0.320475	-0.136862	-0.136862	-1.353035
	+3.124714i	+1.396735i	+1.396735i	+0.79173i
809421	-0.320492	-0.136862	-0.136862	-1.353058
	+3.124700i	+1.396736i	+1.396736i	+0.791726i

element and the conforming element are getting closer.

5.2 Numerical experiments on adaptive meshes

In practical finite element computations, it is desirable to carry out the computations in an adaptive fashion (see, e.g., [1, 9, 18, 48, 50] and references cited therein). For the C-R element approximation of Steklov eigenvalue problem, the a posteriori error estimates has been developed by [46]. Referring to [46] in this subsection we give the a posteriori error estimators by formal deduction, and implement adaptive computation for (1.1).

Let $\ell \in \mathcal{E}_h^i$ shared by elements κ_1 and κ_2 , i.e., $\ell = \partial\kappa_1 \cap \partial\kappa_2$. We choose a unit normal vector γ_ℓ , pointing outwards κ_2 , and we set the jumps of the normal derivatives of v_h across ℓ as

$$[[\nabla v_h]]_\gamma = \nabla v_h|_{\kappa_2} \cdot \gamma_\ell - \nabla v_h|_{\kappa_1} \cdot \gamma_\ell.$$

Denote $\gamma_\ell = (\gamma_{\ell 1}, \gamma_{\ell 2})$, then the tangent $t_\ell = (-\gamma_{\ell 2}, \gamma_{\ell 1})$ on ℓ , and we write the jumps of the tangential derivatives of v_h across ℓ as

$$[[\nabla v_h]]_t = \nabla v_h|_{\kappa_2} \cdot t_\ell - \nabla v_h|_{\kappa_1} \cdot t_\ell.$$

Notice that these values are independent of the chosen direction of the normal vector γ_ℓ .

Now we define the a posteriori error indicators $\eta_\kappa(u_h)$ on κ and $\eta(u_h)$ on Ω for the primal eigenfunction u_h :

For each $\ell \in \mathcal{E}$, let

$$J_{\ell,t}(u_h) = \begin{cases} [[\nabla u_h]]_t, & \text{if } \ell \in \mathcal{E}^i, \\ 0 & \text{if } \ell \in \mathcal{E}^b, \end{cases} \quad J_{\ell,\gamma}(u_h) = \begin{cases} [[\nabla u_h]]_\gamma, & \text{if } \ell \in \mathcal{E}^i, \\ 2(\nabla u_h \cdot \gamma_\ell - \lambda_h u_h)|_\ell & \text{if } \ell \in \mathcal{E}^b, \end{cases}$$

and let

$$\eta_\kappa(u_h)^2 = |\kappa| \|k^2 n u_h\|_{0,\kappa}^2 + \frac{1}{2} \sum_{\ell \in \partial\kappa} |\ell| \|J_{\ell,\gamma}(u_h)\|_{0,\ell}^2 + \frac{1}{2} \sum_{\ell \in \partial\kappa} |\ell| \|J_{\ell,t}(u_h)\|_{0,\ell}^2,$$

$$\eta^2(u_h) = \sum_{\kappa \in \pi_h} \eta_\kappa(u_h)^2.$$

Table 9: The second eigenvalues on adaptive meshes on the L-shaped domain.

l	dof	$\lambda_{2,h}(n=4)$	l	dof	$\lambda_{2,h}(n=4+4i)$
1	9344	0.859246	1	9344	0.398302+1.459749i
2	10022	0.858839	2	9494	0.398303+1.459755i
25	202370	0.857844	37	216512	0.397113+1.459061i
26	225490	0.857841	38	242302	0.397112+1.459059i
27	249481	0.857838	39	286483	0.397108+1.459048i
28	276807	0.857832	40	310425	0.397092+1.459023i
29	331662	0.857821	41	340309	0.397092+1.459023i
30	387329	0.857817	42	391833	0.397084+1.459016i

Table 10: The second eigenvalues on adaptive meshes on the square with a slit.

l	dof	$\lambda_{2,h}(n=4)$	l	dof	$\lambda_{2,h}(n=4+4i)$
1	12448	0.469884	1	12448	0.299812+1.003523i
2	12472	0.467948	2	12607	0.298475+1.002875i
25	170854	0.461819	60	241012	0.292782+0.999958i
26	192640	0.461803	61	250930	0.292782+0.999958i
27	222566	0.461788	62	260992	0.292761+0.999943i
28	261309	0.461786	63	295455	0.292741+0.999931i
29	298511	0.461783	64	311350	0.292738+0.999931i
30	335598	0.461779	65	338930	0.292725+0.999926i

Similarly, we define the a posteriori error indicators $\eta_\kappa(u_h^*)$ on κ and $\eta(u_h^*)$ on Ω for the dual eigenfunction u_h^* .

We use $\sum_{\kappa \in \pi_h} (\eta_\kappa^2(u_h) + \eta_\kappa^2(u_h^*))$ as the a posteriori error indicator of λ_h . Using the indicator and consulting the existing standard adaptive algorithms (see, e.g., [27, 31, 44]), we solve (1.1). From Figs. 2-3 we find that the eigenfunction associated with λ_2 is singular, so in our numerical experiments we compute the approximation of the second eigenvalue λ_2 , and the numerical results on the L-shaped domain and the slit domain are listed in Table 9 and Table 10, respectively.

We show the curves of error and the a posteriori error estimators obtained by adaptive computing for the eigenvalue $\lambda_{2,h}$ in Figs. 5-6. It can be seen from them that the error curves and the error estimators' curves are both basically parallel to the line with slope -1, which indicate that the a posteriori error estimators of numerical eigenvalues are reliable and efficient and $\lambda_{2,h}$ achieves the convergence rate $O(h^2)$.

From tables and figures we also see that under the same dof, the accuracy of approximate eigenvalues computed on adaptive meshes is far higher than that of approximate eigenvalues computed on uniform meshes.

Remark 3(The lower/upper bound of the Stekloff eigenvalues).

We find in Tables 1-4 and 9-10 that when the index of refraction $n(x)$ is




Figure 4: The error curves of the eigenvalues for the unit disk on uniform meshes (left: $n = 4$, right: $n = 4 + 4i$)




Figure 5: The error curves of the second eigenvalues on the L-shaped domain (left: $n = 4$, right: $n = 4 + 4i$)




Figure 6: The error curves of the second eigenvalues on the square with a slit (left: $n = 4$, right: $n = 4 + 4i$)

Table 11: The reference eigenvalues $\lambda_j(L)$ on the L-shaped domain and $\lambda_j(\text{Slit})$ on the square with a slit: $n = 4$.

j	$\lambda_{j,h}$	$\lambda_{j,h}^C$	$\lambda_j(L)$	$\lambda_{j,h}$	$\lambda_{j,h}^C$	$\lambda_j(\text{Slit})$
1	2.533219	2.533209	2.533214	1.484716	1.484710	1.484713
2	0.8578847	0.8577495	0.8578171	0.4627589	0.4612150	0.4619870
3	0.1245261	0.1245229	0.1245245	-0.1841737	-0.1841765	-0.1841751
4	-1.085287	-1.085303	-1.085295	-0.6900708	-0.6900769	-0.6900738
5	-1.091173	-1.091207	-1.091190	-1.899854	-1.899878	-1.899866
6	-1.416868	-1.416912	-1.416890	-1.928554	-1.928784	-1.928669

Table 12: The reference eigenvalues $\lambda_j(L)$ on the L-shaped domain and $\lambda_j(\text{Slit})$ on the square with a slit: $n = 4 + 4i$.

j	$\lambda_{j,h}$	$\lambda_{j,h}^C$	$\lambda_j(L)$	$\lambda_{j,h}$	$\lambda_{j,h}^C$	$\lambda_j(\text{Slit})$
1	0.5142799	0.5143106	0.5142952	0.9192965	0.9193164	0.9193065
	+2.882308i	+2.882334i	+2.882321i	+1.770789i	+1.770782i	+1.770786i
2	0.3971057	0.3969716	0.3970387	0.2935223	0.2917372	0.2926298
	+1.459043i	+1.458911i	+1.458977i	+1.000356i	+0.9993946i	+0.9998754i
3	-0.0771754	-0.0771792	-0.0771773	-0.2626120	-0.2626151	-0.2626135
	+1.042677i	+1.042673i	+1.042675i	+0.7574501i	+0.7574481i	+0.7574491i
4	-1.440479	-1.440535	-1.440507	-0.7420884	-0.7420981	-0.7420933
	+0.8046784i	+0.8047093i	+0.8046939i	+0.6087744i	+0.6087755i	+0.6087749i
5	-1.657258	-1.657409	-1.657333	-2.619310	-2.619442	-2.619376
	+0.7665341i	+0.7664933i	+0.7665137i	+0.5626232i	+0.5626581i	+0.5626407i
6	-2.517610	-2.517765	-2.517687	-2.848830	-2.850239	-2.849534
	+0.5715365i	+0.5715597i	+0.5715481i	+0.4933060i	+0.4929997i	+0.4931528i

real, all series of eigenvalues computed by C-R element show the tendency to decrease as the increase of dof except the first one λ_1 on the square and $\lambda_1 \sim \lambda_3$ on the unit disk. Note that the numerical results in [21, 43] indicate that the conforming finite element eigenvalues approximate the exact ones from below when $n(x)$ is real. So we also use the P1 conforming element to compute, and obtain reference values of the exact eigenvalues by averaging the P1 conforming eigenvalues $\lambda_{j,h}^C$ and the C-R element eigenvalues $\lambda_{j,h}$. We list them in Tables 11-12. The property of monotone convergence of the conforming finite element eigenvalues is easy to prove. However, the property of monotone convergence of the C-R nonconforming finite element eigenvalues is a non-trivial result. In 2014, Carstensen and Gedicke [25] prove rigorously the monotonicity for the classical Laplace eigenvalue problem. For the Stekloff eigenvalue problem considered in this paper, it is meaningful to study the monotone convergence of C-R element eigenvalues which is our next goal.

Acknowledgments This work is supported by the National Natural Science Foundation of China (Grant Nos.11561014,11761022).

References

- [1] Ainsworth, M., Oden, J.T.: A posteriori error estimates in the finite element analysis. Wiley-Inter science, New York (2011)
- [2] Ainsworth, M.: Robust a posteriori error estimation for nonconforming finite element approximation. *SIAM J. Numer. Anal.* 42, 2320-2341 (2005)
- [3] Alonso, A., Russo, A.D.: Spectral approximation of variationally-posed eigenvalue problems by nonconforming methods. *J. Comput. Appl. Math.* 223, 177-197 (2009)
- [4] Andreev, A.B., Todorov, T.D.: Isoparametric finite element approximation of a Steklov eigenvalue problem. *IMA. J. Numer. Anal.* 24, 309-322 (2004)
- [5] Armentano, M.G.: The effect of reduced integration in the Steklov eigenvalue problem. *Math. Mod. and Numer. Anal. (M^2AN)* 38, 27-36 (2004)
- [6] Armentano, M.G., Padra, C.: A posteriori error estimates for the Steklov eigenvalue problem. *Appl. Numer. Math.* 58, 593-601 (2008)
- [7] Armentano, M.G., Duran, R.G.: Asymptotic lower bounds for eigenvalues by nonconforming finite element methods. *Electron. Trans. Numer. Anal.* 17, 92-101 (2004)
- [8] Babuska, I., Osborn, J.E.: Eigenvalue Problems. in: P. G. Ciarlet, J. L. Lions(Eds), *Finite Element Methods (Part I)*, pp. 641-787, in: *Handbook of Numerical Analysis*, Vol. 2, Elsevier Science Publishers, North-Holand (1991)
- [9] Babuska, I., Rheinboldt, W.C.: Error estimates for adaptive finite element computations. *SIAM J. Numer. Anal.* 15, 736-754 (1978)
- [10] Bergman, S., Schiffer, M.: *Kernel Functions and Elliptic Differential Equations in Mathematical Physics*. Academic Press, New York (1953)
- [11] Bermudez, A., Rodriguez, R., Santamarina, D.: A finite element solution of an added mass formulation for coupled fluid-solid vibrations. *Numer. Math.* 87, 201-227 (2000)
- [12] Bernardi, C., Hecht, F.: Error indicators for the mortar finite element discretization of Laplace equation. *Math. Comp.* 71(240), 1371-1403 (2001)
- [13] Bi, H., Yang, Y.: A two-grid method of the non-conforming Crouzeix-Raviart element for the Steklov eigenvalue problem. *Appl. Math. Comput.* 217, 9669-9678 (2011)
- [14] Boffi, D.: Finite element approximation of eigenvalue problems. *Acta Numerica*, 1-120 (2010)

- [15] Bramble, J.H., Osborn, J.E.: Approximation of Steklov eigenvalues of non-selfadjoint second order elliptic operators. in: A. K. Aziz, (Ed.), Math.Foundations of the Finite Element Method with Applications to PDE, PP.387-408, Academic, New York (1972)
- [16] Brenner, S.C., Sung, L.Y.: Linear finite element methods for planar linear elasticity. Math. Comp. 59, 321-338 (1992)
- [17] Brenner, S.C., Scott, L.R.: The Mathematical Theory of Finite Element Methods. 2nd ed.. Springer-Verlag, New york (2002)
- [18] Brenner, S.C.: C^0 interior penalty methods. In Frontiers in Numerical Analysis-Durham 2010, Lecture Notes in Computational Science and Engineering 85, pp. 79-147, Springer-Verlag (2012)
- [19] Bucur, D., Ionescu, I.R.: Asymptotic analysis and scaling of friction parameters. Z. Angew. Math. Phys. (ZAMP) 57, 1042-1056 (2006)
- [20] Cai, Z., Ye, X., Zhang, S.: Discontinuous Galerkin finite element methods for interface problems: a priori and a posteriori error estimations. SIAM J. Numer. Anal. 49, 1761-1787 (2011)
- [21] Cakoni, F., Colton, D., Meng, S., Monk, P.: Stekloff eigenvalues in inverse scattering. SIAM J. Appl. Math. 76(4), 1737-1763 (2016)
- [22] Cao, L., Zhang, L., Allegretto, W., Lin, Y.: Multiscale asymptotic method for Steklov eigenvalue equations in composite media. SIAM J. Numer. Anal. 51, 273-296 (2013)
- [23] Carstensen, C., Hu, J., Orlando, A.: Framework for the a posteriori error analysis of nonconforming finite elements. SIAM J. Numer. Anal. 45, 68-82 (2007)
- [24] Carstensen, C., Hoppe, R.H.W.: Convergence analysis of an adaptive non-conforming finite element method. Numer. Math. 103, 251-266 (2006)
- [25] Carstensen, C., Gedicke, J.: Guaranteed lower bounds for eigenvalues. Math. Comp. 83, 2605-2629 (2014)
- [26] Chatelin, F.: Spectral Approximations of Linear Operators. Academic Press, New York, (1983)
- [27] Chen, L.: iFEM: an innovative finite element methods package in MATLAB. Technical Report, University of California at Irvine (2009)
- [28] Ciarlet, P.G.: Basic error estimates for elliptic problems. in: P. G. Ciarlet, J. L. Lions(Eds), Finite Element Methods (Part I), pp.21-343, in: Handbook of Numerical Analysis, Vol. 2, Elsevier Science Publishers, North-Holland (1991)

- [29] Conca, C., Planchard, J., Vanninathanm, M.: Fluid and Periodic Structures. John Wiley & Sons, New York (1995)
- [30] Crouzeix, M., Raviart, P.A.: Conforming and nonconforming finite element methods for solving the stationary stokes equations. RAIRO. Anal. Numer. 3, 33-75 (1973)
- [31] Dai, X., Xu, J., Zhou, A.: Convergence and optimal complexity of adaptive finite element eigenvalue computations. Numer. Math. 110, 313-355 (2008)
- [32] Dauge, M.: Elliptic boundary value problems on corner domains: smoothness and asymptotics of solutions. in: Lecture Notes in Mathematics, vol.1341. Springer, Berlin (1988)
- [33] Dari, E., Durán, R., Padra, C., Vampa, V.: A posteriori error estimators for noconforming finite element methods. RAIRO Model. Math. Anal. Numer. 30, 385-400 (1996)
- [34] Dunford, N., Schwartz, J. T.: Linear Operators, Vol.2: Spectral Theory, Selfadjoint Operators in Hilbert Space. Interscience, New York, (1963)
- [35] Falk, R.S.: Nonconforming finite element methods for the equations of linear elasticity. Math. Comp. 57, 529-550 (1991)
- [36] Garau, E.M., Morin, P.: Convergence and quasi-optimality of adaptive FEM for Steklov eigenvalue problems. IMA J. Numer. Anal. 31(3), 914-946 (2011)
- [37] Girault, V., Raviart, P.A.: Finite Element Approximation of the Navier-Stokes Equations. Lecture Notes in Mathematics 749, Springer-Verlag, Berlin Heidelberg, New York, (1981)
- [38] Hu, J., Huang, Y., Lin, Q.: The lower bounds for eigenvalues of elliptic operators by Nonconforming finite element methods. J. Sci Comput. 61, 196-221 (2014)
- [39] Jerison, D.S., Kenig, C.E.: The Neumann problem on Lipschitz domains, Bull. Amer. Math. Soc. 4, 203-207 (1981)
- [40] Kufner, A., John, O., Fučík, S.: Function Spaces, Academia Publishing House, Prague (1977)
- [41] Li, Q., Lin, Q., Xie, H.: Nonconforming finite element approximations of the Steklov eigenvalue problems and its lower bound approximations. Appl. Math. 58, 129-151 (2013)
- [42] Li, M., Lin, Q., Zhang, S.: Extrapolation and superconvergence of the Steklov eigenvalue problems. Adv. Comput. Math. 33, 25-44 (2010)

- [43] Liu, J., Sun, J., Turner, T.: Spectral indicator method for a non-selfadjoint Steklov eigenvalue problem. ar Xiv: 1804.02582V1 [math. NA] 7 Apr (2018)
- [44] Morin, P., Nochetto, R.H., Siebert, K.: Convergence of adaptive finite element methods. *SIAM Rev.* 44, 631-658 (2002)
- [45] Oden, J.T., Reddy, J.N.: An Introduction to the Mathematical Theory of Finite Elements. Courier Dover Publications, New York (2012)
- [46] Russo, A.D., Alonso, A.E.: A posteriori error estimates for nonconforming approximations of Steklov eigenvalue problems. *Comput. Math. Appl.* 62(11), 4100-4117 (2011)
- [47] Savaré, G.: Regularity results for elliptic equations in Lipschitz domains. *J. Funct. Anal.* 152, 176-201 (1998)
- [48] Shi, Z., Wang, M.: Finite Element Methods. Science Press, Beijing (2013)
- [49] Strang, G., Fix, G.J.: An analysis of the finite element method. Prentice-Hall, New York (1973)
- [50] Verfürth, R.: A review of a posteriori error estimates and adaptive mesh-refinement techniques. Wiley-Teubner, New York (1996)
- [51] Xie, H.: A type of multilevel method for the Steklov eigenvalue problem. *IMA J. Numer. Anal.* 34, 592-608 (2014)
- [52] Yang, Y., Li, Q., Li, S.: Nonconforming finite element approximations of the Steklov eigenvalue problem. *Appl. Numer. Math.* 59, 2388-2401 (2009)