

**ADDENDUM TO THE PAPER
“ON THE SETS OF n POINTS FORMING $n + 1$ DIRECTIONS”**

CÉDRIC PILATTE

In the paper “On the sets of n points forming $n + 1$ directions” [5] attached to this addendum, it is proven that every finite set R of $n \geq 7$ points in general position in the plane that determines $m = n + 1$ distinct directions is a subset of the vertices of an affinely-regular polygon.¹

This is a special case of a conjecture of Jamison [2], who conjectured that the above result still holds for $m \leq 2n - O(1)$ directions (it is easy to see that $m \geq n$ for any set R in general position). Jamison proved the case $m = n$. The case $m = n$ was independently considered by Wade and Chu, who obtained a slightly weaker result [7, Theorem 2].

Recently, Makhul and Pinchasi observed that Jamison’s conjecture is false for $m > \frac{3}{2}n$ and Pinchasi announced without proof the resolution of Jamison’s conjecture “when $m - n$ is in the order of magnitude of $O(\sqrt{n})$ ” with Alexandr Polyanskii [4].

In this addendum, we show that, for $m \leq n + c\sqrt[6]{n}$, Jamison’s conjecture follows from a powerful structure theorem of Green and Tao on the sets of points defining few ordinary lines [1] (for some absolute constant $c > 0$). This significantly improves the case $m = n + 1$ proved in our paper [5].

Notation. We denote by L_∞ the line at infinity inside the real projective plane \mathbb{RP}^2 . If $x \neq y$ are two points in \mathbb{RP}^2 , we write xy for the line passing through x and y .

Definition. Let P be a finite set of points in \mathbb{RP}^2 . A projective line is an ordinary line for P if it contains exactly two points of P .

In [1, Proposition 5.3], Green and Tao proved the following “intermediate structure theorem”.

Theorem 1 (Green-Tao). *Suppose that P is a finite set of p points in the projective plane that spans Kp ordinary lines for some $K \geq 1$. Let $K' = 2^{88}K^6$. Then one of the following alternatives holds:*

- (1) P is contained in the union of an irreducible cubic γ and an additional K' points.
- (2) there is an irreducible conic σ containing between $p/2 - K'$ and $p/2 + K'$ points of P ; the other points of P are contained in a union of K' lines.
- (3) P is contained in the union of K' lines.

In fact, they give three structure theorems with increasingly strong conclusions, but the “full structure theorem” has weaker explicit constants and we will therefore not use it.

Corollary 2. *There exist absolute constants $c, C > 0$ such that the following holds. Every finite set R of $n \geq C$ points in \mathbb{R}^2 that determines $m \leq n + cn^{1/6}$ distinct directions is a subset of the vertices of an affinely regular m -gon.*

Proof of Corollary 2. Let R_∞ be the set of points at infinity corresponding to the directions of R , i.e.

$$R_\infty = \{xy \cap L_\infty \mid x, y \in R, x \neq y\}.$$

Let $P = R \sqcup R_\infty$. Thus $|R| = n$, $|R_\infty| = m$ and $|P| = n + m$.

Date: January 2021.

¹The image of a regular polygon by an invertible affine transformation.

Let x and y be two distinct points of P . If $x, y \in R$, the line xy passes through a point of R_∞ , while if $x, y \in R_\infty$, $xy = L_\infty$ passes through all the points of R_∞ . In both cases, xy is not an ordinary line for P . Therefore, the set of ordinary lines spanned by P is

$$\{xy \mid (x, y) \in R \times R_\infty \text{ and } xy \cap R = \{x\}\}.$$

Since R is in general position, the number of ordinary lines determined by P can thus be computed as

$$|R| \cdot |R_\infty| - 2 \cdot \binom{|R|}{2} = n(m - n + 1).$$

Hence, we can apply Theorem 1 with the set $P = R \sqcup R_\infty$ and $K = m - n + 1$. Let ε be a small positive constant ($\varepsilon = 1/10$ suffices). The constants c and C may be chosen such that $K' \leq \varepsilon n$ and $n \leq m \leq (1 + \varepsilon)n$.

- Case (1) is impossible. Indeed, the irreducible cubic γ cannot contain more than three points of R_∞ , since the points of R_∞ are collinear.
- Case (3) is also impossible, because at least $n/2$ lines are required to cover R , since R is in general position.

So, Case (2) occurs and there exists an irreducible conic σ that contains at least

$$\frac{1}{2}(n + m) - K' \geq (1 - \varepsilon)n$$

points of P . Partition R as $R = R_\sigma \sqcup R_{-\sigma}$, where $R_\sigma := R \cap \sigma$. We have $|R_\sigma| \geq |P \cap \sigma| - 2 \geq (1 - 2\varepsilon)n$ since at most two points of σ lie on the line at infinity.

Let $\bar{\sigma} = \sigma \cap \mathbb{R}^2$ be the conic obtained by removing the possible points at infinity of σ . Let o be a point of R_σ . By [5, Proposition 2.2], we can give the conic $\bar{\sigma}$ the structure of an abelian group $(\bar{\sigma}, +, o)$. The group law has the property that $p + q = r + s$ if and only if the lines pq and rs are parallel. Therefore, the number of directions determined by R_σ is exactly the size of the restricted sumset

$$H := R_\sigma \dot{+} R_\sigma = \{p + q \mid p, q \in R_\sigma, p \neq q\}.$$

By assumption, $|R_\sigma \dot{+} R_\sigma| \leq m < (1 + 4\varepsilon)|R_\sigma|$. Standard results in additive combinatorics imply that $H = R_\sigma \dot{+} R_\sigma = R_\sigma + R_\sigma$ (see [3, Theorem 3]) and H is a coset of a subgroup of $\bar{\sigma}$ (see e.g. [6, Proposition 1.1]). Since $o \in H$, H is actually a subgroup of $\bar{\sigma}$, and $H \supseteq R_\sigma$. It is easy to prove that $\bar{\sigma}$ is isomorphic to \mathbb{R} , $\mathbb{R} \times \mathbb{Z}/2$ or \mathbb{R}/\mathbb{Z} , when $\bar{\sigma}$ is a parabola, a hyperbola or an ellipse, respectively (for example, see [1, Proposition 7.3]). Neither \mathbb{R} nor $\mathbb{R} \times \mathbb{Z}/2$ have finite subgroups of size bigger than 2, so $\bar{\sigma}$ must be an ellipse. The finite subgroups of $\bar{\sigma}$ are, algebraically, cyclic groups and, geometrically, affinely-regular polygons ([5, Lemma 4]).

It remains to prove that $R_{-\sigma}$ is empty. By applying a projective transformation fixing the line at infinity, we may suppose that σ is the unit circle and $R_\sigma + R_\sigma$ the set of m -th roots of unity. Suppose by contradiction that there is a point $p \in R_{-\sigma}$. For all $x \in R_\sigma$, the line px passes through a point in R_∞ . This is impossible by [1, Corollary 7.6], since p is neither a root of unity ($p \notin \sigma$) nor a point at infinity, nor the origin (since R is in general position).²

So $R = R_\sigma$ is a subset of the affinely-regular polygon H , which has size $|R_\sigma \dot{+} R_\sigma| = |R \dot{+} R| = m$, as desired. \square

Remark 3. The bound $m \leq n + cn^{1/6}$ in Corollary 2 can probably be improved by applying the techniques of Green and Tao [1] to the specific setup of Jamison's conjecture, instead of using their structure theorem as a black box.

Remark 4. The main difficulty in Jamison's conjecture is to prove that the set of points R is a subset of a conic $\bar{\sigma}$. Once this step is done, the ambient group is available and the remaining task

²In fact, since R is in general position, we only need a much weaker version of [1, Corollary 7.6].

is a standard question in additive combinatorics, namely to describe the structure of a subset A of $\bar{\sigma}$, given that the restricted sumset $A \dot{+} A$ is small.

As observed by Makhul and Pinchasi [4], Jamison’s conjecture does not hold for $m \geq \frac{3}{2}n$, because $|A \dot{+} A| = \frac{3}{2}|A|$ when $A = H \sqcup (H + g)$ is a union of two cosets of a cyclic subgroup H of $\bar{\sigma}$, and gH has order at least three in the quotient group $\bar{\sigma}/H$.

In the previous example, the set A is again a subset of an ellipse. Even if the bound $2n - O(1)$ does not hold for the original conjecture of Jamison, the following is, to our knowledge, still unsolved.

Open Problem. For $m \leq 2n - O(1)$, every finite set R of n points in general position that determines m distinct directions is contained in an ellipse.

If we know that R is contained in an ellipse, we automatically get a more precise description of the structure of R by well-known results in additive combinatorics such as [6, Proposition 1.1] or [6, Proposition 1.2], as in the proof of Corollary 2.

REFERENCES

- [1] Ben Green and Terence Tao, *On sets defining few ordinary lines*, Discrete & Computational Geometry **50** (2013), no. 2, 409–468.
- [2] Robert E. Jamison, *Few slopes without collinearity*, Discrete mathematics **60** (1986), 199–206.
- [3] Vsevolod F. Lev, *Restricted set addition in groups I: The classical setting*, Journal of the London Mathematical Society **62** (2000), no. 1, 27–40.
- [4] Mehdi Makhul and Rom Pinchasi, *On sets of points in general position that lie on a cubic curve in the plane and determine lines that can be pierced by few points*, http://www2.math.technion.ac.il/~room/ps_files/MP_cubic.pdf, 2021, [Online; accessed 19 January 2021].
- [5] Cédric Pilatte, *On the sets of n points forming $n + 1$ directions*, The Electronic Journal of Combinatorics **27** (2020), no. 1, P1.24.
- [6] Tom Sanders, *The structure theory of set addition revisited*, Bulletin of the American Mathematical Society **50** (2013), no. 1, 93–127.
- [7] G. A. Wade and Jiang-Hsing Chu, *Drawability of complete graphs using a minimal slope set*, The Computer Journal **37** (1994), 139–142.

DEPARTMENT OF MATHEMATICS AND THEIR APPLICATIONS, ÉCOLE NORMALE SUPÉRIEURE (ENS), RUE D’ULM, 45, 75005 PARIS, FRANCE — DEPARTMENT OF MATHEMATICS, UNIVERSITY OF MONS (UMONS), PLACE DU PARC 20, 7000 MONS, BELGIUM.

Email address: `cedric.pilatte@ens.fr`

ON THE SETS OF n POINTS FORMING $n + 1$ DIRECTIONS

CÉDRIC PILATTE

ABSTRACT. Let S be a set of $n \geq 7$ points in the plane, no three of which are collinear. Suppose that S determines $n + 1$ directions. That is to say, the segments whose endpoints are in S form $n + 1$ distinct slopes. We prove that S is, up to an affine transformation, equal to n of the vertices of a regular $(n + 1)$ -gon. This result was conjectured in 1986 by R. E. Jamison.

Keywords. Combinatorics, Discrete Geometry.

1. INTRODUCTION

In 1970, inspired by a problem of Erdős, Scott [15] asked the following question, now known as *the slope problem*: what is the minimum number of directions determined by a set of n points in \mathbb{R}^2 , not all on the same line? By the number of directions (or slopes) of a set S , we mean the size of the quotient set $\{PQ \mid P, Q \in S, P \neq Q\}/\sim$, where \sim is the equivalence relation given by parallelism: $P_1Q_1 \sim P_2Q_2 \iff P_1Q_1 \parallel P_2Q_2$.

Scott conjectured that n points, not all collinear, determine at least $2\lfloor \frac{n}{2} \rfloor$ slopes. This bound can be achieved, for even n , by a regular n -gon; and for odd n , by a regular $(n - 1)$ -gon with its center. After some initial results of Burton and Purdy [2], this conjecture was proven by Ungar [16] in 1982, using techniques of Goodman and Pollack [5]. His beautiful proof is also exposed in the famous *Proofs from the Book* [1, Chapter 11]. Recently, Pach, Pinchasi and Sharir solved the tree-dimensional analogue of this problem, see [12, 13].

A lot of work has been done to determine the configurations where equality in Ungar's theorem is achieved. A *critical set* (respectively *near-critical set*) is a set of n non-collinear points forming $n - 1$ slopes (respectively n slopes). Jamison and Hill described four infinite families and 102 sporadic critical configurations [6, 7, 10]. It is conjectured that this classification is accurate for $n \geq 49$. No classification is known in the near-critical case. See [8] for a survey of these questions, and other related ones.

In this paper, we suppose that no three points of S are collinear (we say that S is in general position). This situation was first investigated by Jamison [9], who proved that S must determine at least n slopes. As above, equality is possible with a regular n -gon. It is a well-known fact that affine transformations preserve parallelism. Therefore, the image of a regular n -gon under an affine transformation also determines exactly n slopes.¹ Jamison proved the converse, i.e. that the affinely regular polygons are the only configurations forming exactly n slopes.

A much more general statement is believed to be true: for some constant c_1 , if a set of n points in general position forms $m \leq 2n - c_1$ slopes, then it is affinely equivalent to n of the vertices of a regular m -gon (see [9]). This would imply, in particular, that for every $c \geq 0$ and n sufficiently large, every simple configuration of n points determining $n + c$ slopes arises from an affinely regular $(n + c)$ -gon, after deletion of c points. Jamison's result thus shows it for $c = 0$. Here, we will prove the case $c = 1$. The general conjecture is still open. In fact, for $c \geq 2$, it is not even known whether the points of S form a convex polygon.

Date: October 2018.

¹A polygon obtained as the image of a regular polygon by an affine transformation is sometimes called an *affinely-regular* or *affinely regular* polygon.

Every affinely regular polygon is inscribed in an ellipse. Conics will play an important role in our proof. Another problem of Elekes [3] is the following: for all $m \geq 6$ and $C > 0$, there exists some $n_0(m, C)$ such that every set $S \subset \mathbb{R}^2$ with $|S| \geq n_0(m, C)$ forming at most $C|S|$ slopes contains m points on a (possibly degenerate) conic. It is still unsolved, even for $m = 6$.

2. RESULTS

2.1. Preliminary Remarks. Let S be a set of n points in the plane, in general position, that determines exactly $n + 1$ slopes. If S had a point lying strictly inside its convex hull, there would be at least $n + 2$ slopes, as was proved by Jamison [9, Theorem 7]. Therefore, we know that we can label the points of S as A_1, \dots, A_n , such that $A_1A_2 \dots A_n$ is a convex polygon.

For every point $A_i \in S$, there are $n - 1$ segments, with distinct slopes, joining A_i to the other points of S . We will say that a slope is *forbidden* at A_i if it is not the slope of any segment A_iA_j , for $j \neq i$. Since S determines $n + 1$ slopes, *there are exactly two forbidden slopes* at each point of S .

We will denote by ∇A_iA_j the slope of the line A_iA_j . Thus, an equality like $\nabla A_{i_1}A_{i_2} = \nabla A_{i_3}A_{i_4}$ is equivalent to $A_{i_1}A_{i_2} \parallel A_{i_3}A_{i_4}$. Throughout our main proof, we will repeatedly make use of the next lemma. It will be particularly useful to prove that a slope is forbidden at a point or that two slopes are equal. As an obvious corollary, we have that $\nabla A_{i-1}A_{i+1}$ is forbidden at A_i for all $i \in \mathbb{Z}$. Throughout the paper, when we say “for all $i \in \mathbb{Z}$ ”, we consider the indices modulo n , so that $A_{n+1} := A_1$, and so on.

Lemma 2.1. *Let $1 \leq i < j < k \leq n$. Exactly one of the following is true:*

- *the slope of A_iA_k is forbidden at A_j ;*
- *$\exists p, i < p < k$ such that $A_iA_k \parallel A_jA_p$.*

Moreover, in the second case, $\nabla A_jA_p \notin \{\nabla A_iA_l \mid l \neq j, k\} \cup \{\nabla A_lA_j \mid l \neq i, k\}$.

Proof. This is almost immediate from the definition of a forbidden slope. In the second case, if $p \in \{1, \dots, n\}$ were not between i and k , the segments A_iA_k and A_jA_p would intersect. Finally, if ∇A_jA_p were equal to some ∇A_iA_l , then $A_iA_k \parallel A_jA_p \parallel A_iA_l$, so A_i, A_k and A_l would be aligned, a contradiction. The same is true for the segments A_lA_j . \square

We will also need the following result, which can be found in [14, Chapter 1].

Proposition 2.2. *Let \mathcal{C} be a non-degenerate conic and O a point on \mathcal{C} . If P, Q are two points on \mathcal{C} , define $P + Q$ to be the unique point R on \mathcal{C} such that $RO \parallel PQ$ (with the convention that XX is the tangent to \mathcal{C} at X , for $X \in \mathcal{C}$). This addition turns \mathcal{C} into an abelian group, of which O is the identity element.*

In particular, for P, Q, R, S four points on \mathcal{C} , we have $P + Q = R + S$ if and only if $PQ \parallel RS$. Five points in general position determine a unique conic. Lemma 2.3 will enable us to introduce conics in the proof, in order to use Proposition 2.2.

Lemma 2.3. *Suppose P_1, \dots, P_6 are points in the plane such that $P_1P_6 \parallel P_2P_5$, $P_2P_3 \parallel P_1P_4$ and $P_4P_5 \parallel P_3P_6$. Then P_1, \dots, P_6 lie on a common conic.*

Proof. This follows immediately from Pascal’s theorem, with the hexagon $\mathcal{H} = P_1P_4P_5P_2P_3P_6$. Indeed, the intersections of the opposite sides of \mathcal{H} are collinear on the line at infinity. \square




FIGURE 1. Illustration of Lemma 2.3.

For the reader's convenience, we reproduce here a result of Korchmáros [11] (which is also discussed in [4]), that we will use twice in the proof.

Lemma 2.4. *Let P_1, \dots, P_n be distinct points on a non-degenerate conic. Suppose that, for all $j \in \mathbb{Z}$, $P_{j+1}P_{j+2} \parallel P_jP_{j+3}$. Then, P is affinely equivalent to a regular n -gon.*

2.2. Main Theorem.

Theorem 2.5. *Any set S of $n \geq 7$ points in the plane, in general position, that determines exactly $n + 1$ slopes, is affinely equivalent to n of the vertices of a regular $(n + 1)$ -gon.*

Proof. We use the notations of Section 2.1: $S = \{A_1, A_2, \dots, A_n\}$ where $A_1A_2 \dots A_n$ is a convex polygon. We will split the proof into two cases. In the first case, we suppose that, for every $i \in \mathbb{Z}$, $A_{i+1}A_{i+2} \parallel A_iA_{i+3}$. If this fails for some i , we can assume that this i is 1.

Case 1 For every $i \in \mathbb{Z}$, $A_{i+1}A_{i+2} \parallel A_iA_{i+3}$.

We will distinguish subcases according to which segments are parallel to A_iA_{i+5} . As we will see, none of the subcases is actually possible.

Case 1.1 For all $i \in \mathbb{Z}$, $A_iA_{i+5} \parallel A_{i+1}A_{i+4}$.

Let A_{k+1}, \dots, A_{k+6} be any six consecutive points of S . We have $A_{k+1}A_{k+6} \parallel A_{k+2}A_{k+5}$, $A_{k+2}A_{k+3} \parallel A_{k+1}A_{k+4}$ and $A_{k+4}A_{k+5} \parallel A_{k+3}A_{k+6}$ from our two assumptions. Thus, Lemma 2.3 implies that the six points lie on a common conic. As this is true for any six consecutive points, and since five points in general position (i.e. no three collinear) determine a unique conic, all the A_i 's lie on the same conic. Together with the fact that $\forall i, A_{i+1}A_{i+2} \parallel A_iA_{i+3}$, this implies that $A_1A_2 \dots A_n$ is affinely equivalent to a regular n -gon, by Lemma 2.4. Therefore, S determines exactly n directions, which is a contradiction.

Case 1.2 For some $i \in \mathbb{Z}$, we have $A_iA_{i+5} \parallel A_{i+2}A_{i+4}$.

Say $i = 1$, meaning $A_1A_6 \parallel A_3A_5$. By Lemma 2.1 applied three times, we see that ∇A_2A_6 is forbidden at A_3, A_4 and A_5 (here, we have used that $A_2A_6 \not\parallel A_3A_5$ and, for A_4 , that $A_3A_4 \parallel A_2A_5$ and $A_4A_5 \parallel A_3A_6$). For $l = 3, 4, 5$, we know that ∇A_2A_6 and $\nabla A_{l-1}A_{l+1}$ are exactly the two forbidden slopes at A_l . Therefore, ∇A_1A_5 is not forbidden at A_4 , hence, by Lemma 2.1 again, we conclude that $A_1A_5 \parallel A_2A_4$. Similarly, ∇A_3A_7 is not forbidden at A_4 , so $A_3A_7 \parallel A_4A_6$. As the slope of A_2A_7 is not forbidden at A_5 , we conclude $A_2A_7 \parallel A_4A_5$ ($\parallel A_3A_6$). We have $A_3A_4 \parallel A_2A_5$, $A_3A_6 \parallel A_2A_7$ and we just showed that $A_2A_7 \parallel A_3A_6$. By Lemma 2.3, A_2, A_3, \dots, A_7 lie on a common conic.




FIGURE 2. Case 1.2.

We will equip this conic with the group structure described in Proposition 2.2, with A_7 the zero element. We will write $A_7 = 0$ and $A_6 = x$. Then, $A_5A_6 \parallel A_4A_7$, $A_4A_5 \parallel A_3A_6$ and $A_4A_6 \parallel A_3A_7$ together imply $A_5 = 2x$, $A_4 = 3x$ and $A_3 = 4x$. Also, $A_3A_4 \parallel A_2A_5$ gives $A_2 = 5x$. Let B be the point on the conic with $B = 6x$. We thus have $A_2A_3 \parallel BA_4$ and $A_2A_4 \parallel BA_5$. However, there can only be one point P with $A_2A_3 \parallel PA_4$ and $A_2A_4 \parallel PA_5$. As A_1 is such a point, $A_1 = B = 6x$. This contradicts $A_1A_6 \parallel A_3A_5$, as $A_1 + A_6 = 6x + x \neq 4x + 2x = A_3 + A_5$.

Case 1.3 For some $i \in \mathbb{Z}$, we have $A_iA_{i+5} \parallel A_{i+1}A_{i+3}$.

This is exactly the previous case after having relabelled every A_i as A_{n+1-i} .

Case 1.4 The previous cases do not apply.

If none of the previous cases is possible, there must be some i , say $i = 1$, for which A_1A_6 is not parallel to any of A_2A_5 , A_3A_5 and A_2A_4 . Then, ∇A_1A_6 is forbidden at A_2, A_3, A_4 and A_5 . Once again, we deduce that the forbidden slopes at A_l , $2 \leq l \leq 5$, are ∇A_1A_6 and $\nabla A_{l-1}A_{l+1}$. We use Lemma 2.1 to find $A_2A_6 \parallel A_3A_5$ (applied with $A_k = A_4$) and $A_1A_5 \parallel A_2A_4$ ($A_k = A_2$).

Let \mathcal{C} be the conic passing through A_1, A_2, \dots, A_5 . We use Proposition 2.2 to define a group structure on \mathcal{C} , with $A_1 = 0$. Let $A_2 = x$ and $A_3 = y$. From $A_2A_3 \parallel A_1A_4$ and $A_3A_4 \parallel A_2A_5$, we have $A_4 = x + y$ and $A_5 = 2y$. But $A_2A_4 \parallel A_1A_5$ implies $y = 2x$, so $A_i = (i - 1)x$ for $1 \leq i \leq 5$. We use the same argument as before. Let $B = 5x$, then $A_4A_5 \parallel A_3B$ and $A_3A_5 \parallel A_2B$, so $B = A_6 = 5x$. We deduce $A_1A_6 \parallel A_2A_5$, a contradiction.




FIGURE 3. Case 1.4.

Case 2 We have $A_2A_3 \not\parallel A_1A_4$ (without loss of generality).

Without loss of generality, we can also suppose that the point A_4 is closer to the line A_2A_3 than is A_1 . In this situation, the line parallel to A_2A_3 passing through A_4 intersects the segment $[A_1A_2]$ in its relative interior, and the line parallel to A_2A_3 passing through A_1 does not intersect the segment $[A_3A_4]$.

From $A_2A_3 \not\parallel A_1A_4$, we deduce that the forbidden slopes at A_2 and A_3 are ∇A_1A_3 , ∇A_1A_4 and ∇A_2A_4 , ∇A_1A_4 , respectively. Thus, $A_1A_2 \parallel A_nA_3$ and $A_2A_5 \parallel A_3A_4$. We now show that A_2A_3 is forbidden at A_4 . Suppose, for some k , that $A_2A_3 \parallel A_4A_k$. Then, k has to be between 5 and n , so $A_1A_2A_3A_4A_k$ must be a convex polygon, with $A_2A_3 \parallel A_4A_k$. We can see that this contradicts the fact that A_4 is closer than A_1 to the line A_2A_3 .

Case 2.1 $A_{n-1}A_2 \parallel A_nA_1$.

We want to show that this case is impossible. From Lemma 2.1, we find $A_{n-1}A_3 \parallel A_nA_2$. When we apply this lemma again with the slope of A_nA_4 , we find that A_nA_4 is parallel to A_1A_3 , because A_2A_3 is forbidden at A_4 . In the same way, we get $A_{n-1}A_4 \parallel A_nA_3$.




FIGURE 4. Case 2.1.

Let \mathcal{C} be the conic passing through A_3, A_2, A_1, A_n and A_{n-1} . Again, we use Proposition 2.2, setting $A_{n-1} = 0$. Let $A_n = x$ and $A_2 = y$. From $A_{n-1}A_3 \parallel A_nA_2$ we deduce $A_3 = x + y$, and from $A_{n-1}A_4 \parallel A_nA_3$ we get $A_4 = y + 2x$. Let $B = 2x$. Then $A_nA_4 \parallel BA_3$ and $A_nA_3 \parallel BA_2$. This means that B belongs to the line parallel to A_nA_4 through A_3 and to the line parallel to A_nA_3 through A_2 . So $B = A_1$, i.e. $A_1 = 2x$. On the one hand, the relation $A_{n-1}A_2 \parallel A_nA_1$ gives $0 + y = x + 2x$. On the other hand, $A_2A_3 \not\parallel A_1A_4$ yields $y + (y + x) \neq 2x + (y + 2x)$. This is absurd.

Case 2.2 $A_{n-1}A_2 \not\parallel A_nA_1$.

This is the last case of the proof, and the only case that produces valid configurations of points. As $A_{n-1}A_2 \not\parallel A_nA_1$, $\nabla A_{n-1}A_2$ is forbidden at A_1 . With ∇A_0A_2 , those are the two forbidden slopes at A_1 . Therefore, none of ∇A_2A_i , $3 \leq i \leq n-2$ is forbidden at A_1 . So, every ∇A_2A_i , $3 \leq i \leq n-2$, corresponds to a unique ∇A_1A_j for some j .




FIGURE 5. Case 2.2.

A simple but important observation is that, for all $3 \leq i_1, i_2 \leq n - 2$ and $4 \leq j_1, j_2 \leq n$,

$$\begin{cases} A_2A_{i_1} \parallel A_1A_{j_1} \\ A_2A_{i_2} \parallel A_1A_{j_2} \end{cases} \implies (i_1 < i_2 \iff j_1 < j_2).$$

That is, the assignment f that maps every $3 \leq i \leq n - 2$ to the unique $4 \leq j \leq n$ such that $A_2A_i \parallel A_1A_j$ must be strictly increasing. Moreover, it has to satisfy $f(3) \neq 4$ as we assumed $A_2A_3 \not\parallel A_1A_4$. The unique possibility is then $f(i) = i + 2$ for every i . We have proven that, for $5 \leq i \leq n$, $A_2A_{i-2} \parallel A_1A_i$.

Claim 2.2.1. For every $i \in \{5, \dots, n\}$,

- (1) $A_{i-2}A_i$ and A_2A_{i-2} are the two forbidden slopes at A_{i-1} , and;
- (2) $\forall k \in \{3, \dots, i - 2\}$, $A_{i-1}A_k \parallel A_iA_{k-1}$.

Proof of claim. For $i = 5$, we have already proven those two statements. We will prove them for $i = j$, assuming it has already been proven for all $5 \leq i \leq j - 1$.

- (1) We have to show that A_2A_{j-2} is forbidden at A_{j-1} . This is clear as $A_2A_{j-2} \parallel A_1A_j$ and there is no point of S between A_1 and A_2 .
- (2) Since we know the forbidden slopes at A_{j-1} , we can use Lemma 2.1 at the point A_{j-1} several times, with different slopes. First, ∇A_jA_{j-3} is not forbidden, so $A_jA_{j-3} \parallel A_{j-1}A_{j-2}$. Then ∇A_jA_{j-4} is not forbidden, and is distinct from $\nabla A_{j-1}A_{j-2} = \nabla A_jA_{j-3}$, so $A_jA_{j-4} \parallel A_{j-1}A_{j-3}$. We can continue this way, until we get $A_jA_2 \parallel A_{j-1}A_3$. This proves the claim. \square

In particular, for every $i \in \{6, \dots, n - 1\}$, we have $A_3A_i \parallel A_2A_{i+1}$, $A_5A_i \parallel A_4A_{i+1}$. As $A_3A_4 \parallel A_2A_5$, we can use Lemma 2.1, which shows that A_2, A_3, A_4, A_5, A_i and A_{i+1} lie on a conic. As this is true for every $6 \leq i \leq n - 1$, we know that the A_i 's, for $2 \leq i \leq n$, all lie on a common conic (because there is a unique conic passing through five points in general position).

As we have done several times in this proof, we use the group structure on the conic given by parallelism. Choose A_2 to be the identity element, let $A_3 = x$. Solving

$$\begin{cases} A_3A_4 \parallel A_2A_5 \\ A_3A_6 \parallel A_4A_5 \\ A_3A_5 \parallel A_2A_6 \end{cases}$$

gives $A_4 = 2x$, $A_5 = 3x$ and $A_6 = 4x$. Then, a simple induction (using $A_{i-1}A_3 \parallel A_iA_2$) gives $A_i = (i-2)x$ for all $i \in \{2, \dots, n\}$. Let B be the point on the conic with $B = -2x$. Then $A_2A_3 \parallel BA_5$ and $A_2A_4 \parallel BA_6$. However, we proved before that $A_2A_3 \parallel A_1A_5$ and $A_2A_4 \parallel A_1A_6$, so $A_1 = B = -2x$.

To summarize, we know that all the n points of S are on a conic, $A_i = (i-2)x$ for $i \in \{2, \dots, n\}$ and $A_1 = -2x$. We use the group structure one last time: $A_3A_n \parallel A_1A_2$ implies $x + (n-2)x = -2x + 0$, so $(n+1)x = 0$. Therefore, the subgroup generated by $A_3 = x$ is a finite cyclic group of order $n+1$:

$$\langle A_3 \rangle = \left\{ \underbrace{A_2}_0, \underbrace{A_3}_x, \underbrace{A_4}_{2x}, \dots, \underbrace{A_{n-1}}_{(n-3)x}, \underbrace{A_n}_{(n-2)x}, \underbrace{A_1}_{(n-1)x}, -x \right\}.$$

To finish the proof, we use the more convenient notations $P_j := jx$ for $0 \leq j \leq n$ (so that every A_i is a P_j). If the indices are considered modulo $n+1$, we have, for all $j \in \mathbb{Z}$, $P_{j+1}P_{j+2} \parallel P_jP_{j+3}$, because $(j+1)x + (j+2)x = jx + (j+3)x$. By Lemma 2.4, $P_0P_1P_2 \dots P_n$ is, up to an affine transformation, a regular $(n+1)$ -gon. \square

3. ACKNOWLEDGEMENTS

I would like to thank Christian Michaux, without whom this work would not have been possible.

REFERENCES

- [1] Martin Aigner and Günter Ziegler. *Proofs from the Book*, volume 274. Springer, 2010.
- [2] G. R. Burton and G. B. Purdy. The directions determined by n points in the plane. *Journal of the London Mathematical Society*, 2(1):109–114, 1979.
- [3] György Elekes. On linear combinatorics III. Few directions and distorted lattices. *Combinatorica*, 19(1):43–53, 1999.
- [4] J. Chris Fisher and Robert E. Jamison. Properties of affinely regular polygons. *Geometriae Dedicata*, 69(3):241–259, 1998.
- [5] Jacob E. Goodman and Richard Pollack. A combinatorial perspective on some problems in geometry. *Congressus Numerantium*, 32(0981):383–394, 1981.
- [6] Robert E. Jamison. Planar configurations which determine few slopes. *Geometriae Dedicata*, 16(1):17–34, 1984.
- [7] Robert E. Jamison. Structure of slope-critical configurations. *Geometriae Dedicata*, 16(3):249–277, 1984.
- [8] Robert E. Jamison. A survey of the slope problem. *Annals of the New York Academy of Sciences*, 440(1):34–51, 1985.
- [9] Robert E. Jamison. Few slopes without collinearity. *Discrete mathematics*, 60:199–206, 1986.
- [10] Robert E. Jamison and Dale Hill. A catalogue of sporadic slope-critical configurations. *Congressus Numerantium*, 40:101–125, 1983.
- [11] G. Korchmáros. Poligoni affini-regolari dei piani di galois d'ordine dispari. *Atti Accad. Naz. Lincei Rend. Cl. Sci. Fis. Mat. Natur. (8)*, 56(5):690–697, 1974.
- [12] János Pach, Rom Pinchasi, and Micha Sharir. On the number of directions determined by a three-dimensional points set. *Journal of Combinatorial Theory, Series A*, 108(1):1–16, 2004.
- [13] János Pach, Rom Pinchasi, and Micha Sharir. Solution of scott's problem on the number of directions determined by a point set in 3-space. *Discrete & Computational Geometry*, 38(2):399–441, 2007.
- [14] Viktor V. Prasolov and Yuri P. Solov'yev. *Elliptic functions and elliptic integrals*, volume 170. American Mathematical Soc., 1997.
- [15] Paul R. Scott. On the sets of directions determined by n points. *The American Mathematical Monthly*, 77(5):502–505, 1970.
- [16] Peter Ungar. $2n$ noncollinear points determine at least $2n$ directions. *Journal of Combinatorial Theory, Series A*, 33(3):343–347, 1982.

DEPARTMENT OF MATHEMATICS, UNIVERSITY OF MONS (UMONS), PLACE DU PARC 20, 7000 MONS, BELGIUM.