

Fault-Tolerant Metric Dimension of $P(n, 2)$ with Prism Graph ^{*}

Z. Ahmad¹, M. O. Ahmad², A. Q. Baig³, M. Naeem³

¹Department of Mathematics, The University of Lahore, Pakpattan Campus, Pakistan

²Department of Mathematics and Statistics, The University of Lahore, Lahore, Pakistan

³Department of Mathematics, The University of Lahore, Pakpattan Campus, Pakistan

E-mail: zubair786ppn@gmail.com, drchadury@yahoo.com, aqbaig1@gmail.com, naeem@gmail.com

2010 Mathematics Subject Classification: 05C12

Abstract. Let G be a connected graph and $d(a, b)$ be the distance between the vertices a and b . A subset $U = \{u_1, u_2, \dots, u_k\}$ of the vertices is called a resolving set for G if for every two distinct vertices $a, b \in V(G)$, there is a vertex $u_\xi \in U$ such that $d(a, u_\xi) \neq d(b, u_\xi)$. A resolving set containing a minimum number of vertices is called a metric basis for G and the number of vertices in a metric basis is its metric dimension denoted by $dim(G)$. A resolving set U for G is fault-tolerant if $U \setminus \{u\}$ is also a resolving set, for each $u \in U$, and the fault-tolerant metric dimension of G is the minimum cardinality of such a set. In this paper we introduce the study of the fault-tolerant metric dimension of $P(n, 2)$ with prism graph.

Keywords: *Fault-tolerant, Metric dimension, Basis, Resolving set, Petersen graph, Prism Graph*

1 Notation and preliminary results

The length of path between two vertices $a, b \in V(G)$ in a connected graph is the shortest distance $d(a, b)$ between them. Let c be a vertex of G and $U = \{u_1, u_2, \dots, u_k\}$ be an ordered set of vertices of G . The representation $r(c|U)$ of c with respect to U is the k -tuple $(d(c, u_1), d(c, u_2), \dots, d(c, u_k))$. If distinct vertices of G have distinct representations with respect to U , then U is called a resolving set. If the resolving set has minimum number of elements then the set is called a basis for G and the cardinality of U is called the metric dimension of G and is denoted by $dim(G)$. For further study about basis and

^{*} This research is partially supported by The University of Lahor, Pakpattan Campus, Pakistan

metric dimension see [1,2,5,7,9,10,11,12].

Generalized Petersen graph $P(n, m)$ is an undirected graph having size $3m$ and order $2n$. It is 3 connected graph which is 3 partite and has independence number 4. If $U = \{u_1, u_2, \dots, u_k\}$ is the ordered set of vertices of a graph G , then ξ^{th} component of $r(c|U)$ is $0 \Leftrightarrow c = u_\xi$. Thus, in order to show that U is a resolving set it suffices to verify that $r(a|U) \neq r(b|U)$ for each pair of distinct vertices $a, b \in V(G) \setminus U$.

The following property is very useful in order to calculate the dimension of a graph G .

Lemma 1. *Let U be a resolving set for a connected graph G and $a, b \in V(G)$. If $d(a, u) = d(b, u)$ for all vertices $u \in V(G) \setminus \{a, b\}$, then $\{a, b\} \cap U \neq \emptyset$.*

Slater [10] gave the idea of metric dimension in 1975. This concept was applied in chemistry in [3], this invariant was applied to the navigation of robots in networks in [8]. The dimension of a path graph in a connected graph G is 1 but if G is a cycle C_n with $n \geq 3$ then dimension of G is 2.

2 $P(n, 2)$ with Prism Graph




Fig. 1. $P(8, 2)$ with prism graph

Let us denote the inner vertices of G by a_1, a_2, \dots, a_n , central vertices by b_1, b_2, \dots, b_n and outer vertices by c_1, c_2, \dots, c_n . We call the cycle induced by $\{a_\xi : 1 \leq \xi \leq n\}$, the inner cycle, the vertices $\{b_\xi : 1 \leq \xi \leq n\}$, the middle cycle and the vertices $\{c_\xi : 1 \leq \xi \leq n\}$ the outer cycle. We define $V(G) =$

$\{a_\xi, b_\xi, c_\xi; 1 \leq \xi \leq n\}$ and $E(G) = \{a_i b_{\xi+2}; 1 \leq \xi \leq n \pmod{n}\} \cup \{b_\xi b_{\xi+1}; 1 \leq \xi \leq n \pmod{n}\} \cup \{c_\xi c_{\xi+1}; 1 \leq \xi \leq n \pmod{n}\} \cup \{a_\xi b_\xi; b_\xi c_\xi; 1 \leq \xi \leq n \pmod{n}\}$

Theorem 1. For every $n \geq 5; n \equiv 0, 1, 2, 3 \pmod{4}$ we have

$$G = 3.$$

Proof. We will show that only three vertices appropriately chosen suffice to resolve all vertices. We have the following cases:

Case 1: $n \equiv 0 \pmod{4}$

For $n = 4k$ with $k > 1$ and $k \in \mathbb{Z}^+$. In this case, $\{a_1, a_2, b_{2k+1}\}$ resolves $V(G)$. Indeed, a_1 and a_2 distinguish the vertices in the inner cycle, middle cycle and outer cycle. To show that $\{a_1, a_2, b_{2k+1}\}$ resolves vertices of G , first we give representations of the vertices in G with respect to W .

Representations of the inner cycle are

$$r(a_{2\xi+1}|W) = \begin{cases} (\xi, \xi + 2, \xi - 1, k + 1 - \xi), & 2 \leq \xi \leq k; \\ (2k - \xi, 2k + 3 - \xi, 2k + 1 - \xi, \xi + 1 - k), & \xi = k + 1; \\ (2k - \xi, 2k + 3 - \xi, 2k + 1 - \xi, \xi + 1 - k), & k + 2 \leq \xi \leq 2k - 1. \end{cases}$$

$$r(a_{2\xi}|W) = \begin{cases} (\xi + 2, \xi - 1, \xi + 1, k + 2 - \xi), & 2 \leq \xi \leq k; \\ (2k + 3 - \xi, 2k + 1 - \xi, 2k + 3 - \xi, \xi + 1 - k), & \xi = k + 1; \\ (2k + 3 - \xi, 2k + 1 - \xi, 2k + 4 - \xi, \xi + 1 - k), & k + 2 \leq \xi \leq 2k. \end{cases}$$

Representations of the middle cycle are

$$r(b_{2\xi+1}|W) = \begin{cases} (1, 2, 2, k + 2), & \xi = 0; \\ (\xi + 1, \xi + 1, \xi, k + 2 - \xi), & 1 \leq \xi \leq k - 2; \\ (\xi + 1, \xi + 1, \xi, 2), & \xi = k - 1; \\ (\xi - 1, \xi, \xi, 2), & \xi = k + 1; \\ (2k + 1 - \xi, 2k + 2 - \xi, 2k + 2 - \xi, \xi + 2 - k), & k + 2 \leq \xi \leq 2k - 1. \end{cases}$$

$$r(b_{2\xi}|W) = \begin{cases} (2, 1, 2, k + 2), & \xi = 1; \\ (\xi + 1, \xi, \xi, k + 3 - \xi), & 2 \leq \xi \leq k - 2; \\ (\xi + 1, \xi, \xi, 3), & \xi = k - 1; \\ (\xi + 1, \xi, \xi, 1), & \xi = k; \\ (\xi, \xi, \xi + 1, 1), & \xi = k + 1; \\ (\xi - 2, \xi - 2, \xi, 3), & \xi = k + 2; \\ (2k + 2 - \xi, 2k + 2 - \xi, 2k + 3 - \xi, \xi + 2 - k), & k + 3 \leq \xi \leq 2k. \end{cases}$$

Representations of the outer cycle are

$$r(c_{2\xi+1}|W) = \begin{cases} (2, 3, 3, k + 3), & \xi = 0; \\ (\xi + 2, \xi + 2, \xi + 1, k + 3 - \xi), & 1 \leq \xi \leq k - 2; \\ (\xi + 2, \xi + 2, \xi + 1, 3), & \xi = k - 1; \\ (\xi + 2, \xi + 2, \xi + 1, 1), & \xi = k; \\ (\xi, \xi + 1, \xi + 1, 3), & \xi = k + 1; \\ (2k + 2 - \xi, 2k + 3 - \xi, 2k + 3 - \xi, \xi + 3 - k), & k + 2 \leq \xi \leq 2k - 1. \end{cases}$$

$$r(c_{2\xi}|W) = \begin{cases} (3, 2, 3, k+4), & \xi = 1; \\ (\xi + 2, \xi + 1, \xi + 1, k + 4 - \xi), & 2 \leq \xi \leq k - 2; \\ (\xi + 2, \xi + 1, \xi + 1, 4), & \xi = k - 1; \\ (\xi + 2, \xi + 1, \xi + 1, 2), & \xi = k; \\ (\xi + 1, \xi + 1, \xi + 2, 2), & \xi = k + 1; \\ (\xi - 1, \xi - 1, \xi, 4), & \xi = k + 2; \\ (2k + 3 - \xi, 2k + 3 - \xi, 2k + 4 - \xi, \xi + 3 - k), & k + 3 \leq \xi \leq 2k. \end{cases}$$

From the above discussion it follows that $\dim(G) = 4$ in this case.

Case 2: $n \equiv 1 \pmod{4}$

For $n = 4k + 1$ with $k \geq 1$ and $k \in \mathbb{Z}^+$. When $n = 5$ then $W = \{a_1, a_2, b_3\}$ is the resolving set for $V(G)$. In $V(G)$ the representations of the vertices are $r(a_4|W) = (1, 1, 2, 2)$, $r(a_5|W) = (2, 1, 1, 2)$, $r(b_1|W) = (1, 2, 2, 2)$, $r(b_2|W) = (2, 1, 2, 1)$, $r(b_4|W) = (2, 2, 2, 1)$, $r(b_5|W) = (2, 2, 2, 2)$, $r(c_1|W) = (2, 3, 3, 3)$, $r(c_2|W) = (3, 2, 3, 2)$, $r(c_3|W) = (3, 3, 2, 1)$, $r(c_4|W) = (3, 3, 3, 2)$ and $r(c_5|W) = (3, 3, 3, 3)$.

When $n = 9$ then $W = \{a_1, a_2, b_5\}$ is the resolving set for $V(G)$. In $V(G)$ the representations of the vertices are

$r(a_4|W) = (3, 1, 3, 2)$, $r(a_5|W) = (2, 3, 1, 1)$, $r(a_6|W) = (2, 2, 3, 2)$, $r(a_7|W) = (3, 2, 2, 2)$, $r(a_8|W) = (1, 3, 2, 3)$, $r(a_9|W) = (3, 1, 4, 3)$, $r(b_1|W) = (1, 2, 2, 4)$, $r(b_2|W) = (2, 1, 2, 3)$, $r(b_3|W) = (2, 2, 1, 2)$, $r(b_4|W) = (3, 2, 2, 1)$, $r(b_6|W) = (3, 3, 3, 1)$, $r(b_7|W) = (3, 3, 3, 2)$, $r(b_8|W) = (2, 3, 3, 3)$, $r(b_9|W) = (2, 2, 3, 4)$, $r(c_1|W) = (2, 3, 3, 5)$, $r(c_2|W) = (3, 2, 3, 4)$, $r(c_3|W) = (3, 3, 2, 3)$, $r(c_4|W) = (4, 3, 3, 2)$, $r(c_5|W) = (4, 4, 3, 1)$, $r(c_6|W) = (4, 4, 4, 2)$, $r(c_7|W) = (4, 4, 4, 3)$, $r(c_8|W) = (3, 4, 4, 4)$ and $r(c_9|W) = (3, 3, 4, 5)$.

When $n \geq 13$ then in this case, $\{a_1, a_2, b_{2k+1}\}$ resolves $V(G)$. Indeed, a_1 and a_2 distinguish the vertices in the inner cycle, inner pendent cycle, outer pendent cycle and outer cycle. To show that $\{a_1, a_2, b_{2k+1}\}$ resolves vertices of G , first we give representations of the vertices in G with respect to W .

Representations of the inner cycle are

$$r(a_{2\xi+1}|W) = \begin{cases} (\xi, \xi + 2, \xi - 1, k + 1 - \xi), & 2 \leq \xi \leq k - 1; \\ (\xi, \xi + 1, \xi - 1, 1), & \xi = k; \\ (\xi, \xi - 1, \xi - 1, 2), & \xi = k + 1; \\ (2k + 3 - \xi, 2k + 1 - \xi, 2k + 3 - \xi, \xi + 1 - k), & \xi = k + 2; \\ (2k + 3 - \xi, 2k + 1 - \xi, 2k + 4 - \xi, \xi + 1 - k), & k + 3 \leq \xi \leq 2k. \end{cases}$$

$$r(a_{2\xi}|W) = \begin{cases} (\xi + 2, \xi - 1, \xi + 1, k + 2 - \xi), & 2 \leq \xi \leq k - 1; \\ (\xi + 1, \xi - 1, \xi + 1, 2), & \xi = k; \\ (\xi - 1, \xi - 1, \xi, 2), & \xi = k + 1; \\ (\xi - 3, \xi - 1, \xi - 2, 3), & \xi = k + 2; \\ (2k + 1 - \xi, 2k + 4 - \xi, 2k + 2 - \xi, \xi + 1 - k), & k + 3 \leq \xi \leq 2k. \end{cases}$$

Representations of the outer pendent cycle are

$$r(b_{2\xi+1}|W) = \begin{cases} (1, 2, 2, k+2), & \xi = 0; \\ (\xi + 1, \xi + 1, \xi, 2k - 4 - \xi), & 1 \leq \xi \leq k - 2; \\ (\xi + 1, i + 1, \xi, 2k - 2\xi), & k - 1 \leq \xi \leq k; \\ (2k + 2 - \xi, 2k + 2 - \xi, 2k + 2 - \xi, 2), & \xi = k + 1; \\ (2k + 2 - \xi, 2k + 2 - \xi, 2k + 3 - \xi, \xi + 2 - k), & k + 2 \leq \xi \leq 2k. \end{cases}$$

$$r(b_{2\xi}|W) = \begin{cases} (2, 1, 2, 2k - 4), & \xi = 1; \\ (\xi + 1, \xi, \xi, 2k - 3 - \xi), & 2 \leq \xi \leq k - 2; \\ (\xi + 1, \xi, \xi, 2k + 1 - 2\xi), & k - 1 \leq \xi \leq k; \\ (\xi, \xi, \xi, 1), & \xi = k + 1; \\ (\xi - 2, \xi - 1, \xi - 1, 3), & \xi = k + 2; \\ (2k + 2 - \xi, 2k + 3 - \xi, 2k + 3 - \xi, \xi + 2 - k), & k + 3 \leq \xi \leq 2k. \end{cases}$$

Representations of the outer cycle are

$$r(c_{2\xi+1}|W) = \begin{cases} (2, 3, 3, k + 3), & \xi = 0; \\ (\xi + 2, \xi + 2, \xi + 1, 2k - 3 - \xi), & 1 \leq \xi \leq k - 2; \\ (\xi + 2, \xi + 2, \xi + 1, 3), & \xi = k - 1; \\ (\xi + 2, \xi + 2, \xi + 1, 1), & \xi = k; \\ (2k + 3 - \xi, 2k + 3 - \xi, 2k + 3 - \xi, 3), & \xi = k + 1; \\ (2k + 3 - \xi, 2k + 3 - \xi, 2k + 4 - \xi, \xi + 3 - k), & k + 2 \leq \xi \leq 2k. \end{cases}$$

$$r(c_{2\xi}|W) = \begin{cases} (3, 2, 3, 2k - 3), & \xi = 1; \\ (\xi + 2, \xi + 1, \xi + 1, 2k - 2 - \xi), & 2 \leq \xi \leq k - 2; \\ (\xi + 2, \xi + 1, \xi + 1, 2k + 2 - 2\xi), & k - 1 \leq \xi \leq k; \\ (\xi + 1, \xi + 1, \xi + 1, 2), & \xi = k + 1; \\ (\xi - 1, \xi, \xi, 4), & \xi = k + 2; \\ (2k + 3 - \xi, 2k + 4 - \xi, 2k + 4 - \xi, \xi + 3 - k), & k + 3 \leq \xi \leq 2k. \end{cases}$$

From the above discussion it follows that $\dim((G)) = 4$ in this case.

Case 3: $n \equiv 2 \pmod{4}$

For $n = 4k + 2$ with $k \geq 1$ and $k \in \mathbb{Z}^+$. In this case, $\{a_1, a_2, b_{2k+2}\}$ resolves $V(G)$. Indeed, a_1 and a_2 distinguish the vertices in the inner cycle, inner pendent cycle, outer pendent cycle and outer cycle. To show that $\{a_1, a_2, b_{2k+2}\}$ resolves vertices of G , first we give representations of the vertices in G with respect to W .

Representations of the inner cycle are

$$r(a_{2\xi+1}|W) = \begin{cases} (\xi, \xi + 2, \xi - 1, k + 2 - \xi), & 2 \leq \xi \leq k; \\ (2k + 1 - \xi, 2k + 4 - \xi, 2k + 1 - \xi, \xi + 1 - k), & \xi = k + 1; \\ (2k + 1 - \xi, 2k + 4 - \xi, 2k + 2 - \xi, \xi + 1 - k), & k + 2 \leq \xi \leq 2k. \end{cases}$$

$$r(a_{2\xi}|W) = \begin{cases} (\xi + 2, \xi - 1, \xi + 1, k + 2 - \xi), & 2 \leq \xi \leq k + 1; \\ (2k + 4 - \xi, 2k + 2 - \xi, 2k + 5 - \xi, \xi - k), & k + 2 \leq \xi \leq 2k + 1. \end{cases}$$

Representations of the middle cycle are

$$r(b_{2\xi+1}|W) = \begin{cases} (1, 2, 2, k+3), & \xi = 0; \\ (\xi+1, \xi+1, \xi, k+3-\xi), & 1 \leq \xi \leq k-2; \\ (\xi+1, \xi+1\xi, , 3), & \xi = k-1; \\ (\xi+1, \xi+1, \xi, 1), & \xi = k; \\ (\xi, \xi+1, \xi, 1), & \xi = k+1; \\ (\xi-2, \xi-1, \xi-1, 3), & \xi = k+2; \\ (2k+2-\xi, 2k+3-\xi, 2k+3-\xi, \xi+2-k), & k+3 \leq \xi \leq 2k. \end{cases}$$

$$r(b_{2\xi}|W) = \begin{cases} (2, 1, 2, k+2), & \xi = 1; \\ (\xi+1, \xi, \xi, k+3-\xi), & 2 \leq \xi \leq k-1; \\ (\xi+1, \xi, \xi, 2), & \xi = k; \\ (\xi-1, \xi-1, \xi, 2), & \xi = k+2; \\ (2k+3-\xi, 2k+3-\xi, 2k+4-\xi, \xi+1-k), & k+3 \leq \xi \leq 2k+1. \end{cases}$$

Representations of the outer cycle are

$$r(c_{2\xi+1}|W) = \begin{cases} (2, 3, 3, k+4), & \xi = 0; \\ (\xi+2, \xi+2, \xi+1, k+4-\xi), & 1 \leq \xi \leq k-2; \\ (\xi+2, \xi+2, \xi+1, 4), & \xi = k-1; \\ (\xi+2, \xi+2, \xi+1, 2), & \xi = k; \\ (\xi+1, \xi+2, \xi+1, 2), & \xi = k+1; \\ (\xi-1, \xi, \xi, 4), & \xi = k+2; \\ (2k+3-\xi, 2k+4-\xi, 2k+4-\xi, \xi+3-k), & k+3 \leq \xi \leq 2k. \end{cases}$$

$$r(c_{2\xi}|W) = \begin{cases} (3, 2, 3, k+3), & \xi = 1; \\ (\xi+2, \xi+1, \xi+1, k+4-\xi), & 2 \leq \xi \leq k-1; \\ (\xi+2, \xi+1, \xi+1, 3), & \xi = k; \\ (\xi+2, \xi+1, \xi+1, 1), & \xi = k+1; \\ (\xi, \xi, \xi+1, 3), & \xi = k+2; \\ (2k+4-\xi, 2k+4-\xi, 2k+5-\xi, \xi+2-k), & k+3 \leq \xi \leq 2k+1. \end{cases}$$

From the above discussion it follows that $\dim((G) = 4$ in this case.

Case 4: $n \equiv 3 \pmod{4}$

For $n = 4k+3$ with $k \geq 1$ and $k \in \mathbb{Z}^+$. When $n = 7$ then $W = \{a_1, a_2, b_4\}$ is the resolving set for $V(G)$. In $V(G)$ the representations of the vertices are

$$\begin{aligned} r(a_4|W) &= (2, 1, 3, 1), r(a_5|W) = (2, 2, 1, 2), r(a_6|W) = (1, 2, 2, 2), r(a_7|W) = \\ &= (3, 1, 2, 3), r(b_1|W) = (1, 2, 2, 3), r(b_2|W) = (2, 1, 2, 2), r(b_3|W) = (2, 2, 1, 1), r(b_5|W) = \\ &= (3, 3, 2, 1), r(b_6|W) = (2, 3, 3, 2), r(b_7|W) = (2, 2, 3, 3), r(c_1|W) = (2, 3, 3, 4), \\ r(c_2|W) &= (3, 2, 3, 3), r(c_3|W) = (3, 3, 2, 2), r(c_4|W) = (4, 3, 3, 1), r(c_5|W) = \\ &= (4, 4, 3, 2), r(c_6|W) = (3, 4, 4, 3) \text{ and } r(c_7|W) = (3, 3, 4, 4). \end{aligned}$$

When $n = 11$ then $W = \{a_1, a_2, b_6\}$ is the resolving set for $V(G)$. In $V(G)$ the representations of the vertices are

$$\begin{aligned} r(a_4|W) &= (4, 1, 3, 2), r(a_5|W) = (2, 4, 1, 2), r(a_6|W) = (3, 2, 4, 1), r(a_7|W) = \\ &= (3, 3, 2, 2), r(a_8|W) = (2, 3, 3, 2), r(a_9|W) = (4, 2, 3, 3), r(a_{10}|W) = (1, 4, 2, 3), \end{aligned}$$

$r(a_{11}|W) = (3, 1, 4, 4)$, $r(b_1|W) = (1, 2, 2, 5)$, $r(b_2|W) = (2, 1, 2, 4)$, $r(b_3|W) = (2, 2, 1, 3)$, $r(b_4|W) = (3, 2, 2, 2)$, $r(b_5|W) = (3, 3, 2, 1)$, $r(b_7|W) = (4, 4, 3, 1)$, $r(b_8|W) = (3, 4, 4, 2)$, $r(b_9|W) = (3, 3, 4, 3)$, $r(b_{10}|W) = (2, 3, 3, 4)$, $r(b_{11}|W) = (2, 2, 3, 5)$, $r(c_1|W) = (2, 3, 3, 6)$, $r(c_2|W) = (3, 2, 3, 5)$, $r(c_3|W) = (3, 3, 2, 4)$, $r(c_4|W) = (4, 3, 3, 3)$, $r(c_5|W) = (4, 4, 3, 2)$, $r(c_6|W) = (5, 4, 4, 1)$, $r(c_7|W) = (5, 5, 4, 2)$, $r(c_8|W) = (4, 5, 5, 3)$, $r(c_9|W) = (4, 4, 5, 4)$, $r(c_{10}|W) = (3, 4, 4, 5)$ and $r(c_{11}|W) = (3, 3, 4, 6)$.

When $n \geq 15$ then in this case, $\{a_1, a_2, b_{2k+2}\}$ resolves $V(G)$. Indeed, a_1 and a_2 distinguish the vertices in the inner cycle, inner pendent cycle, outer pendent cycle and outer cycle. To show that $\{a_1, a_2, b_{2k+2}\}$ resolves vertices of G , first we give representations of the vertices in G with respect to W .

Representations of the inner cycle are

$$r(a_{2\xi+1}|W) = \begin{cases} (\xi, \xi + 2, \xi - 1, k + 2 - \xi), & 2 \leq \xi \leq k; \\ (\xi, \xi, \xi - 1, 2), & \xi = k + 1; \\ (2k + 4 - \xi, 2k + 2 - \xi, 2k + 4 - \xi, \xi + 1 - k), & \xi = k + 2; \\ (2k + 4 - \xi, 2k + 2 - \xi, \xi + 1 - k), & k + 3 \leq \xi \leq 2k. \end{cases}$$

$$r(a_{2\xi}|W) = \begin{cases} (\xi + 2, \xi - 1, \xi + 1, k + 2 - \xi), & 2 \leq \xi \leq k; \\ (\xi, \xi - 1, \xi, 1), & \xi = k + 1; \\ (\xi - 2, \xi - 1, \xi - 1, 2), & \xi = k + 2; \\ (2k + 2 - \xi, 2k + 5 - \xi, 2k + 3 - \xi, \xi - k), & k + 3 \leq \xi \leq 2k + 1. \end{cases}$$

Representations of the outer pendent cycle are

$$r(b_{2\xi+1}|W) = \begin{cases} (1, 2, 2, k + 3), & \xi = 0; \\ (\xi + 1, \xi + 1, \xi, k + 3 - \xi), & 1 \leq \xi \leq k - 1; \\ (\xi + 1, \xi + 1, \xi, 1), & k \leq \xi \leq k + 1; \\ (\xi - 1, \xi - 1, \xi, 3), & \xi = k + 2; \\ (2k + 3 - \xi, 2k + 3 - \xi, 2k + 4 - \xi, \xi + 2 - k), & k + 3 \leq \xi \leq 2k + 1. \end{cases}$$

$$r(b_{2\xi}|W) = \begin{cases} (2, 1, 2, k + 2), & \xi = 1; \\ (\xi + 1, \xi, \xi, k + 3 - \xi), & 2 \leq \xi \leq k - 1; \\ (\xi + 1, \xi, \xi, 2), & \xi = k; \\ (\xi - 1, \xi, \xi, 2), & \xi = k + 2; \\ (2k + 3 - \xi, 2k + 4 - \xi, 2k + 4 - \xi, \xi + 1 - k), & k + 3 \leq \xi \leq 2k + 1. \end{cases}$$

Representations of the outer cycle are

$$r(c_{2\xi+1}|W) = \begin{cases} (2, 3, 3, k + 4), & \xi = 0; \\ (\xi + 2, \xi + 2, \xi + 1, k + 4 - \xi), & 1 \leq \xi \leq k - 1; \\ (\xi + 2, \xi + 2, \xi + 1, 2), & k \leq \xi \leq k + 1; \\ (\xi, \xi, \xi + 1, 4), & \xi = k + 2; \\ (2k + 4 - \xi, 2k + 4 - \xi, 2k + 5 - \xi, \xi + 3 - k), & k + 3 \leq \xi \leq 2k + 1. \end{cases}$$

$$r(c_{2\xi}|W) = \begin{cases} (3, 2, 3, k+3), & \xi = 1; \\ (\xi + 2, \xi + 1, \xi + 1, k + 4 - \xi), & 2 \leq \xi \leq k - 1; \\ (\xi + 2, \xi + 1, \xi + 1, 3), & \xi = k; \\ (\xi + 2, \xi + 1, \xi + 1, 1), & \xi = k + 1; \\ (\xi, \xi + 1, \xi + 1, 3), & \xi = k + 2; \\ (2k + 4 - \xi, 2k + 5 - \xi, 2k + 5 - \xi, \xi + 2 - k), & k + 3 \leq \xi \leq 2k + 1. \end{cases}$$

From the above discussion it follows that $\dim((G)) = 4$ in this case.

3 Relation between $\beta(G)$ and $\hat{\beta}(G)$

In this section we prove that fault-tolerant metric dimension is bounded by a function of metric dimension (independent of the graph). As usual, $N(v)$ and $N[v]$ denote vertex v s open and closed neighborhoods, respectively.

Lemma 2. *Let W be a resolving set of G . For each vertex $w \in W$, let $T(w) = \{x \in V(G) : N(w) \subseteq N(x)\}$. Then $\hat{W} = \bigcup_{w \in W} (N[w] \cup T(w))$ is a fault-tolerant resolving set of G .*

Proof. Consider a vertex $w \in \hat{W}$. If $w \notin W$ then $\hat{W} \setminus \{w\}$ resolves G since $W \subseteq \hat{W} \setminus \{w\}$. Now assume that $w \in W$.

Let m and n be distinct vertices of G . We must show that m and n are resolved by some vertex in $\hat{W} \setminus \{w\}$. If not, then w must resolve m and n since W resolves m and n . Without loss of generality $d(w, m) = d(w, n) - 1$.

First suppose that $m \neq w$. Let x be the neighbour of w on a shortest path between w and m . Then $x \in \hat{W} \setminus \{w\}$ and $d(w, m) = d(x, m) + 1$. Thus $d(x, m) + 1 \leq d(w, n) - 1$. Now $d(x, n) + 1 \leq d(x, n)$. Hence $d(x, m) + 1 \leq d(w, n)$. Thus x resolves m and n .

Now assume that $m = w$. If $n \notin \hat{W}$ then $n \in \hat{W} \setminus \{w\}$ and n resolves m and n . Otherwise $d(w, n) \geq 2$ and n is not adjacent to some neighbour x of w . Then $d(w, x) = 1$ and $d(n, x) \geq 2$. Thus $x \in \hat{W} \setminus \{w\}$ resolves $w (= m)$ and n . Hence \hat{W} is a fault-tolerant resolving set of G .

Lemma 3. *Let W be a resolving set in a graph G . Then for each vertex $w \in \hat{W}$, the number of vertices of G at distance at most k from w is at most $1 + k(2k + 1)^{|W|-1}$.*

Proof. Let $1 \leq d(w, x) \leq k$. For every vertex $u \in \hat{W}$ with $u \neq w$, we have $|d(x, u) - d(x, w)| \leq k$. Thus there are $2k + 1$ possible values for $d(x, u)$ and there are at most k possible values for $d(x, w)$. Thus the vector of distances from x to W has $1 + k(2k + 1)^{|W|-1}$ possible values. The result follows, since the vertices at distance at most k are resolved by W .

Theorem 2. *Fault-tolerant metric dimension is bounded by a function of the metric dimension (independent of the graph). In particular, $\hat{\beta}(G) \leq \beta(G)(1 + 2.5^{\beta(G)-1})$ for every graph G .*

Proof. Let W be a metric basis for a graph G . Lemma 3 with $k = 2$ implies that $|N[w] \cup T(w)| \leq 1 + 2.5^{\beta(G)-1}$ for each vertex $w \in W$. Thus $\hat{\beta}(G) \leq \beta(G)(1 + 2.5^{\beta(G)-1})$.

4 Conclusion

The concept of resolving set and that of metric dimension is defined in monograph. Graphs are special examples of metric spaces with their intrinsic path metric. In this paper the metric dimension of the G , graph has been studied and proved that graph has a constant metric dimension 3.

References

1. J. Caceres, C. Hernando, M. Mora, I. M. Pelayo, M. L. Puertas, C. Seara, D. R. Wood, On the metric dimension of some families of graphs, *Electronic Notes in Disc. Math.*, 22(2005), 129 – 133.
2. J. Caceres, C. Hernando, M. Mora, I. M. Pelayo, M. L. Puertas, C. Seara, D. R. Wood, On the metric dimension of Cartesian product of graphs, *SIAM J. Disc. Math.*, 2(21), (2007), 423 – 441.
3. G. Chartrand, L. Eroh, M. A. Johnson, O. R. Oellermann, Resolvability in graphs and metric dimension of a graph, *Disc. Appl. Math.*, 105(2000), 99 – 113.
4. F. Harary, R. A. Melter, On the metric dimension of a graph, *Ars Combin.*, 2(1976), 191 – 195.
5. M. Imran, A. Q. Baig, A. Ahmad, Families of plane graphs with constant metric dimension, *Utilitas Math.*, 88(2012), 43 – 57.
6. M. Imran, A. Q. Baig, M. K. Shafiq, I. Tomescu, On metric dimension of generalized Petersen graphs $P(n, 3)$, to appear in *Ars Combin.*
7. I. Javaid, M. T. Rahim, K. Ali, Families of regular graphs with constant metric dimension, *Utilitas Math.*, 75(2008), 21 – 33.
8. S. Khuller, B. Raghavachari, A. Rosenfeld, Localization in graphs, Technical Report CS-TR-3326, University of Maryland at College Park, 1994.
9. R. A. Melter, I. Tomescu, Metric bases in digital geometry, *Computer Vision, Graphics, and Image Processing*, 25(1984), 113 – 121.
10. P. J. Slater, Leaves of trees, *Congress. Numer.*, 14(1975), 549 – 559.
11. I. Tomescu, I. Javaid, On the metric dimension of the Jahangir graph, *Bull. Math. Soc. Sci. Math. Roum.*, 50(98), 4(2007), 371 – 376.
12. I. Tomescu, M. Imran, On metric and partition dimensions of some infinite regular graphs, *Bull. Math. Soc. Sci. Math. Roum.*, 52(100), 4(2009), 461 – 472.