

MONOCHROMATIC SCHUR TRIPLES IN RANDOMLY PERTURBED DENSE SETS OF INTEGERS

ELAD AIGNER-HOREV AND YURY PERSON

ABSTRACT. Given a dense subset A of the first n positive integers, we provide a short proof showing that for $p = \omega(n^{-2/3})$ the so-called *randomly perturbed* set $A \cup [n]_p$ a.a.s. has the property that any 2-colouring of it has a monochromatic Schur triple, i.e. a triple of the form $(a, b, a+b)$. This result is optimal since there are dense sets A , for which $A \cup [n]_p$ does not possess this property for $p = o(n^{-2/3})$.

§1. INTRODUCTION

The model of randomly perturbed graphs was introduced by Bohman, Frieze and Martin in [5], where they considered for which $p = p(n)$ a Hamilton cycle appears a.a.s. if one studies the model $G_\alpha \cup G(n, p)$, where G_α is a graph on n vertices with minimum degree $\delta(G) \geq \alpha n$ (for a fixed $\alpha > 0$) and $G(n, p)$ is the usual binomial random graph, i.e. each of the $\binom{n}{2}$ possible edges appears independently with probability p . Their result was a discovery of a phenomenon, that it suffices to take $p = C(\alpha)/n$, which is lower by a log-term needed for the appearance in the random graph alone, see e.g. [6]. Further properties were studied in [4, 17, 25] and, even more recently, various spanning structures in randomly perturbed graphs and hypergraphs were investigated in [1, 2, 3, 7, 8, 10, 13, 15, 16, 19].

The study of Ramsey properties in random graphs was initiated by Łuczak, Ruciński and Voigt [18] and the so-called symmetric edge Ramsey problem was settled completely by Rödl and Ruciński in a series of papers [20, 21, 22]. Ramsey properties in randomly perturbed graphs were investigated first by Krivelevich, Sudakov, and Tetali [17], who proved that whenever one perturbs a sufficiently large n -vertex dense graph G by adding to it $\omega(n^{2-2/(t-1)})$ edges chosen uniformly at random then the resulting graph almost surely possesses the property that any 2-colouring of its edges admits either a monochromatic triangle (in the first colour) or a monochromatic copy of K_t (in the second colour), where here $t \geq 3$. Generally, one writes $G \rightarrow (K_3, K_t)$ to denote this fact (and shortens it to $G \rightarrow (K_t)_r$ in the symmetric case and r colors). Moreover, the result of [17] is asymptotically best possible in terms of the number of random edges added.

In this short note we will study randomly perturbed dense sets of integers. More precisely, we denote by $[n]_p$ the model where each of the integers

YP is supported by the Carl Zeiss Foundation.

from $[n] := \{1, 2, \dots, n\}$ is chosen independently with probability $p = p(n)$. The model $[n]_p$ itself was thoroughly investigated with respect to extremal properties (e.g. Szemerédi's theorem [9, 14, 23]) and Ramsey-type properties (partition regularity of equations [11, 12]).

Given a dense set $A \subseteq [n]$, i.e. with at least αn elements for some fixed positive α , we will study for which $p = p(n)$, the set $A \cup [n]_p$ a.a.s. satisfies the property that no matter how one colours $A \cup [n]_p$ with two colours, there will always be a monochromatic Schur triple, i.e. a triple of integers x, y, z with $x + y = z$. An old result of Schur [24] states that if \mathbb{N} is finitely coloured, then there is always such a monochromatic triple. Moreover, the threshold $\Theta(n^{-1/2})$ for the random set $[n]_p$ was determined by Graham, Rödl and Ruciński [12] (for two colours) and by Friedgut, Rödl and Schacht [11] (for any constant number of colours). Our main result on Schur triples in the model $A \cup [n]_p$ and two colours is as follows.

THEOREM 1. *For every $\alpha > 0$ and every $p = \omega(n^{-2/3})$ it holds that whenever $A \subseteq [n]$ has $|A| \geq \alpha n$ then a.a.s. $A \cup [n]_p$ has the property that in any 2-colouring of it a monochromatic Schur triple appears.*

Let us say a few words about the nature of the threshold $\omega(n^{-2/3})$. The point is that, above $n^{-2/3}$, the set $A \cup [n]_p$ contains a.a.s. some 10-element set, which, no matter how coloured by red and blue, always contains a monochromatic Schur triple. A similar situation occurs for $G_\alpha \cup G(n, p) \rightarrow (K_3)_2$: it was shown in [4] that the threshold for the appearance of K_6 is then $p = \omega(1/n)$ and noted in [17] that, since $K_6 \rightarrow (K_3)_2$, this is already the threshold.

The condition $p = \omega(n^{-2/3})$ from Theorem 1 is asymptotically best possible. The expected number of Schur triples in the random set $[n]_p$ is $O(p^3 n^2)$; this quantity vanishes whenever $p = o(n^{-2/3})$. A typical set at this density is a.a.s. *sum-free then* (i.e., has no Schur triples). This in turn implies that the perturbation of $[n/2 + 1, n]$, namely $[n/2 + 1, n] \cup [n]_p$, a.a.s. admits a 2-colouring with no monochromatic Schur triples. Indeed, one may colour the (dense) sum-free set $[n/2 + 1, n]$ with one colour and all elements (not already in $[n/2 + 1, n]$) coming from $[n]_p$ with the complementary colour. Moreover, for $r > 2$ colours the colouring in which the set $A := [n/2 + 1, 1]$ is coloured with one colour and $[n]_p$ with the remaining $r - 1$ shows that, asymptotically, one does not gain on probability since the threshold for $[n]_p$ having the property that any $(r - 1)$ -colouring (for any constant $r \geq 3$) of it admits a monochromatic Schur triple is $\Theta(n^{-1/2})$ [11, 12].

§2. PROOF OF THEOREM 1

We refer to a set of integers as *2-Schur-Ramsey* if any 2-colouring of it admits a monochromatic Schur triple. For $x, y, d \in [n]$ satisfying $y > x + d$ and $x > d$ we define the 10-tuple $\mathcal{L}(x, y, d) \in \mathbb{N}^{10}$ as follows

$$(2) \quad \mathcal{L}(x, y, d) := (x - d, x, x + d, x + 2d, y - d, y, y + d, d, y - x - d, y - x).$$

Its first four entries form a 4-term arithmetic progression (4AP, hereafter), the next three form a 3-term arithmetic progression (3AP, hereafter), and its last three points form a Schur triple denoted $\mathcal{S}(x, y, d) := (d, y - x - d, y - x)$.

LEMMA 3. *Let $n \geq 1$ be an integer and let $x, y, d \in [n]$ satisfy $y > x + d$, $x > d$. Then $\mathcal{L}(x, y, d)$ is 2-Schur-Ramsey.*

Proof. The proof proceeds via a simple case analysis. Fix a 2-colouring of $\mathcal{L}(x, y, d)$ and assume towards a contradiction that it admits no monochromatic Schur triple. We may assume that x, y , and d are not coloured in the same colour; for otherwise the assumption of no monochromatic Schur triple would force a contradiction in the shape of the Schur triple $(y - x, x - d, y - d)$ all coloured with the complimentary colour to the one used for x, y , and d .

Nevertheless, two of x, y , and d are coloured using the same colour. We consider the three possible cases arising here. Suppose, firstly, that y and d are coloured, say, red, and that x is coloured blue. We may assume that $y - d$ and $y + d$ are both coloured blue in this setting. Then the Schur triple $(y - x - d, x, y - d)$ forces $y - x - d$ to be red and consequently $y - x$ is blue owing to $(y - x - d, d, y - x)$. The triple $(y - x, x - d, y - d)$ mandates that $x - d$ is red and then one concludes that $x + d$ is blue due to $(y - x - d, x + d, y)$. At this point one notices that $(y - x, x + d, y + d)$ is a blue Schur triple, a contradiction.

Suppose, secondly, that x and d are coloured, say, red and that y is coloured blue. We may assume that $x - d$ and $x + d$ are both coloured blue. The triple $(y - x - d, x + d, y)$ implies that $y - x - d$ is red and then $y - x$ is blue due to $(d, y - x - d, y - x)$. Consequently, $y - d$ is red owing to $(y - x, x - d, y - d)$ and then one arrives at $(y - x - d, x, y - d)$ as a red Schur triple, a contradiction.

Suppose, thirdly (and finally), that x, y are coloured, say, blue and that d is coloured red. Due to $(y - x, x, y)$ we may assume that $y - x$ is coloured red. Next, we have that $y - x - d$ is blue due to $(d, y - x - d, y - x)$, that $y - d$ is red due to $(y - x - d, x, y - d)$, that $x + d$ is red due to $(y - x - d, x + d, y)$, that $y + d$ is blue due to $(y - x, x + d, y + d)$, that $x - d$ is blue due to $(y - x, x - d, y - d)$, and that $x + 2d$ is blue due to $(x + d, d, x + 2d)$. All this ends with $(y - x - d, x + 2d, y + d)$ being a blue Schur triple; contradiction. ■

Facilitating our proof is the following result due to Varnavides [26]; which roughly speaking asserts that dense sets in $[n]$ have $\Omega(n^2)$ 4APs. Originally Varnavides' result was phrased for 3APs, but a standard supersaturation argument generalizes to arithmetic progressions of any fixed length.

THEOREM 4. (Varnavides [26]) *For every $\alpha > 0$ there exists an $n_0 := n_0(\alpha)$ and a $g(\alpha) > 0$ such that whenever $n \geq n_0$ and $A \subseteq [n]$ satisfy $|A| \geq \alpha n$ then A contains at least $g(\alpha)n^2$ 4APs.*

Dense sets in $[n]$ have an abundance of 4APs then. This in turn implies that there are elements in $[n]$ that are repeatedly being used by 4APs found in the dense set as their common steps; in fact there are many such *popular common steps*. Making this precise, define for a set $A \subseteq [n]$ and a parameter $\varepsilon > 0$ the set

$$\mathcal{D}_\varepsilon(A) := \{d \in [n] : A \text{ has } \geq \varepsilon n \text{ 4APs with common step } d\}$$

to be the set of common steps *popular* in the sense that each of these participates in at least εn 4APs (in A).

LEMMA 5. (Linearly many popular common steps) *For every $\alpha > 0$ there exists an $\varepsilon := \varepsilon(\alpha) > 0$ such that for n sufficiently large $|\mathcal{D}_\varepsilon(A)| \geq \varepsilon n$ holds whenever $A \subseteq [n]$ satisfies $|A| \geq \alpha n$.*

Proof. Given α , let $g(\alpha)$ be as asserted by Theorem 4. Set $\varepsilon := g(\alpha)/2$ and let $\#4AP(A)$ denote the number of 4APs in A . By Theorem 4, the set A has at least $g(\alpha)n^2$ 4APs whenever n is sufficiently large. Then owing to

$$g(\alpha)n^2 \leq \#4AP(A) < |\mathcal{D}_\varepsilon(A)|n + n \cdot \varepsilon n$$

the lemma follows. ■

We are now ready to prove Theorem 1.

PROOF OF THEOREM 1. By Lemma 3 it suffices to show that a.a.s. $A \cup [n]_p$ contains $\mathcal{L}(x, y, d)$ for some $x, y, d \in [n]$ (with $y > x + d$ and $x > d$). By Theorem 4 the set A contains at least $g(\alpha)n^2$ 4APs. Moreover, Lemma 5 asserts that there is an $\varepsilon := \varepsilon(\alpha)$ such that $|\mathcal{D}_\varepsilon(A)| \geq \varepsilon n$. Given $d \in \mathcal{D}_\varepsilon(A)$ and a pair of *distinct* 4APs both with common step d , we may write them parametrically as $(x - d, x, x + d, x + 2d)$ and $(y' - d, y', y' + d, y' + 2d)$ with $y' > x$. Setting $y := y' + d$, we see that the Schur triple $\mathcal{S}(x, y, d) = (d, y - x - d, y - x)$ is well-defined.

Let \mathcal{T} be the set of all such Schur triples $\mathcal{S}(x, y, d)$. Since there are at least εn choices for $d \in \mathcal{D}_\varepsilon(A)$ and A contains at least εn 4APs with common difference d , there are, for a fixed $d \in \mathcal{D}_\varepsilon(A)$, at least $\varepsilon n - 1$ different possible values for $y - x$. Therefore, the number of such Schur triples is at least $\varepsilon^2 n^2 / 4$, i.e. $|\mathcal{T}| \geq \varepsilon^2 n^2 / 4$.

It remains to argue that a.a.s. $[n]_p$ contains a member of \mathcal{T} . The expected number of members of \mathcal{T} captured by $[n]_p$ is at least $\varepsilon^2 p^3 n^2 / 4$. We write X_t for the indicator random variable indicating whether $t = \mathcal{S}(x, y, d) \in \mathcal{T}$ is captured by $[n]_p$ and then put $X := \sum_{t \in \mathcal{T}} X_t$. Using Chebyshev's inequality we arrive at:

$$\Pr[X = 0] \leq \frac{\mathbf{Var}(X)}{(\mathbf{E} X)^2} \leq \frac{1}{\mathbf{E}(X)} + \frac{\sum_{t \neq t' \in \mathcal{T}} \mathbf{E}[X_t X_{t'}]}{\mathbf{E}(X)^2}.$$

The number of pairs $t, t' \in \mathcal{T}$, $t \neq t'$, having a single entry in common is at most $3n^3$, and each such pair satisfies $\mathbf{E}[X_t X_{t'}] \leq p^5$ so in total all such pairs contribute $O(p^5 n^3)$ to the sum appearing on the r.h.s. above. Next,

the number of pairs $t, t' \in \mathcal{T}$, $t \neq t'$, having two entries in common is at most $2n^2$, with the total contribution of $O(n^2p^4)$. Using the lower bound on $\mathbf{E}(X) \geq \varepsilon^2 p^3 n^2 / 2$ and $p = \omega(n^{-2/3})$ we have

$$\Pr[X = 0] \leq \frac{1}{\Omega(p^3 n^2)} + \frac{O(p^5 n^3)}{\Omega(p^6 n^4)} + \frac{O(p^4 n^2)}{\Omega(p^6 n^4)} = o(1),$$

finishing the proof. ■

§3. CONCLUDING REMARKS

In this note we studied the first Ramsey-type question for randomly perturbed sets of integers, i.e. the model $A \cup [n]_p$. It would be interesting to determine the thresholds for more general partition regular systems of equations. For the threshold results in random sets see the work of Friedgut, Rödl and Schacht [11].

REFERENCES

1. J. Balogh, A. Treglown, and A. Z. Wagner, *Tilings in randomly perturbed dense graphs*, arXiv:1708.09243, 2017.
2. W. Bedenknecht, J. Han, Y. Kohayakawa, and G. O. Mota, *Powers of tight Hamilton cycles in random perturbed hypergraphs*, arXiv:1802.08900, 2018.
3. P. Bennett, A. Dudek, and A. M. Frieze, *Adding random edges to create the square of a Hamilton cycle*, arXiv:1710.02716, 2017.
4. T. Bohman, A. Frieze, M. Krivelevich, and R. Martin, *Adding random edges to dense graphs.*, Random Struct. Algorithms **24** (2004), no. 2, 105–117.
5. T. Bohman, A. Frieze, and R. Martin, *How many random edges make a dense graph Hamiltonian?*, Random Structures & Algorithms **22** (2003), no. 1, 33–42.
6. B. Bollobás, *Random graphs*, Modern Graph Theory, Springer, 1998.
7. J. Böttcher, J. Han, Y. Kohayakawa, R. Montgomery, O. Parczyk, and Y. Person, *Universality of bounded degree spanning trees in randomly perturbed graphs*, arXiv:1802.04707, 2018.
8. J. Böttcher, R. H. Montgomery, O. Parczyk, and Y. Person, *Embedding spanning bounded degree graphs in randomly perturbed graphs*, arXiv:1802.04603, 2018.
9. D. Conlon and W. T. Gowers, *Combinatorial theorems in sparse random sets.*, Ann. Math. (2) **184** (2016), no. 2, 367–454.
10. A. Dudek, C. Reiher, A. Ruciński, and M. Schacht, *Powers of Hamiltonian cycles in randomly augmented graphs*, arXiv preprint arXiv:1805.10676, 2018.
11. E. Friedgut, V. Rödl, and M. Schacht, *Ramsey properties of random discrete structures.*, Random Struct. Algorithms **37** (2010), no. 4, 407–436.
12. R. L. Graham, V. Rödl, and A. Ruciński, *On Schur properties of random subsets of integers.*, J. Number Theory **61** (1996), no. 2, 388–408.
13. J. Han and Y. Zhao, *Embedding Hamilton ℓ -cycles in randomly perturbed hypergraphs*, arXiv:1802.04586, 2018.
14. Y. Kohayakawa, T. Łuczak, and V. Rödl, *Arithmetic progressions of length three in subsets of a random set.*, Acta Arith. **75** (1996), no. 2, 133–163.
15. M. Krivelevich, M. Kwan, and B. Sudakov, *Cycles and matchings in randomly perturbed digraphs and hypergraphs*, Combinatorics, Probability and Computing **25** (2016), no. 6, 909–927.

16. ———, *Bounded-degree spanning trees in randomly perturbed graphs*, SIAM Journal on Discrete Mathematics **31** (2017), no. 1, 155–171.
17. M. Krivelevich, B. Sudakov, and P. Tetali, *On smoothed analysis in dense graphs and formulas*, Random Structures Algorithms **29** (2006), no. 2, 180–193.
18. T. Łuczak, A. Ruciński, and B. Voigt, *Ramsey properties of random graphs*, Journal of Combinatorial Theory, Series B **56** (1992), no. 1, 55–68.
19. A. McDowell and R. Mycroft, *Hamilton ℓ -cycles in randomly perturbed hypergraphs*, arXiv:1802.04242, 2018.
20. V. Rödl and A. Ruciński, *Lower bounds on probability thresholds for Ramsey properties*, Combinatorics, Paul Erdős is eighty, 1, Bolyai Society Mathematical Studies, vol. 1, János Bolyai Mathematical Society, 1993, pp. 317–346.
21. ———, *Random graphs with monochromatic triangles in every edge coloring*, Random Structures & Algorithms **5** (1994), no. 2, 253–270.
22. ———, *Threshold functions for Ramsey properties*, Journal of the American Mathematical Society **8** (1995), no. 4, 917–942.
23. M. Schacht, *Extremal results for random discrete structures.*, Ann. Math. (2) **184** (2016), no. 2, 333–365.
24. I. Schur, *Über die Kongruenz $x^m + y^m \equiv z^m \pmod{p}$.*, Jahresber. Dtsch. Math.-Ver. **25** (1916), 114–117.
25. B. Sudakov and J. Vondrák, *How many random edges make a dense hypergraph non-2-colorable?*, Random Structures & Algorithms **32** (2008), no. 3, 290–306.
26. P. Varnavides, *On certain sets of positive density*, J. London Math. Soc. **34** (1959), 358–360. MR 0106865

DEPARTMENT OF MATHEMATICS AND COMPUTER SCIENCE, ARIEL UNIVERSITY, ARIEL,
ISRAEL

E-mail address: horev@ariel.ac.il

INSTITUT FÜR MATHEMATIK, TECHNISCHE UNIVERSITÄT ILMENAU, 98684 ILMENAU,
GERMANY

E-mail address: yury.person@tu-ilmenau.de