

# LEAF-INDUCED SUBTREES OF LEAF-FIBONACCI TREES

AUDACE A. V. DOSSOU-OLORY

**ABSTRACT.** In analogy to a concept of Fibonacci trees, we define the leaf-Fibonacci tree of size  $n$  and investigate its number of nonisomorphic leaf-induced subtrees. Denote by  $f_0$  the one vertex tree and  $f_1$  the tree that consists of a root with two leaves attached to it; the leaf-Fibonacci tree  $f_n$  of size  $n \geq 2$  is the binary tree whose branches are  $f_{n-1}$  and  $f_{n-2}$ . We derive a nonlinear difference equation for the number  $N(f_n)$  of nonisomorphic leaf-induced subtrees (subtrees induced by leaves) of  $f_n$ , and also prove that  $N(f_n)$  is asymptotic to  $1.00001887227319 \dots (1.48369689570172 \dots)^\phi$  ( $\phi =$  golden ratio) as  $n$  grows to infinity.

## 1. INTRODUCTION

Fibonacci trees are an alternative approach to a binary search in computer science and information processing [10, p. 417]. The Fibonacci tree of order  $n$  is defined as the binary tree whose left branch is the Fibonacci tree of order  $n-1$  and right branch is the Fibonacci tree of order  $n-2$ , while the Fibonacci tree of order 0 or 1 is the tree with only one vertex [10]. We show in Figure 1 the Fibonacci tree of order 5.




FIGURE 1. The Fibonacci tree of order 5.

Thus, the Fibonacci tree of order  $n$  has precisely  $F_{n+1}$  leaves (so  $2F_{n+1} - 1$  vertices), where  $F_n$  denotes the  $n$ -th Fibonacci number:

$$F_0 = 0, F_1 = 1, F_n = F_{n-1} + F_{n-2} \text{ for } n > 1.$$

Fibonacci trees are also a special case of so-called AVL (“Adel’son-Vel’skii and Landis”—named after the inventors) trees [1]; these trees have the defining property that for every

---

2010 *Mathematics Subject Classification.* Primary 05C05; secondary 05C30.

*Key words and phrases.* leaf-induced subtrees, nonisomorphic subtrees, leaf-Fibonacci trees.

The author was supported by Stellenbosch University in association with African Institute for Mathematical Science (AIMS) South Africa.

internal vertex  $v$ , the heights (i.e., the greatest distance of a leaf from the root) of the left and right branches of the subtree rooted at  $v$  (consisting of  $v$  and all its descendants) differ by at most one. According to [1], AVL trees are the first data structure to be invented. Figure 2 shows an AVL tree of height 3. For more information on Fibonacci trees and their uses, we refer to [13, 8, 9, 7].




FIGURE 2. An AVL tree of height 3.

In analogy to the concept of Fibonacci trees from [10], we define the *leaf-Fibonacci tree of size (height)  $n$*  as follows:

- Denote by  $f_0$  the tree with only one vertex and  $f_1$  the tree that consists of a root with two leaves attached to it;
- For  $n \geq 2$ , connect the roots of the trees  $f_{n-1}$  and  $f_{n-2}$  to a new common vertex to obtain the tree  $f_n$ .

In other words, the leaf-Fibonacci tree  $f_n$  of size  $n \geq 2$  is the binary tree whose branches are the leaf-Fibonacci trees  $f_{n-1}$  and  $f_{n-2}$ . Hence,  $f_n$  has precisely  $F_{n+2}$  leaves, where  $F_n$  is the  $n$ -th Fibonacci number ( $F_1 = 1, F_2 = 1, F_3 = 2, F_4 = 3, F_5 = 5, F_6 = 8, F_7 = 13, \dots$ ). Figure 3 shows the leaf-Fibonacci tree of size 5.




FIGURE 3. The leaf-Fibonacci tree of size 5.

In this note, we shall be interested in the number of nonisomorphic subtrees induced by leaves (henceforth, *leaf-induced subtrees*) of a leaf-Fibonacci tree of size  $n$ .

Let  $T$  be a rooted tree without vertices of outdegree 1 (also known as *topological* or *series-reduced* or *homeomorphically irreducible* trees [3, 2, 5, 12]). Every choice of  $k$  leaves

of  $T$  induces another topological tree, which is obtained by extracting the minimal subtree of  $T$  that contains all the  $k$  leaves and suppressing (if any) all vertices of outdegree 1; see Figure 4 for an illustration. Every subtree obtained through this operation is sometimes




FIGURE 4. A topological tree (on the left) and the subtree induced by the leaves  $l_1, l_2, l_3, l_4$  (on the right).

referred to as a *leaf-induced subtree* [4, 5]. We note the study of subtrees induced by leaves of binary trees finds a noteworthy relevance in phylogenetics—see Semple and Steel’s book [11] which describes the mathematical foundations of phylogenetics.

Two rooted trees are said to be isomorphic if there is a graph isomorphism (preserving adjacency) between them that maps the root of one to the root of the other. It is important to note that the problem of enumerating leaf-induced subtrees becomes trivial if isomorphisms are not taken into account: in fact, it is clear that every topological tree with  $n$  leaves has exactly  $2^n - 1$  leaf-induced subtrees.

We mention that nonisomorphic leaf-induced subtrees of a topological tree have been studied only very recently: Wagner and the present author [6] obtained exact and asymptotic enumeration results on the number of nonisomorphic leaf-induced subtrees of two classes of  $d$ -ary trees, namely so-called  $d$ -ary caterpillars and even  $d$ -ary trees. In [6], they also derived extremal results for the number of root containing leaf-induced subtrees of a topological tree.

We shall denote the number of nonisomorphic leaf-induced subtrees of the leaf-Fibonacci tree  $f_n$  by  $N(f_n)$ . Our main results are a recurrence relation and an asymptotic formula for  $N(f_n)$ . As it turns out, the plan to compute  $N(f_n)$  will be to consider the number of root containing leaf-induced subtrees of  $f_n$ .

In [14], Wagner alone studied the number of independent vertex subsets (set of vertices containing no pair of adjacent vertices) of a Fibonacci tree of order  $n$  with the notable difference that in his context, the Fibonacci tree of order 0 has no vertices. Wagner

derived a system of recurrence relations for the number of independent vertex subsets of a Fibonacci tree of an arbitrary order  $n$ , and also proved that there are positive constants  $A, B > 0$  such that the number of independent vertex subsets of a Fibonacci tree of order  $n$  is asymptotic to  $A \cdot B^{F_n}$  as  $n$  grows to infinity. In the present study, we obtain a similar asymptotic formula for the number  $N(f_n)$  of nonisomorphic leaf-induced subtrees of the leaf-Fibonacci tree of size  $n$ : we demonstrate—not expectedly—that for some effectively computable constants  $A_1, A_2 > 0$ ,

$$N(f_n) \sim A_1 \cdot A_2^{F_n} \text{ as } n \rightarrow \infty.$$

## 2. MAIN RESULTS

We note from the recursive definition of the tree  $f_n$  that  $f_m$  is a leaf-induced subtree of  $f_n$  for every  $m \leq n$ . However, not every leaf-induced subtree of  $f_n$  is again a leaf-Fibonacci tree: in fact, by repeatedly removing leaves from  $f_n$ , one easily sees that  $f_n$  has leaf-induced subtrees of every number of leaves  $k$  between 1 and  $n$ .

As mentioned in the introduction, the plan to compute  $N(f_n)$  will be to consider the number of root containing leaf-induced subtrees of  $f_n$ .

**Lemma 1.** *All nonisomorphic leaf-induced subtrees with two or more leaves of  $f_n$  can be identified as containing the root of  $f_n$ .*

*Proof.* The tree  $f_0$  has only one vertex which is also its leaf and root, so the statement holds vacuously for  $n = 0$ . The statement is trivial for  $n = 1$  ( $f_1$  is the only leaf-induced subtree in this case). Let  $n > 1$  and consider a subset of  $k > 1$  leaves of  $f_n$ . We argue by double induction on  $n$  and  $k$ :

- If all  $k$  leaves belong to  $f_{n-1}$  then by the induction hypothesis on  $n$ , the induced subtree with  $k$  leaves contains the root of  $f_{n-1}$ . Moreover, by the induction hypothesis on  $k$ , the tree  $f_{n-1}$  can be identified as containing the root of  $f_n$  (as  $f_{n-1}$  is clearly a leaf-induced subtree of  $f_n$ ). Hence, the induced subtree with  $k$  leaves contains the root of  $f_n$ .
- If all  $k$  leaves belong to  $f_{n-2}$ , then we also deduce by the induction hypothesis that the induced subtree with  $k$  leaves is a root containing leaf-induced subtree of  $f_n$ .
- If  $k_1$  leaves belong to  $f_{n-1}$  and  $k - k_1$  leaves belong to  $f_{n-2}$ , then by the induction hypothesis, the induced subtrees with  $k_1$  and  $k - k_1$  leaves are root containing leaf-induced subtrees of  $f_{n-1}$  and  $f_{n-2}$ , respectively. Consequently, the root of the induced subtree with  $k$  leaves coincides with the root of  $f_n$ .

This completes the induction step as well as the proof of the lemma. □

We then obtain the following proposition:

**Proposition 2.** *The number  $N(f_n)$  of nonisomorphic leaf-induced subtrees of the leaf-Fibonacci tree  $f_n$  satisfies the following nonlinear recurrence relation:*

$$(1) \quad N(f_n) = 1 + \frac{1}{2}N(f_{n-2}) - \frac{1}{2}N(f_{n-2})^2 + N(f_{n-2}) \cdot N(f_{n-1})$$

with initial values  $N(f_0) = 1, N(f_1) = 2$ .

*Proof.* It is obvious that  $N(f_0) = 1$  and  $N(f_1) = 2$ . Let  $n > 1$ . By Lemma 1,  $N(f_n)$  is precisely one more the number of nonisomorphic root containing leaf-induced subtrees of  $f_n$  (the subtree with only one vertex has been included). Since all leaf-induced subtrees of the leaf-Fibonacci tree  $f_{n-2}$  are again leaf-induced subtrees of  $f_{n-1}$ , the nonisomorphic root containing leaf-induced subtrees of  $f_n$  can be categorised by two types of enumeration:

- Both branches of the induced subtree are leaf-induced subtrees of  $f_{n-2}$ . The total number of these possibilities is  $\binom{1+N(f_{n-2})}{2}$  as the induced subtrees have to be nonisomorphic.
- One of the branches of the induced subtree is a leaf-induced subtree of  $f_{n-2}$  while the other branch is a leaf-induced subtree of  $f_{n-1}$  but does not belong to the set of leaf-induced subtrees of  $f_{n-2}$ . The total number of these possibilities is  $N(f_{n-2})(N(f_{n-1}) - N(f_{n-2}))$ .

Therefore, we obtain

$$\begin{aligned} N(f_n) &= 1 + \binom{1 + N(f_{n-2})}{2} + N(f_{n-2})(N(f_{n-1}) - N(f_{n-2})) \\ &= 1 + \frac{1}{2}N(f_{n-2}) - \frac{1}{2}N(f_{n-2})^2 + N(f_{n-2}) \cdot N(f_{n-1}), \end{aligned}$$

which completes the proof of the proposition.  $\square$

The sequence  $(N(f_n))_{n \geq 0}$  starts as

$$N(f_0) = 1, N(f_1) = 2, N(f_2) = 3, N(f_3) = 6, N(f_4) = 16, N(f_5) = 82, N(f_6) = 1193, \\ N(f_7) = 94506, N(f_8) = 112034631, \dots$$

We remark that recursion (1) cannot be solved explicitly. Therefore, finding an asymptotic formula should be in order. In the following theorem, we show—not expectedly—that  $N(f_n)$  grows doubly exponentially in  $n$ .

**Theorem 3.** *There are two positive constants  $K_1, K_2 > 0$  (both solely depending on the first terms of  $(N(f_n))_{n \geq 0}$ ) such that*

$$N(f_n) = (1 + o(1))K_1 \cdot K_2^{\left(\frac{1+\sqrt{5}}{2}\right)^n}$$

as  $n \rightarrow \infty$ .

*Proof.* For ease of notation, set  $A_n := N(f_n)$ . Then we have

$$A_n = 1 + \frac{1}{2}A_{n-2} - \frac{1}{2}A_{n-2}^2 + A_{n-2} \cdot A_{n-1}$$

with initial values  $A_0 = 1, A_1 = 2$ . Since the sequence  $(A_n)_{n \geq 0}$  increases with  $n$ , it is not difficult to note that

$$A_n \geq \frac{1}{2}A_{n-1} \cdot A_{n-2}$$

for all  $n \geq 2$ . Also, since  $A_n \geq A_2 = 3$  for all  $n \geq 2$  and  $1 + A_1/2 - A_1^2/2 = 0$ , it is not difficult to see that

$$A_n \leq A_{n-1} \cdot A_{n-2}$$

for all  $n \geq 3$ . Thus, we have

$$(2) \quad \lim_{n \rightarrow \infty} \frac{A_{n-1}}{A_n} = 0,$$

which also implies that the sequence  $(A_{n-1}/A_n)_{n \geq 1}$  is bounded for every  $n \geq 1$ . Let us use  $Q_n$  as a shorthand for  $\log(A_n)$  and  $E_n$  as a shorthand for

$$(3) \quad \log \left( 1 + \frac{1}{2A_{n-1}} - \frac{A_{n-2}}{2A_{n-1}} + \frac{1}{A_{n-1} \cdot A_{n-2}} \right).$$

With these notations, we have

$$Q_n = Q_{n-1} + Q_{n-2} + E_n.$$

By setting  $R_{n-1} := Q_{n-2}$ , we obtain the following system (written in matrix form) of two linear difference equations:

$$\begin{pmatrix} Q_n \\ R_n \end{pmatrix} = \begin{pmatrix} 1 & 1 \\ 1 & 0 \end{pmatrix} \begin{pmatrix} Q_{n-1} \\ R_{n-1} \end{pmatrix} + \begin{pmatrix} E_n \\ 0 \end{pmatrix}.$$

By iteration on  $n$ , one gets

$$\begin{pmatrix} Q_n \\ R_n \end{pmatrix} = \begin{pmatrix} 1 & 1 \\ 1 & 0 \end{pmatrix}^{n-1} \begin{pmatrix} Q_1 \\ Q_0 \end{pmatrix} + \sum_{i=2}^n \begin{pmatrix} 1 & 1 \\ 1 & 0 \end{pmatrix}^{n-i} \begin{pmatrix} E_i \\ 0 \end{pmatrix}$$

for all  $n \geq 2$ , as  $R_1 = Q_0$ . The eigenvalue decomposition gives us

$$\begin{pmatrix} 1 & 1 \\ 1 & 0 \end{pmatrix} = \frac{1}{\lambda_1 - \lambda_2} \begin{pmatrix} \lambda_1 & \lambda_2 \\ 1 & 1 \end{pmatrix} \begin{pmatrix} \lambda_1 & 0 \\ 0 & \lambda_2 \end{pmatrix} \begin{pmatrix} 1 & -\lambda_2 \\ -1 & \lambda_1 \end{pmatrix}$$

with

$$\lambda_1 = \frac{1 - \sqrt{5}}{2} \quad \text{and} \quad \lambda_2 = \frac{1 + \sqrt{5}}{2}.$$

It follows that

$$\begin{aligned} \begin{pmatrix} 1 & 1 \\ 1 & 0 \end{pmatrix}^m &= \frac{1}{\lambda_1 - \lambda_2} \begin{pmatrix} \lambda_1 & \lambda_2 \\ 1 & 1 \end{pmatrix} \begin{pmatrix} \lambda_1^m & 0 \\ 0 & \lambda_2^m \end{pmatrix} \begin{pmatrix} 1 & -\lambda_2 \\ -1 & \lambda_1 \end{pmatrix} \\ &= \frac{1}{\lambda_1 - \lambda_2} \begin{pmatrix} \lambda_1^{m+1} - \lambda_2^{m+1} & \lambda_1 \cdot \lambda_2^{m+1} - \lambda_1^{m+1} \cdot \lambda_2 \\ \lambda_1^m - \lambda_2^m & \lambda_1 \cdot \lambda_2^m - \lambda_1^m \cdot \lambda_2 \end{pmatrix} \end{aligned}$$

for all integer values of  $m$ . Consequently, we have

$$\begin{pmatrix} Q_n \\ R_n \end{pmatrix} = \frac{\log(2)}{\lambda_1 - \lambda_2} \begin{pmatrix} \lambda_1^n - \lambda_2^n \\ \lambda_1^{n-1} - \lambda_2^{n-1} \end{pmatrix} + \sum_{i=2}^n \frac{E_i}{\lambda_1 - \lambda_2} \begin{pmatrix} \lambda_1^{n-i+1} - \lambda_2^{n-i+1} \\ \lambda_1^{n-i} - \lambda_2^{n-i} \end{pmatrix}$$

for all  $n \geq 2$  as  $Q_0 = 0$  and  $Q_1 = \log(2)$ . Therefore, we obtain

$$Q_n = \frac{\log(2)}{\lambda_2 - \lambda_1} (\lambda_2^n - \lambda_1^n) + \frac{1}{\lambda_2 - \lambda_1} \sum_{i=2}^n E_i (\lambda_2^{n-i+1} - \lambda_1^{n-i+1})$$

for all  $n \geq 2$ . Since the sequence  $(E_n)_{n \geq 2}$  is bounded for every  $n \geq 2$  (as  $\lim_{n \rightarrow \infty} E_n = 0$  by virtue of (3) and (2)), we derive that

$$\sum_{i=2}^n |E_i| \cdot |\lambda_1|^{n-i+1} \leq \frac{|\lambda_1|^n - |\lambda_1|}{|\lambda_1| - 1} \cdot \sup_{2 \leq m \leq n} |E_m|$$

for all  $n \geq 2$ . This implies that the quantity

$$\frac{1}{\lambda_2 - \lambda_1} \sum_{i=2}^n E_i \cdot \lambda_1^{n-i+1}$$

converges to a definite limit as  $n \rightarrow \infty$  (note that  $|\lambda_1| < 1$  and  $\sup_{2 \leq m \leq n} |E_m|$  is finite for every  $n \geq 2$ ). On the other hand, we have

$$0 \leq \left| \sum_{i=n+1}^{\infty} E_i \cdot \lambda_2^{n-i+1} \right| \leq \frac{\lambda_2}{\lambda_2 - 1} \cdot \sup_{m \geq n+1} |E_m|$$

for all  $n \geq 2$  (note that  $\lambda_2 > 1$ ), which implies that

$$\frac{1}{\lambda_2 - \lambda_1} \sum_{i=n+1}^{\infty} E_i \cdot \lambda_2^{n-i+1} = \mathcal{O}\left(\sup_{m \geq n+1} |E_m|\right) = o(1)$$

as  $n \rightarrow \infty$ . Putting everything together, we arrive at

$$\begin{aligned} Q_n &= \frac{\lambda_2^n}{\lambda_2 - \lambda_1} \left( \log(2) + \sum_{i=2}^{\infty} E_i \cdot \lambda_2^{-i+1} \right) - \frac{1}{\lambda_2 - \lambda_1} \sum_{i=2}^n E_i \cdot \lambda_1^{n-i+1} \\ &\quad + \mathcal{O}\left(\sup_{m \geq n+1} |E_m|\right) + \mathcal{O}(\lambda_1^n) \\ &= \frac{\lambda_2^n}{\lambda_2 - \lambda_1} \left( \log(2) + \sum_{i=2}^{\infty} E_i \cdot \lambda_2^{-i+1} \right) - \frac{1}{\lambda_2 - \lambda_1} \sum_{i=2}^n E_i \cdot \lambda_1^{n-i+1} + o(1) \end{aligned}$$

as  $n \rightarrow \infty$ . We deduce that

$$\begin{aligned} A_n &= \left( 1 + \mathcal{O}\left(\lambda_1^n + \sup_{m \geq n+1} |E_m|\right) \right) \\ &\quad \cdot \exp\left( \frac{\lambda_2^n}{\lambda_2 - \lambda_1} \left( \log(2) + \sum_{i=2}^{\infty} E_i \cdot \lambda_2^{-i+1} \right) - \frac{1}{\lambda_2 - \lambda_1} \sum_{i=2}^n E_i \cdot \lambda_1^{n-i+1} \right) \\ &= (1 + o(1)) \cdot \exp\left( \frac{\lambda_2^n}{\lambda_2 - \lambda_1} \left( \log(2) + \sum_{i=2}^{\infty} E_i \cdot \lambda_2^{-i+1} \right) - \frac{1}{\lambda_2 - \lambda_1} \sum_{i=2}^n E_i \cdot \lambda_1^{n-i+1} \right) \end{aligned}$$

as  $n \rightarrow \infty$ . Call  $K_2$  the quantity

$$\exp\left(\frac{1}{\lambda_2 - \lambda_1}\left(\log(2) + \sum_{i=2}^{\infty} E_i \cdot \lambda_2^{-i+1}\right)\right),$$

and  $K_1$  the quantity

$$\exp\left(-\frac{1}{\lambda_2 - \lambda_1} \cdot \lim_{n \rightarrow \infty} \left(\sum_{i=2}^n E_i \cdot \lambda_1^{n-i+1}\right)\right).$$

Thus,

$$A_n = N(f_n) = (1 + o(1))K_1 \cdot K_2^{\lambda_2^n} = (1 + o(1))K_1 \cdot K_2^{\left(\frac{1+\sqrt{5}}{2}\right)^n}$$

as  $n \rightarrow \infty$ , where  $K_1$  and  $K_2$  can now be written as

$$K_2 = \exp\left(\frac{1}{\sqrt{5}}\left(\log(2) + \sum_{i=2}^{\infty} \left(\frac{1+\sqrt{5}}{2}\right)^{-i+1} \cdot \log\left(1 + \frac{1}{2N(f_{i-1})} - \frac{N(f_{i-2})}{2N(f_{i-1})} + \frac{1}{N(f_{i-1}) \cdot N(f_{i-2})}\right)\right)\right)$$

and

$$K_1 = \exp\left(-\frac{1}{\sqrt{5}} \cdot \lim_{n \rightarrow \infty} \left(\sum_{i=2}^n \left(\frac{1-\sqrt{5}}{2}\right)^{n-i+1} \cdot \log\left(1 + \frac{1}{2N(f_{i-1})} - \frac{N(f_{i-2})}{2N(f_{i-1})} + \frac{1}{N(f_{i-1}) \cdot N(f_{i-2})}\right)\right)\right).$$

By (numerically) evaluating  $K_1$  and  $K_2$ , we obtain that the number of nonisomorphic leaf-induced subtrees of the leaf-Fibonacci tree  $f_n$  is asymptotically

$$1.00001887227319 \dots (1.48369689570172 \dots)^{\left(\frac{1+\sqrt{5}}{2}\right)^n}$$

as  $n \rightarrow \infty$ . This completes the proof of the theorem.  $\square$

This asymptotic formula can also be written in terms of the Fibonacci number  $F_n$ : the number of leaves of  $f_n$  is given by

$$|f_n| = F_{n+2} = \frac{1}{\sqrt{5}} \left( \left(\frac{1+\sqrt{5}}{2}\right)^{2+n} - \left(\frac{1-\sqrt{5}}{2}\right)^{2+n} \right)$$

for every  $n$ ; so we deduce that

$$\frac{10}{5 + 3\sqrt{5}} \cdot |f_n| \sim \left(\frac{1+\sqrt{5}}{2}\right)^n$$



as  $n \rightarrow \infty$ . This implies that

$$\begin{aligned} N(f_n) &\sim K_1 \cdot K_2^{\frac{10}{5+3\sqrt{5}} \cdot |f_n|} \\ &= 1.00001887227319 \cdots (1.48369689570172 \cdots)^{\frac{-5+3\sqrt{5}}{2} \cdot |f_n|} \end{aligned}$$

as  $n \rightarrow \infty$ .

#### ACKNOWLEDGMENTS

The author thanks Stephan Wagner for a suggestion in the proof of the asymptotic formula.

#### REFERENCES

- [1] Georgii M. Adel'son-Vel'skii and Evgenii M. Landis. An algorithm for the organization of information. *Doklady Akademii Nauk SSSR*, 146(2):263–266, 1962.
- [2] Elizabeth S. Allman and John A. Rhodes. *Mathematical models in biology: an introduction*. Cambridge University Press, Cambridge, 2004.
- [3] François Bergeron, Gilbert Labelle, and Pierre Leroux. *Combinatorial species and tree-like structures*. Cambridge University Press, Cambridge, 1998.
- [4] Éva Czabarka, Audace A.V. Dossou-Olory, László A. Székely and Stephan Wagner. Inducibility of  $d$ -ary trees. Preprint, <https://arxiv.org/abs/1802.03817>, 2018.
- [5] Audace A. V. Dossou-Olory and Stephan Wagner. Inducibility of topological trees. Accepted for publication in *Quaestiones Mathematicae*, 2018. <https://www.tandfonline.com/doi/abs/10.2989/16073606.2018.1497725>
- [6] Audace A. V. Dossou-Olory and Stephan Wagner. On the number of leaf-induced subtrees of a topological tree. To be submitted, 2018.
- [7] Ralph P. Grimaldi. Properties of Fibonacci trees. *Congressus Numerantium*, 84:21–32, 1991.
- [8] Yasuichi Horibe. An entropy view of Fibonacci trees. *The Fibonacci Quarterly*, 20(2):168–178, 1982.
- [9] Yasuichi Horibe. Notes on Fibonacci trees and their optimality. *The Fibonacci Quarterly*, 21(2):118–128, 1983.
- [10] Donald E. Knuth. *The art of computer programming (Volume 3): sorting and searching (second edition)*. Addison-Wesley Series in Computer Science and Information Processing, 791 pp., 1998.
- [11] Charles Semple and Mike Steel. *Phylogenetics*. Oxford University Press, Oxford, 2003.
- [12] Neil J. A. Sloane. The On-Line Encyclopedia of Integer Sequences. Published electronically at <https://oeis.org>, 2018.
- [13] Peter S. Stevens. *Patterns in nature*. Little, Brown and Company, United States of America, 240 pp., 1974.
- [14] Stephan Wagner. The Fibonacci number of Fibonacci trees and a related family of polynomial recurrence systems. *The Fibonacci Quarterly*, 45(3):247–253, 2007.

AUDACE A. V. DOSSOU-OLORY, DEPARTMENT OF MATHEMATICAL SCIENCES, STELLENBOSCH UNIVERSITY, PRIVATE BAG X1, MATIELAND 7602, SOUTH AFRICA

*E-mail address:* [audace@aims.ac.za](mailto:audace@aims.ac.za)