

On enumerating factorizations in reflection groups

Theo Douvropoulos*

Abstract

We describe an approach, via Malle’s permutation Ψ on the set of irreducible characters $\text{Irr}(W)$, that gives a uniform derivation of the Chapuy-Stump formula for the enumeration of reflection factorizations of the Coxeter element. It also recovers its weighted generalization by delMas, Reiner, and Hameister, and further produces structural results for factorization formulas of arbitrary regular elements.

1 Introduction

A famous theorem of Cayley states that there are n^{n-2} vertex-labeled trees on n vertices. The same number,¹ as Hurwitz knew [Hur91] already by the end of the 19th century, enumerates the set of shortest length factorizations $t_1 \cdots t_{n-1} = (12 \cdots n) \in S_n$ of the long cycle into transpositions t_i . A natural generalization of this problem, that Hurwitz himself had later considered [Hur01], is to enumerate such factorizations of arbitrary length.

It took almost a hundred years for the community to return to this question, but by the end of the 80’s Jackson [Jac88, Corol. 4.2] had computed an explicit answer. If $\text{FAC}_{S_n}(t)$ denotes the exponential generating function for the number of arbitrary length factorizations of the long cycle in transpositions (see (9)), then Jackson’s result can be reinterpreted as follows:

$$\text{FAC}_{S_n}(t) = \frac{e^{t\binom{n}{2}}}{n!} (1 - e^{-tn})^{n-1}. \quad (1)$$

As it often happens with some of the most fascinating properties of the symmetric group, the previous statements are special cases of more general theorems that hold for all reflection groups W . A natural analog of the long cycle is the Coxeter element $c \in W$, while transpositions are replaced by reflections. Then, if W is of rank n , \mathcal{R} denotes its set of reflections, and h is the order of c , Bessis [Bes15, Prop. 7.6] proved the following enumeration:

$$\#\{(t_1, \dots, t_n) \in \mathcal{R}^n \mid t_1 \cdots t_n = c\} = \frac{h^n n!}{|W|}. \quad (2)$$

The W -analog of Jackson’s formula (1) regarding arbitrary length factorizations was discovered (and proved) by Chapuy and Stump [CS14] soon after. If $\text{FAC}_W(t)$ denotes the corresponding exponential generating function, they showed that

$$\text{FAC}_W(t) = \frac{e^{t|\mathcal{R}|}}{|W|} (1 - e^{-th})^n. \quad (3)$$

*This work was supported by the European Research Council, grant ERC-2016- STG 716083 “CombiTop”.

¹The two objects are naturally related via a satisfying overcounting argument due to Dénes [D59].

The reduced case (2), which can easily be derived by calculating the leading term of $\text{FAC}_W(t)$, has a long history and appears in connection to many a mathematical endeavour. It originated in singularity theory [Loo74, Conj. (3.5); Del], in combinatorics it appeared as the number of maximal chains in the noncrossing lattice $NC(W)$ [Cha04, Prop. 9], and more importantly it was essential in Bessis’ proof of the $K(\pi, 1)$ -conjecture [Bes15] (see [Dou17, § 1] for a detailed presentation).

A uniform argument

Neither (3) nor (2) are well understood. Although the statements are uniform for all well-generated groups, the proofs of Bessis and Chapuy-Stump have relied on the Shephard-Todd classification (a common misfortune for theorems regarding reflection groups). As it happens, the main goal of this paper is to provide a case-free explanation for these formulas.

The standard approach towards results like (1) and (3) is via the Frobenius lemma (Thm. 3.1), which involves summing over all irreducible characters of a group W . For that matter, one of the main obstacles to producing a conceptual proof for (3) lies in that we have no nice, uniform construction of irreducible characters for complex reflection groups. Only for Weyl groups there is Springer’s correspondence [Spr78], which is however technically difficult for computations.

In this work we also start with the Frobenius Lemma, but instead of explicitly computing the characters $\chi \in \text{Irr}(W)$, we group them together with respect to an invariant called the Coxeter number c_χ (see Defn. 3.3). Then, Malle’s cyclic action Ψ on $\text{Irr}(W)$ allows us to cancel the contribution of those χ for which c_χ is not a multiple of h . The resulting expression is very rigid (Thm. 3.6) and the mere knowledge of bounds for the c_χ allows us to complete the proof.

Ours is not the first approach towards a uniform proof of (3). In [Mic16], Michel also considers a grouping of the characters; the partition given by Lusztig’s families. This is finer (and much more technologically advanced) and although the argument gives a very satisfying connection between (1) and (3), it requires the existence of the elusive “spets” [BMM14] when W is not a Weyl group.

Moreover, our strategy applies in further generality and produces structural results for any regular element $g \in W$ (which become explicit formulas for a larger class of groups than the well-generated ones, see Corol. 3.9). In addition, a refined version in Section 5 recovers (uniformly) and extends the main result of [dHR18] on a weighted version of the Chapuy-Stump formula (3).

When W is a real reflection group, all our theorems are completely case-free. In the complex case, although our approach is indeed uniform, it relies on the BMR-freeness theorem, a property of the Hecke algebra $\mathcal{H}(W)$ that is currently proven in a case-by-case way (see §4.5 for details).

Summary

The main results of this paper (Thm. 3.6 and Thm. 5.5) are presented in Sections 3 and 5 which can be read essentially independently of the rest. They rely on a key technical lemma (Prop. 4.19) that describes how Malle’s permutation Ψ (Defn. 4.17) affects character values on regular elements. The material in Sections 2 and 4 essentially builds up to the proof of that lemma.

In particular, the two theorems are valid for all regular elements due to a characterization of the latter ones as those that have lifts in the braid group that are roots of the full twist (see Prop. 2.9). For this reason, we have reviewed in some detail in §2 the various statements about the topological definition of the braid group and its abelianization, the full twist and the lifts of regular elements.

Similarly in Section 4, building towards the technical lemma, we recall the definition of the Hecke algebras given at [BMR98], and reproduce some key character calculations from [BM97]. The reader who is comfortable with these concepts might skip the bulk of these sections, but we hope the presentation will prove sufficient for those unfamiliar with Hecke algebras, but who might want to further pursue their combinatorial consequences.

2 Complex reflection groups and regular elements

Given a complex vector space $V \cong \mathbb{C}^n$, we call a *finite* subgroup $W \leq \mathrm{GL}(V)$ a *complex reflection group* if it is generated by unitary reflections. These are \mathbb{C} -linear maps t whose fixed spaces $V^t := \ker(t - \mathrm{id})$ are hyperplanes (i.e. $\mathrm{codim}(V^t) = 1$). We further say that W is *irreducible* if it has no stable linear subspaces apart from V and $\{0\}$. Shephard and Todd [ST54] classified irreducible complex reflection groups into an infinite 3-parameter family $G(r, p, n)$ and 34 exceptional cases indexed G_4 to G_{37} . The reader may consult the classical references [Kan01; Bro10; LT09] for the material in this section.

We denote by \mathcal{R} the set of reflections of W and we write \mathcal{A} for the associated arrangement of fixed hyperplanes. For such a hyperplane H , let W_H be its pointwise stabilizer. It consists of the identity and the reflections that fix H . Furthermore, because unitary reflections are semisimple, W_H is cyclic.

Now, if $e_H := |W_H|$ is the size of this cyclic group and t_H is one of its generators, the set of reflections \mathcal{R} can be partitioned as:

$$\mathcal{R} = \bigcup_{H \in \mathcal{A}} \{t_H, \dots, t_H^{e_H-1}\}. \quad (4)$$

The reflection group W acts on \mathcal{A} determining orbits of hyperplanes which we will denote by $\mathcal{C} \in \mathcal{A}/W$. The size $\omega_{\mathcal{C}}$ of an orbit \mathcal{C} is given by $\omega_{\mathcal{C}} := [W : N_W(H)]$ (for any $H \in \mathcal{C}$). All elements $H \in \mathcal{C}$ have conjugate stabilizers W_H and we write $e_{\mathcal{C}}$ for their common order.

With this notation, the cardinalities of the set of reflections \mathcal{R} and of the set of reflecting hyperplanes \mathcal{A} are given by

$$|\mathcal{R}| = \sum_{\mathcal{C} \in \mathcal{A}/W} \omega_{\mathcal{C}}(e_{\mathcal{C}} - 1) \quad \text{and} \quad |\mathcal{A}| = \sum_{\mathcal{C} \in \mathcal{A}/W} \omega_{\mathcal{C}}.$$

Notice that if some $e_{\mathcal{C}} \neq 2$, then $|\mathcal{R}|$ and $|\mathcal{A}|$ are not equal.

2.1 Braid groups and braid reflections

We say that a vector $v \in V$ is *regular* if it is not contained in any reflection hyperplane and we write $V^{\mathrm{reg}} := V \setminus \mathcal{A}$ for the set of regular vectors. We define the *pure braid group* $P(W) := \pi_1(V^{\mathrm{reg}})$ to be the fundamental group of the regular space V^{reg} . It is a theorem of Steinberg that the action of W on V is free precisely on V^{reg} .

Steinberg's theorem implies that the restriction of the quotient map $\rho : V \rightarrow V/W$ on V^{reg} is a Galois covering. We define the *braid group* $B(W) := \pi_1(V^{\mathrm{reg}}/W)$ to be the fundamental group of the base of this covering and use the following short exact sequence [Hat02, Prop. 1.40] to obtain a surjection $\pi : B(W) \rightarrow W$:

$$\begin{array}{ccccccc} 1 & \rightarrow & \pi_1(V^{\mathrm{reg}}) & \xrightarrow{\rho_*} & \pi_1(V^{\mathrm{reg}}/W) & \xrightarrow{\pi} & W \rightarrow 1. \\ & & \Downarrow & & \Downarrow & & \\ & & P(W) & & B(W) & & \end{array} \quad (5)$$

Given a choice of a basepoint $x_0 \in V^{\mathrm{reg}}$, a loop $\mathbf{b} \in B(W)$ lifts to a path that connects x_0 to $\mathbf{b}_*(x_0)$ (we call this the *Galois action* of \mathbf{b}). Then, we define $w := \pi(\mathbf{b})$ to be the *unique* element $w \in W$ such that $w \cdot x_0 = \mathbf{b}_*(x_0)$. The significance of (5) lies in that it gives a topological interpretation of W as the group of deck transformations of the covering map $\rho : V^{\mathrm{reg}} \rightarrow V^{\mathrm{reg}}/W$.

A reflection group W acts on the polynomial algebra $\mathbb{C}[V] := \text{Sym}(V^*)$ of the space V by precomposition (i.e. $w * f(\mathbf{v}) := f(w^{-1} \cdot \mathbf{v})$). The Shephard-Todd-Chevalley theorem [ST54; Che55] states then that the algebra of invariant polynomials $\mathbb{C}[V]^W := \{f \in \mathbb{C}[V] : w * f = f \ \forall w \in W\}$ is itself a polynomial algebra. We choose homogeneous generators for it, which we denote by f_i and order them by increasing degree $\deg(f_i) =: d_i$. The numbers d_i are independent of the choice of the f_i 's and are called the *fundamental degrees* of W .

In this setting, we can further understand the quotient morphism $\rho : V \rightarrow V/W$ by studying its algebro-geometric structure. In particular (and this holds for any finite subgroup of $\text{GL}(V)$) the map ρ is a finite morphism and the quotient V/W can be realized as the affine variety $\text{Spec}(\mathbb{C}[V]^W)$ [see Eis95, Exer. 13.2-4 and Sec. 1.7]. The Shephard-Todd-Chevalley theorem states then that for reflection groups W , the quotient V/W is itself an affine space, so that we may write:

$$\mathbb{C}^n \cong V \ni \mathbf{x} := (x_1, \dots, x_n) \xrightarrow{\rho} \mathbf{f}(\mathbf{x}) := (f_1(\mathbf{x}), \dots, f_n(\mathbf{x})) \in W \backslash V \cong \mathbb{C}^n \quad (6)$$

Now the hyperplane arrangement \mathcal{A} (which is the zero set of a collection of linear forms) is an affine variety, stable under the action of W . Another consequence of the above is then that its image $\mathcal{H} := \rho(\mathcal{A}) \subset V/W$ is itself a variety; we call it the *discriminant hypersurface* of W . The braid group becomes thus the fundamental group of a hypersurface complement $B(W) = \pi_1(V/W - \mathcal{H})$.

Such groups have a special set of generators called *generators of the monodromy* [BMR98, Appendix 1]. These are loops that descend from the basepoint following a path γ , approach a smooth point of an irreducible component of the hypersurface and make a counterclockwise² loop around it, and finally return following the same path γ backwards.

In our case, the irreducible components of \mathcal{H} are the images $\rho(\mathcal{C})$ of the hyperplane orbits $\mathcal{C} \in \mathcal{A}/W$ (again a consequence of the discussion before (6)). We will therefore denote the generators of the monodromy for $B(W)$ by $\mathbf{s}_{\mathcal{C},\gamma}$. They map (via (5)) to a subset of reflections $s_H \in W$ which have determinant $\zeta_{e_{\mathcal{C}}} := \exp(2\pi i/e_{\mathcal{C}})$ and are called *distinguished reflections*. In fact, for this reason, we follow the terminology suggested by Broué, Malle, Rouquier, Michel, and Bessis, (see for instance [Bes15, Defn. 1.6]):

Definition 2.1. The generators of the monodromy of $B(W)$ are called *braid reflections*.

The powers $\mathbf{s}_{\mathcal{C},\gamma}^{e_{\mathcal{C}}}$ are generators of the monodromy for the pure braid group:

Proposition 2.2. [BMR98, Prop. 2.18] *After a choice of basepoint $v \in V^{\text{reg}}$, we can lift the $\mathbf{s}_{\mathcal{C},\gamma}$ to paths in $P(W)$. Then the pure braid group $P(W)$ is generated by $\langle \mathbf{s}_{\mathcal{C},\gamma}^{e_{\mathcal{C}}} \rangle$ (for all \mathcal{C}, γ) and we have*

$$W \cong B(W) / \langle \mathbf{s}_{\mathcal{C},\gamma}^{e_{\mathcal{C}}} \rangle,$$

where the isomorphism is the same as the one induced by the choice of v via (5).

2.2 The full twist and the abelianization of $B(W)$

Broué-Malle-Rouquier considered [BMR98, Notation 2.3] a particular element of the pure braid group $P(W)$; it is fundamental in what follows and for the results in Sections 3 and 5. For an arbitrary regular vector $v \in V^{\text{reg}}$, we define $\pi_v \in \pi_1(V^{\text{reg}}, v)$ as the loop given by:

$$[0, 1] \ni t \rightarrow e^{2\pi i t} \cdot v. \quad (7)$$

²Near a smooth point, an irreducible codimension 1 divisor in \mathbb{C}^n looks like a line in \mathbb{R}^3 ; there is a well-defined way to go around it.

If $\gamma \subset V^{\text{reg}}$ is any path between points $v, v' \in V^{\text{reg}}$, then the cylinder (or torus if γ is a loop) $S^1 \cdot \gamma$ lies completely inside V^{reg} . This is because V^{reg} , the complement of a central hyperplane arrangement, is stable under multiplication by $\mathbb{C}^\times \supset S^1$. It is immediate from this that:

Lemma 2.3. [BMR98, Lemma 2.5]

For v, v' , and γ as above, the loops $\gamma^{-1} \cdot \pi_{v'} \cdot \gamma$ and π_v in $P(W, v)$ are homotopic.

This in particular implies that π_v is always central in $P(W, v)$. Furthermore, if v and v' have the same image in V^{reg}/W , and since ρ is quasihomogeneous (6), the loops $\rho_*(\pi_v)$ and $\rho_*(\pi_{v'})$ are identical. Now, this along with the previous lemma immediately gives:

Corollary 2.4. [BMR98, from Lemma 2.22: (2)]

For any regular vector $v \in V^{\text{reg}}$, the element $\rho_*(\pi_v) \in B(W, \rho(v))$ is central.

For any two basepoints v and v' of V^{reg} and a path γ between them, there are canonical isomorphisms between the fundamental groups $P(W, v)$ and $P(W, v')$, and between $B(W, \rho(v))$ and $B(W, \rho(v'))$. Since π_v and $\rho_*(\pi_v)$ are central, their images will also be central and moreover independent of the path γ (in fact, the previous lemma shows that they will be homotopic to $\pi_{v'}$ and $\rho_*(\pi_{v'})$ respectively). We therefore drop the basepoint from the notation, and for convenience we use the same symbol for the image in $B(W)$ as well:

Definition 2.5. [Bes15, Defn. 6.12]

We call this element π defined in (7) the *full twist*. It is central in $B(W)$ and lies in $P(W)$.

Broué-Malle-Rouquier also consider [BMR98, Defn. 2.15] length functions $l_{\mathcal{C}} : B(W) \rightarrow \mathbb{Z}$, given as periods of the differential forms $d \text{Log}(\delta_{\mathcal{C}})$ associated to discriminant polynomials $\delta_{\mathcal{C}}$ that cut out the strata \mathcal{C} of \mathcal{H} [BMR98, Defn. 2.15]. For a loop $\mathbf{g} \in B(W)$, they essentially record how many radians any of its lifts $\mathbf{g}' \in P(W)$ wraps around each hyperplane in the orbit $\mathcal{C} \in \mathcal{A}/W$, and weigh the result by $e_{\mathcal{C}}$ (see [ibid, Thm. 2.17: Remark]). In particular, they satisfy [ibid, Prop. 2.16]

$$l_{\mathcal{C}}(\mathbf{s}_{\mathcal{C}', \gamma}) = \delta_{\mathcal{C}, \mathcal{C}'},$$

which, since the $\mathbf{s}_{\mathcal{C}, \gamma}$ generate $B(W)$ (see discussion before Prop. 2.2), implies that in fact these length functions completely determine the abelianization B^{ab} of $B(W)$:

Theorem 2.6. [BMR98, Thm. 2.17:(2)]

If $\mathbf{s}_{\mathcal{C}}^{\text{ab}}$ denotes the image of any $\mathbf{s}_{\mathcal{C}, \gamma}$ in the abelianization B^{ab} , then

$$B^{\text{ab}} = \prod_{\mathcal{C} \in \mathcal{A}/W} \langle \mathbf{s}_{\mathcal{C}}^{\text{ab}} \rangle,$$

where each $\langle \mathbf{s}_{\mathcal{C}}^{\text{ab}} \rangle$ is infinite cyclic. Moreover, for an element $\mathbf{g} \in B(W)$, we have

$$\mathbf{g}^{\text{ab}} = \prod_{\mathcal{C} \in \mathcal{A}/W} (\mathbf{s}_{\mathcal{C}}^{\text{ab}})^{l_{\mathcal{C}}(\mathbf{g})}.$$

By definition the full twist π rotates once around each of the $\omega_{\mathcal{C}}$ -many hyperplanes in any orbit \mathcal{C} :

Corollary 2.7. [BMR98, Cor. 2.26 and Lemma 2.22:(2)]

Let π^{ab} be the image in B^{ab} of the full twist π . Then we have

$$\pi^{\text{ab}} = \prod_{\mathcal{C} \in \mathcal{A}/W} (\mathbf{s}_{\mathcal{C}}^{\text{ab}})^{e_{\mathcal{C}} \cdot \omega_{\mathcal{C}}}.$$

2.3 Regular elements and roots of the full twist

Although our initial purpose for this project was to give a uniform proof of the Chapuy-Stump formula (3) which regards Coxeter elements, it soon became clear that the techniques developed (see Lemma 3.5) apply to the larger class of Springer-regular elements. The crucial property these elements share is that they lift to roots of (powers of) the full twist π (Defn. 2.5).

Definition 2.8. [Spr74] Recall the space V^{reg} of *regular* vectors; namely those that do not lie in any hyperplane $H \in \mathcal{A}$. We say that an element $g \in W$ is ζ -regular if it has a regular ζ -eigenvector; all ζ -regular elements are conjugate [LT09, Corol. 11.25]. The order d of a ζ -regular element g is equal to the order of ζ [ibid] and is called a *regular number*.

For real reflection groups W , the product c of the simple generators (in any order) is called a Coxeter element, after Coxeter who first computed its order h and eigenvalues [Cox51]. At the same paper, Coxeter observed (and Steinberg later [Ste59] gave a uniform proof of the fact) that h determines the number of hyperplanes N via the equation $nh = 2N$, where n is the dimension of the ambient space V . Steinberg's work easily implies also that c is an $e^{2\pi i/h}$ -regular element.

Building on that, Gordon and Griffeth (but see also the beginning of §4.4) define a Coxeter number³ for all complex reflection groups as $h = (|\mathcal{R}| + |\mathcal{A}|)/n$. Then, for an arbitrary W , a *Coxeter element* is defined as a $e^{2\pi i/h}$ -regular element. It turns out that Coxeter elements exist precisely when W is *well-generated*; namely when it is generated by n reflections.

It is easy to produce lifts $\mathbf{g} \in B(W)$ of regular elements $g \in W$. Indeed, let g be a ζ -regular element, with $\zeta = \exp(2\pi im/d)$, $(m, d) = 1$, and let x_0 be one of its ζ -eigenvectors. Consider now the path $\pi_{x_0, \zeta}$ in V^{reg} that connects x_0 and ζx_0 and is defined by

$$[0, 1] \ni t \rightarrow e^{2\pi itm/d} x_0. \quad (8)$$

Since $\zeta x_0 = g \cdot x_0$, this determines a loop in V^{reg}/W that would lift the element $g \in W$, if x_0 was the basepoint for $P(W)$. We can easily adjust the construction to deal with a basepoint that is not an eigenvector, and comparing (7) and (8) gives:

Proposition 2.9. [Bro10, Prop. 5.24] *Let $\zeta = \exp(2\pi im/d)$ be a primitive d^{th} root of unity, and let g be a ζ -regular element of W . Then, g has a lift $\mathbf{g} \in B(W)$ such that $\mathbf{g}^d = \pi^m$.*

Proof. Let $v \in V^{\text{reg}}$ be the basepoint of $P(W)$ and γ an arbitrary path in V^{reg} that connects v with a ζ -eigenvector x_0 of g . We view g as a deck transformation of the covering $\rho : V^{\text{reg}} \rightarrow V^{\text{reg}}/W$ and consider the path $(g \cdot \gamma^{-1}) \cdot \pi_{x_0, \zeta} \cdot \gamma$. It connects the points v and $g \cdot v$ and hence determines the following element of the braid group $B(W)$:

$$\mathbf{g} := \rho(\gamma)^{-1} \cdot \rho(\pi_{x_0, \zeta}) \cdot \rho(\gamma).$$

Because g acts on the line $\mathbb{C} \cdot x_0$ as multiplication by ζ , we can see that the loop $\rho(\pi_{x_0, \zeta})^d$ lifts to the element $\pi_{x_0}^m = \pi_{\zeta^{d-1} \cdot x_0, \zeta} \cdots \pi_{x_0, \zeta}$ (recall the definition of π_{x_0} in (7)). This immediately gives

$$\mathbf{g}^d = \rho(\gamma)^{-1} \cdot \rho(\pi_{x_0})^m \cdot \rho(\gamma),$$

which after the discussion before Defn. 2.5 completes the proof. □

³It is not a priori clear that h is an integer; see Corol. 4.16.

Remark 2.10. The converse of the previous theorem is still true; that is, d^{th} roots of the full twist exist precisely when d is a regular number [Bes15, Thm. 12.4]. Moreover, as with Springer-regular elements, Bessis has shown [ibid] that all d^{th} roots of π^m are conjugate. Such results essentially “lift” Springer theory to braid groups; they rely on garside-like structures in [Bes06].

However, we should warn the reader that this does not imply the existence of nice sections from W to $B(W)$. Moreover, even for Coxeter groups, where the existence of simple systems allows us to lift W in $B^+(W)$, these lifts do not satisfy the previous properties. That is, conjugate regular elements (in particular, Coxeter elements) lift to not necessarily conjugate elements in $B(W)$.

If g is a d^{th} root of the full twist, Thm. 2.6 and Corol. 2.7 imply that $l_C(g) \cdot d = e_C \omega_C$. This proves the following as in [dHR18, Thm. 3.2] (but see also [Bro10, Prop. 5.17:(2)]):

Corollary 2.11. *For any orbit $C \in \mathcal{A}/W$, a regular number d always divides the quantity $e_C \cdot \omega_C$.*

In Sections 3 and 5 we prove some structural results for factorization enumeration formulas for arbitrary regular elements. When the order of these elements equals the highest fundamental degree d_n , this structural information is in fact sufficient to determine explicit formulas. We list here the corresponding types:

Proposition 2.12. [Bes01, Prop. 4.1] *Let W be an irreducible complex reflection group and let d_n be its largest degree. Then, d_n is a regular number precisely when W is a Coxeter group, or $G(r, 1, n)$, $G(r, r, n)$ and $G(2r, 2, 2)$, or any exceptional group other than G_{15} .*

Remark 2.13. We have tried to carefully show in this section that the choice of the basepoint $v \in V^{\text{reg}}$ does not affect the theorems regarding the full twist, the abelianization, and the regular elements. At this point we choose a basepoint v , once and for all, and in what follows we consider the surjection $B(W) \rightarrow W$ in (5) fixed.

3 Frobenius lemma via Coxeter numbers

The lemma of Frobenius, which does in fact go back to Frobenius and 1896 [Fro68], gives a representation theoretic formula for enumerating factorizations of group elements, when the factors belong to given (unions of) conjugacy classes:

Theorem 3.1. [LZ04, App. A.1.3] *Let G be a finite group and $A_i \subset G$, $i = 1 \dots l$, subsets that are closed under conjugation. Then the number of factorizations $t_1 \cdots t_l = g$ of an element $g \in W$, where each factor t_i belongs to A_i , is given by*

$$\frac{1}{|G|} \sum_{\chi \in \widehat{G}} \chi(1) \cdot \chi(g^{-1}) \cdot \frac{\chi(A_1)}{\chi(1)} \cdots \frac{\chi(A_l)}{\chi(1)},$$

where \widehat{G} denotes the (complete) set of irreducible characters of G and $\chi(A) := \sum_{g \in A} \chi(g)$.

For a reflection group W , the set of reflections \mathcal{R} is indeed closed under conjugation. This lemma of Frobenius implies then a simple finite-sum form for the exponential generating function of reflection factorizations of elements of W . If we write $\text{Fact}_{W,g}(l)$ for the number of such factorizations of length l , i.e.:

$$\text{Fact}_{W,g}(l) := \#\{(t_1, \dots, t_l) \in \mathcal{R}^l \mid t_1 \cdots t_l = g\},$$

then the lemma of Frobenius implies that

$$\text{Fact}_{W,g}(l) = \frac{1}{|W|} \sum_{\chi \in \widehat{W}} \chi(1) \cdot \chi(g^{-1}) \cdot \left[\frac{\chi(\mathcal{R})}{\chi(1)} \right]^l.$$

After this, the exponential generating function for reflection factorizations of g is given by:

$$\text{FAC}_{W,g}(t) := \sum_{l \geq 0} \text{Fact}_{W,g}(l) \cdot \frac{t^l}{l!} = \frac{1}{|W|} \sum_{\chi \in \widehat{W}} \chi(1) \cdot \chi(g^{-1}) \cdot \exp \left[t \cdot \frac{\chi(\mathcal{R})}{\chi(1)} \right]. \quad (9)$$

Notice that, remarkably, this observation that such generating functions will be expressible as finite sums of exponentials appears already in Hurwitz's paper [Hur01, §3:(15)].

Now, a priori the evaluations $\chi(\mathcal{R})$ are complex numbers, but the special structure of the set of reflections \mathcal{R} forces them to in fact be integers (recall that \mathcal{A} denotes the set of fixed hyperplanes):

Proposition 3.2. *The numbers $\chi(\mathcal{R})$ are integers, and they further satisfy:*

$$-|\mathcal{A}| \cdot \chi(1) \leq \chi(\mathcal{R}) \leq |\mathcal{R}| \cdot \chi(1).$$

Both bounds are achieved; the higher only for the trivial representation, and the lower at least for the det representation.

Proof. Recall the decomposition of the set of reflections with respect to their fixed hyperplanes $H \in \mathcal{A}$ as described in (4). Keeping that notation, we choose a generator t_H for each of the cyclic groups W_H and write $e_H := |W_H|$ for its order.

For each eigenvalue λ of t_H in the representation U_χ associated to χ , the contribution of the set of reflections $\{t_H, \dots, t_H^{e_H-1}\}$ in the evaluation of $\chi(\mathcal{R})$ equals $\sum_{k=1}^{e_H-1} \lambda^k$. Since $\lambda^{e_H} = 1$, this quantity is either $e_H - 1$ or -1 depending on whether λ itself is 1 or not.

This implies the first two statements of the proposition, after noticing that the multiset of eigenvalues of t_H acting on U_χ has $\chi(1)$ -many elements. In particular, in order to recover the second inequality we use that $\sum_{H \in \mathcal{A}} (e_H - 1) = |\mathcal{R}|$ which is immediate after the partitioning (4).

For the last statement, the higher bound is achieved when each eigenvalue of each t_H equals 1; of course this happens only in the trivial representation. For the lower bound, we need all $\lambda \neq 1$, which happens for instance in the (1-dimensional) det representation. \square

The character values $\chi(\mathcal{R})$ on the sum of reflections are related to an statistic of the associated representation called the *Coxeter number* and denoted by c_χ . We postpone to §4.4 the discussion about its origin and for now we only give the definition:

Definition 3.3. [GG12, §1.3] We define the Coxeter number c_χ associated to the character χ , as the normalized trace of the central element $\sum_{t \in \mathcal{R}} (\mathbf{1} - t)$. That is,

$$c_\chi := \frac{1}{\chi(1)} \cdot (|\mathcal{R}| \chi(1) - \chi(\mathcal{R})) = |\mathcal{R}| - \frac{\chi(\mathcal{R})}{\chi(1)}.$$

After Prop. 3.2 the numbers c_χ are rational, but as we will see in Corol. 4.16 they are in fact integers.

It is easy now to reinterpret formula (9) in terms of the Coxeter numbers c_χ . We record the following as a corollary of Thm. 3.1:

Corollary 3.4. *The exponential generating function $\text{FAC}_{W,g}(t)$ for arbitrary length reflection factorizations of an element $g \in W$ is given by:*

$$\text{FAC}_{W,g}(t) = \frac{e^{t|\mathcal{R}|}}{|W|} \sum_{\chi \in \widehat{W}} \chi(1) \cdot \chi(g^{-1}) \cdot e^{-t \cdot c_\chi}. \quad (10)$$

The following lemma is the main technical ingredient for the proof of Thm. 3.6. Its derivation, which we postpone until Section 4 (see after Prop. 4.19), relies on a cyclic action on the set $\text{Irr}(W)$ of irreducible representations of W which is induced by a Galois action (see Defn. 4.17) on the modules of the Hecke algebra. Recall Defn. 2.8 for the concept of a regular element.

Lemma 3.5. *For a complex reflection group W , and a **regular** element $g \in W$, the total contribution in (10) of those characters $\chi \in \widehat{W}$ for which c_χ is not a multiple of $|g|$ is 0.*

The following is an essentially immediate application of Lemma 3.5. We state it as a theorem as all explicit formulas that come after (3.8-3.12) are derived as its corollaries:

Theorem 3.6. *For a complex reflection group W , and a regular element $g \in W$, the exponential generating function $\text{FAC}_{W,g}(t)$ of reflection factorizations of g takes the following form:*

$$\text{FAC}_{W,g}(t) = \frac{e^{t|\mathcal{R}|}}{|W|} \cdot \left[(1 - X)^{l_R(g)} \cdot \Phi(X) \right] \Big|_{X=e^{-t|g|}}.$$

Here $l_R(g)$ is the reflection length of g and $\Phi(X)$ is a polynomial in X that has degree $\frac{|\mathcal{R}|+|\mathcal{A}|}{|g|} - l_R(g)$, is not further divisible by $(1 - X)$, and has constant term equal to 1.

Proof. After Lemma 3.5 we only need to consider terms of the form $\chi(1) \cdot \chi(g^{-1}) \cdot e^{-t \cdot k|g|}$, $k \in \mathbb{Z}$ in the evaluation of (10). Furthermore, rephrasing Prop. 3.2 in terms of the Coxeter numbers (via Defn. 3.3) forces $k \in \{0, \dots, \frac{|\mathcal{R}|+|\mathcal{A}|}{|g|}\}$. This means that if we set $X = e^{-t|g|}$, we can rewrite (10) as

$$\text{FAC}_{W,g}(t) = \frac{e^{t|\mathcal{R}|}}{|W|} \cdot \tilde{\Phi}(X),$$

where $\tilde{\Phi}(X)$ is a priori a polynomial in $\mathbb{C}[X]$ of degree $(|\mathcal{R}| + |\mathcal{A}|)/|g|$. The last statement of Lemma 3.5 implies also that the constant term of $\tilde{\Phi}(X)$ is equal to $\chi_{\text{triv}}(1) \cdot \chi_{\text{triv}}(g^{-1}) = 1$.

Now, since $\tilde{\Phi}(X)$ essentially encodes the generating function $\text{FAC}_{W,g}(t)$, the combinatorial properties of the latter impose restrictions on its structure. In particular, consider the root factorization of the polynomial:

$$\tilde{\Phi}(X) = a(\alpha_1 - X)(\alpha_2 - X) \cdots (\alpha_r - X).$$

If we revert to $X = e^{-t|g|}$, each of the linear terms above has a Taylor expansion that starts with $(\alpha_i - 1) + t|g| + \dots$. This means that it contributes to the leading term of $\text{FAC}_{W,g}(t)$ either by a factor of $(\alpha_i - 1)$ or by a factor of $t|g|$, depending on whether α_i equals 1 or not.

On the other hand, the combinatorial definition of $\text{FAC}_{W,g}(t)$ in (9) implies that its leading term is a multiple of $t^{l_R(g)}$. Therefore, exactly $l_R(g)$ -many of the roots of $\tilde{\Phi}$ must be equal to 1 and this completes the proof. The statements about the degree and the constant term follow from the analogous results for $\tilde{\Phi}$ described previously. \square

Remark 3.7. In the previous argument, the existence of a reflection length and therefore the knowledge that the first few terms of the generating function $\text{FAC}_{W,g}(t)$ are zero, came for free but

was very useful nonetheless. This sort of reasoning has appeared already in [MN10, end of proof of Thm. 2]. It is hoped that similar ideas might apply to other groups with natural length functions, such as $\mathrm{GL}_n(\mathbb{F}_q)$ (see [LRS14; LM16]). Moreover, one might construct special length functions to support different enumerative questions (as we pursue in Prop. 3.11 and in Defn. 5.3).

Corollary 3.8. *For a complex reflection group W , and a regular element $g \in W$, the number of reduced reflection factorizations of g is an integer multiple of the quantity $\frac{|g|^{l_R(g)}(l_R(g))!}{|W|}$.*

Proof. The leading coefficient of $\mathrm{FAC}_{W,g}(t)$ is given, after Thm. 3.6, by

$$\Phi(1) \cdot \frac{|g|^{l_R(g)}(l_R(g))!}{|W|}.$$

It suffices then, to show that $\Phi(1)$ is an integer. By definition, the coefficients of the polynomial $\tilde{\Phi}(X)$ are algebraic integers and so the same is true for $\Phi(X)$. The quantity $\Phi(1)$ is thus an algebraic integer, and since it also has to be a rational number (because an integer multiple of it enumerates factorizations), it must be an integer. \square

Corollary 3.9. *For a complex reflection group W and a regular element $g \in W$ of order $|g| = d_n$, the exponential generating function for reflection factorizations of g is given by:*

$$\mathrm{FAC}_{W,g}(t) = \frac{e^{t|\mathcal{R}|}}{|W|} \cdot (1 - e^{-t|g|})^{l_R(g)}.$$

Proof. After Thm. 3.6 it is sufficient to show that for such an element g , the polynomial $\Phi(X)$ is equal to the scalar 1, or equivalently that its degree is 0 (notice that then, $\Phi(X)$ cannot be any other scalar since, again by Thm. 3.6, its constant term is always 1).

The degree of $\Phi(X)$ is also given in the theorem; it equals $\frac{|\mathcal{R}|+|\mathcal{A}|}{|g|} - l_R(g)$. Now, Bessis has shown [Bes01, Prop. 4.2] that when d_n is a regular number, the quantity $(|\mathcal{R}| + |\mathcal{A}|)/d_n$ is equal to the minimum number of reflections needed to generate W (either n or $n + 1$). Therefore, if the degree of $\Phi(X)$ is not 0, the d_n -regular element g must live in a reflection subgroup W' of W .

If this were indeed the case, g would still be regular in W' and Springer's theorem [Kan01, §32-2] would allow us to list its eigenvalues in two ways:

$$\{\zeta^{1-d_1}, \dots, \zeta^{1-d_n}\} = \{\zeta^{1-d'_1}, \dots, \zeta^{1-d'_n}\},$$

where the d'_i are the invariant degrees of W' and ζ is a primitive d_n -th root of unity. This would force the two (multi-)sets of residues $\{d_i \bmod(d_n)\}$ and $\{d'_i \bmod(d_n)\}$ to be equal, but since $0 \leq d_i \leq d_n$ and $\prod_{i=1}^n d_i = |W| > |W'| = \prod_{i=1}^n d'_i$, this is impossible. \square

Remark 3.10. When W is a well-generated group and c a Coxeter element of W , we always have $|c| = d_n$. The previous corollary therefore completes a proof of the Chapuy-Stump formula (3) and extends it to the groups listed in Prop. 2.12.

In Thm. 3.6 the knowledge of the reflection length of an element provides structural information for a factorization enumeration formula. Here, we show an example where we can push this slightly further by considering a different length function, namely the transitive factorization length:

Proposition 3.11. *The exponential generating function for transitive reflection factorizations of the regular element $g = (12 \cdots n - 1)(n) \in S_n$ is given by*

$$\mathrm{TR}\text{-FAC}_{S_n,g}(t) = \frac{e^{t\binom{n}{2}}}{n!} \cdot (1 - e^{-t(n-1)})^n.$$

Proof. Since S_{n-1} is the only reflection subgroup of S_n that contains the element g , we can enumerate the transitive reflection factorizations of the latter by subtracting from all possible factorizations, those that live in S_{n-1} :

$$\text{TR-FAC}_{S_n,g}(t) = \text{FAC}_{S_n,g}(t) - \text{FAC}_{S_{n-1},g}(t).$$

If we apply Thm. 3.6 and Corol. 3.9 to the two terms above, we get, for $X = e^{-t(n-1)}$:

$$\begin{aligned} \text{FAC}_{S_n,g}(t) - \text{FAC}_{S_{n-1},g}(t) &= \frac{e^{t\binom{n}{2}}}{n!} \cdot (1 - e^{-t(n-1)})^{n-2} \cdot \Phi(X) - \frac{e^{t\binom{n-1}{2}}}{(n-1)!} \cdot (1 - e^{-t(n-1)})^{n-2} \\ &= \frac{e^{t\binom{n}{2}}}{n!} \cdot (1 - e^{-t(n-1)})^{n-2} \cdot (\Phi(X) - nX), \end{aligned}$$

where $\Phi(X)$ has degree $2 = \frac{2\binom{n}{2}}{n-1} - (n-2)$ and constant term equal to 1.

Notice now that the leading term of the generating function $\text{TR-FAC}_{S_n,g}(t)$ needs to be a multiple of t^n . Indeed, n is a lower bound for the length of transitive reflection factorizations of g , since at least $n-1$ reflections are needed to generate S_n , but since also g cannot be written as a product of $n-1$ reflections as it has parity $(-1)^{n-2}$.

Of course, $(1 - e^{-t(n-1)})^{n-2}$ contributes a factor of t^{n-2} to the leading term of the generating function, so $(\Phi(X) - nX)$ must contribute a multiple of t^2 . As in the proof of Thm. 3.6, and because $\deg(\Phi(X)) = 2$, this implies that

$$\Phi(X) - nX = (1 - X)^2,$$

which completes the argument. \square

Corollary 3.12. *For the symmetric group S_n and the regular element $g = (12 \cdots n-1)(n) \in S_n$, the polynomial $\Phi(X)$ from Thm. 3.6 is given by:*

$$\Phi(X) = 1 + (n-2)X + X^2.$$

Remark 3.13. It is not clear whether one should expect a nice formula for the polynomials $\Phi_g(X)$. They don't seem to factor in small order terms and their coefficients, although integers, are not always positive (an example being the regular class of order 3 in E_6). It might be however that a better answer exists for the infinite family $G(r, p, n)$ (or even just the symmetric group S_n), where the regular elements have simple cycle types.

Question 3.14. For Weyl groups W , one can easily see [Spr74, Prop. 4.10] that any regular element of order d divides the set of roots in orbits of size d . Perhaps this could be used in a fashion similar to the recursion in [Del; Rea08] and, possibly assuming the Lemma of Frobenius (10), give a combinatorial proof of our technical Lemma 3.5.

4 Hecke algebras and the technical lemma

Iwahori-Hecke algebras associated to Weyl groups W appear naturally as endomorphism algebras of certain induced modules in the representation theory of finite groups of Lie type. They can also be seen as deformations of the corresponding group ring $\mathbb{Z}[W]$. This second interpretation has been extended for all complex reflection groups:

Let $\mathcal{C} \in \mathcal{A}/W$ denote an orbit of hyperplanes, and $e_{\mathcal{C}}$ the common order of the pointwise stabilizers W_H (for $H \in \mathcal{C}$). Consider now a set of $\sum_{\mathcal{C} \in \mathcal{A}/W} e_{\mathcal{C}}$ many variables $\mathbf{u} := (u_{\mathcal{C},j})_{(\mathcal{C} \in \mathcal{A}/W), (0 \leq j \leq e_{\mathcal{C}}-1)}$ and write $\mathbb{Z}[\mathbf{u}, \mathbf{u}^{-1}]$ for the Laurent polynomial ring on the $u_{\mathcal{C},j}$'s.

Definition 4.1. [BMR98, Defn. 4.21] The *generic Hecke algebra* $\mathcal{H}(W)$ associated to W is the quotient of the group ring $\mathbb{Z}[\mathbf{u}, \mathbf{u}^{-1}]B(W)$ of the braid group, over the ideal generated by the elements of the form

$$(\mathbf{s} - u_{\mathcal{C},0})(\mathbf{s} - u_{\mathcal{C},1}) \cdots (\mathbf{s} - u_{\mathcal{C},e_{\mathcal{C}}-1}), \quad (11)$$

which we call *deformed order relations* (see (12)). Here \mathbf{s} runs over all possible braid reflections (see §2.1) around the stratum \mathcal{C} of \mathcal{H} . Notice that for each orbit \mathcal{C} one such relation is in fact sufficient since all corresponding elements $\mathbf{s}_{\mathcal{C},\gamma}$ are conjugate in $B(W)$.

Notation 4.2. For an element \mathbf{g} of the braid group $B(W)$, we denote the corresponding element in the Hecke algebra by $T_{\mathbf{g}}$.

Any ring map $\theta : \mathbb{Z}[\mathbf{u}, \mathbf{u}^{-1}] \rightarrow R$ defines an R -module structure on the Hecke algebra. We write $\mathcal{H}_R(W) := \mathcal{H}(W) \otimes_{\mathbb{Z}[\mathbf{u}, \mathbf{u}^{-1}]} R$ and call $\mathcal{H}_R(W)$ a *specialization* of $\mathcal{H}(W)$. The map θ induces thus a canonical map $\tilde{\theta} : \mathcal{H}(W) \rightarrow \mathcal{H}_R(W)$ via $T_{\mathbf{g}} \rightarrow T_{\mathbf{g}} \otimes 1$.

The Hecke algebra is by construction a deformation of the group algebra of W . Indeed, the specialization (recall $\zeta_n := \exp(2\pi i/n)$)

$$u_{\mathcal{C},j} \xrightarrow{\sigma} \zeta_{e_{\mathcal{C}}}^j \quad (12)$$

transforms the defining relations (11) to order relations of the form $\mathbf{s}^{e_{\mathcal{C}}} = 1$. Then, by Prop. 2.2 $\mathcal{H}(W)$ reduces to the group ring $\mathbb{Z}[(\zeta_{e_{\mathcal{C}}})]_{(\mathcal{C} \in \mathcal{A}/W)}[W]$ and the map $\tilde{\theta}$ agrees with the fixed (see Rem. 2.13) surjection $B(W) \twoheadrightarrow W$. That is, if $g \in W$ is the image of $\mathbf{g} \in B(W)$ under (5), then

$$\tilde{\theta}(T_{\mathbf{g}}) = g.$$

Definition 4.3. A specialization θ will be called *admissible* if it factors through (12); in other words if there is a map $f : R \rightarrow \mathbb{Z}[(\zeta_{e_{\mathcal{C}}})]$ such that $f \circ \theta(u_{\mathcal{C},j}) = \zeta_{e_{\mathcal{C}}}^j$.

Two particular specializations are fundamental in what follows. We first pick a set of parameters $\mathbf{x} := (x_{\mathcal{C}})_{\mathcal{C} \in \mathcal{A}/W}$ and the single parameter x and define the following ring maps:

$$\begin{aligned} \theta_{\mathbf{x}} : \mathbb{Z}[\mathbf{u}, \mathbf{u}^{-1}] &\rightarrow \mathbb{Z}[\mathbf{x}, \mathbf{x}^{-1}] & \text{and} & & \theta_x : \mathbb{Z}[\mathbf{u}, \mathbf{u}^{-1}] &\rightarrow \mathbb{Z}[x, x^{-1}] \\ \theta_{\mathbf{x}}(u_{\mathcal{C},j}) &= \begin{cases} x_{\mathcal{C}} & \text{if } j = 0 \\ \zeta_{e_{\mathcal{C}}}^j & \text{if } j \neq 0 \end{cases} & & & \theta_x(u_{\mathcal{C},j}) &= \begin{cases} x & \text{if } j = 0 \\ \zeta_{e_{\mathcal{C}}}^j & \text{if } j \neq 0 \end{cases} \end{aligned} \quad (13)$$

Both $\theta_{\mathbf{x}}$ and θ_x are admissible specializations (as seen by further sending $x_{\mathcal{C}}$ or x to 1). We write $\mathcal{H}_{\mathbf{x}}(W)$ and $\mathcal{H}_x(W)$ for the corresponding Hecke algebras, while noting that the latter is the analogue of the 1-parameter Iwahori-Hecke algebra of real reflection groups W .

Artin-like presentations and the BMR-freeness theorem

Bessis [Bes01] has shown that the braid groups $B(W)$ always have ‘‘Artin-like’’ presentations. These are presentations of the form

$$\langle \mathbf{s}_1, \dots, \mathbf{s}_n \mid p_j(\mathbf{s}_1, \dots, \mathbf{s}_n) = q_j(\mathbf{s}_1, \dots, \mathbf{s}_n) \rangle,$$

where the \mathbf{s}_i ’s are braid reflections (so they equal $\mathbf{s}_{\mathcal{C},\gamma}$ for suitable \mathcal{C} and γ) and their images $s_H \in W$ form a minimal generating set of (distinguished) reflections. Furthermore, the relations (p_j, q_j) encode positive words of equal length in the \mathbf{s}_i ’s and are such so that by adding the order relations $\mathbf{s}_i^{e_H} = 1$, one obtains a presentation of the group W .

By now, such Artin-like presentations have been found for all braid groups $B(W)$ (see [Bro10, Appendix A.2]). With access to these, one can write down explicit presentations for the Hecke algebras and with them attempt to study their various structural properties and invariants.

Example 4.4. The generic Hecke algebra of G_{26} (over the ring $\mathbb{Z}[x_0^{\pm 1}, \dots, y_2^{\pm 1}]$) is:

$$\begin{aligned} \mathcal{H}(G_{26}) = \langle \mathbf{s}, \mathbf{t}, \mathbf{u} \mid & \mathbf{stst} = \mathbf{tsts}, \mathbf{su} = \mathbf{us}, \mathbf{tut} = \mathbf{utu}, \\ & (\mathbf{s} - x_0)(\mathbf{s} - x_1) = 0 \\ & (\mathbf{t} - y_0)(\mathbf{t} - y_1)(\mathbf{t} - y_2) = 0 \\ & (\mathbf{u} - y_0)(\mathbf{u} - y_1)(\mathbf{u} - y_2) = 0 \rangle \end{aligned}$$

The braid reflections \mathbf{t} and \mathbf{u} are conjugate (although this is a bit hard to see from the given presentation of $B(G_{26})$), so we use the same set of variables for their deformed order relations. After the specializations $(x_0, x_1) = (1, -1)$, $(y_0, y_1, y_2) = (1, \zeta_3, \zeta_3^2)$, we obtain the following Coxeter-like presentation of G_{26} :

$$G_{26} = \langle s, t, u \mid stst = tsts, su = us, tut = utu, s^2 = t^3 = u^3 = 1 \rangle.$$

This definition of Hecke algebras, which recovers the usual Iwahori-Hecke algebras when W is a Coxeter group, is due to Broué, Malle, and Rouquier, and was introduced in their seminal paper [BMR98]. There, they also made various conjectures about these Hecke algebras, the most important of which was until recently known as “*The BMR freeness conjecture*”:

Theorem. [Eti17][Bou+18, after Thm. 3.5]

The algebra $\mathcal{H}(W)$ is a free $\mathbb{Z}[\mathbf{u}, \mathbf{u}^{-1}]$ -module of rank $|W|$.

4.1 Tits’ deformation theorem for admissible specializations

For this work, the first important consequence of the BMR-freeness theorem is that it determines, via Tits’ deformation theorem, a bijection between the irreducible representations of W and those of the Hecke algebra. The reader might refer to [GP00, §7] for proofs and terminology.

To apply Tits’ deformation theorem, we first have to move to split extensions of $\mathcal{H}(W)$ and of the group algebra of W . For the latter, we could simply work over $\mathbb{C}[W]$, but it takes little effort to describe its minimal splitting field. To begin with, it is easy to see [Bes97, Corol. 3.2] that the reflection representation V of W can be realized over the field K generated by the traces of the elements of W on V . It is a theorem of Benard and Bessis [Ben76; Bes97] that in fact all representations of W can be realized over K .

We henceforth call K the *field of definition* of W ; it equals \mathbb{Q} when W is a Weyl group and satisfies $K \leq \mathbb{R}$ when W is a finite Coxeter group. One might then hope that $K(\mathbf{u})$ is a splitting field for $\mathcal{H}(W)$. Although this is not the case, the answer is only slightly more complicated. Assuming the BMR-freeness conjecture, Malle proved (with further case-specific arguments, but see §4.5):

Proposition 4.5. [Mal99, Thm. 5.2] *Let K be the field of definition of W as above. Then, there exists a number N_W such that for a set of parameters $\mathbf{v} := (v_{\mathcal{C},j})_{(\mathcal{C} \in A/W), (0 \leq j \leq e_{\mathcal{C}}-1)}$ that satisfy*

$$v_{\mathcal{C},j}^{N_W} = \zeta_{e_{\mathcal{C}}}^{-j} u_{\mathcal{C},j},$$

the field $K(\mathbf{v})$ is a splitting field for $\mathcal{H}(W)$. We write $\mathcal{H}_{K(\mathbf{v})}(W) := \mathcal{H}(W) \otimes_{\mathbb{Z}[\mathbf{u}, \mathbf{u}^{-1}]} K(\mathbf{v})$.

Of course, after the BMR-freeness conjecture, $\mathcal{H}_{K(\mathbf{v})}(W)$ will also be a free $K[\mathbf{v}, \mathbf{v}^{-1}]$ -module and we may extend the specialization (12) to a map $K[\mathbf{v}, \mathbf{v}^{-1}] \rightarrow K$, which we also call σ and is given by

$$v_{\mathcal{C},j} \xrightarrow{\sigma} 1. \tag{14}$$

Notice that, just as in (12), the induced map $\tilde{\sigma} : \mathcal{H}_{K(\mathbf{v})}(W) \rightarrow K[W]$ agrees with the fixed surjection $B(W) \twoheadrightarrow W$. The freeness over $\mathbb{Z}[\mathbf{v}, \mathbf{v}^{-1}]$, the fact that $K(\mathbf{v})$ and K are splitting fields for $\mathcal{H}(W)$ and W respectively, and the semisimplicity of $K[W]$, constitute the assumptions of Tits' deformation theorem (see [GP00, §7.3-4]). Its conclusion is then:

Theorem 4.6. *The algebra $\mathcal{H}_{K(\mathbf{v})}(W)$ is also semisimple and the specialization map σ induces a bijection*

$$d_\sigma : \text{Irr}(\mathcal{H}_{K(\mathbf{v})}(W)) \xrightarrow{\sim} \text{Irr}(K[W]),$$

between the irreducible modules of the two algebras, that respects the spectra of elements. That is, if U and $d_\sigma(U)$ are irreducible modules matched by d_σ , then the following diagram commutes:

$$\begin{array}{ccc} \mathcal{H}_{K(\mathbf{v})}(W) \ni T_{\mathbf{g}} & \xrightarrow{\mathfrak{p}_U} & K[\mathbf{v}, \mathbf{v}^{-1}][X] \\ \tilde{\sigma} \downarrow & & \downarrow t_\sigma \\ K[W] \ni g & \xrightarrow{\mathfrak{p}_{d_\sigma(U)}} & K[X] \end{array} \quad (15)$$

The horizontal maps \mathfrak{p}_M send an element $T_{\mathbf{g}}$ or g to its characteristic polynomial under the representation M , while the vertical maps are naturally induced by σ . In particular, since character values are determined by the spectra of elements, if $\chi_{\mathbf{v}}$ and χ are the characters associated to U and $d_\sigma(U)$ respectively, we will have

$$\chi(g) = \sigma(\chi_{\mathbf{v}}(T_{\mathbf{g}})). \quad (16)$$

Remark: It is not a priori clear that the characteristic polynomials of elements $T_{\mathbf{g}}$ live in $K[\mathbf{v}, \mathbf{v}^{-1}][X]$ (instead of just $K(\mathbf{v})[X]$); this is shown in [GP00, Prop. 7.3.8]. The existence of the map d_σ and that it respects spectra is proved in [ibid, Thm. 7.4.3], and the fact that it is a bijection in [ibid, Thm. 7.4.6]. \square

We can apply Tits' deformation theorem on any admissible (see Defn. 4.3) specialization of $\mathcal{H}(W)$ by first moving to a splitting field as prescribed by Prop. 4.5. In particular, for the algebras $\mathcal{H}_{\mathbf{x}}(W)$ and $\mathcal{H}_x(W)$ from (13), the corresponding splitting fields have to be $K(\mathbf{y})$ and $K(y)$ respectively for parameters $\mathbf{y} := (y_{\mathcal{C}})_{\mathcal{C} \in \mathcal{A}/W}$ and y that satisfy $y_{\mathcal{C}}^{N_W} = x_{\mathcal{C}}$ and $y^{N_W} = x$.

Now Thm. 4.6 implies that we can simultaneously index the characters of $\mathcal{H}(W)$, $\mathcal{H}_{\mathbf{x}}(W)$, and $\mathcal{H}_x(W)$ by characters $\chi \in \widehat{W}$. Indeed, if say $f_{\mathbf{x}}$ is the factoring morphism of Defn. 4.3, we have

$$\text{Irr}(\mathcal{H}(W)) \xrightarrow{d_{\theta_{\mathbf{x}}}} \text{Irr}(\mathcal{H}_{\mathbf{x}}(W)) \xrightarrow{d_{f_{\mathbf{x}}}} \text{Irr}(K[W]), \quad (17)$$

where $d_{\theta_{\mathbf{x}}}$ and $d_{f_{\mathbf{x}}}$ are bijections which satisfy $d_\sigma = d_{\theta_{\mathbf{x}}} \circ d_{f_{\mathbf{x}}}$ and moreover respect spectra as in (15). We will therefore denote the characters of the three Hecke algebras by $\chi_{\mathbf{v}}$, $\chi_{\mathbf{y}}$, and χ_y respectively, using the parameters $\mathbf{v}, \mathbf{y}, y$ that define the splitting fields.

Definition 4.7. We say that a character of the Hecke algebra $\mathcal{H}(W)$ is *rational* with respect to the specializations $\theta_{\mathbf{x}}$ or θ_x (respectively *generically rational*) if its values lie in $K(\mathbf{x})$ or $K(x)$ (respectively in $K(\mathbf{u})$), as opposed to the splitting fields. Similarly we talk of a rational spectrum of some element $T_{\mathbf{g}}$ for a given representation and specialization.

Remark 4.8. Notice that a character might be rational for the specialization θ_x but not for $\theta_{\mathbf{x}}$. This is for instance the case when a monomial of the form $\sqrt{x_{\mathcal{C},0} x_{\mathcal{C}',0}}$ appears as its value (which is not rational for $\theta_{\mathbf{x}}$ but becomes x for θ_x). For example, the group G_6 has 6 characters that are not generically rational (see [Mal99, Table 8.1]) but a CHEVIE [Gec+96; Mic15] calculation shows only 2 irrational characters for θ_x .

4.2 Character values on roots of the full twist

For a character $\chi_{\mathbf{v}}$ of the generic Hecke algebra $\mathcal{H}_{K(\mathbf{v})}(W)$, let $m_{\mathcal{C},j}^{\chi_{\mathbf{v}}}$ denote the multiplicity of $u_{\mathcal{C},j}$ as an eigenvalue of any braid reflection $s_{\mathcal{C},\gamma}$ in the representation U associated with $\chi_{\mathbf{v}}$. After Tits' deformation theorem (in particular, after (15)) this equals the multiplicity of $\zeta_{e_{\mathcal{C}}}^j = \sigma(u_{\mathcal{C},j})$ as an eigenvalue of any *distinguished* reflection s_H , $H \in \mathcal{C}$, in the representation $d_{\sigma}(U)$.

The same is true for any admissible specialization θ (notice that since $f \circ \theta(u_{\mathcal{C},j}) = \zeta_{e_{\mathcal{C}}}^j$, the elements $\theta(u_{\mathcal{C},j})$ cannot be equal), so for the analogously defined numbers $m_{\mathcal{C},j}^{\chi_{\mathbf{v}}}$, $m_{\mathcal{C},j}^{\chi_{\mathbf{y}}}$, $m_{\mathcal{C},j}^{\chi}$, we have

$$m_{\mathcal{C},j}^{\chi_{\mathbf{v}}} = m_{\mathcal{C},j}^{\chi_{\mathbf{y}}} = m_{\mathcal{C},j}^{\chi_{\mathbf{y}}} = m_{\mathcal{C},j}^{\chi}.$$

In view of this, we will only use the latter notation $m_{\mathcal{C},j}^{\chi}$ from now on. Notice finally that by the defining relations (11), the only possible eigenvalues for any $s_{\mathcal{C},\gamma}$ are precisely the $u_{\mathcal{C},j}$'s. We therefore have (for any $\mathcal{C} \in \mathcal{A}/W$)

$$\sum_{j=0}^{e_{\mathcal{C}}-1} m_{\mathcal{C},j}^{\chi} = \chi(1). \quad (18)$$

The following proposition is essential for the proof of our technical lemma (Prop. 4.19). For completion, we reproduce here its proof following [BM97, §4: E] very closely. To simplify its statement we first introduce the following notation (recall also that $\omega_{\mathcal{C}} = |\mathcal{C}|$ for an orbit $\mathcal{C} \in \mathcal{A}/W$):

Definition 4.9. Consider⁴ the element of $K[\mathbf{u}^{1/|W|}]$ given as

$$z_{\chi_{\mathbf{v}}}(\boldsymbol{\pi}) := \prod_{\mathcal{C} \in \mathcal{A}/W} \prod_{j=0}^{e_{\mathcal{C}}-1} u_{\mathcal{C},j}^{(1/\chi(1))m_{\mathcal{C},j}^{\chi} e_{\mathcal{C}} \omega_{\mathcal{C}}},$$

and, for a regular number d (see Defn. 2.8), write

$$z_{\chi_{\mathbf{v}}}(\boldsymbol{\pi})^{1/d} := \prod_{\mathcal{C} \in \mathcal{A}/W} \prod_{j=0}^{e_{\mathcal{C}}-1} u_{\mathcal{C},j}^{(1/d\chi(1))m_{\mathcal{C},j}^{\chi} e_{\mathcal{C}} \omega_{\mathcal{C}}}.$$

Finally, denote by $N(\chi)$ the quantity

$$N(\chi) := \sum_{\mathcal{C} \in \mathcal{A}/W} \omega_{\mathcal{C}} \cdot \sum_{j=0}^{e_{\mathcal{C}}-1} j m_{\mathcal{C},j}^{\chi}.$$

Remark 4.10. $N(\chi)$ usually denotes the sum of the χ^* -exponents (see [LT09, Chapter 4: §4]) of the representation that affords χ . This in fact agrees with the definition above (see [BM97, Prop. 4.1], or [LT09, Lemma 10.15 and Remark 10.12] which includes Gutkin's theorem). We are only going to use it as a symbol (but see also Remark 4.15).

Proposition 4.11. [BM97, Prop. 4.16]

For a character $\chi_{\mathbf{v}}$ of the generic Hecke algebra, the values on the full twist $T_{\boldsymbol{\pi}}$ are given by

$$\chi_{\mathbf{v}}(T_{\boldsymbol{\pi}}) = \chi(1) e^{-2i\pi N(\chi)/\chi(1)} z_{\chi_{\mathbf{v}}}(\boldsymbol{\pi}).$$

Moreover, if \mathbf{w} is a d -th root of some power $\boldsymbol{\pi}^l$ and its image in W under the fixed surjection (5) is w , we have

$$\chi_{\mathbf{v}}(T_{\mathbf{w}}) = \chi(w) e^{-2i\pi l N(\chi)/d\chi(1)} z_{\chi_{\mathbf{v}}}(\boldsymbol{\pi})^{l/d}.$$

⁴We move to a larger ring, so that the roots $u_{\mathcal{C},j}^{1/\chi(1)}$ are well defined. Shortly however, Prop. 4.11 will show that these monomials actually live in $K[\mathbf{v}]$.

Proof. We are only going to prove the second statement, which reduces to the first for $d = l = 1$.

Consider the determinant character \det_{χ_v} associated to χ_v . Because it is linear, it factors through the abelianization B^{ab} and its values on powers of the full twist are given after Corol. 2.7 by

$$\det_{\chi_v}(T_{\boldsymbol{\pi}}^l) = \prod_{\mathcal{C} \in \mathcal{A}/W} \prod_{j=0}^{e_{\mathcal{C}}-1} u_{\mathcal{C},j}^{m_{\mathcal{C},j}^{\chi} \omega_{\mathcal{C}} e_{\mathcal{C}} l} = z_{\chi_v}(\boldsymbol{\pi})^{\chi(1)l}.$$

Now, since $\boldsymbol{\pi}^l$ is central in $B(W)$ (and therefore also in $\mathcal{H}_{K(v)}(W)$), it acts on irreducible representations as a scalar. That is, its spectrum is given by

$$\text{Spec}_{\chi_v}(T_{\boldsymbol{\pi}}^l) = \{\xi z_{\chi_v}(\boldsymbol{\pi})^l \text{ (}\chi(1)\text{-many times)}\},$$

where ξ is a $\chi(1)$ -th root of unity. Since $\boldsymbol{w}^d = \boldsymbol{\pi}^l$, we further have that the spectrum of $T_{\boldsymbol{w}}$ is

$$\text{Spec}_{\chi_v}(T_{\boldsymbol{w}}) = \{\xi_k z_{\chi_v}(\boldsymbol{\pi})^{l/d} \mid (1 \leq k \leq \chi(1))\},$$

where the ξ_k are d -th roots of ξ , which of course means that

$$\chi_v(T_{\boldsymbol{w}}) = z_{\chi_v}(\boldsymbol{\pi})^{l/d} \sum_{k=1}^{\chi(1)} \xi_k.$$

We are only left with computing the sum of the ξ_k . Notice that after Tits' deformation theorem (in particular, the statement (16) that character values are respected), we will have for the specialization σ of (14) that $\sigma(\chi_v(T_{\boldsymbol{w}})) = \chi(w)$. Since the right hand side of the previous equation will then also have to evaluate to $\chi(w)$ under σ , we will have that

$$\sum_{k=1}^{\chi(1)} \xi_k = \chi(w) \sigma(z_{\chi_v}(\boldsymbol{\pi})^{l/d})^{-1}.$$

Finally, recalling Defn. 4.9 and that $\sigma(u_{\mathcal{C},j}) = \zeta_{e_{\mathcal{C}}}^j = \exp(2i\pi j/e_{\mathcal{C}})$, it is easy to see that

$$\sigma(z_{\chi_v}(\boldsymbol{\pi})^{l/d}) = \exp\left(2i\pi l/(d\chi(1)) \sum_{\mathcal{C} \in \mathcal{A}/W} \omega_{\mathcal{C}} \sum_{j=1}^{e_{\mathcal{C}}-1} j m_{\mathcal{C},j}^{\chi}\right) = e^{2i\pi l N(\chi)/d\chi(1)}.$$

□

By applying the specialization $\theta_{\boldsymbol{x}}$ from (13) on the previous proposition, we easily get:

Corollary 4.12. *Let \boldsymbol{w} be a d -th root of some power $\boldsymbol{\pi}^l$ as above and let $\chi_{\boldsymbol{y}}$ be a character of the specialization $\mathcal{H}_{\boldsymbol{x}}(W)$ as in (17). We have*

1.
$$\chi_{\boldsymbol{y}}(T_{\boldsymbol{\pi}}) = \chi(1) \prod_{\mathcal{C} \in \mathcal{A}/W} x_{\mathcal{C}}^{(1/\chi(1))m_{\mathcal{C},0}^{\chi} e_{\mathcal{C}} \omega_{\mathcal{C}}}.$$
2.
$$\chi_{\boldsymbol{y}}(T_{\boldsymbol{w}}) = \chi(w) \prod_{\mathcal{C} \in \mathcal{A}/W} x_{\mathcal{C}}^{(l/d\chi(1))m_{\mathcal{C},0}^{\chi} e_{\mathcal{C}} \omega_{\mathcal{C}}}.$$

4.3 Local Coxeter numbers

We are now going to define a local version of Coxeter numbers (see Defn. 3.3) and study how they are precisely related to the exponents that appear in the character calculation of the previous Corol. 4.12.

Definition 4.13. We define the *local Coxeter number* $c_{\chi,c}$ associated to the character χ and the hyperplane orbit $\mathcal{C} \in \mathcal{A}/W$, as the normalized trace

$$c_{\chi,c} := \frac{1}{\chi(\mathbf{1})} \cdot \chi\left(\sum_{V^t \in \mathcal{C}} (\mathbf{1} - t)\right).$$

Here, the sum is taken over all reflections t whose fixed hyperplane $H = V^t$ belongs to the orbit \mathcal{C} . Notice that these numbers are a refinement of the Coxeter numbers in the sense that $c_\chi = \sum c_{\chi,c}$

Proposition 4.14. *The local Coxeter numbers satisfy*

$$c_{\chi,c} = e_c \cdot \omega_c \cdot \left(1 - \frac{m_{\mathcal{C},0}^\chi}{\chi(\mathbf{1})}\right).$$

Proof. As we saw in (4), because the parabolic groups for hyperplanes are cyclic, the set of reflections can be partitioned into sets of the form $\{t_H, \dots, t_H^{e_H-1}\}$. Moreover, recalling the definition of $m_{\mathcal{C},j}^\chi$ from the beginning of this section, we see that the spectrum of t_H^k (for $H \in \mathcal{C}$) is given by

$$\text{Spec}_\chi(t_H^k) = \{\zeta_{e_c}^{jk} (m_{\mathcal{C},j}^\chi\text{-many times}) \mid 0 \leq j \leq e_c - 1\}.$$

We can then pick an $H \in \mathcal{C}$ and a generator t_H of W_H , and start the evaluation by computing

$$\begin{aligned} \sum_{V^t \in \mathcal{C}} \chi(\mathbf{1} - t) &= \chi(\mathbf{1})(e_c - 1)\omega_c - \omega_c \sum_{k=1}^{e_c-1} \chi(t_H^k) \\ &= \chi(\mathbf{1})(e_c - 1)\omega_c - \omega_c \sum_{k=1}^{e_c-1} \sum_{j=0}^{e_c-1} m_{\mathcal{C},j}^\chi \zeta_{e_c}^{jk}. \end{aligned}$$

Now, notice that the sum $\sum_{k=1}^{e_c-1} \zeta_{e_c}^{jk}$ equals $e_c - 1$ or -1 depending on whether $j = 0$ or not. So, after changing the order of summation, we have

$$\begin{aligned} \sum_{V^t \in \mathcal{C}} \chi(\mathbf{1} - t) &= \chi(\mathbf{1})(e_c - 1)\omega_c + \sum_{j=1}^{e_c-1} \omega_c m_{\mathcal{C},j}^\chi - (e_c - 1)\omega_c m_{\mathcal{C},0}^\chi \\ &= \chi(\mathbf{1})e_c \omega_c - e_c \omega_c m_{\mathcal{C},0}^\chi, \end{aligned}$$

where the second equation is because of (18). This completes the proof. \square

We can now rewrite the character calculation from Corol. 4.12 replacing the quantities in the exponents with equivalent ones in terms of the Coxeter numbers $c_{\chi,c}$ (and via Prop. 4.14). With the notation being the same as in the statement of the Corollary, we have:

$$\chi_{\mathbf{y}}(T_{\boldsymbol{\pi}}) = \chi(\mathbf{1}) \prod_{\mathcal{C} \in \mathcal{A}/W} x_{\mathcal{C}}^{e_c \omega_c - c_{\chi,c}} \quad \text{and} \quad \chi_{\mathbf{y}}(T_{\mathbf{w}}) = \chi(w) \prod_{\mathcal{C} \in \mathcal{A}/W} x_{\mathcal{C}}^{(e_c \omega_c - c_{\chi,c})l/d}. \quad (19)$$

Moreover, after the further specialization $x_{\mathcal{C}} \rightarrow x$ of θ_x from (13) and for the characters $\chi_{\mathbf{y}}$ of $\mathcal{H}_x(W)$ as in (17), we have (recalling that $\sum e_c \omega_c = |\mathcal{R}| + |\mathcal{A}|$ and that $\sum c_{\chi,c} = c_\chi$):

$$\chi_{\mathbf{y}}(T_{\boldsymbol{\pi}}) = \chi(\mathbf{1}) \cdot x^{|\mathcal{R}| + |\mathcal{A}| - c_\chi} \quad \text{and} \quad \chi_{\mathbf{y}}(T_{\mathbf{w}}) = \chi(w) \cdot x^{(|\mathcal{R}| + |\mathcal{A}| - c_\chi)l/d}. \quad (20)$$

Remark 4.15. This last equation is precisely what appears in [BM97, Prop. 4.18] but with an equivalent expression for the Coxeter numbers:

$$c_\chi = \frac{N(\chi) + N(\chi^*)}{\chi(1)},$$

where the numbers $N(\chi)$ are given in Defn. 4.9 (see also Rem. 4.10). This expression also appears in [Mic16, Lemma 1] but the statement of that Lemma might be misleading as it holds regardless of the values e_C . For completion, we include the calculation:

$$\chi(1)c_\chi = \chi(1) \sum_{c \in \mathcal{A}/W} c_{\chi,c} = \sum_{c \in \mathcal{A}/W} \omega_c \sum_{j=1}^{e_c-1} e_c m_{c,j}^\chi = N(\chi) + N(\chi^*).$$

In fact, Michel later on [Mic16, Rem. 2] notes that for all groups W one has (see Defn. 4.17)

$$c_\chi = \frac{N(\chi) + N(\Psi(\chi^*))}{\chi(1)},$$

which is equivalent to the first statement as $N(\Psi(\chi)) = N(\chi)$ after Prop. 4.18.

For the proof of the integer property in the following corollary, we again follow [BM97] closely and reproduce the argument here for completion.

Corollary 4.16. [BM97, Corol. 4.17] *The Coxeter numbers $c_{\chi,c}$ are integers and they satisfy*

$$0 \leq c_{\chi,c} \leq e_c \cdot \omega_c.$$

Proof. The inequalities are immediate from Prop. 4.14, since $0 \leq m_{c,0}^\chi \leq \chi(1)$. To see that the numbers $c_{\chi,c}$ are integers, it is enough to show that the values $\chi_{\mathbf{y}}(T_\pi)$ given in (19) belong to $K(\mathbf{x})$ (as opposed to the splitting field $K(\mathbf{y})$ of $\mathcal{H}_{\mathbf{x}}(W)$, see above (17)). In other words, we must show that the characters $\chi_{\mathbf{y}}$ take rational values (see Defn. 4.7) on the full twist T_π .

Consider any Galois automorphism $\sigma \in \text{Gal}(K(\mathbf{y})/K(\mathbf{x}))$ of the field extension. Then the Galois-conjugate character should satisfy $\sigma(\chi_{\mathbf{y}})(T_\pi) = \zeta_\sigma \chi_{\mathbf{y}}(T_\pi)$ for some root of unity ζ_σ , because as we see in (19) $\chi_{\mathbf{y}}(T_\pi)$ is a monomial in \mathbf{y} (recall $y_c^{N_W} = x_c$). Now if we also call $(\sigma\chi)$ the irreducible character of W that corresponds to $\sigma(\chi_{\mathbf{y}})$ via Tits' deformation theorem (but keep in mind that $(\sigma\chi)$ is *not* necessarily a Galois conjugate of χ), the previous equation implies

$$(\sigma\chi)(1) = \zeta_\sigma \chi(1),$$

which of course can only be true if $\zeta_\sigma = 1$. The only way this can be true for *any* choice of σ is if the character value was rational to begin with. \square

4.4 Malle's character permutations and the technical lemma

The fake degree $P_\chi(q) := \sum q^{e_i(\chi)}$ of an irreducible character $\chi \in \widehat{W}$ is a polynomial that records the exponents $e_i(\chi)$ of the character (see [LT09, §4.4]). Beynon and Lusztig [BL78, Prop. A] had observed a remarkable reciprocity property for these polynomials. They satisfy

$$P_\chi(q) = q^{e_\chi} P_{i(\chi)}(q^{-1}),$$

where c_χ is the Coxeter number⁵ as given in Defn. 3.3 and ι is a permutation of the irreducible characters that for Weyl groups is the identity apart from two characters of E_7 and four of E_8 .

Malle later on [Mal99, Thm. 6.5] extended this reciprocity result for all complex reflection groups, defining a permutation of the characters Ψ that is induced by a Galois action on the irreducible characters of the Hecke algebra (the two permutations satisfy $\iota(\chi) = \Psi(\chi^*)$). This permutation of Malle is exactly the missing ingredient for the proof of Lemma 3.5; the characters χ for which c_χ is not a multiple of $|g|$ are grouped together by Ψ and their contributions cancel.

A Galois action on the characters

Recall (see (13) and (17)) the specializations of the Hecke algebra $\mathcal{H}_x(W)$ and $\mathcal{H}_y(W)$ that have coefficient fields $K(\mathbf{x})$ and $K(x)$, and splitting fields $K(\mathbf{y})$ and $K(y)$ respectively. Recall also that, after Prop. 4.5 the parameters satisfy $y_c^{N_W} = x_c$ and $y^{N_W} = x$.

Definition 4.17. We consider the permutations $\Psi_{\mathcal{C}}$ and Ψ acting on the sets $\text{Irr}(\mathcal{H}_x(W))$ and $\text{Irr}(\mathcal{H}_y(W))$ that are respectively induced by the Galois automorphisms $\Sigma_{\mathcal{C}}$ (for $\mathcal{C} \in \mathcal{A}/W$) and Σ :

$$\begin{aligned} \Sigma_{\mathcal{C}} \in \text{Gal}(K(\mathbf{y})/K(\mathbf{x})) & & \Sigma \in \text{Gal}(K(y)/K(x)) \\ y_{\mathcal{C}} \rightarrow e^{2\pi i/N_W} \cdot y_{\mathcal{C}} & & y \rightarrow e^{2\pi i/N_W} \cdot y \end{aligned}$$

In particular, they are defined via $\Psi_{\mathcal{C}}(\chi_{\mathbf{y}})(T_{\mathbf{g}}) := \Sigma_{\mathcal{C}}(\chi_{\mathbf{y}}(T_{\mathbf{g}}))$ and similarly for Ψ . By Tits' deformation theorem, they induce permutations on the set \widehat{W} of irreducible characters of W , which we also denote by $\Psi_{\mathcal{C}}$ and Ψ .

The permutations $\Psi_{\mathcal{C}}$ and Ψ satisfy a set of properties with respect to the Coxeter numbers and other statistics of the characters $\chi \in \widehat{W}$:

Proposition 4.18. *For any character $\chi \in \widehat{W}$ and orbits $\mathcal{C}, \mathcal{C}' \in \mathcal{A}/W$, the following are true:*

1. $\Psi_{\mathcal{C}}(\chi)(1) = \chi(1)$
2. $m_{\mathcal{C}',j}^{\Psi_{\mathcal{C}}(\chi)} = m_{\mathcal{C}',j}^{\chi}$
3. $c_{\Psi_{\mathcal{C}}(\chi),\mathcal{C}'} = c_{\chi,\mathcal{C}'}$

Proof. Since $\Psi_{\mathcal{C}}$ is induced by a Galois automorphism, it has to respect the degree of the character $\chi_{\mathbf{y}}$, hence also of χ ; this proves part 1. The spectrum of any braid reflection $\mathbf{s}_{\mathcal{C}',\gamma}$ is generically rational (see Defn. 4.7) by the defining relations (11). This means that the eigenvalues of any $\mathbf{s}_{\mathcal{C}',\gamma}$ in the representation that affords $\chi_{\mathbf{y}}$ live in the coefficient field $K(\mathbf{x})$ and are therefore fixed by $\Psi_{\mathcal{C}}$. This proves part 2. after recalling the definition of $m_{\mathcal{C}',j}^{\chi}$ from the start of §4.2 and also part 3. after Prop. 4.14. The same results are of course true for Ψ . \square

The following is the key technical lemma that we have been building towards through all of Section 4. The character calculations of Prop. 4.11 were included just so that the argument presented here is self-contained.

Proposition 4.19 (The key technical lemma).

Let g be a ζ -regular element of W , with $\zeta = e^{2\pi i l/d}$ of order d , $\chi \in \widehat{W}$ an irreducible character, and $\mathcal{C} \in \mathcal{A}/W$ an orbit of hyperplanes. Then, we have

$$\Psi_{\mathcal{C}}(\chi)(g) = \exp\left(-2\pi i \cdot \frac{lc_{\chi,\mathcal{C}}}{d}\right) \cdot \chi(g) \quad \text{and} \quad \Psi(\chi)(g) = \exp\left(-2\pi i \cdot \frac{lc_{\chi}}{d}\right) \cdot \chi(g).$$

⁵However, Beynon and Lusztig, and later Malle, did not assign an epithet for these numbers; the mathematical godfathers were Gordon and Griffeth [GG12] who named them after Coxeter.

Proof. By Prop. 2.9, we can lift g to some element $\mathbf{g} \in B(W)$ that is commensurable with the full twist (i.e. it satisfies $\mathbf{g}^d = \boldsymbol{\pi}^l$ with $(l, d) = 1$). Now, replacing $x_{\mathcal{C}}$ with $y_{\mathcal{C}}^{N_W}$ we can rewrite the character evaluations from (19) as

$$\chi_{\mathbf{y}}(T_{\mathbf{g}}) = \chi(g) \cdot \prod_{\mathcal{C}' \in \mathcal{A}/W} y_{\mathcal{C}'}^{N_W(e_{\mathcal{C}'}\omega_{\mathcal{C}'} - c_{\chi, \mathcal{C}'})l/d},$$

which, after applying the Galois automorphism $\Sigma_{\mathcal{C}}$, becomes

$$\Psi_{\mathcal{C}}(\chi_{\mathbf{y}})(T_{\mathbf{g}}) = \chi(g) \cdot e^{2\pi i(e_{\mathcal{C}}\omega_{\mathcal{C}} - c_{\chi, \mathcal{C}})l/d} \cdot \prod_{\mathcal{C}' \in \mathcal{A}/W} y_{\mathcal{C}'}^{N_W(e_{\mathcal{C}'}\omega_{\mathcal{C}'} - c_{\chi, \mathcal{C}'})l/d}.$$

Now, this is really

$$\Psi_{\mathcal{C}}(\chi_{\mathbf{y}})(T_{\mathbf{g}}) = e^{2\pi i(e_{\mathcal{C}}\omega_{\mathcal{C}} - c_{\chi, \mathcal{C}})l/d} \cdot \chi_{\mathbf{y}}(T_{\mathbf{g}}),$$

which completes the proof after applying Tits' deformation theorem and recalling that $e_{\mathcal{C}}\omega_{\mathcal{C}}$ is a multiple of d by Corol. 2.11. The same argument of course works for Ψ . \square

We are now ready to prove Lemma 3.5. Only Malle's permutation Ψ is sufficient for that, while the "local" version $\Psi_{\mathcal{C}}$ will be used in Section 5 to deduce similar results for generating functions of weighted reflection factorizations.

Proof of Lemma 3.5. We consider the partition of the set of irreducible characters $\chi \in \widehat{W}$ into orbits under the action of Ψ . We will show that the total contribution of the characters in any orbit that is not a singleton is 0.

Consider a character χ in such an orbit and let k be the smallest number such that $\Psi^k(\chi) = \chi$. Notice, that after Prop. 4.19 we must have $k = \frac{d}{\gcd(c_{\chi}, d)}$. Since by Prop. 4.18 the degrees $\chi(1)$ as well as the Coxeter numbers c_{χ} are not affected by Ψ , it is sufficient to show that

$$\sum_{j=1}^k \Psi^j(\chi)(g^{-1}) = 0.$$

But if $\xi = \exp(-2\pi i c_{\chi, \mathcal{C}}/d)$, we have by Prop. 4.19 that $\Psi^j(\chi)(g^{-1}) = \xi^j \chi(g^{-1})$ after which the above is immediate (indeed, ξ is also a k^{th} root of unity). \square

Remark 4.20. Notice that Prop. 4.19 gives some insight on why in Weyl groups the orbits under Ψ can have at most two elements. Indeed, every regular element g will come with (at least) a pair of regular eigenvalues $e^{\pm 2\pi i l/d}$. Since we can lift g to d^{th} roots of either powers $\boldsymbol{\pi}^l$ and $\boldsymbol{\pi}^{d-l}$, the only way the proposition is valid for both lifts is if $\gcd(c_{\chi, \mathcal{C}}, d) \leq 2$ or $\chi(g) = 0$.

More generally, for a given χ and \mathcal{C} , Prop. 4.19 implies that if $\chi(g) \neq 0$, then $l \cdot c_{\chi, \mathcal{C}} \pmod{d}$ is constant for all l such that $\zeta = e^{2\pi i l/d}$ is a regular eigenvalue of g .

4.5 On the uniformity of the proofs

Our proofs rely so far mainly on two properties that are known in a case-by-case fashion; the BMR-freeness theorem and the structure of the splitting fields for the Hecke algebras. Both of those are known uniformly for real reflection groups ([GP00, Thm. 4.4.6] and [Opd95, Thm. 5]).

In fact, we could do away with the second reliance. Opdam's work [Opd98, Thm. 6.7] is sufficient information for the structure of the group $\text{Gal}(\mathbb{C}(\mathbf{v})/\mathbb{C}(\mathbf{u}))$ which in turn is all we need to define the permutations $\Psi_{\mathcal{C}} \in \text{Perm}(\text{Irr}(W))$. In fact Opdam's elements $g_{\mathcal{C}, 0}$ of this Galois

group correspond precisely to our $\Sigma_{\mathcal{C}}$ of Defn. 4.17 (see [ibid, Prop. 7.1] and the discussion before [ibid, Prop. 7.4]). We have chosen not to follow Opdam’s presentation here (which involves the KZ-connection, a much more complicated beast) even if it is more uniform, as it does not eventually illuminate Prop. 4.19 much better.

As far as the BMR-freeness theorem goes, and again because we are really interested in the “geometric” Galois group $\text{Gal}(\mathbb{C}(\mathbf{v})/\mathbb{C}(\mathbf{u}))$, it is possible that we could replace it by Losev’s weaker but uniform theorem [Los15]. We hope to be able to clarify this in the future.

5 The weighted enumeration

The following section studies the weighted enumeration of reflection factorizations as considered in [dHR18], where each reflection $t \in \mathcal{R}$ is weighted by the orbit $\mathcal{C} \in \mathcal{A}/W$ of its fixed hyperplane V^t . It provides a uniform proof of their result and extends it in a similar direction as with the Chapuy-Stump formula (3). Again we assume that W is irreducible (but see §5.1).

Definition 5.1. Consider a set of variables $\mathbf{w} := (w_{\mathcal{C}})_{(\mathcal{C} \in \mathcal{A}/W)}$ and a weight function

$$\text{wt} : \mathcal{R} \rightarrow \{w_{\mathcal{C}} \mid \mathcal{C} \in \mathcal{A}/W\},$$

such that $\text{wt}(t) = w_{\mathcal{C}}$ if \mathcal{C} is the orbit that contains the fixed hyperplane V^t . Then, the weighted enumeration of reflection factorizations of some element $g \in W$ is encoded via the following generating function:

$$\text{FAC}_{W,g}(\mathbf{w}, z) := \sum_{\substack{(t_1, \dots, t_N) \in \mathcal{R}^N \\ t_1 \cdots t_N = g}} \text{wt}(t_1) \cdots \text{wt}(t_N) \cdot \frac{z^N}{N!}.$$

Because the sets $\mathcal{C}^{\text{ref}} := \{t \in \mathcal{R} \mid V^t \in \mathcal{C}\}$ are closed under conjugation, the Lemma of Frobenius can again be used to express $\text{FAC}_{W,g}(\mathbf{w}, z)$ as a finite sum of exponentials. Notice first, that the order of the subsets A_i in Thm. 3.1 does not affect the enumeration as the different sets of factorizations have the same size. Indeed, one can easily construct a bijective map by considering a sequence of *Hurwitz moves*:

$$(t_1, t_2, \dots, t_k, t_{k+1}, \dots, t_l) \rightarrow (t_1, t_2, \dots, t_k t_{k+1} t_k^{-1}, t_k, \dots, t_l).$$

Having said that, and assuming there are $r = |\mathcal{A}/W|$ different orbits of hyperplanes, denoted $\mathcal{C}_1, \dots, \mathcal{C}_r$, Thm. 3.1 now implies that

$$\begin{aligned} \text{FAC}_{W,g}(\mathbf{w}, z) &= \sum_{\substack{N \geq 0 \\ l_1 + \dots + l_r = N}} \binom{N}{l_1, \dots, l_r} \times \\ &\quad \times \frac{1}{|W|} \sum_{\chi \in \widehat{W}} \chi(1) \cdot \chi(g^{-1}) \cdot \left[\frac{\chi(\mathcal{C}_1^{\text{ref}})}{\chi(1)} \right]^{l_1} \cdots \left[\frac{\chi(\mathcal{C}_r^{\text{ref}})}{\chi(1)} \right]^{l_r} \cdot w_{\mathcal{C}_1}^{l_1} \cdots w_{\mathcal{C}_r}^{l_r} \frac{z^N}{N!}. \end{aligned}$$

Using standard properties of exponential generating functions, we can rewrite the sum as

$$\text{FAC}_{W,g}(\mathbf{w}, z) = \frac{1}{|W|} \sum_{\chi \in \widehat{W}} \chi(1) \cdot \chi(g^{-1}) \cdot \exp \left[z w_{\mathcal{C}_1} \cdot \frac{\chi(\mathcal{C}_1^{\text{ref}})}{\chi(1)} \right] \cdots \exp \left[z w_{\mathcal{C}_r} \cdot \frac{\chi(\mathcal{C}_r^{\text{ref}})}{\chi(1)} \right].$$

Finally, notice that by Defn. 4.13 we can rewrite the quantities in the exponentials in terms of local Coxeter numbers. Indeed, we have $c_{\chi, \mathcal{C}} = |\mathcal{C}^{\text{ref}}| - \chi(\mathcal{C}^{\text{ref}})/\chi(1)$ and if we define $\text{wt}(\mathcal{R}) := \sum_{t \in \mathcal{R}} \text{wt}(t)$, the previous expression becomes a direct analog of (10) :

$$\text{FAC}_{W,g}(\mathbf{w}, z) = \frac{e^{z \cdot \text{wt}(\mathcal{R})}}{|W|} \sum_{\chi \in \widehat{W}} \chi(1) \cdot \chi(g^{-1}) \cdot (e^{-z w_{c_1}})^{c_{\chi, c_1}} \dots (e^{-z w_{c_r}})^{c_{\chi, c_r}}. \quad (21)$$

Lemma 5.2. *For a complex reflection group W , and a **regular** element $g \in W$, the total contribution in (21) of those characters $\chi \in \widehat{W}$ for which **any** $c_{\chi, c}$ is not a multiple of $|g|$ is 0.*

Proof. The proof is essentially the same as for Lemma 3.5. However, we first need to order the orbits $\mathcal{C} \in \mathcal{A}/W$ (arbitrarily) and then apply the same idea sequentially.

We start by partitioning the set of irreducible characters $\chi \in \widehat{W}$ into orbits under the action of $\Psi_{\mathcal{C}_1}$. Pick a character χ whose orbit is *not* a singleton and let k be the smallest number such that $\Psi_{\mathcal{C}_1}^k(\chi) = \chi$ (again, we will have $k = \frac{|g|}{\gcd(c_{\chi, \mathcal{C}_1}, |g|)}$). Now, since by Prop. 4.18 the degrees of characters and the (local) Coxeter numbers are respected by $\Psi_{\mathcal{C}_1}$, it is enough to show that

$$\sum_{j=1}^k \Psi_{\mathcal{C}_1}^j(\chi)(g^{-1}) = 0.$$

Indeed, this follows immediately from Prop. 4.19 as $\Psi_{\mathcal{C}_1}^j(\chi)(g^{-1}) = \xi^j \chi(g^{-1})$ for some k^{th} root of unity ξ . Notice now that we can continue with the *remaining* characters and the orbit \mathcal{C}_2 without worrying that we might eventually cancel the same character twice. \square

Before we proceed with our structural result for weighted enumeration formulas, we introduce the following combinatorial generalizations of the length function $l_R(g)$:

Definition 5.3. For an arbitrary element $g \in W$ and an orbit $\mathcal{C} \in \mathcal{A}/W$, we define $n_{\mathcal{C}}(g)$ to be the smallest number of reflections in \mathcal{C}^{ref} that may appear in *any* reflection factorization of g (i.e. not necessarily reduced).

Remark 5.4. Notice that it is not always true that $\sum n_{\mathcal{C}}(g) = l_R(g)$. Indeed, the element $g := (12\bar{1}2) = -\mathbf{1}$ in B_2 (which is the square of the Coxeter element) can be written both as $g = (12)(\bar{1}2)$ and as $g = (1\bar{1})(2\bar{2})$, so that $n_1(g) = n_2(g) = 0$.

Theorem 5.5. *For a complex reflection group W and a regular element $g \in W$, the exponential generating function $\text{FAC}_{W,g}(\mathbf{w}, z)$ of weighted reflection factorizations of g takes the form:*

$$\text{FAC}_{W,g}(\mathbf{w}, z) = \frac{e^{z \cdot \text{wt}(\mathcal{R})}}{|W|} \cdot \left[\Phi(\mathbf{X}) \cdot \prod_{\mathcal{C} \in \mathcal{A}/W} (1 - X_{\mathcal{C}})^{n_{\mathcal{C}}(g)} \right] \Big|_{X_{\mathcal{C}} = e^{-z w_{\mathcal{C}} |g|}}.$$

Here, $\Phi(\mathbf{X})$ is a polynomial of degree $(e_{\mathcal{C}} \cdot w_{\mathcal{C}})/|g| - n_{\mathcal{C}}(g)$ on each of its variables $X_{\mathcal{C}}$, it has constant term $\Phi(\mathbf{0}) = 1$, and it is not further divisible by $(1 - X_{\mathcal{C}})$ for any $X_{\mathcal{C}}$. The exponents satisfy

$$\frac{e_{\mathcal{C}} w_{\mathcal{C}}}{|g|} \geq n_{\mathcal{C}}(g) \geq l_R(g) - \frac{|\mathcal{R}| + |\mathcal{A}| - e_{\mathcal{C}} w_{\mathcal{C}}}{|g|}.$$

Proof. The proof is very similar to that of Thm. 3.6. After Lemma 5.2, we need only consider in (21) those characters χ for which all $c_{\chi, \mathcal{C}}$ are multiples of $|g|$. This allows us to write the exponential function as

$$\text{FAC}_{W,g}(\mathbf{w}, z) = \frac{e^{z \cdot \text{wt}(\mathcal{R})}}{|W|} \cdot \tilde{\Phi}(\mathbf{X}),$$

for a polynomial $\tilde{\Phi}$ on variables $\mathbf{X} := (X_{\mathcal{C}})_{\mathcal{C} \in \mathcal{A}/W}$, by setting $X_{\mathcal{C}} = (e^{-z w_{\mathcal{C}}})^{|g|}$. By Corol. 4.16 the polynomial $\tilde{\Phi}(\mathbf{X})$ has degree $(e_{\mathcal{C}} \omega_{\mathcal{C}})/|g|$ on each of its variables $X_{\mathcal{C}}$, and it has constant term 1 since all $c_{\chi, \mathcal{C}}$ can be *simultaneously* 0 only for the trivial representation.

To find the largest power of $(1 - X_{\mathcal{C}})$ that divides $\tilde{\Phi}(\mathbf{X})$, we view $\tilde{\Phi}$ as a polynomial in the single variable $X_{\mathcal{C}}$ and treat the other $X_{\mathcal{C}'}$'s as complex scalars. This is equivalent to assigning arbitrary values on all variables $w_{\mathcal{C}'} \neq w_{\mathcal{C}}$ of the weight function in Defn. 5.1. If we further fix $z = 1$, the enumerative interpretation of $(e^{\text{wt}(\mathcal{R})}/|W|) \cdot \tilde{\Phi}(X_{\mathcal{C}})$ is then that it counts weighted reflection factorizations of g keeping track only of the number of reflections that fix a hyperplane in \mathcal{C} .

Now, as in Thm. 3.6 consider the root factorization of $\tilde{\Phi}(X_{\mathcal{C}})$:

$$\tilde{\Phi}(X_{\mathcal{C}}) = a(\alpha_1 - X_{\mathcal{C}})(\alpha_2 - X_{\mathcal{C}}) \cdots (\alpha_r - X_{\mathcal{C}}),$$

with $r = (e_{\mathcal{C}} \omega_{\mathcal{C}})/|g|$. We see again that by plugging back $X_{\mathcal{C}} = e^{-w_{\mathcal{C}}|g|}$ each root contributes a factor of either $(\alpha_i - 1)$ or $w_{\mathcal{C}}|g|$ to the leading term of the generating function. Since by Defn. 5.3 this must be a scalar multiple of $w_{\mathcal{C}}^{n_{\mathcal{C}}(g)}$, we have that $(1 - X_{\mathcal{C}})^{n_{\mathcal{C}}(g)}$ divides $\tilde{\Phi}(X_{\mathcal{C}})$ and is the largest power that does so (this furthermore proves the first inequality). Since this is true for a dense set of the complex values $X_{\mathcal{C}'}$, we in fact have that $(1 - X_{\mathcal{C}})^{n_{\mathcal{C}}(g)}$ is a maximal factor of $\tilde{\Phi}(\mathbf{X})$.

The only thing left to show is the second inequality for the $n_{\mathcal{C}}(g)$'s. To see this, we now identify all weights $w_{\mathcal{C}'}$, $\mathcal{C}' \neq \mathcal{C}$ to a single weight w , set again $z = 1$, and treat $\tilde{\Phi}$ as a polynomial on two variables $X = e^{-w|g|}$ and $X_{\mathcal{C}} = e^{-w_{\mathcal{C}}|g|}$. The general argument about $\tilde{\Phi}(\mathbf{X})$ implies that we can consider the polynomial $\Phi'(X, X_{\mathcal{C}})$ defined by

$$\Phi'(X, X_{\mathcal{C}}) := \frac{\tilde{\Phi}(X, X_{\mathcal{C}})}{(1 - X_{\mathcal{C}})^{n_{\mathcal{C}}(g)}}.$$

Now, the generating function

$$\frac{e^{\text{wt}(\mathcal{R})}}{|W|} \cdot \Phi'(X, X_{\mathcal{C}}) \cdot (1 - X_{\mathcal{C}})^{n_{\mathcal{C}}(g)}$$

counts reflection factorizations of g weighing reflections in \mathcal{C}^{ref} by $w_{\mathcal{C}}$ and the rest by w . We want to enumerate factorizations that have exactly the minimal number $n_{\mathcal{C}}(g)$ of reflections of type \mathcal{C} . Since the term $(1 - X_{\mathcal{C}})^{n_{\mathcal{C}}(g)}$ always contributes a factor of $(w_{\mathcal{C}}|g|)^{n_{\mathcal{C}}(g)}$ to the Taylor expansion, the answer to the previous question would be given by

$$\frac{|g|^{n_{\mathcal{C}}(g)}}{|W|} \times e^{\text{wt}(\mathcal{R})} \Big|_{w_{\mathcal{C}}=0} \times \Phi'(X, X_{\mathcal{C}}) \Big|_{\substack{X_{\mathcal{C}}=1 \\ X=e^{-w|g|}}}.$$

The leading term of this exponential generating function should clearly be a multiple of $w^{l_R(g) - n_{\mathcal{C}}(g)}$. As in the previous argument, this implies that $\Phi'(X, 1)$ is a multiple of $(1 - X)^{l_R - n_{\mathcal{C}}(g)}$, but since by construction its degree is equal to $\sum_{\mathcal{C}' \neq \mathcal{C}} e_{\mathcal{C}'} \omega_{\mathcal{C}'} = |\mathcal{R}| + |\mathcal{A}| - e_{\mathcal{C}} \omega_{\mathcal{C}}$, we must have

$$l_R(g) - n_{\mathcal{C}}(g) \leq |\mathcal{R}| + |\mathcal{A}| - e_{\mathcal{C}} \omega_{\mathcal{C}},$$

which completes the proof. \square

Corollary 5.6. *For a complex reflection group W and a regular element $g \in W$ of order $|g| = d_n$, the weighted reflection factorizations of g are counted by the formula:*

$$\text{FAC}_{W,g}(\mathbf{w}, z) = \frac{e^{z \cdot \text{wt}(\mathcal{R})}}{|W|} \cdot \prod_{\mathcal{C} \in \mathcal{A}/W} (1 - e^{-z w_{\mathcal{C}} |g|})^{n_{\mathcal{C}}(g)},$$

where the exponents are explicitly given by $n_{\mathcal{C}}(g) = (e_{\mathcal{C}} \omega_{\mathcal{C}}) / |g|$.

Proof. As we showed in the proof of Corol. 3.9, when g is some d_n -regular element we must have $l_R(g) = (|\mathcal{R}| + |\mathcal{A}|) / |g|$. Then the previous theorem implies that $n_{\mathcal{C}}(g) = (e_{\mathcal{C}} \omega_{\mathcal{C}}) / |g|$, which further forces the equality $\Phi(\mathbf{X}) = 1$ and hence completes the argument. \square

Remark 5.7. For well-generated groups W , we always have $|c| = d_n$ so that the previous Corollary recovers the main theorem of [dHR18] and extends it to the groups of Prop. 2.12. Notice that while in well-generated groups we have at most two orbits of hyperplanes, the groups $G_7, G_{11}, G_{15}, G_{19}$ have three orbits. For all of them but G_{15} , d_n is regular.

5.1 When W is reducible

So far to simplify the arguments, we have silently assumed everywhere that W is irreducible. This is not a real restriction though and in fact the statement of Thm. 5.5 remains true essentially as is.

Indeed, assume that $W = W_1 \times \cdots \times W_k$ acts on the space $V = V_1 \oplus \cdots \oplus V_k$, with W_i acting irreducibly on V_i . Then, a regular eigenvector $\mathbf{v} = (v_1, \dots, v_k)$ must have all v_i 's regular in their respective groups too and hence a regular element $W \ni g = g_1 \cdots g_k$ must have all g_i 's regular in the W_i 's. Moreover since reflections from different W_i 's commute, the corresponding weighted generating function would just be the product

$$\text{FAC}_{W,g}(\mathbf{w}, z) = \prod_{i=1}^k \text{FAC}_{W_i, g_i}(\mathbf{w}, z).$$

Since the hyperplane orbits $\mathcal{C} \in \mathcal{A}/W$ are the disjoint union of the orbits $\mathcal{C}' \in \mathcal{A}_i/W_i$ the statement of Thm. 5.5 remains valid if we only change the evaluation of $X_{\mathcal{C}}$ from $e^{-z w_{\mathcal{C}} |g|}$ to $e^{-z w_{\mathcal{C}} |g_i|}$, where g_i is the regular element in the group W_i that contains the orbit \mathcal{C} .

Acknowledgments

We would like to thank Jean Michel and Maria Chlouveraki for interesting discussions and their guidance à propos the Hecke algebras. We would also like to thank Guillaume Chapuy for his detailed explanation of some calculations in [CS14] and Christian Stump for showing us how to use SAGE [The18] to derive formulas like (9). It was at this vantage point that the existence of a statement like Lemma 3.5 became clear.

We would also like to thank Ivan Marin for his detailed suggestions vis-à-vis the uniformity of the proofs and Losev's theorem.

References

- [Ben76] Mark Benard. "Schur indices and splitting fields of the unitary reflection groups". In: *J. Algebra* 38.2 (1976), pp. 318–342.

- [Bes01] David Bessis. “Zariski theorems and diagrams for braid groups”. In: *Invent. Math.* 145.3 (2001), pp. 487–507.
- [Bes15] David Bessis. “Finite complex reflection arrangements are $K(\pi, 1)$ ”. In: *Ann. of Math. (2)* 181.3 (2015), pp. 809–904.
- [Bes97] David Bessis. “Sur le corps de définition d’un groupe de réflexions complexe”. In: *Comm. Algebra* 25.8 (1997), pp. 2703–2716.
- [BL78] W. M. Beynon and G. Lusztig. “Some numerical results on the characters of exceptional Weyl groups”. In: *Math. Proc. Cambridge Philos. Soc.* 84.3 (1978), pp. 417–426.
- [BM97] Michel Broué and Jean Michel. “Sur certains éléments réguliers des groupes de Weyl et les variétés de Deligne-Lusztig associées”. In: *Finite reductive groups (Luminy, 1994)*. Vol. 141. Progr. Math. Birkhäuser Boston, Boston, MA, 1997, pp. 73–139.
- [BMM14] Michel Broué, Gunter Malle, and Jean Michel. “Split spetses for primitive reflection groups”. In: *Astérisque* 359 (2014). With an erratum to [MR1712862], pp. vi+146.
- [BMR98] Michel Broué, Gunter Malle, and Raphaël Rouquier. “Complex reflection groups, braid groups, Hecke algebras”. In: *J. Reine Angew. Math.* 500 (1998), pp. 127–190.
- [Bro10] Michel Broué. *Introduction to complex reflection groups and their braid groups*. Vol. 1988. Lecture Notes in Mathematics. Springer-Verlag, Berlin, 2010, pp. xii+138.
- [Cha04] Frédéric Chapoton. “Enumerative properties of generalized associahedra”. In: *Sém. Lothar. Combin.* 51 (2004/05), Art. B51b, 16.
- [Che55] Claude Chevalley. “Invariants of finite groups generated by reflections”. In: *Amer. J. Math.* 77 (1955), pp. 778–782.
- [Cox51] H. S. M. Coxeter. “The product of the generators of a finite group generated by reflections”. In: *Duke Math. J.* 18 (1951), pp. 765–782.
- [CS14] Guillaume Chapuy and Christian Stump. “Counting factorizations of Coxeter elements into products of reflections”. In: *J. Lond. Math. Soc. (2)* 90.3 (2014), pp. 919–939.
- [D59] József Dénes. “The representation of a permutation as the product of a minimal number of transpositions, and its connection with the theory of graphs”. In: *Magyar Tud. Akad. Mat. Kutató Int. Közl.* 4 (1959), pp. 63–71.
- [Del] Pierre Deligne. *Letter to E. Looijenga, 9/3/1974*. available at <http://homepage.univie.ac.at/christian.looi>
- [dHR18] Elise delMas, Thomas Hameister, and Victor Reiner. “A refined count of Coxeter element reflection factorizations”. In: *Electron. J. Combin.* 25.1 (2018), Paper 1.28, 11.
- [Dou17] Theodosios Douvropoulos. *Applications of Geometric Techniques in Coxeter-Catalan Combinatorics*. Thesis (Ph.D.)—University of Minnesota. ProQuest LLC, Ann Arbor, MI, 2017, p. 106.
- [Eis95] David Eisenbud. *Commutative algebra*. Vol. 150. Graduate Texts in Mathematics. With a view toward algebraic geometry. Springer-Verlag, New York, 1995, pp. xvi+785.
- [Eti17] Pavel Etingof. “Proof of the Broué-Malle-Rouquier conjecture in characteristic zero (after I. Losev and I. Marin—G. Pfeiffer)”. In: *Arnold Math. J.* 3.3 (2017), pp. 445–449.
- [Gec+96] M. Geck et al. “CHEVIE – A system for computing and processing generic character tables for finite groups of Lie type, Weyl groups and Hecke algebras”. In: *Appl. Algebra Engrg. Comm. Comput.* 7 (1996), pp. 175–210.

- [GG12] Iain G. Gordon and Stephen Griffeth. “Catalan numbers for complex reflection groups”. In: *Amer. J. Math.* 134.6 (2012), pp. 1491–1502.
- [GP00] Meinolf Geck and Götz Pfeiffer. *Characters of finite Coxeter groups and Iwahori-Hecke algebras*. Vol. 21. London Mathematical Society Monographs. New Series. The Clarendon Press, Oxford University Press, New York, 2000, pp. xvi+446.
- [Hat02] Allen Hatcher. *Algebraic topology*. Cambridge University Press, Cambridge, 2002, pp. xii+544.
- [Hur01] A. Hurwitz. “Ueber die Anzahl der Riemann’schen Flächen mit gegebenen Verzweigungspunkten”. In: *Math. Ann.* 55.1 (1901), pp. 53–66.
- [Hur91] A. Hurwitz. “Ueber Riemann’sche Flächen mit gegebenen Verzweigungspunkten”. In: *Math. Ann.* 39.1 (1891), pp. 1–60.
- [Jac88] D. M. Jackson. “Some combinatorial problems associated with products of conjugacy classes of the symmetric group”. In: *J. Combin. Theory Ser. A* 49.2 (1988), pp. 363–369.
- [Kan01] Richard Kane. *Reflection groups and invariant theory*. Vol. 5. CMS Books in Mathematics/Ouvrages de Mathématiques de la SMC. Springer-Verlag, New York, 2001, pp. x+379.
- [LM16] Joel Brewster Lewis and Alejandro H. Morales. “ $GL_n(\mathbf{F}_q)$ -analogues of factorization problems in the symmetric group”. In: *European J. Combin.* 58 (2016), pp. 75–95.
- [Loo74] Eduard Looijenga. “The complement of the bifurcation variety of a simple singularity”. In: *Invent. Math.* 23 (1974), pp. 105–116.
- [Los15] Ivan Losev. “Finite-dimensional quotients of Hecke algebras”. In: *Algebra Number Theory* 9.2 (2015), pp. 493–502.
- [LRS14] J. B. Lewis, V. Reiner, and D. Stanton. “Reflection factorizations of Singer cycles”. In: *J. Algebraic Combin.* 40.3 (2014), pp. 663–691.
- [LT09] Gustav I. Lehrer and Donald E. Taylor. *Unitary reflection groups*. Vol. 20. Australian Mathematical Society Lecture Series. Cambridge University Press, Cambridge, 2009, pp. viii+294.
- [LZ04] Sergei K. Lando and Alexander K. Zvonkin. *Graphs on surfaces and their applications*. Vol. 141. Encyclopaedia of Mathematical Sciences. With an appendix by Don B. Zagier, Low-Dimensional Topology, II. Springer-Verlag, Berlin, 2004, pp. xvi+455.
- [Mal99] Gunter Malle. “On the rationality and fake degrees of characters of cyclotomic algebras”. In: *J. Math. Sci. Univ. Tokyo* 6.4 (1999), pp. 647–677.
- [Mic15] Jean Michel. “The development version of the CHEVIE package of GAP3”. In: *J. Algebra* 435 (2015), pp. 308–336.
- [Mic16] Jean Michel. “Deligne-Lusztig theoretic derivation for Weyl groups of the number of reflection factorizations of a Coxeter element”. In: *Proc. Amer. Math. Soc.* 144.3 (2016), pp. 937–941.
- [MN10] Sho Matsumoto and Jonathan Novak. “Unitary matrix integrals, primitive factorizations, and Jucys-Murphy elements”. In: *22nd International Conference on Formal Power Series and Algebraic Combinatorics (FPSAC 2010)*. Discrete Math. Theor. Comput. Sci. Proc., AN. Assoc. Discrete Math. Theor. Comput. Sci., Nancy, 2010, pp. 403–411.
- [Opd95] E. M. Opdam. “A remark on the irreducible characters and fake degrees of finite real reflection groups”. In: *Invent. Math.* 120.3 (1995), pp. 447–454.

- [Rea08] Nathan Reading. “Chains in the noncrossing partition lattice”. In: *SIAM J. Discrete Math.* 22.3 (2008), pp. 875–886.
- [Spr74] T. A. Springer. “Regular elements of finite reflection groups”. In: *Invent. Math.* 25 (1974), pp. 159–198.
- [Spr78] T. A. Springer. “A construction of representations of Weyl groups”. In: *Invent. Math.* 44.3 (1978), pp. 279–293.
- [ST54] G. C. Shephard and J. A. Todd. “Finite unitary reflection groups”. In: *Canadian J. Math.* 6 (1954), pp. 274–304.
- [Ste59] Robert Steinberg. “Finite reflection groups”. In: *Trans. Amer. Math. Soc.* 91 (1959), pp. 493–504.
- [Bes06] D. Bessis. “Garside categories, periodic loops and cyclic sets”. In: *ArXiv Mathematics e-prints* (Oct. 2006). eprint: [math/0610778](https://arxiv.org/abs/math/0610778).
- [Bou+18] C. Boura et al. “The BMM symmetrising trace conjecture for groups G_4, G_5, G_6, G_7, G_8 ”. In: *ArXiv e-prints* (Feb. 2018). arXiv: [1802.07482](https://arxiv.org/abs/1802.07482) [[math.RT](#)].
- [Fro68] Ferdinand Georg Frobenius. *Gesammelte Abhandlungen. Bd. I–III. Herausgegeben von J.-P. Serre*. German. Berlin-Heidelberg-New York: Springer-Verlag 1968. vii, 650 p.; iii, 733 p.; iv, 740 p. (1968). 1968.
- [Opd98] E. M. Opdam. “Complex Reflection Groups and Fake Degrees”. In: *ArXiv Mathematics e-prints* (Aug. 1998). eprint: [math/9808026](https://arxiv.org/abs/math/9808026).
- [The18] The Sage Developers. *SageMath, the Sage Mathematics Software System (Version 8.3.beta3)*. <http://www.sagemath.org>. 2018.

Theo Douvropoulos, IRIF, UMR CNRS 8243, UNIVERSITÉ PARIS DIDEROT, PARIS 7, FRANCE
E-mail address: douvr001@irif.fr