

# WEIERSTRASS SEMIGROUPS AT EVERY POINT OF THE SUZUKI CURVE

DANIELE BARTOLI, MARIA MONTANUCCI AND GIOVANNI ZINI

**ABSTRACT.** In this article we explicitly determine the structure of the Weierstrass semigroups  $H(P)$  for any point  $P$  of the Suzuki curve  $\mathcal{S}_q$ . As the point  $P$  varies, exactly two possibilities arise for  $H(P)$ : one for the  $\mathbb{F}_q$ -rational points (already known in the literature), and one for all remaining points. For this last case a minimal set of generators of  $H(P)$  is also provided. As an application, we construct dual one-point codes from an  $\mathbb{F}_{q^4} \setminus \mathbb{F}_q$ -point whose parameters are better in some cases than the ones constructed in a similar way from an  $\mathbb{F}_q$ -rational point.

**Keywords:** Suzuki curve, Weierstrass semigroups, algebraic-geometric codes

**2010 MSC:** Primary: 11G20. Secondary: 11R58, 14H05, 14H55.

## 1. INTRODUCTION

Let  $\mathcal{X}$  be a nonsingular, projective algebraic curve of genus  $g$  defined over a field  $\mathbb{F}$ . Let  $P$  be a point of  $\mathcal{X}$ . The *Weierstrass semigroup*  $H(P)$  at  $P$  is defined as the set of integers  $k$  such that there exists a function on  $\mathcal{X}$  having pole divisor exactly  $kP$ . It is clear that  $H(P)$  is a subset of natural numbers  $\mathbb{N} = \{0, 1, 2, \dots\}$ . By the Weierstrass gap Theorem [14, Theorem 1.6.8], the set  $G(P) := \mathbb{N} \setminus H(P)$  contains exactly  $g$  elements called *gaps*. The structure of  $H(P)$  depends on the choice of  $P$ . However, it is well known that  $H(P)$  is the same for all but a finite number of points  $P$ , namely the *Weierstrass points* of  $\mathcal{X}$ . On one hand, such points are of intrinsic interest, for example in Stöhr–Voloch Theory [15]. On the other hand, in the finite fields setting, the parameters of algebraic-geometric (AG) codes constructed from  $\mathcal{X}$  rely on the inner structure of the semigroup  $H(P)$ ; see e.g. [16].

In this context, *maximal curves*, that is algebraic curves defined over a finite field  $\mathbb{F} = \mathbb{F}_q$  having as many rational points as possible according to the Hasse–Weil bound, have been widely investigated. More precisely, an algebraic curve  $\mathcal{X}$  of genus  $g$  and defined over  $\mathbb{F}_q$  is said to be an  $\mathbb{F}_q$ -maximal curve if it has  $q + 1 + 2g\sqrt{q}$  points defined over  $\mathbb{F}_q$ . Clearly, this can only be the case if either the cardinality  $q$  of the finite field is a square or  $g = 0$ .

Among maximal curves, the most studied are the so called *Deligne–Lusztig curves* [3], that is, the Hermitian curve, the Ree curve in characteristic 3, and the Suzuki curve

$$\mathcal{S}_q : \quad Y^q + Y = X^{q_0}(X^q + X)$$

with  $q = 2q_0^2$ ,  $q_0 = 2^h$ ,  $h \geq 0$ .<sup>1</sup>

Other examples of maximal curves are the GK curve [7], the GGS curve [5], the Klein quartic when  $\sqrt{q} \equiv 6 \pmod{7}$  [12], the BM curve [1], together with their quotient curves. Maximal curves often have large automorphism groups which in many cases can be inherited by the AG code itself: this can bring good performances in encoding [11] and decoding [8].

In this paper we focus on Weierstrass semigroups at points  $P$  of the Suzuki curve  $\mathcal{S}_q$ . In particular, we consider the case  $P \in \mathcal{S}_q(\mathbb{F}_{q^4}) \setminus \mathcal{S}_q(\mathbb{F}_q)$ ; that is,  $P \in \mathcal{S}_q$  is  $\mathbb{F}_{q^4}$ -rational but not  $\mathbb{F}_q$ -rational. Since it is well known that the set of Weierstrass points of  $\mathcal{S}_q$  coincides with  $\mathcal{S}_q(\mathbb{F}_q)$  (see Lemma 2.3), and  $H(P)$  is known for  $P \in \mathcal{S}_q(\mathbb{F}_q)$  (see Lemma 2.2), our assumption  $P \in \mathcal{S}_q(\mathbb{F}_{q^4}) \setminus \mathcal{S}_q(\mathbb{F}_q)$  is not restrictive. Together with [13, Lemma 3.1], our result provides the Weierstrass semigroup at every point of the Suzuki curve. Our main result is the following.

**Theorem 1.1.** *Let  $P \in \mathcal{S}_q \setminus \mathcal{S}_q(\mathbb{F}_q)$ . The set*

$$(1.1) \quad \mathcal{G} := \left\{ \begin{aligned} &\nu_{h,k} := hq - kq_0 - \left\lfloor \frac{2h - k - 2}{2} \right\rfloor : h \in \{1, \dots, q_0\}, k \in \{0, \dots, 2h - 2\} \\ &\cup \{\mu_h := hq - (2(h - q_0) - 1)q_0 - (q_0 - 1) : h \in \{q_0 + 1, \dots, 2q_0\}\} \end{aligned} \right\}$$

*is a minimal set of generators for  $H(P)$ .*

It is worth pointing out that, unlike the case  $P \in \mathcal{S}_q(\mathbb{F}_q)$ , the structure of  $H(P)$  with  $P \in \mathcal{S}_q \setminus \mathcal{S}_q(\mathbb{F}_q)$  is quite complicated. For instance, the number of generator is far more large, as can be seen comparing Lemma 2.2 and Theorem 1.1.

In the last section we provide examples of dual one-point codes arising from the curve  $\mathcal{S}_q$  and a point  $P \in \mathcal{S}_q(\mathbb{F}_{q^4}) \setminus \mathcal{S}_q(\mathbb{F}_q)$ . We compare such codes with the ones obtained using a point  $P' \in \mathcal{S}_q(\mathbb{F}_q)$ : in some cases improvements on the parameters can be obtained.

## 2. PRELIMINARY RESULTS

Through this section and in the rest of the paper we will use the following notation. Let  $q_0 = 2^s$ , where  $s \geq 1$ , and  $q = 2q_0^2$ . Let us denote the finite field with  $q$  elements and its algebraic closure by  $\mathbb{F}_q$  and  $\mathbb{K}$  respectively. The Suzuki curve  $\mathcal{S}_q$  is given by the affine model

$$(2.1) \quad \mathcal{S}_q : Y^q + Y = X^{q_0}(X^q + X).$$

The function field of  $\mathcal{S}_q$  over  $\mathbb{K}$  is denoted by  $\mathbb{K}(\mathcal{S}_q)$ . The curve  $\mathcal{S}_q$  is  $\mathbb{F}_{q^4}$ -maximal of genus  $g(\mathcal{S}_q) = q_0(q - 1)$ . It has a unique singular point, namely its unique point at infinity  $P_\infty$ , which is a  $q_0$ -fold point and the centre of just one branch of  $\mathcal{S}_q$ . The automorphism group

<sup>1</sup>OK per mettere l'equazione della sola Suzuki.

$\text{Aut}(\mathcal{S}_q)$  of  $\mathcal{S}_q$  over  $\mathbb{K}$  is isomorphic to the simple Suzuki group  ${}^2B_2(q) = \text{Sz}(q)$  and it acts on  $\mathcal{S}_q(\mathbb{F}_q)$  as  $\text{Sz}(q)$  on the Suzuki-Tits ovoid in  $\text{PG}(3, q)$ . The curve  $\mathcal{S}_q$  is  $\mathbb{F}_q$ -optimal and its number of  $\mathbb{F}_q$ -rational points is  $|\mathcal{S}_q(\mathbb{F}_q)| = q^2 + 1$ . For more details on the main properties of  $\mathcal{S}_q$  we refer the readers to [4, 6], and [9, Section 12.2].

**Lemma 2.1.** [9, Example 9.80] *Let  $P_0 \in \mathcal{S}_q(\mathbb{F}_q)$  and  $P \in \mathcal{S}_q$ . Then the following linear equivalence holds:*

$$qP + 2q_0\Phi(P) + \Phi^2(P) \sim (q + 2q_0 + 1)P_0,$$

where  $\Phi$  denotes the  $\mathbb{F}_q$ -Frobenius endomorphism. Then for every  $P \in \mathcal{S}_q$  there exists an element  $f_P \in \mathbb{K}(\mathcal{S}_q)$  with principal divisor

$$(f_P) = qP + 2q_0\Phi(P) + \Phi^2(P) - (q + 2q_0 + 1)P_\infty.$$

The Weierstrass semigroup at every  $\mathbb{F}_q$ -rational point of  $\mathcal{S}_q$  is known, as stated in the following lemma.

**Lemma 2.2.** [13, Lemma 3.1] *The Weierstrass semigroup at every  $\mathbb{F}_q$ -rational point of  $\mathcal{S}_q$  (and hence in particular at its infinite point  $P_\infty$ ) is*

$$H(P_\infty) = \langle q, q + q_0, q + 2q_0, q + 2q_0 + 1 \rangle.$$

The following lemma describes the structure of the set of Weierstrass points of  $\mathcal{S}_q$ . As a direct consequence of Lemmas 2.3 and 2.2, in order to compute the Weierstrass semigroup at every point of  $\mathcal{S}_q$ , it is not reductive to consider a point  $P \in \mathcal{S}_q(\mathbb{F}_{q^4}) \setminus \mathcal{S}_q(\mathbb{F}_q)$ .

**Lemma 2.3.** [4, Remark 5.11] and [4, Remark 5.13] *The curve  $\mathcal{S}_q$  is non-classical with respect to the canonical morphism. Furthermore, the set of Weierstrass points of  $\mathcal{S}_q$  coincides with the set of  $\mathbb{F}_q$ -rational points  $\mathcal{S}_q(\mathbb{F}_q)$  of  $\mathcal{S}_q$ .*

### 3. THE WEIERSTRASS SEMIGROUP AT EVERY POINT $P \in \mathcal{S}_q \setminus \mathcal{S}_q(\mathbb{F}_q)$

As recalled in Lemma 2.2, the structure of  $H(P)$  is known if  $P \in \mathcal{S}_q(\mathbb{F}_q)$ . Also, due to Lemma 2.3, it is not reductive to consider  $P \in \mathcal{S}_q(\mathbb{F}_{q^4}) \setminus \mathcal{S}_q(\mathbb{F}_q)$ . For this reason, for the rest of this section,  $P$  will always stand for an  $\mathbb{F}_{q^4}$ -rational point of  $\mathcal{S}_q$  which is not  $\mathbb{F}_q$ -rational. Clearly  $P$  is an affine point of  $\mathcal{S}_q$ , say  $P = (a, b)$  with  $a, b \in \mathbb{F}_{q^4} \setminus \mathbb{F}_q$ . In order to prove Theorem 1.1, we divide our investigation in three steps.

#### 3.1. The construction.

Let  $\Phi^i(P) = (a^{q^i}, b^{q^i})$  with  $i = 2, 3$  denote the  $i$ -th image of  $P$  with respect to the  $\mathbb{F}_q$ -Frobenius endomorphism  $\Phi : (X, Y) \mapsto (X^q, Y^q)$ . From Lemma 2.1 there exist

four functions  $f_P, f_{\Phi(P)}, f_{\Phi^2(P)}, f_{\Phi^3(P)}$  in  $\mathbb{K}(\mathcal{S}_q)$  such that

$$(3.1) \quad (f_P) = qP + 2q_0\Phi(P) + \Phi^2(P) - (q + 2q_0 + 1)P_\infty,$$

$$(3.2) \quad (f_{\Phi(P)}) = q\Phi(P) + 2q_0\Phi^2(P) + \Phi^3(P) - (q + 2q_0 + 1)P_\infty,$$

$$(3.3) \quad (f_{\Phi^2(P)}) = q\Phi^2(P) + 2q_0\Phi^3(P) + P - (q + 2q_0 + 1)P_\infty,$$

$$(3.4) \quad (f_{\Phi^3(P)}) = q\Phi^3(P) + 2q_0P + \Phi(P) - (q + 2q_0 + 1)P_\infty.$$

Note that since  $P \neq P_\infty$ ,  $P$  is a simple point of  $\mathcal{S}_q$  and the tangent line to  $\mathcal{S}_q$  at  $P$  is

$$\mathcal{T}_P : Y - b - a^{q_0}(X - a) = 0;$$

denote by  $t_P = y - b - a^{q_0}(x - a) \in \mathbb{K}(\mathcal{S}_q)$  the corresponding rational function.

From Bezout's Theorem, the intersection of  $\mathcal{S}_q$  and  $\mathcal{T}_P$  is a set of  $q + q_0$  points (counted with multiplicity) containing both  $P$  and  $\Phi(P)$ . The intersection multiplicity of  $\mathcal{T}_P$  and  $\mathcal{S}_q$  at  $P$  is  $q_0$  while it is equal to 1 at every other point of intersection. In fact, since  $\mathcal{T}_P$  does not contain  $P_\infty$ , all the other points  $Q$  of intersection are simple points of  $\mathcal{S}_q$  and  $\mathcal{T}_P$  is not the tangent line of  $\mathcal{S}_q$  at  $Q$ . Hence,

$$(3.5) \quad (t_P) = q_0P + \Phi(P) + E - (q + q_0)P_\infty,$$

where  $E$  is the effective divisor of degree  $q - 1$  whose support consists on the remaining  $q - 1$  intersection points of  $\mathcal{S}_q$  and  $t_P$ .

For non-negative integers  $h, i, j, k$ , and  $\ell$  we consider the following function

$$(3.6) \quad \theta_{h,i,j,k,\ell} := \frac{f_{\Phi(P)}^i f_{\Phi^2(P)}^j f_{\Phi^3(P)}^k t_P^\ell}{f_P^h} \in \mathbb{K}(\mathcal{S}_q).$$

In what follows, we select suitable functions of type (3.6) to obtain elements in  $H(P)$ .

**Lemma 3.1.** *Let*

$$(3.7) \quad m_{j,k,\ell} = \left\lceil \frac{q_0}{q_0 - 1} \left( j + k + \ell - \frac{k + \ell}{q} \right) \right\rceil,$$

where

$$(3.8) \quad \ell \in \{0, 1\}, \quad k \in \{0, \dots, q_0 - 1\}, \quad j \in \{0, \dots, q_0 - 1\}, \quad h \in \{\max\{1, m_{j,k,\ell}\}, \dots, 2q_0\}.$$

Let

$$(3.9) \quad \mathcal{F}_1 := \{ n_{h,j,k,\ell} := hq - (\ell + 2k)q_0 - j : h, j, k, \ell \text{ satisfy (3.8)} \}.$$

Then  $\mathcal{F}_1 \subseteq H(P)$ .

*Proof.* Let  $i$  be a non-negative integer and  $\theta_{h,i,j,k,\ell}$  be as in (3.6), with  $\ell, k, j, h, m_{j,k,\ell}$  as in (3.7) and (3.8). Then, from (3.1), (3.2), (3.3), (3.4), and (3.5) one gets

$$\begin{aligned}
(\theta_{h,i,j,k,\ell}) &= \left( \frac{f_{\Phi(P)}^i f_{\Phi^2(P)}^j f_{\Phi^3(P)}^k t_P^\ell}{f_P^h} \right) \\
&= i(q\Phi(P) + 2q_0\Phi^2(P) + \Phi^3(P) - (q + 2q_0 + 1)P_\infty) \\
&\quad + j(q\Phi^2(P) + 2q_0\Phi^3(P) + P - (q + 2q_0 + 1)P_\infty) \\
&\quad + k(q\Phi^3(P) + 2q_0P + \Phi(P) - (q + 2q_0 + 1)P_\infty) \\
&\quad + \ell(q_0P + \Phi(P) + E - (q + q_0)P_\infty) \\
&\quad - h(qP + 2q_0\Phi(P) + \Phi^2(P) - (q + 2q_0 + 1)P_\infty) \\
&= \ell E + (iq + \ell + k - 2hq_0)\Phi(P) + (2iq_0 + jq - h)\Phi^2(P) \\
&\quad + (i + 2jq_0 + kq)\Phi^3(P) - (hq - (\ell + 2k)q_0 - j)P \\
&\quad + [(h - \ell - k - i - j)(q + q_0) + (h - j - k - i)(q_0 + 1)]P_\infty.
\end{aligned}$$

Hence, to prove that  $n_{h,j,k,\ell} \in H(P)$  for each  $(h, j, k, \ell)$  satisfying (3.8), it is sufficient to show that  $P$  is the unique pole of  $\theta_{h,i,j,k,\ell}$  for some non-negative integer  $i$ . Since the divisor  $\ell E \geq 0$  and  $i + 2jq_0 + kq \geq 0$ , this is equivalent to show that the following system of inequalities

$$(3.10) \quad \begin{cases} iq + \ell + k - 2hq_0 \geq 0 \\ 2iq_0 + jq - h \geq 0 \\ (h - \ell - k - i - j)(q + q_0) + (h - j - k - i)(q_0 + 1) \geq 0 \\ hq - (\ell + 2k)q_0 - j > 0 \end{cases}$$

is satisfied for some non-negative integer  $i$ . The first inequality is equivalent to  $i \geq h/q_0 - (\ell + k)/q$ , while the second is equivalent to  $i \geq h/2q_0 - j/q_0$ . Since  $0 \leq k + \ell \leq q_0$ ,

$$\frac{h}{q_0} - \frac{\ell + k}{q} \geq \frac{h}{q_0} - \frac{q_0}{q} = \frac{h}{2q_0} + \frac{h-1}{2q_0} \geq \frac{h}{2q_0} + \frac{-2j}{2q_0} = \frac{h}{2q_0} - \frac{j}{q_0}.$$

This means that the first inequality in (3.10) implies the second one.

The third inequality is equivalent to  $i(q + 2q_0 + 1) \leq (h - k - j)(q + 2q_0 + 1) - \ell(q + q_0)$  and, since  $i$  must be an integer,

$$i \leq h - k - j - \left\lceil \frac{\ell(q + q_0)}{q + 2q_0 + 1} \right\rceil = h - k - j - \ell.$$

Finally, the fourth inequality in (3.10) is equivalent to

$$h > \frac{j}{q} + \frac{\ell + 2k}{2q_0}.$$

First, we show that the above condition on  $h$  holds for any  $(h, j, k, \ell)$  satisfying (3.8).

Note that  $\max\{1, m_{j,k,\ell}\} = 1$  if and only if  $j = k = \ell = 0$ : if  $j = k = \ell = 0$  then

$$m_{j,k,\ell} = \left\lceil \frac{q_0}{q_0 - 1} \left( j + k + \ell - \frac{k + \ell}{q} \right) \right\rceil = 0;$$

if  $j + k + \ell \geq 1$  then

$$m_{j,k,\ell} = \left\lceil \frac{q_0}{q_0 - 1} \left( j + k + \ell - \frac{k + \ell}{q} \right) \right\rceil \geq \left\lceil \frac{q_0}{q_0 - 1} \right\rceil \geq 2.$$

Consider now the case  $j + k + \ell \geq 1$ . We have that

$$\begin{aligned} & qq_0(j + k + \ell) - (k + \ell)q_0 - (q_0 - 1)(j + q_0\ell + 2q_0k) \\ &= k(qq_0 - q_0 - q + 2q_0) + j(qq_0 - q_0 + 1) + \ell(qq_0 - q_0^2) \\ &\geq k + j + \ell \geq 1, \end{aligned}$$

and hence

$$\frac{q_0}{q_0 - 1} \left( j + k + \ell - \frac{k + \ell}{q} \right) \geq \frac{j}{q} + \frac{\ell + 2k}{2q_0},$$

which yields

$$m_{j,k,\ell} \geq \frac{j}{q} + \frac{\ell + 2k}{2q_0}$$

and shows that the fourth inequality in (3.10) is satisfied for any  $(h, j, k, \ell)$  satisfying (3.8).

Summing up, we only have to show that there exists at least one non-negative integer  $i$  with

$$\frac{h}{q_0} - \frac{\ell + k}{q} \leq i \leq h - k - j - \ell.$$

Since  $k + j + \ell \leq \max\{1, m_{j,k,\ell}\} \leq h$ , the integer  $h - k - j - \ell$  is non-negative and therefore it is enough to show that  $\frac{h}{q_0} - \frac{\ell + k}{q} \leq h - k - j - \ell$ , which is equivalent to

$$h \geq \frac{q_0}{q_0 - 1} \left( k + \ell + j - \frac{k + \ell}{q} \right).$$

As  $h \geq m_{j,k,\ell}$ , the claim follows.  $\square$

We now construct a second family of nongaps at  $P$  which we will prove to be disjoint from  $\mathcal{F}_1$ .

**Lemma 3.2.** *Let*

$$(3.11) \quad \mathcal{F}_2 := \{ n_{\tilde{h}} = \tilde{h}q - (2\tilde{h} - 2q_0 - 1)q_0 - (q_0 - 1) : \tilde{h} \in \{q_0 + 1, \dots, 2q_0\} \}.$$

*Then  $\mathcal{F}_2 \subseteq H(P)$ .*

*Proof.* Let  $\tilde{i} \geq 0$ ,  $\tilde{h} \in \{q_0 + 1, \dots, 2q_0\}$ , and

$$(3.12) \quad \psi_{\tilde{h}, \tilde{i}} := \frac{f_{\Phi(P)}^{\tilde{i}} f_{\Phi^3(P)}^{\tilde{h} - q_0}}{f_P^{\tilde{h}} f_{\Phi^2(P)}} \in \mathbb{K}(\mathcal{S}_q).$$

Then from (3.1), (3.2), (3.3), and (3.4) follows

$$\begin{aligned}
(\psi_{\tilde{h}, \tilde{i}}) &= \left( \frac{f_{\Phi(P)}^{\tilde{i}} f_{\Phi^3(P)}^{\tilde{h}-q_0}}{f_P^{\tilde{h}} f_{\Phi^2(P)}} \right) \\
&= \tilde{i} (q\Phi(P) + 2q_0\Phi^2(P) + \Phi^3(P) - (q + 2q_0 + 1)P_\infty) \\
&\quad + (\tilde{h} - q_0) (q\Phi^3(P) + 2q_0P + \Phi(P) - (q + 2q_0 + 1)P_\infty) \\
&\quad - \tilde{h} (qP + 2q_0\Phi(P) + \Phi^2(P) - (q + 2q_0 + 1)P_\infty) \\
&\quad - (q\Phi^2(P) + 2q_0\Phi^3(P) + P - (q + 2q_0 + 1)P_\infty) \\
&= (\tilde{i}q + \tilde{h} - q_0 - 2\tilde{h}q_0)\Phi(P) + (2\tilde{i}q_0 - \tilde{h} - q)\Phi^2(P) \\
&\quad + (\tilde{i} + \tilde{h}q - qq_0 - 2q_0)\Phi^3(P) + (-\tilde{i} + q_0 + 1)(q + 2q_0 + 1)P_\infty \\
&\quad - (\tilde{h}q - (2\tilde{h} - 2q_0 - 1)q_0 - (q_0 - 1))P.
\end{aligned}$$

Hence, as in the proof of Lemma 3.1, the claim follows proving that  $\psi_{\tilde{h}, \tilde{i}}$  has a unique pole at  $P$ . This is equivalent to prove that the following system of inequalities

$$(3.13) \quad \begin{cases} \tilde{i}q + \tilde{h} - q_0 - 2\tilde{h}q_0 \geq 0 \\ 2\tilde{i}q_0 - \tilde{h} - q \geq 0 \\ \tilde{i} + \tilde{h}q - qq_0 - 2q_0 \geq 0 \\ -\tilde{i} + q_0 + 1 \geq 0 \\ \tilde{h}q - (2\tilde{h} - 2q_0 - 1)q_0 - (q_0 - 1) > 0 \end{cases}$$

is satisfied for some non-negative integer  $\tilde{i}$ .

The first and the second inequalities in (3.13) are equivalent to  $\tilde{i} \geq \tilde{h}/q_0 + 1/2q_0 - \tilde{h}/q$  and  $\tilde{i} \geq \tilde{h}/2q_0 + 1/q_0$ , respectively. Since by hypothesis  $\tilde{h} \geq q_0 + 1 > 1$ ,

$$\frac{\tilde{h}}{2q_0} + \frac{1}{q_0} = \frac{\tilde{h}}{q_0} + \frac{2 - \tilde{h}}{2q_0} \leq \frac{\tilde{h}}{q_0} + \frac{1 - \tilde{h}/q_0}{2q_0},$$

and the first inequality implies the second one.

The third inequality is equivalent to  $\tilde{i} \geq 2q_0 + qq_0 - \tilde{h}q$ , which is always satisfied since  $\tilde{h} \geq q_0 + 1$ . The fourth inequality is equivalent to  $\tilde{i} \leq q_0 + 1$ . Finally, the last inequality in (3.13) is equivalent to  $\tilde{h}(q - 2q_0) > -(2q_0 + 1)q_0 + (q_0 - 1) = -q - 1$  which is satisfied as  $\tilde{h}$  is strictly positive.

The integer  $\tilde{i} := q_0 + 1$  is positive and clearly satisfies  $\tilde{i} \geq 3 > \tilde{h}/q_0 + 1/2q_0$ . The claim follows.  $\square$

3.2.  $\langle \mathcal{F}_1 \cup \mathcal{F}_2 \rangle$  coincides with  $H(P)$ .

*Remark 3.3.* Let  $S$  be a numerical semigroup and  $s$  be the multiplicity of  $S$ , i.e. its least non-zero element. If an element  $r \in S$  is such that  $[r, \dots, r + s - 1] \subset S$ , then the conductor of  $S$  is at most  $r$ , that is  $m \in S$  for any  $m \geq r$ ; in fact  $m = \lfloor (m - r)/s \rfloor \cdot s + u$  for some  $u \in [r, \dots, r + s - 1]$ .

By Lemma 3.1 and Lemma 3.2 we already know that  $\langle \mathcal{F}_1 \cup \mathcal{F}_2 \rangle \subseteq H(P)$ . In what follows we prove that  $\mathcal{F}_1 \cup \mathcal{F}_2$  contains at least  $g(\mathcal{S}_q)$  elements which are less than or equal to  $2g(\mathcal{S}_q) - 1$ , and that  $[2g(\mathcal{S}_q) - q + 2, \dots, 2g(\mathcal{S}_q) + 1] \subset \mathcal{F}_1 \cup \mathcal{F}_2$ . Therefore  $H(P) = \langle \mathcal{F}_1 \cup \mathcal{F}_2 \rangle$  by Remark 3.3.

**Lemma 3.4.** *Let  $\mathcal{F}_1$  be as in (3.9). Then the elements of  $\mathcal{F}_1$  are pairwise distinct and*

$$|\mathcal{F}_1 \cap [1, \dots, 2g(\mathcal{S}_q) - 1]| = 2q_0^3 - 2q_0 - 1.$$

*Proof.* Suppose  $n_{h,j,k,\ell} = n_{h_1,j_1,k_1,\ell_1} \in \mathcal{F}_1$ , for some  $(h, j, k, \ell), (h_1, j_1, k_1, \ell_1)$  satisfying (3.8). Then  $hq - (\ell + 2k)q_0 - j = h_1q - (\ell_1 + 2k_1)q_0 - j_1$  and  $j \equiv j_1 \pmod{q_0}$ . Since  $0 \leq j, j_1 \leq q_0 - 1$  we have that  $j = j_1$  and hence  $2hq_0 - (\ell + 2k)q_0 = 2h_1q_0 - (\ell_1 + 2k_1)q_0$ . Thus  $\ell \equiv \ell_1 \pmod{2}$  and therefore  $\ell = \ell_1$ . We are left with  $hq_0 - k = h_1q_0 - k_1$ . Considering the congruence modulo  $q_0$  we get that  $k = k_1$  and hence  $h = h_1$ . This shows that there are no repetitions in  $\mathcal{F}_1$ .

Let  $n_{h,j,k,\ell} \in \mathcal{F}_1$  such that  $n_{h,j,k,\ell} \geq 2g(\mathcal{S}_q) = 2q_0q - 2q_0$ . This yields  $h = 2q_0$  and  $2k + \ell \leq 2$ . More precisely, one of the following three possibilities occurs

- $\ell = 0, k = 1$  and  $j = 0$ , or
- $\ell = 1, k = 0$  and  $j = 0, \dots, q_0 - 1$ , or
- $\ell = 0, k = 0$  and  $j = 0, \dots, q_0 - 1$ .

Since  $0 \notin \mathcal{F}_1$ ,

$$(3.14) \quad |\mathcal{F}_1 \cap [1, \dots, 2g(\mathcal{S}_q) - 1]| = |\mathcal{F}_1| - 1 - q_0 - q_0 = |\mathcal{F}_1| - 2q_0 - 1.$$

To compute  $|\mathcal{F}_1|$  it is convenient to divide the elements of  $\mathcal{F}_1$  according to the corresponding value of  $M_{j,k,\ell} := \max\{1, m_{j,k,\ell}\}$ . To this end, three cases are considered.

- (A) Assume that  $j = k = \ell = 0$ . Then, clearly,  $m_{j,k,\ell} = 0$ . In this case  $M_{j,k,\ell} = 1$ .  
 (B) Let  $j + k + \ell \geq 1$  and  $j \leq q_0 - 1 - (k + \ell)$ . Then

$$\begin{aligned} m_{j,k,\ell} &= \left\lceil \frac{q_0}{q_0 - 1} \left( j + k + \ell - \frac{k + \ell}{q} \right) \right\rceil = j + k + \ell + \left\lceil \frac{j + k + \ell}{q_0 - 1} - \frac{k + \ell}{2q_0(q_0 - 1)} \right\rceil \\ &\leq j + k + \ell + \left\lceil \frac{(q_0 - 1 - k - \ell) + k + \ell}{q_0 - 1} - \frac{k + \ell}{2q_0(q_0 - 1)} \right\rceil = j + k + \ell + 1. \end{aligned}$$

Since  $j + k + \ell \geq 1$  and then  $m_{j,k,\ell} \geq j + k + \ell + 1$ , we get  $M_{j,k,\ell} = m_{j,k,\ell} = j + k + \ell + 1$ .



(C) Let  $j \geq q_0 - (k + \ell)$ . Then  $j + k + \ell \geq 1$ . Also,

$$\begin{aligned} m_{j,k,\ell} &= j + k + \ell + \left\lceil \frac{j + k + \ell}{q_0 - 1} - \frac{k + \ell}{2q_0(q_0 - 1)} \right\rceil \\ &\geq j + k + \ell + \left\lceil \frac{q_0}{q_0 - 1} - \frac{1}{2(q_0 - 1)} \right\rceil > j + k + \ell + 1; \\ m_{j,k,\ell} &= j + k + \ell + \left\lceil \frac{j + k + \ell}{q_0 - 1} - \frac{k + \ell}{2q_0(q_0 - 1)} \right\rceil \\ &\leq j + k + \ell + \left\lceil \frac{2q_0 - 1}{q_0 - 1} \right\rceil = j + k + \ell + 3. \end{aligned}$$

Hence either  $m_{j,k,\ell} = j + k + \ell + 2$  or  $m_{j,k,\ell} = j + k + \ell + 3$ . Actually, the case  $m_{j,k,\ell} = j + k + \ell + 3$  cannot occur. Indeed,  $m_{j,k,\ell} = j + k + \ell + 3$  if and only if

$$\left\lceil \frac{j + k + \ell}{q_0 - 1} - \frac{k + \ell}{2q_0(q_0 - 1)} \right\rceil = 3 \iff j + k + \ell > \frac{k + \ell}{2q_0} + 2q_0 - 2 > 2q_0 - 2.$$

This yields  $j + k + \ell = 2q_0 - 1$  and therefore

$$h \geq M_{j,k,\ell} = m_{j,k,\ell} = j + k + \ell + 3 = 2q_0 + 2 > 2q_0,$$

a contradiction to the hypothesis  $h \leq 2q_0$ . So  $M_{j,k,\ell} = m_{j,k,\ell} = j + k + \ell + 2$ .

We proceed computing the number of elements in  $\mathcal{F}_1$  of type (A), (B) and (C).

- Since  $h = 1, \dots, 2q_0$ , there are exactly  $2q_0$  elements of type (A).
- Deleting the  $2q_0$  elements of type (A), for which clearly  $j \leq q_0 - 1 - (k + \ell)$ , and noting that  $k \leq q_0 - 1 - \ell$  as  $j \geq 0$ , we get that the number of elements of type (B) is

$$\begin{aligned} &-2q_0 + \sum_{\ell=0}^1 \sum_{k=0}^{q_0-1-\ell} \sum_{j=0}^{q_0-1-k-\ell} (2q_0 - j - k - \ell) = \\ &-2q_0 + \sum_{\ell=0}^1 \sum_{k=0}^{q_0-1-\ell} \frac{(q_0 - k - \ell)(3q_0 - k - \ell + 1)}{2}. \end{aligned}$$

From the previous equality, defining  $x = q_0 - k - \ell$ , we get that the number of elements of type (B) equals

$$\begin{aligned}
& -2q_0 + \sum_{\ell=0}^1 \sum_{x=1}^{q_0-\ell} \left( \frac{x^2}{2} + x \left( q_0 + \frac{1}{2} \right) \right) = \\
& -2q_0 + \sum_{\ell=0}^1 \left( \left( q_0 + \frac{1}{2} \right) \frac{(q_0 - \ell)(q_0 - \ell + 1)}{2} + \frac{1}{2} \sum_{x=1}^{q_0-\ell} x^2 \right) = \\
& -2q_0 + \sum_{\ell=0}^1 \left[ \left( q_0 + \frac{1}{2} \right) \frac{(q_0 - \ell)(q_0 - \ell + 1)}{2} + \frac{(q_0 - \ell)(q_0 - \ell + 1)(2q_0 - 2\ell + 1)}{12} \right] = \\
& \frac{4}{3}q_0^3 + \frac{1}{2}q_0^2 + \frac{1}{6}q_0 - 2q_0.
\end{aligned}$$

- We divide the elements of type (C) in two classes, namely the ones having  $\ell = 0$  and the ones for which  $\ell = 1$ .

If  $\ell = 0$  then  $1 \leq k \leq q_0 - 1$  as  $q_0 - k \leq j \leq q_0 - 1$ , while  $j + k + \ell + 2 \leq h \leq 2q_0$ . Hence the number  $c_0$  of elements of type (C) with  $\ell = 0$  is

$$\begin{aligned}
c_0 &= \sum_{k=1}^{q_0-1} \sum_{j=q_0-k}^{q_0-1} (2q_0 - j - k - 1) = \sum_{k=1}^{q_0-1} \frac{(2q_0 - k - 1)k}{2} = \\
& \sum_{k=1}^{q_0-1} \left( k \left( q_0 - \frac{1}{2} \right) - \frac{k^2}{2} \right) = \left( q_0 - \frac{1}{2} \right) \frac{(q_0 - 1)q_0}{2} - \frac{q_0(q_0 - 1)(2q_0 - 1)}{12} = \\
& \frac{q_0(q_0 - 1)}{2} \left( q_0 - \frac{1}{2} - \frac{2q_0 - 1}{6} \right).
\end{aligned}$$

If  $\ell = 1$  then  $0 \leq k \leq q_0 - 1$ ,  $q_0 - 1 - k \leq j \leq q_0 - 1$  and  $j + k + 3 \leq h \leq 2q_0$ . Thus the number  $c_1$  of elements of type (C) with  $\ell = 1$  is

$$\begin{aligned}
c_1 &= \sum_{k=0}^{q_0-1} \sum_{j=q_0-k-1}^{q_0-1} (2q_0 - j - k - 2) = \sum_{k=0}^{q_0-1} \frac{(2q_0 - k - 2)(k + 1)}{2} = \\
& \sum_{k=0}^{q_0-1} \left( k \left( q_0 - \frac{3}{2} \right) - \frac{k^2}{2} + q_0 - 1 \right) = \left( q_0 - \frac{3}{2} \right) \frac{(q_0 - 1)q_0}{2} - \frac{q_0(q_0 - 1)(2q_0 - 1)}{12} + q_0^2 - q_0 = \\
& \frac{q_0(q_0 - 1)}{2} \left( q_0 + \frac{1}{2} - \frac{2q_0 - 1}{6} \right).
\end{aligned}$$

By direct computation, we get

$$c_0 + c_1 = \frac{q_0(q_0 - 1)}{2} \left( 2q_0 - \frac{2q_0 - 1}{3} \right) = \frac{2}{3}q_0^3 - \frac{q_0^2}{2} - \frac{q_0}{6}.$$

Summing up the contributions we get that

$$|\mathcal{F}_1| = 2q_0 + \frac{4}{3}q_0^3 + \frac{1}{2}q_0^2 + \frac{1}{6}q_0 - 2q_0 + \frac{2}{3}q_0^3 - \frac{1}{2}q_0^2 - \frac{1}{6}q_0 = 2q_0^3,$$

and hence, by (3.14),

$$|\mathcal{F}_1 \cap [1, \dots, 2g(\mathcal{S}_q) - 1]| = |\mathcal{F}_1| - 2q_0 - 1 = 2q_0^3 - 2q_0 - 1.$$

□

The following lemma shows that  $\mathcal{F}_1$  and  $\mathcal{F}_2$  are disjoint.

**Lemma 3.5.** *Let  $\mathcal{F}_1$  and  $\mathcal{F}_2$  be the sets of non-gaps defined in Lemma 3.1 and Lemma 3.2 respectively. Then  $\mathcal{F}_1 \cap \mathcal{F}_2 = \emptyset$ .*

*Proof.* Assume by contradiction that  $n_{h,j,k,\ell} = n_{\tilde{h}}$  for some  $n_{h,j,k,\ell} \in \mathcal{F}_1$  and  $n_{\tilde{h}} \in \mathcal{F}_2$ . Then

$$hq - (\ell + 2k)q_0 - j = \tilde{h}q - (2\tilde{h} - 2q_0 - 1)q_0 - (q_0 - 1).$$

Since  $j$  must be congruent to  $-1$  modulo  $q_0$  we get that  $j = q_0 - 1$  and

$$2hq_0 - (\ell + 2k) = 2\tilde{h}q_0 - (2\tilde{h} - 2q_0 - 1).$$

Considering the congruence modulo 2 we get that  $\ell = 1$  and

$$hq_0 - k = \tilde{h}q_0 - \tilde{h} + q_0 + 1.$$

Taking the congruence modulo  $q_0$  we get that  $k = \tilde{h} - (q_0 + 1)$  and  $h = \tilde{h}$ ; hence,  $(h, j, k, \ell) = (\tilde{h}, q_0 - 1, \tilde{h} - (q_0 + 1), 1)$ . Conditions (3.8) imply

$$\begin{aligned} h &\geq m_{j,k,\ell} \geq \frac{q_0}{q_0 - 1} \left( j + k + \ell - \frac{k + \ell}{q} \right) \\ &= \frac{q_0}{q_0 - 1} \left( q_0 - 1 + \tilde{h} - (q_0 + 1) + 1 - \frac{\tilde{h} - (q_0 + 1) + 1}{q} \right) \\ &= \frac{q_0}{q_0 - 1} \left( \tilde{h} - 1 - \frac{\tilde{h} - q_0}{q} \right) = \tilde{h} \frac{2q_0^2 - 1}{2q_0(q_0 - 1)} - \frac{q_0 - 1/2}{q_0 - 1} > \tilde{h}, \end{aligned}$$

a contradiction to  $(h, j, k, \ell) = (\tilde{h}, q_0 - 1, \tilde{h} - (q_0 + 1), 1)$ . □

**Lemma 3.6.** *Let  $\mathcal{F}_2$  be the set of non-gaps constructed in Lemma 3.2. Then the elements of  $\mathcal{F}_2$  are pairwise distinct and*

$$|\mathcal{F}_2 \cap [1, \dots, 2g(\mathcal{S}_q) - 1]| = q_0.$$

*Proof.* It is easily seen that  $n_{\tilde{h}} = n_{\tilde{h}_1}$  implies  $\tilde{h} = \tilde{h}_1$  and therefore the elements of  $\mathcal{F}_2$  are pairwise distinct. Also,  $\mathcal{F}_2$  contains exactly  $q_0$  elements since  $\tilde{h} = q_0 + 1, \dots, 2q_0$ . Finally, for each  $\tilde{h} = q_0 + 1, \dots, 2q_0$ ,

$$\begin{aligned} n_{\tilde{h}} &= \tilde{h}q - (2\tilde{h} - 2q_0 - 1)q_0 - (q_0 - 1) \leq 2q_0q - (4q_0 - 2q_0 - 1)q_0 - (q_0 - 1) \\ &= 2q_0q - (2q_0 - 1)q_0 - q_0 + 1 = 2q_0q - (q - 1) < 2g(\mathcal{S}_q) - 1. \end{aligned}$$

The claim follows.  $\square$

**Lemma 3.7.** *The interval  $[2g(\mathcal{S}_q) - q + 2, \dots, 2g(\mathcal{S}_q) + 1]$  is contained in  $\langle \mathcal{F}_1 \cup \mathcal{F}_2 \rangle$ .*

*Proof.* Let  $I = [2g(\mathcal{S}_q) - q + 2, \dots, 2g(\mathcal{S}_q) + 1] = [(2q_0 - 1)q - q_0 - (q_0 - 2), \dots, 2q_0q - 2q_0 + 1]$  and  $t \in I$ .

- If  $t = 2q_0q - q + 1$ , then  $t = n_{2q_0} \in \mathcal{F}_2$ .
- If  $t \in I$  and  $t \leq (2q_0 - 1)q$ , then  $t = n_{2q_0-1,j,0,\ell}$ . Since  $m_{j,0,\ell} \leq j + 0 + \ell + 2 < q_0 + 1 \leq 2q_0 - 1 = h$ , (3.8) is satisfied and  $t \in \mathcal{F}_1$ .
- If  $t \in I \setminus \{2q_0q - q + 1\}$  and  $t > (2q_0 - 1)q$ , then  $t = n_{2q_0,l,k,\ell}$  where  $m_{j,k,\ell} \leq j + k + \ell + 2 \leq 2q_0 = h$  by direct checking. Hence (3.8) is satisfied and  $t \in \mathcal{F}_1$ .

$\square$

*Remark 3.8.* From Lemma 3.7 it follows in particular that  $2g(\mathcal{S}_q) - 1$  is a non-gap at  $P$ , that is the semigroup  $H(P)$  is non-symmetric for any  $P \in \mathcal{S}_q \setminus \mathcal{S}_q(\mathbb{F}_q)$ ; unlikely the Weierstrass semigroup at any  $\mathbb{F}_q$ -rational point of  $\mathcal{S}_q$ , which is symmetric ([4, Lemma 5.7]).

**Theorem 3.9.** *Let  $\mathcal{F}_1$  and  $\mathcal{F}_2$  as in Lemmas 3.1 and 3.2. If  $P \in \mathcal{S}_q$  is not  $\mathbb{F}_q$ -rational then*

$$H(P) = \langle \mathcal{F}_1 \cup \mathcal{F}_2 \rangle.$$

*Proof.* The claim follows immediately from Lemma 3.7 and

$$|(\mathcal{F}_1 \cup \mathcal{F}_2 \cup \{0\}) \cap [0, \dots, 2g(\mathcal{S}_q) - 1]| = 2q_0^3 - q_0 = q_0(q - 1) = g(\mathcal{S}_q).$$

$\square$

### 3.3. A minimal set of generators for $H(P)$ .

In this section we provide a set of minimal generators for the Weierstrass semigroup  $H(P)$ ,  $P \in \mathcal{S}_q(\mathbb{F}_{q^4}) \setminus \mathcal{S}_q(\mathbb{F}_q)$ . The proof of Theorem 1.1 is divided into two steps.

Let us define

$$(3.15) \quad \mathcal{G}_1 := \left\{ \nu_{h,k} = hq - kq_0 - \left\lfloor \frac{2h - k - 2}{2} \right\rfloor : h \in \{1, \dots, q_0\}, k \in \{0, \dots, 2h - 2\} \right\},$$

so that  $\mathcal{G} = \mathcal{G}_1 \cup \mathcal{F}_2$ .

**Step 1:  $\mathcal{G}$  generates  $H(P)$ .**

Consider an element  $n_{h,j,k,\ell} \in \mathcal{F}_1$ . Let  $\delta(n_{h,j,k,\ell}) = \left\lfloor \frac{2h - (\ell + 2k) - 2}{2} \right\rfloor - j = h - 1 - k - j - \left\lceil \frac{\ell}{2} \right\rceil$ . We will relate  $\delta(n_{h,j,k,\ell})$  with the number of elements of  $\mathcal{G}_1$  we need to generate  $n_{h,j,k,\ell}$ .

**Proposition 3.10.** *If  $\delta(n_{h,j,k,\ell}) = 0$  then  $n_{h,j,k,\ell}$  is a generator.*

*Proof.* If  $\delta(n_{h,j,k,\ell}) = 0$  then  $h \leq q_0$ . In fact,

$$\frac{q_0}{q_0 - 1} \left( j + k + \ell - \frac{k + \ell}{q} \right) \leq h.$$

Using  $j = \left\lfloor \frac{2h - (\ell + 2k) - 2}{2} \right\rfloor$  we get

$$\frac{q_0}{q_0 - 1} \left( \left\lfloor \frac{2h - (\ell + 2k) - 2}{2} \right\rfloor + k + \ell - \frac{k + \ell}{q} \right) \leq h,$$

and hence

$$\frac{q_0}{q_0 - 1} \left( h - 1 - \left\lceil \frac{\ell}{2} \right\rceil + \ell - \frac{k + \ell}{q} \right) \leq h.$$

Since  $\ell - \left\lceil \frac{\ell}{2} \right\rceil = 0$ , we get

$$h \leq q_0 + \left\lfloor \frac{k + \ell}{2q_0} \right\rfloor = q_0.$$

Now to show that if  $\delta(n_{h,j,k,\ell}) = 0$  then  $n_{h,j,k,\ell}$  is a generator it is sufficient to note that  $n_{h,j,k,\ell} = \nu_{h,2k+\ell}$ ,  $h \leq q_0$  and  $2h - (\ell + 2k) - 2 \geq 0$  implying  $2k + \ell \leq 2h - 2$ .  $\square$

So we can assume that  $\delta(n_{h,j,k,\ell}) \neq 0$ . We distinguish two cases.

**Proposition 3.11.** *Every element  $n_{h,j,k,0} \in \mathcal{F}_1$  belongs to  $\langle \mathcal{G}_1 \rangle$ .*

*Proof.* We prove that we need  $\delta(n_{h,j,k,0}) + 1$  elements from  $\mathcal{G}_1$  to generate  $n_{h,j,k,0}$ . First, we show that  $0 \leq \delta(n_{h,j,k,0}) \leq h - 1$ . Since  $h - 1 = \delta(n_{h,j,k,0}) + k + j$  the upper bound is clear. To see the lower bound, recall that

$$(3.16) \quad h \geq \left\lceil \frac{q_0}{q_0 - 1} \left( j + k + \ell - \frac{k + \ell}{q} \right) \right\rceil,$$

so

$$\delta(n_{h,j,k,0}) \geq \left\lfloor \frac{q_0}{q_0 - 1} \left( j + k - \frac{k}{q} \right) \right\rfloor - 1 - k - j.$$

Now, apart from the case  $j = k = 0$ , we have that  $\frac{q_0}{q_0 - 1} \left( j + k - \frac{k}{q} \right) > k + j$ , since  $j + k > \frac{k}{2q_0}$ . If  $j = k = 0$  then  $h \geq 1 = \max \left\{ 1, \left\lceil \frac{q_0}{q_0 - 1} \left( j + k + \ell - \frac{k + \ell}{q} \right) \right\rceil \right\}$  and hence  $\delta(n_{h,j,k,0}) \geq 0$ .

Consider a sequence of  $(h_i, k_i)_{i=1, \dots, \delta(n_{h,j,k,0})+1}$  such that

$$\sum_{i=1}^{\delta(n_{h,j,k,0})+1} h_i = h, \quad q_0 \geq h_i \geq 1, \quad \sum_{i=1}^{\delta(n_{h,j,k,0})+1} k_i = 2k, \quad k_i \text{ even.}$$

By (3.16), we can consider  $h_i \geq k_i/2 + 1$ , since  $h \geq k + 1$  and then each  $\nu_{h_i, k_i}$  is an element of  $\mathcal{G}_1$ . Now,

$$\begin{aligned} \sum_{i=1}^{\delta(n_{h,j,k,0})+1} \left( h_i q - k_i q_0 - \left\lfloor \frac{2h_i - k_i - 2}{2} \right\rfloor \right) &= \sum_{i=1}^{\delta(n_{h,j,k,0})+1} \left( h_i q - k_i q_0 - h_i + 1 + \frac{k_i}{2} \right) \\ &= hq - 2kq_0 - h + \delta(n_{h,j,k,0}) + 1 + k \\ &= hq - 2kq_0 - j = n_{h,j,k,0} \end{aligned}$$

and the claim follows.  $\square$

**Proposition 3.12.** *Every element  $n_{h,j,k,1} \in \mathcal{F}_1$  belongs to  $\langle \mathcal{G}_1 \rangle$ .*

*Proof.* We distinguish three subcases.

- (1) Suppose  $h \geq m_{j,k,1} + 1$ . We can also suppose  $h \geq 3$ , since the smallest integer with  $\ell = 1$  which is not in  $\mathcal{G}_1$  is  $3q - q_0$ . Consider the generator  $\nu_{2,1} = 2q - q_0$ . Now

$$n_{h,j,k,1} - \nu_{2,1} = n_{h-2,j,k,0}.$$

We only have to prove that such element is in the semigroup  $H(P)$  so that, by Proposition 3.11, it belongs to  $\langle \mathcal{G}_1 \rangle$ . To this end, observe that

$$h - 2 \geq m_{j,k,1} - 1 \geq m_{j,k,0}$$

because

$$\begin{aligned} m_{j,k,1} - 1 &= \left\lfloor \frac{q_0}{q_0 - 1} \left( j + k + 1 - \frac{k+1}{q} \right) \right\rfloor - 1 \\ &= \left\lfloor \frac{q_0}{q_0 - 1} \left( j + k - \frac{k}{q} \right) + \frac{q_0}{q_0 - 1} - \frac{1}{2q_0(q_0 - 1)} \right\rfloor - 1 \\ &= \left\lfloor \frac{q_0}{q_0 - 1} \left( j + k - \frac{k}{q} \right) + \frac{2q_0 - 1}{q - 2q_0} \right\rfloor \geq m_{j,k,0}. \end{aligned}$$

Therefore  $n_{h,j,k,1} \in \langle \mathcal{G}_1 \rangle$ .

- (2) Suppose  $h = m_{j,k,1}$  and  $h \leq q_0$ ; we show that  $n_{h,j,k,1}$  is a generator. From  $h \leq q_0$  follows  $j + k + 1 \leq q_0 - 1$ . Also,

$$\begin{aligned} \delta &= m_{j,k,1} - k - j - 2 \leq \frac{q_0}{q_0 - 1} \left( j + k + 1 - \frac{k+1}{q} \right) - (k + j + 1) \\ &= \frac{1}{q_0 - 1} (j + k + 1) - \frac{k+1}{2q_0(q_0 - 1)} \leq 1 - \frac{k+1}{2q_0(q_0 - 1)} < 1, \end{aligned}$$

and then  $\delta = 0$ , so that  $n_{h,j,k,1}$  is a generator by Proposition 3.10.

(3) Suppose  $h = m_{j,k,1} = q_0 + i$ ,  $1 \leq i \leq q_0$ .

First, note that  $j + k = q_0 + i - 3$ . In fact,

$$\begin{aligned} q_0 + i - 1 &\leq \frac{q_0}{q_0-1} \left( j + k + 1 - \frac{k+1}{q} \right) \leq q_0 + i, \\ q_0 + i - 1 + \frac{k+1}{2q_0(q_0-1)} &\leq \frac{q_0}{q_0-1} (j + k + 1) \leq q_0 + i + \frac{k+1}{2q_0(q_0-1)}, \\ \frac{q_0-1}{q_0} (q_0 + i - 1) + \frac{k+1}{q} &\leq j + k + 1 \leq \frac{q_0-1}{q_0} (q_0 + i) + \frac{k+1}{q}, \\ q_0 - 1 + \frac{q_0-1}{q_0} (i - 1) + \frac{k+1}{q} &\leq j + k + 1 \leq q_0 - 1 + \frac{q_0-1}{q_0} i + \frac{k+1}{q}. \end{aligned}$$

Now,  $\frac{q_0-1}{q_0} (i - 1) + \frac{k+1}{q} > i - 2$  since  $-\frac{i-1}{q_0} + \frac{k+1}{q} > -1$ , and  $\frac{q_0-1}{q_0} i + \frac{k+1}{q} < i$ , since  $-\frac{i}{q_0} + \frac{k+1}{q} < 0$ . So  $q_0 - 1 + i - 2 < j + k + 1 < q_0 - 1 + i$  and then  $j + k + 1 = q_0 - 2 + i$ , that is  $j + k = q_0 - 3 + i$ .

This is enough to show that actually  $i = 1$  cannot occur. In fact,

$$\begin{aligned} h &= q_0 + 1 > q_0 = \left\lceil \frac{q_0}{q_0-1} \left( q_0 - 1 - \frac{k+1}{q} \right) \right\rceil \\ &= \left\lceil \frac{q_0}{q_0-1} \left( j + k + 1 - \frac{k+1}{q} \right) \right\rceil = m_{j,k,1}, \end{aligned}$$

a contradiction to  $h = m_{j,k,1}$ .

Since  $i \geq 2$ ,  $j = q_0 - 3 + i - k \geq i - 2$ .

We prove now that  $n_{h,j,k,1} - \nu_{i,1}$  belongs to the semigroup. In fact,  $n_{h,j,k,1} - \nu_{i,1} = n_{q_0, j-i+2, k, 0}$  so that, in order to prove that  $n_{q_0, j-i+2, k, 0} \in \mathcal{F}_1$ , we only have to check that  $q_0 \geq m_{j-i+2, k, 0}$ . We have that

$$\begin{aligned} m_{j-i+2, k, 0} &= \left\lceil \frac{q_0}{q_0-1} \left( j - i + 2 + k - \frac{k}{q} \right) \right\rceil = \left\lceil \frac{q_0}{q_0-1} \left( q_0 - 1 - \frac{k}{q} \right) \right\rceil \\ &= \left\lceil q_0 - \frac{k}{2q_0(q_0-1)} \right\rceil = q_0. \end{aligned}$$

Therefore  $n_{h,j,k,1} \in \langle \mathcal{G}_1 \rangle$ .

□

Propositions 3.10, 3.11, and 3.12 show that  $\mathcal{G} = \mathcal{G}_1 \cup \mathcal{F}_2$  is a set of generator for  $H(P) = \langle \mathcal{F}_1 \cup \mathcal{F}_2 \rangle$ .

**Step 2:**  $\mathcal{G}$  is a minimal set of generators for  $H(P)$ .

**Proposition 3.13.** *For any  $\nu_{h,k} \in \mathcal{G}_1$ ,  $\nu_{h,k} \notin \langle \mathcal{G} \setminus \{\nu_{h,k}\} \rangle$ .*

*Proof.* Suppose, by way of contradiction, that  $\nu_{h,k} \in \langle \mathcal{G} \setminus \{\nu_{h,k}\} \rangle$  for some  $h, k$ . Hence,  $\nu_{h,k} = \sum_i n_i \nu_{h_i, k_i} + \sum_j m_j \mu_{h_j}$  for some  $\nu_{h_i, k_i}, \mu_{h_j} \in \mathcal{G}$  and integers  $n_i, m_j \geq 0$ ; also,  $\sum_i n_i + \sum_j m_j \geq 2$ . Since  $k_i \leq 2h_i - 2$ , we have

$$\begin{aligned} \sum_i n_i \nu_{h_i, k_i} &= \sum_i n_i h_i (q-1) - \sum_i n_i \left( k_i q_0 - \left\lceil \frac{k_i}{2} \right\rceil \right) + \sum_i n_i \\ &\geq \sum_i n_i h_i (q-1) - \sum_i n_i \left( (2h_i - 2)q_0 - \left\lceil \frac{2h_i - 2}{2} \right\rceil \right) + \sum_i n_i \\ &= (q - 2q_0) \sum_i n_i h_i + 2q_0 \sum_i n_i. \end{aligned}$$

As  $\nu_{h,k} \leq q_0 q < \mu_{h_j}$  we have  $m_j = 0$  for any  $j$ . If  $\sum_i n_i h_i \leq h - 1$ , then  $\nu_{h,k} > (h - 1)q \geq \sum_i \nu_{h_i, k_i}$ , a contradiction. If  $\sum_i n_i h_i \geq h + 1$ , then

$$\begin{aligned} \sum_i n_i \nu_{h_i, k_i} &\geq (q - 2q_0)(h + 1) + 2q_0 \sum_i n_i \\ &= hq + 2(q_0 - h)q_0 + 2q_0 \left( -1 + \sum_i n_i \right) > hq \geq \nu_{h,k}, \end{aligned}$$

a contradiction. Then  $\sum_i n_i h_i = h$ . By direct computation,  $\nu_{h,k} = \sum_i n_i \nu_{h_i, k_i}$  is equivalent to

$$(3.17) \quad \left( k - \sum_i n_i k_i \right) q_0 = \left\lceil \frac{k}{2} \right\rceil + 1 - \sum_i n_i \left( \left\lceil \frac{k_i}{2} \right\rceil + 1 \right),$$

and hence  $\left\lceil \frac{k}{2} \right\rceil + 1 \equiv \sum_i n_i \left( \left\lceil \frac{k_i}{2} \right\rceil + 1 \right) \pmod{q_0}$ . Since  $1 \leq \left\lceil \frac{k}{2} \right\rceil + 1 \leq q_0$  and  $2 \leq \sum_i n_i \left( \left\lceil \frac{k_i}{2} \right\rceil + 1 \right) \leq \sum_i n_i h_i = h \leq q_0$ , this implies

$$(3.18) \quad \left\lceil \frac{k}{2} \right\rceil + 1 = \sum_i n_i \left( \left\lceil \frac{k_i}{2} \right\rceil + 1 \right);$$

hence, by (3.17),

$$(3.19) \quad k = \sum_i n_i k_i.$$

From (3.18) and (3.19) it follows

$$\frac{k}{2} + \frac{3}{2} \geq \left\lceil \frac{k}{2} \right\rceil + 1 = \sum_i n_i \left( \left\lceil \frac{k_i}{2} \right\rceil + 1 \right) \geq \sum_i n_i \left( \frac{k_i}{2} + 1 \right) = \frac{k}{2} + \sum_i n_i,$$

a contradiction to  $\sum_i n_i \geq 2$ . □

**Proposition 3.14.** *For any  $\mu_h \in \mathcal{F}_2$ ,  $\nu_{h,k} \notin \langle \mathcal{G} \setminus \{\mu_h\} \rangle$ .*



*Proof.* Suppose, by way of contradiction, that  $\mu_h \in \langle \mathcal{G} \setminus \{\mu_h\} \rangle$  for some  $h \in \{q_0 + 1, \dots, 2q_0\}$ . Hence,  $\mu_h = \sum_i n_i \nu_{h_i, k_i} + \sum_j m_j \mu_{h_j}$  for some  $\nu_{h_i, k_i}, \mu_{h_j} \in \mathcal{G}$  and integers  $n_i, m_j \geq 0$ ; also,  $\sum_i n_i + \sum_j m_j \geq 2$ .

- Assume  $\sum_i m_i \geq 2$ . Then

$$\begin{aligned} \sum_i n_i \nu_{h_i, k_i} + \sum_j m_j \mu_{h_j} &\geq 2((q_0 + 1)q - q_0 - (q_0 - 1)) \\ &= (2q_0 + 2)q - 3q_0 - (q_0 - 2) > \mu_{\tilde{h}} \end{aligned}$$

for any  $\tilde{h}$ , a contradiction.

- Assume  $\sum_i m_i = 1$ . Then  $\mu_h = \sum_i n_i \nu_{h_i, k_i} + \mu_{\tilde{h}}$  for some  $\tilde{h} < h$  and  $\sum_i n_i \geq 1$ . Hence,

$$(h - \tilde{h})q - 2(h - \tilde{h})q_0 = \sum_i n_i \left( h_i q - k_i q_0 - \left\lfloor \frac{2h_i - 2 - k_i}{2} \right\rfloor \right).$$

If  $\sum_i n_i h_i \leq (h - \tilde{h}) - 1$ , then

$$(h - \tilde{h})q - 2(h - \tilde{h})q_0 = \sum_i n_i \left( h_i q - k_i q_0 - \left\lfloor \frac{2h_i - 2 - k_i}{2} \right\rfloor \right) < ((h - \tilde{h}) - 1)q,$$

a contradiction. If  $\sum_i n_i h_i \geq (h - \tilde{h}) + 1$ , then

$$\begin{aligned} (h - \tilde{h})q - 2(h - \tilde{h})q_0 &= \sum_i n_i \left( h_i q - k_i q_0 - \left\lfloor \frac{2h_i - 2 - k_i}{2} \right\rfloor \right) \\ &\geq \sum_i n_i (h_i q - (2h_i - 2)q_0) \\ &\geq (h - \tilde{h})q + q - 2q_0(h - \tilde{h} + 1) + 2q_0 \sum_i n_i \\ &\geq (h - \tilde{h})q + q - 2q_0(h - \tilde{h}) > (h - \tilde{h})q, \end{aligned}$$

a contradiction. Therefore  $\sum_i n_i h_i = h - \tilde{h}$ . By direct computation,  $\mu_h = \sum_i n_i \nu_{h_i, k_i} + \mu_{\tilde{h}}$  is equivalent to

$$(3.20) \quad \left( 2(h - \tilde{h}) - \sum_i n_i k_i \right) q_0 = (h - \tilde{h}) - \sum_i n_i \left( \left\lfloor \frac{k_i}{2} \right\rfloor + 1 \right).$$

Hence  $h - \tilde{h} \equiv \sum_i n_i \left( \left\lfloor \frac{k_i}{2} \right\rfloor + 1 \right) \pmod{q_0}$ ; since the integers  $h - \tilde{h}$  and  $\sum_i n_i \left( \left\lfloor \frac{k_i}{2} \right\rfloor + 1 \right)$  are in  $[1, \dots, q_0 - 1]$ , this implies  $h - \tilde{h} = \sum_i n_i \left( \left\lfloor \frac{k_i}{2} \right\rfloor + 1 \right)$ . By (3.20),  $2(h - \tilde{h}) = \sum_i n_i k_i$ . Thus,

$$\sum_i n_i k_i = 2(h - \tilde{h}) = 2 \sum_i n_i \left( \left\lfloor \frac{k_i}{2} \right\rfloor + 1 \right) \geq \sum_i n_i k_i + 2 \sum_i n_i > \sum_i n_i k_i,$$

a contradiction.

- Assume  $\sum_i m_i = 0$ , that is,  $\mu_h = \sum_i n_i \nu_{h_i, k_i}$  with  $\sum_i n_i \geq 2$ . If  $\sum_i n_i h_i \leq h - 1$ , then  $\mu_h > (h - 1)q \geq \sum_i n_i \nu_{h_i, k_i}$ , a contradiction. If  $\sum_i n_i h_i \geq h + 1$ , then

$$\begin{aligned} \sum_i n_i \nu_{h_i, k_i} &\geq (q - 2q_0) \sum_i n_i h_i + 2q_0 \sum_i n_i \\ &= h(q - 2q_0) + q + 2q_0(-1 + \sum_i n_i) \\ &\geq h(q - 2q_0) + q + 1 = \mu_h, \end{aligned}$$

a contradiction. Then  $\sum_i n_i h_i = h$ . Let  $\hat{h} = h - q_0 \in \{1, \dots, q_0\}$ . By direct computation,  $\mu_h = \sum_i n_i \nu_{h_i, k_i}$  is equivalent to

$$(3.21) \quad \left(2\hat{h} - 1 - \sum_i n_i k_i\right) q_0 = \hat{h} + 1 - \sum_i n_i \left(\left\lceil \frac{k_i}{2} \right\rceil + 1\right);$$

hence,

$$(3.22) \quad \hat{h} + 1 \equiv \sum_i n_i \left(\left\lceil \frac{k_i}{2} \right\rceil + 1\right) \pmod{q_0}.$$

Since

$$2 \leq \sum_i n_i \leq \sum_i n_i \left(\left\lceil \frac{k_i}{2} \right\rceil + 1\right) \leq \sum_i n_i h_i = \hat{h} + q_0,$$

(3.22) implies  $\hat{h} + 1 = \sum_i n_i \left(\left\lceil \frac{k_i}{2} \right\rceil + 1\right)$ . By (3.21),  $2\hat{h} - 1 = \sum_i n_i k_i$ . Thus,

$$\begin{aligned} \sum_i n_i k_i = 2\hat{h} - 1 &= 2 \left( \sum_i n_i \left(\left\lceil \frac{k_i}{2} \right\rceil + 1\right) - 1 \right) - 1 \geq 2 \left( \sum_i n_i \left(\frac{k_i}{2} + 1\right) - 1 \right) - 1 \\ &= \sum_i n_i k_i + 2 \sum_i n_i - 3 > \sum_i n_i k_i, \end{aligned}$$

a contradiction. □

#### 4. DUAL ONE-POINT CODES FROM THE SUZUKI CURVE

In this section we construct dual one-point codes from  $\mathcal{S}_q$ . In particular, we consider codes of type  $\mathcal{C}_i = C_{\mathcal{L}}(mP_i, D_i)^\perp$ , where

$$P_1 \in \mathcal{S}_q(\mathbb{F}_{q^4}) \setminus \mathcal{S}_q(\mathbb{F}_q), \quad P_2 \in \mathcal{S}_q(\mathbb{F}_q), \quad D_i = \sum_{P \in \mathcal{S}_q(\mathbb{F}_{q^4}) \setminus P_i} P.$$

For a more detailed introduction on AG codes we refer the readers to [14].

Also, estimates on the minimum distance are obtained using the so-called Feng-Rao function. Let  $\mathcal{X}$  be a nonsingular curve and  $P$  an  $\mathbb{F}_q$ -rational point of  $\mathcal{X}$ . Let the Weierstrass

TABLE 1. Comparison between the codes  $\mathcal{C}_1$  and  $\mathcal{C}_2$ ,  $q = 8$ 

$\rho_\ell$	$n$	$n - \ell$	$d(\mathcal{C}_1)$	$d(\mathcal{C}_2)$
34	4124	4103	10	8
35	4124	4102	12	10
36	4124	4101	12	10

semigroup at  $P$  be given by

$$(4.1) \quad H(P) = \{\rho_1 = 0 < \rho_2 < \rho_3 < \dots\}.$$

For  $\ell \geq 0$ , define the *Feng-Rao function*

$$\nu_\ell := |\{(i, j) \in \mathbb{N}_0^2 : \rho_i + \rho_j = \rho_{\ell+1}\}|.$$

Consider  $C_\ell(P) = C_{\mathcal{L}}(P_1 + P_2 + \dots + P_N, \rho_\ell P)^\perp$ , with  $P, P_1, \dots, P_N$  pairwise distinct points in  $\mathcal{X}(\mathbb{F}_q)$ . The number

$$d_{ORD}(C_\ell(P)) := \min\{\nu_m : m \geq \ell\}$$

is a lower bound for the minimum distance  $d(C_\ell(P))$  of the code  $C_\ell(P)$ , called the *order bound* or the *Feng-Rao designed minimum distance* of  $C_\ell(P)$ ; see [10, Theorem 4.13]. Also, by [10, Theorem 5.24],  $d_{ORD}(C_\ell(P)) \geq \ell + 1 - g$  and equality holds if  $\ell \geq 2c - g - 1$ , where  $c = \max\{m \in \mathbb{Z} : m - 1 \notin H(P)\}$  is the conductor of  $H(P)$ .

Note that length and dimension of  $C_{\mathcal{L}}(D_1, \rho_\ell P_1)^\perp$  and  $C_{\mathcal{L}}(D_2, \rho_\ell P_2)^\perp$  coincide; here, the coefficients  $\rho_\ell$  of  $P_1$  and  $P_2$  are the  $\ell$ -th non-gap at  $P_1$  and the  $\ell$ -th non-gap at  $P_2$ , respectively (see (4.1)).

We list in Tables 1 and 2 the parameters of the codes  $C_{\mathcal{L}}(D_i, \rho_\ell P_i)^\perp$  for which the minimum distance of  $C_{\mathcal{L}}(D_1, \rho_\ell P_1)^\perp$  is larger than the one of  $C_{\mathcal{L}}(D_2, \rho_\ell P_2)^\perp$ , when  $q \in \{8, 32\}$ . Computations have been made comparing the Feng-Rao designed minimum distance of both the codes and have been performed using MAGMA [2].

#### ACKNOWLEDGMENTS

The research of D. Bartoli, M. Montanucci, and G. Zini was partially supported by Ministry for Education, University and Research of Italy (MIUR) (Project PRIN 2012 ‘‘Geometrie di Galois e strutture di incidenza’’ - Prot. N. 2012XZE22K\_005) and by the Italian National Group for Algebraic and Geometric Structures and their Applications (GNSAGA - INdAM).

TABLE 2. Comparison between the codes  $\mathcal{C}_1$  and  $\mathcal{C}_2$ ,  $q = 32$ 

$\rho_\ell$	$n$	$n - \ell$	$d(\mathcal{C}_1)$	$d(\mathcal{C}_2)$	$\rho_\ell$	$n$	$n - \ell$	$d(\mathcal{C}_1)$	$d(\mathcal{C}_2)$
261	1048824	1048686	38	32	262	1048824	1048685	38	32
263	1048824	1048684	38	32	264	1048824	1048683	38	32
265	1048824	1048682	40	32	266	1048824	1048681	40	32
267	1048824	1048680	40	32	268	1048824	1048679	40	32
269	1048824	1048678	40	32	270	1048824	1048677	40	32
271	1048824	1048676	40	32	272	1048824	1048675	40	32
273	1048824	1048674	40	32	274	1048824	1048673	40	32
275	1048824	1048672	40	32	276	1048824	1048671	40	32
277	1048824	1048670	40	32	278	1048824	1048669	40	32
279	1048824	1048668	40	36	280	1048824	1048667	40	36
281	1048824	1048666	40	36	282	1048824	1048665	40	36
285	1048824	1048662	42	40	286	1048824	1048661	42	40
287	1048824	1048660	42	41	297	1048824	1048650	66	64
298	1048824	1048649	66	64	299	1048824	1048648	66	64
300	1048824	1048647	66	64	301	1048824	1048646	66	64
302	1048824	1048645	66	64	303	1048824	1048644	66	64
304	1048824	1048643	66	64	305	1048824	1048642	66	64
306	1048824	1048641	66	64	307	1048824	1048640	66	64
308	1048824	1048639	66	64	309	1048824	1048638	66	64
310	1048824	1048637	66	64	313	1048824	1048634	70	68
314	1048824	1048633	70	68	317	1048824	1048630	73	72
318	1048824	1048629	73	72	321	1048824	1048626	80	76
322	1048824	1048625	82	76	323	1048824	1048624	84	77
324	1048824	1048623	84	80	325	1048824	1048622	88	80
326	1048824	1048621	88	80	327	1048824	1048620	88	81
328	1048824	1048619	88	82	341	1048824	1048606	97	96
342	1048824	1048605	97	96	353	1048824	1048594	109	108
354	1048824	1048593	112	108	355	1048824	1048592	112	109
357	1048824	1048590	114	112	358	1048824	1048589	114	112
359	1048824	1048588	114	113	361	1048824	1048586	118	116
362	1048824	1048585	118	116	363	1048824	1048584	118	117
365	1048824	1048582	121	120	366	1048824	1048581	121	120
369	1048824	1048578	124	123	386	1048824	1048561	142	140
387	1048824	1048560	142	141	389	1048824	1048558	145	144
390	1048824	1048557	145	144					

## REFERENCES

- [1] P. Beelen and M. Montanucci, A new family of maximal curves, *J. London Math. Soc. (2)*, 2018, doi:10.1112/jlms.12144.
- [2] W. Bosma, J. Cannon, and C. Playoust, The Magma algebra system. I. The user language, *J. Symbolic Comput.* **24**, 235–265, 1997.

- [3] P. Deligne and G. Lusztig, Representations of reductive groups over finite fields, *Ann. Math.*, **103**, 103–161, 1976.
- [4] R. Fuhrmann, F. Torres, On Weierstrass points and optimal curves, *Rend. Circ. Mat. Palermo (2)* **51**, 25–46 (1998).
- [5] A. Garcia, C. Güneri, and H. Stichtenoth, A generalization of the Giulietti-Korchmáros maximal curve, *Advances in Geometry*, **10**(3), 427–434, 2010.
- [6] Garcia, A., Stichtenoth, H., Elementary abelian  $p$ -extensions of algebraic functions fields, *Manuscripta Math.* **72**, 67–79 (1991).
- [7] M. Giulietti and G. Korchmáros, A new family of maximal curves over a finite field, *Math. Ann.*, **343**, 229–245, 2009.
- [8] C. Heegard, J. Little, and K. Saints, Systematic encoding via Gröbner bases for a class of algebraic-geometric Goppa codes, *IEEE Trans. Inf. Theory*, **41**, 1752–1761, 1995.
- [9] J.W.P. Hirschfeld, G. Korchmáros, F. Torres, *Algebraic Curves over a Finite Field*. Princeton Series in Applied Mathematics, Princeton (2008).
- [10] T. Høholdt, J. H. van Lint, and R. Pellikaan, Algebraic geometry codes, in: *Handbook of Coding Theory*, V. S. Pless, W. C. Huffman, and R. A. Brualdi, Eds. Amsterdam, The Netherlands: Elsevier, 1998, vol. 1, pp. 871–961.
- [11] D. Joyner, An error-correcting codes package, *SIGSAM Comm. Computer Algebra*, **39**(2), 65–68, 2005.
- [12] S. Meagher and J. Top, Twists of genus three curves over finite fields, *Finite Fields Appl.*, **16**, 347–368, 2010.
- [13] G. Matthews, Codes from the Suzuki function field, *IEEE Trans. Inform. Theory* **50**(12), 3298–3302, (2004).
- [14] H. Stichtenoth, *Algebraic function fields and codes*, Springer, 2009.
- [15] K.O. Stöhr, J.F. Voloch, Weierstrass points and curves over finite fields, *Proc. London Math. Soc.* **52**(3), 1–19 (1986).
- [16] M.A. Tsfasman, G. Vladut, *Algebraic-geometric Codes*, Kluwer, Dordrecht, (1991).

Daniele Bartoli

Università degli Studi di Perugia,  
Dipartimento di Matematica e Informatica,  
Via Luigi Vanvitelli 1,  
06123 Perugia,  
Italy,  
daniele.bartoli@unipg.it

Maria Montanucci

Università degli Studi di Padova,  
Dipartimento di Tecnica e Gestione dei Sistemi Industriali,  
Stradella S. Nicola 3,  
36100 Vicenza,  
Italy,  
montanucci@gest.unipd.it

Giovanni Zini

Università degli Studi di Milano-Bicocca  
Dipartimento di Matematica e Applicazioni,  
Via Cozzi 55,  
20125 Milano,  
Italy,  
[giovanni.zini@unimib.it](mailto:giovanni.zini@unimib.it)