

A note on long rainbow arithmetic progressions

Jesse Geneson

ISU

geneson@gmail.com

March 14, 2022

Abstract

Jungić et al (2003) defined T_k as the minimal number $t \in \mathbb{N}$ such that there is a rainbow arithmetic progression of length k in every equinumerous t -coloring of $[tn]$ for every $n \in \mathbb{N}$. They proved that for every $k \geq 3$, $\lfloor \frac{k^2}{4} \rfloor < T_k \leq \frac{k(k-1)^2}{2}$ and conjectured that $T_k = \Theta(k^2)$. We prove for all $\epsilon > 0$ that $T_k = O(k^{5/2+\epsilon})$ using the Kővári-Sós-Turán theorem and Wigert's bound on the divisor function.

A coloring is called *equinumerous* if each color is used an equal number of times. We call an arithmetic progression *rainbow* if it does not contain any terms of the same color. We use the notation $AP(k)$ for arithmetic progression of length k , so T_k is the minimal number $t \in \mathbb{N}$ such that there is a rainbow $AP(k)$ in every equinumerous t -coloring of $[tn]$ for every $n \in \mathbb{N}$.

Besides the bounds $\lfloor \frac{k^2}{4} \rfloor < T_k \leq \frac{k(k-1)^2}{2}$ for every $k \geq 3$ from [4], it is also known that $T_k > k$ for every $k > 5$ [1], $T_3 = 3$ [1, 5], and $T_4 > 4$ [3]. A variant of T_k in which the coloring is not required to be equinumerous was investigated in [2]. In this note, we prove for all $\epsilon > 0$ that $T_k = O(k^{5/2+\epsilon})$.

Our proof represents the occurrences of a single color in $[tn]$ with a family of 0-1 matrices, and we bound T_k using the Kővári-Sós-Turán theorem. We say that a 0-1 matrix A *avoids* a 0-1 matrix P if there is no submatrix in A that is equal to P or that can be transformed into P by changing some ones to zeroes. Define $ex(m, n, P)$ as the maximum number of ones in an $m \times n$ 0-1 matrix that avoids P . We let $R_{s,t}$ denote the $s \times t$ matrix of all ones.

Theorem 1. [6] For all $m, n, s, t \geq 2$, $ex(m, n, R_{s,t}) \leq (s-1)^{1/t}(n-t+1)m^{1-1/t} + (t-1)m$.

Let $\tau(n)$ be the number of positive integer divisors of n . We use Wigert's upper bound on $\tau(n)$ [7] for our bound on T_k .

Theorem 2. [7] $\tau(n) = n^{O(\frac{1}{\ln \ln n})}$

Before we prove the new bound, we start with the proof of the $O(k^3)$ upper bound from [4] since the beginning of our proof is the same. There was a small typo in the proof in [4] for the value of the number of $AP(k)$ s in $[m]$, which we correct in the version below.

Theorem 3. [4] $T_k = O(k^3)$

Proof. Let $m = a(k-1) + b$ with $k \geq 3$, $a \geq 1$, and $0 \leq b < k-1$. Since there is a bijective correspondence between the set of all $AP(k)$ s and the set of all 2-element sets $\{x, y\} \subset [m]$, $x < y$, with $x \equiv y \pmod{k-1}$, the number of all $AP(k)$ s in $[m]$ is $b \binom{a+1}{2} + (k-b-1) \binom{a}{2}$, implying that the number of $AP(k)$ s in $[tn]$ is greater than $\frac{tn(tn-3(k-1))}{2(k-1)}$.

For any equinumerous t -coloring of $[tn]$, in each color there are $\binom{n}{2}$ pairs of numbers. Each pair can be terms of at most $\binom{k}{2}$ different $AP(k)$ s. Thus for any equinumerous t -coloring of $[tn]$, there are at most $t \binom{k}{2} \binom{n}{2}$ $AP(k)$ s that are not rainbow. Thus T_k is bounded by the smallest t satisfying $\frac{tn(tn-3(k-1))}{2(k-1)} \geq t \binom{k}{2} \binom{n}{2}$ for all n , giving the bound. \square

Theorem 4. $T_k = O(k^{5/2+\epsilon})$ for all $\epsilon > 0$

Proof. Let t be minimal so that there is a rainbow $AP(k)$ in every equinumerous t -coloring of $[tn]$ for all $n \in \mathbb{N}$. From the bounds proved in [4], we may assume that $t < k^3$.

The overall method of the proof is the same as Jungić et al [4], until the point where we find an upper bound on the number of $AP(k)$ s in $[tn]$ that are not rainbow. For this bound, we split into cases depending on whether $n \leq k$.

If $n \leq k$, then $tn < k^4$. In each color, there are $\binom{n}{2}$ pairs of numbers. The difference between the numbers in each pair is less than k^4 , so that difference has at most $k^{o(1)}$ divisors by Theorem 2. Thus each pair can be terms of at most $k^{1+o(1)}$ different $AP(k)$ s. Therefore for any t -regular coloring of $[tn]$, there are at most $t \binom{n}{2} k^{1+o(1)}$ $AP(k)$ s that are not rainbow. The smallest t satisfying $\frac{tn(tn-3(k-1))}{2(k-1)} \geq t \binom{n}{2} k^{1+o(1)}$ for all $n \leq k$ is $O(k^{2+\epsilon})$ for all $\epsilon > 0$.

Now if $n > k$, fix an arbitrary color c_i and let $a_1 < \dots < a_n$ be the n numbers in $[tn]$ colored with c_i . We call the gap between a_j and a_{j+1} *wide* if $a_{j+1} - a_j > k^8$. By the pigeonhole principle, there are at most $\frac{tn}{k^8} < \frac{n}{k^5}$ wide gaps between the consecutive numbers colored with c_i .

Partition the numbers colored with c_i into *blocks* of k consecutive numbers, where only the last block may have fewer than k numbers. We say a block is wide if it contains both of the consecutive numbers in a wide gap. Since there are at most $\frac{n}{k^5}$ wide gaps between the consecutive numbers colored with c_i , there are at most $\frac{n}{k^5}$ wide blocks. In all other blocks, all of the gaps are not wide.

Fix $1 \leq j \leq n-1$. The number of a_r that lie in a wide block is at most $\frac{n}{k^4}$ since there are at most $\frac{n}{k^5}$ wide blocks and each block contains at most k numbers. Thus there are at most $\frac{n}{k^4} \binom{k}{2}$ $AP(k)$ s containing both a_j and a_r for some r such that $j < r$ and a_r lies in a wide block.

Next we bound the number of $AP(k)$ s containing both a_j and a_r for some r such that $j < r$ and a_r lies in a non-wide block. Define a family of 0-1 matrices $\{A^s\}_{1 \leq s \leq \lceil \frac{n}{k} \rceil}$ corresponding to the blocks, each with $k-1$ rows and at most k columns, such that $A^s_{x,y} = 1$ if and only if x divides $a_{(s-1)k+y} - a_j$.

For any fixed r with $j < r$, the number of $AP(k)$ s containing both a_j and a_r is at most $k\tau(a_r - a_j)$. Thus the number of $AP(k)$ s containing both a_j and a_r for some r such that $j < r$ and a_r lies in the block s is at most k times the number of ones in A^s .

By Theorem 2, A^s avoids $R_{q,2}$, where $q = k^{o(1)}$ for non-wide s . By the Kővári-Sós-Turán theorem, A^s has at most $k^{3/2+o(1)} + k$ ones for non-wide s . Thus the number of $AP(k)$ s containing both a_j and a_r for some r such that $j < r$ and a_r lies in the non-wide block s is at most $k^{5/2+o(1)} + k^2$.

Since there are at most $\lceil \frac{n}{k} \rceil \leq \frac{2n}{k}$ non-wide blocks, there are at most $tn(\frac{2n}{k}(k^{5/2+o(1)} + k^2) + \frac{n}{k^4} \binom{k}{2})$ $AP(k)$ s in $[tn]$ that are not rainbow. The smallest t satisfying $\frac{tn(tn-3(k-1))}{2(k-1)} \geq tn(\frac{2n}{k}(k^{5/2+o(1)} + k^2) + \frac{n}{k^4} \binom{k}{2})$ for all $n > k$ is $O(k^{5/2+\epsilon})$ for all $\epsilon > 0$, completing the proof. \square

References

- [1] M. Axenovich, D. Fon-Der-Flaass: On rainbow arithmetic progressions, *Electronic Journal of Combinatorics*, 11, (2004), R1.
- [2] S. Butler, C. Erickson, L. Hogben, K. Hogenson, L. Kramer, R. Kramer, J. Lin, R. Martin, D. Stolee, N. Warnberg, and M. Young, Rainbow arithmetic progressions, *Journal of Combinatorics* 7 (2016), 595-626.
- [3] D. Conlon, V. Jungić, R. Radoičić, On the existence of rainbow 4-term arithmetic progressions, *Graphs and Combinatorics*, 23 (2007), 249-254.
- [4] V. Jungić, J. Licht (Fox), M. Mahdian, J. Nešetřil, R. Radoičić, Rainbow arithmetic progressions and anti-Ramsey results, *Combinatorics, Probability, and Computing - Special Issue on Ramsey Theory*, 12, (2003), 599-620.
- [5] V. Jungić, R. Radoičić, Rainbow 3-term arithmetic progressions, *Integers, The Electronic Journal of Combinatorial Number Theory*, 3, (2003), A18.
- [6] T. Kővári, V. Sós, P. Turán, On a problem of K. Zarankiewicz, *Colloquium Math*, 3:50-57, 1954.
- [7] S. Wigert, Sur l'ordre de grandeur du nombre des diviseurs d'un entier. *Arkiv för matematik, astronomi och fysik* 3, no. 18 (1907), 19.