On the Open Question of The Tracy-Widom Distribution of β -Ensemble With $\beta = 6$

Li YuQi

May 24, 2021

Abstract: We determine completely the Tracy-Widom distribution for Dysons β -ensemble with $\beta = 6$. The problem of the Tracy-Widom distribution of β -ensemble for general $\beta > 0$ has been reduced to find out a bounded solution of the Bloemendal-Virág equation with a specified boundary. Rumanov proposed a Lax pair approach to solve the Bloemendal-Virág equation for even integer β . He also specially studied the $\beta = 6$ case with his approach and found a second order nonlinear ordinary differential equation (ODE) for the logarithmic derivative of the TracyWidom distribution for $\beta = 6$. Grava et al. continued to study $\beta = 6$ and found Rumanov's Lax pair is gauge equivalent to that of Painlevé II in this case. They started with Rumanov's basic idea and came down to two auxiliary functions $\alpha(t)$ and $q_2(t)$, which satisfy a coupled first-order ODE. The open question by Grava et al. asks whether a global smooth solution of the ODE with boundary condition $\alpha(\infty) = 0$ and $q_2(\infty) = -1$ exists. By studying the linear equation that is associated with q_2 and α , we give a positive answer to the open question. Moreover, we find that the solutions of the ODE with $\alpha(\infty) = 0$ and $q_2(\infty) = -1$ are parameterized by c_1 and c_2 . Not all c_1 and c_2 give global smooth solutions. But if $(c_1, c_2) \in R_{smooth}$, where R_{smooth} is a large region containing (0,0), they do give. We prove the constructed solution is a bounded solution of the Bloemendal-Virág equation with the required boundary condition if and only if $(c_1, c_2) = (0, 0).$

1 Introduction

In the one dimensional case, the interaction energy of two point charges is

$$k_e \ln(\frac{r_0}{|x_A - x_B|}),$$

where k_e is the electric force constant, r_0 is the distance that the interaction energy is 0, x_A and x_B are the positions of the two point charges. Dyson's Coulomb gas model is N particles with like charges, i.e., $k_e > 0$, in an external field V = V(x). By the canonical ensemble, the probability that the first particle is in $[x_1, x_1 + dx_1], \dots$, and that N-th particle is in $[x_N, x_N + dx_N]$, is

$$p(x_{1}, x_{2}, \dots, x_{N}) dx_{1} dx_{2} \dots dx_{N} = \frac{1}{Z_{N}} e^{-\frac{1}{k_{B}T} \left(\sum_{1 \leq i < j \leq N} k_{e} \ln \left(\frac{r_{0}}{|x_{i} - x_{j}|} \right) + \sum_{j=1}^{N} V(x_{j}) \right)} dx_{1} dx_{2} \dots dx_{N}$$

$$= \frac{1}{Z_{N}} \left(\prod_{1 \leq i < j \leq N} \left| \frac{x_{i} - x_{j}}{r_{0}} \right|^{\frac{k_{e}}{k_{B}T}} \right) e^{-\frac{1}{k_{B}T} \sum_{j=1}^{N} V(x_{j})} dx_{1} dx_{2} \dots dx_{N},$$

where k_B is the Boltzmann constant, T is the temperature, and Z_N is the normalization constant. Here we assume V is Gaussian, i.e.,

(1.1)
$$V(x) = \frac{1}{2}\nu x^2.$$

Let

$$\beta = \frac{k_e}{k_B T}, \quad \lambda = \sqrt{\frac{\nu}{k_e}} x.$$

Then the particle distribution becomes

$$(1.2) \quad \tilde{p}(\lambda_1, \lambda_2, \cdots, \lambda_N) d\lambda_1 d\lambda_2 \cdots d\lambda_N = \frac{1}{\tilde{Z}_N} \left(\prod_{1 \leq i < j \leq N} |\lambda_i - \lambda_j|^{\beta} \right) e^{-\frac{\beta}{2} \sum\limits_{j=1}^N \lambda_j^2} d\lambda_1 d\lambda_2 \cdots d\lambda_N.$$

A system of random variables $\lambda_1, \lambda_2, \dots, \lambda_N$ with distribution (1.2) is called the β -ensemble. The β -ensemble, $\beta = 1, 2, 4$, describes the joint density of eigenvalues of the three classical matrix models, i.e., the Gaussian orthogonal ensemble(GOE), Gaussian unitary ensemble(GUE) and Gaussian symplectic ensemble(GSE), respectively. For general $\beta > 0$, the β -ensemble can be realised as the joint density of eigenvalues of the *spiked* β -Hermite matrix ensemble [2]. β -ensemble for general β has also other physical applications, for example, it can be mapped to a chiral Liouville theory with central charge [8]. Also, in some sense, the harmonic potential (1.1) is not so serious a limitation since the universality of the β -ensemble had been proved by Bourgade et al. [4].

The interesting case is the thermodynamic limit $N \to \infty$. Almost all particles distribute in $[-\sqrt{2N}, \sqrt{2N}]$ obeying the Wigner semicircle law with an approximate density $\sigma(\lambda) = \pi^{-1}\sqrt{2N-\lambda^2}$ [15], i.e., the particle number in $[\lambda, \lambda + d\lambda]$ is about $\sigma(\lambda)d\lambda$. But few particles may lie outside $[-\sqrt{2N}, \sqrt{2N}]$. It is proved that near the edge a proper scaling limit is the soft edge probability distribution [16]

$$F_{\beta}(t) = \lim_{N \to \infty} E_{\beta N}^{\text{Soft}} \left(0; \left(\sqrt{2N} + \frac{t}{\sqrt{2}N^{1/6}}, \infty \right) \right),$$

where

$$E_{\beta N}^{\text{Soft}}\left(0;(t,\infty)\right) = \int_{\lambda_N = -\infty}^t \cdots \int_{\lambda_1 = -\infty}^t \tilde{p}(\lambda_1, \cdots, \lambda_N) d\lambda_1 \cdots d\lambda_N.$$

 $F_{\beta}(t)$ is called the Tracy-Widom distribution.

The explicit expressions for $F_{\beta}(t)$ for $\beta = 1, 2, 4$ are classical [19, 20, 16]

$$F_{\beta}(t) = \begin{cases} \exp\left(\frac{1}{2}\int_{\infty}^{t}(s-t)u(s)^{2}ds\right)\exp\left(\frac{1}{2}\int_{\infty}^{t}u(s)ds\right), & \beta = 1, \\ \exp\left(\int_{\infty}^{t}(s-t)u(s)^{2}ds\right), & \beta = 2, \\ \exp\left(\frac{1}{2}\int_{\infty}^{2^{2/3}t}(s-2^{2/3}t)u(s)^{2}ds\right)\cosh\left(\frac{1}{2}\int_{\infty}^{2^{2/3}t}u(s)ds\right), & \beta = 4. \end{cases}$$

Also $F_2(t)$ has a Fredholm determinant representation $F_2(t) = \det(I - A_t)$, where A_t is the Fredholm integral operator on (t, ∞) with the Airy kernel $\frac{\operatorname{Ai}(x)\operatorname{Ai}'(y) - \operatorname{Ai}'(x)\operatorname{Ai}(y)}{x-y}$.

The expansions of $F_{\beta}(t)$ at $t = -\infty$ are of special interests. In [19] and [20], Tracy and Widom obtained and proved $F_{\beta}(t)$ for $\beta = 1, 2, 4$ without the constant term. They also conjectured the values of the constant term c_0 for $\beta = 1, 2, 4$. By the Deift-Zhou nonlinear steepest descent method [6], Deift et al. [5] proved the constant term for $\beta = 2$ and Baik et al. [1] proved the constant terms for $\beta = 1, 2, 4$. Finally Borot et al. [3] derived an amazing asymptotic expression of $F_{\beta}(t)$ for general $\beta > 0$ at $t = -\infty$ by the loop-equation technique. Their asymptotic expression is

$$(1.3) F_{\beta}(t) = \exp\left(-\frac{\beta}{24}|t|^3 + \frac{\sqrt{2}}{3}(\frac{\beta}{2} - 1)|t|^{\frac{3}{2}} + \frac{1}{8}(\frac{\beta}{2} + \frac{2}{\beta} - 3)\ln|t| + c_{\beta} + O(|t|^{-\frac{3}{2}})\right),$$

where the constant term c_{β} is

$$c_{\beta} = \frac{\gamma_{E}}{6\beta} + \left(\frac{17}{8} - \frac{25}{24}(\frac{\beta}{2} + \frac{2}{\beta})\right) \ln 2 - \frac{1}{2}\ln(\frac{\beta}{2}) - \frac{\ln(2\pi)}{4} + \frac{\beta}{2}\left(\frac{1}{12} - \zeta'(-1)\right) + \int_{0}^{\infty} \frac{1}{t^{2}} \frac{1}{e^{\beta t/2} - 1} \left(\frac{t}{e^{t} - 1} - 1 + \frac{t}{2} - \frac{t^{2}}{12}\right) dt.$$
(1.4)

Here γ_E denotes the Euler's constant and ζ refers to the Riemann zeta function. Note the prime 'will always be used to denote derivative.

But the asymptotics (1.3) is only valid at $t = -\infty$ and can not be continued to finite t. In fact, even with infinite terms (1.3) can not determine $F_{\beta}(t)$, see Theorem 1.2 below. So we still need the explicit expression for $F_{\beta}(t)$ beyond $\beta = 1, 2, 4$. Following the pioneering work of Dumitriu and Edelman [7], Bloemendal and Virág [2] finally found out a representation of $F_{\beta}(t)$ in terms of the solution of a linear partial differential equation(PDE). They represent $F_{\beta}(t)$ by the limit of $F(\beta; x, t)$

(1.5)
$$F_{\beta}(t) = \lim_{x \to \infty} F(\beta; x, t),$$

where $F(\beta; x, t)$ is a special solution of the linear PDE

(1.6)
$$\frac{\partial F}{\partial t} + \frac{2}{\beta} \frac{\partial^2 F}{\partial x^2} + (t - x^2) \frac{\partial F}{\partial x} = 0.$$

More precisely, they proved the following theorem.

Theorem 1.1. [2] PDE(1.6) with boundary

(1.7)
$$\begin{cases} F(\beta; x, t) \xrightarrow{x \to \infty, t \to \infty} 1 \\ F(\beta; x, t) \xrightarrow{x \to -\infty, t \text{ fixed}} 0 \end{cases}$$

has a unique bounded smooth solution. $F_{\beta}(t)$ is represented by the solution through (1.5).

So the remaining problem is to find a bounded solution of the Bloemendal-Virág equation (1.6) with the boundary condition (1.7). In [17], Rumanov proposed a Lax representation of (1.6) for even integer β . Let

$$\Psi_x = \hat{L}\Psi, \quad \Psi_t = \hat{B}\Psi$$

be Rumanov's Lax pair, where \hat{L} and \hat{B} are 2×2 matrices. Ψ can denote both a 2×2 non-singular matrix or a 2×1 column vector. Here we assume it is a column vector

(1.8)
$$\Psi(x,t) = \begin{pmatrix} \mathcal{F}(x,t) \\ \mathcal{G}(x,t) \end{pmatrix}.$$

The key of Rumanov's scheme is to let \mathcal{F} satisfy the rescaled Bloemendal-Virág equation

(1.9)
$$\frac{\beta}{2} \frac{\partial \mathcal{F}}{\partial t} + \frac{\partial^2 \mathcal{F}}{\partial x^2} + (t - x^2) \frac{\partial \mathcal{F}}{\partial x} = 0.$$

Combining some other considerations, Rumanov concluded

(1.10)
$$F(\beta; x, t) = \mathcal{F}\left(\left(\frac{\beta}{2}\right)^{\frac{1}{3}} x, \left(\frac{\beta}{2}\right)^{\frac{2}{3}} t\right).$$

In [18], Rumanov studied the $\beta = 6$ case and expressed $F_{\beta}(t)$ by an auxiliary function $\eta(t)$ and the solution of Painlevé II

$$(1.11) u''(t) = tu(t) + 2u(t)^3$$

which he deduced as the Hastings-McLeod solution [12]. The auxiliary function $\eta(t)$ satisfies a second order ODE that can be linearized.

Grava, Its, Kapaev and Mezzadri [11] found Rumanov's Lax pair for $\beta = 6$ is gauge equivalent to the Lax pair of Painlevé II. Their gauge transformation is of form

(1.12)
$$\Psi(x,t) = e^{\frac{x^3}{6} - \frac{xt}{2}} \kappa(t) \begin{pmatrix} \frac{1 + q_2(t)}{2} x - \alpha(t) & -1\\ \frac{1 - q_2^2(t)}{4} & 0 \end{pmatrix} \psi(t)^{\sigma_3} \psi_0(x,t),$$

where $\kappa(t)$ and $\psi(t)$ are scalar functions, and $\psi_0(x,t)$ is the 2×1 column vector of the wave function of Painlevé II, and σ_3 is the Pauli matrix $\sigma_3 = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}$. They also suggested $\psi = -\frac{\mathrm{i}}{\sqrt{u}}$, where u is the Hastings-McLeod solution of Painlevé II. Then they showed $q_2(t)$ and $\alpha(t)$ satisfy the ODE

(1.13)
$$q_2'(t) = \frac{2}{3}\alpha q_2 + \frac{u'}{u} \frac{(1+q_2)(2-q_2)}{3},$$

(1.14)
$$\alpha'(t) = \alpha \left(\frac{2}{3}\alpha + \frac{u'}{u}\frac{2-q_2}{3}\right) - \frac{t}{6}(1+q_2) - \frac{u^2}{3}(3+q_2).$$

Moreover, they proved that

$$(1.15) q_2(t) \xrightarrow{t \to \infty} -1 + o(1), \quad \alpha(t) \xrightarrow{t \to \infty} o(1).$$

It is straightforward to verify that

$$(1.16) q_2(t) = \frac{1}{\sqrt{2}}(-t)^{-3/2} + \frac{21}{8}(-t)^{-3} + \frac{1707}{64\sqrt{2}}(-t)^{-9/2} + \frac{49123}{256}(-t)^{-6} + \cdots,$$

(1.17)
$$\alpha(t) = \frac{1}{\sqrt{2}}(-t)^{1/2} - \frac{1}{8}(-t)^{-1} - \frac{37}{64\sqrt{2}}(-t)^{-5/2} - \frac{373}{256}(-t)^{-4} + \cdots$$

is an asymptotic solution of Equation (1.13)-(1.14) at $t = -\infty$.

The open question in [11] contains two parts:

- (1). Prove the system (1.13)-(1.14) with (1.15) has a smooth solution on $(-\infty, \infty)$.
- (2). Assume (1) succeeds. Prove the solution in (1) has expansions (1.16)-(1.17) at $t = -\infty$.

In this paper we will show that there are a 2-parameter family of solutions of (1.13)-(1.14) that satisfy $q_2 \xrightarrow{t \to \infty} -1$ and $\alpha \xrightarrow{t \to \infty} 0$. More precisely, at $t = \infty$ these solutions have asymptotics

(1.18)
$$q_2(t) = -1 + \operatorname{Ai}(t) \left[c_1 \operatorname{Bi}(\frac{t}{3^{2/3}}) + c_2 \operatorname{Ai}(\frac{t}{3^{2/3}}) + c_1^2 \tilde{M}_2(t) \right] + o\left(e^{-\frac{4}{3}t^{3/2}}\right),$$

(1.19)
$$\alpha(t) = -\frac{3^{1/3}}{2} \operatorname{Ai}(t) \left[c_1 \operatorname{Bi}'(\frac{t}{3^{2/3}}) + c_2 \operatorname{Ai}'(\frac{t}{3^{2/3}}) + c_1^2 \tilde{N}_2(t) \right] + o\left(e^{-\frac{4}{3}t^{3/2}}\right),$$

where $\tilde{M}_2(t) = o\left(\operatorname{Ai}\left(\frac{t}{3^{2/3}}\right)\right)$ and $\tilde{N}_2(t) = o\left(\operatorname{Ai}'\left(\frac{t}{3^{2/3}}\right)\right)$. The detailed expressions of $\tilde{M}_2(t)$ and $\tilde{N}_2(t)$ will be given in Section 5. If $c_1 \neq 0$ or $c_2 \neq 0$, the leading terms in the asymptotics (1.18) and (1.19) are obvious. If $c_1 = 0$ and $c_2 = 0$, the leading terms for the asymptotics of $q_2 + 1$ and α are

$$q_{2} + 1 = \frac{4\pi}{3^{4/3}} \operatorname{Ai}(t) \left[\operatorname{Bi}\left(\frac{t}{3^{2/3}}\right) \int_{\infty}^{t} \operatorname{Ai}(s) \operatorname{Ai}\left(\frac{s}{3^{2/3}}\right) ds - \operatorname{Ai}\left(\frac{t}{3^{2/3}}\right) \int_{\infty}^{t} \operatorname{Ai}(s) \operatorname{Bi}\left(\frac{s}{3^{2/3}}\right) ds \right]$$

$$(1.20) + o\left(e^{-\frac{8}{3}t^{3/2}}\right),$$

$$\alpha = \frac{2\pi}{3} \operatorname{Ai}(t) \left[\operatorname{Ai}'\left(\frac{t}{3^{2/3}}\right) \int_{\infty}^{t} \operatorname{Ai}(s) \operatorname{Bi}\left(\frac{s}{3^{2/3}}\right) ds - \operatorname{Bi}'\left(\frac{t}{3^{2/3}}\right) \int_{\infty}^{t} \operatorname{Ai}(s) \operatorname{Ai}\left(\frac{s}{3^{2/3}}\right) ds \right]$$

$$(1.21) + o\left(\sqrt{t}e^{-\frac{8}{3}t^{3/2}}\right).$$

Not all solutions with asymptotics of (1.18) and (1.19) at $t = \infty$ can be smoothly evolved to $t = -\infty$. It may develop to singularity at $t = t_0$, which depends on c_1 and c_2 . Our first main result of this paper is the following.

Theorem 1.2. There is a region R_{smooth} that is the neighbourhood of the positive c_2 -axis including the origin (0,0) in the (c_1,c_2) -plane, such that if (c_1,c_2) is in the region R_{smooth} then the solution defined by the asymptotics of (1.18)-(1.19) at $t=\infty$ is smooth for $t \in (-\infty,\infty)$ and has asymptotics of (1.16)-(1.17) at $t=-\infty$.

In fact, the region R_{smooth} is very large. The numerical results for R_{smooth} are shown in Figure 1.




Figure 1. R_{smooth} and $R_{singular}$. R_{smooth} is the light green region. If (c_1, c_2) belongs to R_{smooth} , the solution defined at $t = \infty$ by this (c_1, c_2) is smooth on $(-\infty, \infty)$. Else if (c_1, c_2) belongs to $R_{singular}$ (the light yellow region), the corresponding solution must have singularity at some finite $t = t_0$. The red curve is the boundary between R_{smooth} and $R_{singular}$. P_c is a special point on the boundary curve: the boundary curve becomes straight on the right of P_c .

Theorem 1.2 gives positive answers to the open questions (1) and (2) of [11]. But the non-uniqueness of $q_2 = q_2(c_1, c_2; t)$ causes the non-uniqueness of $F_6(t)$. In fact, by formula [11]

(1.22)
$$F_6\left(\frac{t}{3^{2/3}}\right) = \frac{q_2 - 1}{2q_2} \exp\left(-\frac{1}{3} \int_{\infty}^t \omega(s) ds + \frac{2}{3} \int_{\infty}^t \frac{u'(s)}{u(s)} \frac{1 + q_2(s)}{q_2(s)} ds\right),$$

where $\omega(s) = u(s)^4 + su(s)^2 - u'(s)^2$, we can verify that $F_6(t)$ is indeed dependent on c_1 and c_2 . So we have to determine the values of c_1 and c_2 to guarantee there is only a unique $F_6(t)$. To determine c_1 and c_2 , we rely on Theorem 1.1. Grava et al. [11] have formulated $F(\beta = 6; x, t)$ from q_2 and α as

$$(1.23) \quad F\left(\beta = 6; \frac{x}{3^{1/3}}, \frac{t}{3^{2/3}}\right) = \kappa u^{\frac{1}{2}} \left[u^{-1} \left(\frac{1 + q_2}{2} x - \alpha \right) Y_{12}^{(6)}(x, t) + Y_{22}^{(6)}(x, t) \right], \quad x \geqslant 0,$$

$$(1.24) \quad F\left(\beta = 6; \frac{x}{3^{1/3}}, \frac{t}{3^{2/3}}\right) = -\kappa u^{\frac{1}{2}} e^{\frac{x^3}{3} - xt} \left[u^{-1} \left(\frac{1 + q_2}{2} x - \alpha \right) Y_{11}^{(3)}(x, t) + Y_{21}^{(3)}(x, t) \right], \quad x \leq 0,$$

where $Y^{(3)}$ and $Y^{(6)}$ are 2×2 matrices of the wave function of the Painlevé II. By (1.23)-(1.24), $F(\beta =$ (6; x, t) contains parameters c_1 and c_2 . By (1.18)-(1.19) and (1.23)-(1.24), we can prove $c_1 = 0$ is enough to guarantee the boundary condition (1.7). But if $c_2 \neq 0$, $F\left(\beta = 6; \frac{x}{3^{1/3}}, \frac{t}{3^{2/3}}\right)$ will grow exponentially near the line $x = -\sqrt{t}$ for $t \to \infty$.

The second main result of this paper is the following:

Theorem 1.3. If and only if $c_1 = c_2 = 0$, the resulting $F(\beta = 6; x, t)$ is bounded at the boundary $x^2 + t^2 = \infty.$

Now all requirements of Theorem 1.1 are satisfied: $F(\beta = 6; x, t)$ given by (1.23)-(1.24) satisfies the Bloemendal-Virág equation (1.6); $c_1 = 0$ guarantees it satisfies the boundary condition (1.7); Theorem 1.3 guarantees it is a bounded solution. So $F(\beta = 6; x, t)$ is indeed given by (1.23) and (1.24) with $c_1 = c_2 = 0$. We also note that $F_6(t)$ is given by (1.22) with the hidden parameters $c_1 = c_2 = 0$.

2 Derivation of the ODEs of q_2 , α and κ

The Flaschka-Newell Lax pair of Painlevé II is [9]

(2.1)
$$\frac{d\psi_0}{dx} = \hat{L}_0\psi_0,$$
(2.2)
$$\frac{d\psi_0}{dt} = \hat{B}_0\psi_0$$

$$\frac{d\psi_0}{dt} = \hat{B}_0 \psi_0$$

where

(2.3)
$$\psi_0 = \begin{pmatrix} \mathcal{F}_0(x,t) \\ \mathcal{G}_0(x,t) \end{pmatrix},$$

(2.4)
$$\hat{L}_0 = \frac{x^2}{2}\sigma_3 + x \begin{pmatrix} 0 & u(t) \\ u(t) & 0 \end{pmatrix} + \begin{pmatrix} -\frac{t}{2} - u(t)^2 & -u'(t) \\ u'(t) & \frac{t}{2} + u(t)^2 \end{pmatrix},$$

(2.5)
$$\hat{B}_0 = -\frac{x}{2}\sigma_3 - \begin{pmatrix} 0 & u(t) \\ u(t) & 0 \end{pmatrix}.$$

By (1.8) and (1.12), we get

$$(2.6) \mathcal{F}(x,t) = \frac{1}{2}e^{\frac{x^3}{6} - \frac{tx}{2}}(q_2(t)x + x - 2\alpha(t))\kappa(t)\psi(t)\mathcal{F}_0(x,t) - e^{\frac{x^3}{6} - \frac{tx}{2}}\frac{\kappa(t)}{\psi(t)}\mathcal{G}_0(x,t).$$

Grava et. al suggested

$$\psi = -\frac{\mathrm{i}}{\sqrt{u}}.$$

Substituting (2.6) and (2.7) into (1.9) with $\beta = 6$, we immediately obtain (1.13)-(1.14) and

(2.8)
$$\frac{\kappa'}{\kappa} = -\frac{2}{3}\alpha - \frac{1}{3}(t+u^2)u^2 + \frac{u'}{6u}(2q_2 - 1) + \frac{1}{3}(u')^2.$$

By requiring $\mathcal{F}(x,t) \xrightarrow{x \to \infty, t \to \infty} 1$, Grava et. al [11] proved (1.15) and

(2.9)
$$\kappa u^{\frac{1}{2}} \xrightarrow{t \to \infty} 1.$$

Equation (2.8) with boundary condition (2.9) determine κ completely if a smooth solution of (α, q_2) has been obtained under the boundary condition (1.15), more precisely, the asymptotics of (1.18)-(1.19).

The expansion of $\kappa(t)$ at $t = -\infty$ can also be obtained. By (2.8), (1.16)-(1.17) and the asymptotics of Hastings-McLeod solution u, the asymptotics of $\frac{\kappa'}{\kappa}$ at $t = -\infty$ is obtained as

$$\frac{\kappa'(t)}{\kappa(t)} = \frac{t^2}{12} - \frac{\sqrt{2}}{3}(-t)^{\frac{1}{2}} + \frac{5}{24}(-t)^{-1} + \frac{7}{32\sqrt{2}}(-t)^{-\frac{5}{2}} + \cdots$$

Therefore

(2.10)
$$\kappa(t) = C_{\kappa} \times e^{-\frac{1}{36}(-t)^3 + \frac{2\sqrt{2}}{9}(-t)^{3/2} - \frac{5}{24}\ln(-t) + \frac{7}{48\sqrt{2}}(-t)^{-3/2} + \cdots}$$

Assuming (1.3), we get

(2.11)
$$\ln C_{\kappa} = c_{\beta=6} - \frac{\ln 3}{36} + \frac{5}{4} \ln 2.$$

For $\beta = 6$, Borot et al. [3] was able to simplify (1.4) to

(2.12)
$$c_{\beta=6} = -\frac{97}{72} \ln 2 - \frac{7}{36} \ln 3 - \frac{\ln(2\pi)}{6} + \frac{\ln\Gamma(\frac{1}{3})}{3} + \frac{\zeta'(-1)}{3}.$$

So we have

(2.13)
$$\ln C_{\kappa} = -\frac{7}{72} \ln 2 - \frac{2}{9} \ln 3 - \frac{\ln(2\pi)}{6} + \frac{\ln \Gamma(\frac{1}{3})}{3} + \frac{\zeta'(-1)}{3}.$$

The value of $\ln C_{\kappa}$ can be obtained by numerical experiments similar to the ones in Section 7 with $c_1 = c_2 = 0$. Our numerical experiments give

$$(2.14) \ln C_{\kappa} = -0.3445050500286934815501994065702518 \cdots$$

In fact, $\ln C_{\kappa}$ from our numerical experiments coincides with (2.13) for more than 100 digits, which gives a numerical verification of (1.3) and (1.4) for $\beta = 6$.

Altogether, the algorithm is as following. First give the ansatz for $\mathcal{F}(x,t)$ as (2.6); then by (1.9) obtain the ODEs for the unknowns; next by the boundary condition for $\mathcal{F}(x,t)$ get all boundary conditions for the unknowns, which should determine all unknowns uniquely; at last prove the obtained $\mathcal{F}(x,t)$ satisfies all the requirements for it.

3 Asymptotics of q_2 and α at $t = -\infty$

In this section we will show by linearization analysis that the asymptotics (1.16)-(1.17) are not the asymptotics of a specific solution of (1.13)-(1.14), but of a general solution of (1.13)-(1.14). More detailed analysis of these asymptotics will be given in Section 6.

Suppose $(q_{20}(t), \alpha_0(t))$ is a smooth solution of (1.13)-(1.14) with asymptotics of (1.16)-(1.17). Let $(q_2(t), \alpha(t))$ be a solution of (1.13)-(1.14) near $(q_{20}(t), \alpha_0(t))$. Then $(q_2(t), \alpha(t))$ can be expressed as

$$q_2(t) = q_{20}(t) + \epsilon \mathcal{Q}(t),$$

$$\alpha(t) = \alpha_0(t) + \epsilon \mathcal{Q}(t),$$

where $\epsilon \to 0$ is infinitesimal.

So $(\mathcal{Q}, \mathcal{A})$ satisfies the ODE

$$\mathscr{Q}'(t) = \left(\frac{2}{3}\alpha_0 + \frac{u'}{u}\frac{1 - 2q_{20}}{3}\right)\mathscr{Q} + \frac{2}{3}q_{20}\mathscr{A},$$

$$\mathscr{A}'(t) = -\frac{1}{3}\left(\frac{u'}{u}\alpha_0 + \frac{t}{2} + u^2\right)\mathscr{Q} + \left(\frac{4}{3}\alpha_0 + \frac{u'}{u}\frac{2 - q_{20}}{3}\right)\mathscr{A}.$$

At $t = -\infty$, the expansions of u, $\frac{u'}{u}$, α_0 and q_{20} are known

$$u = \sqrt{\frac{-t}{2}} + \cdots, \quad \frac{u'}{u} = \frac{1}{2t} + \cdots, \quad q_{20} = \frac{1}{\sqrt{2}} (-t)^{-\frac{3}{2}} + \cdots, \quad \alpha_0 = \sqrt{\frac{-t}{2}} + \cdots.$$

Therefore, \mathcal{Q} and \mathcal{A} satisfy

$$\mathscr{Q}'(t) = \left(\frac{\sqrt{-2t}}{3} + \cdots\right) \mathscr{Q} + \left(\frac{\sqrt{2}}{3}(-t)^{-\frac{3}{2}} + \cdots\right) \mathscr{A},$$
$$\mathscr{A}'(t) = \left(\frac{1}{6}(-2t)^{-\frac{1}{2}} + \cdots\right) \mathscr{Q} + \left(\frac{2\sqrt{-2t}}{3} + \cdots\right) \mathscr{A}.$$

Now it is clear that the solution (q_2, α) is exponentially close to the solution (q_{20}, α_0) in an order of $e^{-\frac{2\sqrt{2}}{9}(-t)^{3/2}}$.

So we reach the following result.

Theorem 3.1. If a solution $(q_{20}(t), \alpha_0(t))$ is nonsingular on $(-\infty, t_0]$ and has asymptotics (1.16)-(1.17), then the general solutions that are close to $(q_{20}(t), \alpha_0(t))$ are all non-singular on $(-\infty, t_0]$ and also have asymptotics (1.16)-(1.17) at $t = -\infty$.

4 The linear variables and the integral equations

At $t = \infty$, it is convenient to work with α and

$$\tilde{q}_2 = 1 + q_2.$$

Also it is helpful to remember

(4.1)
$$u(t) = \operatorname{Ai}(t) + \left(\frac{1}{32\pi^{3/2}}t^{-\frac{7}{4}} + \cdots\right)e^{-\frac{6}{3}t^{3/2}} + \cdots$$

The ODEs for \tilde{q}_2 and α are

(4.2)
$$\tilde{q}_2'(t) = \frac{2}{3}(\tilde{q}_2 - 1)\alpha + \frac{u'}{u}\frac{3 - \tilde{q}_2}{3}\tilde{q}_2,$$

(4.3)
$$\alpha'(t) = \left(\frac{2}{3}\alpha + \frac{u'}{u}\frac{3-\tilde{q}_2}{3}\right)\alpha - \frac{t}{6}\tilde{q}_2 - \frac{2+\tilde{q}_2}{3}u^2.$$

Equations (4.2)-(4.3) are linearized by

(4.4)
$$\tilde{q}_2(t) = \frac{\phi_1(t)}{\phi_0(t)} u(t), \quad \alpha(t) = \frac{\phi_2(t)}{\phi_0(t)} u(t),$$

where ϕ_1 , ϕ_2 and ϕ_0 satisfy

(4.5)
$$\phi_1'(t) = -\frac{2}{3}\phi_2(t),$$

(4.6)
$$\phi_2'(t) = -\frac{2}{3}u(t)\phi_0(t) - \frac{1}{6}(t + 2u(t)^2)\phi_1(t),$$

(4.7)
$$\phi_0'(t) = \frac{1}{3}u'(t)\phi_1(t) - \frac{2}{3}u(t)\phi_2(t).$$

To analyze (4.5)-(4.7), we need the following estimations.

Proposition 4.1. There exists t_0^P , such that for $t \ge t_0^P$:

- $|\text{Bi}(t)| < e^{\frac{2}{3}t^{3/2}}$ and $|\text{Bi}'(t)| < e^{\frac{2}{3}t^{3/2}}\sqrt{t}$;
- $|\operatorname{Ai}(t)| < e^{-\frac{2}{3}t^{3/2}}$ and $|\operatorname{Ai}'(t)| < e^{-\frac{2}{3}t^{3/2}}\sqrt{t}$;
- $\bullet \ |u(t)| < e^{-\frac{2}{3}t^{3/2}} \ and \ |u'(t)| < e^{-\frac{2}{3}t^{3/2}} \sqrt{t}.$

The proof is routine. Thus we omit it.

Remark 4.2. For example, we can take $t_0^P = 1$. By numerical results, $t_0^P = 1$ satisfies all requirements for t_0^P in Proposition 4.1. We assume t_0^P is sufficiently large. Also, the actual value of t_0^P is never needed in the following proofs.

4.1 The three independent solutions defined near $t = \infty$

For convenience, denote

$$g_1(t) = \text{Bi}(\frac{t}{3^{2/3}}), \quad g_2(t) = \text{Ai}(\frac{t}{3^{2/3}}), \quad g_0(t) = 0.$$

We will show, for k = 1, 2, 0, the following integral equations

$$\Phi_{1k}(t) = g_{k}(t) + \frac{2\pi}{3^{1/3}} \left[\operatorname{Ai}(\frac{t}{3^{2/3}}) \int_{\infty}^{t} \operatorname{Bi}(\frac{s}{3^{2/3}}) \left(-\frac{2}{3}u(s)\Phi_{0k}(s) - \frac{1}{3}u(s)^{2}\Phi_{1k}(s) \right) ds \right]
(4.8) - \operatorname{Bi}(\frac{t}{3^{2/3}}) \int_{\infty}^{t} \operatorname{Ai}(\frac{s}{3^{2/3}}) \left(-\frac{2}{3}u(s)\Phi_{0k}(s) - \frac{1}{3}u(s)^{2}\Phi_{1k}(s) \right) ds \right] ,
\Phi_{2k}(t) = -\frac{3}{2}g'_{k}(t) - \pi \left[\operatorname{Ai}'(\frac{t}{3^{2/3}}) \int_{\infty}^{t} \operatorname{Bi}(\frac{s}{3^{2/3}}) \left(-\frac{2}{3}u(s)\Phi_{0k}(s) - \frac{1}{3}u(s)^{2}\Phi_{1k}(s) \right) ds \right] ,
(4.9) - \operatorname{Bi}'(\frac{t}{3^{2/3}}) \int_{\infty}^{t} \operatorname{Ai}(\frac{s}{3^{2/3}}) \left(-\frac{2}{3}u(s)\Phi_{0k}(s) - \frac{1}{3}u(s)^{2}\Phi_{1k}(s) \right) ds \right] ,
(4.10) - \Phi_{0k}(t) = \delta_{k}^{0} + \int_{\infty}^{t} \left(\frac{1}{3}u'(s)\Phi_{1k}(s) - \frac{2}{3}u(s)\Phi_{2k}(s) \right) ds .$$

define three independent solutions for (4.5)-(4.7), i.e., $(\phi_1(t), \phi_2(t), \phi_0(t)) = (\Phi_{1k}(t), \Phi_{2k}(t), \Phi_{0k}(t)), k = 1, 2, 0$, solves (4.5)-(4.7).

For k = 1, 2, 0, define

(4.11)
$$\Phi_{1k}^{(0)}(t) = (1 - \delta_k^0)g_k(t),$$

(4.12)
$$\Phi_{2k}^{(0)}(t) = -\frac{3}{2}(1 - \delta_k^0)g_k'(t),$$

(4.13)
$$\Phi_{0k}^{(0)}(t) = \delta_k^0 + \int_0^t \left(\frac{1}{3}u'(s)\Phi_{1k}^{(0)}(s) - \frac{2}{3}u(s)\Phi_{2k}^{(0)}(s)\right)ds,$$

where $\delta_k^0 = \begin{cases} 1, & k = 0 \\ 0, & k \neq 0 \end{cases}$.

$$\begin{split} \Phi_{1k}^{(j+1)}(t) &= g_k(t) + \frac{2\pi}{3^{1/3}} \left[\operatorname{Ai}(\frac{t}{3^{2/3}}) \int_{\infty}^{t} \operatorname{Bi}(\frac{s}{3^{2/3}}) \left(-\frac{2}{3}u(s)\Phi_{0k}^{(j)}(s) - \frac{1}{3}u(s)^{2}\Phi_{1k}^{(j)}(s) \right) ds \\ (4.14) &\qquad -\operatorname{Bi}(\frac{t}{3^{2/3}}) \int_{\infty}^{t} \operatorname{Ai}(\frac{s}{3^{2/3}}) \left(-\frac{2}{3}u(s)\Phi_{0k}^{(j)}(s) - \frac{1}{3}u(s)^{2}\Phi_{1k}^{(j)}(s) \right) ds \right], \\ \Phi_{2k}^{(j+1)}(t) &= -\frac{3}{2}g_k'(t) - \pi \left[\operatorname{Ai}'(\frac{t}{3^{2/3}}) \int_{\infty}^{t} \operatorname{Bi}(\frac{s}{3^{2/3}}) \left(-\frac{2}{3}u(s)\Phi_{0k}^{(j)}(s) - \frac{1}{3}u(s)^{2}\Phi_{1k}^{(j)}(s) \right) ds \\ (4.15) &\qquad -\operatorname{Bi}'(\frac{t}{3^{2/3}}) \int_{\infty}^{t} \operatorname{Ai}(\frac{s}{3^{2/3}}) \left(-\frac{2}{3}u(s)\Phi_{0k}^{(j)}(s) - \frac{1}{3}u(s)^{2}\Phi_{1k}^{(j)}(s) \right) ds \right], \\ (4.16) &\Phi_{0k}^{(j+1)}(t) &= \delta_k^0 + \int_{\infty}^{t} \left(\frac{1}{3}u'(s)\Phi_{1k}^{(j+1)}(s) - \frac{2}{3}u(s)\Phi_{2k}^{(j+1)}(s) \right) ds. \end{split}$$

Proposition 4.3. For $t \ge t_0^P$,

$$\begin{split} |\Phi_{1k}^{(j+1)}(t) - \Phi_{1k}^{(j)}(t)| &< C_{1k}^{(j+1)} e^{-(\frac{4}{3}j + \frac{4}{9}k + \frac{2}{3})t^{3/2}}, \quad |\Phi_{2k}^{(j+1)}(t) - \Phi_{2k}^{(j)}(t)| < C_{2k}^{(j+1)} t^{\frac{1}{2}} e^{-(\frac{4}{3}j + \frac{4}{9}k + \frac{2}{3})t^{3/2}}, \\ |\Phi_{0k}^{(j+1)}(t) - \Phi_{0k}^{(j)}(t)| &< C_{0k}^{(j+1)} e^{-(\frac{4}{3}j + \frac{4}{9}k + \frac{4}{3})t^{3/2}}, \end{split}$$

where

$$\begin{split} C_{11}^{(j)} &= 2C_{21}^{(j)} = \pi^{j} 3^{\frac{-j}{3}} \prod_{i=1}^{j} \frac{6i-1}{(3i-1)(3i-2)}, \quad C_{01}^{(j)} = \pi^{j} 3^{\frac{-j}{3}} \prod_{i=1}^{j} \frac{6i-1}{(3i-1)(3i+1)}, \\ C_{12}^{(j)} &= 2C_{22}^{(j)} = \pi^{j} 3^{\frac{-j}{3}} \prod_{i=1}^{j} \frac{6i+1}{3i(3i-1)}, \quad C_{02}^{(j)} &= \frac{1}{2} \pi^{j} 3^{\frac{-j}{3}} \prod_{i=1}^{j} \frac{6i+1}{3i(3i+2)}, \\ C_{10}^{(j)} &= 2C_{20}^{(j)} &= 3\pi^{j} 3^{\frac{-j}{3}} \frac{1}{(j-1)!} \prod_{i=1}^{j} \frac{2i-1}{3i-2}, \quad C_{00}^{(j)} &= \pi^{j} 3^{\frac{-j}{3}} \frac{1}{j!} \prod_{i=1}^{j} \frac{2i-1}{3i-2}. \end{split}$$

Theorem 4.4. $(\Phi_{1k}^{(j)}, \Phi_{2k}^{(j)}, \Phi_{0k}^{(j)}), k = 1, 2, 0, defined by (4.11)-(4.16) converge to the solutions of (4.5)-(4.7) for <math>t \ge t_0^P$.

5 Asymptotics of q_2 and α at $t = \infty$

At $t = \infty$, it is straightforward to verify

(5.1)
$$\Phi_{11}(t) = \text{Bi}(\frac{t}{3^{2/3}}) + O(e^{-\frac{10}{9}t^{3/2}}),$$

(5.2)
$$\Phi_{21}(t) = -\frac{3^{1/3}}{2} \operatorname{Bi}'(\frac{t}{3^{2/3}}) + O(\sqrt{t}e^{-\frac{10}{9}t^{3/2}}),$$

(5.3)
$$\Phi_{01}(t) = \int_{\infty}^{t} \left(\frac{1}{3} \operatorname{Ai}'(s) \operatorname{Bi}(\frac{s}{3^{2/3}}) + \frac{1}{3^{2/3}} \operatorname{Ai}(s) \operatorname{Bi}'(\frac{s}{3^{2/3}}) \right) ds + O(e^{-\frac{16}{9}t^{3/2}}),$$

(5.4)
$$\Phi_{12}(t) = \operatorname{Ai}(\frac{t}{3^{2/3}}) + O(e^{-\frac{14}{9}t^{3/2}}),$$

(5.5)
$$\Phi_{22}(t) = -\frac{3^{1/3}}{2} \operatorname{Ai}'(\frac{t}{3^{2/3}}) + O(\sqrt{t}e^{-\frac{14}{9}t^{3/2}}),$$

(5.6)
$$\Phi_{02}(t) = \int_{\infty}^{t} \left(\frac{1}{3} \operatorname{Ai}'(s) \operatorname{Ai}(\frac{s}{3^{2/3}}) + \frac{1}{3^{2/3}} \operatorname{Ai}(s) \operatorname{Ai}'(\frac{s}{3^{2/3}}) \right) ds + O(e^{-\frac{20}{9}t^{3/2}}),$$

$$(5.7) \quad \Phi_{10}(t) = -\frac{4\pi}{3^{4/3}} \left(\operatorname{Ai}(\frac{t}{3^{2/3}}) \int_{\infty}^{t} \operatorname{Bi}(\frac{s}{3^{2/3}}) \operatorname{Ai}(s) ds - \operatorname{Bi}(\frac{t}{3^{2/3}}) \int_{\infty}^{t} \operatorname{Ai}(\frac{s}{3^{2/3}}) \operatorname{Ai}(s) ds \right) + O(e^{-2t^{3/2}})$$

$$(5.8) \quad \Phi_{20}(t) = \frac{2\pi}{3} \left(\operatorname{Ai}'(\frac{t}{3^{2/3}}) \int_{0}^{t} \operatorname{Bi}(\frac{s}{3^{2/3}}) \operatorname{Ai}(s) ds - \operatorname{Bi}'(\frac{t}{3^{2/3}}) \int_{0}^{t} \operatorname{Ai}(\frac{s}{3^{2/3}}) \operatorname{Ai}(s) ds \right) + O(\sqrt{t}e^{-2t^{3/2}}),$$

$$(5.9) \quad \Phi_{00}(t) = 1 + \int_{\infty}^{t} \left(\frac{1}{3} \operatorname{Ai}'(s) \tilde{\Phi}_{10}^{(1)}(s) - \frac{2}{3} \operatorname{Ai}(s) \tilde{\Phi}_{20}^{(1)}(s) \right) ds + O(e^{-\frac{8}{3}t^{3/2}}),$$

where $\tilde{\Phi}_{10}^{(1)}$ and $\tilde{\Phi}_{20}^{(1)}$ are defined by dropping the error terms of (5.7) and (5.8) respectively.

Lemma 5.1. The asymptotics of a solution of (4.2)-(4.3) at $t = \infty$ must belong to one of the following three classes.

Class A:
$$\tilde{q}_2(t) \xrightarrow{t \to \infty} 0$$
 and $\alpha(t) \xrightarrow{t \to \infty} 0$.

(5.10)
$$\tilde{q}_2(t) = \left(c_1 \operatorname{Bi}(\frac{t}{3^{2/3}}) + c_2 \operatorname{Ai}(\frac{t}{3^{2/3}}) + c_1^2 \tilde{M}_2(t) e^{-\frac{8}{9}t^{3/2}}\right) \operatorname{Ai}(t) + o\left(e^{-\frac{4}{3}t^{3/2}}\right),$$

(5.11)
$$\alpha(t) = -\frac{3^{1/3}}{2} \left(c_1 \operatorname{Bi}'(\frac{t}{3^{2/3}}) + c_2 \operatorname{Ai}'(\frac{t}{3^{2/3}}) + c_1^2 \tilde{N}_2(t) e^{-\frac{8}{9}t^{3/2}} \right) \operatorname{Ai}(t) + o\left(e^{-\frac{4}{3}t^{3/2}}\right),$$

where

$$\tilde{M}_{2}(t) = -\text{Bi}(\frac{t}{3^{2/3}}) \int_{\infty}^{t} \left(\frac{1}{3}\text{Ai}'(s)\text{Bi}(\frac{s}{3^{2/3}}) + \frac{1}{3^{2/3}}\text{Ai}(s)\text{Bi}'(\frac{s}{3^{2/3}})\right) ds,
\tilde{N}_{2}(t) = -\text{Bi}'(\frac{t}{3^{2/3}}) \int_{\infty}^{t} \left(\frac{1}{3}\text{Ai}'(s)\text{Bi}(\frac{s}{3^{2/3}}) + \frac{1}{3^{2/3}}\text{Ai}(s)\text{Bi}'(\frac{s}{3^{2/3}})\right) ds.$$

Class B: $\tilde{q}_2(t) \xrightarrow{t \to \infty} \infty$ and $\alpha(t) \xrightarrow{t \to \infty} -\infty$

$$\begin{split} \tilde{q}(t) &= \left[\mathrm{Bi}(\frac{t}{3^{2/3}}) + c_2 \left(\mathrm{Ai}(\frac{t}{3^{2/3}}) - \mathrm{Bi}(\frac{t}{3^{2/3}}) \frac{\Phi_{02}^{(0)}(t)}{\Phi_{01}^{(0)}(t)} \right) \right] \frac{\mathrm{Ai}(t)}{\Phi_{01}^{(0)}(t)} + o\left(e^{-\frac{8}{9}t^{3/2}}\right), \\ \alpha(t) &= -\frac{3^{1/3}}{2} \left[\mathrm{Bi}'(\frac{t}{3^{2/3}}) + c_2 \left(\mathrm{Ai}'(\frac{t}{3^{2/3}}) - \mathrm{Bi}'(\frac{t}{3^{2/3}}) \frac{\Phi_{02}^{(0)}(t)}{\Phi_{01}^{(0)}(t)} \right) \right] \frac{\mathrm{Ai}(t)}{\Phi_{01}^{(0)}(t)} + o\left(e^{-\frac{8}{9}t^{3/2}}\right). \end{split}$$

Class C:
$$\tilde{q}(t) \xrightarrow{t \to \infty} 2$$
 and $\alpha(t) \xrightarrow{t \to \infty} \sqrt{t}$.

Proof. The general solution of (4.5)-(4.7) is $(\phi_1, \phi_2, \phi_0) = (c_1\Phi_{11} + c_2\Phi_{12} + c_0\Phi_{10}, c_1\Phi_{21} + c_2\Phi_{22} + c_0\Phi_{20}, c_1\Phi_{01} + c_2\Phi_{02} + c_0\Phi_{00})$. So $\tilde{q}_2 = \frac{c_1\Phi_{11} + c_2\Phi_{12} + c_0\Phi_{10}}{c_1\Phi_{01} + c_2\Phi_{02} + c_0\Phi_{00}} u$ and $\alpha = \frac{c_1\Phi_{21} + c_2\Phi_{22} + c_0\Phi_{20}}{c_1\Phi_{01} + c_2\Phi_{02} + c_0\Phi_{00}} u$. If $c_0 \neq 0$, c_0 can be taken as 1. By (5.1)-(5.9), we have $\tilde{q}_2 = (c_1\Phi_{11} + c_2\Phi_{12} + o(e^{-\frac{2}{3}t^{3/2}}))(1 - c_1\Phi_{01} + o(e^{-\frac{8}{9}t^{3/2}}))u = (c_1\Phi_{11} + c_2\Phi_{12} - c_1^2\Phi_{11}\Phi_{01} + o(e^{-\frac{2}{3}t^{3/2}}))u$. Also considering $u(t) = \mathrm{Ai}(t) + o(e^{-2t^{3/2}})$ and (5.1)-(5.9), we finally get (5.10). We can verify directly $\tilde{M}_2(t) = o(\mathrm{Ai}(\frac{t}{3^{2/3}}))$. Similarly, (5.11) is obtained. Therefore, the Class A describes the asymptotics of $\tilde{q}_2(t)$ and $\alpha(t)$ at $t = \infty$ for $c_0 \neq 0$. If $c_0 = 0$ and $c_1 \neq 0$, c_1 can be taken as 1. Then we can prove the asymptotics belong to Class B in this case. At last, if $c_0 = c_1 = 0$, c_2 can be taken as 1. Class C describes the asymptotics of this case.

Proposition 5.2. A solution of (1.13)-(1.14), which has property (1.15), must have asymptotics (1.18)-(1.21).

Proof. Let (ϕ_1, ϕ_2, ϕ_0) and (c_1, c_2, c_0) be the ones defined in the proof of Lemma 5.1. By (1.15) and $\tilde{q}_2 = q_2 + 1$, we know $\tilde{q}_2(t) \xrightarrow{t \to \infty} 0$ and $\alpha(t) \xrightarrow{t \to \infty} 0$. This is the Class A case. The only remaining problem is to verify (1.20)-(1.21) for $c_1 = c_2 = 0$. It is straightforward to verify this by (5.7)-(5.9).

Remark 5.3. The error terms of (1.20)-(1.21) are not optimal. In fact, the error terms can be shown to be $O\left(t^{-\frac{5}{2}}e^{-\frac{8}{3}t^{3/2}}\right)$ and $O\left(t^{-2}e^{-\frac{8}{3}t^{3/2}}\right)$ respectively by a tedious calculation from (4.11)-(4.16).

6 Proof of Theorem 1.2

Proposition 6.1. There exists a minimal $k_0 > 0$ so as to $t + k_0 u(t)^2 \ge 0$ for all t.

Proof. It is obvious $k_0 > 0$. Let $f(t) = t + 2u(t)^2$. f(t) has minimum since f(0) > 0, $f(-\infty) = 0$ and f(t) < 0 for large negative enough t. Since $u(t) \neq 0$ for $t < \infty$, there exists k such that $t + 2u(t)^2 + ku(t)^2 \ge 0$ for all t. Obviously, such k has minimum.

Remark 6.2. By the preceding proof, we know $k_0 > 2$. To calculate k_0 numerically, we use $k_0 = -\min\left(\frac{t}{u^2}\right)$, by which k_0 is calculated up to more than 100 digits. Though k_0 is so accurately known, its "closed form" is still unknown. $k_0 \approx 2.1228589561253469$ is achieved at $t \approx -1.188111911480737877$. In the following, $k_0 < \frac{10}{3}$ is needed. The proof of $k_0 < \frac{10}{3}$ is somewhat technical and digressed, so we put it in Appendix A.

A visualized estimation of k_0 is given by Figure 2.




Figure 2. Estimate k_0 by the graph of $\frac{t}{u^2}$. The green curve is the plot of $\frac{t}{u^2}$ and the horizontal red line is y=-2.2. By the graph, it is obvious min $\left(\frac{t}{u^2}\right)>-2.2$, i.e., $k_0<2.2$.

Lemma 6.3. If a solution of (4.5)-(4.7) has properties $\phi_1(t_0) > 0$, $\phi_2(t_0) > 0$ and $\phi_0(t_0) > 0$ and $\frac{2}{3}\phi_0(t_0) - \frac{k_0 - 2}{6}u(t_0)\phi_1(t_0) > 0$, then

(1) $\phi_1(t)$, $\phi_2(t)$, $\phi_0(t)$ and $\frac{2}{3}\phi_0(t) - \frac{k_0-2}{6}u(t)\phi_1(t)$ are all monotonic decreasing on $(-\infty, t_0]$;

(2)
$$\phi_1(t) \xrightarrow{t \to -\infty} \infty$$
, $\phi_2(t) \xrightarrow{t \to -\infty} \infty$ and $\phi_0(t) \xrightarrow{t \to -\infty} \infty$.

(3)

(6.1)
$$\lim_{t \to -\infty} \frac{\phi_1(t)}{\phi_0(t)} u(t) = 1, \quad \lim_{t \to -\infty} \frac{\phi_2(t)}{\phi_0(t)} = 1.$$

Correspondingly, near $t = -\infty$,

(6.2)
$$q(t) = o(1), \quad \alpha(t) = \sqrt{\frac{-t}{2}} + o((-t)^{\frac{1}{2}}).$$

Proof. (1)
$$\phi_1'(t_0) = -\frac{2}{3}\phi_2(t_0) < 0$$
.
 $\phi_2'(t_0) = -\frac{2}{3}u(t_0)\phi_0(t_0) - \frac{1}{6}(t_0 + k_0u(t_0)^2)\phi_1(t_0) + \frac{1}{6}(k_0 - 2)u(t_0)^2\phi_1(t_0) < 0$.
 $\phi_0'(t_0) = \frac{1}{3}u'(t_0)\phi_1(t_0) - \frac{2}{3}u(t_0)\phi_2(t_0) < 0$.

$$\begin{split} &\frac{d}{dt}\left(\frac{2}{3}\phi_0(t_0) - \frac{k_0 - 2}{6}u(t)\phi_1(t_0)\right) \\ &= &\frac{2}{3}\left(\frac{1}{3}u'(t)\phi_1(t_0) - \frac{2}{3}u(t_0)\phi_2(t_0)\right) - \frac{k_0 - 2}{6}u'(t_0)\phi_1(t_0) - \frac{k_0 - 2}{6}u(t_0)\left(-\frac{2}{3}\phi_2(t_0)\right) \\ &= &\frac{10 - 3k_0}{18}u'(t_0)\phi_1(t_0) - \frac{6 - k_0}{9}u(t_0)\phi_2(t_0). \end{split}$$

By Remark 6.2, $k_0 < \frac{10}{3}$. Therefore, $\frac{d}{dt} \left(\frac{2}{3} \varphi_0(t_0) - \frac{k_0 - 2}{6} u(t_0) \varphi_1(t_0) \right) < 0$. So we have $\phi_1(t_0 - \epsilon) > \phi_1(t_0) > 0$, $\phi_2(t_0 - \epsilon) > \phi_2(t_0) > 0$, $\phi_0(t_0 - \epsilon) > \phi_3(t_0) > 0$ and $\frac{2}{3} \phi_0(t) - \frac{k_0 - 2}{6} u(t) \phi_1(t)|_{t=t_0 - \epsilon} > \frac{2}{3} \phi_0(t) - \frac{k_0 - 2}{6} u(t) \phi_1(t)|_{t=t_0} > 0$. This process can be repeated endlessly. So the first statement of the lemma is proved.

- (2) By the preceding proof, $\phi_1(t) > 0$, $\phi_2(t) > 0$, $\phi_1(t) = -\frac{2}{3}\phi_2(t) < 0$ and $\phi_2'(t) < -\frac{1}{6}(t + k_0u(t)^2)\phi_1(t) \le 0$. So both $\phi_1(t)$ and $\phi_2(t)$ grow exponentially to infinity as $t \to -\infty$. By $\phi_0'(t) = \frac{1}{3}u'(t)\phi_1(t) \frac{2}{3}u(t)\phi_2(t)$, $\phi_0(t) \xrightarrow{t \to -\infty} \infty$ is got.

 (3) Let $x_1 = \lim_{t \to -\infty} \frac{\phi_1(t)}{\phi_0(t)}u(t)$ and $x_2 = \lim_{t \to -\infty} \frac{\phi_2(t)}{\phi_0(t)}$. We apply L'Hospital's rule to obtain the values
- (3) Let $x_1 = \lim_{t \to -\infty} \frac{\phi_1(t)}{\phi_0(t)} u(t)$ and $x_2 = \lim_{t \to -\infty} \frac{\phi_2(t)}{\phi_0(t)}$. We apply L'Hospital's rule to obtain the values of x_1 and x_2 . It is legal since $\phi_1(t)u(t) \xrightarrow{t \to -\infty} \infty$, $\phi_2(t) \xrightarrow{t \to -\infty} \infty$, $\phi_0(t) \xrightarrow{t \to -\infty} \infty$ and $\phi_0'(t) < 0$ for $t < t_0$. Then by L'Hospital's rule and (4.5)-(4.7), we obtain the algebraic equations for x_1 and x_2

(6.3)
$$x_1 = \frac{-\frac{2}{3}x_2u(t) + \frac{u'(t)}{u(t)}x_1}{\frac{1}{3}\frac{u'(t)}{u(t)}x_1 - \frac{2}{3}u(t)x_2},$$

(6.4)
$$x_2 = \frac{-\frac{1}{6} \frac{t + 2u(t)^2}{u(t)} x_1 - \frac{2}{3} u(t)}{\frac{1}{3} \frac{u'(t)}{u(t)} x_1 - \frac{2}{3} u(t) x_2}.$$

Note t in (6.3)-(6.4) should be understood as $t \to -\infty$. The algebraic equations (6.3)-(6.4) for x_1 and x_2 have 3 set of solutions. Considering

(6.5)
$$u(t) = \sqrt{\frac{-t}{2}} \left(1 - \frac{1}{8} (-t)^{-3} - \frac{73}{128} (-t)^{-6} - \frac{10657}{1024} (-t)^{-9} + \cdots \right),$$

we can write out explicitly the 3 set of solutions as following.

Set A:
$$x_1 = 1 + \frac{1}{\sqrt{2}}(-t)^{-\frac{3}{2}} + \cdots$$
, $x_2 = 1 - \frac{1}{4\sqrt{2}}(-t)^{-\frac{3}{2}} + \cdots$.
Set B: $x_1 = -8(-t)^3 + \frac{57}{2} + \cdots$, $x_2 = 2\sqrt{2}(-t)^{\frac{3}{2}} - \frac{39}{4\sqrt{2}}(-t)^{-\frac{3}{2}} + \cdots$.
Set C: $x_1 = 1 - \frac{1}{\sqrt{2}}(-t)^{-\frac{3}{2}} + \cdots$, $x_2 = -1 - \frac{1}{\sqrt{2}}(-t)^{-\frac{3}{2}} + \cdots$.

The solutions of Set 2 and Set 3 are contradictory with the fact that $x_1 > 0$ and $x_2 > 0$. So we get (6.1). Considering $q_2(t) = \tilde{q}_2(t) - 1$ and $u(t) \sim \sqrt{\frac{-t}{2}}$, (6.2) is immediately obtained.

By Proposition 6.6, we will see (6.1) is the general case.

Proposition 6.4. For i = 1, 2, 0 and j = 2, 0, Φ_{ij} are all positive and monotonic decreasing. Furthermore, all of them approach to positive infinity as $t \to -\infty$.

Proof. (1) j = 2 case.

Both $\Phi_{12}(t) \xrightarrow{t \to \infty} 0_+$ and $\Phi_{22}(t) \xrightarrow{t \to \infty} 0_+$ are obvious. By $\Phi_{02}(t) \xrightarrow{t \to \infty} \frac{3^{1/6}}{8\pi} e^{-\frac{8}{9}t^{3/2}} t^{-\frac{1}{2}}$, we get $\Phi_{02}(t) \xrightarrow{t \to \infty} 0_+$. Further, $\frac{2}{3}\Phi_{02}(t) - \frac{k_0 - 2}{6}u(t)\Phi_{12}(t) = \frac{4 - k_0}{8 \times 3^{5/6}\pi} e^{-\frac{8}{9}t^{3/2}} (t^{-\frac{1}{2}} + O(t^{-2}))$. Thus $\frac{2}{3}\Phi_{02}(t) - \frac{k_0 - 2}{6}u(t)\Phi_{12}(t) \xrightarrow{t \to \infty} 0_+$. By Lemma 6.3, we proved the proposition for j = 2. (2) j = 0 case.

$$-\frac{4\pi}{3^{4/3}} \left(\operatorname{Ai}(\frac{t}{3^{2/3}}) \int_{-\infty}^{t} \operatorname{Bi}(\frac{s}{3^{2/3}}) u(s) ds - \operatorname{Bi}(\frac{t}{3^{2/3}}) \int_{-\infty}^{t} \operatorname{Ai}(\frac{s}{3^{2/3}}) u(s) ds \right) = e^{-\frac{2}{3}t^{3/2}} \left(\frac{3}{4\sqrt{\pi}} t^{-\frac{5}{4}} + O(t^{-\frac{11}{4}}) \right)$$

and

$$\frac{2\pi}{3} \left(\operatorname{Ai}'(\frac{t}{3^{2/3}}) \int_{\infty}^{t} \operatorname{Bi}(\frac{s}{3^{2/3}}) u(s) ds - \operatorname{Bi}'(\frac{t}{3^{2/3}}) \int_{\infty}^{t} \operatorname{Ai}(\frac{s}{3^{2/3}}) u(s) ds \right) = e^{-\frac{2}{3}t^{3/2}} \left(\frac{3}{8\sqrt{\pi}} t^{-\frac{3}{4}} + O(t^{-\frac{9}{4}}) \right),$$

 $\Phi_{10} \xrightarrow{t \to \infty} 0_+$ and $\Phi_{20} \xrightarrow{t \to \infty} 0_+$ are obtained. Obviously, $\Phi_{00}(t) \xrightarrow{t \to \infty} 1 > 0$. Also, $\frac{2}{3}\Phi_{00}(t) - \frac{k_0 - 2}{6}u(t)\Phi_{10}(t) \xrightarrow{t \to \infty} \frac{2}{3} > 0$. By Lemma 6.3, the proposition is also true for j = 0.

Proposition 6.5. For any fixed finite real t_0 , if $c_2 \ge 0$, $c_0 > 0$ and c_1 is sufficiently small, then $c_1\Phi_{11} + c_2\Phi_{12} + c_0\Phi_{10}$, $c_1\Phi_{21} + c_2\Phi_{22} + c_0\Phi_{20}$ and $c_1\Phi_{01} + c_2\Phi_{02} + c_0\Phi_{00}$ are all monotonic decreasing and positive on $(-\infty, t_0]$. Furthermore, $c_1\Phi_{01} + c_2\Phi_{02} + c_0\Phi_{00} > 0$ for $t \ge t_0$.

Proof. Let $\phi_1(t) = c_1\Phi_{11} + c_2\Phi_{12} + c_0\Phi_{10}$, $\phi_2(t) = c_1\Phi_{21} + c_2\Phi_{22} + c_0\Phi_{20}$ and $\phi_0(t) = c_1\Phi_{01} + c_2\Phi_{02} + c_0\Phi_{00}$. It is obvious there exists $\delta_1 > 0$ such that $\phi_1(t_0) > 0$, $\phi_2(t_0) > 0$, $\phi_0(t_0) > 0$ and $\frac{2}{3}\phi_0(t_0) - \frac{k_0-2}{6}u(t_0)\phi_1(t_0) > 0$ for any $|c_1| < \delta_1$, since they are all greater than 0 for $c_1 = 0$. By Lemma 6.3, $\phi_1(t)$, $\phi_2(t)$ and $\phi_0(t)$ are positive and monotonic decreasing for $t \leq t_0$. By (5.6), there exists $\delta_2 > 0$ such that $\phi_0(t) > 0$ for all $t \in [t_0, \infty)$ for $|c_1| < \delta_2$, since $c_2\Phi_{02}(t) + c_0\Phi_{00}(t) > 0$ for all $t \geq t_0$. Let $\delta = \min(\delta_1, \delta_2)$. Then, if $|c_1| < \delta$, $\phi_1(t)$, $\phi_2(t)$ and $\phi_0(t)$ have all the desired properties.

After changing of variables $s = \sqrt{-t}$, $\tilde{\phi}_1(s) = \phi_1(t)$, $\tilde{\phi}_2(s) = \phi_2(t)$, $\tilde{\phi}_0(s) = \phi_0(t)$, we can see the ODE system for $\tilde{\phi}_1(s)$, $\tilde{\phi}_2(s)$ and $\tilde{\phi}_0(s)$ satisfies all the requirements of Theorem 12.3 of [21]. After changing the variables back, we get the following result.

Proposition 6.6. At $t = -\infty$, $\phi_1(t)$ and $\phi_2(t)$ and $\phi_0(t)$ have asymptotics

$$\phi_1(t) \sim k_P \times \varphi_{1P}(t) + k_O \times \varphi_{1O}(t) + k_N \times \varphi_{1N}(t),$$

(6.7)
$$\phi_2(t) \sim k_P \times \varphi_{2P}(t) + k_O \times \varphi_{2O}(t) + k_N \times \varphi_{2N}(t),$$

(6.8)
$$\phi_0(t) \sim k_P \times \varphi_{0P}(t) + k_O \times \varphi_{0O}(t) + k_N \times \varphi_{0N}(t),$$

where

(6.9)
$$\varphi_{1P}(t) = \left(\sqrt{2}(-t)^{-\frac{1}{2}} + \frac{55}{48}(-t)^{-2} + \frac{9107}{1536\sqrt{2}}(-t)^{-\frac{7}{2}} + \cdots\right)(-t)^{\frac{1}{12}}e^{\frac{2\sqrt{2}}{9}(-t)^{3/2}},$$

$$(6.10) \quad \varphi_{2P}(t) = \left(1 - \frac{5}{48\sqrt{2}}(-t)^{-\frac{3}{2}} - \frac{1013}{3072}(-t)^{-3} - \frac{2547101}{1327104\sqrt{2}}(-t)^{-\frac{9}{2}} + \cdots\right)(-t)^{\frac{1}{12}}e^{\frac{2\sqrt{2}}{9}(-t)^{3/2}},$$

$$(6.11) \quad \varphi_{0P}(t) = \left(1 + \frac{7}{48\sqrt{2}}t^{-\frac{3}{2}} + \frac{145}{1024}(-t)^{-3} + \frac{1496311}{1327104\sqrt{2}}t^{-\frac{9}{2}} + \cdots\right)(-t)^{\frac{1}{12}}e^{\frac{2\sqrt{2}}{9}(-t)^{3/2}},$$

$$(6.12) \quad \varphi_{1O}(t) = \left(1 + \frac{67}{72}(-t)^{-3} + \frac{551671}{10368}(-t)^{-6} + \frac{22894539769}{2239488}(-t)^{-9} + \cdots\right)(-t)^{-\frac{1}{6}},$$

$$(6.13) \quad \varphi_{2O}(t) = \left(-\frac{1}{4}(-t)^{-1} - \frac{1273}{288}(-t)^{-4} - \frac{20411827}{41472}(-t)^{-7} + \cdots\right)(-t)^{-\frac{1}{6}},$$

$$(6.14) \quad \varphi_{0O}(t) = \left(\frac{1}{\sqrt{2}}(-t)^{-\frac{5}{2}} + \frac{1009}{18\sqrt{2}}(-t)^{-\frac{11}{2}} + \frac{6873355}{648\sqrt{2}}(-t)^{-\frac{17}{2}} + \cdots\right)(-t)^{-\frac{1}{6}},$$

$$(6.15) \quad \varphi_{1N}(t) = \left(-\sqrt{2}(-t)^{-\frac{1}{2}} + \frac{55}{48}(-t)^{-2} - \frac{9107}{1536\sqrt{2}}(-t)^{-\frac{7}{2}} + \cdots\right)(-t)^{\frac{1}{12}}e^{-\frac{2\sqrt{2}}{9}(-t)^{3/2}},$$

$$(6.16) \quad \varphi_{2N}(t) = \left(1 + \frac{5}{48\sqrt{2}}(-t)^{-\frac{3}{2}} - \frac{1013}{3072}(-t)^{-3} + \frac{2547101}{1327104\sqrt{2}}(-t)^{-\frac{9}{2}} + \cdots\right)(-t)^{\frac{1}{12}}e^{-\frac{2\sqrt{2}}{9}(-t)^{3/2}},$$

$$(6.17) \quad \varphi_{0N}(t) = \left(-1 + \frac{7}{48\sqrt{2}}t^{-\frac{3}{2}} - \frac{145}{1024}(-t)^{-3} + \frac{1496311}{1327104\sqrt{2}}t^{-\frac{9}{2}} + \cdots\right)(-t)^{\frac{1}{12}}e^{-\frac{2\sqrt{2}}{9}(-t)^{3/2}}.$$

Remark 6.7. Proposition 6.6 gives a straightforward explanation for the 3 sets of solutions appearing in the proof of Lemma 6.3. If $k_P \neq 0$, the limits are given by the Set A. If $k_P = 0$ and $k_O \neq 0$, the limits are given by the Set C. There is no other possibility for the limits. However, the understanding of Proposition 6.6 is subtle: for example, if in case the best approximation (obtained by optimal truncation) of u(t) by its asymptotic series has an error more than the order of $e^{-\frac{\sqrt{2}}{3}(-t)^{3/2}}$, the lower order terms in (6.6)-(6.8) lost their meaning for REAL t. Fortunately, the error order of the best approximation of u(t) by its asymptotic series is $e^{-(-t)^{3/2}}$. So all terms in (6.6)-(6.8) are contributing.

So we have constructed two sets of solutions for (4.5)-(4.7): at $t = \infty$, we have $(\Phi_{1,i}, \Phi_{2,i}, \Phi_{0,i})$, i = 1, 2, 0; and at $t = -\infty$, we have $(\varphi_{1,i}, \varphi_{2,i}, \varphi_{0,i})$, i = P, O, N. Therefore, they only differ by a constant matrix

$$\begin{pmatrix} \Phi_{11}(t) & \Phi_{12}(t) & \Phi_{10}(t) \\ \Phi_{21}(t) & \Phi_{22}(t) & \Phi_{20}(t) \\ \Phi_{01}(t) & \Phi_{02}(t) & \Phi_{00}(t) \end{pmatrix} = \begin{pmatrix} \varphi_{1P}(t) & \varphi_{1O}(t) & \varphi_{1N}(t) \\ \varphi_{2P}(t) & \varphi_{2O}(t) & \varphi_{2N}(t) \\ \varphi_{0P}(t) & \varphi_{0O}(t) & \varphi_{0N}(t) \end{pmatrix} \begin{pmatrix} k_{P1} & k_{P2} & k_{P0} \\ k_{O1} & k_{O2} & k_{O0} \\ k_{N1} & k_{N2} & k_{N0} \end{pmatrix}.$$

 $\Phi_{ij}(t) \xrightarrow{t \to -\infty} \infty$ for i = 2, 0 and j = 1, 2, 0 mean $k_{P2} > 0$ and $k_{P0} > 0$. In fact, their approximate values are $k_{P2} \approx 0.1678571$ and $k_{P0} \approx 0.6235798$. More accurate values of them are given in Section 7, where they are determined up to more than 100 digits.

Now we are able to prove Theorem 1.2.

Proof. Let $c_2 \ge 0$ and c_1 be sufficiently small. Define $\phi_1(t) = c_1\Phi_{11} + c_2\Phi_{12} + \Phi_{10}$, $\phi_2(t) = c_1\Phi_{21} + c_2\Phi_{22} + \Phi_{20}$, $\phi_0(t) = c_1\Phi_{01} + c_2\Phi_{02} + \Phi_{00}$. Then $\phi_1(t)$, $\phi_2(t)$ and $\phi_0(t)$ satisfy (4.5)-(4.7). Next define $\tilde{q}_2(t) = \frac{\phi_1(t)}{\phi_0(t)}u(t)$ and $\alpha(t) = \frac{\phi_2(t)}{\phi_0(t)}u(t)$. By (4.4), $\tilde{q}_2(t)$ and $\alpha(t)$ satisfy (4.2)-(4.3). By Proposition 6.5, $\tilde{q}_2(t)$ and $\alpha(t)$ are smooth on $(-\infty, \infty)$. By Proposition 5.2 and 6.6, $\tilde{q}_2(t)$ and $\alpha(t)$ have desired asymptotics at $t = \infty$ and $t = -\infty$.

7 Numerical experiments about Figure 1

In this section, we give the details to generate Figure 1. A few important data, such as the numerical values of the connection data, are also given, as well as some interesting observations from the numerical experiments.

7.1 Description of the procedure

By Section 6, we know the singularities of $q_2(t)$ and $\alpha(t)$ are completely determined by the zeroes of $\phi_0(t) = \Phi_{00}(t) + c_1\Phi_{01}(t) + c_2\Phi_{02}(t)$. So our first step is to obtain the numerical solutions of $\Phi_{ij}(t)$, i, j = 1, 2, 0, for $t \in [t_N, t_P]$. Since $\phi_0(t) \xrightarrow{t \to \infty} 1 > 0$, we must require $\phi_0(-\infty) \ge 0$ in order that $q_2(t)$ and $\alpha(t)$ have no zeroes for $t \in (-\infty, \infty)$. Therefore, our second step is to compute the matrix elements k_{P1} , k_{P2} and k_{P0} , which will reflect the main behaviors of the solution near $t = -\infty$. For moderate t, we use the numerical solutions to resolve if $\phi_0(t)$ has zeroes, which constitutes our last step. More precisely, we determine the boundary between R_{smooth} and $R_{singular}$ by seeking the minimal c_2 such that $\Phi_{00}(t) + c_1\Phi_{01}(t) + c_2\Phi_{02}(t) \ge 0$ for all $t \in (-\infty, \infty)$ for given $c_1 \in \mathbb{R}$. Since $\Phi_{02}(t) > 0$, the problem is simplified to find the minimum of $\frac{\Phi_{00}(t)}{\Phi_{02}(t)} + c_1\frac{\Phi_{01}(t)}{\Phi_{02}(t)}$ for given c_1 , i.e., $c_2 = -\min_{\forall t \in (-\infty, \infty)} \left(\frac{\Phi_{00}(t)}{\Phi_{02}(t)} + c_1\frac{\Phi_{01}(t)}{\Phi_{02}(t)}\right)$.

Obviously, we have to do numerical integration of ODEs. Currently, the most precise ODE integrator, such as Taylor[14] or high-order Runge-Kutta, can integrate an ODE numerically with precision up to 1000 digits. For convenience, we use the build-in 'NDSolve' of Mathematica to do the numerical integration for (4.2)-(4.3). The default option of 'NDSolve' is inappropriate to do high-precision numerical integration. By explicitly giving the 'Method' option of 'NDSolve', we can force it to use the Gauss-Legendre Runge-Kutta method, which is suitable for the high-precision purpose. To save running time, we manage to let the typical precision be of order 10^{-120-1} . The stages of the Runge-Kutta method are set according to the precision goal of the numerical integration. As a rule, we always let the stages greater than 100, i.e, the order of the numerical scheme is always more than 200. The step-sizes h are chosen as $0.01 \le h \le 0.05$. By rough but careful estimations for each case, we guarantee that the errors

 $^{^{1}}$ It does not mean the final error or final relative error is less than 10^{-120} . It just mean, the relative error is less than 10^{-120} for every step.

generated by the numeric scheme itself are always negligible, comparing to the errors that exist on the boundaries and are propagated by the ODE system.

7.2 Determine T_P

 T_P is determined by two key factors: the truncation orders of Φ_{ij} at $t = \infty$ and the precision goal of the numerical integration. We use (5.1)-(5.9) as the truncation of Φ_{ij} since the higher order truncation will involve multiple integrals, which is difficult to get satisfactory high-precision results.

We demand the error of $\Phi_{ij}(t)$ at t=0 is of order 10^{-120} . For the solution $\Phi_{i1}(t)$, we can show their errors at t=0 are of order $e^{-\frac{8}{9}t_P^{3/2}}$. Solving $e^{-\frac{8}{9}t_P^{3/2}}=10^{-120}$, we get $t_P\approx 45.888$. For convenience, we set $t_P=46$, at which the relative errors are of order $e^{-\frac{4}{3}46^{3/2}}\approx 2.19\times 10^{-181}$. So we set the precision goal of the numeric scheme as 10^{-182} in computing $\Phi_{i1}(t)$. By a similar way, we could show it is appropriate to set $t_P=36$ and the precision goal as 10^{-120} in computing $\Phi_{i2}(t)$. In computing $\Phi_{i0}(t)$, we also use $t_P=36$ and the precision 10^{-120} .

7.3 Determine T_N

By (6.6)-(6.8), each k_O term contributes to a portion of order $e^{-\frac{2\sqrt{2}}{9}(-t)^{3/2}}$. By solving $e^{-\frac{2\sqrt{2}}{9}(-t)^{3/2}} = 10^{-120}$, we get $t \approx -91.7761$. This means the k_O terms can be neglected when t < -91.7761. For convenience, we set $T_N = -92$.

7.4 The numerical solution of u(t)

For computation efficiency, the numerical solution of u(t) is first obtained independently on $[t_M, t_H]$. We demand the max error of u(t) is of order 10^{-120} . Since the best approximation of u(t) by (6.5) has an error of order $e^{-(-t)^{3/2}}$, t_M is obtained by solving $e^{-(-t)^{3/2}} = 10^{-120}$, i.e., $t_M \approx -42.42$. For safety, we set $t_M = -44$. So, for $t_N \leqslant t < t_M$, we use the asymptotic expansion (6.5) up to the $(-t)^{-414}$ term to compute u(t). For $t_M \leqslant t \leqslant t_H$, u(t) is obtained by the high-precision numerical integration of (1.11). Let

$$\tilde{u}(t) = u(t) + \epsilon \mathscr{U}(t),$$

where ϵ is an infinitesimal, and \tilde{u} also satisfies (1.11). Then $\mathcal{U}(t)$ satisfies

$$\mathscr{U}''(t) = (t + 6u(t)^2) \mathscr{U}(t).$$

As $u(t) \xrightarrow{t \to -\infty} \sqrt{\frac{-t}{2}} + \cdots$, $\mathscr{U}(t)$ is of order $e^{\frac{1}{3}(-2t)^{3/2}}$. In the numerical experiments, $e\mathscr{U}(t)$ is understood as the error. So $\epsilon \sim 3 \times 10^{-240}$ in order that at t=-44 the error is of order 10^{-120} . Then the error of u(t) at t=0 should be of order 10^{-240} . The computational error of u(t) behaves like e_u Ai(t) for t>0. So, the relative error at $t=t_H$ should also be of order 10^{-240} . For safety, we manage the relative error at $t=t_H$ to be of order 10^{-250} .

The value of t_H is related to how u(t) is approximated near $t=\infty$. We take

(7.1)
$$u(t) \approx \operatorname{Ai}(t) - 2\pi \operatorname{Ai}(t) \int_{-\infty}^{t} \operatorname{Bi}(s) \operatorname{Ai}(s)^{3} ds + 2\pi \operatorname{Bi}(t) \int_{-\infty}^{t} \operatorname{Ai}(s)^{4} ds$$

as the approximation of u(t). The error order of the approximation (7.1) is about $e^{-\frac{10}{3}t^{3/2}}$. So the relative error is of order $e^{-\frac{8}{3}t^{3/2}}$. By solving $e^{-\frac{8}{3}t^{3/2}}=10^{-250}$, $t_H\approx 35.985$ is obtained. For convenience, we set $t_H=36$. We use 250 digits in computing the numerical solution of u(t).

7.5 Transformations to avoid small step size

Fixing the step size, the Runge-Kutta method will be generally more accurate to integrate a slow-varying system. To see the crux, let us consider the approximation of $e^{-t^{3/2}}$ by polynomials. It is easy to see that the relative error of the approximation on interval [100, 100.01] is almost the same as the one on interval [1,1.1] when using the same degree of approximation polynomials. This means smaller step size is needed for large t if the system increases or decreases too fast. To avoid the small step size for large t, we use Table 1 to transform the fast variables to slow ones.

Table 1: Transformations used to transform the fast variables to the slow ones			
fast variables	t < -1	$-1 \leqslant t \leqslant 1$	t > 1
u	u(t)	u(t)	$\tilde{u}(t) = u(t)e^{\frac{2}{3}t^{3/2}}$
$\Phi_{i1}, i = 1, 2, 0.$	$\tilde{\Phi}_{i1}(t) = \Phi_{i1}(t)e^{-\frac{2\sqrt{2}}{9}(-t)^{3/2}}$	$\Phi_{i1}(t)$	$ \tilde{\Phi}_{11}(t) = \Phi_{11}(t)e^{-\frac{2}{9}t^{3/2}} \tilde{\Phi}_{21}(t) = \Phi_{21}(t)e^{-\frac{2}{9}t^{3/2}} \tilde{\Phi}_{01}(t) = \Phi_{01}(t)e^{\frac{4}{9}t^{3/2}} $
$\Phi_{i2}, i = 1, 2, 0.$	$\tilde{\Phi}_{i2}(t) = \Phi_{i2}(t)e^{-\frac{2\sqrt{2}}{9}(-t)^{3/2}}$	$\Phi_{i2}(t)$	$ \tilde{\Phi}_{12}(t) = \Phi_{12}(t)e^{\frac{2}{9}t^{3/2}} \tilde{\Phi}_{22}(t) = \Phi_{22}(t)e^{\frac{2}{9}t^{3/2}} \tilde{\Phi}_{02}(t) = \Phi_{02}(t)e^{\frac{8}{9}t^{3/2}} $
$\Phi_{i0}, i = 1, 2, 0.$	$\tilde{\Phi}_{i0}(t) = \Phi_{i0}(t)e^{-\frac{2\sqrt{2}}{9}(-t)^{3/2}}$	$\Phi_{i0}(t)$	$\tilde{\Phi}_{10}(t) = \Phi_{10}(t)e^{\frac{2}{3}t^{3/2}}$ $\tilde{\Phi}_{20}(t) = \Phi_{20}(t)e^{\frac{2}{3}t^{3/2}}$ $\tilde{\Phi}_{20}(t) = (\Phi_{20}(t)e^{\frac{2}{3}t^{3/2}}$

7.6 Numerical results

The main numerical results are displayed in Figure 1.

7.6.1 The values of k_{P1} , k_{P2} and k_{P0}

In principle, k_{Pi} , i = 1, 2, 0, can be computed by any of $\lim_{t \to -\infty} \frac{\Phi_{1i}(t)}{\varphi_{1P}(t)}$, $\lim_{t \to -\infty} \frac{\Phi_{2i}(t)}{\varphi_{2P}(t)}$ or $\lim_{t \to -\infty} \frac{\Phi_{0i}(t)}{\varphi_{0P}(t)}$. In our numerical experiments, we use

(7.2)
$$k_{Pi} = \frac{\Phi_{0i}(t_N)}{\varphi_{0P}(t_N)},$$

which is a little more accurate than the other two choices. In (7.2), $\Phi_{0i}(t_N)$ are obtained directly from the numerical integration of ODEs of $\tilde{\Phi}_{ji}$, while $\varphi_{0P}(t_N)$ is calculated by its asymptotic expansion (6.11), where $\varphi_{0P}(t)$ is computed up to the term $c_{354}^0 \times (-t)^{354}(-t)^{\frac{1}{12}}e^{\frac{2\sqrt{2}}{9}(-t)^{3/2}}$. It is not surprising that c_{354}^0 is very large since (6.11) is an asymptotic expansion. In fact, the term $c_{354}^0 \times (-t)^{354}$ contributes about 1.74545×10^{-126} at t=-92. So k_{Pi} is determined with an approximate precision of 10^{-120} . The final numerical results of k_{Pi} are

- $(7.3) k_{P1} = -0.0969123435570255523226380385083332\cdots,$
- $(7.4) k_{P2} = 0.167857102921338590132168687360301197 \cdots,$
- $(7.5) k_{P0} = 0.62357981669501424223251084362366955\cdots.$

7.6.2 $\Phi_{ij}(t)$ near t = 0

By (5.1)-(5.1), (6.9)-(6.11) and (7.3)-(7.5), the main behaviors of $\Phi_{ij}(t)$ at $t = \pm \infty$ have been described. We demonstrate their behaviors on the "transition zone" by Figures 3, 4 and 5.




Figure 3. Plots of Φ_{11} (red), Φ_{21} (green) and Φ_{01} (blue). At any t, $\Phi_{11}(t)$ is always the largest. $\Phi_{01}(t)$ and $\Phi_{21}(t)$ intersect at $t \approx -4.3166745$.




Figure 4. Plots of Φ_{12} (red), Φ_{22} (green) and Φ_{02} (blue). All of them are positive. There are 4 intersections in the figure: $t_1 \approx 3.30090866$, $t_2 \approx -1.7223227$, $t_3 \approx -3.5443904$, $t_4 \approx -3.72852126$. For $t > t_1$, $\Phi_{22}(t) > \Phi_{12}(t) > \Phi_{02}(t)$. For $t_1 > t > t_2$, $\Phi_{12}(t) > \Phi_{22}(t) > \Phi_{02}(t)$. For $t_2 > t > t_3$, $\Phi_{12}(t) > \Phi_{02}(t) > \Phi_{02}(t) > \Phi_{02}(t)$. For $t_3 > t > t_4$, $\Phi_{02}(t) > \Phi_{12}(t) > \Phi_{22}(t)$. For $t < t_4$, $\Phi_{02}(t) > \Phi_{22}(t) > \Phi_{12}(t)$.




Figure 5. Plots of Φ_{10} (red), Φ_{20} (green) and Φ_{00} (blue). All of them are positive. For all t, $\Phi_{00}(t) > \Phi_{20}(t) > \Phi_{10}(t)$.

7.6.3 The critical point P_c

In Section 7.1, we have explained $c_2 = -\min_{\forall t \in (-\infty,\infty)} \left(\frac{\Phi_{00}(t)}{\Phi_{02}(t)} + c_1 \frac{\Phi_{01}(t)}{\Phi_{02}(t)}\right)$ on the boundary between R_{smooth} and $R_{singular}$. Given c_1 , let the minimum is achieved at $t = t_z$. The numerical results show the t_z is unique for any given c_1 . So, on the boundary cure, $t_z = t_z(c_1)$. It is obvious that both t_z and c_2 must approach to ∞ when $c_1 \to -\infty$. As c_1 increases gradually to P_c , t_z decreases and finally approaches to $-\infty$ as displayed by Figure 6.




Figure 6. Plots of c_2 (red) and t_z (green). The c_2 curve, which is the boundary between R_{smooth} and $R_{singular}$, is smooth. Though it looks very like a straight line, it is indeed a curve. The t_z curve has apparently a singularity near $c_1 = -\frac{1}{2}\frac{k_{P0}}{k_{P1}} \approx 3.217236287$.

On the right of P_c , the minimum is always achieved at $t = -\infty$, i.e., $t_z = -\infty$. So we have

(7.6)
$$c_2 = -\left(\frac{k_{P0}}{k_{P2}} + c_1 \frac{k_{P1}}{k_{P2}}\right) = -\frac{k_{P0}}{k_{P2}} - \frac{k_{P1}}{k_{P2}}c_1,$$

which is the straight line right of P_c in Figure 1. For the critical point P_c , the interesting observation from the numerical experiment is $c_1 = -\frac{1}{2}\frac{k_{P0}}{k_{P1}}$. Then, from (7.6), $c_2 = -\frac{1}{2}\frac{k_{P0}}{k_{P2}}$ at P_c .

7.6.4 The values of k_{O1} , k_{O2} , k_{O0} , k_{N1} , k_{N2} and k_{N0}

Integrating u(t) and $\Phi_{ij}(t)$ numerically along the path O-A-B in Figure 7, we have obtained the values of k_{P1} , k_{P2} and k_{P0} with about 120 digits of precision. But k_{O1} , k_{O2} , k_{O0} , k_{N1} , k_{N2} and k_{N0} can not be obtained in this way. To calculate them, we have to extend our numerical integration from the real line to the complex plane of t as displayed by Figure 7.




Figure 7. Paths used to integrate $\Phi_{ij}(t)$. $\angle BOD = \frac{7}{10}$ and $\angle BOE = \frac{2}{3}\arccos(\frac{135\ln 10}{184\sqrt{46}}) \approx 0.879372$. The boundary between the light yellow region and the light green one is $r = \frac{44}{\left(\cos(\frac{3}{2}\theta - \frac{3\pi}{2})\right)^{2/3}}$. Φ_{ij} at point D are used to calculate k_{O1} , k_{O2} and k_{O0} . k_{N1} , k_{N2} and k_{N0} are calculated from Φ_{ij} at point E. For precision reason, we use path O-C-D rather than arc \widehat{BD} to numerically integrate the ODEs for $\widetilde{\Phi}_{ij}$. Path O-E is used for the same reason. u(t) on the path in the light green region is obtained by the numerical integration of (1.11) while on the path in the light yellow region it is calculated by the expansion (6.5) up to the $(-t)^{-414}$ term.

We compute k_{Oi} by $k_{Oi} = \frac{\Phi_{1i}(t) - k_{Pi} \times \varphi_{1P}(t)}{\varphi_{1O}(t)}$, where t is chosen as the point D. The argument of D is chosen by solving $e^{\frac{2\sqrt{2}}{9}(92)^{3/2}\cos(\frac{3}{2}\theta + \frac{3\pi}{2})} = 10^{60}$, i.e., $\pi - \theta \approx 0.699535$. For simplicity, we choose $\pi - \theta = \frac{7}{10}$. It is easy to show that $\tilde{\Phi}_{ij}$ lost their precision when they are integrated numerically along arc $\widehat{\text{BD}}$ starting from B. So we integrate them numerically along the ray O-D, by which $\tilde{\Phi}_{ij}$ can be guaranteed to have about 120 digits of precision. k_{Oi} obtained by this way can be shown to have about 60 digits of precision, which is almost the best that we can expect for the computation of k_{Oi} when u(0) is computed with about 240 digits of precision.

is computed with about 240 digits of precision. To compute k_{Ni} , we use $k_{Ni} = \frac{\Phi_{2i}(t) - k_{Pi} \times \varphi_{2P}(t) - k_{Oi} \times \varphi_{2O}}{\varphi_{2N}(t)}$, where t is chosen as the point E. The argument of E is chosen by solving $e^{\frac{2\sqrt{2}}{9}(92)^{3/2}\cos(\frac{3}{2}\theta + \frac{3\pi}{2})} = 10^{30}$. At the first sight, one may want to evaluate k_{Ni} from Φ_{ij} on the dotted line. But [10] has proved

(7.7)
$$u(t) = \sqrt{\frac{-t}{2}} \left(1 + O((-t)^{-3/2}) \right) + \frac{i}{2^{\frac{7}{4}} \sqrt{\pi}} (-t)^{-\frac{1}{4}} e^{-\frac{2\sqrt{2}}{3}(-t)^{3/2}} \left(1 + O(t^{-\frac{1}{4}}) \right),$$

for $\frac{2\pi}{3} \leq \arg(t) < \frac{4\pi}{3}$. So we should not use the expansions (6.9)-(6.17) near $\theta = \frac{2\pi}{3}$. Considering the exponential term of (7.7), we can show k_{Ni} are best calculated near E. Also, it can be shown that k_{Ni} calculated in this way have about 30 digits of precision.

The final numerical values of k_{ij} , i = O, N, j = 1, 2, 0, are

$$(7.8) k_{O1}^{+} = (k_{O1}^{-})^{*} = 0.474787653555570800096 \dots + i \times 0.091372926529406526556 \dots,$$

(7.9)
$$k_{O2}^{+} = (k_{O2}^{-})^{*} = 0.274118779588219579669 \cdots - i \times 0.158262551185190266698 \cdots,$$

$$(7.10) k_{O0}^+ = \left(k_{O2}^-\right)^* = -1.018336045084649924885 \cdot \cdot \cdot - \mathrm{i} \times 0.58793658975512151298 \cdot \cdot \cdot ,$$

$$(7.11) k_{N1}^+ = (k_{N1}^-)^* = -0.19583328674156168848 \cdots + i \times 0.048456171778512776161 \cdots,$$

$$(7.12) k_{N2}^{+} = (k_{N2}^{-})^{*} = 0.0484561717785127761613 \cdots + i \times 0.083928551460669295066 \cdots,$$

$$(7.13) k_{N0}^+ = (k_{N0}^-)^* = -0.360023975030083963185\cdots.$$

From our numerical results, we observe that $k_{N1}^+ = \left(\frac{7\sqrt{3}}{6} - \frac{1}{2}\mathrm{i}\right)k_{P1}$, $k_{N2}^+ = \left(\frac{\sqrt{3}}{6} + \frac{1}{2}\mathrm{i}\right)k_{P2}$ and $k_{N0}^+ = -\frac{\sqrt{3}}{3}k_{P0}$ with the errors less than 10^{-30} , which are consistent with the estimated precision of the numerical k_{N1}^+ , k_{N2}^+ and k_{N0}^+ . Also it is observed that $\left(\frac{k_{O2}^+}{k_{P2}}\right)^* = -\frac{k_{O0}^+}{k_{P0}}$ with more than 60 digits of precision.

7.6.5 The solution corresponding to P_c

Let us consider the solutions of (4.5)-(4.7) described by Figure 1. We note that the solution corresponding to P_c in Figure 1 has a special property. For simplicity, we scale the solution as

$$\begin{array}{rcl} \Phi_c & = & \left(\Phi_{1c}, \Phi_{2c}, \Phi_{0c}\right) \\ & = & \frac{2}{k_{P0}} \left(\Phi_{10}, \Phi_{20}, \Phi_{00}\right) - \frac{1}{k_{P1}} \left(\Phi_{11}, \Phi_{21}, \Phi_{01}\right) - \frac{1}{k_{P2}} \left(\Phi_{12}, \Phi_{22}, \Phi_{02}\right). \end{array}$$

By the numerical connection data (7.3-7.5) and (7.8)-(7.13), it is easy to verify (within the tolerance of precision) $\Phi_c(t) \xrightarrow{t \to -\infty} -2\sqrt{3} \left(\varphi_{1N}(t), \varphi_{2N}(t), \varphi_{0N}(t)\right)$. So this special solution decreases exponentially to 0 as $t \to -\infty$. We also note the other bounded solutions at $t = -\infty$, which are spanned by Φ_c and $\frac{1}{k_{P1}} \left(\Phi_{11}, \Phi_{21}, \Phi_{01}\right) - \frac{1}{k_{P2}} \left(\Phi_{12}, \Phi_{22}, \Phi_{02}\right)$, decrease algebraically² to 0.

8 The wave function of Painlevé II

The Lax pair of Painlevé II is

$$\frac{d\Psi_0}{dx} = \hat{L}_0 \Psi_0,$$

$$\frac{d\Psi_0}{dt} = \hat{B}_0 \Psi_0,$$

where \hat{L}_0 and \hat{B}_0 are defined by (2.4) and (2.5). Unlike in Section 2 where ψ_0 is vector, here, Ψ_0 is a 2×2 matrix.

Define the six regions in the complex x-plane as

$$\Omega_j = \left\{ x \left| \frac{\pi}{2} + \frac{j-2}{3} \pi < \arg x < \frac{\pi}{2} + \frac{j}{3} \pi \right. \right\}, \quad j = 1, 2, \dots, 6.$$

Equation (8.1) has 6 canonical solutions $\Psi_0^{(j)}(x)$ defined in the regions Ω_j , $j=1,\cdots,6$,

$$\Psi_0^{(j)}(x) \xrightarrow{x \to \infty} \left(I + \frac{m_1}{x} + \cdots \right) e^{\left(\frac{x^3}{6} - \frac{xt}{2}\right)\sigma_3}, \qquad \frac{\pi}{2} + \frac{j-2}{3}\pi < \arg x < \frac{\pi}{2} + \frac{j}{3}\pi.$$

For convenience, we denote $\Omega_7 = \Omega_1$ and $\Psi_0^{(7)} = \Psi_0^{(1)}$. If Ψ_0 is known, then u(t) can be recovered by

$$u = (m_1)_{21} = -(m_1)_{12}.$$

The sector Ω_j overlaps with Ω_{j+1} . In the crossover region,

(8.3)
$$\Psi_0^{(j+1)} = \Psi_0^{(j)} S_0^{(j)}.$$

²Just as the asymptotic series hints, the numerical results show the decrease looks very like $(-t)^{-\frac{1}{6}}$.

For the case that u(t) is the Hastings-McLeod solution,

$$S_0^{(1)} = \begin{pmatrix} 1 & 0 \\ -1 & 1 \end{pmatrix}, \quad S_0^{(2)} = I_{2 \times 2}, \quad S_0^{(3)} = \begin{pmatrix} 1 & 0 \\ 1 & 1 \end{pmatrix},$$
$$S_0^{(4)} = \begin{pmatrix} 1 & -1 \\ 0 & 1 \end{pmatrix}, \quad S_0^{(5)} = I_{2 \times 2}, \quad S_0^{(6)} = \begin{pmatrix} 1 & 1 \\ 0 & 1 \end{pmatrix}.$$

Define $Y^{(j)}(x,t)$ by

(8.4)
$$Y^{(j)}(x,t) = \Psi_0^{(j)}(x,t)e^{-\left(\frac{x^3}{6} - \frac{xt}{2}\right)\sigma_3}.$$

For convenience, we call both $\Psi_0^{(j)}$ and $Y^{(j)}$ as the wave functions of Painlevé II. By (8.3), $Y^{(j)}$ satisfy the Riemann-Hilbert problem illustrated by Figure 8.




Figure 8. The original Riemann-Hilbert Problem.

To prove Theorem 1.3, a detailed analysis for the case $t \to \infty$ is needed. So we deform the original Riemann-Hilbert problem to Figure 9.




Figure 9. The final Riemann-Hilbert Problem.

By solving the Riemann-Hilbert problem of Y, one gets the following result.

Lemma 8.1. $Y^{(6)}$ and $Y^{(3)}$ have the following asymptotics:

(A) For
$$x \to \infty$$
 and fixed t, $Y^{(6)}(x,t)$ has expansion $Y^{(6)}(x,t) = I_{2\times 2} + \frac{m_1(t)}{x} + \cdots$

(B) For
$$x \to -\infty$$
 and fixed t , $Y^{(3)}(x,t)$ has expansion $Y^{(3)}(x,t) = I_{2\times 2} + \frac{m_1(t)}{x} + \cdots$.

(C)
$$\lim_{x \to \infty, t \to \infty, t^{\frac{1}{4}} | x - \sqrt{t}| \to \infty} Y^{(6)}(x, t) = I_{2 \times 2} \text{ and } \lim_{x \to -\infty, t \to \infty, t^{\frac{1}{4}} | x + \sqrt{t}| \to \infty} Y^{(3)}(x, t) = I_{2 \times 2}.$$

(D) For
$$t \to \infty$$
 and $0 \le x < \sqrt{t} - t^{\epsilon - \frac{1}{4}}$, $Y^{(6)} \to \begin{pmatrix} 1 & -e^{\frac{x^3}{3} - xt} \\ 0 & 1 \end{pmatrix}$.

(E) For
$$t \to \infty$$
 and $-\sqrt{t} + t^{\epsilon - \frac{1}{4}} < x \le 0, Y^{(3)} \to \begin{pmatrix} 1 & 0 \\ -e^{xt - \frac{x^3}{3}} & 1 \end{pmatrix}$.

In both cases (A) and (B),
$$m_1(t) = \begin{pmatrix} (u')^2 - u^4 - tu^2 & -u \\ u & -(u')^2 + u^4 + tu^2 \end{pmatrix}$$
.

Lemma 8.1 is already known, see for example [10].

Remark 8.2. Because of (2.10), at $t = -\infty$, $\kappa(t)$ is 'smaller' than other quantities in the formulae. It is unnecessary to estimate $Y^{(3)}$ and $Y^{(6)}$ so accurately at $t = -\infty$.

Lemma 8.1 fulfils parts of our purpose to prove (1.3). In fact we still need more detailed behaviour of $Y^{(3)}$ on $x = k\sqrt{t}$. For completeness, we also give the results for $Y^{(6)}$.

Before we study the asymptotics of $Y^{(6)}(x,t)$ and $Y^{(3)}(x,t)$ along $x = k\sqrt{t}$, let us first write down the ODEs for them, which our study will rely on.

By (8.4), $Y^{(j)}$ satisfies

$$\frac{dY^{(j)}}{dx} = \hat{L}_0 Y^{(k)} + \left(\frac{t}{2} - \frac{x^2}{2}\right) Y^{(j)} \sigma_3,$$
$$\frac{dY^{(j)}}{dt} = \hat{B}_0 Y^{(k)} + \frac{x}{2} Y^{(j)} \sigma_3.$$

The detailed formulae are

(8.5)
$$\frac{d}{dx} \begin{pmatrix} Y_{11}^{(j)} \\ Y_{21}^{(j)} \end{pmatrix} = \begin{pmatrix} -u(t)^2 & xu(t) - u'(t) \\ xu(t) + u'(t) & t - x^2 + u(t)^2 \end{pmatrix} \begin{pmatrix} Y_{11}^{(j)} \\ Y_{21}^{(j)} \end{pmatrix},$$

(8.6)
$$\frac{d}{dt} \begin{pmatrix} Y_{11}^{(j)} \\ Y_{21}^{(j)} \end{pmatrix} = \begin{pmatrix} 0 & -u(t) \\ -u(t) & x \end{pmatrix} \begin{pmatrix} Y_{11}^{(j)} \\ Y_{21}^{(j)} \end{pmatrix},$$

(8.7)
$$\frac{d}{dx} \begin{pmatrix} Y_{12}^{(j)} \\ Y_{22}^{(j)} \end{pmatrix} = \begin{pmatrix} -t + x^2 - u(t)^2 & xu(t) - u'(t) \\ xu(t) + u'(t) & u(t)^2 \end{pmatrix} \begin{pmatrix} Y_{12}^{(k)} \\ Y_{22}^{(k)} \end{pmatrix},$$

(8.8)
$$\frac{d}{dt} \begin{pmatrix} Y_{12}^{(j)} \\ Y_{22}^{(j)} \end{pmatrix} = \begin{pmatrix} -x & -u(t) \\ -u(t) & 0 \end{pmatrix} \begin{pmatrix} Y_{12}^{(j)} \\ Y_{22}^{(j)} \end{pmatrix}.$$

Along the line $x = k\sqrt{t}$, by

$$\frac{dY^{(j)}(k\sqrt{t},t)}{dt} = \frac{k}{2\sqrt{t}} \frac{dY^{(j)}(x,t)}{dx} \bigg|_{x=k\sqrt{t}} + \frac{dY^{(j)}(x,t)}{dt} \bigg|_{x=k\sqrt{t}}, \quad j = 3, 6,$$

we get

$$(8.9) \qquad \frac{d}{dt} \begin{pmatrix} Y_{11}^{(j)} \\ Y_{21}^{(j)} \end{pmatrix} = \begin{pmatrix} \frac{-ku(t)^2}{2\sqrt{t}} & \frac{k^2 - 2}{2}u(t) - \frac{k}{2\sqrt{t}}u'(t) \\ \frac{k^2 - 2}{2}u(t) + \frac{k}{2\sqrt{t}}u'(t) & -k\frac{(k^2 - 3)t - u(t)^2}{2\sqrt{t}} \end{pmatrix} \begin{pmatrix} Y_{11}^{(j)} \\ Y_{21}^{(j)} \end{pmatrix},$$

(8.10)
$$\frac{d}{dt} \begin{pmatrix} Y_{12}^{(j)} \\ Y_{22}^{(j)} \end{pmatrix} = \begin{pmatrix} k \frac{(k^2 - 3)t - u(t)^2}{2\sqrt{t}} & \frac{k^2 - 2}{2}u(t) - \frac{k}{2\sqrt{t}}u'(t) \\ \frac{k^2 - 2}{2}u(t) + \frac{k}{2\sqrt{t}}u'(t) & \frac{ku(t)^2}{2\sqrt{t}} \end{pmatrix} \begin{pmatrix} Y_{12}^{(j)} \\ Y_{22}^{(j)} \end{pmatrix}.$$

8.1 The asymptotics of $Y^{(6)}(x,t)$ for $x\to\infty$ and $t\to\infty$ along $x=k\sqrt{t}$

In this case k > 0. First let us assume $k \neq 1$. By Lemma 8.1, we know $Y_{11}^{(6)} \to 1$ and $Y_{22}^{(6)} \to 1$. Therefore, the approximate differential equations for $Y_{21}^{(6)}$ and $Y_{12}^{(6)}$ along $x = k\sqrt{t}$ are

(8.11)
$$\frac{d}{dt}Y_{21}^{(6)} = \frac{(3-k^2)k}{2}\sqrt{t}Y_{21}^{(6)} + \frac{k^2-2}{2}\operatorname{Ai}(t) + \frac{k}{2\sqrt{t}}\operatorname{Ai}'(t),$$

(8.12)
$$\frac{d}{dt}Y_{12}^{(6)} = \frac{(k^2 - 3)k}{2}\sqrt{t}Y_{12}^{(6)} + \frac{k^2 - 2}{2}\operatorname{Ai}(t) - \frac{k}{2\sqrt{t}}\operatorname{Ai}'(t),$$

while the approximate differential equations for $Y_{21}^{(6)}$ and $Y_{12}^{(6)}$ for fixed but large t are

(8.13)
$$\frac{d}{dx}Y_{21}^{(6)} = (t - x^2)Y_{21}^{(6)} + x\operatorname{Ai}(t) + \operatorname{Ai}'(t),$$

(8.14)
$$\frac{d}{dx}Y_{12}^{(6)} = (-t + x^2)Y_{12}^{(6)} + x\operatorname{Ai}(t) - \operatorname{Ai}'(t).$$

8.1.1 $Y_{21}^{(6)}$

Case 0 < k < 2:

In this case³, the solution of (8.11) is

$$(8.15) Y_{21}^{(6)} = \Upsilon_1(k)e^{\frac{k}{3}(3-k^2)t^{3/2}} + e^{\frac{k}{3}(3-k^2)t^{3/2}} \int_{\infty}^{t} e^{\frac{k}{3}(k^2-3)s^{3/2}} \left(\frac{k^2-2}{2}\operatorname{Ai}(s) + \frac{k}{2\sqrt{s}}\operatorname{Ai}'(s)\right) ds.$$

The solution of (8.13) is

(8.16)
$$Y_{21}^{(6)} = C_1(t)e^{tx-\frac{1}{3}x^3} + e^{tx-\frac{1}{3}x^3} \int_{\sqrt{t}}^x e^{-ts+\frac{1}{3}s^3} \left(s\operatorname{Ai}(t) + \operatorname{Ai}'(t)\right) ds.$$

(8.15) and (8.16) must coincide at $x = k\sqrt{t}$. Therefore, we have

$$(8.17) C_{1}(t) - \Upsilon_{1}(k) = \int_{\infty}^{t} e^{\frac{k}{3}(k^{2}-3)s^{3/2}} \left(\frac{k^{2}-2}{2}\operatorname{Ai}(s) + \frac{k}{2\sqrt{s}}\operatorname{Ai}'(s)\right) ds$$
$$- \int_{\sqrt{t}}^{k\sqrt{t}} e^{-ts + \frac{1}{3}s^{3}} \left(s\operatorname{Ai}(t) + \operatorname{Ai}'(t)\right) ds.$$

Note that the right-side of (8.17) is a solution of $\frac{\partial^2}{\partial k \partial t} f(k,t) = 0$. So (8.17) determines $C_1(t)$ and $\Upsilon_1(k)$ up to a constant. But we know $\Upsilon_1(k) = 0$ for $k \in (0, \sqrt{3}]$. So we get

(8.18)
$$C_1(t) = \int_{\infty}^t e^{-\frac{2}{3}s^{3/2}} \left(-\frac{1}{2} \text{Ai}(s) + \frac{1}{2\sqrt{s}} \text{Ai}'(s) \right) ds.$$

Taking the $t \to \infty$ limit of (8.17) and considering (8.18), we obtain

$$\Upsilon_1(k) = 0, \quad k \in (0, 2).$$

Therefore, the final result for 0 < k < 2 is

$$Y_{21}^{(6)} \approx e^{\frac{k}{3}(3-k^2)t^{3/2}} \int_{\infty}^{t} e^{\frac{k}{3}(k^2-3)s^{3/2}} \left(\frac{k^2-2}{2} \operatorname{Ai}(s) + \frac{k}{2\sqrt{s}} \operatorname{Ai}'(s)\right) ds.$$

Case k=2:

Similar to the case 0 < k < 2, we can also derive

$$\Upsilon_1(2) = 0.$$

Thus

$$Y_{21}^{(6)} \approx e^{-\frac{2}{3}t^{3/2}} \int_{\infty}^{t} e^{\frac{2}{3}s^{3/2}} \left(\operatorname{Ai}(s) + \frac{1}{\sqrt{s}} \operatorname{Ai}'(s) \right) ds.$$

³By the result of Riemann-Hilbert problem, $Y_{21}^{(6)} \rightarrow 0$ and $Y_{22}^{(6)} \rightarrow 1$ are still true for k=1.

Case k > 2:

In this case, the solution of (8.11) is

$$(8.20) Y_{21}^{(6)} = \Upsilon_1(k)e^{\frac{k}{3}(3-k^2)t^{3/2}} + e^{\frac{k}{3}(3-k^2)t^{3/2}} \int_{t_A}^t e^{\frac{k}{3}(k^2-3)s^{3/2}} \left(\frac{k^2-2}{2}\operatorname{Ai}(s) + \frac{k}{2\sqrt{s}}\operatorname{Ai}'(s)\right) ds.$$

Clearly, the first term can be neglected. Thus for k > 2,

$$(8.21) Y_{21}^{(6)} \approx e^{\frac{k}{3}(3-k^2)t^{3/2}} \int_{t_A}^t e^{\frac{k}{3}(k^2-3)s^{3/2}} \left(\frac{k^2-2}{2} \operatorname{Ai}(s) + \frac{k}{2\sqrt{s}} \operatorname{Ai}'(s)\right) ds.$$

Note t_A is a fixed arbitrary real number.

Remark 8.3. As $k \to \infty$, (8.21) is consistent with $Y_{21}^{(6)} \xrightarrow{x \to \infty} \frac{u(t)}{x}$.

8.1.2 $Y_{12}^{(6)}$

For all k > 0, (8.12) has the solution

$$(8.22) Y_{12}^{(6)} = \Upsilon_2(k)e^{\frac{k}{3}(k^2-3)t^{3/2}} + e^{\frac{k}{3}(k^2-3)t^{3/2}} \int_{\infty}^{t} e^{\frac{k}{3}(3-k^2)s^{3/2}} \left(\frac{k^2-2}{2}\operatorname{Ai}(s) - \frac{k}{2\sqrt{s}}\operatorname{Ai}'(s)\right) ds.$$

Since (8.22) has to approach 0 as $t \to \infty$, we have $\Upsilon_2(k) = 0$ for $k \ge \sqrt{3}$. The solution of (8.14) is

(8.23)
$$Y_{12}^{(6)} = C_2(t)e^{-tx + \frac{1}{3}x^3} + e^{-tx + \frac{1}{3}x^3} \int_{\sqrt{t}}^x e^{ts - \frac{1}{3}s^3} \left(s\operatorname{Ai}(s) - \operatorname{Ai}'(s)\right) ds.$$

By the consistence of (8.22) and (8.23) for $x = k\sqrt{t}$, we get

$$(8.24) C_2(t) - \Upsilon_2(k) = \int_{\infty}^{t} e^{\frac{k}{3}(3-k^2)s^{3/2}} \left(\frac{k^2-2}{2}\operatorname{Ai}(s) - \frac{k}{2\sqrt{s}}\operatorname{Ai}'(s)\right) ds$$
$$- \int_{\sqrt{t}}^{k\sqrt{t}} e^{ts-\frac{1}{3}s^3} \left(s\operatorname{Ai}(s) - \operatorname{Ai}'(s)\right) ds.$$

Unlike the case of $Y_{21}^{(6)}$, we can not recklessly take the $k \to 1$ limit of (8.24). Let us fix k, k > 1. Consider the asymptotics of (8.24) as $t \to \infty$. The first term of the right-side of (8.24) can be neglected since it is exponentially small for $t \to \infty$. So we have

$$C_{2}(t) - \Upsilon_{2}(k) \approx -\int_{\sqrt{t}}^{k\sqrt{t}} e^{ts - \frac{1}{3}s^{3}} (\operatorname{Ai}(t)s - \operatorname{Ai}'(t)) ds$$

$$= -\int_{0}^{k-1} e^{\frac{2}{3}t^{3/2} - t^{3/2}r^{2} - \frac{1}{3}t^{3/2}r^{3}} \left(\operatorname{Ai}(t)\sqrt{t}(1+r) - \operatorname{Ai}'(t) \right) \sqrt{t} dr$$

$$\approx -\int_{0}^{\infty} e^{-t^{3/2}r^{2}} \left(1 - \frac{1}{3}t^{3/2}r^{3} + \cdots \right) e^{\frac{2}{3}t^{3/2}} \left(\operatorname{Ai}(t)\sqrt{t}(1+r) - \operatorname{Ai}'(t) \right) \sqrt{t} dr$$

$$= -\frac{1}{2} - \frac{1}{12\sqrt{\pi}}t^{-\frac{3}{4}} + \frac{35}{1728\sqrt{\pi}}t^{-\frac{9}{4}} + \cdots$$
(8.25)

By the condition that $\Upsilon_2(k) = 0$ for $k \ge \sqrt{3}$, we get

(8.26)
$$\Upsilon_2(k) = 0, \quad k > 1,$$

$$C_2(t) = -\frac{1}{2} - \frac{1}{12\sqrt{\pi}} t^{-\frac{3}{4}} + \frac{35}{1728\sqrt{\pi}} t^{-\frac{9}{4}} + \cdots$$

Similarly, for 0 < k < 1, we have

(8.27)
$$C_2(t) - \Upsilon_2(k) = \frac{1}{2} - \frac{1}{12\sqrt{\pi}} t^{-\frac{3}{4}} + \frac{35}{1728\sqrt{\pi}} t^{-\frac{9}{4}} + \cdots$$

Therefore,

$$\Upsilon_2(k) = -1, \quad 0 < k < 1.$$

Thus,

(8.28)
$$\Upsilon_2(k) = \begin{cases} 0, & k > 1, \\ -1, & 0 < k < 1. \end{cases}$$

One should not try to get the expression of $C_2(t)$ by setting k = 1 in (8.24), since $\Upsilon_2(1)$ is not defined. We claim that

(8.29)
$$C_2(t) = -\frac{1}{2} - \frac{1}{2} \int_{\infty}^{t} e^{\frac{2}{3}s^{3/2}} \left(\operatorname{Ai}(s) + \frac{1}{\sqrt{s}} \operatorname{Ai}'(s) \right) ds.$$

Proof. Let $k_P = 1 + \epsilon - \frac{\epsilon^2}{6} + \frac{5}{72}\epsilon^3 + \cdots$ and $k_N = 1 - \epsilon - \frac{\epsilon^2}{6} - \frac{5}{72}\epsilon^3 + \cdots$ be the two roots of $\frac{1}{3}(3 - k^2)k = \frac{2}{3} - \epsilon^2$, $\epsilon > 0$. Then

$$C_{2}(t) + \frac{1}{2} = \frac{1}{2} \left(C_{2}(t) - \Upsilon_{2}(k_{P}) + C_{2}(t) - \Upsilon_{2}(k_{N}) \right)$$

$$= \int_{\infty}^{t} e^{\left(\frac{2}{3} - \epsilon^{2}\right)s^{3/2}} \left(-\frac{1}{2} \left(\operatorname{Ai}(s) + \frac{\operatorname{Ai}'(s)}{\sqrt{s}} \right) + \frac{1}{12} \epsilon^{2} \left(4\operatorname{Ai}(s) + \frac{\operatorname{Ai}'(s)}{\sqrt{s}} \right) + \cdots \right) ds$$

$$- \int_{\sqrt{t}}^{k_{P}\sqrt{t}} \cdots ds - \int_{\sqrt{t}}^{k_{N}\sqrt{t}} \cdots ds.$$

Since $4\text{Ai}(s) + \frac{\text{Ai}'(s)}{\sqrt{s}} = e^{-\frac{2}{3}s^{3/2}} \left(\frac{3}{2\sqrt{\pi}} s^{-\frac{1}{4}} - \frac{9}{32\sqrt{\pi}} s^{-\frac{7}{4}} + \cdots \right)$, we only need to show $\lim_{\epsilon \to 0} \epsilon^2 \int_{\infty}^t e^{-\epsilon^2 s^{3/2}} s^{-\frac{1}{4}} ds = 0$. But $\epsilon^2 \int_{\infty}^t e^{-\epsilon^2 s^{3/2}} s^{-\frac{1}{4}} ds = \frac{2}{3} \epsilon \sqrt{\pi} \left(\text{Erf}(\epsilon t^{\frac{3}{4}}) - 1 \right)$. Therefore, we set $\epsilon = \epsilon(t)$ smaller than t^{-n} for any $n > \frac{3}{4}$, for example, $\epsilon(t) = e^{-t}$. Then the terms $\int_{\sqrt{t}}^{k_P \sqrt{t}} \cdots ds$, $\int_{\sqrt{t}}^{k_N \sqrt{t}} \cdots ds$ and $\int_{\infty}^t \frac{1}{12} \epsilon^2 \left(4 \text{Ai}(s) + \frac{\text{Ai}'(s)}{\sqrt{s}} \right) ds$ can all be neglected. So (8.29) is obtained.

Remark 8.4. For $k \to \infty$, (8.22) with (8.28) is consistent with $Y_{12}^{(6)} \approx -\frac{u}{x}$.

Altogether, (8.22) with (8.28) is convenient for estimating $Y_{12}^{(6)}$ on $x = k\sqrt{t}$, $k \neq 1$, and (8.23) with (8.29) is proper for estimating $Y_{12}^{(6)}$ near $x = \sqrt{t}$.

8.2 The asymptotics of $Y^{(3)}(x,t)$ for $x\to -\infty$ and $t\to \infty$ along $x=k\sqrt{t}$

The behaviour of $Y^{(3)}$ are similar to $Y^{(6)}$ presented in Section 8.1.

In this case, k < 0. $x = -\sqrt{t}$ is the dividing line. By Lemma 8.1, we know $Y_{11}^{(3)} \to 1$ and $Y_{22}^{(3)} \to 1$. Therefore, the approximate differential equations for $Y_{21}^{(3)}$ and $Y_{12}^{(3)}$ along $x = k\sqrt{t}$ are

(8.30)
$$\frac{d}{dt}Y_{21}^{(3)} = \frac{(3-k^2)k}{2}\sqrt{t}Y_{21}^{(3)} + \frac{k^2-2}{2}\operatorname{Ai}(t) + \frac{k}{2\sqrt{t}}\operatorname{Ai}'(t),$$

(8.31)
$$\frac{d}{dt}Y_{12}^{(3)} = \frac{(k^2 - 3)k}{2}\sqrt{t}Y_{12}^{(3)} + \frac{k^2 - 2}{2}\operatorname{Ai}(t) - \frac{k}{2\sqrt{t}}\operatorname{Ai}'(t).$$

The approximate differential equations for $Y_{21}^{(3)}$ and $Y_{12}^{(3)}$ for fixed but large t are

(8.32)
$$\frac{d}{dx}Y_{21}^{(3)} = (t - x^2)Y_{21}^{(3)} + x\operatorname{Ai}(t) + \operatorname{Ai}'(t),$$

(8.33)
$$\frac{d}{dx}Y_{12}^{(3)} = (-t + x^2)Y_{12}^{(3)} + x\operatorname{Ai}(t) - \operatorname{Ai}'(t).$$

8.2.1 $Y_{12}^{(3)}$

The behaviour of $Y_{12}^{(3)}$ is similar to $Y_{21}^{(6)}$. Case -2 < k < 0:

In this case, the solution of (8.31) is

$$(8.34) Y_{21}^{(3)} = \Upsilon_3(k)e^{\frac{k}{3}(k^2-3)t^{3/2}} + e^{\frac{k}{3}(k^2-3)t^{3/2}} \int_{\infty}^{t} e^{\frac{k}{3}(3-k^2)s^{3/2}} \left(\frac{k^2-2}{2}\operatorname{Ai}(s) - \frac{k}{2\sqrt{s}}\operatorname{Ai}'(s)\right) ds.$$

The solution of (8.33) is

$$(8.35) Y_{12}^{(3)} = C_3(t)e^{-tx+\frac{1}{3}x^3} + e^{-tx+\frac{1}{3}x^3} \int_{-\sqrt{t}}^x e^{ts-\frac{1}{3}s^3} \left(s\operatorname{Ai}(t) - \operatorname{Ai}'(t)\right) ds.$$

So we have

$$(8.36) C_3(t) - \Upsilon_3(k) = \int_{\infty}^{t} e^{\frac{k}{3}(3-k^2)s^{3/2}} \left(\frac{k^2-2}{2}\operatorname{Ai}(s) - \frac{k}{2\sqrt{s}}\operatorname{Ai}'(s)\right) ds - \int_{-\sqrt{t}}^{k\sqrt{t}} e^{ts-\frac{1}{3}s^3} \left(s\operatorname{Ai}(t) - \operatorname{Ai}'(t)\right) ds.$$

By $\Upsilon_3(k) = 0$ for $k \in [-\sqrt{3}, 0)$, we get

(8.37)
$$C_3(t) = \int_{-\infty}^{t} e^{-\frac{2}{3}s^{3/2}} \left(-\frac{1}{2} \text{Ai}(s) + \frac{1}{2\sqrt{s}} \text{Ai}'(s) \right) ds.$$

Taking the $t \to \infty$ limit of (8.36) and considering (8.37), we obtain

$$\Upsilon_3(k) = 0, \quad k \in (-2, 0).$$

Therefore, the final result for -2 < k < 0 is

$$Y_{12}^{(3)} \approx e^{\frac{k}{3}(k^2 - 3)t^{3/2}} \int_{\infty}^{t} e^{\frac{k}{3}(3 - k^2)s^{3/2}} \left(\frac{k^2 - 2}{2} \operatorname{Ai}(s) - \frac{k}{2\sqrt{s}} \operatorname{Ai}'(s) \right) ds.$$

Case k = -2:

$$Y_{12}^{(3)} \approx e^{-\frac{2}{3}t^{3/2}} \int_{\infty}^{t} e^{\frac{2}{3}s^{3/2}} \left(\operatorname{Ai}(s) + \frac{1}{\sqrt{s}} \operatorname{Ai}'(s) \right) ds.$$

Case k < -2:

In this case, the solution of (8.31) is

$$(8.39) Y_{12}^{(3)} = \Upsilon_3(k)e^{\frac{k}{3}(k^2-3)t^{3/2}} + e^{\frac{k}{3}(k^2-3)t^{3/2}} \int_{t_A}^t e^{\frac{k}{3}(3-k^2)s^{3/2}} \left(\frac{k^2-2}{2}\operatorname{Ai}(s) - \frac{k}{2\sqrt{s}}\operatorname{Ai}'(s)\right) ds.$$

Thus, for k < -2, $Y_{12}^{(3)}$ is approximated by

$$(8.40) Y_{12}^{(3)} \approx e^{\frac{k}{3}(k^2 - 3)t^{3/2}} \int_{t_A}^{t} e^{\frac{k}{3}(3 - k^2)s^{3/2}} \left(\frac{k^2 - 2}{2} \operatorname{Ai}(s) - \frac{k}{2\sqrt{s}} \operatorname{Ai}'(s)\right) ds.$$

8.2.2 $Y_{21}^{(3)}$

For all k < 0, (8.30) has the solution

$$(8.41) Y_{21}^{(3)} = \Upsilon_4(k)e^{\frac{k}{3}(3-k^2)t^{3/2}} + e^{\frac{k}{3}(3-k^2)t^{3/2}} \int_{\infty}^{t} e^{\frac{k}{3}(k^2-3)s^{3/2}} \left(\frac{k^2-2}{2}\operatorname{Ai}(s) + \frac{k}{2\sqrt{s}}\operatorname{Ai}'(s)\right) ds.$$

The solution of (8.32) is

$$(8.42) Y_{21}^{(3)} = C_4(t)e^{tx-\frac{1}{3}x^3} + e^{tx-\frac{1}{3}x^3} \int_{-\sqrt{t}}^x e^{-ts+\frac{1}{3}s^3} \left(s\operatorname{Ai}(t) + \operatorname{Ai}'(t)\right) ds.$$

By the consistence of (8.41) and (8.42) for $x = k\sqrt{t}$, we get

$$(8.43) C_4(t) - \Upsilon_4(k) = \int_{\infty}^{t} e^{\frac{k}{3}(k^2 - 3)s^{3/2}} \left(\frac{k^2 - 2}{2} \operatorname{Ai}(s) + \frac{k}{2\sqrt{s}} \operatorname{Ai}'(s)\right) ds$$
$$- \int_{-\sqrt{t}}^{k\sqrt{t}} e^{-ts + \frac{1}{3}s^3} \left(s\operatorname{Ai}(t) + \operatorname{Ai}'(t)\right) ds.$$

Let us fix k, k < -1. Near $t = \infty$, the first term of the right-side of (8.43) can be neglected since it is exponentially small. Therefore, we obtain

$$C_{4}(t) - \Upsilon_{4}(k) \approx -\int_{-\sqrt{t}}^{k\sqrt{t}} e^{-ts + \frac{1}{3}s^{3}} (\operatorname{Ai}(t)s + \operatorname{Ai}'(t)) ds$$

$$= -\int_{0}^{k+1} e^{\frac{2}{3}t^{3/2} - t^{3/2}r^{2} + \frac{1}{3}t^{3/2}r^{3}} \left(\operatorname{Ai}(t)\sqrt{t}(-1+r) + \operatorname{Ai}'(t) \right) \sqrt{t} dr$$

$$\approx -\int_{0}^{-\infty} e^{-t^{3/2}r^{2}} \left(1 + \frac{1}{3}t^{3/2}r^{3} + \cdots \right) e^{\frac{2}{3}t^{3/2}} \left(\operatorname{Ai}(t)\sqrt{t}(r-1) + \operatorname{Ai}'(t) \right) \sqrt{t} dr$$

$$= -\frac{1}{2} - \frac{1}{12\sqrt{\pi}}t^{-\frac{3}{4}} + \frac{35}{1728\sqrt{\pi}}t^{-\frac{9}{4}} + \cdots$$
(8.44)

By the condition that $\Upsilon_4(k) = 0$ for $k \leq -\sqrt{3}$, we get

(8.45)
$$\Upsilon_4(k) = 0, \quad k < -1,$$

$$C_4(t) = -\frac{1}{2} - \frac{1}{12\sqrt{\pi}} t^{-\frac{3}{4}} + \frac{35}{1728\sqrt{\pi}} t^{-\frac{9}{4}} + \cdots$$

Similarly, for -1 < k < 0, we have

(8.46)
$$C_4(t) - \Upsilon_4(k) = \frac{1}{2} - \frac{1}{12\sqrt{\pi}} t^{-\frac{3}{4}} + \frac{35}{1728\sqrt{\pi}} t^{-\frac{9}{4}} + \cdots$$

Therefore,

$$\Upsilon_4(k) = -1, \quad -1 < k < 0.$$

Thus.

(8.47)
$$\Upsilon_4(k) = \begin{cases} 0, & k < -1, \\ -1, & -1 < k < 0. \end{cases}$$

We claim that

(8.48)
$$C_4(t) = -\frac{1}{2} - \frac{1}{2} \int_{\infty}^t e^{\frac{2}{3}s^{3/2}} \left(\operatorname{Ai}(s) + \frac{1}{\sqrt{s}} \operatorname{Ai}'(s) \right) ds.$$

The proof is similar to the case in Section 8.1, and thus we omit it.

9 Proof of Theorem 1.3

By (1.10), we only need to prove that $\mathcal{F}(x,t)$ satisfies the Bloemendal-Virág boundary (1.7) and that $\mathcal{F}(x,t)$ is bounded at $x^2 + t^2 = \infty$.




Figure 10. Diagram of the boundary $x^2 + t^2 = \infty$. \widehat{AB} : $x \to \infty$ and $t \to \infty$; \widehat{BC} : fixed x and $t \to \infty$; \widehat{CD} : $x \to -\infty$ and $t \to \infty$; \widehat{DE} : $x \to -\infty$ and fixed t; \widehat{EF} : $x \to -\infty$ and $t \to -\infty$; \widehat{FG} : fixed x and $t \to -\infty$; \widehat{GH} : $x \to \infty$ and $t \to -\infty$; \widehat{HA} : $x \to \infty$ and fixed t.

By considering $\mathcal{F}(x,t)$ on the boundary \widehat{AB} , [11] proved

$$\begin{pmatrix} \mathcal{F}_0(x,t) \\ \mathcal{G}_0(x,t) \end{pmatrix} = i \begin{pmatrix} \Psi_{012}^{(6)}(x,t) \\ \Psi_{022}^{(6)}(x,t) \end{pmatrix},$$

where \mathcal{F}_0 and \mathcal{G}_0 are defined by (2.3), and $\Psi_0^{(6)}$ is a canonical wave solution of Painlevé II. By (2.6), we get

(9.1)
$$\mathcal{F}(x,t) = \kappa u^{\frac{1}{2}} \left[u^{-1} \left(\frac{1+q_2}{2} x - \alpha \right) Y_{12}^{(6)}(x,t) + Y_{22}^{(6)}(x,t) \right].$$

Formula (9.1) is proper for $x \ge 0$.

The expression of $\mathcal{F}(x,t)$ for $x \leq 0$ has also been given by [11]

(9.2)
$$\mathcal{F}(x,t) = -\kappa u^{\frac{1}{2}} e^{\frac{x^3}{3} - xt} \left[u^{-1} \left(\frac{1 + q_2}{2} x - \alpha \right) Y_{11}^{(3)}(x,t) + Y_{21}^{(3)}(x,t) \right].$$

Note a = 1 has been applied to the expression in [11]. By (8.3) and (8.4), (9.1) and (9.2) coincide on x = 0.

9.1 Boundedness of $\mathcal{F}(x,t)$ to $c_1=0$ and $c_2=0$

Let us investigate $\mathcal{F}(x,t)$ at the boundary of \widehat{CD} , i.e., $x \to -\infty$ and $t \to \infty$ simultaneously. It is convenient to study the case along $x = k\sqrt{t}$, $k \in (-\sqrt{3},0)$. In this case, $e^{\frac{x^3}{3}-xt}$ is very large:

$$e^{\frac{x^3}{3} - xt} = e^{\frac{1}{3}k(k^2 - 3)t^{3/2}}.$$

The largest case is k = -1, i.e.,

$$e^{\frac{x^3}{3} - xt} = e^{\frac{2}{3}t^{3/2}}.$$

By (8.41), we know $-\kappa u^{\frac{1}{2}}e^{\frac{x^3}{3}-xt}Y_{21}^{(3)}(x,t)$ only contributes a finite term $-\Upsilon_4(k)$. Thus we can temporally neglect it.

Also we know for t > 0, $Y_{11}^{(3)}(x,t) \xrightarrow{x \to -\infty} 1$. Then by (2.9), we have

$$\lim_{x=k\sqrt{t},k\in(-\sqrt{3},0),t\to\infty}\mathcal{F}(x,t) = -\Upsilon_4(k) + \lim_{k\in(-\sqrt{3},0),t\to\infty} -e^{\frac{1}{3}k(k^2-3)t^{3/2}}u(t)^{-1}\left(k\sqrt{t}\frac{1+q_2(t)}{2} - \alpha(t)\right).$$

Since $\frac{1}{3}k(k^2-3)$ varies in $(0,\frac{2}{3}]$ for $k \in (-\sqrt{3},0)$ and $u(t)^{-1}$ has order of $e^{\frac{2}{3}t^{3/2}}$, we have to require $k\sqrt{t}\frac{1+q_2(t)}{2}-\alpha(t)$ has order of $e^{-\frac{4}{3}t^{3/2}}$. By (1.18) and (1.19), we find it is only possible for $c_1=c_2=0$.

9.2 $c_1 = c_2 = 0$ to boundedness of $F(\beta = 6; x, t)$ at $x^2 + t^2 = \infty$

9.2.1 On \widehat{HA}

In this case, t is fixed, $x \to \infty$,

$$\mathcal{F}(x,t) = \kappa u^{\frac{1}{2}} \left[u^{-1} \left(\frac{1+q_2}{2} x - \alpha \right) Y_{12}^{(6)}(x,t) + Y_{22}^{(6)}(x,t) \right].$$

Recall $Y_{12}^{(6)} \to -\frac{u(t)}{x}$ and $Y_{22}^{(6)} \to 1$ in this case, we get

(9.3)
$$\lim_{x \to \infty} \mathcal{F}(x,t) = -\kappa(t)u(t)^{\frac{1}{2}} \frac{q_2(t)+1}{2} + \kappa(t)u(t)^{\frac{1}{2}} = \frac{1}{2}\kappa(t)u(t)^{\frac{1}{2}} (1 - q_2(t)).$$

It is straightforward to verify that (9.3) is the same as (1.22).

9.2.2 On \widehat{AB}

$$\mathcal{F}(x,t) = \kappa u^{\frac{1}{2}} \left[u^{-1} \left(\frac{1+q_2}{2} x - \alpha \right) Y_{12}^{(6)}(x,t) + Y_{22}^{(6)}(x,t) \right].$$

By $Y_{22}^{(6)}(x,t) \to 1$ on \widehat{AB} , we know

(9.4)
$$\lim_{x \to \infty} \kappa(t) u(t)^{\frac{1}{2}} Y_{22}^{(6)}(x,t) = 1.$$

By $Y_{12}^{(6)}(x,t) \to 0$ on \widehat{AB} , we have

(9.5)
$$\lim_{x \to \infty, t \to \infty} -\kappa(t)u(t)^{-\frac{1}{2}}\alpha(t)Y_{12}^{(6)}(x,t) = 0.$$

Then we will show

(9.6)
$$\lim_{x \to \infty, t \to \infty} \kappa(t) u(t)^{-\frac{1}{2}} \frac{q_2(t) + 1}{2} x Y_{12}^{(6)}(x, t) = 0.$$

To prove (9.6), we divide the problem into 2 cases⁴: (1) $x \ge t$; (2) $x \le t$. In the case (1), by the Riemann-Hilbert problem of Y, it is easy to show $Y_{12}^{(6)} = o(1) \times x^{-1}$, and thus (9.6) is true. In the case (2),

$$\lim_{x \to \infty, t \to \infty} \left| \kappa(t) u(t)^{-\frac{1}{2}} \frac{q_2(t) + 1}{2} x Y_{12}^{(6)}(x, t) \right| \leq \lim_{x \to \infty, t \to \infty} \left| \kappa(t) u(t)^{-\frac{1}{2}} \frac{q_2(t) + 1}{2} \times t \times Y_{12}^{(6)}(x, t) \right|.$$

Considering $Y_{12}^{(6)} \to 0$ for x > 0, we know (9.6) is also true in this case. Thus (9.6) is proved. Gathering (9.4), (9.5) and (9.6), we get

$$\lim_{x \to \infty, t \to \infty} \mathcal{F}(x, t) = 1.$$

Remark 9.1. By Lemma 8.1, (8.22) and Remark 8.4, we know $c_1 = 0$ is enough to guarantee $\lim_{x \to \infty, t \to \infty} \mathcal{F}(x, t) = 1$ on \widehat{AB} .

9.2.3 On \widehat{BC}

Case $x \ge 0$.

$$\mathcal{F}(x,t) = \kappa u^{\frac{1}{2}} \left[u^{-1} \left(\frac{1+q_2}{2} x - \alpha \right) Y_{12}^{(6)}(x,t) + Y_{22}^{(6)}(x,t) \right].$$

In this case, x is finite and t is positive infinite. So, $\left|Y_{12}^{(6)}(x,t)\right| \leq 1$ and $Y_{22}^{(6)}(x,t) \to 1$. Therefore,

$$\lim_{t \to \infty} \mathcal{F}(x,t) = \lim_{t \to \infty} \kappa u^{\frac{1}{2}} \left[u^{-1} \left(\frac{1+q_2}{2} x - \alpha \right) Y_{12}^{(6)}(x,t) + Y_{22}^{(6)}(x,t) \right]$$
$$= 1, \quad x \geqslant 0.$$

⁴The division is at liberty. For example, for given $\epsilon > 0$, any division of $x \ge t^{\frac{1}{2} + \epsilon}$ and $x \le t^{\frac{1}{2} + \epsilon}$ works.

Case $x \leq 0$.

$$\mathcal{F}(x,t) = -\kappa(t)u(t)^{\frac{1}{2}}e^{\frac{x^3}{3}-xt}\left[u(t)^{-1}\left(\frac{1+q_2(t)}{2}x - \alpha(t)\right)Y_{11}^{(3)}(x,t) + Y_{21}^{(3)}(x,t)\right].$$

By Lemma 8.1, we know $Y_{11}^{(3)}(x,t) \approx 1$ and $Y_{21}^{(3)}(x,t) \approx -e^{xt-\frac{1}{3}x^3}$. So we get

$$\lim_{t \to \infty} \mathcal{F}(x,t) = \lim_{t \to \infty} -\kappa(t)u(t)^{\frac{1}{2}} e^{\frac{x^3}{3} - xt} \left[u(t)^{-1} \left(\frac{1 + q_2(t)}{2} x - \alpha(t) \right) Y_{11}^{(3)}(x,t) + Y_{21}^{(3)}(x,t) \right].$$

$$= 1, \quad x \le 0.$$

9.2.4 On \widehat{CD}

$$\mathcal{F}(x,t) = -\kappa(t)u(t)^{\frac{1}{2}}e^{\frac{x^3}{3}-xt}\left[u(t)^{-1}\left(\frac{1+q_2(t)}{2}x - \alpha(t)\right)Y_{11}^{(3)}(x,t) + Y_{21}^{(3)}(x,t)\right].$$

This is the most complicated case. The result is

$$\lim_{x \to -\infty, t \to \infty} \mathcal{F}(x, t) = \begin{cases} 0, & \frac{x + \sqrt{t}}{t^{-\frac{3}{4}}} \to -\infty, \\ 1, & \frac{x + \sqrt{t}}{t^{-\frac{3}{4}}} \to \infty, \\ \in (0, 1), & \text{near} x = -\sqrt{t}. \end{cases}$$

The corresponding proof is given in Appendix C.

On \widehat{DE} 9.2.5

$$\mathcal{F}(x,t) = -\kappa(t)u(t)^{\frac{1}{2}}e^{\frac{x^3}{3}-xt}\left[u(t)^{-1}\left(\frac{1+q_2(t)}{2}x - \alpha(t)\right)Y_{11}^{(3)}(x,t) + Y_{21}^{(3)}(x,t)\right].$$
 Since t is finite, $Y_{11}^{(3)}(x,t) \xrightarrow{x \to -\infty} 1$ and $Y_{21}^{(3)}(x,t) \xrightarrow{x \to -\infty} 0$, we obtain

$$\lim_{x \to -\infty} \mathcal{F}(x, t) = 0.$$

On \widehat{EF} 9.2.6

$$\mathcal{F}(x,t) = -\kappa(t)u(t)^{\frac{1}{2}}e^{\frac{x^3}{3}-xt}\left[u(t)^{-1}\left(\frac{1+q_2(t)}{2}x-\alpha(t)\right)Y_{11}^{(3)}(x,t)+Y_{21}^{(3)}(x,t)\right].$$

Let us first evaluate $Y_{11}^{(3)}(x,t)$ and $Y_{21}^{(3)}(x,t)$ along the curve $x=-\sqrt{2}\sqrt{\Lambda_1+t}$ for $t\in[-\Lambda_1,0]$ with

Along the curve,

$$(9.7) \qquad \frac{d}{dt} \begin{pmatrix} Y_{11}^{(3)}(t) \\ Y_{21}^{(3)}(t) \end{pmatrix} = \frac{1}{\sqrt{2}\sqrt{\Lambda_1 + t}} \begin{pmatrix} u(t)^2 & u'(t) \\ -u'(t) & -t - u(t)^2 \end{pmatrix} \begin{pmatrix} Y_{11}^{(3)}(t) \\ Y_{21}^{(3)}(t) \end{pmatrix}.$$

By (B.6), we get

$$\ln \sqrt{Y_{11}^{(3)}(0)^2 + Y_{21}^{(3)}(0)^2} \geqslant \ln \sqrt{Y_{11}^{(3)}(t)^2 + Y_{21}^{(3)}(t)^2} + \frac{1}{2} \int_t^0 \frac{-s - |s + 2u(s)^2|}{\sqrt{2}\sqrt{\Lambda_1 + s}} ds.$$

But
$$Y_{11}^{(3)}(0) = Y_{11}^{(3)}(x = -\sqrt{2}\sqrt{\Lambda_1}, t = 0) \approx 1$$
 and $Y_{21}^{(3)}(0) = Y_{21}^{(3)}(x = -\sqrt{2}\sqrt{\Lambda_1}, t = 0) \approx 0$. So
$$\sqrt{Y_{11}^{(3)}(t)^2 + Y_{21}^{(3)}(t)^2} < e^{\frac{1}{2}\int_0^t \frac{-s - |s + 2u(s)|^2}{\sqrt{2}\sqrt{\Lambda_1} + s}} ds.$$

Since $s + 2u(s)^2 \approx 0$ for large negative s, we assume $e^{\frac{1}{2}\int_0^t \frac{-s - |s + 2u(s)|^2}{\sqrt{2}\sqrt{\Lambda_1 + s}}ds} < 1$. Thus $\sqrt{Y_{11}^{(3)}(t)^2 + Y_{21}^{(3)}(t)^2} < 1$. 1 for large negative t. Also considering (1.16)-(1.17), (2.10) and (6.5), we immediately obtain

$$\lim_{x \to -\infty, t \to -\infty} \mathcal{F}(x, t) = 0.$$
28

9.2.7 On \widehat{FG}

Case $x \leq 0$.

$$\mathcal{F}(x,t) = -\kappa(t)u(t)^{\frac{1}{2}}e^{\frac{x^3}{3}-xt}\left[u(t)^{-1}\left(\frac{1+q_2(t)}{2}x - \alpha(t)\right)Y_{11}^{(3)}(x,t) + Y_{21}^{(3)}(x,t)\right].$$

$$\frac{dY_{11}^{(3)}(x,t)}{dt} = -u(t)Y_{21}^{(3)}(x,t),$$

$$\frac{dY_{21}^{(3)}(x,t)}{dt} = xY_{21}^{(3)}(x,t) - u(t)Y_{11}^{(3)}(x,t).$$

At $t = -\infty$, by (6.5), we know

$$Y_{11}^{(3)}(x,t) = C_1(x) \times \left(1 + \frac{x}{2\sqrt{2}(-t)^{1/2}} + \frac{x^2}{16t} - \frac{8\sqrt{2} + 9\sqrt{2}x^3}{192(-t)^{3/2}} + \cdots\right) \times e^{\frac{tx}{2} + \frac{1}{6}(x^2 - 2t)^{3/2}} + \cdots,$$

$$Y_{21}^{(3)}(x,t) = C_1(x) \times \left(1 - \frac{x}{2\sqrt{2}(-t)^{1/2}} + \frac{x^2}{16t} + \frac{-8\sqrt{2} + 9\sqrt{2}x^3}{192(-t)^{3/2}} + \cdots\right) \times e^{\frac{tx}{2} + \frac{1}{6}(x^2 - 2t)^{3/2}} + \cdots.$$

Also considering (1.16)-(1.17), (6.5) and (2.10), we finally get

$$\lim_{t \to -\infty} \mathcal{F}(x, t) = 0, \quad x \le 0.$$

which, actually, has been proved in Section 9.2.6.

Case $x \ge 0$.

$$\mathcal{F}(x,t) = \kappa u^{\frac{1}{2}} \left[u^{-1} \left(\frac{1+q_2}{2} x - \alpha \right) Y_{12}^{(6)}(x,t) + Y_{22}^{(6)}(x,t) \right].$$

$$\frac{dY_{12}^{(6)}(x,t)}{dt} = -xY_{12}^{(6)}(x,t) - u(t)Y_{22}^{(6)}(x,t),$$

$$\frac{dY_{22}^{(6)}(x,t)}{dt} = -u(t)Y_{12}^{(6)}(x,t).$$

Similar to the case x < 0, we get

$$Y_{12}^{(6)}(x,t) = C_2(x) \times \left(1 + \frac{x}{2\sqrt{2}(-t)^{1/2}} + \frac{x^2}{16t} - \frac{8\sqrt{2} + 9\sqrt{2}x^3}{192(-t)^{3/2}} + \cdots\right) \times e^{-\frac{tx}{2} + \frac{1}{6}(x^2 - 2t)^{3/2}} + \cdots,$$

$$Y_{22}^{(6)}(x,t) = C_2(x) \times \left(1 - \frac{x}{2\sqrt{2}(-t)^{1/2}} + \frac{x^2}{16t} + \frac{-8\sqrt{2} + 9\sqrt{2}x^3}{192(-t)^{3/2}} + \cdots\right) \times e^{-\frac{tx}{2} + \frac{1}{6}(x^2 - 2t)^{3/2}} + \cdots.$$

Therefore, we also have

$$\lim_{t \to -\infty} \mathcal{F}(x, t) = 0, \quad x \geqslant 0,$$

which can also be inferred from Section 9.2.8.

9.2.8 On \widehat{GH}

$$\mathcal{F}(x,t) = \kappa(t) \left(u(t)^{-\frac{1}{2}} \left(\frac{q_2(t) + 1}{2} x - \alpha(t) \right) Y_{12}^{(6)}(x,t) + u(t)^{\frac{1}{2}} Y_{22}^{(6)}(x,t) \right)$$

Along the curve $x=\sqrt{2}\sqrt{\Lambda_2+t},\,Y_{12}^{(6)}(x,t)$ and $Y_{22}^{(6)}(x,t)$ satisfy

$$(9.8) \qquad \frac{d}{dt} \left(\begin{array}{c} Y_{12}^{(6)}(t) \\ Y_{22}^{(6)}(t) \end{array} \right) = \frac{1}{\sqrt{2}\sqrt{\Lambda_2 + t}} \left(\begin{array}{cc} -\frac{1}{2}(t + 2u(t)^2) & -u'(t) \\ u'(t) & \frac{1}{2}(t + 2u(t)^2) \end{array} \right) \left(\begin{array}{c} Y_{12}^{(6)}(t) \\ Y_{22}^{(6)}(t) \end{array} \right).$$

By (B.5) and (B.6),

$$\ln \sqrt{Y_{12}^{(6)}(0)^2 + Y_{22}^{(6)}(0)^2} \le \ln \sqrt{Y_{12}^{(6)}(t)^2 + Y_{22}^{(6)}(t)^2} + \frac{1}{2} \int_t^0 \frac{|s + 2u(s)^2|}{\sqrt{2}\sqrt{\Lambda_2 + s}} ds,$$

$$\ln \sqrt{Y_{12}^{(6)}(0)^2 + Y_{22}^{(6)}(0)^2} \ge \ln \sqrt{Y_{12}^{(6)}(t)^2 + Y_{22}^{(6)}(t)^2} - \frac{1}{2} \int_t^0 \frac{|s + 2u(s)^2|}{\sqrt{2}\sqrt{\Lambda_2 + s}} ds.$$

Considering (6.5), for a large Λ_2 , we have

$$\int_{t}^{0} \frac{|s + 2u(s)^{2}|}{\sqrt{2}\sqrt{\Lambda_{2} + s}} ds \approx 0.$$

Also we know $Y_{12}^{(6)}(0) = Y_{12}^{(6)}(x = \sqrt{2}\sqrt{\Lambda_2}, t = 0) \approx 0$ and $Y_{22}^{(6)}(0) = Y_{22}^{(6)}(x = \sqrt{2}\sqrt{\Lambda_2}, t = 0) \approx 1$. Therefore,

$$(9.9) Y_{12}^{(6)}(t)^2 + Y_{22}^{(6)}(t)^2 \approx 1$$

on the curve $x = \sqrt{2}\sqrt{\Lambda_2 + t}$, $t \in [-\Lambda_2, 0]$. By (1.16)-(1.17), (2.10), (6.5) and (9.9), we obtain

$$\lim_{x \to \infty, t \to -\infty} \mathcal{F}(x, t) = \lim_{x \to \infty, t \to -\infty} \kappa(t) \left(u(t)^{-\frac{1}{2}} \left(\frac{q_2(t) + 1}{2} x - \alpha(t) \right) Y_{12}^{(6)}(x, t) + u(t)^{\frac{1}{2}} Y_{22}^{(6)}(x, t) \right)$$

$$= \lim_{x \to \infty, t \to -\infty} \frac{1}{2} \kappa(t) u(t)^{-\frac{1}{2}} x Y_{12}^{(6)}(x, t).$$

By (2.10) and (6.5), if $-t \gg (18 \ln \Lambda_2)^{\frac{1}{3}}$, we have $\mathcal{F}(t) \approx 0$.

Now let us prove when $-t < 2 \times (18 \ln \Lambda_2)^{\frac{1}{3}}$, $\mathcal{F}(t) \to 0$ as $t \to -\infty$.

By the mean value theorem,

$$\begin{split} Y_{12}^{(6)}(t) &= Y_{12}^{(6)}(0) + \left(Y_{12}^{(6)}\right)'(\xi)t \\ &= Y_{12}^{(6)}(2\sqrt{\Lambda_2}, 0) + \frac{1}{\sqrt{2}\sqrt{\Lambda_2 + \xi}} \left(-\frac{1}{2}(\xi + 2u(\xi)^2)Y_{12}^{(6)}(\xi) - u'(\xi)Y_{22}^{(6)}(\xi)\right)t, \end{split}$$

where $t < \xi < 0$.

From

$$Y_{12}^{(6)}(2\sqrt{\Lambda_2},0) \approx \frac{-u(0)}{\sqrt{2}\sqrt{\Lambda_2}}, \quad \frac{\sqrt{2}\sqrt{\Lambda_2+t}}{\sqrt{2}\sqrt{\Lambda_2+\xi}} < 1,$$

we know

$$\left|\kappa(t)u(t)^{-\frac{1}{2}}Y_{12}^{(6)}(t)x(t)\right| < \left|\kappa(t)u(t)^{-\frac{1}{2}}\right| \left(|u(0)| + \left|\frac{1}{2}(\xi + 2u(\xi)^2)Y_{12}^{(6)}(\xi) + u'(\xi)Y_{22}^{(6)}(\xi)\right| |t|\right).$$

By $t < \xi < 0$, (2.10), (6.5) and (9.9), we have

$$\lim_{t \to -\infty} \left| \kappa(t) u(t)^{-\frac{1}{2}} \right| \left(|u(0)| + \left| \frac{1}{2} (\xi + 2u(\xi)^2) Y_{12}^{(6)}(\xi) + u'(\xi) Y_{22}^{(6)}(\xi) \right| |t| \right) = 0.$$

Thus

$$\kappa(t) u(t)^{-\frac{1}{2}} Y_{12}^{(6)}(t) x(t) \approx 0,$$

when $t > -2 \times (18 \ln \Lambda_2)^{\frac{1}{3}}$ but large negative enough.

Altogether, we have

$$\lim_{x \to \infty, t \to -\infty} \mathcal{F}(x, t) = 0.$$

Appendix A $k_0 < \frac{10}{3}$

From $k_0 = -\min\left(\frac{t}{u^2}\right)$, we know

(A.1)
$$k_0 = 2 \times \left[\min \left(\frac{u}{\sqrt{\frac{-t}{2}}} \right) \right]^{-2}, \quad t \in (-\infty, 0).$$

Following the original arguments of [12], we give a lower bound for the local minimum of $\frac{u}{\sqrt{\frac{-t}{2}}}$ for large negative t.

Proposition A.1. If there is a local minimum of $\frac{u}{\sqrt{\frac{-t}{2}}}$ for $t < -\frac{11}{8}$, it must be greater than $\sqrt{\frac{1203}{1331}}$.

Proof. Let $u(t) = \sqrt{\frac{-t}{2}}z(t)$. Obviously, z(t) > 0. Then z satisfies $z''(t) + \frac{z'(t)}{t} = \left(\frac{1}{4t^2} - t\left(z(t)^2 - 1\right)\right)z(t)$. At a local minimum, we have u'(t) = 0 and u''(t) > 0. Then, we have $\frac{1}{4t^2} - t(z(t)^2 - 1) > 0$, i.e., $z(t) > \sqrt{1 + \frac{1}{4t^3}}$. Since $t < -\frac{11}{8}$, the local minimum is greater than $\sqrt{\frac{1203}{1331}}$

Note that Proposition A.1 does not mean $\frac{u}{\sqrt{\frac{-t}{2}}} > \sqrt{\frac{1203}{1331}}$ for $t \in (-\infty, -\frac{11}{8})$ since $\frac{u}{\sqrt{\frac{-t}{2}}}$ may be smaller near the boundary $t = -\frac{11}{8}$. But if we could also prove $\frac{u}{\sqrt{\frac{-t}{2}}} > \sqrt{\frac{1203}{1331}}$ for $t \in [-\frac{11}{8}, 0)$, then we can still conclude $\frac{u}{\sqrt{\frac{-t}{2}}} > \sqrt{\frac{1203}{1331}}$ for $t \in (-\infty, 0)$. The next proposition fulfills this aim.

Proposition A.2. For $t \in [-\frac{11}{8}, 0), \frac{u}{\sqrt{\frac{-t}{2}}} > \sqrt{\frac{1203}{1331}}$

Proof. Huang et. al [13] proved

(A.2)
$$\left| u(0) - \frac{98}{267} \right| < 11 \times 10^{-4}, \quad \left| u'(0) + \frac{153}{518} \right| < 12 \times 10^{-4}.$$

They also defined the approximate solution as

$$y_b(t) = \frac{t^{15}}{13206825} + \frac{t^{14}}{717099} + \frac{t^{13}}{81755} + \frac{t^{12}}{15201} + \frac{t^{11}}{47200} + \frac{13t^{10}}{24088} + \frac{39t^9}{53333} + \frac{18t^8}{61523}$$
(A.3)
$$-\frac{17t^7}{20578} - \frac{93t^6}{35396} - \frac{224t^5}{30615} - \frac{360t^4}{36911} + \frac{203t^3}{10806} + \frac{33530t^2}{688889} - \frac{153t}{518} + \frac{98}{267}$$

and the remainder term as

(A.4)
$$R_4(t) = y_b''(t) - ty_b(t) - 2y_b(t)^3.$$

We can verify ⁵

(A.5)
$$|R_4(t)| < 2 \times 10^{-3}, \quad t \in \left[-\frac{11}{8}, 0\right].$$

Let $\delta_4(t) = u(t) - y_b(t)$. It is easy to show

$$\delta_4''(t) = (6y_b(t)^2 + t)\delta_4(t) + 6y_b(t)\delta_4(t)^2 + 2\delta_4(t)^3 - R_4(t).$$

Next, we will show $\delta_4(t)$ is sufficiently small for $t \in [-\frac{11}{8}, 0]$. We can verify $\frac{4}{5} < 6y_b(t)^2 + t < \frac{13}{5}$ and $\frac{11}{5} < 6y_b(t) < \frac{49}{10}$ for $t \in [-\frac{11}{8}, 0]$. Therefore, we have $\delta_4(t) \ge \delta_b(t)$ in the interval, where $\delta_b(t)$ is defined by

(A.6)
$$\delta_b''(t) = \frac{13}{5}\delta_b(t) - \frac{49}{10}\delta_b(t)^2 + 2\delta_b(t)^3 - \frac{1}{500}, \quad \delta_b(0) = -\frac{11}{10000}, \quad \delta_b'(0) = \frac{3}{2500}.$$

So we have

(A.7)
$$\delta_b'(t) = \sqrt{\frac{13}{5}\delta_b(t)^2 - \frac{49}{15}\delta_b(t)^3 + \delta_b(t)^4 - \frac{1}{250}\delta_b(t) - \frac{183310481923}{3} \times 10^{-16}}, \quad \delta_b(0) = -\frac{11}{10000}$$

By

$$\int_{-11\times10^{-4}}^{-120\times10^{-4}} \left(\frac{13}{5} \delta_b^2 - \frac{49}{15} \delta_b^3 + \delta_b^4 - \frac{1}{250} \delta_b - \frac{183310481923}{3} \times 10^{-16} \right)^{-\frac{1}{2}} d\delta_b < -\frac{11}{8},$$

we obtain $\delta_b(t) > -120 \times 10^{-4} = -\frac{3}{250}$ for $t \in [-\frac{11}{8}, 0]$. When $t \in [-\frac{11}{8}, 0]$, we can show $\frac{y_b(t) - \frac{3}{250}}{\sqrt{\frac{-t}{25}}} > \sqrt{\frac{1203}{1331}}$. Therefore, for $t \in [-\frac{11}{8}, 0)$, we have

$$\min\left(\frac{u(t)}{\sqrt{\frac{-t}{2}}}\right) = \min\left(\frac{y_b(t) + \delta_4(t)}{\sqrt{\frac{-t}{2}}}\right) > \min\left(\frac{y_b(t) + \delta_b(t)}{\sqrt{\frac{-t}{2}}}\right) > \min\left(\frac{y_b(t) - \frac{3}{250}}{\sqrt{\frac{-t}{2}}}\right) > \sqrt{\frac{1203}{1331}}$$

Combining Proposition A.1 and A.2, we obtain $\min \left(\frac{u(t)}{\sqrt{-t}} \right) > \sqrt{\frac{1203}{1331}}$. By (A.1), $k_0 < \frac{2662}{1203}$ is

 $^{^{5}}$ Since $R_{4}(t)$ is a polynomial, Sturm's theorem applies. The following several cases of verification can also be done in

Appendix B The growth rate estimate for the solution of a second order linear ODE

For a > 0, c > 0, $b^2 - 4ac < 0$ and $V = (x, y)^T$, let us define

$$|V| = \sqrt{ax^2 + bxy + cy^2},$$

where |V| can be understood as the length of vector V. Denote

$$M = \left(\begin{array}{cc} m_{11} & m_{12} \\ m_{21} & m_{22} \end{array} \right),$$

then

$$r^2 = \frac{|MV|^2}{|V|^2}$$

has both a minimal value $\mathcal{M}_1 = r_1^2$ and a maximal value $\mathcal{M}_2 = r_2^2$. These two extreme values satisfy

(B.1)
$$\Delta \mathcal{M}^2 + 2\delta \mathcal{M} + \Delta D^2 = 0,$$

where

$$\Delta = b^2 - 4ac,$$

$$D = \det M,$$

$$\delta = -b^2(m_{11}m_{22} + m_{12}m_{21}) - 2b(am_{12} - cm_{21})(m_{11} - m_{22}) + 2a^2m_{12}^2 + 2c^2m_{21}^2 + 2ac(m_{11}^2 + m_{22}^2).$$

Assume the linear ODE is of form

(B.2)
$$\frac{d}{dt} \begin{pmatrix} x(t) \\ y(t) \end{pmatrix} = \begin{pmatrix} f_{11}(t) & f_{12}(t) \\ f_{21}(t) & f_{22}(t) \end{pmatrix} \begin{pmatrix} x(t) \\ y(t) \end{pmatrix}.$$

We know

$$\begin{pmatrix} x(t_1) \\ y(t_1) \end{pmatrix} = \lim_{N \to \infty} \begin{pmatrix} 1 + hf_{11}(t_0 + (N-1)h) & hf_{12}(t_0 + (N-1)h) \\ hf_{21}(t_0 + (N-1)h) & 1 + hf_{22}(t_0 + (N-1)h) \end{pmatrix} \cdots \\ \cdots \begin{pmatrix} 1 + hf_{11}(t_0 + h) & hf_{12}(t_0 + h) \\ hf_{21}(t_0 + h) & 1 + hf_{22}(t_0 + h) \end{pmatrix} \begin{pmatrix} 1 + hf_{11}(t_0) & hf_{12}(t_0) \\ hf_{21}(t_0) & 1 + hf_{22}(t_0) \end{pmatrix} \begin{pmatrix} x(t_0) \\ y(t_0) \end{pmatrix},$$

where $h = \frac{t_1 - t_0}{N}$. Then by (B.1), for $t_1 > t_0$, we can prove

(B.3)
$$\ln|(x(t_1), y(t_1))^T| \leq \ln|(x(t_0), y(t_0))^T| + \int_{t_0}^{t_1} \left(\frac{1}{2}(f_{11}(s) + f_{22}(s)) + \frac{\sqrt{H(s)}}{\sqrt{-\Delta}}\right) ds,$$

(B.4)
$$\ln|(x(t_1), y(t_1))^T| \ge \ln|(x(t_0), y(t_0))^T| + \int_{t_0}^{t_1} \left(\frac{1}{2}(f_{11}(s) + f_{22}(s)) - \frac{\sqrt{H(s)}}{\sqrt{-\Delta}}\right) ds,$$

where

$$H(s) = ac(f_{11}(s) - f_{22}(s))^2 - b(af_{12}(s) - cf_{21}(s))(f_{11}(s) - f_{22}(s)) + (af_{12}(s) + cf_{21}(s))^2 - b^2f_{12}(s)f_{21}(s).$$
If $a = c = 1$, $b = 0$, for $t_1 > t_0$, (B.3) and (B.4) are reduced to

$$\ln |(x(t_1), y(t_1))^T| \leq \frac{1}{2} \int_{t_0}^{t_1} \left(f_{11}(s) + f_{22}(s) + \sqrt{(f_{11}(s) - f_{22}(s))^2 + (f_{12}(s) + f_{21}(s))^2} \right) ds$$
(B.5)
$$+ \ln |(x(t_0), y(t_0))^T|,$$

$$\ln |(x(t_1), y(t_1))^T| \geq \frac{1}{2} \int_{t_0}^{t_1} \left(f_{11}(s) + f_{22}(s) - \sqrt{(f_{11}(s) - f_{22}(s))^2 + (f_{12}(s) + f_{21}(s))^2} \right) ds$$
(B.6)
$$+ \ln |(x(t_0), y(t_0))^T|.$$

Appendix C $\mathcal{F}(x,t)$ on \widehat{CD}

$$\mathcal{F}(x,t) = -\kappa(t)u(t)^{\frac{1}{2}}e^{\frac{x^3}{3}-xt}\left[u(t)^{-1}\left(\frac{1+q_2(t)}{2}x - \alpha(t)\right)Y_{11}^{(3)}(x,t) + Y_{21}^{(3)}(x,t)\right].$$

We divide the region \widehat{CD} into two parts: $x \leq -\sqrt{3}\sqrt{t}$ and $x \geq -\sqrt{3}\sqrt{t}$.

C.1 Case $x \leq -\sqrt{3}\sqrt{t}$.

In this case, $Y_{21}^{(3)}(x,t) \rightarrow 0$ and $e^{\frac{x^3}{3}-xt} \leqslant 1$, so

$$\lim_{x \le -\sqrt{3}\sqrt{t}, t \to \infty} -\kappa(t)u(t)^{\frac{1}{2}}e^{\frac{x^3}{3}-xt}Y_{21}^{(3)}(x,t) = 0.$$

By $Y_{11}^{(3)}(x,t) \to 1$, we know

$$\lim_{x \leqslant -\sqrt{3}\sqrt{t}, t \to \infty} \kappa(t) u(t)^{\frac{1}{2}} e^{\frac{x^3}{3} - xt} u(t)^{-1} \alpha(t) Y_{11}^{(3)}(x, t) = 0.$$

Thus

$$\lim_{x \leq -\sqrt{3}\sqrt{t}, t \to \infty} \mathcal{F}(x, t) = \lim_{x \leq -\sqrt{3}\sqrt{t}, t \to \infty} -\kappa(t)u(t)^{\frac{1}{2}} e^{\frac{x^3}{3} - xt} u(t)^{-1} \frac{q_2(t) + 1}{2} x Y_{11}^{(3)}(x, t)$$
(C.1)
$$= \lim_{x \leq -\sqrt{3}\sqrt{t}, t \to \infty} x e^{\frac{x^3}{3} - xt} u(t)^{-1} \frac{q_2(t) + 1}{2}.$$

(C.1) can be proved to be 0 by diving the region $x \leqslant -\sqrt{3}\sqrt{t}$ into two parts, for example $x \leqslant -2\sqrt{t}$ and $-2\sqrt{t} \leqslant x \leqslant -\sqrt{3}\sqrt{t}$. In both parts, $xe^{\frac{x^3}{3}-xt}u(t)^{-1}\frac{q_2(t)+1}{2} \to 0$ is obvious.

Therefore, we have

$$\lim_{x \le -\sqrt{3}\sqrt{t}, t \to \infty} \mathcal{F}(x, t) = 0.$$

C.2 Case $x \ge -\sqrt{3}\sqrt{t}$.

First we show

(C.2)
$$\lim_{x \to -\infty, t \to \infty, x \geqslant -\sqrt{3}\sqrt{t}} -\kappa(t)u(t)^{\frac{1}{2}}e^{\frac{x^3}{3}-xt}u(t)^{-1}\left(\frac{1+q_2(t)}{2}x - \alpha(t)\right)Y_{11}^{(3)}(x,t) = 0.$$

In fact, by (1.20) and (1.21), we obtain their expansions at $t = \infty$ as

(C.3)
$$q_2(t) + 1 = e^{-\frac{4}{3}t^{3/2}} \left(\frac{1}{8\pi} t^{-\frac{3}{2}} - \frac{59}{192\pi} t^{-3} + \cdots \right),$$

(C.4)
$$\alpha(t) = e^{-\frac{4}{3}t^{3/2}} \left(\frac{3}{16\pi} t^{-1} - \frac{29}{128\pi} t^{-\frac{5}{2}} + \cdots \right).$$

Also we have

$$u(t)^{-1} \approx \frac{1}{\operatorname{Ai}(t)} = e^{\frac{2}{3}t^{3/2}} \left(2\sqrt{\pi}t^{\frac{1}{4}} + \frac{5}{24}\sqrt{\pi}t^{-\frac{5}{4}} + \cdots \right).$$

Therefore,

$$\lim_{x \to -\infty, t \to \infty, x \geqslant -\sqrt{3}\sqrt{t}} -\kappa(t)u(t)^{\frac{1}{2}} e^{\frac{x^3}{3} - xt} u(t)^{-1} \left(\frac{1 + q_2(t)}{2}x - \alpha(t)\right) Y_{11}^{(3)}(x, t)$$

$$= \lim_{x \to -\infty, t \to \infty, x \geqslant -\sqrt{3}\sqrt{t}} -e^{\frac{x^3}{3} - xt} u(t)^{-1} \left(\frac{1 + q_2(t)}{2}x - \alpha(t)\right)$$

$$= \lim_{x \to -\infty, t \to \infty, x \geqslant -\sqrt{3}\sqrt{t}} -e^{\frac{x^3}{3} - xt - \frac{2}{3}t^{3/2}} \left(\frac{1}{4\sqrt{\pi}}t^{-\frac{5}{4}}x - \frac{3}{8\sqrt{t}}t^{-\frac{3}{4}}\right).$$

But $\frac{x^3}{3} - xt - \frac{2}{3}t^{\frac{3}{2}} \le 0$ in this case and also $|x| \le \sqrt{3}t^{\frac{1}{2}}$. So we get (C.2). By (8.41), we have

(C.5)
$$\lim_{x=k\sqrt{t}, k<0, k\neq -1, t\to\infty} -\kappa(t)u(t)^{\frac{1}{2}}e^{\frac{x^3}{3}-xt}Y_{21}^{(3)}(x,t) = -\Upsilon_4(k).$$

Near $x = -\sqrt{t}$, by (8.42), we get

$$(C.6) \qquad \kappa(t)u(t)^{\frac{1}{2}}e^{\frac{x^3}{3}-xt}Y_{21}^{(3)}(x,t)$$

$$\approx \frac{1}{2} + \frac{1}{2} \int_{\infty}^{t} e^{\frac{2}{3}s^{3/2}} \left(\operatorname{Ai}(s) + \frac{1}{\sqrt{s}} \operatorname{Ai}'(s) \right) ds - \int_{-\sqrt{t}}^{x} e^{-ts + \frac{1}{3}s^3} \left(s\operatorname{Ai}(t) + \operatorname{Ai}'(t) \right) ds$$

(C.7)
$$\approx \frac{1}{2} - \int_0^{\frac{x}{\sqrt{t}}+1} e^{\frac{2}{3}t^{3/2} - t^{3/2}r^2 + \frac{1}{3}t^{3/2}r^3} \left(\operatorname{Ai}(t)\sqrt{t}(r-1) + \operatorname{Ai}'(t) \right) \sqrt{t} dr.$$

By (C.6), we see $-\kappa(t)u(t)^{\frac{1}{2}}e^{\frac{x^3}{3}-xt}Y_{21}^{(3)}(x,t)$ is monotonic increasing with x. So it must lie in (0,1). (C.7) is convenient for estimating its value.

C.2.1 Case $k\sqrt{t} < x \le 0, k > -1$.

(8.33) is valid for a large positive t. So does (8.35). Along the line that t is fixed, by (8.35) we get

(C.8)
$$\frac{d}{dx} \left(-e^{\frac{x^3}{3} - xt} Y_{21}^{(3)}(x, t) \right) = -e^{-tx + \frac{1}{3}x^3} \left(x \operatorname{Ai}(t) + \operatorname{Ai}'(t) \right).$$

As

$$\frac{\text{Ai}'(t)}{\text{Ai}(t)} = -\sqrt{t} - \frac{1}{4t} + \frac{t}{32}t^{-\frac{5}{2}} + \cdots,$$

(C.8) never vanishes in the region. Thus $-e^{\frac{x^3}{3}-xt}Y_{21}^{(3)}(x,t)$ is monotonic increasing in the region. But it is known

$$-e^{\frac{x^3}{3}-xt}Y_{21}^{(3)}(x,t)|_{x=k\sqrt{t}} \approx 1, \quad k > -1,$$

and

$$-e^{\frac{x^3}{3}-xt}Y_{21}^{(3)}(x,t)|_{x=0} \approx \mathcal{F}(x,t)|_{x=0} \approx 1.$$

We must conclude

$$\lim_{k\sqrt{t} < x \le 0, k > -1, t \to \infty} \mathcal{F}(x, t) = 1.$$

Acknowledgement. Part of this work was done while Y. Li was visiting the Department of Mathematical Sciences of IUPUI. Y. Li would like to thank A. Its for his hospitality, encouragement and suggestions. The work is partly supported by NSFC(11375090, 11675054, 11435005) and Shanghai Collaborative Innovation Center of Trustworthy Software for Internet of Things (ZF1213).

References

- [1] Baik, J.; Buckingham, R.; DiFranco, J. Asymptotics of Tracy-Widom distributions and the total integral of a Painlevé II function. *Comm. Math. Phys.* 280 (2008), no. 2, 463-497.
- [2] A. Bloemendal; B. Virág. Limits of spiked random matrices I. *Probab. Theory Related Fields* 156(2013), 795-825.
- [3] Borot, G.; Eynard, B.; Majumdar, S. N.; Nadal, C. Large deviations of the maximal eigenvalue of random matrices. J. Stat. Mech. Theory Exp. (2011), no.11, P11024.
- [4] Bourgade, P; Erdős, L; Yau, H. T. Universality of general β -ensembles. Duke Math. J. 163(2014), no 6, 1127-1190.
- [5] Deift, P.; Its, A.; Krasovsky, I. Asymptotics of the Airy-kernel determinant. Comm. Math. Phys. 278 (2008), no. 3, 643-678.
- [6] Deift P.; Zhou X. A steepest descent method for oscillatory RiemannHilbert problems, asymptotics for the MKdV equation, *Ann. of Math.* 137 (1993), 295-368.

- [7] Dumitriu I.; Edelman A. Matrix models for beta ensembles, J. Math. Phys. 43(2002), 5830-5847.
- [8] Dijkgraaf, R.; Vafa, C. Toda theories, matrix models ,topological strings and N=2 gauge systems. arXiv:0909.2453.
- [9] Flaschka H.; Newell A.C. Monodromy- and spectrum-preserving deformations. I, Comm. Math. Phys. 76(1980), 65116.
- [10] Foka A.; Its A.; Kapaev A.; Novokshenov V. Painleve Transcendents: The Riemann-Hilbert Approach. AMS, 2006.
- [11] Grava T.; Its A.; Kapaev A.; Mezzadri F. On the Tracy-Widom_{β} distribution for $\beta = 6$, SIGMA 12(2016), 105, 26 pages.
- [12] Hastings S. P.; McLeod J.B. A boundary value problem associated with the second Painlevé transcendent and the Kortewegde Vries equation, *Arch. Rational Mech. Anal.*73 (1980), 3151.
- [13] Huang M.; Xu S.X.; Zhang L. Location of poles for the HastingsMcLeod solution to the second Painlevé equation, *Constr. Approx.*43(2016), 463494.
- [14] Jorba A.; Zou M. A software package for the numerical integration of ODEs by Means of high-order Taylor methods, *Experimental Mathematics*, 14:1(2005), 99-117.
- [15] Mehta M. L. Random matrices, Pure and Applied Mathematics (Amsterdam), Vol. 142, 3rd ed., Elsevier/Academic Press, Amsterdam, 2004.
- [16] Ramírez, J. A.; Rider, B.; Virág, B. Beta ensembles, stochastic Airy spectrum, and a diffusion. J. Amer. Math. Soc. 24 (2011), no. 4, 919-944.
- [17] Rumanov I. Classical integrability for beta-ensembles and general Fokker-Plank equations. *J. Math. Phys.* 56(2015), 013508.
- [18] Rumanov I. Painlevé representation of Tracy-Widom_{β} distribution for $\beta = 6$. Comm. Math. Phys. 342(2016), 843-868.
- [19] Tracy, C. A.; Widom, H. Level-spacing distributions and the Airy kernel, Comm. Math. Phys.159(1994), (1): 151-174.
- [20] Tracy, C. A.; Widom, H. On orthogonal and symplectic matrix ensembles. Comm. Math. Phys. 177 (1996), no. 3, 727-754.
- [21] Wasow W. Asymptotic expansions for ordinary differential equations, Pure and Applied Mathematics, Vol. 14, Interscience Publishers John Wiley & Sons, Inc., New York London Sydney, 1965.

YUQI Li

Institute of Computer Theory, School of Computer Science and Software Engineering East China Normal University

Shanghai, 200062

China

E-mail: yqli@sei.ecnu.edu.cn