

GRAPH CALCULUS AND THE DISCONNECTED-BOUNDARY SCHWINGER-DYSON EQUATIONS OF QUARTIC TENSOR FIELD THEORIES

CARLOS I. PÉREZ-SÁNCHEZ

*Mathematisches Institut der Westfälischen Wilhelms-Universität
Einsteinstraße 62, 48149 Münster, Germany*
&

Faculty of Physics, University of Warsaw
ul. Pasteura 5, 02-093 Warsaw, Poland*

ABSTRACT. Tensor field theory (TFT) focuses on quantum field theory aspects of random tensor models, a quantum-gravity-motivated generalisation of random matrix models. The TFT correlation functions have been shown to be classified by graphs that describe the geometry of the boundary states, the so-called boundary graphs. These graphs can be disconnected, although the correlation functions are themselves connected. In a recent work, the Schwinger-Dyson equations for an arbitrary albeit connected boundary were obtained. Here, we introduce the multivariable graph calculus in order to derive the missing equations for all correlation functions with disconnected boundary, thus completing the Schwinger-Dyson pyramid for quartic melonic (‘pillow’-vertices) models in arbitrary rank. We first study finite group actions that are parametrised by graphs and build the graph calculus on a suitable quotient of the monoid algebra $\mathcal{A}[G]$ corresponding to a certain function space \mathcal{A} and to the free monoid G in finitely many graph variables; a derivative of an element of $\mathcal{A}[G]$ with respect to a graph yields its corresponding group action on \mathcal{A} . The present result and the graph calculus have three potential applications: the non-perturbative large- N limit of tensor field theories, the solvability of the theory by using methods that generalise the topological recursion to the TFT setting and the study of ‘higher dimensional maps’ via Tutte-like equations. In fact, we also offer a term-by-term comparison between Tutte equations and the present Schwinger-Dyson equations.

E-mail address: cperez@fuw.edu.pl.

Date: June 2019.

1991 Mathematics Subject Classification. Primary 81Txx; Secondary 20Nxx, 05Exx.

Key words and phrases. Tensor Models, Quantum Gravity, Quantum Field Theory, Schwinger-Dyson equations, Matrix Models, Random Maps, Algebras and Rings, Tutte Equations.

* Current affiliation.

CONTENTS

1. Introduction and motivation	2
2. Graph calculus	6
2.1. Single variable graph calculus	6
2.2. Examples of graph-group action systems	9
2.3. Multivariable graph calculus	11
2.4. Algebraic structure	12
2.5. Three limit cases and examples	18
3. Tensor models	21
3.1. Tensor Field Theory	21
3.2. From graph calculus to tensor field theory	22
4. Disconnected-boundary Schwinger-Dyson equations	25
4.1. The quartic melonic tensor field theory	25
4.2. Main result	26
5. Four and six point SDE with disconnected boundary	34
5.1. Schwinger-Dyson equations for $G_{\ominus i\boxplus i}^{(6)}$	34
5.2. Schwinger-Dyson equation for $G_{\ominus \ominus}^{(4)}$	37
5.3. Schwinger-Dyson equation for $G_{\ominus \ominus \ominus}^{(6)}$	37
6. Towards higher-dimensional Tutte equations	38
6.1. Tutte equations and matrix models	38
6.2. Parallel between Tutte equations and disconnected- ∂ SDE of TFT	39
7. Conclusions and outlook	44
Acknowledgements	45
Appendix A. The first coefficients of the Y -term	45
References	46

1. INTRODUCTION AND MOTIVATION

The quest for laws of physics near the Planck scale leads some quantum gravitologist and quantum cosmologists to replace the smooth space-time paradigm with new geometrical structures that are suitable for said energy scale. Those new structures include discretisation of space-time (e.g. causal dynamical triangulations [AGJL13]), the algebraisation of space-time (e.g. noncommutative geometry [CC97, Mar18]), just to name

some¹. Already the sole description of a space-time by a single mathematical object is expected to require therefore novel geometrical ideas.

If one adopts a path-integral approach, the exploration of the quantum theory of space-time requires additionally a multi-geometry description, to which ‘off-shell’ geometries also contribute. Each of these geometries ξ is weighted via $\exp(iS(\xi)/\hbar)d\xi$ by a ‘classical’ action $S(\xi)$, bounded to resemble the Einstein-Hilbert action in the classical limit, in which ξ starts looking like a Riemannian or Lorentzian manifold.

Random tensors [ADJ91, GR12, Riv16] and related theories construct these kinds of measures $d\xi$, offer precisely a built-in description of both *random* and *discrete* geometry in arbitrary dimensions, and therefore constitute a tool to test models of background independent quantum gravity (see e.g. [EKLP18]). Interest in the study of Euclidean quantum field theory (QFT) aspects of random tensors leads to *tensor field theory* (TFT) [BGR13, BGS13, OV14, RVT], the matter of this article. Usually TFT fits in a ‘QFT + ϵ ’ framework, that is to say a conservative modification of QFT, which one can pursue in the perturbative or non-perturbative approaches.

Non-perturbative TFT deals with the geometry of boundary states. Single geometries in TFT are represented by certain decorated graphs called *coloured graphs*; this decoration is precisely the information that allows the construction of PL-manifolds from graphs. Bulk-geometry graphs —Feynman diagrams of TFT— have a colour more than the graphs that triangulate the boundary geometries, which are called therefore *boundary graphs* (∂ -graphs, for short). The two-fold purpose of this article is to define in abstract way a calculus with coloured boundary-graph variables, and shortly afterwards, to apply this construct to a particular problem in non-perturbative TFT.

The single-variable graph calculus has been used there as a toolkit for non-perturbative field tensor theory, leading to the full Ward-Takahashi identity [Pér18]. The single-variable graph calculus allows to define each correlation function as a graph derivative of the free energy. Boundary graphs turn out to classify the correlation functions of tensor field theories; these obey analytic² Schwinger-Dyson Equation (SDE).

Each and every analytic SDE for a *connected* correlation function corresponding to a *connected*, but otherwise arbitrary, boundary graph was presented in [PPW17] in terms of a general formula that relates a given correlation function with its neighbourings (relative to the number of points) in terms of simple graph operations. In order to obtain these results, the single-variable graph calculus was a useful tool, which however does not assist any longer in the the derivation of SDE for connected correlation functions with disconnected boundary. This derivation requires a multivariable graph calculus, the variables being the different boundary components.

¹See for instance [MT18] for a thorough classification.

²We write ‘analytic’ as opposed to algebraic SDE for expectation values. We conceive tensor field theory as a discretisation (therefore, 0-dimensional) of a D -dimensional quantum field theory. The $2k$ -point correlation functions are thus functions of $\mathbb{Z}^{k \times D} \rightarrow \mathbb{C}$ which in the continuum limit pass to functions $\mathbb{R}^{k \times D} \rightarrow \mathbb{C}$, and render the SDE integro-differential equations.




FIGURE 1. An example of a -coloured edge swap increasing the genus of a connected graph

We introduce graph-group actions as the basis of the multivariable calculus, and study their generating functionals. Concretely, we obtain formulae for the graph derivative of products of functionals, i.e. the corresponding Leibniz rule that generalises

$$\partial^\alpha(F \cdot G) = \sum_{0 \leq \gamma \leq \alpha} \binom{\alpha}{\gamma} \partial^\gamma F \cdot \partial^{\alpha-\gamma} G \quad (\text{in multi-index notation}), \quad (1.1)$$

when one replaces usual partial derivatives ∂_μ (say, with x^μ a coordinate of \mathbb{R}^n , and F, G real-valued smooth functions there) by derivatives $\partial_g = \partial/\partial g$ with respect to a graph g . The formula for graphs takes a different form, but reduces, as it should, to (1.1) when one replaces functionals with functions and simultaneously considers trivial group actions.

We prove that this abstract structure underlies tensor models functionals and use it to find a general formula for the SDE of the quartic ‘pillow’-model, for the *connected* correlation functions with arbitrary *disconnected boundary* (abbr. disconnected- ∂). This is the missing piece that complements the connected-boundary SDE-pyramid obtained in [PPW17]. To have it complete is important for the analysis of the non-perturbative large- N limit of tensor field theories. Moreover, although it is not clear which recursion should generalise the topological recursion [Eyn14], it is clear that the disconnected- ∂ correlation functions play an important role³.

Taking graph-derivatives can be understood as the tensor model counterpart of ‘taking residues’ in matrix models (cf. [Eyn16, Ch. 1-2])

$$\mathcal{T}_{l_1 \dots l_\kappa} = \text{Res}_{x_1 \rightarrow \infty} \dots \text{Res}_{x_\kappa \rightarrow \infty} [x_1^{l_1} \dots x_\kappa^{l_\kappa} W_\kappa(x_1, \dots, x_\kappa)]$$

in order to project the free energy W_κ (with κ boundaries) of a matrix model onto the generating function $\mathcal{T}_{l_1 \dots l_\kappa}$ of random maps with κ marked faces with fixed perimeter lengths (l_1, \dots, l_κ) . After the well-known equivalence [Eyn16, Thm. 2.5.1] between Tutte equations [Tut62, Tut63] for the enumeration of random maps and the loop equations for a suitable matrix model [Mig83] (also summarised here in in Sect. 6.2), we can state the

³For instance, higher dimensional analogue of the ‘pair of pants’ being represented by a correlation function with three melonic boundary components.

present result as the basis to obtain Tutte-like equations for the higher-dimensional analogue of the generating function of random maps $\mathcal{T}_{l_1 \dots l_\kappa}$. This improvement over [PPW17]—where only the equations analogous to a single-boundary correlator, say \mathcal{T}_{l_1} 's cousin, are presented—requires new developments: A straightforward generalisation of the proof given in [PPW17] is impossible due to the absence of the multi-variable graph calculus. Although it would be possible to state the main result solely in a TFT context, the full notation would be a burden in the proof. We state some results in lighter notation and offer a shorter proof, at expenses of introducing some new concepts. The output of the main theorem is also a set of new graph operations that extend those used to describe the connected- ∂ SDE in [PPW17]. Also the *edge swap*⁴ ς_a (see Fig. 1), as unary operation on a connected graph, is extended to a binary operation implying two connected components; ς_a is then interpreted, following [Pér17], as their connected sum.

This article is divided in an abstract part (Sect. 2, whose main results are lemmas 2.10 and 2.12), and a TFT-part. The next section explains in detail the following mnemonics: for coloured graphs g and h ,

$$\frac{\partial g}{\partial h} = \delta(h, g) \cdot \text{group action of } G(g). \quad (1.2)$$

Here $\delta(g, h) = 1$ if the graphs g and h are isomorphic or otherwise $\delta(g, h)$ vanishes, and $G(g)$ is a group determined by g . In Section 3 we make the connection between the two first sections and model-independent TFT. In Section 4, our quartic model is detailed and the results of the previous sections are applied to the main problem, namely to find the SDE for *connected* correlation functions with arbitrary *disconnected boundary* with arbitrary number of connected components. Section 5 gives explicitly some of the SDE for 4, 6-point functions for rank-3 theories. We highlight in Section 6, preparing an important future task to apply our results, the term-to-term parallel between Tutte equations and those presented here, as well as analogies in the derivation of both sets of equations. Concretely, we compare the new operations on the boundary graphs of TFT (Table 5) with their matrix models counterpart (Table 4). The operations and terms presented in the SDE of [PPW17] are only the counterpart of those matrix models SDE presented in Table 3. The conclusions and outlook are given in Section 7, discussing a potential application related to the higher-dimensional analogue of the topological recursion. The useful coefficients that encode the insertion of the 4-point functions into the 2-point function and the 6-point functions into the 4-point functions are given in the Appendix A.

⁴ This operation has been studied in the literature of Graph-Encoded Manifolds [LM06] and by the Crystallisation Theory [CC15]

2. GRAPH CALCULUS

In this section we explain what we mean by graph calculus. For naturality reasons⁵ we consider the empty graph \emptyset as coloured and add it to the set of (possibly) disconnected, closed, regularly edge- D -coloured, vertex-bipartite graphs (' D -coloured graphs') $\text{Grph}_D^{\text{II,cl}}$, to form $\mathbf{G}_D = \{\emptyset\} \cup \text{Grph}_D^{\text{II,cl}}$. Henceforth, all graphs are coloured, but other types of graphs could be used for the next constructs.

2.1. Single variable graph calculus. We regard \mathbf{G}_D with a monoidal structure, the product and the unit being given by

$$c_1 c_2 = c_1 \amalg c_2 \quad \text{and} \quad c \amalg \emptyset = c = \emptyset \amalg c,$$

respectively, for all $c, c_1, c_2 \in \mathbf{G}_D$. We choose to remember the order of the factors, so this product is generally non-commutative, $c_1 c_2 \neq c_2 c_1$.

Definition 2.1 (System of graph-group actions). For a finite collection $H \subset \mathbf{G}_D$, consider the following structures:

- for each connected graph $c \in H$:
 - a set $\mathcal{V}(c)$ is associated with c ; for the empty graph, $\mathcal{V}(\emptyset) = \{*\}$ is the singleton
 - a finite group $G(c)$ and
 - a group action $G(c) \curvearrowright \mathcal{V}(c)$ of $G(c)$ on $\mathcal{V}(c)$
- if $g = c_1 c_2 \cdots c_n$ is a factorisation in connected components c_i , then $\mathcal{V}(g)$ satisfies⁶

$$\mathcal{V}(g) = \mathcal{V}(c_1) \amalg \cdots \amalg \mathcal{V}(c_n) \tag{2.1}$$

The collection $\{\mathcal{V}(g), G(g)\}_{g \in H}$ is a *system of graph-group actions*.

If additionally, for each graph in $g \in H$, one has functions

$$u_g : \mathcal{V}(g) \rightarrow \mathbb{C} \text{ (or } \mathbb{R}),$$

then one says that $\{u_g, \mathcal{V}(g), G(g)\}_{g \in H}$, or more succinctly, $\{u_g\}_{g \in H}$ is a family of functions *supported on* $\{\mathcal{V}(g), G(g)\}_{g \in H}$.

We are interested in triples $\{u_g, \mathcal{V}(g), G(g)\}_{g \in H}$ and formal sums of the type $U = \sum_{h \in H} u_h h$, which we refer to as their *generating functionals*. In this case we say that (the set of graphs) H *spans* U . Why these are functionals instead of functions will become apparent while addressing the applications. At this point also the following terminology, inherited from the physical significance, might seem mysterious: we call the elements of $\mathcal{V}(g)$ the *momenta* of the graph g . Notice that u_\emptyset is a constant.

As the last reference to TFT in this section, we clarify that the nature of these graphs is not important at this point; examples will be presented in later sections. We just clarify the reader that is well-versed with tensor models, that graphs treated here are

⁵This is in line with other theories (as a matter of fact Topological Quantum Field Theories) that need the empty D -manifold. Also, it is technically advantageous and not the first time it is considered, see for instance [KT16].

⁶One could relax this condition so that there exist domains $\mathcal{U}(g)$ of $\mathcal{V}(g)$ compatible with the $G(g)$ -action and such that $\mathcal{U}(g) \subset \mathcal{U}(c_1) \amalg \cdots \amalg \mathcal{U}(c_n)$. However, we keep the natural condition (2.1)

not Feynman graphs, but boundary graphs of these in a rank- D TFT. In this context, functions u_g are unknown⁷, and one derives equations that they should satisfy. Only after knowing solutions we would be able to fix a function space u_g should belong to, which is for now unspecified. We vaguely refer then to them as ‘functions’.

Next, some words on notation. For a factorisation $g = c_1 \cdots c_n$ in connected graphs $c_j \neq \emptyset$, we let g/c_i be the graph with the i -th connected component deleted,

$$g/c_i := c_1 \cdots \widehat{c_i} \cdots c_n = c_1 \cdots c_{i-1} c_{i+1} \cdots c_n.$$

Notice that this deletion does not only care about the graph-class, but also about its spot in the factorisation, which we can keep track of thanks to the monoidal structure of \mathbf{G}_D .

For $g = c_1 \cdots c_n$ as before, let $Y \in \mathcal{V}(c_r)$, $1 \leq r \leq n$. Given a function $v_g : \mathcal{V}(g) \rightarrow \mathbb{C}$ we define the insertion of Y in the r -th argument of v_g

$$\iota_Y^r v_g : \mathcal{V}(c_1) \amalg \cdots \amalg \mathcal{V}(c_{r-1}) \amalg \mathcal{V}(c_{r+1}) \amalg \cdots \amalg \mathcal{V}(c_n) \rightarrow \mathbb{C},$$

using (2.1) by

$$(\iota_Y^r v_g)(X_1, \dots, X_{r-1}, X_{r+1}, \dots, X_n) = v_g(X_1, \dots, X_{r-1}, Y, X_{r+1}, \dots, X_n),$$

where $X_i \in \mathcal{V}(c_i)$ for each $i \neq r$.

Definition 2.2. Let $H \subset \mathbf{G}_D$ span the functional $U = \sum_g u_g g$. Given any connected graph $h \in \mathbf{G}_D$, $h \neq \emptyset$, and an arbitrary graph $g = c_1 \cdots c_n$ factorised in connected components c_i , we define $I(g, h) = \{i \in \llbracket 1, n \rrbracket \mid h = c_i\}$; that is, $I(g, h)$ is the subset of numbers that indexes the factors of g that coincide with h . For $r \in I(g, h)$, we label by $h^{(r)}$ the appearance of h in the r -th factor of $g = c_1 \cdots c_n$. We define the *functional graph derivative* with respect to h (evaluated at X) as the functional

$$\frac{\delta U}{\delta h(X)} = \sum_{g \in H} \sum_{r \in I(g, h)} \sum_{\sigma \in G(h)} \iota_{\sigma(X)}^r u_g (g/h^{(r)}), \quad X \in \mathcal{U}(h) \subset \mathcal{V}(h).$$

The well-definedness follows from condition (2.1), which implies that in each case $\iota_{\sigma(X)}^r u_g$ is indeed a function on $\mathcal{V}(g/h^{(r)})$. We stress that this derivative could be defined in a proper domain $\mathcal{U}(h)$ of $\mathcal{V}(h)$. Further, if h occurs nowhere as a factor of g the sum is empty, and thus $\delta U / \delta h \equiv 0$. The derivative with respect to \emptyset is the coefficient of that graph, $\delta U / \delta \emptyset := u_\emptyset \in \mathbb{C}$.

To clarify this definition, consider a monomial functional, $U = u_g g$, with $g = h^n$ for some integer $n \geq 1$ and h a connected graph. By definition, one has

$$\frac{\delta U}{\delta h(X)} = \sum_{r=1}^n \sum_{\sigma \in G(h)} (\iota_{\sigma(X)}^r u_{h^n}) h^{n-1},$$

⁷For instance, u_g can be the correlation functions.

which one can rethink as

$$\begin{aligned} \frac{\delta(h^n)}{\delta h} &= \left(\frac{\delta h}{\delta h} h^{n-1} + h \frac{\delta h}{\delta h} h^{n-2} + \dots + h^{n-1} \frac{\delta h}{\delta h} \right) \\ &= \left(G(h) h^{n-1} + h G(h) h^{n-2} + \dots + h^{n-1} G(h) \right), \end{aligned} \quad (2.2)$$

if

$$\left(\frac{\delta h}{\delta h} u_h \right)(X) = \sum_{\sigma \in G(h)} (\sigma \cdot u_h)(X) = \sum_{\sigma \in G(h)} u_h(\sigma X). \quad (2.3)$$

To illustrate the action in slightly more generality, if in $u_{c_1 \dots c_{r-1} h c_{r+1} \dots}$ none of the c_j is isomorphic to h , then

$$\left(\frac{\delta h}{\delta h} u_{c_1 \dots c_{r-1} h c_{r+1} \dots} \right)(X) = \sum_{\sigma \in G(h)} \iota_{\sigma(X)}^r u_{c_1 \dots c_{r-1} h c_{r+1} \dots}. \quad (2.4)$$

On this account, the useful symbolism to keep in mind is that the derivative of a graph with respect to itself is the group action of $G(h)$ on $\mathcal{V}(h)$,

$$\frac{\delta h}{\delta h} = G(h) \curvearrowright \{\text{functions } \mathcal{V}(h) \rightarrow \mathbb{C}\}. \quad (2.5)$$

In the sequel, we will often abuse on notation and write this equality without the curved action-arrow, as we already did above in eq. (2.2). The moral is that each factor h occurring in a term of the type

$$u_{c_1 \dots c_{r-1} h c_{r+1} \dots c_n}(c_1 \dots c_{r-1} h c_{r+1} \dots c_n)$$

is a potential $G(h)$ -orbit of the r -th argument $u_{c_1 \dots c_{r-1} h c_{r+1} \dots c_n}$.

For iterated derivatives with respect to h , one can see by induction that for $n \geq 2$, the iteration of n graph derivatives applied to h^n yields

$$\frac{\delta^n(h^n)}{\delta h \delta h \dots \delta h}(X_1, \dots, X_n) = \sum_{\mu \in \mathfrak{S}(n)} \sum_{(\sigma^1, \dots, \sigma^n) \in G(h)^n} \iota_{\sigma^{\mu(1)}(X_1)}^1 \dots \iota_{\sigma^{\mu(n)}(X_n)}^n. \quad (2.6)$$

If $G_i(h)$ is the i -th factor of the group $G(h)^n$, a more transparent notation of last equation is

$$\frac{\delta^n(h^n)}{\delta h \delta h \dots \delta h} = \sum_{\mu \in \mathfrak{S}(n)} G_{\mu(1)}(h) G_{\mu(2)}(h) \dots G_{\mu(n)}(h), \quad (2.7)$$

where the group $G_{\mu(i)}(h)$ acts on the i -th factor of the set $\mathcal{V}(h) \amalg \dots \amalg \mathcal{V}(h)$. The group corresponding to the n -th derivative of the n -th power of a graph h with respect to itself is

$$\frac{\delta^n(h^n)}{\delta h \delta h \dots \delta h} = G(h) \wr \mathfrak{S}(n). \quad (2.8)$$

In this case, the wreath product $G(h) \wr \mathfrak{S}(n)$ is the semi-direct product $G(h)^n \rtimes_{\psi} \mathfrak{S}(n)$, with the obvious action ψ of $\mathfrak{S}(n)$ on the n copies of $G(h)$.

To give further detail, given a generating system of graph-group actions $\{\mathcal{V}(g), G(g)\}_{g \in H}$ and $h \in H$, consider a function $F : \mathcal{V}(h) \amalg \dots \amalg \mathcal{V}(h) \rightarrow \mathbb{C}$. An element $\Omega = (\sigma; \mu) = (\sigma^1, \dots, \sigma^n; \mu)$ of the group in eq. (2.7) acts as follows:

$$(\Omega \cdot F)(X_1, \dots, X_n) = F(\sigma^1(X_{\mu(1)}), \dots, \sigma^n(X_{\mu(n)})). \quad (2.9)$$

By departing from eq. (2.9), the composition with another element $\Xi = (\tau, \nu)$ in the group (2.7) is easily proven to yield $\Xi \circ \Omega = (\tau\psi_\nu(\sigma); \nu\mu)$, where

$$\psi : \mathfrak{S}(n) \rightarrow \text{Aut}(G(h)^n) \quad \mu \mapsto [\psi_\mu : (\sigma_i)_{i=1}^n \mapsto (\sigma_{\mu(i)})_{i=1}^n],$$

which is the product of $G(h)^n \rtimes_\psi \mathfrak{S}(n)$, as claimed.

2.2. Examples of graph-group action systems. Roughly stated, a multivariable graph calculus (of n graph variables) consists of generating functionals of functions u_g supported on a system of graph-group actions $\{\mathcal{V}(g), G(g)\}_{g \in H}$ that are spanned by a finite set H . We take $H \subset \text{FM}(\{h_1, \dots, h_n\})$, the free monoid generated⁸ by n non-isomorphic graphs $\mathfrak{h} = \{h_1, \dots, h_n\}$. For a multivariable graph calculus the key property is that the graph-group actions $G(h_i)$ are pairwise independent, that is for each $i, j = 1, \dots, n$,

$$\frac{\delta h_i}{\delta h_j} = \delta_j^i G(h_i), \quad h_i, h_j \in \mathfrak{h}. \quad (2.10)$$

For the special element $g = h_1^{\alpha_1} \dots h_n^{\alpha_n}$ the restriction imposed by eq. (2.10) implies

$$\frac{\delta g}{\delta g} = G(g) = G(h_1) \wr \mathfrak{S}(\alpha_1) \times G(h_2) \wr \mathfrak{S}(\alpha_2) \times \dots \times G(h_n) \wr \mathfrak{S}(\alpha_n). \quad (2.11)$$

Before formally defining multivariable graph calculus, the next examples are just meant to illustrate last action (2.11), rather than the role of the graphs in graph-generated actions, and therefore can be skipped (to Sect. 2.3).

Example. Let $\zeta_n \neq 1$ denote a n -th root of unit ($n \geq 2$), and consider the system of graph-group actions with a single graph g . Let $G(g)$ be the group spanned by $\zeta_n \mathcal{V}(g) = \mathbb{C}$ by multiplication. Then the functional graph derivative of g with respect to itself on the identity $\text{id}_{\mathbb{C}}$ vanishes identically:

$$\left(\frac{\delta g}{\delta g} \right) \text{id}_{\mathbb{C}} \equiv 0.$$

The $G(g)$ -orbit of the function $f_n : z \mapsto z^n$ yields

$$\left(\frac{\delta g}{\delta g(z)} \right) (f_n) = n \cdot z^n.$$

Example. Consider a finite set $H \subset \text{FM}(\mathfrak{h})$ and the following graph-group actions system

$$G(g) = \mathfrak{S}(|g^0|), \quad \mathcal{V}(g) = M_{|g^0| \times |g^0|}(\mathbb{R}).$$

Here $|g^0|$ is the number of vertices of g . The action of the symmetric group on the matrices permutes columns (or rows). Then the orbit of the determinant $\det : \mathcal{V}(g) \rightarrow \mathbb{R}$ vanishes

⁸We recall that the *free monoid* generated by $\mathfrak{h} = \{h_1, \dots, h_n\}$ is in this case the following set $\text{FM}(\mathfrak{h}) = \{l_1 \dots l_m : m \in \mathbb{Z}_{\geq 0} \text{ and } l_i \in \mathfrak{h}\}$ endowed with the concatenation operation; containing the empty graph, i.e. the empty word.

identically. This follows from considering, for an arbitrary matrix $X = (X_{ab}) \in \mathcal{V}(g)$,

$$\begin{aligned} \left(\frac{\delta g}{\delta g(X)} \right) \det(\cdot) &= \sum_{\sigma \in \mathfrak{S}(|g^0|)} \det(X_{\sigma(a)b}) \\ &= \sum_{\sigma \in A} \det(X_{\sigma(a)b}) + \sum_{\sigma \in \mathfrak{S}(|g^0|) - A} \det(X_{\sigma(a)b}) \\ &= \sum_{\sigma \in A} \det(X_{ab}) - \sum_{\sigma \in \mathfrak{S}(|g^0|) - A} \det(X_{ab}) = 0, \end{aligned}$$

where A is the alternating subgroup; its complement in the symmetric group consists of odd-degree permutations, whence the common minus sign in the last line. Both have the same order, which explains why the sum vanishes independently of X .

Example. Let K be a finite group that accepts (cf. [Sze16] for a criterion) faithful irreducible representations. Consider n of them $\pi^i : K \rightarrow \text{End}(W_i)$, and set $G(h_i) = K$ for each $i = 1, \dots, n$. Define for each i the momenta of h_i as the matrix space $\mathcal{V}(h_i) = \pi^i(K)$ (since K is finite, irreps are finite-dimensional). The group K acts on $\mathcal{V}(h_i)$ by

$$\mathcal{V}(h_i) \ni \pi^i(m) \xrightarrow{u} \pi^i(u)\pi^i(m) = \pi^i(um), \quad (m, u \in K).$$

Consider the following functions f^i defined in terms of the characters $\chi^i(u) = \text{Tr}(\pi^i(u))$, $u \in K$,

$$f^i : \mathcal{V}(h_i) \rightarrow \mathbb{C}, \quad f^i(X_i) = \chi^i(m)^* \chi^i(m), \quad (X_i = \pi^i(m)).$$

Then for $X_i = \pi^i(m)$, the following holds:

$$\begin{aligned} \left(\frac{\delta h_i}{\delta h_i(X_i)} \right) f^i &= \sum_{u \in K} (u \cdot f^i)(\pi^i(m)) \quad (i \text{ is fixed}) \\ &= \sum_{u \in K} \chi^i(um)^* \chi^i(um) \\ &= \sum_{u \in K} \chi^i(u)^* \chi^i(u) = |K|. \end{aligned}$$

Fix any $g \in \mathfrak{h}$ and let π be the associated representation. Define for any $h_i, h_j \in \mathfrak{h}$, and $X = \pi(m)$,

$$F^{ij} : \mathcal{V}(g) \rightarrow \mathbb{C}, \quad F^{ij}(X) = \chi^i(m)^* \chi^j(m).$$

For $X = \pi(m)$,

$$\left(\frac{\delta g}{\delta g(X)} \right) F^{ij} = \sum_{u \in K} \chi^i(um)^* \chi^j(um) = \sum_{u' \in K} \chi^i(u')^* \chi^j(u') = \delta_{ij} |K|.$$

Last equality is due to Schur orthogonality.

Example. Let $D \in \mathbb{Z}_{\geq 1}$. Let two non-isomorphic graphs $H = \{g, h\} \subset \mathbf{G}_D$ parametrise the system of graph-group actions given by $\{\mathcal{V}(l), G(l)\}_{l \in H}$, being

$$\begin{array}{lll} \mu \in G(g) = \mathfrak{S}(D) & Z = (z_i)_i \in \mathcal{V}(g) = \mathbb{C}^D & \mu : (z_i) \mapsto (z_{\mu(i)}) \\ \tau \in G(h) = \mathbb{Z}_2 & \varepsilon \in \mathcal{V}(h) = \mathbb{Z}_2 & \varepsilon : s \mapsto \tau \varepsilon \end{array}$$

where \mathbb{Z}_2 is written multiplicatively $\{-1, 1\}$. Let $F : \mathcal{V}(g) \amalg \mathcal{V}(g) \amalg \mathcal{V}(h) \rightarrow \mathbb{C}$ be given by, say,

$$F(Z, W, \varepsilon) = \frac{\varepsilon}{2(D!)^2} e^{-|W| + \varepsilon|Z|} =: c(D) \cdot \varepsilon \cdot e^{-|W| + \varepsilon|Z|}.$$

Then the functional graph derivative of $g^2 f$ with respect to itself yields the following group-orbit, when applied to F :

$$\begin{aligned}
 & \frac{\delta^3(g^2 h)}{\delta g(Z_1) \delta g(Z_2) \delta h(\varepsilon)} F \\
 &= \left(\frac{\delta^3 g g h}{\delta g \delta g \delta h} F \right) (Z_1, Z_2, \varepsilon) \\
 &= \left([G(g) \wr \mathfrak{S}(2) \times G(h)] \cdot F \right) (Z_1, Z_2, \varepsilon) \\
 &= \sum_{\mu \in \mathfrak{S}(2)} \sum_{\tau \in \mathbb{Z}_2} \sum_{\sigma \in \mathfrak{S}(D)} \sum_{\rho \in \mathfrak{S}(D)} F(\sigma(Z_{\mu(1)}), \rho(Z_{\mu(2)}), \tau \varepsilon) \\
 &= \sum_{\tau \in \mathbb{Z}_2} \sum_{\sigma \in \mathfrak{S}(D)} \sum_{\rho \in \mathfrak{S}(D)} \left\{ F(\sigma(Z_1), \rho(Z_2), \tau \varepsilon) + F(\sigma(Z_2), \rho(Z_1), \tau \varepsilon) \right\} \\
 &= 2c(D) \cdot (D!)^2 \cdot \varepsilon \cdot (e^{\varepsilon|Z_1|-|Z_2|} + e^{\varepsilon|Z_2|-|Z_1|} \\
 &\quad - e^{-\varepsilon|Z_1|-|Z_2|} - e^{-\varepsilon|Z_2|-|Z_1|}) \\
 &= \varepsilon (e^{-|Z_1|} \sinh |Z_2| + e^{-|Z_2|} \sinh |Z_1|).
 \end{aligned}$$

We have used the invariance under the action of two copies $\mathfrak{S}(D)$, which contributed a factor $(D!)^2$.

2.3. Multivariable graph calculus. Let $\mathfrak{h} = \{h_1, \dots, h_n\} \subset \mathbf{G}_D$ be a set of connected, non-isomorphic graphs. For the basis of the multivariable calculus the free monoid $\text{FM}(\{h_1, \dots, h_n\})$ is too ‘verbose’, and not each one of its elements has the ordered form $h_1^{\alpha_1} \dots h_n^{\alpha_n}$. This could in principle be solved by taking the free commutative monoid $\text{FM}^{\text{ab}}(\mathfrak{h})$ instead, which, however, turns out to be overly restrictive (for our aims). A mild compromise between these two alternatives —the free monoid and its abelianisation— is to allow to permute letters in an arbitrary word, as to make use of the action (2.11), and then in some sense undo the changes. Next definition introduces precisely such reordering.

Definition 2.3. Given a finite set of graphs $\mathfrak{h} = \{h_1, \dots, h_n\}$, the *degree* $|g|$ of an element g in $\text{FM}(\mathfrak{h})$ is the number of factors of g , i.e. the number of connected components g consists of. We let $\mathfrak{S}(|g|)$ act by permuting the factors of g , $g \mapsto \sigma(g)$; notice that $\mathfrak{S}(|g|)$ left-acts naturally as $\sigma \cdot f = f \circ \sigma^{-1}$ on functions $f : \mathcal{V}(g) \rightarrow \mathbb{C}$. Given a family of functions $\{u_g\}_{g \in H}$ supported on a system of graph-group actions $\{\mathcal{V}(g), G(g)\}_{g \in H \subset \text{FM}(\mathfrak{h})}$, and given a $g \in H$, we declare the pairs $(u_g, g) \sim (\sigma \cdot u_g, \sigma(g))$ equivalent for each $\sigma \in \mathfrak{S}(|g|)$. The notation we choose for this equivalence, called *reordering*, is

$$u_g g \sim u_h h \quad \text{if and only if} \quad u_h = \sigma \cdot u_g \text{ and } h = \sigma(g) \text{ for certain } \sigma \in \mathfrak{S}(|g|). \quad (2.12)$$

Definition 2.4. Given a finite set of connected non-isomorphic graphs $\mathfrak{h} = \{h_1, \dots, h_n\} \subset \mathbf{G}_D$, a system of graph-group actions $\mathcal{S} = \{\mathcal{V}(h), G(h)\}_{h \in \mathfrak{h}}$ is said to be *independent* if eq. (2.10) holds. When the context is clear, we just say that ‘ \mathfrak{h} is independent’, or that \mathcal{S} is.

Definition 2.5. A *multivariable graph calculus* $\mathcal{C}(\mathfrak{h})$ or a *graph calculus* with variables \mathfrak{h} consists of two objects:

- the choice of an independent system of graph-group actions $\{\mathcal{V}(h), G(h)\}_{h \in \mathfrak{h}}$ for a finite set $\mathfrak{h} \subset \mathbf{G}_D$ and
- the set of finite formal sums in elements of $g \in \text{FM}(\mathfrak{h})$ having each of these a function of the form $v_g : \mathcal{V}(g) \rightarrow \mathbb{C}$ as coefficient, modulo reordering. That is,

$$\mathcal{C}(\mathfrak{h}) = \left\{ \sum_g v_g g \mid v_g \equiv 0 \text{ for almost all } g \in \text{FM}(\mathfrak{h}) \right\} / \sim \quad (2.13)$$

where \sim is the linear extension of relation (2.12), abusing on the same symbol.

2.4. Algebraic structure. We now explore the structure of a graph calculus $\mathcal{C}(\mathfrak{h})$ with variables $\mathfrak{h} = \{h_1, \dots, h_n\}$. The elements of $\mathcal{C}(\mathfrak{h})$, called also *functionals*, have a non-unique representation, since $\sum_g v_g g = \sum_{\tilde{g}} v_{\tilde{g}} \tilde{g}$ where $\tilde{g} = \tau_g(g)$ and $v_{\tilde{g}} = \tau_g \cdot (v_g)$ for an arbitrary $\tau_g \in \mathfrak{S}(|g|)$. For sake of computability, it will be helpful to be able to fix representing elements g that span a functional, and subordinate the order of the arguments of the coefficient-functions to that choice.

We write $g \stackrel{\sim}{\sim} h$ for any $g, h \in \text{FM}(\mathfrak{h})$ if $g = h$ in the free commutative monoid $\text{FM}^{\text{ab}}(\mathfrak{h})$ spanned by \mathfrak{h} . In other words, $g \stackrel{\sim}{\sim} h$ if and only if h and g match in $\text{FM}(\mathfrak{h})$ up to a rearranging $\sigma \in \mathfrak{S}(|g|)$, i.e. if $\sigma(g) = h$.

Definition 2.6. Given a family of functions $\{v_l\}_l$ supported on $\{\mathcal{V}(l), G(l)\}_{l \in H}$ where $H \subset \text{FM}(\mathfrak{h})$. Let $g, h \in H$ be such that $h \stackrel{\sim}{\sim} g$. We define for the *reordering of a function* $v_h : \mathcal{V}(h) \rightarrow \mathbb{C}$ with respect to g by

$$\langle v_h \rangle_g = \sigma \cdot v_g, \text{ if } \sigma(g) = h \text{ as elements of } \text{FM}(\mathfrak{h}),$$

being σ the rearranging element $\sigma \in \mathfrak{S}(|g|) = \mathfrak{S}(|h|)$.

We shall drop the subindex g in $\langle \cdot \rangle_g$ when the context is clear. If one factors g as $g_1 g_2$ with respect to an ‘abelianised’ product, an element $\sigma \in \mathfrak{S}(|g|)$ serves as correction, so that $\sigma(g_1 g_2) = g$. Their rearranging yields $\langle u_{g_1} t_{g_2} \rangle_g = \sigma \cdot (u_{g_1} t_{g_2})$ for suitably chosen functions u_{g_1} and t_{g_2} . In general, the collection of graphs g_1, \dots, g_r is not required to be connected. If the context is clear, we pick this rearranging element in a smaller group $\sigma \in \mathfrak{S}(r)$ that only permutes the arguments of $\mathcal{V}(g_i)$.

Definition 2.7. Denote by $H_1 H_2$ the subset $\{g_1 g_2 \mid g_a \in H_a\}$ in the free commutative monoid $\text{FM}^{\text{ab}}(\mathfrak{h})$ spanned by an independent set of graphs \mathfrak{h} . Given two functionals in $\mathcal{C}(\mathfrak{h})$, $U = \sum_{h \in H_1} u_h h$ and $T = \sum_{h \in H_2} t_h h$, we define their product $V = (U \cdot T)$ as the functional

$$V = U \cdot T = \sum_{g \in H_1 H_2 \subset \text{FM}^{\text{ab}}(\mathfrak{h})} v_g g,$$

whose coefficients v_g are given by the ‘ordered convolution’

$$v_g = \sum_{\substack{(g_1, g_2) \in H_1 \amalg H_2 \\ g_1 g_2 \stackrel{\sim}{\sim} g}} \langle u_{g_1} t_{g_2} \rangle_g. \quad (2.14)$$

Lemma 2.8. *This product on $\mathcal{C}(\mathfrak{h})$ is commutative.*

Proof. Let $U = \sum_{j \in J} u_j j$ and $T = \sum_{l \in L} t_l l$ be in $\mathcal{C}(\mathfrak{h})$. Given $g \in \text{FM}(\mathfrak{h})$ with $g \stackrel{\circ}{\sim} jl$ for some $j \in J$ and some $l \in L$ we show that the components v_g of $V = U \cdot T$ and \tilde{v}_g of $\tilde{V} = T \cdot U$ satisfy $v_g g = \tilde{v}_g g$ in $\mathcal{C}(\mathfrak{h})$. It suffices to exhibit an element $\nu \in \mathfrak{S}(|g|)$ that satisfies $\langle u_j t_l \rangle_g = \langle t_l u_j \rangle_g \nu(g)$. This ν will be next constructed.


We have the freedom to assume that $g = h_1^{\alpha_1} \cdots h_n^{\alpha_n}$. Since $j, l \in \text{FM}(\mathfrak{h})$,

$$j \underset{\circ}{\sim} h_1^{\theta_1} h_2^{\theta_2} \cdots h_n^{\theta_n} \quad \text{and} \quad l \underset{\circ}{\sim} h_1^{\lambda_1} h_2^{\lambda_2} \cdots h_n^{\lambda_n} \quad (2.15)$$


for some $0 \leq \lambda_i, \theta_i \leq \alpha_i$ that satisfy $\alpha_i = \theta_i + \lambda_i$, $i = 1, \dots, n$. In the notation introduced above, $\#I(j, h_i) = \theta_i$ and $\#I(l, h_i) = \lambda_i$. We begin by assuming that the relations above are equalities,

$$j = h_1^{\theta_1} h_2^{\theta_2} \cdots h_n^{\theta_n} \quad \text{and} \quad l = h_1^{\lambda_1} h_2^{\lambda_2} \cdots h_n^{\lambda_n}, \quad (2.16)$$

and restore towards the end the more general form (2.15). Let $|j| = \theta_1 + \dots + \theta_n$ and $|l| = \lambda_1 + \dots + \lambda_n$ be the orders of j and l . We define first $\sigma \in \mathfrak{S}(|g|)$ as the $(|j|, |l|)$ -shuffle determined by



Each h_i before double bar in the first row is a factor of j ; after the double bar, the h_i 's represent the factors of l . The lower is a factorisation of g . Thus the diagram states that $\sigma(jl) = g$. Analogously, we can define a $(|l|, |j|)$ -shuffle ρ that satisfies $\rho(lj) = g$. This is depicted in the following diagram, in which we represent l to the left of the double bar and j to the right.



One has $\rho(lj) = g = \sigma(jl)$ but this still does not guarantee that $\rho^{-1}\sigma(v_{jl}) = \tilde{v}_{lg}$. In order to correct this, we define certain permutations τ_i that are constant everywhere except in the elements pertaining a particular h_i for fixed i . This embeds $\mathfrak{S}(\#I(g; h_i)) \subset \mathfrak{S}(|g|)$. Notice first that for each such element $\mu \in \mathfrak{S}(\#I(g; h_i))$

$$w_g g = (\mu \cdot w_g) \mu(g) = (\mu \cdot w_g) g$$

holds in $\mathcal{C}(\mathfrak{h})$ for any $w_g : \mathcal{V}(g) \rightarrow \mathbb{C}$. For each $i = 1, \dots, n$, define τ_i as certain permutation β_i (given below) in the range $[\alpha_{i-1} + 1, \alpha_i]$ and constant outside it:

$$\tau_i(x) = \begin{cases} \beta_i(x) & 0 < x - s_i < \alpha_i \\ x & \text{otherwise} \end{cases},$$

where $s_i := (\alpha_1 + \dots + \alpha_{i-1})$

$$\beta_i(x) = \begin{cases} x + \lambda_i & s_i < x \leq s_i + \theta_i, \\ x - \theta_i & \theta_i < x \leq \theta_i + \lambda_i = \alpha_i. \end{cases}$$


Then the sought-after ν is

$$\nu = \rho^{-1} \circ (\tau_1 \circ \dots \circ \tau_n) \circ \sigma,$$

which by construction satisfies

$$(\nu \cdot v_{jl}) \nu(jl) = \tilde{v}_{lj} l_j. \quad (2.17)$$

The map ν is thus given by



We now come back to the strongest, original statement, in which l and j have the form (2.15), instead of (2.16). This means that there are permutations $\gamma \in \mathfrak{S}(|j|) \subset \mathfrak{S}(|j| + |l|)$ and $\delta \in \mathfrak{S}(|l|) \subset \mathfrak{S}(|j| + |l|)$ with

$$\gamma(j) = h_1^{\theta_1} h_2^{\theta_2} \dots h_n^{\theta_n} \quad \text{and} \quad \delta(l) = h_1^{\lambda_1} h_2^{\lambda_2} \dots h_n^{\lambda_n}. \quad (2.18)$$

Then we correct ν by these two elements:

$$\nu = \rho^{-1} \circ (\tau_1 \circ \dots \circ \tau_n) \circ \sigma \circ (\gamma, \delta)$$

which satisfies, in the most general case, eq. (2.17). The statement follows by linear extension of it. \square

Example. To illustrate this notation, consider the sets $H = \{fg, f^2\}$ and $K = \{fg, g^2\}$ of coloured graphs and let $U = \sum_{e \in K} u_e e$ and $T = \sum_{e \in H} t_e e$. If X_A are momenta of f and Z_A of g (for $A = 1, 2$), we pick a particular graph $l = f^2 g^2$, and define the following permutations in $\mathfrak{S}(4)$

$$\sigma = (13)(24) = \begin{array}{c} \text{---} \\ \diagdown \quad \diagup \\ \text{---} \\ \diagup \quad \diagdown \\ \text{---} \end{array} \quad \tau = (1)(23)(4) = \begin{array}{c} \text{---} \\ \diagdown \quad \diagup \\ \text{---} \\ \text{---} \\ \text{---} \end{array} \quad (2.19)$$

The coefficient $v_l = v_{f^2g^2}$ is given by

$$\begin{aligned} v_{f^2g^2}(X_1, X_2, Z_1, Z_2) &= (\langle u_{fgtfg} \rangle_l + \langle u_{g^2tfg^2} \rangle_l)(X_1, X_2, Z_1, Z_2) \\ &= (\tau \cdot (u_{fgtfg}))(X_1, X_2, Z_1, Z_2) \\ &\quad + (\sigma \cdot (u_{g^2tfg^2}))(X_1, X_2, Z_1, Z_2) \\ &= u_{fgtfg}(X_1, Z_1, X_2, Z_2) + u_{g^2tfg^2}(Z_1, Z_2, X_1, X_2). \end{aligned}$$

Structures appearing in the graph calculus resemble the monoid ring. Given a commutative unit ring R and a monoid M , the monoid ring [Lan02, Ch. II] is built by formal finite sums in M with coefficients in R ,

$$R[M] = \left\{ \sum_m r_m m \mid r_m \in R, m \in M \text{ \& } r_n \neq 0 \text{ for finitely many } n \in M \right\},$$

and endowed with the convolution product. The structure of the graph calculus generated by n variables $\mathfrak{h} = \{h_1, \dots, h_n\}$ requires to define, instead of the ring R , the collection \mathcal{A} of algebras of functions

$$\mathcal{A} = \prod_{g \in \text{FM}(\mathfrak{h})} A_g, \quad \text{where } A_g = \{\text{functions } \mathcal{V}(g) \rightarrow \mathbb{C}\},$$

and then to consider the following restricted version of the monoid algebra

$$\mathcal{A}[\text{FM}(\mathfrak{h})] := \left\{ \sum_{g \in \text{FM}(\mathfrak{h})} u_g g \mid u_g \in A_g, u_h = 0 \text{ for almost all } h \in \text{FM}(\mathfrak{h}) \right\}.$$

Then we see that

$$\mathcal{C}(\mathfrak{h}) = \mathcal{A}[\text{FM}(\mathfrak{h})] / \sim.$$

Given functionals $U = \sum_{h \in H} u_h h$, $T = \sum_{l \in L} t_l l$ in $\mathcal{C}(\mathfrak{h})$ one defines their sum by

$$U + T = \sum_{m \in H \cup L} \left(\sum_{h \overset{\sim}{\sim} m, h \in H} \langle u_h \rangle_m + \sum_{l \overset{\sim}{\sim} m, l \in L} \langle t_l \rangle_m \right) m;$$

for a scalar $x \in \mathbb{C}$, the functional xU is defined by componentwise multiplication by x ,

$$xU = \sum_{h \in H} (x \cdot u_h) h.$$

The in-depth study of the structure of $\mathcal{C}(\mathfrak{h})$ is beyond the scope of this paper. Here, we will prove the properties that are useful to in later sections.

Definition 2.9 (Coloured Borel transformation). Let $V = \sum_{h \in H} v_g g$ be a generating functional of graph-group actions. Then define its *coloured Borel transformation* by

$$B_c(V) := \sum_{g \in H \subset G_D} \frac{1}{|G(g)|} v_g g.$$

If $H \subset G_D \ni h$, the following set

$$H_h := \{g \in G_D \mid hg \overset{\sim}{\sim} j \text{ for some element } j \text{ of } H\},$$

will be relevant for the next lemma.

Lemma 2.10. *Consider the generating functional of a system of graph-group actions $V = \sum_{g \in H} v_g g$ belonging to a graph calculus in (connected graph) variables $\mathfrak{h} = \{h_1, \dots, h_n\} \subset$*

\mathcal{G}_D . Suppose that the coefficients $v_{c_1 \dots c_a \dots c_b \dots c_p}$ obey the following rule under the transposition $(ab) \in \mathfrak{S}(p)$ of graphs:

$$v_{c_1 \dots c_a \dots c_b \dots c_p} = (ab) \cdot v_{c_1 \dots c_b \dots c_a \dots c_p} \quad (2.20)$$

for each $c_a, c_b \in \mathfrak{h}$, where a and b denote the number of factor (connected component) where c_a and c_b are located, respectively. Moreover, assume that $v_{c_1 \dots c_r \dots c_p}$ is invariant under $G(c_r)$ for each factor c_r , $1 \leq r \leq p$. Then, one has for $h = c_r$,

$$\frac{\delta B_c(V)}{\delta h} = \sum_{g \in H_h} \frac{1}{|G(g)|} v_{hg} g,$$

which means that for each $X \in \mathcal{V}(h)$,

$$\frac{\delta B_c(V)}{\delta h(X)} = \sum_{g \in H_h} \frac{1}{|G(g)|} (\iota_X^1 v_{hg}) g = \sum_{g \in H_h} \frac{1}{|G(g)|} v_{hg}(X, \cdot) g,$$

being ι_X^1 the insertion of momentum X of h at the first argument of the function v_{hg} .

Proof. By assumption, one can bring every element $g \in H$ to the form $g = h_1^{\alpha_1} \dots h_n^{\alpha_n}$, and change the coefficients v_g accordingly without alteration. Notice that each $g \in H$ can be further factorised as $h^\ell f$, where $\delta f / \delta h \equiv 0$, and ℓ depends on g . Through direct computation,

$$\begin{aligned} \frac{\delta B_c(V)}{\delta h(X)} &= \sum_{g=h^\ell f \in H} \sum_{r=1, \dots, \ell} \sum_{\sigma \in G(h)} \frac{1}{|G(h^\ell f)|} \iota_{\sigma(X)}^r v_{h^\ell f} h^{\ell-1} f \\ &= \sum_{g=h^\ell f \in H} \sum_{\sigma \in G(h)} \frac{\ell}{|G(h^\ell f)|} \iota_{\sigma(X)}^1 v_{h^\ell f} h^{\ell-1} f \\ &= \sum_{g=h^\ell f \in H} \frac{\ell \cdot |G(h)|}{|G(h^\ell f)|} \iota_X^1 v_{h^\ell f} h^{\ell-1} f. \end{aligned} \quad (2.21)$$

First, we used the invariance (2.20), i.e. $\iota_X^m v_{h^\ell f} = \iota_X^p v_{h^\ell f}$ for each $1 \leq p, m \leq \ell$; then, the invariance under $G(h)$. Throughout, we can assume $\ell \geq 1$, since this is required for a summand in the first equality of eq. (2.21) not to vanish. Also, since

$$G(h^\ell f) = G(h^\ell) \wr \mathfrak{S}(\ell) \times G(f),$$

the orders of the groups should satisfy

$$|G(h^\ell f)| = \ell! \cdot |G(h)|^\ell \cdot |G(f)| = \ell |G(h)| \cdot |G(h^{\ell-1} f)|.$$

Hence, after cancellation one gets

$$\frac{\delta B_c(V)}{\delta h(X)} = \sum_{g \in H_h} \frac{1}{|G(g)|} \iota_X^1 v_{hg} g. \quad \square$$

Definition 2.11. The graph derivative $\partial U / \partial h$ of a generating functional of group actions U is given by the coefficient v_\emptyset of the empty graph \emptyset of the functional derivative of U with respect to h , $\delta U / \delta h =: \sum_g v_g g$, to wit

$$\frac{\partial U}{\partial h} = \left(\frac{\delta U}{\delta h} \right)_\emptyset.$$

Parenthetically, the difference in notations for ‘partial derivative’ and ‘functional derivative’ does not intend to mirror any difference between multivariable ordinary and functional calculi.

The next result is simple and useful at the same time:

Lemma 2.12 (Graph calculus Leibniz product rule). *Consider a graph calculus $\mathcal{C}(\mathfrak{h})$ and let $J, L \subset \text{FM}(\mathfrak{h})$ span functionals U and T in $\mathcal{C}(\mathfrak{h})$,*

$$U = \sum_{j \in J} u_j j, \quad T = \sum_{l \in L} t_l l. \quad (2.22)$$

Then the graph derivative of the product $V = U \cdot T = \sum_g v_g g$ is

$$\frac{\partial(U \cdot T)}{\partial g} = \sum_{\Omega \in G(g)} \Omega \cdot v_g = \sum_{\Omega \in G(g)} \sum_{\substack{(j,l) \in J \amalg L \\ j l \sim g}} \Omega \cdot \langle u_j t_l \rangle_g.$$

Proof. We compute directly the derivative of the product $U \cdot T$ with respect to g from the \emptyset -coefficient of $\delta(U \cdot T)/\delta g$:

$$\begin{aligned} \frac{\partial(U \cdot T)}{\partial g} &= \left(\frac{\delta(U \cdot T)}{\delta g} \right)_{\emptyset} \\ &= \left(\frac{\delta}{\delta g} \sum_{f \in JLC\text{FM}^{\text{ab}}(\mathfrak{h})} \sum_{\substack{(j,l) \in J \amalg L \\ j l \sim f}} \langle u_j t_l \rangle_f f \right)_{\emptyset} \\ &= \sum_{f \in JLC\text{FM}^{\text{ab}}(\mathfrak{h})} \frac{\delta f}{\delta g} \sum_{\substack{(j,l) \in J \amalg L \\ j l \sim f}} \langle u_j t_l \rangle_f \\ &= G(g) \curvearrowright \left[\sum_{\substack{(j,l) \in J \amalg L \\ j l \sim g}} \langle u_j t_l \rangle_g \right] \\ &= \sum_{\Omega \in G(g)} \sum_{\substack{(j,l) \in J \amalg L \\ j l \sim g}} \Omega \cdot \langle u_j t_l \rangle_g. \end{aligned}$$

For the second equality we inserted the coefficients explicitly, according to the definition of the product. The fourth equality holds by graph independence, eq. (2.10), guaranteed for a graph calculus. \square

From now on let $2k(g) = |g^0|$ denote the number of vertices of $g \in \mathbf{G}_D$. Our canonical example of system of graph-group actions have the form

$$\{u_g, \mathcal{V}(g) = M_{D \times k(g)}(\mathbb{Z}), G(g) = \text{Aut}_c(g)\}_{g \in H}. \quad (2.23)$$

Due to the rigidity of a coloured graph, each automorphism⁹ π of a connected graph g is determined by a permutation σ of the black (or white) vertices of g ; we write $\pi = \hat{\sigma}$ for such $\sigma \in \mathfrak{S}(k(g))$. The action of the coloured automorphisms $\hat{\sigma} \in \text{Aut}_c(g) \subset \mathfrak{S}(k(g))$ on $M_{D \times k(g)}(\mathbb{Z})$ is by permutation of the matrix columns, $\mathbf{y}^i \mapsto \mathbf{y}^{\sigma(i)}$, $i = 1, \dots, k$. As a notational remark, we will often write k instead of $k(g)$, as we just did, if the context is clear.

2.5. Three limit cases and examples. The previous lemma implies the Leibniz multivariable rule. Before elaborating on it, for the case of graph-group actions by automorphisms, $G(g) = \text{Aut}_c(g)$, it will be helpful to exhibit this group action on a function v_g in three limit cases, according to the graph type of g .

Consider $\{\mathcal{V}(g), \text{Aut}_c(g)\}_{g \in \mathfrak{h}}$, an independent system graph-group actions, $\mathfrak{h} = \{h_1, \dots, h_n\} \subset \mathbf{G}_D$. Let $g = h_1^{\alpha_1} \cdots h_n^{\alpha_n}$ and let U and T graph-generated functionals by J and L , respectively,¹⁰ being these subsets of the the monoid generated by $\{h_1, \dots, h_n\}$. Then according to Lemma 2.12,

$$\frac{\partial(U \cdot T)}{\partial g} = \sum_{\Omega \in \text{Aut}_c(g)} \sum_{\substack{(j,l) \in JL \subset \text{FM}^{\text{ab}}(\mathfrak{h}) \\ j \cdot l \sim g}} \Omega \cdot \langle u_j t_l \rangle_g. \quad (2.24)$$

This relation holds in a subdomain $\mathcal{U}(g) = \mathcal{F}_{D,k(g)} \subset \mathcal{V}(g) = M_{D \times k(g)}(\mathbb{Z})$ to be justified later (see Sect. 3.2):

$$\mathcal{F}_{D,k} := \{(\mathbf{y}^1, \dots, \mathbf{y}^k) \in M_{D \times k}(\mathbb{Z}) \mid y_c^\alpha \neq y_c^\nu \text{ for all } c = 1, \dots, D \\ \text{and } \alpha, \nu = 1, \dots, k, \alpha \neq \nu\}.$$

The canonical action of $\text{Aut}_c(g)$ obviously restricts to this set $\mathcal{U}(g) = \mathcal{F}_{D,k(g)}$. Recalling that $\text{Aut}_c(g) = \text{Aut}_c(h_1^{\alpha_1} \amalg h_2^{\alpha_2} \amalg \dots \amalg h_n^{\alpha_n}) = \prod_{i=1}^n \text{Aut}_c(h_i) \wr \mathfrak{S}(\alpha_i)$, first we elaborate on three simple cases:

- **Case I:** if $n = 1$. Then $g = h^\alpha$, any $\Omega \in \text{Aut}_c(g) = \text{Aut}_c(h) \wr \mathfrak{S}(\alpha)$ is given by $\sigma = (\sigma^1, \dots, \sigma^\alpha) \in \text{Aut}_c(h)^\alpha$ and $\mu \in \mathfrak{S}(\alpha)$, yielding for eq. (2.24)

$$\Omega^{-1} \cdot (v_g)(X_1, \dots, X_\alpha) = v_g(\sigma^1 X_{\mu(1)}, \dots, \sigma^\alpha X_{\mu(\alpha)}).$$

Here each $X_A = (\mathbf{x}_A^1, \dots, \mathbf{x}_A^k) \in \mathcal{F}_{D,k(h)}$ and the action of the automorphism group $\tau \in \text{Aut}_c(h)$ is given by $\tau(X_A) = (\mathbf{x}_A^{\tau(1)}, \dots, \mathbf{x}_A^{\tau(k)})$.

- **Case II:** if $n \neq 1$ but $\alpha_A = 1$ for all $A = 1, \dots, n$. In this case, $g = h_1 h_2 \cdots h_n$. Then $\text{Aut}_c(g) = \prod_i \text{Aut}_c(h_i) \ni \Omega = (\sigma_1, \dots, \sigma_n)$, which acts like

$$\Omega^{-1} \cdot (v_g)(X^1, \dots, X^n) = v_g(\sigma_1(X^1), \dots, \sigma_n(X^n)).$$

⁹There are more than one definition of ‘automorphism of a coloured graph’. The one used here is introduced in [Pér18]. In this setting, an automorphism of a coloured graph is a graph-morphism that preserves the colouring of the edges and bipartiteness of the vertex-set in a strict way (not up to a permutation of colours as the factor 1/3 in the action of the quartic rank-3 model in [CT16] suggests). That is, edges of colour a have to be mapped to edges of colour a ; black (resp. white) vertices to black (resp. white) vertices.

¹⁰Of course, one could just take the union of the both spanning sets of graphs, if they do would not a priori coincide.

- **Case III:** If $g = h_1^{\alpha_1} \cdots h_n^{\alpha_n}$, but all automorphisms $\text{Aut}_c(h_i)$ are trivial. Then

$$\text{Aut}_c(g) = \prod_i \text{Aut}_c(h_i) \wr \mathfrak{S}(\alpha_i) = \mathfrak{S}(\alpha_1) \times \cdots \times \mathfrak{S}(\alpha_n) \ni (\mu_1, \dots, \mu_n).$$

We use now multi-index notation $\alpha = (\alpha_1, \dots, \alpha_n)$ and $\gamma = (\gamma_1, \dots, \gamma_n)$ and abbreviate $u_{\gamma_1, \dots, \gamma_n} = u_{h_1^{\gamma_1} \cdots h_n^{\gamma_n}}$, and similarly for $t_{\gamma_1, \dots, \gamma_n}$. One can rewrite then

$$\begin{aligned} \frac{\partial(U \cdot T)}{\partial g} &= \sum_{\mu_1 \in \mathfrak{S}(\alpha_1)} \cdots \sum_{\mu_n \in \mathfrak{S}(\alpha_n)} \sum_{\substack{(\gamma_i, \nu_i) \\ \gamma_i, \nu_i \geq 0 \\ \gamma_i + \nu_i = \alpha_i \\ i=1, \dots, n}} (\mu_1, \dots, \mu_n)^* \langle u_{\gamma_1, \dots, \gamma_n} t_{\nu_1, \dots, \nu_n} \rangle g \\ &= \sum_{\mu_1 \in \mathfrak{S}(\alpha_1)} \cdots \sum_{\mu_n \in \mathfrak{S}(\alpha_n)} \sum_{\substack{0 \leq \gamma_i \leq \alpha_i \\ i=1, \dots, n}} (\mu_1, \dots, \mu_n)^* \langle u_{\gamma_1, \dots, \gamma_n} t_{\alpha_1 - \gamma_1, \dots, \alpha_n - \gamma_n} \rangle g \end{aligned}$$

in multi-index notation as

$$\frac{\partial(U \cdot T)}{\partial g} = \sum_{\mu \in \prod_{i=1}^n \mathfrak{S}(\alpha_i)} \sum_{\gamma \leq \alpha} (\mu)^* \langle u_{\gamma} t_{\alpha - \gamma} \rangle g.$$

For constant functions $u_{\gamma_1, \dots, \gamma_n}, t_{\gamma_1, \dots, \gamma_n}$, this should reduce to the multivariable product formula. Indeed,

$$\frac{\partial(U \cdot T)}{\partial g} = \sum_{\gamma \leq \alpha} \binom{\alpha_1}{\gamma_1} \cdots \binom{\alpha_n}{\gamma_n} u_{\gamma} t_{\alpha - \gamma}$$

which is just the Leibniz rule eq. (1.1).

Conveniently, lower-case (super)indices ($i = 1, \dots, n$) of momenta label the graph type, whereas upper-case (sub)indices indicate the number of copy ($A = 1, \dots, \alpha_i$) of the i -th graph type.

We describe now the action of $\text{Aut}_c(g)$ on a general function $v_g : M_{D \times k(g)}(\mathbb{C}) \rightarrow \mathbb{C}$.

Let $\mathbf{X} = ((X_1^1, \dots, X_{\alpha_1}^1) \cdots (X_1^n, \dots, X_{\alpha_n}^n)) \in M_{D \times k(g)}(\mathbb{Z})$, being $X_A^i \in M_{D \times k(h_i)}(\mathbb{Z})$ the momentum of the A -th copy of h_i , for $A = 1, \dots, \alpha_i$. Picking an element $\Omega = (\sigma, \mu) \in \text{Aut}_c(g)$, with

$$(\sigma_1, \dots, \sigma_n) \in \prod_{i=1}^n \text{Aut}(h_i)^{\alpha_i}, \quad \sigma_i = (\sigma_i^1, \dots, \sigma_i^{\alpha_i}), \quad (2.25)$$

$$\mu = (\mu_1, \dots, \mu_n) \in \prod_{i=1}^n \mathfrak{S}(\alpha_i), \quad (2.26)$$

the following holds:

$$(\Omega^{-1} \cdot v_g)(X_A^i) = v_g((\sigma_i^A X_{\mu_i(A)}^i)^i), \quad (2.27)$$

which is short-hand notation for

$$\begin{aligned} &\Omega^{-1} \cdot v_g(\mathbf{X}) \\ &= v_g((\sigma_1^1 X_{\mu_1(1)}^1, \sigma_1^2 X_{\mu_1(2)}^1, \dots, \sigma_1^{\alpha_1} X_{\mu_1(\alpha_1)}^1), \dots, (\sigma_n^1 X_{\mu_n(1)}^n, \dots, \sigma_n^{\alpha_n} X_{\mu_n(\alpha_n)}^n)). \end{aligned}$$

Usually, it is summed all over $\Omega \in \text{Aut}_c(g)$, which being a group, allows us to choose whether we put the inverse in (2.27).

Example. For $e = \ominus$, $f = 1\text{I}\text{I}_1$ and $g = \text{I}\text{I}\text{I}$, let the subsets

$$H = \{\emptyset, e^2, f^2\} \quad \text{and} \quad L = \{e^3 f, e^2 g, e^2 f^2 g\}$$

span the functionals $U = \sum_{h \in H} u_h h$ and $T = \sum_{l \in L} t_l l$, and consider their product $V = U \cdot T$. According to the action (2.27), one has the following formulae:

- For $b = e^5 f$,

$$\begin{aligned} \frac{\partial V}{\partial b} &= \sum_{\Omega \in \text{Aut}_c(b)} \Omega \cdot \langle u_h t_l \rangle_b \\ &= \sum_{\Omega \in \text{Aut}_c(e^5 f)} \Omega \cdot \langle u_{e^2 t_{e^3 f}} \rangle_b \\ &= \sum_{(\sigma, \epsilon) \in \mathfrak{S}(5) \times \mathbb{Z}_2} (\sigma, \epsilon) \cdot (u_{e^2 t_{e^3 f}}), \end{aligned} \quad (2.28)$$

since $\text{Aut}_c(e^5 f) = \text{Aut}_c(\ominus) \wr \mathfrak{S}(5) \times \text{Aut}_c(1\text{I}\text{I}_1) \wr \mathfrak{S}(1) = \{1\} \wr \mathfrak{S}(5) \times \mathbb{Z}_2 \wr \{1\}$. Also the ordering $\langle \cdot \rangle_b$ is trivial. This in turn means that for the five momenta $X_A \in \mathcal{F}_{1,D=3} = \mathbb{Z}^3$ of e^5 and the momentum $Z = (\mathbf{z}^1, \mathbf{z}^2) \in \mathcal{F}_{2,3}$,

$$\frac{\partial V}{\partial e^5 | f}(X_A, Z) = \sum_{(\sigma, \epsilon) \in \mathfrak{S}(5) \times \mathbb{Z}_2} (u_{e^2 t_{e^3 | f}})(X_{\sigma(A)}, \mathbf{z}^{\epsilon(1)}, \mathbf{z}^{\epsilon(2)})$$

in abstract notation, or displaying the graphs:

$$\begin{aligned} \frac{\partial V}{\partial \ominus^5 | 1\text{I}\text{I}_1}(X_A, Z) &= \sum_{(\sigma, \epsilon) \in \mathfrak{S}(5) \times \mathbb{Z}_2} u_{\ominus | \ominus}(X_{\sigma(1)}, X_{\sigma(2)}) \\ &\quad \times t_{\ominus | \ominus | \ominus | 1\text{I}\text{I}_1}(X_{\sigma(3)}, X_{\sigma(4)}, X_{\sigma(5)}, \mathbf{z}^{\epsilon(1)}, \mathbf{z}^{\epsilon(2)}). \end{aligned}$$

- For $b' = e^2 f^2 g$,

$$\frac{\partial V}{\partial b'} = \sum_{\Omega \in \text{Aut}_c(e^2 f^2 g)} \Omega \cdot \langle u_{\emptyset t_{e^2 f^2 g}} \rangle + \Omega \cdot \langle u_{f^2 t_{e^2 g}} \rangle.$$

For X_A momenta of e^2 , Z_A momenta of f^2 and total momentum $\mathbf{X} = (X_1, X_2, Z_1, Z_2, W)$, one sums over elements $\Omega = (\epsilon, (\sigma^1, \sigma^2; \mu), \tau) \in \text{Aut}_c(e^2 f^2 g) = \mathfrak{S}(2) \times (\mathbb{Z}_2 \wr \mathfrak{S}(2)) \times \mathbb{Z}_3$, which yields for $(\partial V / \partial b')(\mathbf{X}) = (\partial V / \partial (\ominus^2 | 1\text{I}\text{I}_1^2 | \text{I}\text{I}\text{I}))(\mathbf{X})$ the expression

$$\begin{aligned} &\sum_{\Omega} [\Omega \cdot (u_{\emptyset t_{e^2 f^2 g}}) + \Omega \cdot ((13)(24)]^* u_{f^2 t_{e^2 g}}](\mathbf{X}) \\ &= u_{\emptyset} \cdot \left\{ \sum_{(\epsilon, (\sigma^1, \sigma^2; \mu), \tau)} [t_{e^2 f^2 g}(X_{\epsilon(1)}, X_{\epsilon(2)}, (\sigma^1 Z_{\mu(1)}, \sigma^2 Z_{\mu(2)}, \tau(W))] \right\} \\ &\quad + \sum_{(\epsilon, (\sigma^1, \sigma^2; \mu), \tau)} u_{f^2} (\sigma^1 Z_{\mu(1)}, \sigma^2 Z_{\mu(2)}) t_{e^2 g}(X_{\epsilon(1)}, X_{\epsilon(2)}, \tau(W)) \\ &= u_{\emptyset} \cdot \left\{ \sum_{(\epsilon, (\sigma^1, \sigma^2; \mu), \tau)} [t_{\ominus^2 | 1\text{I}\text{I}_1^2 | \text{I}\text{I}\text{I}}(X_{\epsilon(1)}, X_{\epsilon(2)}, (\sigma^1 Z_{\mu(1)}, \sigma^2 Z_{\mu(2)}, \tau(W))] \right\} \\ &\quad + \sum_{(\epsilon, (\sigma^1, \sigma^2; \mu), \tau)} u_{1\text{I}\text{I}_1^2} (\sigma^1 Z_{\mu(1)}, \sigma^2 Z_{\mu(2)}) \times t_{\ominus^2 | \text{I}\text{I}\text{I}}(X_{\epsilon(1)}, X_{\epsilon(2)}, \tau(W)). \end{aligned}$$

One should still insert the explicit momenta $Z_A = (\mathbf{z}_A^1, \mathbf{z}_A^2) \in \mathcal{F}_{2,3}$, $W = (\mathbf{w}^1, \mathbf{w}^2, \mathbf{w}^3) \in \mathcal{F}_{3,3}$, $\tau(W) = (\mathbf{w}^{\tau(1)}, \mathbf{w}^{\tau(2)}, \mathbf{w}^{\tau(3)})$, and $\sigma^A Z_{\mu(A)} = (\mathbf{z}_{\mu(A)}^{\sigma^A(1)}, \mathbf{z}_{\mu(A)}^{\sigma^A(2)})$.

3. TENSOR MODELS

In this section, we implement the graph calculus for TFT.

3.1. Tensor Field Theory. The main idea to use the Ward-identity [DGMR07] to decouple the Schwinger-Dyson equations (at a planar sector) and to obtain a master integral equation for the 2-point functions for matrix models [GW14] can be extended to the TFT setting. Some progress along these lines has been made for complex tensor field theory and consists in the further study [Pér18] of the Ward-Takahashi identity of Ousmane-Samary [Ous14] in order to descend the SDE tower [PPW17] and eventually find closed equations. This led lately to the large- N limit [PPTW18] of the connected- ∂ SDE. We treat a TFT as inspired by group field theory [BG12, BGS13, COR14, OV14].

Unlike matrix models, where there is a canonical way of forming a scalar, for tensor models a specific trace $\text{Tr}_{\mathcal{B}}$, indicating how to contract the indices, should be specified. These traces $\text{Tr}_{\mathcal{B}}$ are indexed by D -coloured graphs \mathcal{B} , where D is the rank of the tensors $\varphi_{x_1 \dots x_D}, \bar{\varphi}_{x_1 \dots x_D}$.

The graphical representation of these traces derives from the independence of the imposed transformation rules under the action of $U(N)$ on the spaces corresponding to each index x_a of $\varphi_{x_1 \dots x_D}, \bar{\varphi}_{x_1 \dots x_D}$, for $a = 1, \dots, D$, deemed *colouring*. That is to say, to form invariants only indices of identical colour can be contracted.

Therefore, a trace corresponding to a quartic interaction would be, say, $\mathbb{1}_{\square_1}$ formed by colour-wise contracting the indices with deltas, as follows:

$$\text{Tr}_{\mathbb{1}_{\square_1}}(\varphi, \bar{\varphi}) = \sum_{\mathbf{x}, \mathbf{y}} \varphi_{x_1 y_2 x_3} \bar{\varphi}_{x_1 y_2 y_3} \varphi_{y_1 y_2 y_3} \bar{\varphi}_{y_1 x_2 x_3}.$$

The actual index-set of the tensors is $\{1, \dots, N\}$, but thinking of N as a large integer, we typically write these sums over \mathbb{N} or \mathbb{Z} , or omit the domains in the sums. Although orthogonal groups [CT16], compact symplectic groups [CP18] and mixed symmetries [Tan16] define other classes of tensor models, we restrict our discussion to the $U(N)$ -tensor models we just introduced.

A tensor model is thus determined by a dimension D (the rank of the tensors) and an action $S[\varphi, \bar{\varphi}]$ given by a finite sum of traces indexed by connected D -coloured graphs. The partition function is given by

$$Z[J, \bar{J}] = Z[0, 0] \frac{\int \mathcal{D}[\varphi, \bar{\varphi}] e^{\text{Tr}(\bar{J}\varphi) + \text{Tr}(\bar{\varphi}J) - N^{D-1}S[\varphi, \bar{\varphi}]}{\int \mathcal{D}[\varphi, \bar{\varphi}] e^{-N^{D-1}S[\varphi, \bar{\varphi}]}} \tag{3.1}$$

$$\mathcal{D}[\varphi, \bar{\varphi}] := \prod_{\mathbf{x}} N^{D-1} \frac{d\varphi_{\mathbf{x}} d\bar{\varphi}_{\mathbf{x}}}{2\pi i}.$$

Its logarithm, $W[J, \bar{J}] = \log Z[J, \bar{J}]$ is the *free energy* and generates the connected correlation functions, which as pointed out before, are classified by boundary (possibly disconnected D -coloured) graphs.

3.2. From graph calculus to tensor field theory. For a deeper exposition and motivation of the terminology and proofs of the results exposed in this section, we refer to [Pér18].

Some objects of interest in tensor models are functionals generated by graphs (e.g. the free energy). By this we mean expansions in graphs with functions (or distributions) as coefficients. For a graph $g \in H \subset \mathbf{G}_D$ we recall that $2k(g)$ denotes its number of vertices. One is interested in collections of functions

$$\{u_g : \mathbb{Z}^{D \times k(g)} \rightarrow \mathbb{C}\}_{g \in H}$$

and in their generating functionals

$$U[J, \bar{J}] = \sum_{g \in H \subset \mathbf{G}_D} u_g \star \mathbb{J}(g), \quad \text{where } u_g \star \mathbb{J}(g) := \sum_{\mathbf{X} \in \mathbb{Z}^{D \cdot k}} (u_g(\mathbf{X})) \mathbb{J}(g)(\mathbf{X}).$$

Here $\mathbb{J}(g)(\mathbf{X}) = \prod_{\alpha=1}^{k(g)} J_{\mathbf{x}^\alpha} \bar{J}_{\mathbf{y}^\alpha}$, where $\{\mathbf{y}^\alpha\}_\alpha$ are determined by g and \mathbf{X} through $g_*(\mathbf{X}) = (\mathbf{y}^1, \dots, \mathbf{y}^{k(g)})$. The induced map g_* is defined as follows. The D -tuple \mathbf{x}^α (resp. the \mathbf{y}^α) indexes white (resp. black) vertices in a graph. Then $g_* : M_{D \times k(g)}(\mathbb{Z}) \rightarrow M_{D \times k(g)}(\mathbb{Z})$ is given by $\mathbf{X} = (\mathbf{x}^1, \dots, \mathbf{x}^{k(g)}) \mapsto g_*(\mathbf{X}) = (\mathbf{y}^1, \dots, \mathbf{y}^k)$, where $y_c^\alpha = x_c^\nu$ (for $\alpha = 1, \dots, k$) if and only in the graph g there exists a c -coloured edge starting at \mathbf{x}^α and ending at \mathbf{y}^ν . As before, \mathbf{X} is called momentum, but also each one of these arguments \mathbf{x}^α is referred to as (*entering*) *momentum* of the white vertex $J_{\mathbf{x}^\alpha}$. Similarly \mathbf{y}^ν is the (*outgoing*) *momentum* at the black vertex $\bar{J}_{\mathbf{y}^\nu}$; the terminology relies on Figure 2. Although an ordering of the vertices is assumed, notice that $\mathbb{J}(g)$ is independent of it.

Consider the canonical system of graph-group actions introduced in eq. (2.23)

$$\{u_g, M_{D \times k(g)}(\mathbb{Z}), \text{Aut}_c(g)\}_{g \in H}.$$

A very important domain where the graph derivatives shall be defined is the subspace $\mathcal{F}_{D,k(g)}$ of $\mathbb{Z}^{D \cdot k(g)} \simeq M_{D \times k(g)}(\mathbb{Z})$ consisting of points outside all the coloured diagonals, i.e.

$$\begin{aligned} \mathcal{F}_{D,k} := \{(\mathbf{y}^1, \dots, \mathbf{y}^k) \in M_{D \times k}(\mathbb{Z}) \mid y_c^\alpha \neq y_c^\nu \text{ for all } c = 1, \dots, D \\ \text{and } \alpha, \nu = 1, \dots, k, \alpha \neq \nu\}. \end{aligned}$$

For a *connected* graph g , the elements of $\text{Aut}_c(g)$ are a lift $\hat{\sigma}$ of an element σ of the symmetric group $\mathfrak{S}(k(g))$ (see discussion below eq. (2.23), or [Pér18] for details). Defining

$$\frac{\partial U[J, \bar{J}]}{\partial g(\mathbf{X})} = \prod_{\alpha=1}^{k(g)} \frac{\delta}{\delta J_{\mathbf{x}^\alpha}} \frac{\delta}{\delta \bar{J}_{\mathbf{y}^\alpha}} U[J, \bar{J}] \Big|_{J=\bar{J}=0}, \quad \mathbf{X} = (\mathbf{x}^1, \dots, \mathbf{x}^{k(g)}) \in \mathcal{F}_{D,k(g)}, \quad (3.2)$$

one can give this derivative the the meaning of eq. (1.2) as a permutation of the arguments of u_g , that is

$$\frac{\partial U[J, \bar{J}]}{\partial g(\mathbf{X})} = \sum_{\hat{\sigma} \in \text{Aut}_c(g)} (\sigma \cdot u_g)(\mathbf{X}), \quad \text{where} \quad (3.3)$$

$$(\sigma \cdot u_g)(\mathbf{x}^1, \dots, \mathbf{x}^{k(g)}) := u_g(\mathbf{x}^{\sigma^{-1}(1)}, \dots, \mathbf{x}^{\sigma^{-1}(k(g))}), \quad (3.4)$$

for all $\mathbf{X} = (\mathbf{x}^1, \dots, \mathbf{x}^{k(g)}) \in \mathcal{F}_{D,k(g)}$. This statement is [Pér18, Lem. 4.1].




FIGURE 2. *Left.* On the interpretation of the induced map g_* . One takes any representing graph \mathcal{G} such that $\partial\mathcal{G} = g$. For the quartic melodic model (pillow-interactions) we can do so because it has been proven in [Pér18] that the spectrum of boundary states is full, which is to say, the boundary graphs are all of \mathbb{G}_D . Assuming an ordering on the white and on the black vertices, the components \mathbf{y}^α of $g_*(\mathbf{x}^1, \dots, \mathbf{x}^k) = (\mathbf{y}^1, \dots, \mathbf{y}^k)$ are determined by this picture, seeing the \mathbf{x}^μ 's as independent momenta entering the graph, with output $g_*(\mathbf{x}^1, \dots, \mathbf{x}^k)$. *Right.* If $g = \heartsuit$, and we choose the numeration $x_1^\alpha = y_1^\alpha$ for $\alpha = 1, 2, 3$; there that \mathcal{G} satisfying $\partial\mathcal{G} = \heartsuit$ is shown. The map $(\heartsuit)_*(\mathbf{x}^1, \mathbf{x}^2, \mathbf{x}^3) = (\mathbf{y}^1, \mathbf{y}^2, \mathbf{y}^3) = ((x_1^1, x_2^2, x_3^3)^t, (x_1^2, x_2^2, x_3^2)^t, (x_1^3, x_2^1, x_3^2)^t)$ is determined by following the $0a$ momenta lines for each colour a

Remark 3.1. Notice that in eq. (3.3) it is summed over the group $\text{Aut}_c(g)$. We are therefore entitled to drop the inverse in the RHS in (3.4). However, if the sum is not over all the group, we will keep the right ‘orientation’ of the action, for the convention in single terms $(\sigma \cdot u_g)$ is important in that case.

For graphs h, g , functions u_g and functionals $U[J, \bar{J}]$ we abbreviate the usual notation as follows:

$$U = U[J, \bar{J}], \quad u_{gg} := u_g \star \mathbb{J}(g) \quad \text{and} \quad gh := \mathbb{J}(g)\mathbb{J}(h), \quad (3.5)$$

and treat the latter as a product of graphs. This product is not considered commutative, since the star \star implies an ordering in the arguments of a function u_{gh} , which need not satisfy $u_{gh} = u_{hg}$. Now we exhibit the relation to the generating functional of group actions. With the product defined above, consider the functional $V^{\leq n}$ that generates the *connected* correlation functions of TFT

$$V^{\leq n} = \sum_{\mathcal{B} \in \mathbb{G}_D^n} G_{\mathcal{B}} \mathcal{B}.$$

(with $G_\emptyset = 0$.) Here n is a large integer, and \mathbb{G}_D^n is a finite set of coloured graphs whose elements \mathcal{D} satisfy

$$\#(\text{vertices of } \mathcal{D}) \leq 2n. \quad (3.6)$$

In tensor models, the subindices of the functions corresponding to the graph $\mathcal{A} \amalg \mathcal{B}$ (also written as juxtaposition $\mathcal{A}\mathcal{B}$) are rather denoted by $G_{\mathcal{A}|\mathcal{B}}$. These particular functions $G_{\mathcal{D}}$ satisfy invariance under $\text{Aut}_c(\mathcal{D})$; this means invariance under the Aut_c -groups of the connected components of \mathcal{D} and, for any isomorphic connected components \mathcal{A} and \mathcal{B} of \mathcal{D} , $G_{\dots|\mathcal{A}|\dots|\mathcal{B}|\dots} = G_{\dots|\mathcal{B}|\dots|\mathcal{A}|\dots}$.

The truncation (3.6) would declare vanishing all the floors above the n -th floor of the SDE-tower, but we can increase n at desired accuracy. It bounds any graph \mathcal{D} appearing in a non-identically vanishing correlator $G_{\mathcal{D}}$ to have n components at most. In rank 3, since the canonical (optimal in number of vertices) 3-coloured graph of genus g has $4g + 2$ vertices, this truncation bounds the genus through $2g + 1 \leq n$, making higher-genera boundary states vanish. We write the infinite sums keeping in mind that we mean their $n \rightarrow \infty$ limit. The coloured Borel transform of $V^\infty = \lim_{n \rightarrow \infty} V^{\leq n}$ is called the *free energy*:

$$W = \text{B}_c(V^\infty) = \sum_{\mathcal{D} \in \mathcal{G}_D} \frac{1}{|\text{Aut}_c(\mathcal{D})|} G_{\mathcal{D}} \mathcal{D}. \quad (3.7)$$

This equation holds in ‘ $N = 1$ -units’ and this assumption is innocuous within the scope of this article. However, if one plans to proceed perturbatively in $1/N$, the realistic case that drops this simplification ought to be addressed. Adding the power counting conjectured in [PPTW18] that scales $G_{\mathcal{D}} \rightarrow N^{\gamma(\mathcal{D})} G_{\mathcal{D}}$, where $\gamma(\mathcal{D})$ is certain factor already determined for the 2-pt and 4-pt functions, would help analysing the convergence of W (see Sect. 7).

This free energy functional (but not only this) corresponds to a system of graph-group actions that has the following constituents:

- For each graph \mathcal{B} , $\mathcal{V}(\mathcal{B}) = M_{D \times k(\mathcal{B})}(\mathbb{Z})$.
- For a disconnected graph $\mathcal{D} = \amalg_p \mathcal{B}_p$ one has

$$\mathcal{V}(\mathcal{D}) = M_{D \times k(\mathcal{D})}(\mathbb{Z}) = \prod_p \mathcal{V}(\mathcal{B}_p).$$

- There is an action of each $\text{Aut}_c(\mathcal{B}_p)$ on $M_{D \times k(\mathcal{B}_p)}(\mathbb{Z})$. Since $\sigma \in \text{Aut}_c(\mathcal{B}_p) \subset \mathfrak{S}(k(\mathcal{B}_p))$, precomposition by a function by σ permuting the columns of $M_{D \times k(\mathcal{B}_p)}(\mathbb{Z})$ gives this action.

For connected graphs \mathcal{B}_i and \mathcal{B}_j , the following holds [Pér18, Lemma 4]:

$$\frac{\delta \mathcal{B}_i}{\delta \mathcal{B}_j} = \delta_{ij} \text{Aut}_c(\mathcal{B}_i),$$

on the domain $\mathcal{F}_{D, k(\mathcal{B}_j)} \subset M_{D, k(\mathcal{B}_j)}$. For general disconnected graphs $\mathcal{D} = \mathcal{B}_1^{\alpha_1} \amalg \dots \amalg \mathcal{B}_m^{\alpha_m}$

$$\frac{\delta \mathcal{D}}{\delta \mathcal{D}} = \text{Aut}_c(\mathcal{D}) = \text{Aut}_c(\mathcal{B}_1) \wr \mathfrak{S}(\alpha_1) \times \dots \times \text{Aut}_c(\mathcal{B}_m) \wr \mathfrak{S}(\alpha_m) \quad (3.8)$$

on $\mathcal{F}_{D, k(\mathcal{D})} \subset M_{D \times k(\mathcal{D})}(\mathbb{Z})$. One can then operate with functionals, now without needing to evaluate the sources at 0. That is,

$$\frac{\delta U[J, \bar{J}]}{\delta g(\mathbf{X})} = \prod_{\alpha=1}^{k(g)} \frac{\delta}{\delta J_{\mathbf{x}^\alpha}} \frac{\delta}{\delta \bar{J}_{\mathbf{y}^\alpha}} U.$$

This object is, unlike the function $\partial U/\partial g$, a functional that can be graph-derived again (without getting something a trivial result) and get again a functional generated by graphs. Contrast this with eq. (3.2), whose result is a function.

Another important functional in the next derivation is the so-called Y -term that emerged in the derivation of the Ward-Takahashi identity [Pér18] and which encodes all the pertinent insertions of $2p$ point functions into $2p - 2$ point functions for all p .

The expression to order six is given in [PPW17, Lemma 4.1], but this article only will evoke the Y -term up to order four, located in Appendix A. For this paper, it is sufficient to additionally know the expression

$$Y_x^{(c)} = \sum_{\mathcal{B} \in \mathcal{G}_D} \mathfrak{f}_{\mathcal{B},x}^{(c)} \mathcal{B},$$

where $c \in \{1, \dots, D\}$ is a colour and $x \in \mathbb{Z}$. It is important to notice that unlike W (for which we set $G_\emptyset = 0$), there is a non-vanishing constant term $\mathfrak{f}_{\emptyset,x}^{(c)}$ in $Y_x^{(c)}$. Each function coefficient $\mathfrak{f}_{\mathcal{B},x}^{(c)}$ of a graph \mathcal{B} denotes a triple propagator contraction of vertices from ${}^c\mathbb{Q}^c$ with vertices of the graph \mathcal{C} having $2k(\mathcal{C}) = 2 + 2k(\mathcal{B})$ vertices and such that the whole contraction's boundary is \mathcal{B} . Although $G_{\mathcal{C}}$ is symmetric with respect to action of $\text{Aut}_c(\mathcal{C})$, the resulting insertion need not to be $\text{Aut}_c(\mathcal{B})$ -symmetric.

In order to derive any of the $2p$ -point SDE¹¹ we shall employ the graph calculus $\mathcal{C}(\mathfrak{h})$ with variables $\mathfrak{h} = \{\text{connected, closed, } D\text{-coloured graphs with } \leq 2(p+1) \text{ vertices}\}$, being the system of graph-group actions the canonical one given by automorphism groups (see (2.23) above for details).

4. DISCONNECTED-BOUNDARY SCHWINGER-DYSON EQUATIONS

The next section introduces the model whose SDE are found in Section 4.2.

4.1. The quartic melonic tensor field theory. The $\varphi_{D,m}^4$ -theory is the model with quartic interaction vertices $V[\varphi, \bar{\varphi}] = \lambda \sum_{a=1}^D \text{Tr}_{V_a}(\varphi, \bar{\varphi})$. These vertices are sometimes called pillows, since the graphs they correspond to have that appearance:

$$V_a = \begin{array}{c} a \\ \circ \text{---} \circ \\ | \quad | \\ \circ \text{---} \circ \\ a \end{array} \quad a = 1, \dots, D. \quad (4.1)$$

We analyse this theory with an abstract Laplacian $E : \mathbb{Z}^D \rightarrow \mathbb{R}_{\geq 0}$ as propagator, $S_0[\varphi, \bar{\varphi}] = \text{Tr}_{\ominus}(\bar{\varphi}, E\varphi) = \sum_{\mathbf{x}} \bar{\varphi}_{\mathbf{x}} E_{\mathbf{x}} \varphi_{\mathbf{x}}$, assumed here to satisfy the following technical assumption: for each colour c , the difference

$$E(t_c, s_c) := E_{p_1 \dots t_c \dots p_D} - E_{p_1 \dots s_c \dots p_D} \quad (4.2)$$

is independent of the fixed momenta in colours different from c . Such kind of technical conditions permit to exploit the Ward-Identity and are common. In matrix field theory [GW14, Thm. 2.3], this is analogous to the assumed injectivity of $n \mapsto E_n$ for the generalised matrix Laplacian $E_{\text{matrix}} = \text{diag}(E_n)_{n \in \mathbb{N}}$ there.

¹¹Here we mean the melonic quartic model. The bound on the vertices is model dependent and justified in the statement of Theorem 4.1.

4.2. Main result. We prepare¹² now some notions and notations needed for the formulation of the main result. Let \mathcal{R} be a connected graph, $2r$ its number of vertices, and $X \in \mathcal{F}_{r,D}$. Given a colour c and a numeration w^1, \dots, w^r of the black vertices of \mathcal{R} we set, for any $1 \leq \rho \leq r$,

$$\text{Br}^{(2)}(\mathcal{R}, \rho, c) := \{\tau \in \{1, \dots, r\} \mid \varsigma_c(\mathcal{R}, w^\rho, w^\tau) \text{ is disconnected}\}.$$

Thus, τ belonging to this set indexes the vertices w^τ at which a swap of the colour c edge at w^τ and w^ρ disconnects \mathcal{R} . In other words, $\text{Br}^{(2)}(\mathcal{R}, \rho, c)$ indexes c -coloured edges of \mathcal{R} that, paired with the c -coloured edge at w^ρ , form a 2-bridge. If $\text{Br}^{(2)}(\mathcal{R}, \rho, c) \neq \emptyset$, then \mathcal{R} must be, in particular, 3-edge connected.

Example. The coloured utility graph \heartsuit satisfies $\text{Br}^{(2)}(\heartsuit, \rho, c) = \emptyset$ for any colour c and vertex ρ ('no edge-swap separates it'). On the other hand, if $\{\rho, \mu\} = \{1, 2\}$ label the two black vertices of the pillow $1\heartsuit 1$, then

$$\text{Br}^{(2)}(1\heartsuit 1, \rho, c = 1) = \{\mu\} \text{ and } \text{Br}^{(2)}(1\heartsuit 1, \rho, c) = \emptyset \text{ for } c = 2, 3.$$

Notation of the theorem. Let \mathcal{D} be a D -coloured graph with $2d$ vertices. By [Pér18, Thm. 1] \mathcal{D} is a boundary graph of the $\varphi_{D,m}^4$ -theory and $G_{\mathcal{D}} \neq 0$. Given any $\mathbf{X} \in \mathcal{F}_{d,D}$, we select an outgoing momentum $\mathbf{s} = \mathbf{y}^\beta$ listed in

$$(\mathbf{y}^1, \dots, \mathbf{y}^\beta, \dots, \mathbf{y}^d) = \mathcal{D}_*(\mathbf{X}) \in \mathcal{F}_{d,D}. \quad (4.3)$$

This \mathbf{s} determines both a connected graph component \mathcal{R} of \mathcal{D} , and an r -tuple $X_0 \in \mathcal{F}_{r,D}$ of momenta, being $2r$ the number of vertices in \mathcal{R} , by asking that \mathbf{s} appears listed in the r -tuple $\mathcal{R}_*(X_0)$, particularly. Different choices of the distinguished momentum variable \mathbf{s} —say $\mathbf{s} = \mathbf{y}^{\beta_1}$ and $\mathbf{s} = \mathbf{y}^{\beta_2}$ —lead to a different SDE whenever the respective β_1 -th and β_2 -th black vertices lie on different connected components; or, less obviously, when such vertices do lie in the same connected graph, \mathcal{R} , but they are not related by a non-trivial element of $\text{Aut}_c(\mathcal{R})$. In particular, if the distinguished connected component \mathcal{R} has no symmetries, then there are exactly half as many SDE's as vertices of \mathcal{R} , to wit $k(\mathcal{R})$ equations (for that connected component).

For the rest of components of \mathcal{D} we write \mathcal{Q} ($\mathcal{D} = \mathcal{R} \amalg \mathcal{Q}$). We factorise \mathcal{Q} in α_u copies of pairwise non-isomorphic connected graphs $\{Q_u\}_{u=1, \dots, n}$:

$$\mathcal{Q} = Q_1^{\alpha_1} \amalg \dots \amalg Q_n^{\alpha_n}, \quad (Q_i^{\alpha_i} = \amalg_{A=1}^{\alpha_i} Q_i).$$

We split the d -tuple \mathbf{X} into momenta X_0 of \mathcal{R} and momenta \mathbb{X} of \mathcal{Q} , so that $(X_0, \mathbb{X}) = \mathbf{X}$, up to reordering. For $\tau \in \text{Br}^{(2)}(\mathcal{R}, \beta, c)$ we can therefore write

$$\varsigma_c(\mathcal{R}; \beta, \tau) = \mathcal{R}' \amalg \mathcal{R}'',$$

and accordingly split the momentum X_0 in momenta X'_0 of \mathcal{R}' and X''_0 of \mathcal{R}'' , $X_0 = (X'_0, X''_0)$. Furthermore, for any factorising pair of graphs

$$\mathcal{C}, \mathcal{B} \in \mathcal{G}_D \text{ with } \mathcal{C} \amalg \mathcal{B} \stackrel{\sim}{\simeq} \mathcal{Q},$$

¹²We come back to the usual notation for graphs or 'bubbles'.

we define the two functions $H_{\mathcal{C},\mathcal{B}}^{(c,\tau)}$ and $I_{\mathcal{C},\mathcal{B}}^{(c)}$ by the following products:

$$\begin{aligned} H_{\mathcal{C},\mathcal{B}}^{(c,\tau)} &= \langle G_{\mathcal{R}'|\mathcal{C}}(X'_0; \bullet) \times G_{\mathcal{R}''|\mathcal{B}}(X''_0; \bullet) \rangle_{\mathcal{Q}}, \quad (\tau \in \text{Br}^{(2)}(\mathcal{R}, \beta, c)) \\ I_{\mathcal{C},\mathcal{B}}^{(c)} &= \frac{1}{|\text{Aut}_c(\mathcal{B})|} \langle \mathfrak{f}_{\mathcal{C},s_c}^{(c)} \times G_{\mathcal{R}|\mathcal{B}}(X_0; \bullet) \rangle_{\mathcal{Q}}, \end{aligned}$$

where the reorderings refer only to the graph components of the graphs in the pair $(\mathcal{C}, \mathcal{B})$. The pivotal term, appearing in each equation, is

$$\mathfrak{f}_{\emptyset, s_c}^{(c)} = Y_{s_c}^{(c)}[0, 0] = \sum_{\mathbf{q}_{\hat{c}}} G_{\odot}^{(2)}(s_c, \mathbf{q}_{\hat{c}}).$$

Theorem 4.1. *For the $\varphi_{D,m}^4$ -theory with kinetic term (4.2), the Schwinger-Dyson equations for the disconnected graph \mathcal{D} read, for the particular vertex choice $\mathbf{s} = \mathbf{y}^\beta$, as follows:*

$$\begin{aligned} &G_{\mathcal{D}}^{(2k)}(\mathbf{X}) \\ &= \frac{(-2\lambda)}{E_{\mathbf{s}}} \sum_{c=1}^D \left\{ \sum_{\hat{\sigma} \in \text{Aut}_c(\mathcal{D})} \sigma \cdot \mathfrak{f}_{\mathcal{D}, s_c}^{(c)}(\mathbf{X}) \right. \\ &+ \sum_{\rho \neq \beta} \frac{1}{E(y_c^\rho, s_c)} \left[\frac{\partial W[J, \bar{J}]}{\partial \varsigma_c(\mathcal{D}; \beta, \rho)}(\mathbf{X}) - \frac{\partial W[J, \bar{J}]}{\partial \varsigma_c(\mathcal{D}; \beta, \rho)}(\mathbf{X}|_{s_c \rightarrow y_c^\rho}) \right] \\ &- \sum_{b_c} \frac{1}{E(s_c, b_c)} [G_{\mathcal{D}}^{(2k)}(\mathbf{X}) - G_{\mathcal{D}}^{(2k)}(\mathbf{X}|_{s_c \rightarrow b_c})] \\ &+ \sum_{\tau \in \text{Br}^{(2)}(\mathcal{R}; \beta, c)} \left[\sum_{\substack{(\mathcal{C}, \mathcal{B}) \in \mathbb{G}_D \amalg \mathbb{G}_D \\ (\mathcal{C} \amalg \mathcal{B}) \sim \mathcal{Q}}} \sum_{\Omega \in \text{Aut}_c(\mathcal{Q})} \frac{(\Omega \cdot H_{\mathcal{B}, \mathcal{C}}^{(c,\tau)})(\mathbb{X}) - (\Omega \cdot H_{\mathcal{B}, \mathcal{C}}^{(c,\tau)})(\mathbb{X}|_{s_c \rightarrow y_c^\tau})}{E(y_c^\tau, s_c)} \right] \\ &+ \left. \sum_{\substack{(\mathcal{C}, \mathcal{B}) \in \mathbb{G}_D \amalg \mathbb{G}_D \\ (\mathcal{C} \amalg \mathcal{B}) \sim \mathcal{Q}}} \sum_{\Omega \in \text{Aut}_c(\mathcal{Q})} (\Omega \cdot I_{\mathcal{B}, \mathcal{C}}^{(c)})(\mathbb{X}) \right\}. \end{aligned} \tag{4.4}$$

Proof. By definition,

$$G_{\mathcal{D}} = \frac{\partial W}{\partial \mathcal{D}} = \left[\frac{\delta}{\delta \mathcal{Q}} \left(\frac{\delta W}{\delta \mathcal{R}} \right) \right] \Big|_{J=\bar{J}=0}.$$

Spelled out, this means that

$$G_{\mathcal{D}}(\mathbf{X}) = \left\{ \left[\prod_{i=1}^n \frac{\delta^{\alpha_i}}{\delta \mathcal{Q}_1^i \delta \mathcal{Q}_2^i \cdots \delta \mathcal{Q}_{\alpha_i}^i}(\mathbb{X}) \left(\prod_{\alpha=1}^r \frac{\delta}{\delta J_{\mathbf{x}_0^\alpha}} \frac{\delta}{\delta \bar{J}_{\mathbf{y}_0^\alpha}} \right) \right] W[J, \bar{J}] \right\} \Big|_{J=\bar{J}=0}. \tag{4.5}$$

To compute $G_{\mathcal{D}}$, one needs to start then with a functional derivative of W with respect to a source, which we choose to be $\bar{J}_{\mathbf{s}} = \bar{J}_{\mathbf{y}^\beta}$. The partition function $Z[J, \bar{J}] =$

$\exp(W[J, \bar{J}])$ has been shown [Pér18] to satisfy

$$\frac{\delta W}{\delta \bar{J}_s} = \frac{1}{E_s} \left\{ J_s - e^{-W} \left(\frac{\partial V(\varphi, \bar{\varphi})}{\partial \bar{\varphi}_s} \right) \Big|_{\substack{\varphi \rightarrow \delta/\delta \bar{J} \\ \bar{\varphi} \rightarrow \delta/\delta J}} e^{+W} \right\}. \quad (4.6)$$

The colour- c -WTI leads to

$$\left(\frac{\partial V(\varphi, \bar{\varphi})}{\partial \bar{\varphi}_s} \right) \Big|_{\substack{\varphi \rightarrow \delta/\delta \bar{J} \\ \bar{\varphi} \rightarrow \delta/\delta J}} Z[J, \bar{J}] \quad (4.7)$$

$$= 2\lambda \sum_{c=1}^D (A_c(\mathbf{s}) - B_c(\mathbf{s}) + C_c(\mathbf{s}) + D_c(\mathbf{s}) + F_c(\mathbf{s})), \quad (4.8)$$

where each of the summands is given by

$$\begin{aligned} A_c(\mathbf{s}) &= Y_{s_c}^{(c)}[J, \bar{J}] \cdot \frac{\delta Z[J, \bar{J}]}{\delta \bar{J}_s}, \\ B_c(\mathbf{s}) &= \sum_{\mathbf{b}} \frac{J_{\mathbf{b}_{\hat{c}s_c}}}{E(b_c, s_c)} \frac{\delta^2 Z[J, \bar{J}]}{\delta \bar{J}_{s_{\hat{c}b_c}} \delta J_{\mathbf{b}}}, \quad (\mathbf{s}_{\hat{c}b_c} = (s_1, \dots, s_{a-1}, b_c, s_{c+1}, \dots, s_D)) \\ C_c(\mathbf{s}) &= \sum_{b_c} \frac{1}{E(b_c, s_c)} \frac{\delta Z[J, \bar{J}]}{\delta \bar{J}_s}, \\ D_c(\mathbf{s}) &= \sum_{\mathbf{b}} \frac{\bar{J}_{\mathbf{b}}}{E(b_c, s_c)} \frac{\delta^2 Z[J, \bar{J}]}{\delta \bar{J}_{s_{\hat{c}b_c}} \delta \bar{J}_{\mathbf{b}_{\hat{c}s_c}}}, \quad (\mathbf{b}_{\hat{c}s_c} = (b_1, \dots, b_{c-1}, s_c, b_{c+1}, \dots, b_D)) \\ F_c(\mathbf{s}) &= \frac{\delta Y_{s_c}^{(c)}[J, \bar{J}]}{\delta \bar{J}_s} \cdot Z[J, \bar{J}], \end{aligned}$$

for any $c = 1, \dots, D$. In this new derivation, it is convenient to work with

$$e^{-W} A_c(\mathbf{s}) = Y_{s_c}^{(c)}[J, \bar{J}] \cdot \frac{\delta W[J, \bar{J}]}{\delta \bar{J}_s}, \quad (4.9)$$

$$e^{-W} B_c(\mathbf{s}) = \sum_{\mathbf{b}} \frac{J_{\mathbf{b}_{\hat{c}s_c}}}{E(b_c, s_c)} \left[\frac{\delta^2 W[J, \bar{J}]}{\delta \bar{J}_{s_{\hat{c}b_c}} \delta J_{\mathbf{b}}} + \frac{\delta W[J, \bar{J}]}{\delta J_{\mathbf{b}}} \frac{\delta W[J, \bar{J}]}{\delta \bar{J}_{s_{\hat{c}b_c}}} \right], \quad (4.10)$$

$$e^{-W} C_c(\mathbf{s}) = \sum_{b_c} \frac{1}{E(b_c, s_c)} \frac{\delta W[J, \bar{J}]}{\delta \bar{J}_s}, \quad (4.11)$$

$$e^{-W} D_c(\mathbf{s}) = \sum_{\mathbf{b}} \frac{\bar{J}_{\mathbf{b}}}{E(b_c, s_c)} \left[\frac{\delta^2 W[J, \bar{J}]}{\delta \bar{J}_{s_{\hat{c}b_c}} \delta \bar{J}_{\mathbf{b}_{\hat{c}s_c}}} + \frac{\delta W[J, \bar{J}]}{\delta \bar{J}_{s_{\hat{c}b_c}}} \frac{\delta W[J, \bar{J}]}{\delta \bar{J}_{\mathbf{b}_{\hat{c}s_c}}} \right], \quad (4.12)$$

$$e^{-W} F_c(\mathbf{s}) = \frac{\delta Y_{s_c}^{(c)}[J, \bar{J}]}{\delta \bar{J}_s}. \quad (4.13)$$

Notice the presence of the product of derivatives in the B_c and D_c terms. These are called $B_c^{(\text{prod})}$ and $D_c^{(\text{prod})}$, respectively. The other two terms (which already appeared on the SDE's for connected boundaries) containing a double derivative are denoted by $B_c^{(\text{dd})}$ and $D_c^{(\text{dd})}$, respectively.

Next, we use the freedom to numerate momenta starting with the component \mathcal{R} ,

$$X_0 = (\mathbf{x}_0^1, \dots, \mathbf{x}_0^r) \in \mathcal{F}_{D,r}, \quad \mathcal{R}_*(X_0) = (\mathbf{y}_0^1, \dots, \mathbf{y}_0^r),$$

and $\mathbf{s} = \mathbf{y}^\beta$. For each item in the list

$$(\mathfrak{m}, M) \in \{(\mathfrak{a}, A), (\mathfrak{b}, B), (\mathfrak{c}, C), (\mathfrak{d}, D), (\mathfrak{f}, F)\},$$

we define the following functions:

$$\mathfrak{m}_c(\mathbf{X}; \mathbf{s}; \mathcal{D}) := \frac{\delta^\alpha}{\delta \mathcal{Q}^\alpha(\mathbb{X})} \prod_{\substack{\alpha \neq \beta \\ 0 \leq \alpha \leq r \\ \nu=1, \dots, r}} \frac{\delta}{\delta \bar{J}_{\mathbf{y}_0^\alpha}} \frac{\delta}{\delta J_{\mathbf{x}_0^\nu}} \left[e^{-W[J, \bar{J}]} M_c(\mathbf{s}) \right] \Big|_{J=\bar{J}=0}. \quad (4.14)$$

The splitting of \mathfrak{b} and \mathfrak{d} into terms $\mathfrak{b}^{(\text{dd})}$, $\mathfrak{b}^{(\text{prod})}$, $\mathfrak{d}^{(\text{dd})}$, $\mathfrak{d}^{(\text{prod})}$, respectively, still makes sense. We now determine all coefficients, beginning with the easiest.

The \mathfrak{c}_c and \mathfrak{f}_c terms are readily computed:

$$\begin{aligned} \mathfrak{c}_c(\mathbf{X}; \mathbf{s}; \mathcal{D}) &= \sum_{b_c} \frac{1}{E(b_c, s_c)} G_{\mathcal{D}}^{(2k)}(\mathbf{X}), \\ \mathfrak{f}_c(\mathbf{X}; \mathbf{s}; \mathcal{D}) &= \sum_{\hat{\pi} \in \text{Aut}_c(\mathcal{D})} \pi^* \mathfrak{f}_{\mathcal{D}}^{(c)}(\mathbf{X}). \end{aligned}$$

The term \mathfrak{c}_c itself is not finite, but a term arising from one of the three functions will render it finite. The three remaining \mathfrak{m}_c -functions involve derivatives of products of functionals. We first observe that the $2r - 1$ derivatives in the sources complete the graph derivative $\delta W / \delta \mathcal{R}$, so the \mathfrak{a}_c yields


$$\begin{aligned} \mathfrak{a}_c(\mathbf{X}; \mathbf{s}; \mathcal{D}) &= \frac{\delta^\alpha}{\delta \mathcal{Q}^\alpha(\mathbb{X})} \prod_{\substack{\alpha \neq \beta \\ 0 \leq \alpha \leq r \\ \nu=1, \dots, r}} \frac{\delta}{\delta \bar{J}_{\mathbf{y}_0^\alpha}} \frac{\delta}{\delta J_{\mathbf{x}_0^\nu}} \left(Y_{s_c}^{(c)}[J, \bar{J}] \cdot \frac{\delta W[J, \bar{J}]}{\delta \bar{J}_{\mathbf{s}}} \right) \Big|_{J=\bar{J}=0} \\ &= \frac{\delta^\alpha}{\delta \mathcal{Q}^\alpha(\mathbb{X})} \left(Y_{s_c}^{(c)}[J, \bar{J}] \cdot \frac{\delta W[J, \bar{J}]}{\delta \mathcal{R}(X_0)} \right) \Big|_{J=\bar{J}=0}. \end{aligned} \quad (4.15)$$

Notice that, after evaluation in the sources at 0,

$$\frac{\partial}{\partial \mathcal{A}} \cdots \frac{\partial}{\partial \mathcal{A}'} \frac{\partial}{\partial \mathcal{R}(X_0)} W[J, \bar{J}] = G_{\mathcal{R}|\mathcal{A}|\dots|\mathcal{A}'}(X_0, \bullet)$$

holds for any boundary graphs $\mathcal{A}, \dots, \mathcal{A}'$. Using the formula for the graph derivative of products (Lem. 2.12) and subsequently Lemma 2.10 one deduces

$$\begin{aligned} \mathfrak{a}_c(\mathbf{X}; \mathbf{s}; \mathcal{D}) &= \sum_{\substack{\mathcal{B}, \mathcal{C} \\ \mathcal{B} \amalg \mathcal{C} \sim \mathcal{Q}}} \sum_{\Omega \in \text{Aut}_c(\mathcal{Q})} \sum_{\sigma \in \text{Aut}_c(\mathcal{R})} \\ &\quad \frac{1}{|\text{Aut}_c(\mathcal{R} \amalg \mathcal{B})|} \Omega \cdot \langle G_{\mathcal{R}|\mathcal{B}}(\sigma(X_0); \bullet) \times \mathfrak{f}_{\mathcal{C}, s_c}^{(c)} \rangle_{\mathcal{Q}} \\ &= \sum_{\substack{\mathcal{B}, \mathcal{C} \\ \mathcal{B} \amalg \mathcal{C} \sim \mathcal{Q}}} \sum_{\Omega \in \text{Aut}_c(\mathcal{Q})} \frac{1}{|\text{Aut}_c(\mathcal{B})|} \Omega \cdot \langle G_{\mathcal{R}|\mathcal{B}}(X_0; \bullet) \times \mathfrak{f}_{\mathcal{C}, s_c}^{(c)} \rangle_{\mathcal{Q}} \end{aligned}$$

FIGURE 3. The swapping operation ς_c ; variables are relevant in the proof

$$= \sum_{\substack{\mathcal{B}, \mathcal{C} \\ \mathcal{B} \cup \mathcal{C} \sim \mathcal{Q}}} \sum_{\Omega \in \text{Aut}_c(\mathcal{Q})} \Omega \cdot I_{\mathcal{C}, \mathcal{B}}^{(c)}(\mathbb{X}).$$

We compute now¹³ the term \mathfrak{d}_c . We can split D_c into the double derivative contribution $D_c^{(\text{dd})}$ and the product of single derivatives $D_c^{(\text{prod})}$. If we decompose $\delta^\alpha / \delta \mathcal{Q}^\alpha$ and the rest of derivatives implied in \mathcal{R} into single functional derivatives,

$$\begin{aligned} & \mathcal{O}(\bar{J}) + \prod_{\substack{\alpha \neq \beta \\ \nu=1, \dots, d}} \frac{\delta}{\delta \bar{J}_{\mathbf{y}^\alpha}} \frac{\delta}{\delta J_{\mathbf{x}^\nu}} (e^{-W} D_c^{(\text{dd})}(\mathbf{s})) \\ &= \prod_{\alpha \neq \beta; \nu} \frac{\delta}{\delta \bar{J}_{\mathbf{y}^\alpha}} \frac{\delta}{\delta J_{\mathbf{x}^\nu}} \left[\sum_{\mathbf{b}} \frac{1}{E(b_c, s_c)} \bar{J}_{\mathbf{b}} \frac{\delta^2 W[J, \bar{J}]}{\delta \bar{J}_{s_{\hat{c}} b_c} \delta \bar{J}_{\mathbf{b}_{\hat{c}} s_c}} \right] \\ &= \sum_{\substack{\rho=1 \\ \rho \neq \beta}}^d \prod_{\substack{\beta \neq \alpha \neq \rho \\ \nu=1, \dots, d}} \frac{\delta}{\delta \bar{J}_{\mathbf{y}^\alpha}} \frac{\delta}{\delta J_{\mathbf{x}^\nu}} \left[\sum_{\mathbf{b}} \frac{\delta_{\mathbf{y}^\rho}^{\mathbf{b}}}{E(b_c, s_c)} \frac{\delta^2 W[J, \bar{J}]}{\delta \bar{J}_{s_{\hat{c}} b_c} \delta \bar{J}_{\mathbf{b}_{\hat{c}} s_c}} \right] \\ &= \sum_{\substack{\rho=1 \\ \rho \neq \beta}}^d \prod_{\substack{\alpha; (\beta \neq \alpha \neq \rho) \\ \nu=1, \dots, k}} \frac{\delta}{\delta \bar{J}_{\mathbf{y}^\alpha}} \frac{\delta}{\delta J_{\mathbf{x}^\nu}} \left[\frac{1}{E(y_c^\rho, s_c)} \frac{\delta^2 W[J, \bar{J}]}{\delta \bar{J}_{s_{\hat{c}} y_c^\rho} \delta \bar{J}_{\mathbf{y}_c^\rho s_c}} \right]. \end{aligned} \quad (4.16)$$

This is, after evaluation at $\bar{J} = J = 0$, all the (coloured) graphs obtained from \mathcal{D} (also implying the other connected components) after the colour- c swapping at $\mathbf{s} = \mathbf{y}^\beta$ and \mathbf{y}^ρ : for $\rho \neq \beta$, i.e. running over black vertices skipping $\bar{J}_{\mathbf{s}} = \bar{J}_{\mathbf{y}^\beta}$. Thus the derivatives on $D_c^{(\text{dd})}$ contribute

$$\sum_{\substack{\rho \neq \beta \\ \rho, \text{any black vertex}}} \frac{1}{E(y_c^\rho, s_c)} \frac{\partial W[J, J]}{\partial \varsigma_c(\mathcal{D}; \beta, \rho)(\mathbf{X})} \text{ for } \rho \neq \beta.$$

To fully compute D_c , we add now the product of derivatives, $D_c^{(\text{prod})}$. Unlike $D_c^{(\text{dd})}$, with $D_c^{(\text{prod})}$ one can first see the effect of applying the rest of the derivatives of the graph \mathcal{R} , that is those with momenta $(\mathbf{x}_0^\alpha)_\alpha$ and $(\mathbf{y}_0^\alpha)_{\alpha \neq \beta}$. This is due to the product of derivatives

¹³The part of the calculation of the contributions of the double derivatives $\delta^2 W / \delta J \delta \bar{J}$ or $\delta^2 W / \delta \bar{J} \delta \bar{J}$ to \mathfrak{b}_c and \mathfrak{d}_c is shortened, due to the very similar derivation provided already in [PPW17].

of W , which forces both factors to be derived with respect to momenta in the same graph in order not to vanish. Thus,

$$\begin{aligned}
 & \prod_{\substack{\alpha \neq \beta \\ \nu=1, \dots, r}} \frac{\delta}{\delta \bar{J}_{\mathbf{y}_0^\alpha}} \frac{\delta}{\delta J_{\mathbf{x}_0^\nu}} (e^{-W} D_c^{(\text{prod})}(\mathbf{s})) + \mathcal{O}(\bar{J}) \\
 &= \prod_{\alpha \neq \beta; \nu} \frac{\delta}{\delta \bar{J}_{\mathbf{y}_0^\alpha}} \frac{\delta}{\delta J_{\mathbf{x}_0^\nu}} \left[\sum_{\mathbf{b}} \frac{\bar{J}_{\mathbf{b}}}{E(b_c, s_c)} \frac{\delta W}{\delta \bar{J}_{s_c b_c}} \frac{\delta W}{\delta \bar{J}_{\mathbf{b}_c s_c}} \right] \\
 &= \sum_{\rho \neq \beta}^r \prod_{\substack{\alpha \neq \beta (\alpha \neq \rho) \\ \nu=1, \dots, r}} \frac{\delta}{\delta \bar{J}_{\mathbf{y}_0^\alpha}} \frac{\delta}{\delta J_{\mathbf{x}_0^\nu}} \left[\sum_{\mathbf{b}} \frac{\delta_{\mathbf{y}^\rho}^{\mathbf{b}}}{E(b_c, s_c)} \frac{\delta W}{\delta \bar{J}_{s_c b_c}} \frac{\delta W}{\delta \bar{J}_{\mathbf{b}_c s_c}} \right] \\
 &= \sum_{\rho \neq \beta}^r \frac{1}{E(y_c^\rho, s_c)} \prod_{\substack{\alpha \neq \beta (\alpha \neq \rho) \\ \nu=1, \dots, r}} \frac{\delta}{\delta \bar{J}_{\mathbf{y}_0^\alpha}} \frac{\delta}{\delta J_{\mathbf{x}_0^\nu}} \left[\frac{\delta W}{\delta \bar{J}_{\mathbf{y}_c^\beta y_c^\rho}} \frac{\delta W}{\delta \bar{J}_{\mathbf{y}_c^\rho y_c^\beta}} \right],
 \end{aligned} \tag{4.17}$$

where in the last step we only used that $\mathbf{s} = \mathbf{y}^\beta = \mathbf{y}_0^\beta$. It is evident that these two derivatives in the square bracket form of a colour- c edge swap at \mathbf{y}_0^β and \mathbf{y}_0^ρ , but in order not to lead to a vanishing term, they also have to lie on a different component of a graph (as otherwise a graph derivative would be incomplete). It is therefore additionally required that the graph is disconnected after the swapping at \mathbf{y}_0^β and \mathbf{y}_0^ρ , that is that there are connected graphs \mathcal{R}_1 and \mathcal{R}_2 , such that

$$s_c(\mathcal{R}; \beta, \rho) = \mathcal{R}_1 \amalg \mathcal{R}_2 \quad \text{if and only if} \quad \rho \in \text{Br}(\mathcal{R}, \beta, c).$$

Therefore

$$\begin{aligned}
 & \frac{\delta^\alpha}{\delta \mathcal{Q}^\alpha(\mathbb{X})} \prod_{\substack{\alpha \neq \beta \\ \nu=1, \dots, r}} \frac{\delta}{\delta \bar{J}_{\mathbf{y}_0^\alpha}} \frac{\delta}{\delta J_{\mathbf{x}_0^\nu}} (e^{-W} D_c^{(\text{prod})}(\mathbf{s})) + \mathcal{O}(\bar{J}) \\
 &= \sum_{\tau \in \text{Br}(\mathcal{R}, \beta, c)} \frac{1}{E(y_c^\tau, s_c)} \frac{\delta^\alpha}{\delta \mathcal{Q}^\alpha(\mathbb{X})} \prod_{\substack{\alpha \neq \beta (\alpha \neq \rho) \\ \nu=1, \dots, r}} \frac{\delta}{\delta \bar{J}_{\mathbf{y}_0^\alpha}} \frac{\delta}{\delta J_{\mathbf{x}_0^\nu}} \left[\frac{\delta W}{\delta \bar{J}_{y_c^\beta y_c^\tau}} \frac{\delta W}{\delta \bar{J}_{y_c^\tau y_c^\beta}} \right]. \\
 &= \sum_{\tau \in \text{Br}(\mathcal{R}, \beta, c)} \frac{1}{E(y_c^\tau, s_c)} \frac{\delta^\alpha}{\delta \mathcal{Q}^\alpha(\mathbb{X})} \left(\frac{\delta W}{\delta \mathcal{R}_1(X'_0)} \frac{\delta W}{\delta \mathcal{R}_2(X''_0)} \right).
 \end{aligned}$$

At this place we use the multivariable graph calculus Leibniz rule (Lemma 2.12), namely

$$\begin{aligned}
 & \frac{\delta^\alpha}{\delta \mathcal{Q}^\alpha(\mathbb{X})} \prod_{\substack{\alpha \neq \beta \\ \nu=1, \dots, r}} \frac{\delta}{\delta \bar{J}_{\mathbf{y}_0^\alpha}} \frac{\delta}{\delta J_{\mathbf{x}_0^\nu}} (e^{-W} D_c^{(\text{prod})}(\mathbf{s})) \Big|_{J=\bar{J}=0} \\
 &= \sum_{\tau \in \text{Br}(\mathcal{R}, \beta, c)} \frac{1}{E(y_c^\tau, s_c)} \sum_{\substack{\mathcal{B}, \mathcal{C} \\ \mathcal{B} \amalg \mathcal{C} \sim \mathcal{Q}^\alpha}} \sum_{\Omega \in \text{Aut}_c(\mathcal{Q}^\alpha)} \\
 & \quad \Omega \cdot [\langle G_{\mathcal{R}_1|\mathcal{B}}(X'_0, \cdot) \times G_{\mathcal{R}_1|\mathcal{C}}(X''_0, \cdot) \rangle_{\mathcal{Q}^\alpha}](\mathbb{X})
 \end{aligned}$$

$$= \sum_{\tau \in \text{Br}(\mathcal{R}, \beta, c)} \frac{1}{E(y_c^\tau, s_c)} \sum_{\substack{\mathcal{B}, \mathcal{C} \\ \mathcal{B} \amalg \mathcal{C} \sim \mathcal{Q}^\alpha}} \sum_{\Omega \in \text{Aut}_c(\mathcal{Q}^\alpha)} (\Omega \cdot H_{\mathcal{B}, \mathcal{C}}^{(c, \tau)})(\mathbb{X}).$$

In summary, the \mathfrak{d}_c -term is given by

$$\begin{aligned} \mathfrak{d}_c(\mathbf{X}; \mathbf{s}; \mathcal{D}) &= \sum_{\substack{\rho \neq \beta \\ \rho, \text{any black vertex}}} \frac{1}{E(y_c^\rho, s_c)} \frac{\partial W[J, \bar{J}]}{\partial \varsigma_c(\mathcal{D}; 1, \rho)}(\mathbf{X}) \\ &+ \sum_{\tau \in \text{Br}(\mathcal{R}, \beta, c)} \frac{1}{E(y_c^\tau, s_c)} \sum_{\substack{\mathcal{B}, \mathcal{C} \\ \mathcal{B} \amalg \mathcal{C} \sim \mathcal{Q}^\alpha}} \sum_{\Omega \in \text{Aut}_c(\mathcal{Q}^\alpha)} (\Omega \cdot H_{\mathcal{B}, \mathcal{C}})(\mathbb{X}). \end{aligned} \quad (4.18)$$

As for the derivatives on $B_c(\mathbf{s})$, we divide the derivation in two parts. One concerns the double derivative, $B_c^{(\text{dd})}$:

$$\begin{aligned} \mathcal{O}(J) &+ \prod_{\substack{\alpha \neq \beta \\ \nu=1, \dots, d}} \frac{\delta}{\delta \bar{J}_{\mathbf{y}^\alpha}} \frac{\delta}{\delta J_{\mathbf{x}^\nu}} B_c^{(\text{dd})}(\mathbf{s}) \\ &= \prod_{\alpha \neq \beta; \nu} \frac{\delta}{\delta \bar{J}_{\mathbf{y}^\alpha}} \frac{\delta}{\delta J_{\mathbf{x}^\nu}} \left[\sum_{\mathbf{b}} \frac{1}{E(b_c, s_c)} J_{\mathbf{b} \varepsilon s_c} \frac{\delta^2 W[J, \bar{J}]}{\delta \bar{J}_{\mathbf{s}_c b_c} \delta J_{\mathbf{b}}} \right] \\ &= \sum_{\theta=1}^d \prod_{\alpha \neq \beta; \nu \neq \theta} \frac{\delta}{\delta \bar{J}_{\mathbf{y}^\alpha}} \frac{\delta}{\delta J_{\mathbf{x}^\nu}} \left[\sum_{b_c} \frac{1}{E(b_c, s_c)} \delta_{x_c^{s_c}} \frac{\delta^2 W[J, \bar{J}]}{\delta \bar{J}_{\mathbf{s}_c b_c} \delta J_{\mathbf{x}_c^\beta b_c}} \right] \\ &= \prod_{\alpha \neq \beta; \nu \neq \gamma} \frac{\delta}{\delta \bar{J}_{\mathbf{y}^\alpha}} \frac{\delta}{\delta J_{\mathbf{x}^\nu}} \left[\sum_{b_c} \frac{1}{E(b_c, x_c^\gamma)} \frac{\delta^2 W[J, \bar{J}]}{\delta \bar{J}_{\mathbf{s}_c b_c} \delta J_{\mathbf{x}_c^\beta b_c}} \right]. \end{aligned} \quad (4.19)$$

since there is a single vertex \mathbf{x}^γ , $\gamma = \gamma(c)$, with $x_c^\gamma = s_c$, so $\delta_{x_c^{s_c}} = \delta_{x_c^\theta}^{s_c} \delta_\theta^\gamma$. The term $\delta^2 W[J, \bar{J}] / \delta \bar{J}_{\mathbf{s}_c b_c} \delta J_{\mathbf{x}_c^\beta b_c}$, is selected by $\delta_{x_c^{s_c}}^{s_c}$ leads to $\partial Z / \partial \mathcal{D}(\mathbf{X})|_{x_c^\gamma \rightarrow b_c}$, after taking all the rest of derivatives, with the single coordinate x_c^γ being substituted by (the running) b_c . But when

$$b_c \in \{y_c^1, y_c^2, \dots, \widehat{y_c^\beta}, \dots, y_c^d\},$$

one does not have exactly a ‘graph derivative’, since we are evaluating it not in $\mathcal{F}_{D,d}$, but in one of its diagonals of colour c . A direct computation yields then a second contribution to $\mathfrak{b}_c^{(\text{dd})}$ (the third and fourth lines below):

$$\begin{aligned} \mathfrak{b}_c^{(\text{dd})}(\mathbf{X}; \mathbf{s}; \mathcal{D}) &= \sum_{b_c} \frac{1}{E(b_c, x_c^\gamma)} \\ &\quad \times G_{\mathcal{D}}^{(2k)}(\mathbf{x}^1, \dots, \mathbf{x}^{\gamma-1}, (x_1^\gamma, \dots, x_{a-1}^\gamma, b_c, x_{a+1}^\gamma, \dots, x_D^\gamma), \mathbf{x}^{\gamma+1}, \dots, \mathbf{x}^d) \\ &+ \sum_{\rho > 1} \frac{1}{E(x_c^{\kappa(\rho)}, x_c^\gamma)} \\ &\quad \times \frac{\partial W[J, \bar{J}]}{\partial \varsigma_c(\mathcal{D}; 1, \rho)}(\mathbf{x}^1, \dots, (x_1^\gamma, \dots, x_{a-1}^\gamma, x_c^{\kappa(\rho)}, x_{a+1}^\gamma, \dots, x_D^\gamma), \dots, \mathbf{x}^d). \end{aligned} \quad (4.20)$$

where $\kappa(\rho)$ is defined in Figure 3 (i.e. $x_c^{\kappa(\rho)} = y_c^\rho$).

The last computation is $\mathfrak{b}_c^{(\text{prod})}$,

$$\begin{aligned} & \mathfrak{b}_c^{(\text{prod})}(\mathbf{X}; \mathbf{s}; \mathcal{D}) \\ &= \frac{\delta^\alpha}{\delta \mathcal{Q}^\alpha(\mathbb{X})} \prod_{\substack{\alpha \neq \beta \\ 0 \leq \alpha \leq r \\ \nu=1, \dots, r}} \frac{\delta}{\delta \bar{J}_{\mathbf{y}_0^\alpha}} \frac{\delta}{\delta J_{\mathbf{x}_0^\nu}} \left[\sum_{\mathbf{b}} \frac{J_{\mathbf{b}_c s_c}}{E(b_c, s_c)} \frac{\delta W[J, \bar{J}]}{\delta J_{\mathbf{b}}} \frac{\delta W[J, \bar{J}]}{\delta \bar{J}_{s_c b_c}} \right] \Big|_{J=\bar{J}=0}. \end{aligned}$$

This computation is quite similar to the one for the $\mathfrak{d}_c^{(\text{prod})}$ -term, presented above, with the only difference that the evaluation is not at \mathbb{X} , but at $\mathbb{X}|_{s_c \rightarrow y_c^\tau}$. The less trivial part in that derivation is to figure out, which non-zero contributions come from

$$\frac{\delta W[J, \bar{J}]}{\delta J_{b_c \mathbf{x}_c^\gamma}} \frac{\delta W[J, \bar{J}]}{\delta \bar{J}_{s_c b_c}}. \quad (4.21)$$

Because of the repetition of b_c in both factors, it seems that the term vanishes after deriving it. However, b_c runs, and it does so also through the particular c -coloured entries of the black vertices of \mathcal{R} ,

$$b_c \in \{y_{0,c}^1, y_{0,c}^2, \dots, \widehat{y_{0,c}^\beta}, \dots, y_{0,c}^d\}.$$

Only if we also require that $b_c \in \{y_c^\tau | \tau \in \text{Br}(\mathcal{R}, \beta, c)\}$, we guarantee that each one of those factors forms a graph derivative. However, notice that momentum in the white vertex $\mathbf{x}^\gamma = (s_c \mathbf{x}_c^\gamma)$ has changed to $(y_c^\tau \mathbf{x}_c^\gamma)$. Thus, one changes X_0 into $X_0|_{s_c \rightarrow y_c^\tau}$. Therefore

$$\mathfrak{b}_c^{(\text{prod})}(\mathbf{X}; \mathbf{s}; \mathcal{D}) = \sum_{\tau \in \text{Br}(\mathcal{R}, \beta, c)} \frac{1}{E(y_c^\tau, s_c)} \frac{\delta^\alpha}{\delta \mathcal{Q}^\alpha(\mathbb{X})} \frac{\delta W}{\delta \mathcal{R}_1(X_0')} \frac{\delta W}{\delta \mathcal{R}_2(X_0'')} \Big|_{\substack{J=\bar{J}=0 \\ s_c \rightarrow y_c^\tau}}.$$

We apply again Lemma 2.12 and find that

$$\mathfrak{b}_c^{(\text{prod})}(\mathbf{X}; \mathbf{s}; \mathcal{D}) = \sum_{\tau \in \text{Br}^{(2)}(\mathcal{R}; \beta, c)} \sum_{\substack{(\mathcal{C}, \mathcal{B}) \in \mathbb{G}_D \amalg \mathbb{G}_D \\ (\mathcal{C} \amalg \mathcal{B}) \sim \mathcal{Q}}} \sum_{\Omega \in \text{Aut}_c(\mathcal{Q})} \frac{(\Omega \cdot H_{\mathcal{B}, \mathcal{C}})(\mathbb{X})|_{s_c \rightarrow y_c^\tau}}{E(y_c^\tau, s_c)}.$$

Due to eqs. (4.7), one has

$$\begin{aligned} \frac{\partial W[J, \bar{J}]}{\partial \mathcal{D}(\mathbf{X})} &= \prod_i \frac{\delta^{\alpha_i}}{\delta \mathcal{Q}_i(X_1^i) \cdots \mathcal{Q}_i(X_{\alpha_i}^i)} \left\{ \prod_{\substack{\alpha \neq \beta \\ \nu=1, \dots, k}} \frac{\delta}{\delta \bar{J}_{\mathbf{y}_0^\alpha}} \frac{\delta}{\delta J_{\mathbf{x}_0^\nu}} \left((-2\lambda E_{\mathbf{s}}^{-1}) \right. \right. \\ &\quad \left. \left. \times \sum_{c=1}^D e^{-W} [A_c(\mathbf{s}) + C_c(\mathbf{s}) + D_c(\mathbf{s}) + F_c(\mathbf{s}) - B_c(\mathbf{s})] \right) \right\} \Big|_{\substack{J=0 \\ \bar{J}=0}} \\ &= \frac{(-2\lambda)}{E_{\mathbf{s}}} \sum_c \left(\mathfrak{a}_c(\mathbf{X}; \mathbf{s}; \mathcal{D}) + \mathfrak{c}_c(\mathbf{X}; \mathbf{s}; \mathcal{D}) + \mathfrak{d}_c^{(\text{dd})}(\mathbf{X}; \mathbf{s}; \mathcal{D}) \right. \\ &\quad \left. + \mathfrak{d}_c^{(\text{prod})}(\mathbf{X}; \mathbf{s}; \mathcal{D}) + \mathfrak{f}_c(\mathbf{X}; \mathbf{s}; \mathcal{D}) \right. \\ &\quad \left. - \mathfrak{b}_c^{(\text{dd})}(\mathbf{X}; \mathbf{s}; \mathcal{D}) - \mathfrak{b}_c^{(\text{prod})}(\mathbf{X}; \mathbf{s}; \mathcal{D}) \right). \end{aligned}$$

This is precisely $G_{\mathcal{D}}(\mathbf{X})$ and the result follows. \square

5. FOUR AND SIX POINT SDE WITH DISCONNECTED BOUNDARY

Concrete SDE's for the rank-3 φ_3^4 -theory are presented next. Recall, the interaction in this case is $\lambda(1\boxed{1} + 2\boxed{2} + 3\boxed{3})$. We display some of the equations in traditional notation with explicit graphs, which allows to see immediately the graph operations. In other equations we use the simplification clarified in Table 1.

CORRELATION FUNCTIONS		
Order	Graph notation	Simplified notation
2-pt function	$G_{\ominus}^{(2)}$	$G_m^{(2)}$ or $G^{(2)}$
4-pt functions	$G_{a\boxed{a}}^{(4)}$	$G_{V_a}^{(4)}$
	$G_{\ominus \ominus}^{(4)}$	$G_{m m}^{(4)}$
6-pt functions	$G_{\text{graph}}^{(6)}$	$G_{Q_a}^{(6)}$
	$G_{\text{graph}}^{(6)}$	$G_{K_{3,3}}^{(6)}$
	$G_{\text{graph}}^{(6)}$	$G_{F_{a;bc}}^{(6)}$
	$G_{\ominus a\boxed{a}}^{(6)}$	$G_{m V_a}^{(6)}$
	$G_{\ominus \ominus \ominus}^{(6)}$	$G_{m m m}^{(6)}$

TABLE 1. Two notations for the correlation functions. Here a, b, c are colours bound to satisfy $\{a, b, c\} = \{1, 2, 3\}$. The subindex m originates from ‘melon’.

5.1. Schwinger-Dyson equations for $G_{\ominus|i\boxed{i}}^{(6)}$. We single out the terms in the derivation of the SDE for this case, which is the most complicated presented here. The rest of the results are obtained in a similar and more direct way.

There are two equations, depending on whether one chooses \mathbf{s} (cf. Theorem 4.1 above) as a component of the outgoing momentum in \ominus or in $i\boxed{i}$. We choose this last vertex to be $V_1 = 1\boxed{1}$, for sake of clarity (since the model is colour-invariant, SDE for the other colours are readily obtained from it).

- *If $\mathbf{s} = \mathbf{x}$ is outgoing momentum of the graph \ominus .* In the notation of the theorem, here $\mathcal{D} = m|V_1$ being $\mathcal{R} = m$, since $\mathbf{y}^1 = \mathbf{s} = X_0$ is the momentum of the black vertex of m . Therefore $\mathcal{Q} = V_1$. The remaining momenta \mathbb{X} equal (\mathbf{y}, \mathbf{z}) . The $I_{\mathcal{C}, \mathcal{B}}^{(c)}$ -terms are then computed as follows, from any of the factorisations $\mathcal{C}, \mathcal{B} = (\emptyset, V_1)$ or (V_1, \emptyset) and read

$$I_{\mathcal{C}, \mathcal{B}}^{(c)}(\mathbb{X}) = \frac{1}{|\text{Aut}_c(\mathcal{B})|} \langle \mathfrak{f}_{\mathcal{C}; s_c}^{(c)} \times G_{m|\mathcal{B}}(X_0; \cdot) \rangle_{V_1}(\mathbb{X}) \quad (5.1)$$

$$= \begin{cases} \mathfrak{f}_{V_1; s_c}^{(c)}(\mathbf{y}, \mathbf{z}) G_m^{(2)}(\mathbf{x}) & (\mathcal{C}, \mathcal{B}) = (V_1, \emptyset), \\ \frac{1}{2} \mathfrak{f}_{\emptyset; s_c}^{(c)} G_{m|V_1}^{(6)}(\mathbf{x}, \mathbf{y}, \mathbf{z}) & (\mathcal{C}, \mathcal{B}) = (\emptyset, V_1). \end{cases} \quad (5.2)$$

On these functions $\hat{\sigma} \in \text{Aut}_c(V_1)$ acts exchanging \mathbf{y} with \mathbf{z} ; just as on the terms coming from the derivative of the Y -term with respect to $(\mathfrak{m}|V_1)$:

$$\sum_{\hat{\sigma} \in \mathbb{Z}_2} \sigma \cdot \mathfrak{f}_{\mathfrak{m}|V_1; x_c}^{(c)}(\mathbf{x}, \mathbf{y}, \mathbf{z}) = \mathfrak{f}_{\mathfrak{m}|V_1; x_c}^{(c)}(\mathbf{x}, \mathbf{y}, \mathbf{z}) + \mathfrak{f}_{\mathfrak{m}|V_1; x_c}^{(c)}(\mathbf{y}, \mathbf{x}, \mathbf{z}). \quad (5.3)$$

Next, we obtain the second line in eq. (4.4) (the ‘swap-term’). We have chosen $\mathbf{y}^1 = \mathbf{s}$ ($\beta = 1$), so

$$\frac{\partial W[J, \bar{J}]}{\partial \varsigma_{c=1}(\mathfrak{m}|V_1; \beta = 1, \rho = 2, 3)}(\mathbf{x}, \mathbf{y}, \mathbf{z}) = G_{Q_1}^{(6)}(\mathbf{x}, \mathbf{y}, \mathbf{z}) \quad (5.4a)$$

$$\frac{\partial W[J, \bar{J}]}{\partial \varsigma_{c=2}(\mathfrak{m}|V_1; \beta = 1, \rho = 2)}(\mathbf{x}, \mathbf{y}, \mathbf{z}) = G_{F_{3;21}}^{(6)}(\mathbf{x}, \mathbf{z}, \mathbf{y}) \quad (5.4b)$$

$$\frac{\partial W[J, \bar{J}]}{\partial \varsigma_{c=2}(\mathfrak{m}|V_1; \beta = 1, \rho = 3)}(\mathbf{x}, \mathbf{y}, \mathbf{z}) = G_{F_{3;21}}^{(6)}(\mathbf{x}, \mathbf{y}, \mathbf{z}) \quad (5.4c)$$

$$\frac{\partial W[J, \bar{J}]}{\partial \varsigma_{c=3}(\mathfrak{m}|V_1; \beta = 1, \rho = 2)}(\mathbf{x}, \mathbf{y}, \mathbf{z}) = G_{F_{2;31}}^{(6)}(\mathbf{x}, \mathbf{z}, \mathbf{y}) \quad (5.4d)$$

$$\frac{\partial W[J, \bar{J}]}{\partial \varsigma_{c=2}(\mathfrak{m}|V_1; \beta = 1, \rho = 3)}(\mathbf{x}, \mathbf{y}, \mathbf{z}) = G_{F_{2;31}}^{(6)}(\mathbf{x}, \mathbf{y}, \mathbf{z}) \quad (5.4e)$$

In this case the set $\text{Br}(\ominus, \rho, c)$ is empty, for any values of ρ and c . Therefore, the sum over the H -terms vanishes. For each $(\mathbf{x}, \mathbf{y}, \mathbf{z}) \in \mathcal{F}_{3,3}$,

$$\begin{aligned} & G_{\mathfrak{m}|V_1}^{(6)}(\mathbf{x}, \mathbf{y}, \mathbf{z}) \quad (5.5) \\ &= \left(\frac{-2\lambda}{E_{\mathbf{x}}} \right) \times \left\{ \sum_{c=1}^3 \mathfrak{f}_{\mathfrak{m}|V_1; x_c}^{(c)}(\mathbf{x}, \mathbf{y}, \mathbf{z}) + \mathfrak{f}_{\mathfrak{m}|V_1; x_c}^{(c)}(\mathbf{y}, \mathbf{x}, \mathbf{z}) \right. \\ &+ \frac{1}{E(y_1, x_1)} [G_{Q_1}^{(6)}(\mathbf{x}, \mathbf{y}, \mathbf{z}) - G_{Q_1}^{(6)}(y_1, x_2, x_3, \mathbf{y}, \mathbf{z})] \\ &+ \frac{1}{E(z_1, x_1)} [G_{Q_1}^{(6)}(\mathbf{x}, \mathbf{y}, \mathbf{z}) - G_{Q_1}^{(6)}(z_1, x_2, x_3, \mathbf{y}, \mathbf{z})] \\ &+ \frac{1}{E(z_2, x_2)} [G_{F_{3;21}}(\mathbf{x}, \mathbf{z}, \mathbf{y}) - G_{F_{3;21}}(x_1, z_2, x_3, \mathbf{z}, \mathbf{y})] \\ &+ \frac{1}{E(y_2, x_2)} [G_{F_{3;21}}(\mathbf{x}, \mathbf{y}, \mathbf{z}) - G_{F_{3;21}}(x_1, y_2, x_3, \mathbf{y}, \mathbf{z})] \\ &+ \frac{1}{E(z_3, x_3)} [G_{F_{2;31}}(\mathbf{x}, \mathbf{z}, \mathbf{y}) - G_{F_{2;31}}(x_1, x_2, z_3, \mathbf{z}, \mathbf{y})] \\ &+ \frac{1}{E(y_3, x_3)} [G_{F_{2;31}}(\mathbf{x}, \mathbf{y}, \mathbf{z}) - G_{F_{2;31}}(x_1, x_2, y_3, \mathbf{y}, \mathbf{z})] \\ &- \sum_{c=1}^3 \left[\sum_{b_c} \frac{1}{E(x_c, b_c)} [G_{\mathfrak{m}|V_1}^{(6)}(\mathbf{x}, \mathbf{y}, \mathbf{z}) - G_{\mathfrak{m}|V_1}^{(6)}(x_c b_c, \mathbf{y}, \mathbf{z})] \right. \\ &\left. + [(\mathfrak{f}_{V_1; s_c}^{(c)}(\mathbf{z}, \mathbf{y}) + \mathfrak{f}_{V_1; s_c}^{(c)}(\mathbf{y}, \mathbf{z})) \cdot G^{(2)}(\mathbf{x}) + \mathfrak{f}_{\emptyset; s_c}^{(c)} G_{\mathfrak{m}|V_1}^{(6)}(\mathbf{x}, \mathbf{y}, \mathbf{z})] \right\}. \end{aligned}$$

- If $\mathbf{s} = \mathbf{x}$ is outgoing momentum of the boundary graph $1\overline{\square}1$. For $\mathbf{s} = (x_1, y_2, y_3)$ an outgoing momentum of V_1 we derive now the SDE for $G_{\mathbf{m}|V_1}^{(6)}$. Then $\mathcal{R} = V_1$, $\mathcal{Q} = \mathbf{m}$, by definition. The $I_{\mathcal{C},\mathcal{B}}^{(c)}$ -coefficients are given by

$$\begin{aligned} I_{\mathbf{m},\emptyset}^{(c)}(\mathbf{x}, \mathbf{y}, \mathbf{z}) &= \mathfrak{f}_{\mathbf{m};s_c}^{(c)}(\mathbf{z}) \cdot G_{V_1}(\mathbf{x}, \mathbf{y}), \\ I_{\emptyset,\mathbf{m}}^{(c)}(\mathbf{x}, \mathbf{y}, \mathbf{z}) &= \mathfrak{f}_{\emptyset;s_c}^{(c)} \cdot G_{V_1|\mathbf{m}}(\mathbf{x}, \mathbf{y}, \mathbf{z}). \end{aligned}$$

The $H_{\mathcal{C},\mathcal{B}}^{(c,\tau)}$ -terms are computed from the set

$$\text{Br}(1\overline{\square}1, \beta = 1, c) = \begin{cases} \{2\} & c = 1, \\ \emptyset & \text{otherwise,} \end{cases}$$

since only the colour-1 swap at the vertex \mathbf{s} with the vertex with outgoing momentum \mathbf{y}^2 (also in V_1) separates $1\overline{\square}1$. The only contributions are therefore

$$\begin{aligned} H_{\emptyset,\mathbf{m}}^{(c=1,\tau=2)}(\mathbf{X}) &= G^{(2)}(\mathbf{x}) \cdot G_{\mathbf{m}|\mathbf{m}}^{(4)}(\mathbf{y}, \mathbf{z}), \\ H_{\mathbf{m},\emptyset}^{(c=1,\tau=2)}(\mathbf{X}) &= G_{\mathbf{m}|\mathbf{m}}^{(4)}(\mathbf{x}, \mathbf{z}) \cdot G^{(2)}(\mathbf{y}). \end{aligned}$$

Thus $G_{V_1|\mathbf{m}}^{(6)}$ satisfies, for all $(\mathbf{x}, \mathbf{y}, \mathbf{z}) \in \mathcal{F}_{3,3}$,

$$\begin{aligned} &G_{V_1|\mathbf{m}}^{(6)}(\mathbf{x}, \mathbf{y}, \mathbf{z}) \tag{5.6} \\ &= \left(\frac{-2\lambda}{E_s} \right) \times \left\{ \sum_{c=1}^3 \mathfrak{f}_{\mathbf{m}|V_1;s_c}^{(c)}(\mathbf{z}, \mathbf{x}, \mathbf{y}) + \mathfrak{f}_{\mathbf{m}|V_1;s_c}^{(c)}(\mathbf{z}, \mathbf{y}, \mathbf{x}) \right. \\ &+ \frac{1}{E(y_1, x_1)} [G_{\mathbf{m}|\mathbf{m}|\mathbf{m}}^{(6)}(\mathbf{x}, \mathbf{y}, \mathbf{z}) - G_{\mathbf{m}|\mathbf{m}|\mathbf{m}}^{(6)}(y_1, x_2, x_3, \mathbf{y}, \mathbf{z})] \\ &+ \frac{1}{E(z_1, x_1)} [G_{Q_1}^{(6)}(\mathbf{x}, \mathbf{y}, \mathbf{z}) - G_{Q_1}^{(6)}(z_1, x_2, x_3; \mathbf{y}, \mathbf{z})] \\ &+ \frac{1}{E(x_2, y_2)} [G_{V_3|\mathbf{m}}^{(6)}(\mathbf{x}, \mathbf{y}, \mathbf{z}) - G_{V_3|\mathbf{m}}^{(6)}(x_1, y_2, x_3; \mathbf{y}, \mathbf{z})] \\ &+ \frac{1}{E(z_2, y_2)} [G_{F_{3;12}}^{(6)}(\mathbf{x}, \mathbf{y}, \mathbf{z}) - G_{F_{3;12}}^{(6)}(\mathbf{x}; y_1, x_2, y_3; \mathbf{z})] \\ &+ \frac{1}{E(x_3, y_3)} [G_{V_2|\mathbf{m}}^{(6)}(\mathbf{x}, \mathbf{y}, \mathbf{z}) - G_{V_2|\mathbf{m}}^{(6)}(\mathbf{x}; y_1, y_2, x_3; \mathbf{z})] \\ &+ \frac{1}{E(z_3, y_3)} [G_{F_{2;13}}^{(6)}(\mathbf{x}, \mathbf{y}, \mathbf{z}) - G_{F_{2;13}}^{(6)}(\mathbf{x}; y_1, y_2, z_3; \mathbf{z})] \\ &- \sum_{c=1}^3 \left[\sum_{b_c} \frac{1}{E(s_c, b_c)} [G_{V_1|\mathbf{m}}^{(6)}(\mathbf{x}, \mathbf{y}, \mathbf{z}) - G_{V_1|\mathbf{m}}^{(6)}([\mathbf{x}, \mathbf{y}, \mathbf{z}]|_{s_c \rightarrow b_c})] \right. \\ &\left. + [(\mathfrak{f}_{V_1;s_c}^{(c)}(\mathbf{z}, \mathbf{y}) + \mathfrak{f}_{V_1;s_c}^{(c)}(\mathbf{y}, \mathbf{z})) \cdot G^{(2)}(\mathbf{x}) + \mathfrak{f}_{\emptyset;s_c}^{(c)} G_{\mathbf{m}|V_1}^{(6)}(\mathbf{x}, \mathbf{y}, \mathbf{z})] \right\}, \end{aligned}$$

5.2. **Schwinger-Dyson equation for $G_{\Theta|\Theta}^{(4)}$.** There is only one SDE for the ‘disconnected- ∂ ’ 4-point function. For every $(\mathbf{x}, \mathbf{y}, \mathbf{z}) \in \mathcal{F}_{3,3}$,

$$\begin{aligned}
 & G_{\Theta|\Theta}^{(4)}(\mathbf{x}, \mathbf{y}) \\
 &= \left(\frac{-2\lambda}{E_{\mathbf{x}}} \right) \times \sum_{c=1}^3 \left\{ \sum_{\mathbf{q}_{\hat{c}}} G_{\Theta}^{(2)}(x_c, \mathbf{q}_{\hat{c}}) \cdot G_{\Theta|\Theta}^{(4)}(\mathbf{x}, \mathbf{y}) + G_{\Theta}^{(2)}(\mathbf{x}) f_{\Theta, x_c}^{(c)}(\mathbf{y}) \right. \\
 &+ \sum_{b_c} \frac{1}{E(b_c, x_c)} [G_{\Theta|\Theta}^{(4)}(\mathbf{x}, \mathbf{y}) - G_{\Theta|\Theta}^{(4)}(b_c \mathbf{x}_{\hat{c}}, \mathbf{y})] \\
 &+ \frac{1}{E(y_c, x_c)} [G_{c|\square_c}^{(4)}(\mathbf{x}, \mathbf{y}) - G_{c|\square_c}^{(4)}(b_c \mathbf{x}_{\hat{c}}, \mathbf{y})] \\
 &\left. + f_{x_c, \Theta|\Theta}^{(c)}(\mathbf{x}, \mathbf{y}) + f_{x_c, \Theta|\Theta}^{(c)}(\mathbf{y}, \mathbf{x}) \right\}. \tag{5.7}
 \end{aligned}$$

Only this equation is not new, but was already (directly) derived in [PPTW18], in notation of Table 1.

5.3. **Schwinger-Dyson equation for $G_{\Theta|\Theta|\Theta}^{(6)}$.** Similarly, since one can permute the arguments of $G_{\mathbf{m}|\mathbf{m}|\mathbf{m}}^{(6)}$, it satisfies only one SDE:

$$\begin{aligned}
 & \left(1 + \frac{2\lambda}{E_{\mathbf{x}}} \sum_{c=1}^3 \sum_{\mathbf{q}_{\hat{c}}} G_{\Theta}^{(2)}(x_c, \mathbf{q}_{\hat{c}}) \right) \times G_{\Theta|\Theta|\Theta}^{(6)}(\mathbf{x}, \mathbf{y}, \mathbf{z}) \\
 &= \left(\frac{-2\lambda}{E_{\mathbf{x}}} \right) \sum_{c=1}^3 \left\{ f_{\Theta; x_c}^{(c)}(\mathbf{y}) G_{\Theta|\Theta}^{(4)}(\mathbf{x}, \mathbf{z}) + f_{\Theta; x_c}^{(c)}(\mathbf{z}) G_{\Theta|\Theta}^{(4)}(\mathbf{x}, \mathbf{y}) \right. \\
 &+ G_{\Theta}^{(2)}(\mathbf{x}) \cdot f_{\Theta|\Theta}^{(c)}(\mathbf{y}, \mathbf{z}) \\
 &- \sum_{b_c} \frac{1}{E(x_c, b_c)} [G_{\Theta|\Theta|\Theta}^{(6)}(\mathbf{x}, \mathbf{y}, \mathbf{z}) - G_{\Theta|\Theta|\Theta}^{(6)}(b_c \mathbf{x}_{\hat{c}}, \mathbf{y}, \mathbf{z})] \\
 &+ \frac{1}{E(y_c, x_c)} [G_{\Theta|\square|\square}^{(6)}(\mathbf{z}, \mathbf{x}, \mathbf{y}) - G_{\Theta|\square|\square}^{(6)}(\mathbf{z}, y_c \mathbf{x}_{\hat{c}}, \mathbf{y})] \\
 &+ \frac{1}{E(z_c, x_c)} [G_{\Theta|\square|\square}^{(6)}(\mathbf{y}, \mathbf{x}, \mathbf{z}) - G_{\Theta|\square|\square}^{(6)}(\mathbf{y}, y_c \mathbf{x}_{\hat{c}}, \mathbf{z})] \\
 &\left. + \sum_{\sigma \in \mathfrak{S}(3)} \sigma \cdot f_{\Theta|\Theta|\Theta}^{(c)}(\mathbf{x}, \mathbf{y}, \mathbf{z}) \right\}. \tag{5.8}
 \end{aligned}$$

We kept the graph notation in order to ease the reading of the graph movements. Equivalently,

$$\begin{aligned}
 & \left(1 + \frac{2\lambda}{E_{\mathbf{x}}} \sum_{c=1}^3 \sum_{\mathbf{q}_{\hat{c}}} G^{(2)}(x_c, \mathbf{q}_{\hat{c}}) \right) \times G_{\mathbf{m}|\mathbf{m}|\mathbf{m}}^{(6)}(\mathbf{x}, \mathbf{y}, \mathbf{z}) \\
 &= \left(\frac{-2\lambda}{E_{\mathbf{x}}} \right) \sum_{c=1}^3 \left\{ f_{\mathbf{m}; x_c}^{(c)}(\mathbf{y}) G_{\mathbf{m}|\mathbf{m}}^{(4)}(\mathbf{x}, \mathbf{z}) + f_{\mathbf{m}; x_c}^{(c)}(\mathbf{z}) G_{\mathbf{m}|\mathbf{m}}^{(4)}(\mathbf{x}, \mathbf{y}) \right.
 \end{aligned}$$

$$\begin{aligned}
& + G_{\ominus}^{(2)}(\mathbf{x}) \cdot \mathfrak{f}_{\mathbf{m}|\mathbf{m}}^{(c)}(\mathbf{y}, \mathbf{z}) \\
& - \sum_{b_c} \frac{1}{E(x_c, b_c)} [G_{\mathbf{m}|\mathbf{m}|\mathbf{m}}^{(6)}(\mathbf{x}, \mathbf{y}, \mathbf{z}) - G_{\mathbf{m}|\mathbf{m}|\mathbf{m}}^{(6)}(b_c \mathbf{x}_{\hat{c}}, \mathbf{y}, \mathbf{z})] \\
& + \frac{1}{E(y_c, x_c)} [G_{\mathbf{m}|V_c}^{(6)}(\mathbf{z}, \mathbf{x}, \mathbf{y}) - G_{\mathbf{m}|V_c}^{(6)}(\mathbf{z}, y_c \mathbf{x}_{\hat{c}}, \mathbf{y})] \\
& + \frac{1}{E(z_c, x_c)} [G_{\mathbf{m}|V_c}^{(6)}(\mathbf{y}, \mathbf{x}, \mathbf{z}) - G_{\mathbf{m}|V_c}^{(6)}(\mathbf{y}, y_c \mathbf{x}_{\hat{c}}, \mathbf{z})] \\
& + \sum_{\sigma \in \mathfrak{S}(3)} \sigma \cdot \mathfrak{f}_{\mathbf{m}|\mathbf{m}|\mathbf{m}}^{(c)}(\mathbf{x}, \mathbf{y}, \mathbf{z}) \Big\}.
\end{aligned} \tag{5.9}$$

6. TOWARDS HIGHER-DIMENSIONAL TUTTE EQUATIONS

We compare the result with matrix models loop equations and, their equivalent, Tutte equations that count discrete surfaces.

6.1. Tutte equations and matrix models. This material is based on the exposition by Eynard [Eyn16, Ch. 1 & 2]. There, three facts are proven:

- (1) The generating functions $\mathcal{T}_{l_1+1, l_2, \dots, l_\kappa}$ of connected *maps*¹⁴ with κ marked faces (boundaries) of perimeters $l_1 + 1, \dots, l_\kappa$ satisfy Tutte equations:

$$\mathcal{T}_{l_1+1}^{(0)} = \sum_{j=3}^d \lambda_j \mathcal{T}_{l_1-j-1}^{(0)} + \sum_{l_1+l_2=l-1} \mathcal{T}_{l_1}^{(0)} \mathcal{T}_{l_2}^{(0)} \tag{6.1a}$$

in the planar, single-boundary ($\kappa = 1$) case, while for all $g \geq 1$, setting $K = \{l_2, \dots, l_\kappa\}$,

$$\begin{aligned}
\mathcal{T}_{l_1+1, K}^{(g)} & = \sum_{\alpha=3}^d \lambda_\alpha \mathcal{T}_{l_1+\alpha-1, K}^{(g)} + \sum_{m=2}^\kappa l_m \mathcal{T}_{l_1+l_m-1, K \setminus \{l_m\}}^{(g)} \\
& + \sum_{j=0}^{l_1-1} \left[\mathcal{T}_{j, l_1-1-j, K}^{(g-1)} + \sum_{\substack{g_1+g_2=g \\ J \subset K}} \mathcal{T}_{j, J}^{(g_1)} \times \mathcal{T}_{l_1-1-j, K \setminus J}^{(g_2)} \right].
\end{aligned} \tag{6.1b}$$

In the last two equations the superindex h in $\mathcal{T}_{\dots}^{(h)}$ means the restriction to genus- h maps. Also the formal variables λ_α count the number n_α of α -gons in each map. That is, the $t^{v(\mathbf{m})} \lambda_3^{n_3(\mathbf{m})} \dots \lambda_d^{n_d(\mathbf{m})}$ -coefficient of $\mathcal{T}_{l_1, \dots, l_\kappa}^{(h)}$ counts, modulo automorphisms, how many genus- h maps \mathbf{m} are there, having precisely $n_\alpha(\mathbf{m})$ α -gons with marked faces of lengths $l_m > 0$ ($\alpha = 3, \dots, d$; $m = 1, \dots, \kappa$). To render this number finite, the variable t counts the number of vertices $v(\mathbf{m})$ of the map \mathbf{m} . These \mathcal{T} -generating functions are not independent but related via Tutte equations — and in fact obey a rather universal relation known as Topological Recursion.

¹⁴A *map* is a concept slightly more general than a gluing of a collection of n_α α -gons by their sides ($\alpha \geq 3$). Maps might have certain number κ of marked faces of perimeters $l_1, \dots, l_\kappa \geq 0$. Their Euler characteristic is $\chi = \#\text{vertices} - \#\text{edges} + \#\text{unmarked faces} = 2 - 2g - \kappa$, being g the *genus* of the map. The precise concept will not be needed here and we refer to [Eyn16, Sect. 1.1.2] for the definition in terms of permutations.

(2) The matrix model

$$Z = \int_{\text{formal}} dM e^{-\frac{N}{t} \left[\frac{M^2}{2} - V(M) \right]}, \quad V(x) = \sum_{\alpha=3}^d \lambda_\alpha \frac{x^\alpha}{\alpha}, \quad (6.2)$$

satisfies Migdal's loop equations [Mig83]

$$\begin{aligned} & \sum_{j=0}^{l_1-1} \langle \text{Tr}(M^j) \text{Tr}(M^{l_1-1-j}) \rangle + \sum_{m=2}^{\kappa} l_m \langle \text{Tr}(M^{l_1-1+l_j} \prod_{\substack{i=2 \\ i \neq j}}^{\kappa} \text{Tr}(M^{l_i})) \rangle \\ &= \frac{N}{t} \left\langle \text{Tr}(M^{l_1} \cdot [M - V'(M)]) \times \text{Tr}(M^{l_2}) \cdots \text{Tr}(M^{l_\kappa}) \right\rangle. \end{aligned} \quad (6.3)$$

These expressions are the SDE for the matrix-valued correlator

$$\langle M^{l_1} \text{Tr}(M^{l_2}) \cdots \text{Tr}(M^{l_\kappa}) \rangle, \quad (6.4)$$

defined for a function $\Phi : M_N(\mathbb{R}) \rightarrow M_N(\mathbb{R})$ by

$$\langle \Phi(M) \rangle = \frac{\int dM \Phi(M) e^{-\frac{N}{t} \text{Tr}(M^2/2 - V(M))}}{\int dM e^{-\frac{N}{t} \text{Tr}(M^2/2 - V(M))}}.$$

(3) The crux of the matter is that Tutte Equations (6.1) hold if and only if the SDE (6.3) for the matrix model (6.2) do. The bridge is the following. For closed maps $\log Z$, the logarithm of the formal integral (6.2), is well-known to yield the generating function of connected closed maps (cf. [BIPZ78]). The formal variables λ_α in both cases coincide: if only maps consisting of, say, triangulations and quadrangulations are to be counted, one sets a cubic and quartic interaction in the matrix model (in eq. (6.2) $\lambda_\alpha = \delta_3^\alpha \lambda_3 + \delta_4^\alpha \lambda_4$). For maps with κ marked faces, if x_1, \dots, x_κ are formal variables and one defines W_κ by

$$W_\kappa(x_1, \dots, x_\kappa) = \left\langle \text{Tr} \frac{1}{x_1 - M} \cdots \text{Tr} \frac{1}{x_\kappa - M} \right\rangle_{\text{connected}}, \quad (6.5)$$

in the sense of Neumann series, then $\mathcal{T}_{l_1 \dots l_\kappa}$ can be recovered by taking residues at $x_i \rightarrow \infty$ as follows:

$$\mathcal{T}_{l_1 \dots l_\kappa} = (-1)^\kappa \text{Res}_{x_1 \rightarrow \infty} \cdots \text{Res}_{x_\kappa \rightarrow \infty} [x_1^{l_1} \cdots x_\kappa^{l_\kappa} W_\kappa(x_1, \dots, x_\kappa)]. \quad (6.6)$$

Tutte equations (6.1) for all genera $g \in \mathbb{Z}_{\geq 0}$ emerge by taking the small- t expansion¹⁵ $\mathcal{T}_{l_1+1, \dots, l_\kappa} = \sum_g (N/t)^{2-2g-\kappa} \mathcal{T}_{l_1+1, \dots, l_\kappa}^{(g)}$.

6.2. Parallel between Tutte equations and disconnected- ∂ SDE of TFT. We contrast now elements appearing in the SDE of Theorem 4.1 with Tutte equations, as well as the derivation of both sets.

The first parallel, depicted in Table 2, concerns the role of the boundaries in each framework.

Moreover, we can regard each Tutte equation as a set of operations in the input, namely the perimeters $l_1 + 1, \dots, l_\kappa$ of the marked faces, and distinguish the two cases $\kappa = 1$ and $\kappa > 1$. The output for the *connected- ∂* Tutte equations (6.1a) is illustrated in Table 3;

¹⁵To see the subtleties between small- t expansion and an $1/N$ -expansion we refer to [Eyn16, Sect. 1.2.4]

PARALLEL BETWEEN TUTTE EQUATIONS AND TFT SDE: ROLE OF BOUNDARIES		
Concept	Variable in Tutte Equations	Variable in TFT
number of boundary components	κ	k
parameters of the boundaries	κ lengths	D -coloured graph \mathcal{D}
boundary decomposition	$(l_1 + 1, l_2, \dots, l_\kappa)$	$\mathcal{D} = \mathcal{R} \sqcup Q_1^{\sqcup \alpha_1} \sqcup \dots \sqcup Q_n^{\sqcup \alpha_n}$ ($k = \alpha_1 + \dots + \alpha_n + 1$)
distinguished boundary component in the derivation of the equations	$l_1 + 1$	\mathcal{R} (cf. Notation of the theorem)

TABLE 2. Boundaries in both frameworks

similarly, for the disconnected- ∂ Tutte equations Table 4 depicts the types of operations on the list of perimeters.

A second similitude is the role played by a distinguished boundary in each case. Tutte equations can be derived [Eyn16, Ch. 1] by distinguishing a single boundary length, say $l_1 + 1$, among the list of perimeters $l_1 + 1, \dots, l_\kappa$, and seeing the possible effects¹⁶ that the removal of a marked edge of this $(l_1 + 1)$ -agon has at the level of the \mathcal{T} -generating functions. In our TFT SDE, the role of that marked face is taken by the boundary component denoted \mathcal{R} . This undergoes the transformations explained in Table 5.

This new operations are an improvement of description given in [PPW17]. Put into the Tutte Equations perspective, the analogue of Table 3 are precisely those operations on a *connected* ∂ -graph of [PPW17]. The contribution of Theorem 4.1 is to give the full set of operations described and interpreted in Table 5.

One can compare term by term Tutte vs. Schwinger-Dyson equations. The terminology refers to Tables 4 and 5. They do not match 1:1, due to the fact that in TFT the number of operations on boundaries increases.

Finally, boundaries —and therefore SDE— of TFT are more complex than their matrix counterparts due to their lack of symmetry. As pointed out in Section 4.2, the choice of vertex \mathbf{s} does matter in the sense that different choices unrelated by non-trivial graph automorphisms of the distinguished boundary \mathcal{R} yield different SDE.

¹⁶Incidentally, this amounts to four possible scenarios that account for each one of the terms in eq. (6.1b):

- I. if the marked edge separates an ordinary face from a marked face: this accounts for the term $\lambda_\alpha \mathcal{T}_{l_1 + \alpha - 1, K}^{(g)}$
- II. if the marked edge separates two marked faces $\rightarrow l_m \mathcal{T}_{l_1 + l_m - 1, K \setminus \{l_m\}}^{(g)}$
- III. if the marked edge bounds twice the same face:
 - III.a and the marked edge is not a bridge $\rightarrow \mathcal{T}_{j, J}^{(g_1)} \times \mathcal{T}_{l_1 - 1 - j, K \setminus J}^{(g_2)}$
 - III.b and the marked edge is a bridge $\rightarrow \mathcal{T}_{j, l_1 - 1 - j, K}^{(g-1)}$.

SINGLE BOUNDARY TUTTE / SCHWINGER DYSON EQUATIONS		
Input	Outputs	Interpretation
$l + 1$	$\rightarrow \{l - 1 + \alpha\}_{\alpha \geq 3}$ if $[x^\alpha]V(x) \neq 0$	increase by α length of boundary
(boundary perimeter)	$\searrow \{j, l - 1 - j\}_{j=0, \dots, l-1}$	splitting in 2 lengths, adding up to $l - 1$

TABLE 3. Operations on the boundaries (marked face) for the connected boundary Tutte for random maps / Schwinger-Dyson equations for the matrix model $Z = \int dM e^{-\frac{N}{t}(\frac{M^2}{2} - V(M))}$ with $V(x) = \sum_{\alpha=3}^d \lambda_\alpha x^\alpha / \alpha$

TUTTE / SCHWINGER-DYSON EQUATIONS FOR MULTIPLE MARKED FACES/BOUNDARIES OF LENGTHS $l_1 + 1, l_2, \dots, l_\kappa$			
Input	Type	Output: perimeter of marked face	Interpretation
$l_1^+, l_2, \dots, l_\kappa$ the boundary perimeters	(i)	$\{l_1^- + \alpha, l_2, \dots, l_\kappa\}_{\alpha \geq 3}$	increase by α the perimeter of marked face whenever $[x^\alpha]V(x) \neq 0$
	(ii)	$L_1 = \{j\} \cup J; L_2 = \{l_1^- - j\} \cup (K \setminus J)$ for $j = 0, \dots, l_1^-, J \subset K$	$1 + \kappa$ marked faces split into a product of two correlators of length lists L_1 and L_2
(Here $K := \{l_2, \dots, l_\kappa\}$ and $l_1^\pm := l_1 \pm 1$)	(iii)	$\{l_1^- + l_\alpha\} \cup K \setminus \{l_\alpha\}$	coalescence of two marked faces \Rightarrow reduction to $\kappa - 1$ marked faces
	(iv)	$\{j, l_1^- - j\} \cup K$ for $j = 0, \dots, l_1^-$	increased number of marked faces to $\kappa + 1$

TABLE 4. Operations on the lengths on the boundaries of disconnected boundary Tutte / Schwinger-Dyson equations

DISCONNECTED- ∂ SCHWINGER DYSON EQUATIONS		
Term in SDE	Type	Transformation of \mathcal{D} and interpretation
$G_{\mathcal{D}}$	0.	None, \mathcal{D} is input
$f_{\mathcal{D},s_c}^{(c)}$	1.	Contraction of three vertices of ${}^c\mathbb{Q}^c$ with $(2 + \#\text{Vertices } \mathcal{D})$ -pt functions, effectively leading to a boundary graph \mathcal{D}
$\Delta\left(\frac{\partial W}{\partial \varsigma_c(\mathcal{D};\beta,\rho)}\right)$	2A.	Contraction of two white vertices of ${}^c\mathbb{Q}^c$ with $\varsigma_c(\mathcal{D};\beta,\rho)$ at the black vertices β and $\rho \Rightarrow$ effectively \mathcal{D} ; accounts for splitting and merging of boundaries. Here ρ is a vertex of the graph \mathcal{R}
$\Delta\left(\frac{\partial W}{\partial \varsigma_c(\mathcal{D};\beta,\rho)}\right)$	2B.	Idem, but ρ is not a vertex of the graph \mathcal{R}
$\Delta G_{\mathcal{D}}$	3.	The tadpole-like term contracting both (say) lower vertices of \mathbb{Q}^c with two vertices of \mathcal{D} corresponding to \mathbf{s} and the white vertex of \mathcal{D} connected to \mathbf{s} by a c -coloured edge
$\Delta H_{\mathcal{B},\mathcal{C}}^{(c,\tau)}$	4.	For $\tau \in \text{Br}^{(2)}(\mathcal{R},\beta,c)$, splitting of \mathcal{R} into two connected components, \mathcal{R}' and \mathcal{R}'' ; splitting of the remaining components \mathcal{B},\mathcal{C} of $\mathcal{D} \stackrel{c}{\sim} (\mathcal{B} \mathcal{C} \mathcal{R})$, as product of correlators, $G_{\mathcal{R}' \mathcal{B}} \times G_{\mathcal{R}'' \mathcal{C}}$, one containing \mathcal{R}' , the other \mathcal{R}''
$I_{\mathcal{B},\mathcal{C}}^{(c)}$	5.	Splitting into product of correlators, one with an insertion of \mathbb{Q}^c , namely $f_{\mathcal{C},s_c}^{(c)}$, and $G_{\mathcal{R} \mathcal{B}}$; altogether the components of $\mathcal{R},\mathcal{C},\mathcal{B}$ must be exactly those of \mathcal{D}

TABLE 5. Tensor model analogue of Table 4. The notation in the column ‘Term in SDE’ refers each row in the Eq. (4.4). In types 2 through 4, ΔL refers to the line of Eq. (4.4) containing a term of the form $[L(\mathbf{X}) - L(\mathbf{X}|_{s_c \rightarrow \xi})]$ that can be easily read off (together with its factors and sums over automorphisms and, as appropriate, over momenta)

INTERPRETATION	TUTTE EQUATIONS	SCHWINGER-DYSON EQUATIONS IN TFT
<i>Interaction vertex insertion in boundary component</i>	<p><i>Type (i)</i></p> <p>This is a connected sum of polygons by a segment $(\alpha, l_1 + 1) \mapsto (\alpha + l_1 - 1)$, since two identified edges disappear.</p>	<p><i>Types 1, 2A and 3</i></p> <p>In type 1, the propagator-contraction of three (out of four) fields is represented. One could as well contract only two vertices of $\overline{\mathfrak{Q}}$, horizontally, and this leads to type 3 terms. A contraction of two white vertices of $\overline{\mathfrak{Q}}$ and the black vertex β (fixed) and another ρ in the same graph \mathcal{R}, leads to type 2A terms.</p>
<i>Coalescence of two boundary components</i>	<p><i>Type (iii)</i></p> <p>Again a connected sum, but now between two different boundary polygons</p>	<p><i>Type 2B</i></p> <p>A contraction of two white vertices of $\overline{\mathfrak{Q}}$ and the black vertex β and a black vertex ρ in a connected component of $\mathcal{D} \setminus \mathcal{R} = \mathcal{Q}$, leads to type 2B terms.</p>
<i>Splitting of a boundary</i>	<p><i>Types (ii) and (iv)</i></p> <p>In type (ii) the splitting implies a product of correlators; in type (iv) it does not. In both the splitting increases $\kappa \mapsto \kappa + 1$</p>	<p><i>Types 4 and 5</i></p> <p>In type 4 the product of correlators occurs. In type 5 one encounters a product of an \mathfrak{f}-function (i.e. an insertion of $\overline{\mathfrak{Q}}$ into a component of \mathcal{Q} and a correlator that contains the distinguished boundary \mathcal{R}</p>

TABLE 6. Term-by-term comparison

7. CONCLUSIONS AND OUTLOOK

We introduced the multi-variable graph calculus—a tool to prove a general formula for the disconnected- ∂ Schwinger-Dyson for the most general quartic-melonic TFT in arbitrary rank (Thm 4.1). In their description, a new set of graph operations on an input graph has been exposed in Table 5. This list is the tensorial equivalent of the matrix model operations on the boundary graphs of matrix models (cf. Table 4).

The well-known dictionary between the theory of enumeration of random maps and matrix models allows to pose two uses our SDE could be useful for. The theories of graph-encoded manifolds [Pez75, LM06, CC15] bring tensor models into prominence as a theory of ‘random higher-dimensional maps’. For instance, gluings of octahedra [BL16] are studied from the tensor model with (non-melonic) interaction vertex



which is the dual graph-representation to an octahedron. The boundary-completeness of the quartic-melonic models [Pér18, Thm 1] further studied here shows a path to the theory of higher-dimensional maps made of gluings of particular triangulations of D -balls, in the rank- D case. The natural candidate for higher-dimensional Tutte equations is the set of relations derived from the $1/N$ -sectors of the present SDE that have the same boundary as well as value of Gurău degree. This is based on the fact (recalled in Sect. 6) that Tutte equations hide in the $N^{2-2g-\kappa}$ -sectors of Migdal’s SDE for matrix models.

The second guess is the existence of a recursion allowing to compute higher-point TFT-correlators from some small number of lower-point ones, analogous to the Topological Recursion (TR) [ACP⁺13, Bor17, Eyn14, EO07, Suh18] that satisfy matrix model correlators. What is not speculative is that, assuming the dictionary of last paragraph, such tensorial TR would require the disconnected- ∂ SDE, since it is also a recursion in the number of boundaries (presumably also in their Gurău’s degree). The blobbed TR for quartic *tensor models* has already been obtained [BD18]—yet it would be interesting to develop the purely tensorial¹⁷ cousin of the TR for *tensor field theory*¹⁸.

The large- N limit of the disconnected- ∂ SDE should be analysed in order to access also their physical significance. At leading order, their *melonic approximation* [OPVW15] is expected to yield closed equations. A significant progress in this direction has been undertaken in [PPTW18], whose techniques could [Pas] be extended to the present disconnected- ∂ SDE in a next project.

¹⁷The Bonzom-Dartois TR bases on an initial Hubbard-Stratonovich transformation. Vertically cutting the pillow vertices, as the rank-3 $\mathfrak{c}\mathfrak{Q}^c$, they “map” quartic melonic tensor models to a suitable intermediate field multi-matrix model, for which the authors develop a TR analogous to the introduced by Borot [Bor15].

¹⁸The difference between tensor models and tensor field theory is here substantial. The former usually focuses on numerical observables $\langle \text{Tr}_{\mathcal{B}}(\varphi, \bar{\varphi}) \rangle \in \mathbb{C}$ and the latter on functions (or distributions) $G_{\mathcal{B}} : \mathcal{V}(\mathcal{B}) \rightarrow \mathbb{C}$. In contrast, the loop equations, Ward Identities [IMM17] and SDE [Gur12] are algebraic in the for tensor models, whereas for tensor field theory ‘loop equations’ [Pér18, PPW17] are integro-differential, as shown also in [PPTW18] explicitly.

Heading towards a quantum gravity perspective, objects appearing in the Functional Renormalization Group [BGKOP18] —or Ward-constrained flows [LO19b]— are expected to be described in terms of graph-generated functionals studied here. Together with the boundary-completeness of the quartic-melonic models [Pér18, Thm 1], this motivates us to study the geometric nature of the flow from a simple quartic model.

On the purely mathematical side, systems of graph-group actions can be extended to Lie groups actions and to calculi in infinitely many graph-variables by using rigorous analytic tools. It would be also interesting to consider the coefficient functions u_g directly in certain algebra of functions. One could dispense with the functions $\{\mathcal{V}(g) \rightarrow \mathbb{C}\}$ by using instead directly (non-commutative) algebras.

Finally, the (symmetric) monoidal structure on the set of boundary graphs emerges in a natural way. This guides us towards the language of Topological Quantum Field Theories (TQFT) [Ati89]. Since these boundary graphs triangulate boundary states, an interesting program would be to obtain discrete TQFT from matrix models and ‘TQFT with observables’ [Oec16] from tensor models, or enframe these in Oeckl’s positive boundary formalism (*op. cit.*), which also facilitates the gluing-boundary procedure that TQFT provides. In the tensor and matrix models case, the gluing of boundaries should be implemented as an operation $\wedge_{\mathcal{A}}$ on two correlation functions sharing a boundary state \mathcal{A} , $G_{\mathcal{A}|\mathcal{B}|\dots|\mathcal{C}'} \wedge_{\mathcal{A}} G_{\mathcal{A}|\mathcal{B}'|\dots|\mathcal{C}'}$, which should be related to $G_{\mathcal{B}|\mathcal{B}'|\dots|\mathcal{C}'}$ due to their geometrical interpretation.

Acknowledgements. The author thanks Raimar Wulkenhaar and Adrian Tanasă for hospitality; and Romain Pascalie for helpful hints and carefully reading the draft (any error is the author’s responsibility). Thanks to the Faculty of Physics, Astronomy and Applied Computer Science, Jagiellonian University (Cracow, Poland), where part of this article was written, for hospitality. The author acknowledges the Short-Term Scientific Mission program of the COST Action MP 1405 for this mobility opportunity.

This research was funded by the Deutsche Forschungsgemeinschaft, SFB 878 (Mathematical Institute of the University of Münster, Germany). Subsequently it was carried out at the Institute of Theoretical Physics, University of Warsaw and has been supported by the TEAM programme of the Foundation for Polish Science co-financed by the European Union under the European Regional Development Fund (POIR.04.04.00-00-5C55/17-00).

APPENDIX A. THE FIRST COEFFICIENTS OF THE Y -TERM

For completeness, we give the first coefficients of the Y -term, keeping in mind the notation simplification (Table 1). The computation of these functions is presented in detail in [Pér18]. As before, the set equality $\{a, b, c\} = \{1, 2, 3\}$ holds.

$$\begin{aligned} \mathfrak{f}_{\mathbf{m};s_a}^{(a)}(\mathbf{x}) &= G_{V_a}^{(4)}(\mathbf{x}, s_a, x_b, x_c) \\ &+ \sum_{c \neq a} \sum_{q_b \in \mathbb{Z}} G_{V_c}^{(4)}(\mathbf{x}; s_a, q_b, x_c) + \sum_{q_b, q_c} G_{\mathbf{m}|\mathbf{m}}^{(4)}(\mathbf{x}; s_a, q_b, q_c), \end{aligned} \tag{A.1a}$$

$$\begin{aligned}
\mathfrak{f}_{V_a; s_a}^{(a)}(\mathbf{x}, \mathbf{y}) &= \frac{1}{3} \left(G_{Q_a}^{(6)}(s_a, x_b, x_c, \mathbf{x}, \mathbf{y}) + \text{cyclic perm. in } (s_a, x_b, x_c), \mathbf{x} \text{ and } \mathbf{y} \right) \\
&\quad + \frac{1}{3} \left(G_{K_{3,3}}^{(6)}(s_a, x_b, y_c; \mathbf{x}, \mathbf{y}) + \text{cyclic perm.} \right) \\
&\quad + \sum_{q_b} G_{F_{b;ac}}^{(6)}(\mathbf{x}; \mathbf{y}; s_a, q_b, y_c) + \sum_{q_c} G_{F_{c;ab}}^{(6)}(\mathbf{x}; \mathbf{y}; s_a, q_c, y_b) \\
&\quad + \frac{1}{2} \sum_{q_b, q_c} G_{m|V_a}^{(6)}(s_a, q_b, q_c; \mathbf{x}, \mathbf{y}),
\end{aligned} \tag{A.1b}$$

$$\begin{aligned}
\mathfrak{f}_{V_b; s_a}^{(a)}(\mathbf{x}, \mathbf{y}) &= \frac{1}{3} \left(\sum_{q_b} G_{Q_b}^{(6)}(s_a, q_b, y_c; \mathbf{x}, \mathbf{y}) + \text{cyclic perm.} \right) + G_{F_{c;ab}}^{(6)}(s_a, y_b, x_c; \mathbf{x}; \mathbf{y}) \\
&\quad + G_{F_{c;ab}}^{(6)}(\mathbf{x}; s_a, x_b, x_c; \mathbf{y}) + \sum_{q_b} G_{F_{a;bc}}^{(6)}(\mathbf{x}; \mathbf{y}; s_a, q_b, y_c) \\
&\quad - \frac{1}{2} \sum_{q_b, q_c} G_{m|V_b}^{(6)}(s_a, q_b, q_c; \mathbf{x}; \mathbf{y}),
\end{aligned} \tag{A.1c}$$

$$\begin{aligned}
\mathfrak{f}_{m|m; s_a}^{(a)}(\mathbf{x}, \mathbf{y}) &= \left(\sum_{q_b, q_c} G_{m|m|m}^{(6)}(s_a, q_b, q_c, \mathbf{x}, \mathbf{y}) + \text{cyclic perm.} \right) \\
&\quad + G_{m|V_a}^{(6)}(\mathbf{x}, s_a, y_b, y_c, \mathbf{y}) + \sum_{q_c} G_{m|V_b}^{(6)}(\mathbf{x}, s_a, y_b, q_c, \mathbf{y}) \\
&\quad + \sum_{q_b} G_{m|V_c}^{(6)}(\mathbf{x}, s_a, q_b, y_c, \mathbf{y}) + \sum_{q_c} G_{m|V_b}^{(6)}(\mathbf{x}, \mathbf{y}, s_a, y_b, q_c) \\
&\quad + \sum_{q_b} G_{m|V_c}^{(6)}(\mathbf{x}, \mathbf{y}, s_a, q_b, y_c) + G_{m|V_a}^{(6)}(\mathbf{x}, \mathbf{y}, s_a, y_b, y_c) \\
&\quad + G_{F_{a;bc}}^{(6)}(\mathbf{x}, s_a, x_b, y_c, \mathbf{y}).
\end{aligned} \tag{A.1d}$$

REFERENCES

- [ADJ91] Jan Ambjørn, Bergfinnur Durhuus, and Thordur Jonsson. Three-dimensional simplicial quantum gravity and generalized matrix models. *Mod. Phys. Lett.*, A6:1133–1146, 1991.
- [AGJL13] Jan Ambjørn, Andrzej Görlich, Jerzy Jurkiewicz, and Renate Loll. Causal dynamical triangulations and the search for a theory of quantum gravity. *Int. J. Mod. Phys.*, D22:1330019, 2013.
- [ACP⁺13] Jorgen E. Andersen, Leonid O. Chekhov, R. C. Penner, Christian M. Reidys, and Piotr Sulkowski. Topological recursion for chord diagrams, RNA complexes, and cells in moduli spaces. *Nucl. Phys.*, B866:414–443, 2013.
- [Ati89] Michael Atiyah. Topological quantum field theories. *Inst. Hautes Etudes Sci. Publ. Math.*, 68:175–186, 1989.
- [BD18] Valentin Bonzom and Stephane Dartois. Blobbed topological recursion for the quartic melonic tensor model. *J. Phys.*, A51(32):325201, 2018.
- [BG12] Joseph Ben Geloun. Asymptotic Freedom of Rank 4 Tensor Group Field Theory. In *29th International Colloquium on Group-Theoretical Methods in Physics (GROUP 29) Tianjin, China, August 20-26, 2012*, 2012. arXiv:1210.5490.

- [BG18] Dario Benedetti and Razvan Gurau. 2PI effective action for the SYK model and tensor field theories. *JHEP*, 05:156, 2018.
- [BGKOP18] Joseph Ben Geloun, Tim A. Koslowski, Daniele Oriti, and Antonio D. Pereira. Functional Renormalization Group analysis of rank 3 tensorial group field theory: The full quartic invariant truncation. *Phys. Rev.*, D97(12):126018, 2018.
- [BGR13] Joseph Ben Geloun and Vincent Rivasseau. A Renormalizable 4-Dimensional Tensor Field Theory. *Commun. Math. Phys.*, 318:69–109, 2013. arXiv:1111.4997.
- [BGR11] Valentin Bonzom, Răzvan Gurău, Aldo Riello, and Vincent Rivasseau. Critical behavior of colored tensor models in the large N limit. *Nucl. Phys.*, B853:174–195, 2011. arXiv:1105.3122.
- [BGS13] Joseph Ben Geloun and Dine Ousmane Samary. 3D Tensor Field Theory: Renormalization and One-loop β -functions. *Annales Henri Poincaré*, 14:1599–1642, 2013. arXiv:1201.0176.
- [BIPZ78] Édouard Brezin, Claude Itzykson, Giorgio Parisi, and Jean-B. Zuber. Planar Diagrams. *Commun. Math. Phys.*, 59:35, 1978.
- [BL16] Valentin Bonzom and Luca Lionni. Counting gluings of octahedra. *Elec. J. Combin.*, 24, 2016. arXiv:1608.00347.
- [Bor15] Gaëtan Borot. Blobbed topological recursion. *Theor. Math. Phys.*, 185(3):1729–1740, 2015. [Teor. Mat. Fiz.185,no.3,423(2015)].
- [Bor17] Gaëtan Borot. Lecture notes on topological recursion and geometry. 2017. arXiv:1705.09986.
- [CC97] Ali H. Chamseddine and Alain Connes. The Spectral action principle. *Commun. Math. Phys.*, 186:731–750, 1997.
- [COR14] Sylvain Carrozza, Daniele Oriti, and Vincent Rivasseau. Renormalization of Tensorial Group Field Theories: Abelian U(1) Models in Four Dimensions. *Commun. Math. Phys.*, 327:603–641, 2014.
- [CP18] Sylvain Carrozza and Victor Pozsgay. SYK-like tensor quantum mechanics with Sp(N) symmetry. arXiv:1809.07753 2018.
- [CT16] Sylvain Carrozza and Adrian Tanasă. $O(N)$ Random Tensor Models. *Lett. Math. Phys.*, 106(11):1531–1559, 2016.
- [CC15] Rita Cassali and Paola Cristofori. *Cataloguing PL 4-manifolds by gem-complexity*, *Electron. J. Combin.*, **22**(4) #P4.25, (2015) 1–25.
- [DGMR07] Margherita Disertori, Răzvan Gurău, Jacques Magnen, and Vincent Rivasseau. Vanishing of Beta Function of Non Commutative Φ_4^4 Theory to all orders. *Phys. Lett.*, B649:95–102, 2007. arXiv:hep-th/0612251.
- [Eic19] Astrid Eichhorn. An asymptotically safe guide to quantum gravity and matter. *Front. Astron. Space Sci.*, 5:47, 2019.
- [EKLP18] Astrid Eichhorn, Tim Koslowski, Johannes Lumma, and Antonio D. Pereira. Towards background independent quantum gravity with tensor models. 2018. arXiv:1811.00814.
- [EKP19] Astrid Eichhorn, Tim Koslowski, and Antonio D. Pereira. Status of background-independent coarse-graining in tensor models for quantum gravity. *Universe*, 5(2):53, 2019.
- [EO07] Bertrand Eynard and Nicolas Orantin. Invariants of algebraic curves and topological expansion. *Commun. Num. Theor. Phys.*, 1:347–452, 2007.
- [Eyn14] Bertrand Eynard. A short overview of the “Topological recursion”. 2014. arXiv:1412.3286.
- [Eyn16] Bertrand Eynard. *Counting Surfaces*, volume 70 of *Progress in Mathematical Physics*. Springer, 2016.
- [GR12] Răzvan Gurău and James P. Ryan. Colored Tensor Models - a review. *SIGMA*, 8:020, 2012. arXiv:1109.4812.

- [Gur12] Răzvan Gurău. The Schwinger Dyson equations and the algebra of constraints of random tensor models at all orders. *Nucl. Phys.*, B865:133–147, 2012. arXiv:1203.4965 [hep-th].
- [GW14] Harald Grosse and Raimar Wulkenhaar. Self-Dual Noncommutative φ^4 -Theory in Four Dimensions is a Non-Perturbatively Solvable and Non-Trivial Quantum Field Theory. *Commun. Math. Phys.*, 329:1069–1130, 2014. arXiv:1205.0465.
- [IMM17] Hiroshi Itoyama, Andrei Mironov, and Alexei Morozov. Ward identities and combinatorics of rainbow tensor models. *JHEP*, 06:115, 2017.
- [KT16] Thomas Krajewski and Reiko Toriumi. Exact Renormalisation Group Equations and Loop Equations for Tensor Models. *SIGMA*, 12:068, 2016. arXiv:1603.00172.
- [Lan02] Serge Lang. *Algebra*. Springer, Graduate Texts in Mathematics, 2002.
- [LM06] Sostenes Lins and Michele Mulazzani, Blobs and flips on gems, *J. Knot Theory Ramif.*, **15** (2006) 1001–1035.
- [LO19] Vincent Lahoche and Dine Ousmane Samary. Progress in the solving nonperturbative renormalization group for tensorial group field theory. *Universe*, 5:86, 2019.
- [LO19b] Vincent Lahoche and Dine Ousmane-Samary. Ward-constrained melonic renormalization group flow. arXiv:1904.05655, 2019.
- [Mar18] Matilde Marcolli. *Noncommutative Cosmology*. WSP, 2018.
- [MT18] Jakub Mielczarek and Tomasz Trześniewski. Towards the map of quantum gravity. *Gen. Rel. Grav.*, 50(6):68, 2018.
- [Mig83] Alexander A. Migdal. Loop equations and $1/N$ expansion. *Phys. Rep.*, 102:199–290, 1983.
- [Oec16] Robert Oeckl. A local and operational framework for the foundations of physics. 2016. arXiv:1610.09052.
- [OPVW15] Dine Ousmane Samary, Carlos I. Pérez-Sánchez, Fabien Vignes-Tourneret, and Raimar Wulkenhaar. Correlation functions of a just renormalizable tensorial group field theory: the melonic approximation. *Class. Quant. Grav.*, 32(17):175012, 2015. arXiv:1411.7213.
- [Ous14] Dine Ousmane Samary. Closed equations of the two-point functions for tensorial group field theory. *Class. Quant. Grav.*, 31:185005, 2014. arXiv:1401.2096.
- [Pas19] Romain Pascalie. A Solvable Tensor Field Theory. 2019. arXiv:1903.02907.
- [Pas] Romain Pascalie (*et. al?*). Work in progress. 201?.
- [Pér17] Carlos I. Pérez-Sánchez. Surgery in colored tensor models. *J. Geom. Phys.*, 120:262–289, 2017.
- [Pér18] Carlos I. Pérez-Sánchez. The full Ward-Takahashi Identity for colored tensor models. *Commun. Math. Phys.*, 358(2):589–632, 2018. arXiv:1608.08134.
- [Pez75] Mario Pezzana. Sulla struttura topologica delle varietà compatte. *Ati Sem. Mat. Fis. Univ. Modena*, 23(1):269–277, 1975.
- [PPTW18] Romain Pascalie, Carlos I Pérez-Sánchez, Adrian Tanasă, and Raimar Wulkenhaar. The large- N limit of the Schwinger-Dyson equations of Tensor Field Theory. 2018. arXiv:1810.09867.
- [PPW17] Romain Pascalie, Carlos I. Pérez-Sánchez, and Raimar Wulkenhaar. Correlation functions of $U(N)$ -tensor models and their Schwinger-Dyson equations. 2017. arXiv:1706.07358.
- [Riv16] Vincent Rivasseau. Random Tensors and Quantum Gravity. *SIGMA*, 12:069, 2016. arXiv:1603.07278.
- [RVT] Vincent Rivasseau and Fabien Vignes-Tourneret. Constructive tensor field theory: The t_4^4 model. *Commun. Math. Phys.*, 366(2):567–646.
- [Suł18] Piotr Sułkowski. Topological recursion . *Snapshots of modern mathematics from Oberwolfach*, (2), 2018.

- [OV14] Dine Ousmane-Samary and Fabien Vignes-Tourneret. Just Renormalizable TGFT's on $U(1)^d$ with Gauge Invariance. *Commun. Math. Phys.*, 329:545–578, 2014. arXiv:1211.2618.
- [Sze16] Fernando Szechtman. Groups having a faithful irreducible representation. *Journal of Algebra*, 454:292 – 307, 2016.
- [Tan16] Adrian Tanasa. The Multi-Orientable Random Tensor Model, a Review. *SIGMA*, 12:056, 2016.
- [Tut62] William T. Tutte. A census of planar triangulations. *Can. J. Math.*, 14:21–38, 1962.
- [Tut63] William T. Tutte. A census of planar maps. *Can. J. Math.*, 15:249–271, 1963.