

Improved bounds for box dimensions of potential singular points to the Navier–Stokes equations

Yanqing Wang* & Minsuk Yang†

Abstract

In this paper, we study the potential singular points of interior and boundary suitable weak solutions to the 3D Navier–Stokes equations. It is shown that upper box dimension of interior singular points and boundary singular points are bounded by $7/6$ and $10/9$, respectively. Both proofs rely on recent progress of ε -regularity criteria at one scale.

MSC(2000): 76D03, 76D05, 35B33, 35Q35

Keywords: Navier–Stokes equations; suitable weak solutions; singular points; regularity criteria

1 Introduction

We consider the following three-dimensional incompressible Navier–Stokes equations

$$\begin{cases} u_t - \Delta u + u \cdot \nabla u + \nabla \Pi = 0, \\ \operatorname{div} u = 0, \\ u|_{t=0} = u_0, \end{cases} \quad (1.1)$$

where u is the velocity field, Π is the scalar pressure, and the initial velocity u_0 satisfies the condition $\operatorname{div} u_0 = 0$. We investigate the regularity problem of the Navier–Stokes equations. Our main objective of this paper is to further lower the box dimension of potential singular points of suitable weak solutions to the Navier–Stokes equations.

A point $z = (x, t)$ is said to be regular if u is Hölder continuous at a neighborhood of z . Otherwise, it is called singular. In a celebrated work [1], Caffarelli, Kohn and Nirenberg obtained two ε -regularity criteria to suitable weak solutions of (1.1): $z = 0$ is regular point provided that one of the following conditions holds for an absolute positive constant ε ,

$$\|u\|_{L^3(Q(1))} + \|u\Pi\|_{L^1(Q(1))} + \|\Pi\|_{L^{1,5/4}(Q(1))} < \varepsilon. \quad (1.2)$$

and

$$\limsup_{r \rightarrow 0} \|\nabla u\|_{L^2(Q(r))} < \varepsilon. \quad (1.3)$$

*Department of Mathematics and Information Science, Zhengzhou University of Light Industry, Zhengzhou, Henan 450002, P. R. China. Email: wangyanqing20056@gmail.com

†Correspondence author, Department of Mathematics, Yonsei University, 50 Yonsei-ro Seodaemun-gu, Seoul, Republic of Korea. Email: m.yang@yonsei.ac.kr

Lin [13], Ladyzhenskaya and Seregin [12] gave an alternative condition

$$\|u\|_{L^3(Q(1))} + \|\Pi\|_{L^{3/2}(Q(1))} < \varepsilon. \quad (1.4)$$

instead of (1.2). Recently, Guevara and Phuc [6] improved (1.2) and (1.4) to

$$\|u\|_{L^{2p,2q}(Q(1))} + \|\Pi\|_{L^{p,q}(Q(1))} < \varepsilon, \quad (1.5)$$

for some p, q satisfying $3/p + 2/q = 7/2$ and $1 \leq q \leq 2$. Subsequently, authors in [8] found that (1.5) can be replaced by

$$\|u\|_{L^{p,q}(Q(1))} + \|\Pi\|_{L^1(Q(1))} < \varepsilon \quad (1.6)$$

for some p, q satisfying $1 \leq 2/p + 3/q < 2$ and $1 \leq p, q \leq \infty$. Other interior regularity criteria similar to (1.2) can be found in [3, 17, 19]. Since the gradient of the pressure appears in (1.1), one can replace Π as $\Pi - \bar{\Pi}_{B(1)}$ by subtracting an average in (1.2), (1.4), (1.5), and (1.6). Let \mathcal{S}_i denote the possible interior singular points of suitable weak solutions to the 3D Navier–Stokes equations. One can use the condition (1.2) and (1.3) to obtain that

$$\dim_H(\mathcal{S}_i) \leq 1 \quad \text{and} \quad \dim_B(\mathcal{S}_i) \leq 5/3,$$

where $\dim_H(S)$ and $\dim_B(S)$ denote the Hausdorff dimension and box dimension of a set S , respectively. For the background of fractal dimension, we refer the reader to [4]. The relationship between Hausdorff dimension and box dimension is that the former is less than the latter.

In the past decade, starting from Kukavica's work [10], several authors try to lower the box dimension of potential interior singular points for suitable weak solutions to the 3D Navier–Stokes equations to 1 (Kukavica [10] $135/82 \approx 1.65$); Kukavica and Pei [11] $45/29 (\approx 1.55)$; Koh and Yang [9] $95/63 (1.51)$; Wang and Wu $(360/277 \approx 1.30)$; Ren, Wang and Wu [14] $(975/758 \approx 1.29)$; He, Wang and Zhou [8] $(2400/1903 \approx 1.261)$).

In this paper, we also consider the potential boundary singular points of boundary suitable weak solutions to the Navier–Stokes equations. In particular, the Navier–Stokes equations with no-slip conditions are given by

$$\begin{cases} u_t - \Delta u + u \cdot \nabla u + \nabla \Pi = 0, \mathbb{R}_+^3 \times (0, T) \\ \operatorname{div} u = 0, \quad u|_{\partial \mathbb{R}_+^3} = 0 \\ u|_{t=0} = u_0. \end{cases} \quad (1.7)$$

For the boundary regularity problem of the Navier–Stokes equations (1.7), Seregin [16] obtained the following two ε -regularity conditions

$$\limsup_{\mu \rightarrow 0} \|\nabla u\|_{L^2(Q^+(\mu))} < \varepsilon, \quad (1.8)$$

$$\|u\|_{L^3(Q^+(1))} + \|\Pi\|_{L^{3/2}(Q^+(1))} < \varepsilon. \quad (1.9)$$

After that, Gustafson, Kang and Tsai [7] obtained the ε -regularity condition

$$\|u\|_{L^3(Q^+(1))} + \|\Pi\|_{L^{\lambda, \kappa^*}(Q^+(1))} < \varepsilon, \quad (1.10)$$

where λ and κ^* are defined in (3.3). Let \mathcal{S}_b denote the possible boundary singular points of boundary suitable weak solutions to the 3D Navier–Stokes equations. Using these ε -regularity

conditions, Choe and Yang [2] recently obtained a result about upper box dimension $\dim_B(\mathcal{S}_b)$. In this paper we improve all the previous bounds for the interior and boundary singular points.

The rest of this paper is organized as follows. In Section 2, we give the precise statement of our main results. In Section 3, we introduce relevant notation and definitions of suitable weak solutions to (1.1) and (1.7), respectively. In Section 4, we present a few auxiliary lemmas which are interpolation inequalities and decay estimates for the pressure. In Section 5, we prove Proposition 2.2 and deduce Theorem 2.1, which is an estimation of box dimension of \mathcal{S}_i in (1.1). In Section 6, we prove Theorem 2.6 and Proposition 2.4 and deduce Theorem 2.3, which is an estimation of box dimension of \mathcal{S}_b in (1.7).

2 Main results

In this section we state our main results of this paper.

Theorem 2.1. The upper box dimension of \mathcal{S}_i in (1.1) is at most $7/6$.

Remark 2.1. The bound $7/6$ is better than the previous results obtained in [8–11, 14, 15, 18]. It is a direct consequence of the following regularity criterion.

Proposition 2.2. Suppose that (u, Π) is a suitable weak solution to (1.1). Then for each $\gamma < 1/2$ there exist positive numbers ε_1 and $r_0 < 1$ such that $z = (x, t)$ is a regular point if for some $r < r_0$

$$\int_{Q(z,r)} |\nabla u|^2 + |u|^{10/3} + |\Pi - \bar{\Pi}_{B(x,r)}|^{5/3} + |\nabla \Pi|^{5/4} dx ds < r^{5/3-\gamma} \varepsilon_1 \quad (2.1)$$

where $Q(z, r)$ denotes a parabolic cylinder (see the next section for notations and definitions).

The proof of Proposition 2.2 is different from recent arguments in [8, 9, 14, 18]. The key point is to apply the following ε -regularity criteria, for any $\delta > 0$,

$$\|u\|_{L^{2p}(Q(1))} + \|\Pi\|_{L^p(Q(1))} < \varepsilon \quad \text{and} \quad p > \frac{5}{2} + \delta,$$

to establish an iteration scheme. Our starting point is the following ε -regularity criterion

$$\|\nabla u\|_{L^2(Q(1))} + \|u\|_{L^2(Q(1))} + \|\nabla \Pi\|_{L^{1,3/2}(Q(1))} < \varepsilon, \quad (2.2)$$

which is derived from

$$\|u\|_{L^{2,6}(Q(1))} + \|\nabla \Pi\|_{L^{1,3/2}(Q(1))} < \varepsilon. \quad (2.3)$$

It is worth noting that we get (2.3) by (1.6) and the Poincaé–Sobolev inequality.

We briefly illustrate our strategy in the proof of Proposition 2.2, see Section 5 for its detailed proof. We bound

$$(\theta\rho)^{-3/2} \|u\|_{L^2(Q(\theta\rho))}$$

via interpolation inequality (4.1) and hypothesis (2.1). And then making use of the divergence free condition, we establish a pressure decay estimate (4.3) in terms of $\nabla \Pi$. Using the same scaled quantities, we get smallness of $(\theta\rho)^{-3/2} \|u\|_{L^2(Q(\theta\rho))}$ as well as

$$(\theta\rho)^{-1} (\|\nabla \Pi\|_{L^{5/4}(Q(\theta\rho))} + \|\nabla u\|_{L^2(Q(\theta\rho))}).$$

This together with (2.2) completes the proof of Proposition 2.2.

Before we turn our attention to the boundary case, we give one more remark.

Remark 2.2. As was observed in [18, Remark 1.4, p1762], it seems that it is useful to use the quantity $\|\nabla u\|_{L^2}$ instead of $\|u\|_{L^\infty,2}$ in estimating box dimension of the singular set. The advantage of $\|u\|_{L^2(Q(1))}$ and $\|\nabla \Pi\|_{L^{1,3/2}(Q(1))}$ in (2.2) is the absence of $\|u\|_{L^\infty,2}$ in first part of estimate (4.1) and (4.3). This helps us to make full use of $\|\nabla u\|_{L^2}$ in our proof. One can show that $\dim_B(\mathcal{S}_i) \leq 37/30$ via a combination of the proof described above and the ε -regularity criterion

$$\|u\|_{L^{5/2+\delta}(Q(1))} + \|\nabla \Pi\|_{L^{5/4}(Q(1))} < \varepsilon. \quad (2.4)$$

For the boundary case (1.7) we have the following theorem.

Theorem 2.3. The upper box dimension of \mathcal{S}_b in (1.7) is at most 10/9.

Remark 2.3. The bound 10/9 is better than the previous result obtained in [2]. It is a direct consequence of the following regularity criterion.

Proposition 2.4. Suppose that (u, Π) is a suitable weak solution to (1.7). Then there exist positive numbers ε_2 and $r_0 < 1$ such that z is a regular point if for some $0 < r < r_0$

$$\int_{Q^+(x,t,r)} |\nabla u|^2 + |u|^{10/3} + |\Pi - \bar{\Pi}_{B^+(x,r)}|^{5/3} + |\nabla \Pi|^{5/4} dx ds < r^{10/9} \varepsilon_2. \quad (2.5)$$

Compared with the proof of Proposition 2.2, the proof of Proposition 2.4 includes a new ingredient of an application of new ε -regularity criteria below.

Theorem 2.5. Suppose that (u, Π) is a suitable weak solution to the 3D Navier–Stokes equations (1.7) in $Q^+(1)$. Then there exists an absolute positive constant ε such that $z = 0$ is a regular point if

$$\|\nabla u\|_{L^{p,q}(Q^+(1))} + \|\Pi - \bar{\Pi}_{B_{x,r}^+}(s)\|_{L^{\lambda,\kappa^*}(Q^+(1))} < \varepsilon \quad (2.6)$$

for some p, q satisfying $2 \leq 2/q + 3/p < 3$ and $1 \leq p, q \leq \infty$.

Remark 2.4. A special case is

$$\|\nabla u\|_{L^2(Q^+(1))} + \|\nabla \Pi\|_{L^{5/4,5/4}(Q^+(1))} < \varepsilon,$$

which can be used to show that $\dim_B(\mathcal{S}_b) \leq 5/4$.

The Poincaré–Sobolev inequality guarantees Theorem 2.5 follows from the next theorem, which is of independent interest.

Theorem 2.6. Suppose that (u, Π) is a suitable weak solution to the 3D Navier–Stokes equations (1.7) in $Q^+(1)$. Then there exists an absolute positive constant ε such that $z = 0$ is a regular point if

$$\|u\|_{L^{p,q}(Q^+(1))} + \|\Pi - \bar{\Pi}_{B_{x,r}^+}(s)\|_{L^{\lambda,\kappa^*}(Q^+(1))} < \varepsilon \quad (2.7)$$

for some p, q satisfying $1 \leq 2/q + 3/p < 2$ and $1 \leq p, q \leq \infty$.

Remark 2.5. Theorem 2.6 is an improvement of (1.9) and (1.10). The proof is in part inspired by recent results (1.5) and (1.6).

3 Notations and definitions

We denote by $L^q(-T, 0; X)$, $1 \leq q \leq \infty$, the set of measurable functions on the interval $(-T, 0)$ with values in X and $\|f(t, \cdot)\|_X \in L^q(-T, 0)$. We denote by $z = (x, t) \in \mathbb{R}^3 \times (-T, 0)$ a space-time point and denote balls and cylinders by

$$\begin{aligned} B(x, r) &= \{y \in \mathbb{R}^3 : |x - y| \leq r\}, \\ Q(z, r) &= B(x, r) \times (t - r^2, t). \end{aligned}$$

We recall the definition of suitable weak solutions to (1.1).

Definition 3.1. A pair (u, Π) is called a suitable weak solution to the Navier–Stokes equations (1.1) provided the following conditions are satisfied:

- (1) $u \in L^\infty(-T, 0; L^2(\mathbb{R}^3)) \cap L^2(-T, 0; \dot{H}^1(\mathbb{R}^3))$, $\Pi \in L^{3/2}(-T, 0; L^{3/2}(\mathbb{R}^3))$.
- (2) (u, Π) solves (1.1) in $\mathbb{R}^3 \times (-T, 0)$ in the sense of distributions.
- (3) (u, Π) satisfies the following inequality, for a.e. $t \in (-T, 0)$,

$$\begin{aligned} & \int_{\mathbb{R}^3} |u(x, t)|^2 \phi(x, t) dx + 2 \int_{-T}^t \int_{\mathbb{R}^3} |\nabla u|^2 \phi dx ds \\ & \leq \int_{-T}^t \int_{\mathbb{R}^3} |u|^2 (\partial_s \phi + \Delta \phi) dx ds + \int_{-T}^t \int_{\mathbb{R}^3} u \cdot \nabla \phi (|u|^2 + 2\Pi) dx ds, \end{aligned} \tag{3.1}$$

where non-negative function $\phi(x, s) \in C_0^\infty(\mathbb{R}^3 \times (-T, 0))$.

In the light of the natural scaling property of the Navier–Stokes equations, we introduce the following scaling invariant quantities;

$$\begin{aligned} A(r) &= \sup_{t-r^2 \leq s < t} r^{-1} \int_{B(x, r)} |u(y, s)|^2 dy, \\ E(r) &= r^{-1} \int_{Q(z, r)} |\nabla u(y, s)|^2 dy ds, \\ E_p(r) &= r^{p-5} \int_{Q(z, r)} |u(y, s)|^p dy ds, \\ P_{5/4}(\nabla \Pi, r) &= r^{-1} \left(\int_{Q(z, r)} |\nabla \Pi|^{5/4} dy ds \right)^{4/5}, \\ P_{1,5/4}(\nabla \Pi, r) &= r^{-1} \int_{t-r^2}^t \left(\int_{B(x, r)} |\nabla \Pi|^{5/4} dy \right)^{4/5} ds. \end{aligned} \tag{3.2}$$

For any boundary point $z = (x, t) = (x_1, x_2, 0, t) \in \partial \mathbb{R}_+^3 \times (0, T)$, we denote half balls and half cylinders by

$$\begin{aligned} B^+(x, r) &= \{y \in B(x, r) : y_3 > 0\}, \\ Q^+(z, r) &= \{(y, t) \in Q(z, r) : y_3 > 0\}. \end{aligned}$$

We recall the definition of boundary suitable weak solutions to (1.7).

Definition 3.2. Let $\Omega = \mathbb{R}_+^3$ and $Q_T = \mathbb{R}_+^3 \times [-T, 0)$. A pair of (u, Π) is a suitable weak solution to (1.7) if the following conditions are satisfied:

(1) The functions $u : Q_T \rightarrow \mathbb{R}^3$ and $\Pi : Q_T \rightarrow \mathbb{R}$ satisfy

$$\begin{aligned} u &\in L^\infty(I; L^2(\Omega)) \cap L^2(I; W^{1,2}(\Omega)), \quad \Pi \in L^\lambda(I; L^{\kappa^*}(\Omega)), \\ \nabla^2 u &\in L^\lambda(I; L^\kappa(\Omega)), \quad \nabla \Pi \in L^\lambda(I; L^\kappa(\Omega)), \end{aligned}$$

where λ , κ and κ^* are fixed numbers satisfying

$$1 < \lambda < 2, \quad \frac{2}{\lambda} + \frac{3}{\kappa} = 4, \quad \frac{1}{\kappa^*} = \frac{1}{\kappa} - \frac{1}{3}. \quad (3.3)$$

(2) (u, Π) solves the Navier–Stokes equations in Q_T in the sense of distributions and u satisfy the boundary conditions in the sense of traces.

(3) u and Π satisfy the local energy inequality

$$\begin{aligned} &\int_{B^+(x,r)} |u(x,t)|^2 \phi(x,t) dx + 2 \int_{t_0}^t \int_{B^+(x,r)} |\nabla u(x,s)|^2 \phi(x,s) dx ds \\ &\leq \int_{t_0}^t \int_{B^+(x,r)} |u|^2 (\partial_s \phi + \Delta \phi) dx ds + \int_{t_0}^t \int_{B^+(x,r)} (|u|^2 + 2\Pi) u \cdot \nabla \phi dx ds \end{aligned}$$

for all $t \in I = (0, T)$ and for all nonnegative function $\phi \in C_0^\infty(\mathbb{R}^3 \times \mathbb{R})$.

We shall use the same scaling invariant quantities (3.2) for the boundary case replacing $B(x, r)$ and $Q(z, r)$ by $B^+(x, r)$ and $Q^+(z, r)$ in (3.2). Readers can tell them in the context. In addition, we need the following scaling invariant quantities involving the pressure

$$\tilde{D}(r) = r^{-1} \left(\int_{t-r^2}^t \left(\int_{B^+(x,r)} |\Pi(y,s) - \bar{\Pi}_{B^+(x,r)}(s)|^{\kappa^*} dy \right)^{\frac{\lambda}{\kappa^*}} ds \right)^{\frac{1}{\lambda}}, \quad (3.4)$$

$$\tilde{D}_1(r) = r^{-1} \left(\int_{t-r^2}^t \left(\int_{B^+(x,r)} |\nabla \Pi(y,s)|^\kappa dy \right)^{\frac{\lambda}{\kappa}} ds \right)^{\frac{1}{\lambda}}, \quad (3.5)$$

where we have used the notation \bar{f}_E which is the average of f over the set E .

For κ^* and λ in (3.3), we denote their Hölder conjugate i and j through the relations

$$\frac{1}{i} = 1 - \frac{1}{\lambda}, \quad \frac{1}{j} = 1 - \frac{1}{\kappa^*}.$$

Hence, it follows from (3.3) that

$$\frac{2}{i} + \frac{3}{j} = 2, \quad 2 < i < \infty. \quad (3.6)$$

We end this section by giving a few more notations. When the center of a ball or a cylinder is located at the origin, we put them simply as

$$\begin{aligned} B(r) &= B(0, r) & \text{and} & & Q(r) &= Q(0, r) \\ B^+(r) &= B^+(0, r) & \text{and} & & Q^+(r) &= Q^+(0, r). \end{aligned}$$

For simplicity, we also write $\bar{f}_r = \bar{f}_{B(r)}$ and

$$\begin{aligned} \|f\|_{L^{p,q}(Q(r))} &= \|f\|_{L^p(-r^2, 0; L^q(B(r)))}, \\ \|f\|_{L^p(Q(r))} &= \|f\|_{L^{p,p}(Q(r))}. \end{aligned}$$

We shall use the summation convention on repeated indices. We shall use the notation $A \lesssim B$ if there is a generic positive constant C such that $|A| \leq C|B|$. The generic positive constants C may be different from line to line unless otherwise stated.

4 Auxiliary lemmas

In this section, we present interpolation inequalities and decay estimates for the pressure.

Lemma 4.1 ([8, 14]). *For $0 < r \leq \rho/2$ and $2 \leq p \leq 10/3$,*

$$E_p(r) \lesssim \left(\frac{\rho}{r}\right)^{p/2} A^{(p-2)/2}(\rho) E(\rho) + \left(\frac{r}{\rho}\right)^p A^{p/2}(\rho) \quad (4.1)$$

for the interior case and

$$E_p(r) \lesssim \left(\frac{\rho}{r}\right)^{p/2} A^{(p-2)/2}(\rho) E(\rho) \quad (4.2)$$

for the boundary case, where the implied positive constants does not depend on r and ρ .

We derive decay estimates in terms of $\nabla\Pi$. See [11] for a different version.

Lemma 4.2. *For $0 < r \leq \rho/8$,*

$$P_{1,3/2}(\nabla\Pi, r) \lesssim \left(\frac{\rho}{r}\right) E(\rho) + \left(\frac{r}{\rho}\right) P_{5/4}(\nabla\Pi, \rho) \quad (4.3)$$

for the interior case, where the implied positive constant does not depend on r and ρ .

Proof. Without loss of generality, we assume that $z = 0$. We consider the usual cut-off function $\phi \in C_0^\infty(B(\rho/2))$ such that $\phi \equiv 1$ on $B(3\rho/8)$ with $0 \leq \phi \leq 1$ and

$$|\nabla\phi|^2 + |\nabla^2\phi| \leq C\rho^{-2}.$$

Due to the incompressible condition, we infer that

$$\partial_i \partial_i (\partial_k \Pi \phi) = -\phi \partial_i \partial_j [\partial_k U_{i,j}] + 2\partial_i \phi \partial_i \partial_k \Pi + \partial_k \Pi \partial_i \partial_i \phi$$

where $U_{i,j} = (u_j - \bar{u}_{j,\rho/2})(u_i - \bar{u}_{i,\rho/2})$. Let K denote the normalized fundamental solution of Laplace equation. Then for $x \in B(3\rho/8)$

$$\begin{aligned}
\partial_k \Pi(x) &= K * \{-\phi \partial_i \partial_j [\partial_k U_{i,j}] + 2\partial_i \phi \partial_i \partial_k \Pi + \partial_k \Pi \partial_i \partial_i \phi\} \\
&= -\partial_i \partial_j K * (\phi [\partial_k U_{i,j}]) \\
&\quad + 2\partial_i K * (\partial_j \phi [\partial_k U_{i,j}]) - K * (\partial_i \partial_j \phi [\partial_k U_{i,j}]) \\
&\quad + 2\partial_k \partial_i K * (\partial_i \phi \Pi) + 2\partial_i K * (\partial_k \partial_i \phi \Pi) + \partial_k K * (\partial_i \partial_i \phi \Pi) + K * (\partial_i \partial_i \partial_k \phi \Pi) \\
&=: \partial_k P_1(x) + \partial_k P_2(x) + \partial_k P_3(x).
\end{aligned} \tag{4.4}$$

Since $\phi(x) = 1$, where $x \in B(r)$ ($0 < r \leq \rho/4$), we know that there is no singularity in $\partial_k P_2(x)$ and $\partial_k P_3(x)$. As a consequence, we have

$$\begin{aligned}
|\partial_k P_2(x)| &\leq \rho^{-3} \int_{B(\rho/2)} |\partial_k U_{i,j}| dx, \\
|\partial_k P_3(x)| &\leq \rho^{-4} \int_{B(\rho/2)} |\Pi| dx,
\end{aligned}$$

and by Hölder's inequality

$$\|\partial_k P_2(x)\|_{L^{3/2}(B(r))} \lesssim \left(\frac{r}{\rho}\right)^2 \|\partial_k U_{i,j}\|_{L^{3/2}(B(\rho/4))}, \tag{4.5}$$

$$\|\partial_k P_3(x)\|_{L^{3/2}(B(r))} \lesssim \left(\frac{r}{\rho}\right)^2 \|\Pi\|_{L^{5/4,15/7}(B(\rho/4))}. \tag{4.6}$$

Using the Hölder inequality and the Poincaré-Sobolev inequality, we see that

$$\|\partial_k U_{i,j}\|_{L^{1,3/2}(Q(\rho/2))} \leq \|u - \bar{u}_{\rho/2}\|_{L^{2,6}(Q(\rho/2))} \|\nabla u\|_{L^2(Q(\rho/2))} \lesssim \|\nabla u\|_{L^2(Q(\rho/2))}^2. \tag{4.7}$$

It follows from (4.5) and (4.7) that

$$\|\partial_k P_2(x)\|_{L^{1,3/2}(Q(r))} \lesssim \left(\frac{r}{\rho}\right)^2 \|\nabla u\|_{L^2(Q(\rho/2))}^2. \tag{4.8}$$

Notice that $\Pi - \bar{\Pi}_{\frac{\rho}{2}}$ also satisfies (4.4), we derive from (4.6) that

$$\begin{aligned}
\|\partial_k P_3(x)\|_{L^{1,3/2}(Q(r))} &\lesssim \left(\frac{r}{\rho}\right)^2 \|\Pi - \bar{\Pi}_{\frac{\rho}{2}}\|_{L^{5/4,15/7}(Q(\rho/2))} \\
&\lesssim \left(\frac{r}{\rho}\right)^2 \|\nabla \Pi\|_{L^{5/4}(Q(\rho/2))}.
\end{aligned} \tag{4.9}$$

According to the Calderón-Zygmund theorem and (4.7), we get

$$\|\partial_k P_1(x)\|_{L^{1,3/2}(Q(r))} \lesssim \|\partial_k U_{i,j}\|_{L^{1,3/2}(Q(\rho/2))} \lesssim \|\nabla u\|_{L^2(Q(\rho))}^2. \tag{4.10}$$

Combining the estimates (4.8), (4.9), (4.10), we get

$$\begin{aligned}
\|\partial_k \Pi\|_{L^{1,3/2}(Q(r))} &\lesssim \|\partial_k P_1(x)\|_{L^{1,3/2}(Q(r))} + \|\partial_k P_2(x)\|_{L^{3/2}(Q(r))} + \|\partial_k P_3(x)\|_{L^{1,3/2}(Q(r))} \\
&\lesssim \|\nabla u\|_{L^2(Q(\rho))}^2 + \left(\frac{r}{\rho}\right)^2 \|\partial_k \Pi\|_{L^{5/4}(Q(\rho))}.
\end{aligned} \tag{4.11}$$

Hence

$$P_{1,3/2}(\nabla\Pi, r) \lesssim \left(\frac{\rho}{r}\right)E(\rho) + \left(\frac{r}{\rho}\right)P_{5/4}(\nabla\Pi, \rho).$$

□

Lemma 4.3. For $0 < r \leq \rho/8$,

$$\tilde{D}_1(r) \lesssim \left(\frac{\rho}{r}\right)E^{1/\lambda}(\rho)A^{1-1/\lambda}(\rho) + \left(\frac{r}{\rho}\right)\tilde{D}_1(r), \quad (4.12)$$

where λ is defined in (3.3) and the implied positive constant does not depend on r and ρ . In addition, this lemma remains valid for $\kappa = 1$.

Proof. We get the result by replacing (4.4) with

$$\begin{aligned} \partial_k \Pi(x) &= K * \{-\phi \partial_i \partial_j [\partial_k U_{i,j}] + 2\partial_i \phi \partial_i \partial_k \Pi + \partial_k \Pi \partial_i \partial_i \phi\} \\ &= -\partial_i \partial_j K * (\phi [\partial_k U_{i,j}]) \\ &\quad + 2\partial_i K * (\partial_j \phi [\partial_k U_{i,j}]) - K * (\partial_i \partial_j \phi [\partial_k U_{i,j}]) \\ &\quad - 2\partial_i K * (\partial_i \phi \partial_k \Pi) - K * (\partial_i \partial_i \phi \partial_k \Pi) \\ &=: \partial_k P_1(x) + \partial_k P_2(x) + \partial_k P_3(x). \end{aligned} \quad (4.13)$$

and by making a slight variant of the proof of Lemma 4.2. □

Remark 4.1. This lemma gives an improvement of [7, Lemma 17, p617]. Since the standard parabolic regularization theory are not applicable to the case L_t^1 , a particularly interesting case in this lemma is $\kappa = 1$,

In the spirit of above Lemma 4.2 and [2, Lemma 4, p11], we obtain the following result for the boundary case.

Lemma 4.4. If $z = (x, t)$, $x \in \partial\mathbb{R}_+^3$, $t - \rho^2 > 0$, and $t < T$, then for $0 \leq r \leq \rho/4$, $\lambda \leq 5/4$, and $\kappa \leq 15/7$,

$$\tilde{D}_1(r) \lesssim \left(\frac{\rho}{r}\right)E^{1/\lambda}(\rho)A^{1-1/\lambda}(\rho) + \left(\frac{r}{\rho}\right)(E^{1/2}(\rho) + P_{5/4}(\nabla\Pi, \rho)), \quad (4.14)$$

where λ is defined in (3.3) and the implied positive constant does not depend on r and ρ .

Proof. A slight variant of the proof [7, Lemma 11, p608] provides the estimates

$$\tilde{D}_1(r) \lesssim \left(\frac{\rho}{r}\right)E^{1/\lambda}(\rho)A^{1-1/\lambda}(\rho) + \left(\frac{r}{\rho}\right)(E^{1/2}(\rho) + \rho^{-2}\|\Pi - \bar{\Pi}_\rho\|_{L^{\lambda,\kappa}(Q^+(\rho))}).$$

From $\lambda \leq 5/4$ and $\kappa \leq 15/7$ and Hölder's inequality, we see that

$$\rho^{-2}\|\Pi - \bar{\Pi}_\rho\|_{L^{\lambda,\kappa}(Q^+(\rho))} \lesssim \rho^{-1}\|\Pi - \bar{\Pi}_\rho\|_{L^{5/4,15/7}(Q^+(\rho))} \lesssim \rho^{-1}\|\nabla\Pi\|_{L^{5/4}(Q^+(\rho))}.$$

Combining the two estimates yields the result. □

5 Proof of Theorem 2.1

In this section, we consider the upper box-dimension of potential interior singular points for suitable weak solutions to the Navier–Stokes equations (1.1). Actually, Theorem 2.1 is a direct consequence of Proposition 2.2. The proof is based the contradiction argument, which is very well-known (see, for example, [2, 11, 18]). Thus, we suffices to prove Proposition 2.2. We divide its proof into a few steps.

Step 1) We may prove the theorem with assuming that $z = 0$. We shall show that the quantities on the left of (5.5) can be controlled by the following assumption (5.1), which was used in [18, Page, 1762, inequality (3.2)]. Assume that for some fixed $\rho \leq \rho_0$

$$\int_{Q(2\rho)} |\nabla u|^2 + |u|^{10/3} + |\Pi - \bar{\Pi}_{2\rho}|^{5/3} + |\nabla \Pi|^{5/4} dx dt \leq (2\rho)^{5/3-\gamma} \varepsilon_1 \quad (5.1)$$

From the local energy inequality we obtain that

$$\begin{aligned} & \sup_{-\rho^2 \leq t < 0} \int_{B(\rho)} |u|^2 dx + \int_{Q(\rho)} |\nabla u|^2 dx dt \\ & \lesssim \left(\int_{Q(2\rho)} |u|^{10/3} dx dt \right)^{3/5} + \int_{Q(2\rho)} |u|^2 u \cdot \nabla \phi dx dt \\ & \quad + \int_{Q(2\rho)} (\Pi - \bar{\Pi}_{2\rho}) u \cdot \nabla \phi dx dt \end{aligned} \quad (5.2)$$

by fixing $\phi(x, t)$, which is a smooth positive function supported in $Q(2\rho)$ and with value 1 on the ball $Q(\rho)$.

Step 2) Using the divergence free condition, Hölder's inequality, and the Gagliardo–Nirenberg inequality, we estimate

$$\begin{aligned} & \int_{Q(2\rho)} |u|^2 u \cdot \nabla \phi dx dt \\ & \lesssim \rho^{-1} \left\| |u|^2 - \overline{|u|^2}_{2\rho} \right\|_{L^{10/7,15/8}(Q(2\rho))} \|u\|_{L^{10/3,15/7}(Q(2\rho))} \\ & \lesssim \rho^{-1} \|u^2\|_{L^{5/3}(Q(2\rho))}^{1/2} \|u \nabla u\|_{L^{5/4}(Q(2\rho))}^{1/2} \|u\|_{L^{10/3,15/7}(Q(2\rho))} \\ & \lesssim \rho^{-1/2} \|u\|_{L^{10/3}(Q(2\rho))}^{5/2} \|\nabla u\|_{L^2(Q(2\rho))}^{1/2}, \end{aligned} \quad (5.3)$$

and

$$\begin{aligned} & \int_{Q(2\rho)} (\Pi - \bar{\Pi}_{2\rho}) u \cdot \nabla \phi dx dt \\ & \lesssim \rho^{-1} \|\Pi - \bar{\Pi}_{2\rho}\|_{L^{10/7,15/8}(Q(2\rho))} \|u\|_{L^{10/3,15/7}(Q(2\rho))} \\ & \lesssim \rho^{-1} \|\Pi - \bar{\Pi}_{2\rho}\|_{L^{5/3}(Q(2\rho))}^{1/2} \|\Pi - \bar{\Pi}_{2\rho}\|_{L^{5/4,15/7}(Q(2\rho))}^{1/2} \|u\|_{L^{10/3,15/7}(Q(2\rho))} \\ & \lesssim \rho^{-1/2} \|\Pi - \bar{\Pi}_{2\rho}\|_{L^{5/3}}^{1/2} \|\nabla \Pi\|_{L^{5/4}(Q(2\rho))}^{1/2} \|u\|_{L^{10/3}(Q(2\rho))}. \end{aligned} \quad (5.4)$$

Combining (5.2), (5.3), and (5.4) and using the assumption (5.1), we get

$$\begin{aligned}
& \sup_{-\rho^2 \leq t < 0} \int_{B(\rho)} |u|^2 dx + \int_{Q(\rho)} |\nabla u|^2 dx dt \\
& \lesssim \|u\|_{L^{10/3}(Q(2\rho))}^2 + \rho^{-1/2} \|u\|_{L^{10/3}(Q(2\rho))}^{5/2} \|\nabla u\|_{L^2(Q(2\rho))}^{1/2} \\
& \quad + \rho^{-1/2} \|u\|_{L^{10/3}(Q(2\rho))} \|\Pi - \bar{\Pi}_{2\rho}\|_{L^{5/3}(Q(2\rho))}^{1/2} \|\nabla \Pi\|_{L^{5/4}(Q(2\rho))}^{1/2} \\
& \lesssim \varepsilon_1^{3/5} \rho^{1-3\gamma/5} + \varepsilon_1 \rho^{7/6-\gamma} \\
& \lesssim \varepsilon_1^{3/5} \rho^{7/6-\gamma},
\end{aligned} \tag{5.5}$$

where we have used $\gamma \geq 5/12$. Hence we get

$$A(u, \rho) \lesssim \varepsilon_1^{3/5} \rho^{1/6-\gamma}. \tag{5.6}$$

Step 3) If we take

$$\beta = 1/6 \quad \text{and} \quad \theta = \rho^\beta < 1/8,$$

then from (4.1), (5.1), and (5.6) we get

$$\begin{aligned}
E_2(\theta\rho) & \lesssim \theta^{-1} E(\rho) + \theta^2 A(\rho) \\
& \lesssim \varepsilon_1 \theta^{-1} \rho^{2/3-\gamma} + \varepsilon_1^{3/5} \theta^2 \rho^{1/6-\gamma} \\
& \lesssim \varepsilon_1^{3/5} \rho^{-\beta+2/3-\gamma} + \varepsilon_1^{3/5} \rho^{2\beta+1/6-\gamma} \\
& \lesssim \varepsilon_1^{3/5} \rho^{1/2-\gamma}.
\end{aligned}$$

If $\gamma \leq 1/2$, then

$$E_2(\theta\rho) \lesssim \varepsilon_1^{3/5}. \tag{5.7}$$

Step 4) From Lemma 4.2 and (5.1) we have

$$\begin{aligned}
& E(\theta\rho) + P_{1,3/2}(\nabla \Pi, \theta\rho) \\
& \lesssim \theta^{-1} E(\rho) + \theta P_{5/4}(\nabla \Pi, \rho) \\
& \lesssim \varepsilon_1 \theta^{-1} \rho^{2/3-\gamma} + \varepsilon_1^{4/5} \theta \rho^{1/3-4\gamma/5} \\
& \lesssim \varepsilon_1^{4/5} \rho^{-1/6+2/3-\gamma} + \varepsilon_1^{4/5} \rho^{1/6+1/3-4\gamma/5} \\
& \lesssim \varepsilon_1^{4/5} \rho^{1/2-\gamma}.
\end{aligned} \tag{5.8}$$

Combining (5.7) and (5.8), we get

$$E(\theta\rho) + E_2(\theta\rho) + P_{5/4}(\nabla \Pi, \theta\rho) \lesssim \varepsilon_1^{3/5}.$$

This together with (2.2) yields $z = 0$ is a regular point. This completes the proof of Proposition 2.2.

□

6 Proof of Theorem 2.3

In the first place, we prove Proposition 6.1. Then Theorem 2.5 and Theorem 2.6 follow from Proposition 6.1 and (1.10). After that, we shall give the proof of Proposition 2.4. Then Theorem 2.3 is a direct consequence of Proposition 2.4 by the contradiction argument.

Proposition 6.1. *Let p, q be defined in Theorem 2.6 and denote $\alpha = \frac{2}{\frac{2}{p} + \frac{3}{q}}$. Suppose that (u, Π) is a suitable weak solution to the Navier–Stokes equations (1.7) in $Q^+(R)$. Then there holds, for any $R > 0$,*

$$\begin{aligned} & \|u\|_{L^\infty,2(Q^+(R/2))}^2 + \|\nabla u\|_{L^2(Q^+(R/2))}^2 \\ & \lesssim R^{(3\alpha-4)/\alpha} \|u\|_{L^{p,q}(Q^+(R))}^2 + R^{(3\alpha-5)/(\alpha-1)} \|u\|_{L^{p,q}(Q^+(R))}^{2\alpha/(\alpha-1)} \\ & \quad + R^{-1} \|\Pi - \bar{\Pi}_{B^+(R)}\|_{L^{\lambda,k^*}(Q^+(R))}^2 \end{aligned} \quad (6.1)$$

where the implied positive constants does not depend on R .

We divide its proof into a few steps.

Proof. **Step 1)** Fix r and ρ satisfying

$$0 < R/2 \leq r < \rho \leq R.$$

Let $\phi(x, t)$ be non-negative smooth function supported in $Q(\rho)$ such that $\phi(x, t) = 1$ on $Q(r)$ and

$$|\partial_t \phi| + |\nabla \phi|^2 + |\nabla^2 \phi| \lesssim (\rho - r)^{-2}.$$

From the local energy inequality (3.1) we have

$$\int_{B^+(\rho)} |u(x, t)|^2 \phi(x, t) dx + 2 \int_{Q^+(\rho)} |\nabla u|^2 \phi dx ds \lesssim L_1 + L_2 + L_3 \quad (6.2)$$

where

$$\begin{aligned} L_1 &= \frac{1}{(\rho - r)^2} \int_{Q^+(\rho)} |u|^2 dx ds, \\ L_2 &= \frac{1}{(\rho - r)} \int_{Q^+(\rho)} |u|^3 \phi dx ds, \\ L_3 &= \frac{1}{(\rho - r)} \int_{Q^+(\rho)} u(\Pi - \bar{\Pi}_{B^+(\rho)}) dx ds. \end{aligned}$$

In order to estimate L_1, L_2 , and L_3 , we shall use the following interpolation inequality. If k and l satisfy

$$\frac{2}{l} + \frac{3}{k} = \frac{3}{2} \quad \text{and} \quad 2 \leq l \leq \infty,$$

then

$$\begin{aligned} \|u\|_{L^{k,l}(Q(\mu))} & \lesssim \|u\|_{L^\infty,2(Q(\mu))}^{1-2/l} \|u\|_{L^{2,6}(Q(\mu))}^{2/l} \\ & \lesssim \|u\|_{L^\infty,2(Q(\mu))}^{1-2/l} (\|u\|_{L^\infty,2(Q(\mu))} + \|\nabla u\|_{L^2(Q(\mu))})^{2/l} \\ & \lesssim \|u\|_{L^\infty,2(Q(\mu))} + \|\nabla u\|_{L^2(Q(\mu))}, \end{aligned} \quad (6.3)$$

Step 2) By Hölder's inequality and (6.3), we have

$$\begin{aligned} L_1 &\lesssim \frac{\rho^{5/3}}{(\rho-r)^2} \left(\int_{Q^+(\rho)} |u|^3 dx ds \right)^{2/3} \\ &\lesssim \frac{\rho^{5/3}}{(\rho-r)^2} \rho^{(\alpha-1)} \|u\|_{L^{p,q}(Q^+(\rho))}^{2\alpha/3} \left(\|u\|_{L^\infty,2(Q^+(\rho))}^2 + \|\nabla u\|_{L^2(Q^+(\rho))}^2 \right)^{(3-\alpha)/3}. \end{aligned}$$

By Young's inequality, there is a positive constant C such that

$$L_1 \leq \frac{1}{16} \left(\|u\|_{L^\infty,2(Q^+(\rho))}^2 + \|\nabla u\|_{L^2(Q^+(\rho))}^2 \right) + \frac{C\rho^{5/\alpha}}{(\rho-r)^{6/\alpha}} \rho^{3(\alpha-1)/\alpha} \|u\|_{L^{p,q}(Q^+(\rho))}^2. \quad (6.4)$$

Similarly, by Hölder's inequality and (6.3), we have

$$\begin{aligned} L_2 &= \frac{1}{(\rho-r)} \int_{Q^+(\rho)} |u|^\alpha |u|^{3-\alpha} dx dt \\ &\lesssim \frac{1}{(\rho-r)} \|u\|_{L^{p,q}(Q^+(\rho))}^\alpha \|u\|_{L^{(3-\alpha)(\frac{p}{\alpha}), (3-\alpha)(\frac{q}{\alpha})^*}(Q^+(\rho))}^{3-\alpha} \\ &\lesssim \frac{\rho^{3(\alpha-1)/2}}{(\rho-r)} \|u\|_{L^{p,q}(Q^+(\rho))}^\alpha \|u\|_{L^{2(\frac{p}{\alpha})^*, 2(\frac{q}{\alpha})^*}(Q^+(\rho))}^{3-\alpha} \\ &\lesssim \frac{\rho^{3(\alpha-1)/2}}{(\rho-r)} \|u\|_{L^{p,q}(Q^+(\rho))}^\alpha \left(\|u\|_{L^\infty,2(Q^+(\rho))}^2 + \|\nabla u\|_{L^2(Q^+(\rho))}^2 \right)^{(3-\alpha)/2}. \end{aligned}$$

By Young's inequality, there is a positive constant C such that

$$L_2 \leq \frac{1}{16} \left(\|u\|_{L^\infty,2(Q^+(\rho))}^2 + \|\nabla u\|_{L^2(Q^+(\rho))}^2 \right) + \frac{C\rho^3}{(\rho-r)^{\frac{2}{\alpha-1}}} \|u\|_{L^{p,q}(Q^+(\rho))}^{2\alpha/(\alpha-1)}. \quad (6.5)$$

Finally, by Hölder's inequality and (6.3), we have

$$\begin{aligned} L_3 &\lesssim \frac{1}{(\rho-r)} \|\Pi - \bar{\Pi}_{B^+(\rho)}\|_{L^{\lambda,\kappa^*}(Q^+(\rho))} \|u\|_{L^{i,j}(Q^+(\rho))} \\ &\lesssim \frac{\rho^{1/2}}{(\rho-r)} \|\Pi - \bar{\Pi}_{B^+(\rho)}\|_{L^{\lambda,\kappa^*}(Q^+(\rho))} \left(\|u\|_{L^\infty,2(Q^+(\rho))}^2 + \|\nabla u\|_{L^2(Q^+(\rho))}^2 \right)^{1/2} \end{aligned}$$

where κ^* , λ , i , and j are numbers in (3.3) and (3.6). By Young's inequality, there is a positive constant C such that

$$L_3 \leq \frac{1}{16} \left(\|u\|_{L^\infty,2(Q^+(\rho))}^2 + \|\nabla u\|_{L^2(Q^+(\rho))}^2 \right) + \frac{C\rho}{(\rho-r)^2} \|\Pi - \bar{\Pi}_{B^+(\rho)}\|_{L^{\lambda,\kappa^*}(Q^+(\rho))}^2. \quad (6.6)$$

Step 3) Comgining the inequalities (6.2), (6.5), (6.6), and (6.4), we obtain that

$$\begin{aligned} &\sup_{-r^2 \leq t \leq 0} \int_{B(r)} |u|^2 dx + \int_{Q^+(r)} |\nabla u|^2 dx d\tau \\ &\leq \frac{1}{4} \left(\|u\|_{L^\infty,2(Q^+(\rho))}^2 + \|\nabla u\|_{L^2(Q^+(\rho))}^2 \right) + \frac{C\rho^{5/\alpha}}{(\rho-r)^{6/\alpha}} \rho^{3(\alpha-1)/\alpha} \|u\|_{L^{p,q}(Q^+(\rho))}^2 \\ &\quad + \frac{C\rho^3}{(\rho-r)^{2/(\alpha-1)}} \|u\|_{L^{p,q}(Q^+(\rho))}^{2\alpha/(\alpha-1)} + \frac{C\rho}{(\rho-r)^2} \|\Pi - \bar{\Pi}_{B^+(\rho)}\|_{L^{\lambda,\kappa^*}(Q^+(\rho))}^2. \end{aligned}$$

Applying the standard iteration argument, Lemma V.3.1 in [5], we get the result. \square

We end this section by giving the proof of Proposition 2.4.

Proof of Proposition 2.4. We may assume $z = 0$. First we assume that

$$\int_{Q^+(2\rho)} |\nabla u|^2 + |u|^{10/3} + |\Pi - \bar{\Pi}_{2\rho}|^{5/3} + |\nabla \Pi|^{5/4} dx dt \leq (2\rho)^{5/3-\gamma} \varepsilon_2. \quad (6.7)$$

Following the argument in [9] we shall determine a suitable parameter γ . From (6.7)

$$\rho^{-1} \|\nabla \Pi\|_{L^{5/4}(Q^+(\rho))} \lesssim \varepsilon_2^{4/5} \rho^{1/3-4\gamma/5}. \quad (6.8)$$

By the Poincaré inequality

$$\int_{Q^+(2\rho)} |u|^2 (\partial_t \phi + \Delta \phi) \lesssim \int_{Q^+(2\rho)} |\nabla u|^2 \lesssim \varepsilon_2 \rho^{5/3-\gamma}.$$

This and (5.3) and (5.4) implies that

$$\sup_{-\rho^2 \leq t < 0} \int_{B^+(\rho)} |u|^2 dx + \int_{Q^+(\rho)} |\nabla u|^2 dx dt \lesssim \rho^{5/3-\gamma} + \rho^{7/6-\gamma} \lesssim \rho^{7/6-\gamma}. \quad (6.9)$$

Consequently,

$$A(u, \rho) \lesssim \varepsilon_2 \rho^{1/6-\gamma}. \quad (6.10)$$

If we take

$$\beta = \frac{1}{4\lambda} - \frac{1}{12} - \frac{\gamma}{10} \quad \text{and} \quad \theta = \rho^\beta < \frac{1}{4},$$

then from (4.2), (4.14), and (6.8) we conclude that

$$\begin{aligned} & E(\theta\rho) + \tilde{D}_1(\theta\rho) \\ & \lesssim \theta^{-1} E(\rho) + \theta^{-1} E^{1/\lambda}(\rho) A^{1-1/\lambda}(\rho) + \theta E^{1/2}(\rho) + \theta P_{5/4}(\nabla \Pi, \rho) \\ & \lesssim \varepsilon_2 \rho^{-\beta+2/3-\gamma} + \varepsilon_2 \rho^{-\beta+(1-1/\lambda)(1/6-\gamma)+(2/3-\gamma)/\lambda} + \varepsilon_2^{1/2} \rho^{\beta+(2/3-\gamma)/2} + \varepsilon_2^{4/5} \rho^{\beta+1/3-4\gamma/5} \\ & \lesssim \varepsilon_2 \rho^{-\beta+2/3-\gamma} + \varepsilon_2^{1/2} \rho^{\beta+(2/3-\gamma)/2} + \varepsilon_2^{4/5} \rho^{\beta+1/3-4\gamma/5} \end{aligned} \quad (6.11)$$

since our choice of β satisfies

$$-\beta + (1 - 1/\lambda)(1/6 - \gamma) + (2/3 - \gamma)/\lambda = \beta + 1/3 - 4\gamma/5.$$

All the exponents of ρ on the right of (6.11) are nonnegative if

$$\gamma \leq \min \left\{ \frac{5(3\lambda - 1)}{18\lambda}, \frac{5(1 + \lambda)}{12\lambda}, \frac{5(1 + \lambda)}{18\lambda} \right\}.$$

Thus, we can choose λ sufficiently close to 1 to obtain $\gamma \leq 5/9$ and

$$E(\theta\rho) + \tilde{D}_1(\theta\rho) \lesssim \varepsilon_2^{1/2}.$$

Hence Theorem 2.5 implies that $z = 0$ is a regular point. \square

Acknowledgement

The research of Wang was partially supported by the National Natural Science Foundation of China under grant No. 11601492 and the the Youth Core Teachers Foundation of Zhengzhou University of Light Industry. Yang has been supported by the National Research Foundation of Korea (NRF) grant funded by the Korea government(MSIP) (No. 2016R1C1B2015731) and (No. 2015R1A5A1009350).

References

- [1] L. Caffarelli, R. Kohn and L. Nirenberg, Partial regularity of suitable weak solutions of the Navier–Stokes equations, *Comm. Pure. Appl. Math.*, **35** (1982), 771–831.
- [2] H. J. Choe and M. Yang, The Minkowski dimension of boundary singular points in the Navier–Stokes equations. arXiv:1805.04724
- [3] K. Choi and A. Vasseur, Estimates on fractional higher derivatives of weak solutions for the Navier–Stokes equations. *Ann. Inst. H. Poincaré Anal. Non Linéaire*, **31** (2014), 899–945.
- [4] K. Falconer, *Fractal Geometry: Mathematical Foundations and Applications* (New York: Wiley) 1990.
- [5] M. Giaquinta, *Multiple integrals in the calculus of variations and nonlinear elliptic systems*. Annals of Mathematics Studies, 105. Princeton University Press, Princeton, NJ, 1983.
- [6] C. Guevara and N. C. Phuc, Local energy bounds and ε -regularity criteria for the 3D Navier–Stokes system. *Calc. Var.*, (2017) 56:68.
- [7] S. Gustafson, K. Kang and T. Tsai, Regularity criteria for suitable weak solutions of the Navier–Stokes equations near the boundary. *J. Differential Equations.*, **226**, (2006) 594–618.
- [8] C. He, Y. Wang and D. Zhou, New ε -regularity criteria and application to the box dimension of the singular set in the 3D Navier–Stokes equations, arxiv: 1709.01382.
- [9] Y. Koh and M. Yang, The Minkowski dimension of interior singular points in the incompressible Navier–Stokes equations. *J. Differential Equations.*, **261** (2016), 3137–3148.
- [10] I. Kukavica, The fractal dimension of the singular set for solutions of the Navier–Stokes system. *Nonlinearity.*, **22** (2009), 2889–2900.
- [11] I. Kukavica and Y. Pei, An estimate on the parabolic fractal dimension of the singular set for solutions of the Navier–Stokes system. *Nonlinearity.*, **25** (2012), 2775–2783.
- [12] O. Ladyzenskaja and G. Seregin, On partial regularity of suitable weak solutions to the three-dimensional Navier–Stokes equations, *J. Math. Fluid Mech.*, **1** (1999), 356–387.
- [13] F. Lin, A new proof of the Caffarelli–Kohn–Nirenberg Theorem, *Comm. Pure Appl. Math.*, **51** (1998), 241–257.

- [14] W. Ren, Y. Wang and G. Wu, Remarks on the singular set of suitable weak solutions for the three-dimensional Navier–Stokes equations. *J. Math. Anal. Appl.* **467** (2018), 807-824.
- [15] J. Robinson and W. Sadowski, On the dimension of the singular set of solutions to the Navier–Stokes equations, *Comm. Math. Phys.*, **309** (2012), 497–506.
- [16] G. Seregin, *Local regularity of suitable weak solutions to the Navier–Stokes equations near the boundary*, *J. Math. Fluid Mech.*, **4** (2002) 1-29.
- [17] W. Wang and Z. Zhang, On the interior regularity criteria and the number of singular points to the Navier–Stokes equations, *J. Anal. Math.* **123** (2014), 139-170.
- [18] Y. Wang and G. Wu, On the box-counting dimension of potential singular set for suitable weak solutions to the 3D Navier–Stokes equations, *Nonlinearity*, **30** (2017), 1762–1772.
- [19] A. Vasseur, A new proof of partial regularity of solutions to Navier–Stokes equations, *NoDEA Nonlinear Differential Equations Appl.*, **14** (2007), 753–785.