

A better method than t-free for Robin's hypothesis

Xiaolong Wu

Ex. Institute of Mathematics, Chinese Academy of Sciences

xwu622@comcast.net

November 29, 2018

Abstract

Robin made hypothesis that $\sigma(n) < e^\gamma n \log \log n$ for all integers $n > 5040$. For a positive integer $t > 1$, an integer N is called t -free if N cannot be divided by prime power p^t for any prime p . Some works have shown if N is t -free, then N satisfies Robin's inequality, for $t=5, 7, 11, 16$. This article shows that the condition of t -free can be reduced to " N cannot be divided by 2^t ". I proved that if N cannot be divided by 2^{17} , then N satisfies Robin's inequality.

Robin made a hypothesis [Robin 1984] that the Robin's inequality

$$\sigma(n) < e^\gamma n \log \log n, \tag{RI}$$

holds for all integers $n > 5040$. Here $\sigma(n) = \sum_{d|n} d$ is the divisor sum function, γ is the Euler-Mascheroni constant, \log is the nature logarithm.

Robin also proved that his hypothesis is true if and only if Riemann hypothesis is true.

For calculation convenience, we define

$$\rho(n) := \frac{\sigma(n)}{n}.$$

Then Robin's inequality can also be written as

$$\rho(n) < e^\gamma \log \log n. \quad (\text{RI})$$

Let $N > 5040$ be an integer. Write the factorization of N as

$$N = \prod_{i=1}^r p_i^{a_i},$$

where r is the number of distinct prime factors of N , p_i are listed in increasing orders. So p_r is the largest prime factor of N .

According to [Morrill; Platt 2018], (RI) holds for all integers n , $5040 < n \leq 10^{(10^{13})}$. So, we assume $N > 10^{(10^{13})}$.

For a positive integer $t > 1$, an integer N is called t -free if N cannot be divided by prime power p^t for any prime p . Some works have shown if N is t -free, then N satisfies Robin's inequality, for $t=5$ [CLMS 2007], 7 [Solé; Planat 2012], 11 [Broughan; Trudgian 2015], 16 [Morrill; Platt 2018]. [Morrill; Platt 2018] also shows 25-free integers satisfy (RI). But that is by a different method of numerical calculation as that article said "We calculated that $R_{25}(N_{177\,244\,758\,016}) < 1 - 10^{-16}$ ".

According to abundant number theory, any possible counter-example of (RI) has its exponents of primes in decreasing order, so, the condition "t-free" is basically equivalent to exponent of prime 2 is less than t .

The previous works on t -free used $\zeta(t)$. However, we have $2^{-t} < \zeta(t) < 2^{-t+1}$ for integer $t \geq 2$, and

$$\lim_{t \rightarrow \infty} 2^t \zeta(t) = 1, \quad t \text{ runs over integers } \geq 2.$$

Hence, using $\zeta(t)$ has almost the same effect as using 2^{-t} .

This article shows that the condition of t -free can be reduced to "N cannot be divided by 2^t ". In particular, I proved: if N cannot be divided by 2^{17} , then N satisfies Robin's inequality.

Using a method similar to abundant number theory, I can improve the result to 2^{42} . I will post that improvement later.

Lemma 1. (Mertens' third theorem) *For any integer $n > 7\,713\,133\,853$, we have*

$$\sum_{p \leq n} \log \left(\frac{p}{p-1} \right) = \log \log n + \gamma + R(n), \quad (\text{L2.1})$$

where γ is the Euler-Mascheroni constant, $R(n)$ is the remainder such that

$$-\frac{0.005586}{(\log n)^2} < R(n) < \frac{0.005586}{(\log n)^2}. \quad (\text{L2.2})$$

Proof. Just follow the proof of Theorem 5.6 of [Dusart 2018] with $k = 2, \eta_2 = 0.01$.

$$\begin{aligned} |R(n)| &< \frac{0.01}{2(\log n)^2} + \frac{4}{3} \cdot \frac{0.01}{(\log n)^3} \\ &= \frac{0.01}{(\log n)^2} \left(\frac{1}{2} + \frac{4}{3 \log n} \right) < \frac{0.005586}{(\log n)^2}. \end{aligned} \quad (\text{L2.3})$$

□

Definition. Let N be an integer, p be its largest prime factor. We define a **generalized factorization** of N as the product

$$N = \prod_{p_i \leq p} p_i^{a_i}$$

where p_i runs through all primes $p_i \leq p$. If a prime $p_i < p$ does not divide N , we set $a_i = 0$, i.e. $p_i^{a_i} = p_i^0 = 1$ if p_i is not a factor of N .

Theorem 1. *Let $N > 10^{(10^{13})}$ be an integer. Define*

$$T := \frac{0.005586}{(\log \log (10^{(10^{13})}))^2}, \quad E(p) := \left\lfloor \frac{-\log T}{\log p} \right\rfloor,$$

here $\lfloor x \rfloor$ denotes the largest integer $\leq x$. Write the generalized factorization of N as

$$N = \prod_{p_i \leq p_r} p_i^{a_i},$$

where p_i are listed in increasing order and p_r is the largest prime factor of N . If $p_s^{E(p_s)}$ does not divide N for some p_s , then N satisfies (RI).

In particular, since

$$\log T = -12.04, \quad E(2) = \left\lfloor \frac{12.04}{\log 2} \right\rfloor = 17,$$

if 2^{17} does not divide N , then N satisfies (RI).

Proof. We use induction by assuming that all integers n in $(5040, N)$ with exponent of p_s less than $E(p_s)$ satisfy (RI). We know

$$\rho(N) = \prod_{p_i \leq p_r} \frac{p_i - p_i^{-a_i}}{p_i - 1} = \prod_{p_i \leq p_r} \frac{1 - p_i^{-a_i-1}}{1 - p_i^{-1}}. \quad (1.1)$$

Define

$$P(N) := \prod_{p_i \leq p_r} \frac{p_i}{p_i - 1} = \prod_{p_i \leq p_r} \frac{1}{1 - p_i^{-1}}. \quad (1.2)$$

Define

$$\begin{aligned} S(N) &:= \log P(N) - \log \rho(N) \\ &= - \sum_{p_i \leq p_r} \log(1 - p_i^{-a_i-1}) = \sum_{p_i \leq p_r} \sum_{k=1}^{\infty} \frac{1}{k p_i^{(a_i+1)k}}, \end{aligned} \quad (1.3)$$

Then apply Lemma 1 to $\log P(N)$,

$$\log \rho(N) = \log P(N) - S(N) < \log \log p_r + \gamma + R(p_r) - S(N). \quad (1.4)$$

where

$$R(p_r) = \frac{0.005586}{(\log p_r)^2}. \quad (1.5)$$

By method of Lemma 11 of [CLMS 2007] or Theorem 26 of [NY 2014], we may assume $p_r < \log N$.

Since $\log \log x + \gamma + R(x)$ is an increasing function in x , we may substitute p_r with $\log N$ in (1.4),

$$\log \rho(N) < \log \log \log N + \gamma + R(\log N) - S(N), \quad (1.6)$$

Now,

$$S(N) > \sum_{p_i \leq p_r} p_s^{-a_s-1} \geq p_s^{-a_s-1} \geq p_s^{-E(p_s)}. \quad (1.7)$$

Combine (1.5) and (1.7), we have

$$\begin{aligned} R(\log N) - S(N) &< \frac{0.005586}{(\log \log(10^{(10^{13})}))^2} - p_s^{-E(p_s)} \\ &\leq T - p_s^{\frac{\log T}{\log p_s}} = T - e^{(\log p_s) \left(\frac{\log T}{\log p_s} \right)} = T - T = 0. \end{aligned} \quad (1.8)$$

Then from (1.6) we get $\log \rho(N) < \log \log \log N + \gamma$. Take exponential and we are done. \square

Corollary 1. *Let $N > 10^{(10^{13})}$ be an integer. Define*

$$L := \prod_{p \leq p_r} p^{E(p)}.$$

If N is not divisible by L , then N satisfies (RI).

Proof. If N is not divisible by L , there must exist a prime power, say $p_s^{E(p_s)}$, of L that cannot divide N . Then N satisfies (RI) by Theorem 1. \square

Remark: The size of L can be estimated as $L \approx 10^{(10^{4.868})}$.

References

- [Broughan;Trudgian 2015] K.A. Broughan and T. Trudgian. *Robin's inequality for 11-free integers*. Integers, 15: Paper No. A12, 5, 2015.
- [CLMS 2007] Y.-J. Choie, N. Lichiardopol, P. Moree, and P. Solé. *On Robin's criterion for the Riemann hypothesis*. J. Théor. Nombres Bordeaux, 19(2):357–372, 2007.
- [Dusart 2018] P. Dusart. *Explicit estimates of some functions over primes*. Ramanujan J., 45(1):227–251, 2018.
- [Morrill;Platt 2018] T. Morrill, D. Platt. *Robin's inequality for 25-free integers and obstacles to analytic improvement*
<https://arxiv.org/abs/1809.10813>

[NY 2014] S. Nazardonyavi and S. Yakubovich. *Extremely Abundant Numbers and the Riemann Hypothesis* Journal of Integer Sequences, Vol. 17 (2014), Article 14.2.8.

[Robin 1984] G. Robin. *Grandes valeurs de la fonction somme des diviseurs et hypothèse de Riemann.* Journal de mathématiques pures et appliquées. (9), 63(2):187–213, 1984.

[Solé; Planat 2012] P. Solé and M. Planat. *The Robin inequality for 7-free integers.* Integers, 12(2):301–309, 2012.