

THE KERNEL $\frac{1}{2} + \lfloor \frac{1}{xy} \rfloor - \frac{1}{xy}$ ($0 < x, y \leq 1$) AND MERTENS SUMS

N. Watt

§1. Introduction.

In our recent joint work [HW] with Huxley, it was observed that when N is a positive integer one has

$$M(N^2) = 2M(N) - \left(\sum_{n=1}^N \frac{\mu(n)}{n} \right)^2 N^2 + \frac{(M(N))^2}{2} - \sum_{m=1}^N \sum_{n=1}^N K\left(\frac{m}{N}, \frac{n}{N}\right) \mu(m)\mu(n), \quad (1.1)$$

where $\mu(n)$ is the Möbius function, $M(x)$ is the Mertens function (given by $M(x) = \sum_{n \leq x} \mu(n)$ when $x \in \mathbb{R}$) and

$$K(x, y) = \frac{1}{2} - \left\{ \frac{1}{xy} \right\} \quad \text{for } 0 < x, y \leq 1, \quad (1.2)$$

with $\{\alpha\} := \alpha - \lfloor \alpha \rfloor = \alpha - \max\{n \in \mathbb{Z} : n \leq \alpha\}$, for $\alpha \in \mathbb{R}$: see, in particular, [HW, Equations (1.3), (1.5) and (3.5)].

Assume N is given. The final sum in (1.1) (the sum over m and n) is a quadratic form in $\mu(1), \dots, \mu(N)$: we may write it as $\mathbf{m}^T Z \mathbf{m}$, where \mathbf{m} is the column-matrix $(\mu(1), \dots, \mu(N))^T$ and $Z = Z(N)$ is the $N \times N$ matrix with elements $z_{mn} = K(\frac{m}{N}, \frac{n}{N})$. If $\Lambda \neq 0$ is an eigenvalue of Z (so that $\det(Z - \Lambda I) = 0$ when I is the $N \times N$ identity matrix) then, for the kernel $K^\dagger = K^{\dagger, N}$ given by

$$K^\dagger(x, y) = \begin{cases} \frac{1}{2} - \left\{ \frac{N^2}{\lceil Nx \rceil \lceil Ny \rceil} \right\} & \text{if } 0 < x, y \leq 1, \\ 0 & \text{if } 0 \leq x, y \leq 1 \text{ and } xy = 0, \end{cases} \quad (1.3)$$

where $\lceil \alpha \rceil := \min\{n \in \mathbb{Z} : n \geq \alpha\}$, there exists some eigenfunction $\phi^\dagger(x)$ satisfying both

$$\infty > \int_0^1 |\phi^\dagger(x)|^2 dx > 0 \quad (1.4)$$

and

$$\phi^\dagger(x) = \lambda^\dagger \int_0^1 K^\dagger(x, y) \phi^\dagger(y) dy \quad (0 \leq x \leq 1), \quad (1.5)$$

where λ^\dagger is the non-zero constant $\frac{N}{\Lambda}$, and so, by the relevant definition given in [Tr], the number $\lambda^\dagger = \frac{N}{\Lambda}$ is an eigenvalue of the kernel K^\dagger . Indeed, given the definition (1.3), it is a corollary of the discussion in [Tr, Section 1.2] (concerning kernels that are step-functions) that (1.4) and (1.5) hold if and only if, for some choice of constants b_1, \dots, b_N such that

$$Z \mathbf{b}^\dagger = \left(\frac{N}{\lambda^\dagger} \right) \mathbf{b}^\dagger \neq \mathbf{0} \quad (\mathbf{b}^\dagger = (b_1, \dots, b_N)^T), \quad (1.6)$$

one has, for $0 \leq x \leq 1$,

$$\phi^\dagger(x) = \begin{cases} b_{\lceil Nx \rceil} & \text{if } 0 < x \leq 1, \\ 0 & \text{if } x = 0. \end{cases} \quad (1.7)$$

We therefore have a one-to-one mapping, $\lambda^\dagger \mapsto \frac{N}{\lambda^\dagger}$, from the set of eigenvalues of the kernel $K^\dagger = K^{\dagger, N}$ onto the set of non-zero eigenvalues of the matrix $Z = Z(N)$, and, together with that, a one-to-one correspondence (made explicit by (1.7)) between the set of eigenspaces of K^\dagger and the set of those eigenspaces of Z that are associated with non-zero eigenvalues. In this paper we deal mainly with kernels, and matrices, that are real and symmetric: the eigenvalues of such kernels, or matrices, are real numbers (and in the case of the kernels, the eigenvalues are, by definition, non-zero).

We are interested in the eigenvalues and eigenvectors of the matrix Z due to their relevance in respect of (1.1): it being natural to consider the application, there, of the spectral decomposition of the quadratic form $\mathbf{m}^T Z \mathbf{m}$ (see [HW, Section 3] for some discussion of this matter). The eigenvalues and eigenfunctions of the kernel $K^\dagger = K^{\dagger, N}$ are of equal interest to us, for any information regarding these is translatable (by means of the correspondences noted in the previous paragraph) into information concerning the eigenvalues and eigenvectors of Z . When N is large, one may view the relevant kernel $K^{\dagger, N} : [0, 1] \times [0, 1] \rightarrow \mathbb{R}$ as being a step-function approximating the unique kernel $K : [0, 1] \times [0, 1] \rightarrow \mathbb{R}$ defined by (1.2) and the relations $K(x, 0) = K(0, x) = 0$ ($0 \leq x \leq 1$). One of our aims, in this paper, is to investigate the approximation of eigenvalues of this symmetric kernel K by non-zero eigenvalues of $Z(N)$.

We also investigate how the eigenvalues of K are distributed, as regards their size. This leads us to consider an alternative family of step-function approximations to the kernel K , different from those in the sequence $K^{\dagger, 1}, K^{\dagger, 2}, K^{\dagger, 3}, \dots$: we construct these new approximations to K in the next paragraph. The results of this second investigation will be shown to have implications concerning the distribution of the eigenvalues of the matrix $Z(N)$ ($N = 1, 2, \dots$).

Let N be a positive integer (preferably quite a large one). We take $\varepsilon = \varepsilon(N)$ to be the unique real solution of the equation

$$2 \sinh(\varepsilon) = e^{-N\varepsilon}. \quad (1.8)$$

This implies $\varepsilon > 0$. Next, we put

$$\delta = e^{-\varepsilon}, \quad (1.9)$$

and

$$x_i = \delta^{i-1} \quad \text{for } i = 1, \dots, N+1. \quad (1.10)$$

Then, putting $x_{N+2} = 0$, we have:

$$1 = x_1 > x_2 > \dots > x_{N+1} > x_{N+2} = 0. \quad (1.11)$$

When $(x, y) \in [0, 1]^2$ is such that the sets $\{x, y\}$ and $\{x_1, x_2, \dots, x_{N+2}\}$ are not disjoint, we put $K^\triangleleft(x, y) = K^{\triangleleft, N}(x, y) = 0$. In all other cases (where, still, $(x, y) \in [0, 1]^2$), there will exist a unique ordered pair $(i, j) \in \{1, 2, \dots, N+1\}^2$ such that the open rectangle

$$\mathcal{R}_{i,j} := (x_{i+1}, x_i) \times (x_{j+1}, x_j) \quad (1.12)$$

contains the point (x, y) . If, here, $i + j > N + 1$, then we put $K^\triangleleft(x, y) = 0$ (as before). If, however, $i + j \leq N + 1$, then we put $K^\triangleleft(x, y) = \mu_{i,j}$, where

$$\mu_{i,j} = \mu_{i,j}(N) := \left(\iint_{\mathcal{R}_{i,j}} dx dy \right)^{-1} \iint_{\mathcal{R}_{i,j}} K(x, y) dx dy \quad (1.13)$$

(so that $\mu_{i,j}$ is a real number, and depends only on i, j and N). Since the kernel K is symmetric, the substitution of x for y (and vice versa) in (1.13) shows that, given (1.12), we have

$$\mu_{i,j} = \mu_{j,i} \quad (i, j \in \mathbb{N} \text{ and } i + j \leq N + 1). \quad (1.14)$$

In order to make clear the reason for (1.8), and for the distinction (in the preceding) between the cases with $i + j > N + 1$ and those with $i + j \leq N + 1$, we remark that (1.8) and (1.9) ensure that $1 - \delta^2$ is equal to δ^{N+1} , so that it follows by (1.10) that one has

$$\frac{1}{x_{i+1}x_{j+1}} - \frac{1}{x_i x_j} = \frac{1 - \delta^2}{\delta^{i+j}} = \delta^{N+1-(i+j)} \leq 1 \Leftrightarrow i + j \leq N + 1, \quad (1.15)$$

when $i, j \in \mathbb{N}$. Thus, when i and j are positive integers satisfying $i + j \leq N + 1$, the set $\mathcal{S}_{i,j} := \{\lfloor \frac{1}{xy} \rfloor : (x, y) \in \mathcal{R}_{i,j}\}$ will contain no more than two elements (i.e. either $\mathcal{S}_{i,j}$ is a singleton, and the kernel $K(x, y)$ is continuous on the open rectangle $\mathcal{R}_{i,j}$, or else $\mathcal{S}_{i,j} = \{m, m - 1\}$ for some integer $m > 1$, and the set of points of $\mathcal{R}_{i,j}$ at which $K(x, y)$ is not continuous is the intersection of the hyperbola $xy = \frac{1}{m}$ with $\mathcal{R}_{i,j}$).

In Section 6 we shall discuss the kernel $K^\triangleleft = K^{\triangleleft, N}$ in greater detail. Here we make just a few observations.

Our first observation is that K^\triangleleft is a symmetric kernel (as follows by (1.16)-(1.18), below, and (1.14)).

Secondly, we note the existence of a correspondence between eigenvalues and eigenfunctions of $K^{\triangleleft, N}$ and eigenvalues and eigenvectors of a certain $N \times N$ matrix $H = H(N)$, similar to the correspondence between eigenvalues and eigenfunctions of $K^{\dagger, N}$ and eigenvalues and eigenvectors of the $N \times N$ matrix $Z(N)$ that was mentioned below (1.7). Indeed, by our above definition of the kernel K^\triangleleft , we have

$$K^\triangleleft(x, y) = \sum_{i=1}^N \sum_{j=1}^N h_{ij} \psi_i(x) \psi_j(y) \quad (0 \leq x, y \leq 1), \quad (1.16)$$

where

$$\psi_i(x) := \begin{cases} \frac{1}{\sqrt{x_i - x_{i+1}}} & \text{if } x_{i+1} < x < x_i, \\ 0 & \text{otherwise,} \end{cases} \quad (1.17)$$

and

$$h_{ij} := \begin{cases} \sqrt{(x_i - x_{i+1})(x_j - x_{j+1})} \cdot \mu_{i,j} & \text{if } i + j \leq N + 1, \\ 0 & \text{otherwise,} \end{cases} \quad (1.18)$$

and so any eigenfunction of K^\triangleleft is necessarily of the form

$$\phi^\triangleleft(x) = \sum_{i=1}^N v_i \psi_i(x) \quad (1.19)$$

(where v_1, \dots, v_N denote real constants), and must be associated with a real eigenvalue $\lambda^\triangleleft \neq 0$ such that

$$\begin{aligned} \sum_{i=1}^N v_i \psi_i(x) &= \lambda^\triangleleft \int_0^1 \sum_{i=1}^N \sum_{j=1}^N h_{ij} \psi_i(x) \psi_j(y) \sum_{k=1}^N v_k \psi_k(y) dy \\ &= \lambda^\triangleleft \sum_{i=1}^N \psi_i(x) \sum_{j=1}^N h_{ij} \sum_{k=1}^N v_k \int_0^1 \psi_j(y) \psi_k(y) dy = \lambda^\triangleleft \sum_{i=1}^N \psi_i(x) \sum_{j=1}^N h_{ij} v_j \cdot 1, \end{aligned}$$

for $0 \leq x \leq 1$ (the last equality following by virtue of (1.17) and (1.11)). It follows (using, again, (1.17) and (1.11)) that ϕ^\triangleleft , as given by (1.19), will be an eigenfunction of K^\triangleleft if and only if one has

$$\sum_{j=1}^N h_{ij} v_j = \frac{v_i}{\lambda^\triangleleft} \quad \text{for } i = 1, \dots, N, \quad (1.20)$$

so that $\mathbf{v}^\triangleleft := (v_1, \dots, v_N)^\top$ is an eigenvector of the $N \times N$ matrix $H = H(N)$ with elements h_{ij} , and the corresponding eigenvalue of H is the real number $1/\lambda^\triangleleft \neq 0$. We therefore have a one-to-one mapping, $\lambda^\triangleleft \mapsto 1/\lambda^\triangleleft$, from the set of eigenvalues of the kernel $K^\triangleleft = K^{\triangleleft, N}(x, y)$ onto the set of non-zero real eigenvalues of the $N \times N$ matrix $H = H(N)$.

By (1.18) and (1.14), the matrix H is symmetric, and so does not have any eigenvalues that are not real; it is, in fact, easily seen to be a Hankel matrix (see Section 6). This fact is undoubtedly related to the (quite straightforward) connection that exists between K and the Hankel operator Γ_h with kernel

$$h(s) = e^{-\frac{1}{2}s} K(e^{-s}, 1) = e^{-\frac{1}{2}s} \left(\frac{1}{2} - \{e^s\} \right) \quad (0 \leq s < \infty). \quad (1.21)$$

Given any (Lebesgue) measurable function $v : [0, \infty) \rightarrow \mathbb{C}$ that satisfies $\int_0^\infty |v(s)|^2 ds < \infty$, the Hankel operator Γ_h maps v to the function $\Gamma_h v : [0, \infty) \rightarrow \mathbb{C}$ given by:

$$(\Gamma_h v)(s) = \int_0^\infty h(s+t) v(t) dt \quad (0 \leq s < \infty). \quad (1.22)$$

In order to obtain the connection with the kernel K , one need only substitute $e^{-t} = y$ in (1.22): given (1.2) and (1.21), one finds thereby that the relation (1.22) is equivalent to having

$$(\Gamma_h v)(s) = e^{-\frac{1}{2}s} \int_0^1 K(e^{-s}, y) y^{-\frac{1}{2}} v(-\log y) dy \quad (0 \leq s < \infty).$$

Thus, if $\phi(x)$ is an eigenfunction of the kernel K , and λ the relevant eigenvalue of K , then, by putting $v(t) = e^{-t/2}\phi(e^{-t})$ ($0 \leq t < \infty$), one finds that

$$(\Gamma_h v)(s) = e^{-\frac{1}{2}s} \int_0^1 K(e^{-s}, y) \phi(y) dy = e^{-\frac{1}{2}s} \lambda^{-1} \phi(e^{-s}) = \lambda^{-1} v(s) \quad (1.23)$$

for $s \geq 0$, so that $\frac{1}{\lambda}$ is an eigenvalue of the operator Γ_h , while v lies in the associated eigenspace (i.e. one can verify, directly, that $v(t)$ does have the requisite property of being ‘square integrable’). We remark that K is obviously a real symmetric kernel, and that Γ_h is Hermitian, in that one has $\int_0^\infty (\Gamma_h v_1)(s) \overline{v_2(s)} ds = \int_0^\infty v_1(s) \overline{(\Gamma_h v_2)(s)} ds$ for all square integrable v_1, v_2 (this following by virtue of (1.22) and the fact that h is real valued).

The above mentioned connection between K and Γ_h is something of an afterthought in this paper: the author currently knows very little about Hankel operators. The discussion of this connection will therefore be limited to the few brief observations which follow.

Our first observation is that (1.21) implies that one has

$$\int_0^\infty |h(s)| ds \leq \int_0^\infty \frac{1}{2} e^{-\frac{1}{2}s} ds < \infty, \quad (1.24)$$

so that it follows by [Pa, Corollary 7.2] that Γ_h is compact. By (1.21) and [Pa, Theorem 7.3], we find also that Γ_h is a ‘Hilbert-Schmidt’ operator, with Hilbert-Schmidt norm:

$$\|\Gamma_h\|_{\text{HS}} := \left(\sum_i \sigma_i^2 \right)^{\frac{1}{2}} = \left(\int_0^\infty |h(s)|^2 ds \right)^{\frac{1}{2}} = \|K\| < \infty, \quad (1.25)$$

where σ_i is the i -th singular value of Γ_h , while $\|K\|$ is the positive number defined in (2.1), below (note that the rightmost equality in (1.25) is obtained by substituting $x = e^{-\frac{1}{2}(s+t)}$, $y = e^{-\frac{1}{2}(s-t)}$ in the double integral there). Since Γ_h is Hermitian, and since its eigenvalues are the reciprocals of those of K , the result (1.25) has an equivalent formulation that is the case $K^* = K$ of Lemma 2.1 (below).

By (1.24) and [Pa, Corollary 7.10] (a corollary of a result that V.V. Peller [Pe] and R.R. Coifman and R. Rochberg [C-R] discovered), it follows that if Γ_h were a ‘trace-class’ operator (i.e. if it were ‘nuclear’) then there would exist both a set $\mathcal{E} \subset (0, \infty)$ having Lebesgue measure 0 and a continuous function $h_0 : (0, \infty) \rightarrow \mathbb{C}$ satisfying $h_0(s) = h(s)$ for all $s \in (0, 1) \setminus \mathcal{E}$. If this were the case then, for any given $s \in (0, \infty)$, one could find a pair of convergent real sequences, $(s'_n), (s''_n)$ (say), that each have s as their (common) limit, and that satisfy the conditions $s'_n, s''_n \in (0, \infty) \setminus \mathcal{E}$ and $s'_n < s < s''_n$, for all $n \in \mathbb{N}$: considering (for example) the special case $s = \log 2$, one would then have $\lim_{n \rightarrow \infty} h(s'_n) = \lim_{n \rightarrow \infty} h_0(s'_n) = h_0(\log 2)$ and (similarly) $h_0(\log 2) = \lim_{n \rightarrow \infty} h_0(s''_n) = \lim_{n \rightarrow \infty} h(s''_n)$, and so would obtain $\lim_{n \rightarrow \infty} h(s'_n) = \lim_{n \rightarrow \infty} h(s''_n)$, which is absurd, since (1.21) implies that $\lim_{s \rightarrow (\log 2)^-} h(s)$ and $\lim_{s \rightarrow (\log 2)^+} h(s)$ differ (the first being $-1/\sqrt{8}$; the second: $1/\sqrt{8}$). By this *reductio ad absurdum* we may conclude that the kernel $h(s)$ in (1.21)-(1.22) is discontinuous in a way that precludes Γ_h from being a trace-class operator. One therefore has $\sum_i \sigma_i = \infty$, and so (bearing in mind our earlier observations) we deduce that

any sequence $\lambda_1, \lambda_2, \dots$ of eigenvalues of K in which each eigenvalue of K occurs just as many times as there are elements in a basis of the corresponding eigenspace is necessarily an infinite sequence satisfying

$$\sum_{i=1}^{\infty} |\lambda_i|^{-1} = \infty. \quad (1.26)$$

This observation supplies some context for our result in Theorem 6 (below): see also the Remark following that theorem.

The principal tools used in our proofs, in this paper, are certain well-known elements of the theory of linear homogenous Fredholm equations (of the second kind) that involve a real symmetric L_2 kernel: see especially [Tr, Chapter III] for the relevant theory and terminology. We need, in particular, the result from the Hilbert-Schmidt theory that is stated in Lemma 2.3 (below). Also very significant for our work are the inequalities (involving eigenvalues of real symmetric kernels) that were obtained by Weyl [We]: see Lemma 2.2, (2.7) and the beginning of Section 3 (up to Lemma 3.1) regarding these inequalities.

Machine computations, especially of approximations to eigenvalues and eigenvectors of the matrices Z and H (for selected values of N), have stimulated and informed the (mostly theoretical) work reported in this paper. Much of our earlier computational work, however, has been superseded by our (later) theoretical results: further such computational work might well be worthwhile, but we feel that this would require either additional theoretical input, or else the ability to carry out those computations for some choice of N considerably greater than 10^6 (something that this author has little hope of doing, at present). We therefore report just those numerical results that are mentioned in Section 7.

The organisation of the paper is as follows. In Section 2 we define some notation and terminology (to be used in stating our results, and throughout the paper) and state three previously known results that have a fundamental part to play in our proofs. Our principal new results (Theorems 1 to 7) are stated in Section 3. All but three of these are either corollaries, or else have short proofs that we include in Section 3: the three exceptions are Theorems 1, 3 and 4, which are proved in Sections 4, 5 and 6 (respectively). In Section 7 we briefly discuss a computational application of certain of our lemmas, along with some of the numerical results that are thereby obtained.

§2. Notation and Lemmas.

The kernel K defined in our Introduction is (in the terminology of [Tr]) a real symmetric, non-null L_2 -kernel. Thus the function $K(x, y)$ is real and measurable on $[0, 1] \times [0, 1]$, and one has both $K(x, y) = K(y, x) \in \mathbb{R}$, for $0 \leq x, y \leq 1$, and

$$0 < \|K\| := \left(\int_0^1 \int_0^1 (K(x, y))^2 dx dy \right)^{1/2} < \infty. \quad (2.1)$$

It is known (see [Tr, Section 3.8], for example) that the set of eigenvalues of such a kernel is a non-empty countable subset of $\mathbb{R} \setminus \{0\}$ that has no accumulation point in \mathbb{R} . Moreover, for each eigenvalue of such a kernel, the associated set of eigenfunctions spans a vector

space (over \mathbb{C}) of some finite dimension $m = m(\lambda, K) \in \mathbb{N}$: Tricomi [Tr] calls this m the ‘index’ of the eigenvalue, and we shall follow him in this.

Let $K^* = K^*(x, y)$ be an arbitrary real symmetric L_2 kernel with domain $[0, 1] \times [0, 1]$. We take λ_{k+}^* ($k \in \mathbb{N}$ and $k \leq p^*$) to be that monotonically increasing sequence of positive eigenvalues of K^* in which each $\lambda > 0$ that is an eigenvalue of K^* occurs exactly $m(\lambda, K^*)$ times. It may be supposed that we have here: $p^* = \infty$, if K^* has infinitely many positive eigenvalues; $p^* \in \mathbb{N} \cup \{0\}$, otherwise. Similarly, we take λ_{k-}^* ($k \in \mathbb{N}$ and $k \leq q^*$) to be that monotonically decreasing sequence of negative eigenvalues of K^* in which each $\lambda < 0$ that is an eigenvalue of K^* occurs exactly $m(\lambda, K^*)$ times. We may assume that $q^* \in \mathbb{N} \cup \{0, \infty\}$.

Definition 2.1. Let $k \in \mathbb{N}$. We put: $\nu_{k+}^* = 1/\lambda_{k+}^*$, if the condition $k \leq p^*$ is satisfied; $\nu_{k+}^* = 0$, otherwise; $\nu_{k-}^* = 1/\lambda_{k-}^*$, if the condition $k \leq q^*$ is satisfied; $\nu_{k-}^* = 0$, otherwise. The number ν_{k+}^* (resp. ν_{k-}^*) is the k -th *reciprocal positive eigenvalue* (resp. *reciprocal negative eigenvalue*) of K^* : note that we use this terminology even in cases where ν_{k+}^* , or ν_{k-}^* , has the value zero, which is never the reciprocal of an eigenvalue.

The notation used in Definition 2.1 is easily adapted for use in connection with a variety of differently named real symmetric L_2 kernels, including the kernels K , K^\dagger (or $K^{\dagger, N}$) and K^\triangleleft (or $K^{\triangleleft, N}$) that were defined in Section 1, as well as several other such kernels K' , K'' , \dots that are yet to be defined: in each case the superscript to the kernel (or the absence thereof) transfers to the ‘ λ ’, ‘ ν ’, ‘ p ’, or ‘ q ’ in question. For example, the reciprocal positive eigenvalues of $K''(x, y)$ will be ν_{1+}'' , ν_{2+}'' , \dots , while those of K are ν_{1+} , ν_{2+} , \dots .

One immediate consequence of Definition 2.1 is that, when $k \in \mathbb{N}$ one has:

$$(0, \infty) \ni \nu_{k+}^* \geq \nu_{(k+1)+}^*, \text{ if } 1 \leq k \leq p^*; \quad \nu_{k+}^* = 0, \text{ if } k > p^*; \quad (2.2)$$

$$(-\infty, 0) \ni \nu_{k-}^* \leq \nu_{(k+1)-}^*, \text{ if } 1 \leq k \leq q^*; \quad \nu_{k-}^* = 0, \text{ if } k > q^*. \quad (2.3)$$

Another consequence is that one can find two sequences of eigenfunctions, ϕ_{k+}^* ($k \in \mathbb{N}$ and $k \leq p^*$) and ϕ_{k-}^* ($k \in \mathbb{N}$ and $k \leq q^*$), say, such that, for $k \in \mathbb{N}$, $x \in [0, 1]$, one has: $(\nu_{k+}^*)\phi_{k+}^*(x) = \int_0^1 K^*(x, y)\phi_{k+}^*(y)dy$, if the condition $k \leq p^*$ is satisfied; $(\nu_{k-}^*)\phi_{k-}^*(x) = \int_0^1 K^*(x, y)\phi_{k-}^*(y)dy$, if $k \leq q^*$. Clearly we may assume that these eigenfunctions are real, for if one is not, then either its real part, or else its imaginary part, will be a satisfactory replacement for it. As is shown in [Tr, Section 3.8], these eigenfunctions $\phi_{k\pm}^*$ may be selected in such a way that the set $\{\phi_{k+}^* : k \in \mathbb{N} \text{ and } k \leq p^*\} \cup \{\phi_{k-}^* : k \in \mathbb{N} \text{ and } k \leq q^*\}$ is orthonormal with respect to the (semi-definite) inner-product

$$\langle \alpha, \beta \rangle = \int_0^1 \alpha(x)\beta(x)dx$$

(defined whenever α and β lie in $L^2([0, 1])$, the semimetric space of those Lebesgue measurable functions $f : [0, 1] \rightarrow \mathbb{R}$ that satisfy the condition $\int_0^1 (f(x))^2 dx < \infty$).

Wherever it might be relevant in what follows, we shall assume (without further comment) that the eigenfunctions $\phi_{k\pm}^*$ are chosen in just such a way. Note that, unless K^* be a null kernel (i.e. one with $\|K^*\| = 0$), the pair of sequences ϕ_{k+}^* ($k \in \mathbb{N}$ and $k \leq p^*$) and

ϕ_{k-}^* ($k \in \mathbb{N}$ and $k \leq q^*$) will not be uniquely determined by K^* (even if all eigenvalues of K^* have index equal to 1), for any member of either sequence is capable of being replaced by -1 times itself.

In [Wa] (to appear) we show that the kernel K , in particular, has infinitely many eigenvalues that are positive, and also infinitely many that are negative. Thus we have $p = q = \infty$, and so

$$0 < \lambda_{1+} \leq \lambda_{2+} \leq \cdots \quad \text{and} \quad 0 > \lambda_{1-} \geq \lambda_{2-} \geq \cdots, \quad (2.4)$$

and, by (2.2) and (2.3) (for $K^* = K$),

$$\infty > \nu_{k+} \geq \nu_{(k+1)+} > 0 > \nu_{(k+1)-} \geq \nu_{k-} > -\infty \quad (k \in \mathbb{N}). \quad (2.5)$$

Lemma 2.1. *We have*

$$\sum_{k=1}^{\infty} \left((\nu_{k+}^*)^2 + (\nu_{k-}^*)^2 \right) = \int_0^1 \int_0^1 (K^*(x, y))^2 dx dy = \|K^*\|^2,$$

where the definition of $\|K^*\|$ is analogous to that of $\|K\|$, in (2.1).

Proof. The second equality is trivial. The first only restates (using our terminology) a well-known result, the case $m = 2$ of [Tr, Section 3.10, (8)], and may also be obtained as a fairly direct consequence of the (more fundamental) relation [Tr, Section 3.9, (3)]. ■

Remark. By (2.2), (2.3) and Lemma 2.1 one has, when $m \in \mathbb{N}$,

$$(\nu_{m+}^*)^2 + (\nu_{m-}^*)^2 \leq \frac{1}{m} \sum_{k=1}^m \left((\nu_{k+}^*)^2 + (\nu_{k-}^*)^2 \right) \leq \frac{\|K^*\|^2}{m},$$

and so

$$\|K^*\| m^{-\frac{1}{2}} \geq \nu_{m+}^* \geq 0 \geq \nu_{m-}^* \geq -\|K^*\| m^{-\frac{1}{2}}. \quad (2.6)$$

Lemma 2.2 (Weyl, [We, Satz III]). *Let the functions K^* , K' and K'' , defined on the common domain $[0, 1] \times [0, 1]$, each be a real symmetric L_2 kernel for that domain. Suppose moreover that*

$$K^*(x, y) = K'(x, y) + K''(x, y) \quad (0 \leq x, y \leq 1).$$

The relevant reciprocal positive eigenvalues then satisfy the relations

$$\nu_{(m+n-1)+}^* \leq \nu_{m+}' + \nu_{n+}'' \quad (m, n \in \mathbb{N}).$$

Proof. The proof is sketched in Weyl's paper [We]: it is (essentially) an elaboration of the detailed proof that Weyl does supply for the result [We, Satz I]. ■

Remark. Subject to the hypotheses of the above lemma, one has also

$$\nu_{(m+n-1)-}^* \geq \nu'_{m-} + \nu''_{n-} \quad (m, n \in \mathbb{N}), \quad (2.7)$$

which follows by Lemma 2.2, the observation that one has $\nu_{k+}^\bullet = -\nu_{k-}^*$ ($k \in \mathbb{N}$) when $K^\bullet = -K^*$, and similar observations in respect of the reciprocal eigenvalues of the kernels $-K'$ and $-K''$.

Lemma 2.3 (Hilbert-Schmidt). *Let $\phi, \psi \in L^2([0, 1])$ and put*

$$J^* = J^*(\phi, \psi) = \int_0^1 \int_0^1 K^*(x, y) \phi(x) \psi(y) dx dy.$$

Then

$$J^* = \sum_{\mathbb{N} \ni k \leq p^*} \langle \phi, \phi_{k+}^* \rangle \langle \phi_{k+}^*, \psi \rangle \nu_{k+}^* + \sum_{\mathbb{N} \ni k \leq q^*} \langle \phi, \phi_{k-}^* \rangle \langle \phi_{k-}^*, \psi \rangle \nu_{k-}^*$$

and the sums over k that appear in this equality are absolutely convergent (when not finite or empty).

Proof. See [Tr, Section 3.11], where this result is deduced from a theorem of Hilbert and Schmidt (for which see [Tr, Section 3.10]). ■

Definitions 2.2. When $\phi \in L^2([0, 1])$ we define $\|\phi\|$ (the ‘norm’ of ϕ) by: $\|\phi\| = \sqrt{\langle \phi, \phi \rangle}$. This $\|\phi\|$ should not be confused with the notation $\|K^*\|$ defined earlier, in which $K^* = K^*(x, y)$, being a function of two variables, is not a member of $L^2([0, 1])$.

Note that one has $\|\phi\| = 0$ if and only if the set $\{x \in [0, 1] : \phi(x) \neq 0\}$ has Lebesgue measure 0: thus $\|\phi\|$ is, in fact, only a seminorm on $L^2([0, 1])$.

When $\phi \in L^2([0, 1])$ and $\|\phi\| > 0$, we put $\hat{\phi} = \|\phi\|^{-1} \phi$ (so that $\|\hat{\phi}\| = 1$).

§3. Statements of the main results.

By (2.1) and the definition of K given in Section 1, one certainly has $\|K\| < \frac{1}{2}$. We may therefore deduce from (2.5) and the case $K^* = K$ of (2.6) that one has:

$$\frac{1}{2\sqrt{m}} > \nu_{m+} > 0 > \nu_{m-} > -\frac{1}{2\sqrt{m}} \quad (m \in \mathbb{N}). \quad (3.1)$$

Note that it can be shown that $\|K\|^2 = \frac{3}{4} - \frac{\pi^2}{24} - \frac{(\log(2\pi)-1)^2}{2} + \frac{\gamma_0^2}{2} + \gamma_1$, where $(-1)^n \gamma_n/n!$ is the coefficient of $(s-1)^n$ in the Laurent series expansion of the Riemann zeta function, $\zeta(s)$, about the pole at $s=1$. We omit the proof of this result, since our main results in this paper are obtained independently of it. By this result, and known approximations for π , γ_1 and γ_0 (Euler’s constant), we estimate that $\|K\|$ is approximately 0.286. This estimate would therefore yield, via (2.6), a modest strengthening of the first and last bounds in (3.1). In respect of cases where the positive integer m is sufficiently large, we obtain, by Theorem 6 (below), a more significant strengthening of the same bounds.

Let N be a positive integer, and let the kernels K , $K^\dagger = K^{\dagger, N}$ and $K^\triangleleft = K^{\triangleleft, N}$ be as defined in Section 1. Put $K^\ddagger = K^{\ddagger, N} = K - K^\dagger$ and $K^\bowtie = K^{\bowtie, N} = K - K^\triangleleft$. We suppose henceforth that $K^\square = K^{\square, N}$ is one member of the pair $K^\dagger, K^\triangleleft$, and that we have $K^\# = K^{\#, N} = K - K^\square$: we therefore have $K^\# \in \{K^\ddagger, K^\bowtie\}$, with $K^\# = K^\ddagger$ holding if and only if $K^\square = K^\dagger$. Since $K = K^\square + K^\#$, it follows by Weyl's inequalities, Lemma 2.2 and (2.7), that we have:

$$\nu_{m_+}^\square + \nu_{n_+}^\# \geq \nu_{(m+n-1)_+} \quad \text{and} \quad \nu_{(m+n-1)_-} \geq \nu_{m_-}^\square + \nu_{n_-}^\# \quad (m, n \in \mathbb{N}). \quad (3.2)$$

Since we have $K^\square = K + (-K^\#)$, it therefore follows (by the observation used in justifying (2.7)) that Weyl's inequalities imply also that one has

$$\nu_{m_+} - \nu_{n_-}^\# \geq \nu_{(m+n-1)_+}^\square \quad \text{and} \quad \nu_{(m+n-1)_-}^\square \geq \nu_{m_-} - \nu_{n_+}^\# \quad (m, n \in \mathbb{N}). \quad (3.3)$$

Now, by (1.6) and (1.7), if $K^\square = K^\dagger$ (and otherwise by (1.19) and (1.20)), we must have $p^\square + q^\square \leq N$, with $p^\square + q^\square = N$ holding if and only if the relevant $N \times N$ matrix (which is either $Z(N)$, or $H(N)$) is non-singular. Since p^\square and q^\square have been defined (just prior to Definition 2.1) in such way as to ensure that each is a member of the set $\mathbb{N} \cup \{0, \infty\}$, we may conclude now that they are non-negative integers satisfying $p^\square, q^\square \leq N$. By this conclusion and (2.2) and (2.3), it is certainly the case that

$$\nu_{m_\pm}^\square = 0 \quad \text{for } m = N + 1, N + 2, \dots \quad (3.4)$$

By the special case $m = N + 1$ of (3.4), combined with the same case of (3.2), we find that, for all $n \in \mathbb{N}$, one has both $\nu_{n_+}^{\#, N} \geq \nu_{(N+n)_+}$ and $\nu_{n_-}^{\#, N} \leq \nu_{(N+n)_-}$. By this and (2.5), it follows that

$$\nu_{n_+}^\# > 0 > \nu_{n_-}^\# \quad (n \in \mathbb{N}). \quad (3.5)$$

Thus $K^\#$ shares with the kernel K the property of having infinitely many positive eigenvalues and infinitely many negative eigenvalues, and so we have $p^\# = q^\# = \infty$.

By (3.2) and (3.3), with $n = 1$, we have (immediately) the following result.

Lemma 3.1. *Let $N \in \mathbb{N}$. Suppose that $K^\square \in \{K^{\dagger, N}, K^{\triangleleft, N}\}$, and that $K^\# = K - K^\square$. Then, for either (consistent) choice of sign (\pm), one has*

$$\nu_{1_+}^\# \geq \nu_{m_\pm} - \nu_{m_\pm}^\square \geq \nu_{1_-}^\# \quad (m \in \mathbb{N}).$$

By Lemma 3.1 and the case $K^* = K^\#$, $m = 1$ of (2.6), we obtain the bounds

$$|\nu_{m_\pm} - \nu_{m_\pm}^\square| \leq \|K^\#\| \quad (m \in \mathbb{N}). \quad (3.6)$$

In view of (3.6), it is natural to seek a good estimate for $\|K^\#\|$: our first theorem deals with the cases in which $K^\square = K^{\dagger, N}$, with $N \in \mathbb{N}$ sufficiently large.

Theorem 1. *Let N be an integer satisfying $N \geq 25$. Then*

$$\|K^{\dagger, N}\|^2 < \left(2\sqrt{\log N} + \left(1 + \frac{1}{\sqrt{N}}\right)^8 \left(\frac{4}{9} + \frac{23}{30}N^{-1/6}\right) + \frac{\log \log(N) + 9}{\sqrt{\log N}} \right) N^{-1/2}.$$

By Theorem 1 and (3.6), we immediately obtain the following result.

Theorem 2. *Let $m \geq 1$ and $N \geq 25$ be integers. Then, for either (consistent) choice of sign (\pm), one has*

$$|\nu_{m\pm} - \nu_{m\pm}^{\dagger, N}| < \left(2\sqrt{\log N} + \left(1 + \frac{1}{\sqrt{N}}\right)^8 \left(\frac{4}{9} + \frac{23}{30}N^{-\frac{1}{6}}\right) + \frac{\log \log(N) + 9}{\sqrt{\log N}} \right)^{\frac{1}{2}} N^{-\frac{1}{4}}.$$

Our proof of Theorem 1 requires a certain elementary estimate, which we obtain in Lemma 4.2. Aided by the same elementary estimate, we are able to deduce from (1.1) the following result (to be proved in Section 5).

Theorem 3. *Let $N \in \mathbb{N}$ and $\delta \in \mathbb{R}$ satisfy the condition*

$$\delta > \sqrt{\frac{1 + \log N}{N}}. \quad (3.7)$$

Then

$$\begin{aligned} M(N^2) &= \sum_{\substack{j \in \mathbb{N} \\ |\nu_{j-}| > \delta}} \left(\sum_{n=1}^N \mu(n) \phi_{j-} \left(\frac{n}{N} \right) \right)^2 |\nu_{j-}| - \sum_{\substack{j \in \mathbb{N} \\ \nu_{j+} > \delta}} \left(\sum_{n=1}^N \mu(n) \phi_{j+} \left(\frac{n}{N} \right) \right)^2 \nu_{j+} \\ &\quad + \frac{(M(N))^2}{2} - \left(\sum_{n=1}^N \frac{\mu(n)}{n} \right)^2 N^2 + O(\delta N^2). \end{aligned}$$

Remarks.

1. By Theorem 6 (below), the sums over j occurring in Theorem 3 contain no more than $O(1 + \delta^{-1} |\log \delta|^{3/2})$ summands.
2. In [Wa2] (in preparation) we show that each eigenfunction $\phi_{j\pm}(x)$ is continuous on the interval $[0, 1]$, and differentiable at every point of the set $(0, 1] \setminus \{\frac{1}{2}, \frac{1}{3}, \frac{1}{4}, \dots\}$.

Our next theorem is an estimate for the size of $\|K^\#\|$ in cases where, for some sufficiently large positive integer N , one has $K^\square = K^{\triangleleft, N}$ (and so $K^\# = K - K^{\triangleleft, N} = K^{\triangleright, N}$).

Theorem 4. *Let N be an integer satisfying $N \geq 3$. Then*

$$\|K^{\triangleright, N}\|^2 \leq \left(\frac{7}{60} + O\left((\log N)^{-\frac{2}{3}}\right) \right) \cdot N^{-1} \log^3 N.$$

We prove Theorem 4 in Section 6. By Theorem 4 and (3.6), we have the following result (analogous to Theorem 2).

Theorem 5. *Let $m \geq 1$ and $N \geq 3$ be positive integers. Then, for either (consistent) choice of sign (\pm) , one has*

$$|\nu_{m\pm} - \nu_{m\pm}^{\triangleleft, N}| \leq \left(\sqrt{\frac{7}{60}} + O\left((\log N)^{-\frac{2}{3}}\right) \right) \cdot N^{-\frac{1}{2}} \log^{\frac{3}{2}} N .$$

By (2.5), Lemma 2.1 and the case $K^\# = K^{\bowtie, N}$ ($K^\square = K^{\triangleleft, N}$) of the same inequalities, $\nu_{n+}^{\#, N} \geq \nu_{(N+n)+}$ and $\nu_{n-}^{\#, N} \leq \nu_{(N+n)-}$, that led us to (3.5) we have

$$\|K^{\bowtie, N}\|^2 = \sum_{n=1}^{\infty} \left((\nu_{n+}^{\bowtie, N})^2 + (\nu_{n-}^{\bowtie, N})^2 \right) \geq \sum_{n=1}^{\infty} \left((\nu_{(N+n)+})^2 + (\nu_{(N+n)-})^2 \right) . \quad (3.8)$$

Note that this is just a special case of the result [We, Satz VIII] (attributed to Schmidt [Sch, Chapter 4, Section 18], in [We]). Using one of the simpler ideas occurring in the proof of [We, Theorem IX], we deduce from (3.8) and Theorem 5 the following result.

Theorem 6. *Let $k \geq 9$ be an integer. Then*

$$(\nu_{k+})^2 + (\nu_{k-})^2 \leq \left(\frac{21}{40} + O\left((\log k)^{-\frac{2}{3}}\right) \right) \cdot k^{-2} \log^3 k .$$

Proof. Let k , M and N , be positive integers, with $k \geq 9$. By (3.8) and (2.5), we have

$$\|K^{\bowtie, N}\|^2 > \sum_{n=1}^M \left((\nu_{(N+n)+})^2 + (\nu_{(N+n)-})^2 \right) \geq M \left((\nu_{(N+M)+})^2 + (\nu_{(N+M)-})^2 \right) .$$

By this and Theorem 4, it follows that when $N \geq 3$ one has

$$(\nu_{(N+M)+})^2 + (\nu_{(N+M)-})^2 \leq \left(\frac{7}{60} + O\left((\log N)^{-\frac{2}{3}}\right) \right) \cdot \left(\frac{\log^3 N}{MN} \right) ,$$

for both choices of sign (\pm) .

By specialising the above to the cases in which one has $N = \lceil \frac{k}{3} \rceil \geq \frac{k}{3} \geq 3$ and $M = k - N \geq \frac{2}{3}(k-1) > 5$, we find that $(\nu_{k+})^2 + (\nu_{k-})^2 \leq \left(\frac{7}{60} + O\left((\log k)^{-\frac{2}{3}}\right) \right) \cdot (\log k)^3 / \left(\frac{2}{9} k^2 \right)$, which is equivalent to the result stated in the theorem ■

Remark. Theorem 6 is, in a certain sense, nearly best-possible. For if one had, say,

$$(\nu_{k+})^2 + (\nu_{k-})^2 = O\left(k^{-2}(\log k)^{-(2+2\epsilon)}\right) \quad (k = k_0 + 1, k_0 + 2, \dots) ,$$

for some positive integer $k_0 \geq 8$, then this, together with (3.1), would imply that

$$\begin{aligned} \sum_{k=1}^{\infty} (|\nu_{k+}| + |\nu_{k-}|) &\leq \sum_{k=1}^{k_0} (|\nu_{k+}| + |\nu_{k-}|) + O\left(\sum_{k=k_0+1}^{\infty} k^{-1}(\log k)^{-(1+\varepsilon)}\right) \\ &\leq \sum_{k=1}^{k_0} k^{-\frac{1}{2}} + O\left(\int_{k_0}^{\infty} x^{-1}(\log x)^{-(1+\varepsilon)}\right) \leq k_0 + O\left(\frac{1}{\varepsilon(\log k_0)^\varepsilon}\right) < \infty, \end{aligned}$$

and so would contradict (1.26) (given that the infinite series occurring there is, by Definition 2.1, just some rearrangement of the series $|\nu_{1+}| + |\nu_{1-}| + |\nu_{2+}| + |\nu_{2-}| + \dots$). Since we trust our earlier derivation of (1.26) (based on [Pa, Corollary 7.10]), we therefore conclude that, for any fixed choice of C, ε with $C \geq 1$ and $\varepsilon > 0$, there must exist infinitely many distinct positive integers k for which the sum $(\nu_{k+})^2 + (\nu_{k-})^2$ is greater than $Ck^{-2}(\log k)^{-(2+2\varepsilon)}$. Thus the upper bound in Theorem 6 falls short of being best-possible by no more than a factor $(\log k)^{5+o(1)}$.

Our final result is a consequence of the bounds obtained in Theorems 1, 4 and 6.

Theorem 7. *Let the symbol \pm denote an arbitrary choice of sign. Let $c > 0$. Then there exists integers $k_1 = k_1(c) \geq 25$, $k_2 = k_2(c) \geq 9$ (each depending only on c) such that, whenever the kernel K^\square and numbers N, k satisfy*

$$k, N \in \mathbb{N}, \quad K^\square \in \{K^{\dagger, N}, K^{\triangleleft, N}\} \quad (3.9)$$

and

$$N \geq k \geq \begin{cases} k_1 & \text{if } K^\square = K^{\dagger, N}, \\ k_2 & \text{if } K^\square = K^{\triangleleft, N}, \end{cases} \quad (3.10)$$

one has

$$|\nu_{k\pm}^\square| \leq e^c \cdot \min \left\{ \frac{S}{\theta k} + \frac{1}{\sqrt{(1-\theta)Tk}} : 0 < \theta < 1 \right\}, \quad (3.11)$$

where

$$S = S(k) := \sqrt{\frac{21}{40}} \cdot (\log k)^{\frac{3}{2}} \quad (3.12)$$

and

$$T = T(K^\square) := \begin{cases} \sqrt{\frac{N}{4 \log N}} & \text{if } K^\square = K^{\dagger, N}, \\ \frac{60N}{7 \log^3 N} & \text{if } K^\square = K^{\triangleleft, N}. \end{cases} \quad (3.13)$$

The minimum in (3.11) is attained when $\theta = \theta_1 = \theta_1(\kappa)$, where

$$\kappa = \frac{k}{4S^2T} > 0 \quad (3.14)$$

and $\theta_1 \in (0, 1)$ is determined (uniquely) by the equation

$$(1 - \theta_1)^3 \theta_1^{-4} = \kappa. \quad (3.15)$$

When (3.9), (3.10) and (3.12)-(3.15) hold, and $0 < \kappa \leq 2$, one has:

$$|\nu_{k\pm}^\square| \leq \frac{e^c S}{k} \cdot \left(\frac{1 + 2(\kappa\theta_1)^{\frac{1}{3}}}{\theta_1} \right) \leq \frac{e^c S}{k} \cdot \min \left\{ 6, \exp \left(3\kappa^{\frac{1}{3}} \right) \right\}. \quad (3.16)$$

When (3.9), (3.10) and (3.12)-(3.15) hold, and $\kappa \geq 2$, one has:

$$|\nu_{k\pm}^\square| \leq \frac{e^c}{\sqrt{Tk}} \cdot \left(\frac{1 + \frac{1}{2}(\kappa(1-\theta_1))^{-\frac{1}{4}}}{(1-\theta_1)^{\frac{1}{2}}} \right) \leq \frac{e^c}{\sqrt{Tk}} \cdot \min \left\{ \frac{3}{\sqrt{2}}, \exp \left((2/\kappa)^{\frac{1}{4}} \right) \right\}. \quad (3.17)$$

Proof. We begin by taking note of certain relevant consequences of results that were stated earlier in this section. By Theorem 1, and a short calculation, one finds that

$$\|K^{\ddagger, N}\|^2 < \left(1 + \frac{4}{\sqrt{\log N}} \right) \left(\frac{N}{4 \log N} \right)^{-\frac{1}{2}} \quad (N = 25, 26, 27, \dots), \quad (3.18)$$

while Theorem 2 implies (similarly) the existence of an absolute constant $A_0 > 0$ such that

$$\|K^{\bowtie, N}\|^2 \leq \left(1 + \frac{A_0}{(\log N)^{\frac{2}{3}}} \right) \left(\frac{60N}{7 \log^3 N} \right)^{-1} \quad (N = 3, 4, 5, \dots). \quad (3.19)$$

Theorem 6 implies the existence of an absolute constant $A_1 > 0$ such that

$$\begin{aligned} (\nu_{m\pm})^2 &\leq \frac{21}{40} \left(1 + \frac{A_1}{(\log m)^{\frac{2}{3}}} \right) \cdot m^{-2} \log^3 m \\ &\leq \frac{21}{40} \left(1 + \frac{A_1}{(\log k)^{\frac{2}{3}}} \right) \cdot m^{-2} \log^3 k \quad (m, k \in \mathbb{N} \text{ and } 9 \leq m \leq k). \end{aligned} \quad (3.20)$$

In cases where $m < 9$ we may appeal instead to (3.1), which implies that

$$(\nu_{m\pm})^2 < \frac{1}{4m} < \frac{\log^3 k}{4m^2} \quad (m, k \in \mathbb{N} \text{ and } m < 9 \leq k). \quad (3.21)$$

We put $A_2 = \max\{A_0, A_1\}$, $k_1 = \left\lceil \exp \left(\max \left\{ \log(25), 4c^{-2}, (2A_1^{-1}c)^{-\frac{3}{2}} \right\} \right) \right\rceil$ and $k_2 = \left\lceil \exp \left(\max \left\{ \log(9), (2A_2^{-1}c)^{-\frac{3}{2}} \right\} \right) \right\rceil$. By (3.18)-(3.21) (and the definitions of the kernels $K^{\ddagger, N}$ and $K^{\bowtie, N}$) we find that, subject to both (3.9) and (3.10) holding, one will have

$$\|K^\#\|^2 < \frac{e^{2c}}{T} \quad (3.22)$$

and

$$(\nu_{m\pm})^2 \leq \frac{e^{2c} S^2}{m^2} \quad (m = 1, \dots, k), \quad (3.23)$$

where $K^\# = K - K^\square$, and S and T are as stated in (3.12) and (3.13). We shall assume henceforth that k , N and K^\square satisfy the conditions (3.9) and (3.10) (there is, otherwise, nothing for us to prove), that $K^\# = K - K^\square$, and also that S , T , κ and θ_1 are as specified in (3.12)-(3.15).

By (3.3) and (3.5), one has

$$|\nu_{k\pm}^\square| \leq |\nu_{m\pm}| + |\nu_{n\mp}^\#| \quad (m, n \in \mathbb{N} \text{ such that } m + n = k + 1), \quad (3.24)$$

where $\mp(\pm 1) = -1$. By (2.6) and (3.22), one obtains the bounds

$$|\nu_{n\mp}^\#| \leq \frac{\|K^\#\|}{\sqrt{n}} < \frac{e^c}{\sqrt{Tn}} \quad (n \in \mathbb{N}). \quad (3.25)$$

By (3.23), (3.24) and (3.25) applied with $m = \lceil \theta k \rceil \geq \theta k > 0$ and $n = k + 1 - m > (1 - \theta)k > 0$ (θ here denoting an arbitrary member of the open interval $(0, 1)$), one obtains a set of upper bounds for $|\nu_{k\pm}^\square|$ from which the result (3.11) directly follows.

As for the remainder of the proof, we note firstly that elementary calculus allows one to determine that the minimum of the function $f(\theta) := (S/k)\theta^{-1} + (Tk)^{-1/2}(1 - \theta)^{-1/2}$ on the interval $(0, 1)$ is attained when $\theta = \theta_1$: the details of this are straightforward, and so are omitted here. It is similarly straightforward to show by elementary algebra that, in both (3.16) and (3.17), the first inequality is equivalent to the inequality $|\nu_{k\pm}| \leq e^c f(\theta_1)$ (with $f(\theta)$ as defined in the preceding sentence): those inequalities therefore follow immediately from (3.11), since θ_1 must (by its definition) satisfy $0 < \theta_1 < 1$.

We consider now the final inequalities in (3.16) and (3.17). If $0 < \kappa \leq 2$ then it follows by (3.15) that the number $\theta_1 \in (0, 1)$ must lie in the shorter interval $[\frac{1}{2}, 1)$: for the function of θ_1 on the left-hand side of Equation (3.15) is strictly monotonic decreasing for $0 < \theta_1 < 1$, and is equal to 2 when $\theta_1 = \frac{1}{2}$. For the same reasons, one has $\theta_1 \in (0, \frac{1}{2}]$ if $\kappa \geq 2$. Thus, if one supposes that $\kappa \leq 2$, then (using (3.15)) one finds that

$$\frac{1}{\theta_1} = 1 + \frac{1 - \theta_1}{\theta_1} = 1 + (\kappa\theta_1)^{\frac{1}{3}} \leq \min \left\{ 2, 1 + \kappa^{\frac{1}{3}} \right\},$$

and so (also) that $1 + 2(\kappa\theta_1)^{\frac{1}{3}} \leq 2 \cdot \min \{ 2, 1 + \kappa^{\frac{1}{3}} \} - 1 = \min \{ 3, 1 + 2\kappa^{\frac{1}{3}} \}$; since $(2)(3) = 6$ and $(1 + \kappa^{\frac{1}{3}})(1 + 2\kappa^{\frac{1}{3}}) < \exp(\kappa^{\frac{1}{3}}) \exp(\kappa^{\frac{2}{3}}) = \exp(3\kappa^{\frac{1}{3}})$, we may therefore conclude that the final inequality in (3.16) is valid. Suppose now that one has instead $\kappa \geq 2$, so that $\frac{1}{2} \leq 1 - \theta_1 < 1$. Then one has both

$$\frac{1}{1 - \theta_1} = 1 + \frac{\theta_1}{1 - \theta_1} \leq 1 + \frac{\theta_1}{(\frac{1}{2})^{\frac{1}{4}}(1 - \theta_1)^{\frac{3}{4}}} = 1 + \left(\frac{2}{\kappa} \right)^{\frac{1}{4}} \leq \min \left\{ 2, \exp \left(2^{\frac{1}{4}} \kappa^{-\frac{1}{4}} \right) \right\}$$

(using (3.15) for the second equality here) and $1 + \frac{1}{2}\kappa^{-\frac{1}{4}}(1 - \theta_1)^{-\frac{1}{4}} \leq 1 + 2^{-\frac{3}{4}}\kappa^{-\frac{1}{4}} \leq \min \left\{ \frac{3}{2}, \exp(2^{-\frac{3}{4}}\kappa^{-\frac{1}{4}}) \right\}$; since $(\frac{3}{2})(2^{\frac{1}{2}}) = 3/\sqrt{2}$ and $\exp(2^{-\frac{3}{4}}\kappa^{-\frac{1}{4}}) = (\exp(2^{\frac{1}{4}}\kappa^{-\frac{1}{4}}))^{\frac{1}{2}}$, we therefore are able to conclude that the final inequality in (3.17) is valid. This completes the proof of the theorem \blacksquare

Remarks.

1. The bounds in (3.17) (when applicable) will imply

$$|\nu_{k\pm}^\square| \leq \frac{e^{2c}}{\sqrt{Tk}}, \quad (3.26)$$

provided that the number κ (defined in the theorem) satisfies $\kappa \geq 2c^{-4}$. Given (3.14), this last lower bound on κ is equivalent to the relation $k/\log^3 k \geq \frac{21}{5}c^{-4}T$, and so is certainly satisfied if, in addition to (3.9) and (3.10), one has $k \geq \frac{21}{5}c^{-4}T \log^3 N$. In the case $K^\square = K^{\triangleleft, N}$, the bound $k \geq \frac{21}{5}c^{-4}T \log^3 N$ would combine with the upper bound on k in (3.10) to give $1 \geq k/N \geq 36c^{-4}$, and so would imply that $\log k$ differs from $\log N$ by at most $O(|\log c|)$. Therefore, given that we expect that in any practical application of Theorem 7 one would fix c , and so have $|\log c| = O(1)$, we find it more interesting to observe that, when one has instead $K^\square = K^{\dagger, N}$, the bound (3.26) becomes

$$|\nu_{k\pm}^{\dagger, N}| \leq e^{2c} \cdot \left(\frac{N}{4 \log N} \right)^{-\frac{1}{4}} k^{-\frac{1}{2}},$$

and is valid for all $N, k \in \mathbb{N}$ satisfying $N \geq k \geq \max\{k_1(c), \frac{21}{10}c^{-4}N^{\frac{1}{2}}(\log N)^{\frac{5}{2}}\}$.

2. The bounds in (3.16) (when applicable) will imply

$$|\nu_{k\pm}^\square| \leq \frac{e^{2c}S}{k} = \frac{e^{2c} \cdot \left(\frac{21}{40}\right)^{\frac{1}{2}} \log^{\frac{3}{2}} k}{k}, \quad (3.27)$$

provided that κ (given by (3.12)-(3.14)) satisfies $\kappa \leq (c/3)^3$. This last upper bound on κ holds if and only if

$$\frac{k}{\log^3 k} \leq \frac{7}{90}c^3T. \quad (3.28)$$

We remark that, given that $\min\{k_1(c), k_2(c)\} \geq 9$, it may be shown that (3.28) will hold if one has both (3.10) and $\exp(c^{-\frac{3}{2}}) \leq k \leq \frac{7}{90}T \log T$.

In the particular case where $c = \left(\frac{3}{2}\right)^{\frac{1}{3}}$ and $K^\square = K^{\triangleleft, N}$, one finds (using (3.13)) that the condition (3.28) becomes $k/\log^3 k \leq \frac{2}{3}c^3N/\log^3 N = N/\log^3 N$, and so is satisfied whenever $e^3 \leq k \leq N$. By this and (3.27), it follows that one has

$$|\nu_{k\pm}^{\triangleleft, N}| \leq \frac{\exp(12^{\frac{1}{3}}) \cdot \left(\frac{21}{40}\right)^{\frac{1}{2}} \log^{\frac{3}{2}} k}{k} < \frac{8 \log^{\frac{3}{2}} k}{k},$$

for all $k, N \in \mathbb{N}$ satisfying $N \geq k \geq \max\{e^3, k_2((3/2)^{\frac{1}{3}})\}$.

§4. Proof of Theorem 1.

We assume that N is a positive integer and that $K^\ddagger = K^{\ddagger, N}$ is the kernel given by $K^\ddagger = K - K^\dagger$, where the kernels K and $K^\dagger = K^{\dagger, N}$ are as defined in Section 1. Our definition of $\|K^\ddagger\|$ is analogous to that of $\|K\|$, in (2.1).

Definition 4.1. We put

$$\mathcal{U}(N) = \bigcup_{m \in \mathbb{N}} \left\{ (p, q) \in \mathbb{Z} \times \mathbb{Z} : 0 \leq p, q < N \text{ and } \frac{pq}{N^2} < \frac{1}{m} < \frac{(p+1)(q+1)}{N^2} \right\}.$$

Remark. This definition implies that when $N = 1$ the set $\mathcal{U}(N)$ is just the singleton $\{(0, 0)\}$. We therefore have $|\mathcal{U}(1)| = 1$.

Lemma 4.1. Let N be an integer satisfying $N \geq 25$. Then

$$\|K^\ddagger\|^2 < |\mathcal{U}(N)|N^{-2} + \left(1 + \frac{1}{\sqrt{N}}\right)^8 \left(\frac{4}{9} + \frac{23}{30}N^{-1/6}\right) N^{-1/2}.$$

Proof. We have

$$\|K^\ddagger\|^2 = \sum_{r=0}^{N-1} \sum_{s=0}^{N-1} \int_{\frac{r}{N}}^{\frac{r+1}{N}} \left(\int_{\frac{s}{N}}^{\frac{s+1}{N}} (K^\ddagger(x, y))^2 dy \right) dx = \sum_{r=0}^{N-1} \sum_{s=0}^{N-1} I_{r,s}(N) \quad (\text{say}). \quad (4.1)$$

Let r and s be integers satisfying $0 \leq r, s < N$. For points (x, y) that lie in the open square

$$\mathcal{Q}_{r,s}(N) = \left(\frac{r}{N}, \frac{r+1}{N}\right) \times \left(\frac{s}{N}, \frac{s+1}{N}\right) \subset (0, 1) \times (0, 1), \quad (4.2)$$

we have, by (1.2) and (1.3),

$$\begin{aligned} K^\ddagger(x, y) &= K(x, y) - K^\dagger(x, y) \\ &= \left\{ \frac{N^2}{\lceil Nx \rceil \lceil Ny \rceil} \right\} - \left\{ \frac{1}{xy} \right\} \\ &= \left\{ \frac{N^2}{(r+1)(s+1)} \right\} - \left\{ \frac{1}{xy} \right\} \\ &= \frac{N^2}{(r+1)(s+1)} - \left[\frac{N^2}{(r+1)(s+1)} \right] - \left(\frac{1}{xy} - \left[\frac{1}{xy} \right] \right) \\ &= \left(\frac{N^2}{(r+1)(s+1)} - \frac{1}{xy} \right) + \left(\left[\frac{1}{xy} \right] - \left[\frac{N^2}{(r+1)(s+1)} \right] \right). \end{aligned} \quad (4.3)$$

Since fractional parts always lie in the interval $[0, 1)$, it follows by (4.3) that we have $-1 < K^\ddagger(x, y) < 1$ for all $(x, y) \in \mathcal{Q}_{r,s}(N)$. It is therefore trivially the case that

$$I_{r,s}(N) \leq \int_{\frac{r}{N}}^{\frac{r+1}{N}} \left(\int_{\frac{s}{N}}^{\frac{s+1}{N}} 1^2 dy \right) dx = \frac{1}{N^2}. \quad (4.5)$$

For $(x, y) \in \mathcal{Q}_{r,s}(N)$, we plainly have $0 \leq rs/N^2 < xy < (r+1)(s+1)/N^2$, and so $\infty > 1/(xy) > N^2/((r+1)(s+1))$. Hence, supposing $(x, y) \in \mathcal{Q}_{r,s}(N)$, the relation $\lfloor \frac{1}{xy} \rfloor \geq \lfloor \frac{N^2}{(r+1)(s+1)} \rfloor$ holds, and we must either have

$$\left\lfloor \frac{1}{xy} \right\rfloor = \left\lfloor \frac{N^2}{(r+1)(s+1)} \right\rfloor, \quad (4.6)$$

or else:

$$\frac{N^2}{(r+1)(s+1)} < m \leq \frac{1}{xy}, \quad \text{for some } m \in \mathbb{Z}.$$

In the latter case we have

$$\frac{(r+1)(s+1)}{N^2} > \frac{1}{m} \geq xy > \frac{rs}{N^2},$$

and so the point $(r, s) \in \mathbb{Z} \times \mathbb{Z}$ is an element of the set $\mathcal{U}(N)$ that is given by Definition 4.1. Therefore, when $(r, s) \notin \mathcal{U}(N)$, the relation (4.6) holds for all $(x, y) \in \mathcal{Q}_{r,s}(N)$.

Suppose now that $(r, s) \notin \mathcal{U}(N)$, so that (4.6) holds when $(x, y) \in \mathcal{Q}_{r,s}(N)$. Then, in view of (4.1) and (4.4), we find that

$$I_{r,s}(N) = \int_{r/N}^{(r+1)/N} \left(\int_{s/N}^{(s+1)/N} \left(\frac{N^2}{(r+1)(s+1)} - \frac{1}{xy} \right)^2 dy \right) dx. \quad (4.7)$$

Note, in addition, that since we suppose $(r, s) \notin \mathcal{U}(N)$ and $r, s \in [0, N] \cap \mathbb{Z}$, it follows from Definition 4.1 that r and s must both be (strictly) positive integers, and that there can be no integer contained within the open interval $(\frac{N^2}{(r+1)(s+1)}, \frac{N^2}{rs})$. We therefore must have

$$\frac{N^2}{rs} - \frac{N^2}{(r+1)(s+1)} \leq 1,$$

and so $(r+s+1)N^2 \leq rs(r+1)(s+1)$, which (as we have $0 < r, s < N$) implies:

$$\min\{r+1, s+1\} > \frac{N}{\sqrt{\max\{r+1, s+1\}}} \geq \sqrt{N} > 1. \quad (4.8)$$

We rewrite the double integral in (4.7), by means of the substitutions:

$$\frac{r+1}{Nx} = 1 + \frac{\theta}{r}, \quad \frac{s+1}{Ny} = 1 + \frac{\phi}{s}.$$

This gives us

$$\begin{aligned} I_{r,s}(N) &= \frac{N^2}{r(r+1)s(s+1)} \int_0^1 \left(\int_0^1 \left(1 - \left(1 + \frac{\theta}{r} \right)^{-1} \left(1 + \frac{\phi}{s} \right)^{-1} \right)^2 d\phi \right) d\theta \\ &< \frac{N^2}{r(r+1)s(s+1)} \int_0^1 \left(\int_0^1 \left(\frac{\theta}{r} + \frac{\phi}{s} \right)^2 d\phi \right) d\theta \\ &= \frac{N^2}{3r^3(r+1)s(s+1)} + \frac{N^2}{2r^2(r+1)s^2(s+1)} + \frac{N^2}{3r(r+1)s^3(s+1)}. \end{aligned}$$

By this and (4.8), we obtain (after putting $m = r + 1$, $n = s + 1$, and then exploiting some symmetries) the upper bound:

$$\sum_{\substack{0 \leq r, s < N \\ (r, s) \notin \mathcal{U}(N)}} I_{r, s}(N) \leq \left(1 - \frac{1}{\sqrt{N}}\right)^{-4} N^2 \left(\frac{2}{3}S_{2,4} + \frac{2}{3}S_{4,2} + S_{3,3}\right), \quad (4.9)$$

where

$$S_{\alpha, \beta} = \sum_{N^{2/3} < n \leq N} \frac{1}{n^\alpha} \sum_{N/\sqrt{n} < m \leq n} \frac{1}{m^\beta}.$$

Using the bounds

$$\sum_{\ell > L} \frac{1}{\ell^\sigma} < \frac{1}{L^\sigma} + \int_L^\infty \frac{dx}{x^\sigma} = \left(1 + \frac{\sigma - 1}{L}\right) \frac{L^{1-\sigma}}{\sigma - 1} \quad (L > 0 \text{ and } \sigma > 1),$$

we find that

$$\begin{aligned} S_{3,3} &< \sum_{N^{2/3} < n \leq N} \frac{1}{n^3} \left(1 + \frac{2}{N/\sqrt{n}}\right) \frac{(N/\sqrt{n})^{-2}}{2} \\ &\leq \left(1 + \frac{2}{\sqrt{N}}\right) \left(\frac{1}{2N^2}\right) \sum_{n > N^{2/3}} \frac{1}{n^2} \\ &< \left(1 + \frac{2}{\sqrt{N}}\right) \left(\frac{1}{2N^2}\right) \left(1 + \frac{1}{N^{2/3}}\right) N^{-2/3} < \left(1 + \frac{1}{\sqrt{N}}\right)^3 \left(\frac{1}{2N^{8/3}}\right) \quad (\text{say}). \end{aligned}$$

We obtain, similarly, the bounds

$$\begin{aligned} S_{4,2} &< \sum_{N^{2/3} < n \leq N} \frac{1}{n^4} \left(1 + \frac{1}{N/\sqrt{n}}\right) (N/\sqrt{n})^{-1} \\ &\leq \left(1 + \frac{1}{\sqrt{N}}\right) \left(\frac{1}{N}\right) \sum_{n > N^{2/3}} \frac{1}{n^{7/2}} \\ &< \left(1 + \frac{1}{\sqrt{N}}\right) \left(\frac{1}{N}\right) \left(1 + \frac{5/2}{N^{2/3}}\right) \left(\frac{1}{(5/2)N^{5/3}}\right) \\ &< \left(1 + \frac{1}{\sqrt{N}}\right)^3 \left(\frac{2}{5N^{8/3}}\right) \end{aligned}$$

and

$$S_{2,4} < \sum_{N^{2/3} < n \leq N} \frac{1}{n^2} \left(1 + \frac{3}{N/\sqrt{n}}\right) \frac{(N/\sqrt{n})^{-3}}{3} < \left(1 + \frac{1}{\sqrt{N}}\right)^3 \left(\frac{1}{3N^3}\right) \sum_{n=1}^N \frac{1}{\sqrt{n}}.$$

Since $N > 1$ we have

$$\sum_{n=1}^N \frac{1}{\sqrt{n}} < (-1) + \sum_{n=1}^N \frac{2}{\sqrt{n} + \sqrt{n-1}} = (-1) + 2 \sum_{n=1}^N (\sqrt{n} - \sqrt{n-1}) = 2\sqrt{N} - 1,$$

and so may deduce from the above bounds for $S_{2,4}$ that

$$S_{2,4} < \left(1 + \frac{1}{\sqrt{N}}\right)^3 \left(\frac{2}{3N^{5/2}}\right).$$

By (4.9) and the estimates just obtained, we find that

$$\begin{aligned} \sum_{\substack{0 \leq r, s < N \\ (r, s) \notin \mathcal{U}(N)}} I_{r,s}(N) &\leq \left(1 - \frac{1}{\sqrt{N}}\right)^{-4} N^2 \left(1 + \frac{1}{\sqrt{N}}\right)^3 \left(\frac{4}{9}N^{-5/2} + \left(\frac{4}{15} + \frac{1}{2}\right)N^{-8/3}\right) \\ &< \left(1 + \frac{1}{\sqrt{N}}\right)^8 \left(\frac{4}{9}N^{-1/2} + \frac{23}{30}N^{-2/3}\right), \end{aligned} \quad (4.10)$$

with the last inequality here holding by virtue of our having $\sqrt{N} \geq 5$, and so:

$$\left(1 + \frac{1}{\sqrt{N}}\right)^5 \left(1 - \frac{1}{\sqrt{N}}\right)^5 = \left(1 - \frac{1}{N}\right)^5 > 1 - \frac{5}{N} \geq 1 - \frac{1}{\sqrt{N}}.$$

By (4.1), (4.5) and (4.10), the lemma follows \blacksquare

Lemma 4.2 *Let N be an integer that is greater than 1. Then*

$$|\mathcal{U}(N)| < \left(2 + \frac{\log \log(N) + 9}{\log N}\right) N^{3/2} \sqrt{\log N}.$$

Proof. We recall the definition, in (4.2) (within our proof of Lemma 4.1), of the open square $\mathcal{Q}_{r,s}(N)$. By Definition 4.1, the number of elements (p, q) of $\mathcal{U}(N)$ is equal to the number of those open squares in the set

$$\mathcal{B}(N) = \{\mathcal{Q}_{r,s}(N) : r, s \in \mathbb{Z} \text{ and } 0 \leq r, s < N\}$$

that are crossed by one or more members of the sequence of hyperbolae

$$\mathcal{C}(m) = \left\{ (x, y) \in \mathbb{R} \times \mathbb{R} : xy = \frac{1}{m} \right\} \quad (m \in \mathbb{N}).$$

We therefore consider, for an arbitrary choice of $m \in \mathbb{N}$, the number

$$\nu(m, N) = |\{\mathcal{Q} \in \mathcal{B}(N) : \mathcal{Q} \cap \mathcal{C}(m) \neq \emptyset\}|.$$

Note firstly that the union of all the open squares in the set $\mathcal{B}(N)$ is a subset of the open square $(0, 1) \times (0, 1)$. Hence we can safely ignore all points (x, y) on the hyperbola $\mathcal{C}(m)$ that have $\max\{x, y\} > 2$ (say). That is, we have only to consider the finite part of this hyperbola that has endpoints $(\frac{1}{2m}, 2)$ and $(2, \frac{1}{2m})$. Let the point P travel at a fixed speed along the hyperbola $\mathcal{C}(m)$, starting at the coordinates $(\frac{1}{2m}, 2)$ and ending its journey at $(2, \frac{1}{2m})$. The number of open squares in the set $\mathcal{B}(N)$ that are crossed by P during its journey will then be equal to $\nu(m, N)$. We note that P starts from a point not lying in any of these open squares, and that in entering any one such square it must cross either the topmost or the leftmost of the four straight line segments forming the boundary of that square (for the slope of the hyperbola is negative at any given point on it). Consequently, given the definition of the open squares $\mathcal{Q}_{r,s}(N)$ that are the elements of $\mathcal{B}(N)$, and given that P moves only within the first quadrant of the x, y -plane (without touching or crossing the y -axis in particular), we find that $\nu(m, N) \leq T_m + L_m$, where $T_m = |\{(x, y) \in \mathcal{C}(m) : 0 < x < 1, 0 < y \leq 1 \text{ and } Ny \in \mathbb{Z}\}|$ and $L_m = |\{(x, y) \in \mathcal{C}(m) : 0 < x < 1, 0 < y < 1 \text{ and } Nx \in \mathbb{Z}\}|$. Given the definition of the hyperbola $\mathcal{C}(m)$, we deduce that $L_m \leq T_m = |\{y \in \mathbb{R} : \frac{1}{m} < y \leq 1 \text{ and } Ny \in \mathbb{Z}\}| = |(\frac{N}{m}, N] \cap \mathbb{Z}| = N - \lfloor \frac{N}{m} \rfloor$ (with the initial inequality following as a consequence of the fact that one has $(x, y) \in \mathcal{C}(m)$ if and only if $(y, x) \in \mathcal{C}(m)$). We conclude that one has:

$$\nu(m, N) \leq 2 \left(N - \left\lfloor \frac{N}{m} \right\rfloor \right) \quad (m \in \mathbb{N}). \quad (4.11)$$

We suppose now that M is some positive integer, and that $M \leq N$. By the preceding discussion we have

$$\begin{aligned} |\mathcal{U}(N)| &= \left| \bigcup_{m \in \mathbb{N}} \{ \mathcal{Q} \in \mathcal{B}(N) : \mathcal{Q} \cap \mathcal{C}(m) \neq \emptyset \} \right| \\ &\leq \left| \bigcup_{\substack{m \in \mathbb{N} \\ m > M}} \{ \mathcal{Q} \in \mathcal{B}(N) : \mathcal{Q} \cap \mathcal{C}(m) \neq \emptyset \} \right| + \sum_{m=1}^M \nu(m, N). \end{aligned} \quad (4.12)$$

When $m > M$ and r and s are integers satisfying $0 \leq r, s < N$ (so that the open square $\mathcal{Q}_{r,s}(N)$ is one of the elements of $\mathcal{B}(N)$), the set $\mathcal{Q}_{r,s}(N) \cap \mathcal{C}(m)$ is non-empty if and only if one has $(\frac{r}{N})(\frac{s}{N}) < \frac{1}{m} < (\frac{r+1}{N})(\frac{s+1}{N})$, and so is empty if one does not have $rs/N^2 < 1/M$. We therefore have

$$\begin{aligned} \left| \bigcup_{\substack{m \in \mathbb{N} \\ m > M}} \{ \mathcal{Q} \in \mathcal{B}(N) : \mathcal{Q} \cap \mathcal{C}(m) \neq \emptyset \} \right| &\leq \sum_{\substack{0 \leq r, s < N \\ rs < N^2/M}} 1 \\ &\leq (2N - 1) + \sum_{\substack{0 < r \leq s < N \\ rs < N^2/M}} 2 \\ &= (2N - 1) + \sum_{0 < r < N/\sqrt{M}} \sum_{r \leq s < \min\{N, N^2/(rM)\}} 2 \\ &\leq 2S_1 + 2S_2 + 2N - 1, \end{aligned}$$

where

$$\begin{aligned} S_1 &= \sum_{0 < r < N/M} \sum_{r \leq s < N} 1 \\ &\leq \sum_{0 < r \leq N/M} (N - r) = \left\lfloor \frac{N}{M} \right\rfloor \left(N - \frac{\lfloor \frac{N}{M} \rfloor + 1}{2} \right) < \frac{N^2}{M} \left(1 - \frac{1}{2M} \right) \end{aligned}$$

and

$$\begin{aligned} S_2 &= \sum_{N/M \leq r < N/\sqrt{M}} \sum_{r \leq s < N^2/(rM)} 1 \\ &< \sum_{N/M \leq r < N/\sqrt{M}} \left(\frac{N^2}{Mr} - (r - 1) \right) \\ &< \int_{N/M}^{N/\sqrt{M}} \left(\frac{N^2}{Mx} - (x - 1) \right) dx + \frac{N^2}{MN/M} - (N/M - 1) \\ &= \frac{N^2 \log M}{2M} - \left(\frac{1}{2} \right) \left(\frac{N}{\sqrt{M}} + \frac{N}{M} - 2 \right) \left(\frac{N}{\sqrt{M}} - \frac{N}{M} \right) + N + 1 - \frac{N}{M} \\ &= \left(\frac{N^2}{2M} \right) \left(\log M + \frac{1}{M} - 1 \right) + N \left(1 + \frac{1}{\sqrt{M}} - \frac{2}{M} \right) + 1, \end{aligned}$$

and so may conclude that

$$\left| \bigcup_{\substack{m \in \mathbb{N} \\ m > M}} \{ \mathcal{Q} \in \mathcal{B}(N) : \mathcal{Q} \cap \mathcal{C}(m) \neq \emptyset \} \right| < \left(\frac{N^2}{M} \right) (1 + \log M) + N \left(4 + \frac{2}{\sqrt{M}} - \frac{4}{M} \right) + 1. \quad (4.13)$$

At the same time, it follows by (4.11) that we have

$$\begin{aligned} \frac{1}{2} \sum_{m=1}^M \nu(m, N) &\leq \sum_{m=1}^M \left(N - \left\lfloor \frac{N}{m} \right\rfloor \right) \\ &\leq (-1) + \sum_{m=1}^M \left(N + 1 - \frac{N}{m} \right) \leq (N + 1)M - N \left(\frac{1}{M} + \log M \right) - 1. \end{aligned}$$

By this, together with (4.12) and (4.13), we obtain the bound

$$\begin{aligned} |\mathcal{U}(N)| &< \left(\frac{N^2}{M} \right) (1 + \log M) + N \left(2M - 2 \log(M) + 4 + \frac{2}{\sqrt{M}} - \frac{6}{M} \right) + 2M \\ &\leq \left(\frac{N^2}{M} \right) (1 + \log M) + 2NM + 8N \end{aligned} \quad (4.14)$$

(with the latter inequality following since we assume that $1 \leq M \leq N$).

In order to extract from this last bound for $|\mathcal{U}(N)|$ a result that is close to optimal, we specify now that

$$M = \left\lceil \left(\frac{1}{2}\right) \sqrt{N \log N} \right\rceil . \quad (4.15)$$

Note that, since $N > 1$, we have here $N \log N > 0$, and so (4.15) makes M a positive integer (as we have been assuming all along). We have also $(1/2)\sqrt{N \log N} \leq (1/2)\sqrt{N(N-1)} < N/2 \leq N-1 \in \mathbb{Z}$ (given that N is a positive integer greater than or equal to 2), and from this it follows that (4.15) makes $M \leq N-1$ (so that we do indeed have $M \leq N$, as was assumed in (4.12) and our subsequent working). We may therefore conclude that, with M as specified in (4.15), the bound (4.14) will hold. We may note also that, since $N \geq 2$, we have $N \log N > 1$, so that the choice (4.15) implies: $(\frac{1}{2})\sqrt{N \log N} \leq M < (\frac{1}{2})\sqrt{N \log N} + 1 < (\frac{3}{2})\sqrt{N \log N}$. Hence, by (4.14) and (4.15), we obtain

$$\begin{aligned} |\mathcal{U}(N)| &< \frac{(\log((\frac{3}{2})\sqrt{N \log N}) + 1) N^2}{(\frac{1}{2})\sqrt{N \log N}} + (\sqrt{N \log N} + 10) N \\ &= \left(2 + \frac{\log \log(N) + 2 + 2 \log(\frac{3}{2})}{\log N} + \frac{10}{\sqrt{N \log N}} \right) N^{3/2} \sqrt{\log N} . \end{aligned}$$

The lemma follows from this, since we have here $(N \log N)/(\log N)^2 = N/\log N \geq 3/\log 3$ and $2 + 2 \log(3/2) + 10/\sqrt{3}/\log 3 < 9$. ■

The result of Theorem 1 follows as an immediate consequence of those of Lemmas 4.1 and 4.2, and so Theorem 1 has now been proved. ■

§5. Proof of Theorem 3.

Let $N \in \mathbb{N}$ be given. For $0 \leq x \leq 1$, we put

$$\theta(x) = \begin{cases} 0 & \text{if } x = 0, \\ \mu(\lceil Nx \rceil) & \text{if } 0 < x \leq 1, \end{cases}$$

where $\mu(n)$ is the Möbius function. The function θ so defined is an element of the space $L^2([0, 1])$ with norm $\|\theta\|$ satisfying:

$$\|\theta\|^2 = \int_0^1 |\theta(x)|^2 dx \leq \int_0^1 1^2 dx = 1 .$$

Taking the kernel $K^\dagger(x, y)$ to be as defined in Section 1, we find that the quantity

$$J^\dagger = J^\dagger(\theta, \theta) = \int_0^1 \int_0^1 K^\dagger(x, y) \theta(x) \theta(y) dx dy$$

satisfies:

$$\begin{aligned} J^\dagger &= \sum_{m=1}^N \sum_{n=1}^N \int_{(m-1)/N}^{m/N} \left(\int_{(n-1)/N}^{n/N} K \left(\frac{m}{N}, \frac{n}{N} \right) \mu(n) dy \right) \mu(m) dx \\ &= \frac{1}{N^2} \sum_{m=1}^N \sum_{n=1}^N K \left(\frac{m}{N}, \frac{n}{N} \right) \mu(n) \mu(m) , \end{aligned}$$

so that $N^2 J^\dagger(\theta, \theta)$ is equal to the sum over m and n that occurs on the right-hand side of the identity (1.1). We therefore may rewrite (1.1) as:

$$M(N^2) = \left(\frac{1}{2}\right) (M(N))^2 + 2M(N) - N^2 \left(\sum_{n=1}^N \frac{\mu(n)}{n} \right)^2 - N^2 J^\dagger(\theta, \theta). \quad (5.1)$$

If we now put also

$$J = J(\theta, \theta) = \int_0^1 \int_0^1 K(x, y) \theta(x) \theta(y) dx dy,$$

then we have

$$J - J^\dagger = \int_0^1 \int_0^1 K^\ddagger(x, y) \theta(x) \theta(y) dx dy, \quad (5.2)$$

where $K^\ddagger(x, y) = K(x, y) - K^\dagger(x, y)$ (as was the case in Sections 3 and 4).

Lemma 5.1. *We have*

$$J - J^\dagger = O\left(\sqrt{\frac{1 + \log N}{N}}\right).$$

Proof. By our definition of $\theta(x)$ we have $|\theta(x)| \leq 1$ for $0 \leq x \leq 1$, and so (5.2) implies that

$$\begin{aligned} |J - J^\dagger| &\leq \int_0^1 \int_0^1 |K^\ddagger(x, y)| dx dy \\ &= \sum_{r=0}^{N-1} \sum_{s=0}^{N-1} \int_{\frac{r}{N}}^{\frac{r+1}{N}} \left(\int_{\frac{s}{N}}^{\frac{s+1}{N}} |K^\ddagger(x, y)| dy \right) dx \\ &= \sum_{r=0}^{N-1} \sum_{s=0}^{N-1} L_{r,s}(N) \quad (\text{say}). \end{aligned} \quad (5.3)$$

We note that the term $L_{r,s}(N)$ in (5.3) is similar to the integral $I_{r,s}(N)$ that occurs in the proof of Lemma 4.1 (though the integrand of $I_{r,s}(N)$ is the square of that of $L_{r,s}(N)$). We are therefore able to make use of some facts already noted in our proof of Lemma 4.1.

By (4.4) and the observations in-between (4.5) and (4.7), we find that when r and s are integers satisfying both $0 \leq r, s < N$ and $(r, s) \notin \mathcal{U}(N)$ (with $\mathcal{U}(N)$ being the set of points given by Definition 4.1) one will then have:

$$\begin{aligned} L_{r,s}(N) &= \int_{\frac{r}{N}}^{\frac{r+1}{N}} \left(\int_{\frac{s}{N}}^{\frac{s+1}{N}} \left(\frac{1}{xy} - \frac{N^2}{(r+1)(s+1)} \right) dy \right) dx \\ &= \log\left(\frac{r+1}{r}\right) \log\left(\frac{s+1}{s}\right) - \frac{1}{(r+1)(s+1)} \end{aligned}$$

(note that one cannot have $rs = 0$ here, else one would have $(r, s) \in \mathcal{U}(N)$ - contrary to the conditions we are assuming here). For $t > 0$, one has

$$\begin{aligned} \log\left(\frac{t+1}{t}\right) &= -\log\left(1 - \frac{1}{t+1}\right) = \sum_{n=1}^{\infty} \frac{1}{n(t+1)^n} < \left(\frac{1}{t+1}\right) \left(1 + \sum_{m=1}^{\infty} \frac{1}{2(t+1)^m}\right) \\ &= \left(\frac{1}{t+1}\right) \left(1 + \frac{1}{2t}\right), \end{aligned}$$

and so we are able to deduce that, subject to the above mentioned conditions on r and s , one has:

$$\begin{aligned} 0 \leq L_{r,s}(N) &< \left(\frac{1}{r+1}\right) \left(\frac{1}{s+1}\right) \left(\left(1 + \frac{1}{2r}\right) \left(1 + \frac{1}{2s}\right) - 1\right) \\ &\leq \frac{(5/2)}{(\min\{r+1, s+1\})^2 \max\{r+1, s+1\}}. \end{aligned}$$

In the remaining cases, where (r, s) is an element of the set $\mathcal{U}(N)$, we simply note that, similarly to (4.5), one has:

$$0 \leq L_{r,s}(N) \leq \int_{\frac{r}{N}}^{\frac{r+1}{N}} \left(\int_{\frac{s}{N}}^{\frac{s+1}{N}} 1 dy \right) dx = \frac{1}{N^2}.$$

By (5.3) and our observations following it, combined with (4.8) (which applies when $(r, s) \notin \mathcal{U}(N)$), we find that

$$\begin{aligned} |J - J^\dagger| &\leq \frac{|\mathcal{U}(N)|}{N^2} + \sum_{\substack{1 \leq m \leq n \leq N \\ m > N/\sqrt{n}}} \frac{5}{m^2 n} \\ &= \frac{|\mathcal{U}(N)|}{N^2} + \sum_{N^{2/3} < n \leq N} \left(\frac{1}{n}\right) \cdot O\left(\frac{\sqrt{n}}{N}\right) \\ &= \frac{|\mathcal{U}(N)|}{N^2} + O\left(\frac{1}{\sqrt{N}}\right). \end{aligned}$$

To complete the proof we need only appeal now to Lemma 4.2 (and, in the case $N = 1$, to our Remark immediately below Definition 4.1). ■

In view of the above lemma, we may modify the equation (5.1) by substituting $-N^2 J(\theta, \theta) + O(N^{3/2} \sqrt{1 + \log N})$ in place of the term $-N^2 J^\dagger(\theta, \theta)$ that occurs there. The next step is to note that, by Lemma 2.3, one has:

$$J(\theta, \theta) = \sum_{j=1}^{\infty} \left(\langle \theta, \phi_{j+} \rangle^2 \nu_{j+} + \langle \theta, \phi_{j-} \rangle^2 \nu_{j-} \right). \quad (5.4)$$

Suppose now that δ is some positive number. By Bessel's inequality,

$$\left| \sum_{\substack{j \in \mathbb{N} \\ \nu_{j+} \leq \delta}} \langle \theta, \phi_{j+} \rangle^2 \nu_{j+} + \sum_{\substack{j \in \mathbb{N} \\ |\nu_{j-}| \leq \delta}} \langle \theta, \phi_{j-} \rangle^2 \nu_{j-} \right| \leq \delta \sum_{j=1}^{\infty} \left(\langle \theta, \phi_{j+} \rangle^2 + \langle \theta, \phi_{j-} \rangle^2 \right) \\ \leq \delta \|\theta\|^2 \leq \delta. \quad (5.5)$$

We have also the following result.

Lemma 5.2. *For $j \in \mathbb{N}$, one has both*

$$\langle \theta, \phi_{j+} \rangle^2 \nu_{j+} = \left(\frac{\nu_{j+}}{N} \right) S_{j+}^2 + 2 \langle \theta, \phi_{j+} \rangle E_{j+} - E_{j+}^2 \nu_{j+}^{-1} \quad (5.6)$$

and

$$\langle \theta, \phi_{j-} \rangle^2 \nu_{j-} = \left(\frac{\nu_{j-}}{N} \right) S_{j-}^2 + 2 \langle \theta, \phi_{j-} \rangle E_{j-} - E_{j-}^2 \nu_{j-}^{-1}, \quad (5.7)$$

where, for either (consistent) choice of sign, one has

$$S_{j\pm} = \frac{1}{\sqrt{N}} \sum_{n=1}^N \mu(n) \phi_{j\pm} \left(\frac{n}{N} \right)$$

and

$$E_{j\pm} = \int_0^1 \int_0^1 K^{\%}(x, y) \theta(x) \phi_{j\pm}(y) dx dy,$$

with $K^{\%}$ being the (non-symmetric) kernel that is given by:

$$K^{\%}(x, y) = K(x, y) - K \left(\frac{[Nx]}{N}, y \right) \quad (0 \leq x, y \leq 1).$$

Proof. It suffices to show (5.6): one obtains (5.7) similarly. We have

$$\langle \theta, \phi_{j+} \rangle^2 \nu_{j+} = \langle \theta, \nu_{j+} \phi_{j+} \rangle^2 \nu_{j+}^{-1} \quad (5.8)$$

and

$$\begin{aligned} \langle \theta, \nu_{j+} \phi_{j+} \rangle &= \int_0^1 \theta(x) (\nu_{j+} \phi_{j+}(x)) dx \\ &= \sum_{n=1}^N \mu(n) \int_{\frac{n-1}{N}}^{\frac{n}{N}} \nu_{j+} \phi_{j+}(x) dx \\ &= \sum_{n=1}^N \mu(n) \left(\frac{\nu_{j+}}{N} \right) \phi_{j+} \left(\frac{n}{N} \right) + \sum_{n=1}^N \mu(n) \int_{\frac{n-1}{N}}^{\frac{n}{N}} \left(\nu_{j+} \phi_{j+}(x) - \nu_{j+} \phi_{j+} \left(\frac{n}{N} \right) \right) dx \\ &= \left(\frac{\nu_{j+}}{\sqrt{N}} \right) S_{j+} + \sum_{n=1}^N \mu(n) \int_{\frac{n-1}{N}}^{\frac{n}{N}} \left(\int_0^1 \left(K(x, y) - K \left(\frac{n}{N}, y \right) \right) \phi_{j+}(y) dy \right) dx \\ &= \left(\frac{\nu_{j+}}{\sqrt{N}} \right) S_{j+} + E_{j+}. \end{aligned}$$

By the last five equalities, one has

$$\left(\frac{\nu_{j_+}}{\sqrt{N}}\right) S_{j_+} = \langle \theta, \nu_{j_+} \phi_{j_+} \rangle - E_{j_+} ,$$

and so

$$\left(\frac{\nu_{j_+}^2}{N}\right) S_{j_+}^2 = \langle \theta, \nu_{j_+} \phi_{j_+} \rangle^2 - 2 \langle \theta, \nu_{j_+} \phi_{j_+} \rangle E_{j_+} + E_{j_+}^2 ,$$

which implies:

$$\langle \theta, \nu_{j_+} \phi_{j_+} \rangle^2 = \left(\frac{\nu_{j_+}^2}{N}\right) S_{j_+}^2 + 2\nu_{j_+} \langle \theta, \phi_{j_+} \rangle E_{j_+} - E_{j_+}^2 .$$

By this last equality and (5.8) we get (5.6), as required. ■

By (5.4), (5.5) and the results of Lemma 5.2, we find that

$$J(\theta, \theta) = F_+ + 2G_+ - H_+ + F_- + 2G_- - H_- + O(\delta) , \quad (5.9)$$

where (for each consistent choice of sign) one has:

$$F_{\pm} = \frac{1}{N} \sum_{\substack{j \in \mathbb{N} \\ |\nu_{j_{\pm}}| > \delta}} \nu_{j_{\pm}} S_{j_{\pm}}^2 , \quad G_{\pm} = \sum_{\substack{j \in \mathbb{N} \\ |\nu_{j_{\pm}}| > \delta}} \langle \theta, \phi_{j_{\pm}} \rangle E_{j_{\pm}} , \quad H_{\pm} = \sum_{\substack{j \in \mathbb{N} \\ |\nu_{j_{\pm}}| > \delta}} \frac{E_{j_{\pm}}^2}{\nu_{j_{\pm}}} .$$

Lemma 5.3. *For either choice of sign, we have*

$$\max \left\{ |G_{\pm}|^2 , \delta |H_{\pm}| \right\} \leq D ,$$

where

$$D = \int_0^1 \left(\int_0^1 |K^{\%}(x, y)| dx \right)^2 dy \quad (5.10)$$

($K^{\%}$ here being the non-symmetric kernel that is defined in the statement of Lemma 5.2).

Proof. Referring back to the definitions made in the statement of Lemma 5.2, we find that, for either choice of sign,

$$\begin{aligned} G_{\pm} &= \sum_{\substack{j \in \mathbb{N} \\ |\nu_{j_{\pm}}| > \delta}} \langle \theta, \phi_{j_{\pm}} \rangle \int_0^1 \int_0^1 K^{\%}(x, y) \theta(x) \phi_{j_{\pm}}(y) dx dy \\ &= \int_0^1 \int_0^1 K^{\%}(x, y) \theta(x) \tilde{\theta}_{\pm}(y) dx dy , \end{aligned}$$

where

$$\tilde{\theta}_{\pm}(y) = \sum_{\substack{j \in \mathbb{N} \\ |\nu_{j\pm}| > \delta}} \langle \theta, \phi_{j\pm} \rangle \phi_{j\pm}(y)$$

(we need not consider the question of convergence here, since we know by (3.1), for example, that the relevant summation is finite). Since $|\theta(x)| \leq 1$ for $0 \leq x \leq 1$, we may deduce (with the help of the Cauchy-Schwarz inequality) that one has

$$|G_{\pm}|^2 \leq \left\langle \kappa, |\tilde{\theta}_{\pm}| \right\rangle^2 \leq \|\kappa\|^2 \|\tilde{\theta}_{\pm}\|^2, \quad (5.11)$$

where

$$\kappa(y) = \int_0^1 |K^{\%}(x, y)| dx \quad (0 \leq y \leq 1),$$

so that, by (5.10),

$$\|\kappa\|^2 = D. \quad (5.12)$$

In connection with the preceding, we note that the assumed orthonormality of the relevant set of eigenfunctions implies that we must have here:

$$\|\tilde{\theta}_{\pm}\|^2 = \langle \tilde{\theta}_{\pm}, \tilde{\theta}_{\pm} \rangle = \sum_{\substack{j \in \mathbb{N} \\ |\nu_{j\pm}| > \delta}} \langle \theta, \phi_{j\pm} \rangle^2 \leq \|\theta\|^2$$

(the final inequality being an instance of Bessel's inequality). We found earlier that one has $\|\theta\| \leq 1$, and so these last observations, together with (5.11) and (5.12), yield the bounds $|G_{\pm}|^2 \leq D$. To complete our proof of the lemma we have now only to show that $|H_{\pm}| \leq D/\delta$.

According to the definitions of Lemma 5.2, and those below (5.9), we have

$$\begin{aligned} |H_{\pm}| &= \left| \sum_{\substack{j \in \mathbb{N} \\ |\nu_{j\pm}| > \delta}} \left(\int_0^1 \int_0^1 K^{\%}(x, y) \theta(x) \phi_{j\pm}(y) dx dy \right)^2 (\nu_{j\pm})^{-1} \right| \\ &\leq \frac{1}{\delta} \sum_{\substack{j \in \mathbb{N} \\ |\nu_{j\pm}| > \delta}} \left(\int_0^1 \int_0^1 K^{\%}(x, y) \theta(x) \phi_{j\pm}(y) dx dy \right)^2 \\ &= \frac{1}{\delta} \sum_{\substack{j \in \mathbb{N} \\ |\nu_{j\pm}| > \delta}} \langle \kappa^*, \phi_{j\pm} \rangle^2, \end{aligned}$$

where

$$\kappa^*(y) = \int_0^1 K^{\%}(x, y) \theta(x) dx \quad (0 \leq y \leq 1).$$

It follows from this, by way of Bessel's inequality, that one has

$$|H_{\pm}| \leq \delta^{-1} \|\kappa^*\|^2. \quad (5.13)$$

Since $|\theta(x)| \leq 1$ for $0 \leq x \leq 1$, we have

$$|\kappa^*(y)| \leq \int_0^1 |K^{\%}(x, y)| dx = \kappa(y) \quad \text{for } 0 \leq y \leq 1$$

(κ here denoting the same function that it does in (5.11)). We therefore find that one has $\|\kappa^*\| \leq \|\kappa\|$, and so, given (5.12) and (5.13), the required bound, $|H_{\pm}| \leq D/\delta$, follows. ■

Lemma 5.4. *The quantity D defined in (5.10) satisfies*

$$0 \leq D \leq \frac{49(1 + \log N)}{N}.$$

Proof. Let

$$\mathcal{V}(y) = \left\{ \left\lfloor \frac{N/y}{\ell} \right\rfloor : \ell \in \mathbb{N} \right\} \quad (0 < y \leq 1),$$

and let $K^{\%}(x, y)$ be the non-symmetric kernel defined in the statement of Lemma 5.2. One then has, for each non-negative integer $r < N$,

$$\int_{\frac{r}{N}}^{\frac{r+1}{N}} |K^{\%}(x, y)| dx \leq \begin{cases} \frac{1}{N} & \text{if } r \in \mathcal{V}(y), \\ \frac{1}{(r+1)ry} & \text{otherwise.} \end{cases} \quad (5.14)$$

Indeed, assuming that $r \in [0, N) \cap \mathbb{Z}$, and that $(x, y) \in (\frac{r}{N}, \frac{r+1}{N}) \times (0, 1)$, one has

$$\begin{aligned} |K^{\%}(x, y)| &= \left| K(x, y) - K\left(\frac{r+1}{N}, y\right) \right| \\ &= \left| \left\{ \frac{N}{(r+1)y} \right\} - \left\{ \frac{1}{xy} \right\} \right| \end{aligned} \quad (5.15)$$

$$\begin{aligned} &= \left| \left(\frac{N}{(r+1)y} - \left\lfloor \frac{N}{(r+1)y} \right\rfloor \right) - \left(\frac{1}{xy} - \left\lfloor \frac{1}{xy} \right\rfloor \right) \right| \\ &\leq \left| \frac{N}{(r+1)y} - \frac{1}{xy} \right| + \left| \left\lfloor \frac{1}{xy} \right\rfloor - \left\lfloor \frac{N}{(r+1)y} \right\rfloor \right| \\ &= \left(\frac{1}{x} - \frac{N}{r+1} \right) y^{-1} + \sum_{\frac{N}{(r+1)y} < \ell \leq \frac{1}{xy}} 1. \end{aligned} \quad (5.16)$$

Given an integer ℓ satisfying the conditions of summation in (5.16), one would have $\ell > 0$ and $r+1 > \frac{N/y}{\ell} \geq Nx > r$, and so $r = \lfloor \frac{N/y}{\ell} \rfloor \in \mathcal{V}(y)$. We therefore find that if $r \notin \mathcal{V}(y)$ then (5.16) leads to the bounds:

$$|K^{\%}(x, y)| \leq \left(\frac{1}{x} - \frac{N}{r+1} \right) y^{-1} \leq \left(\frac{N}{r} - \frac{N}{r+1} \right) y^{-1} = \frac{N}{(r+1)ry}.$$

In respect of the remaining cases, where r is an element of the set $\mathcal{V}(y)$, we simply observe that, by (5.15), one is certain to have $|K^\%(x, y)| < 1$. As we have

$$\int_{\frac{r}{N}}^{\frac{r+1}{N}} c dx = \frac{c}{N} \quad (c \in \mathbb{R} \text{ independent of } x),$$

the preceding two observations lead one immediately to the bound in (5.14).

Now clearly one has, for $0 < y < 1$,

$$\begin{aligned} \mathcal{V}(y) &= \left\{ \left\lfloor \frac{N/y}{\ell} \right\rfloor : \ell \in \mathbb{N} \text{ and } \ell < \sqrt{N/y} \right\} \cup \left\{ \left\lfloor \frac{N/y}{\ell} \right\rfloor : \ell \in \mathbb{N} \text{ and } \ell \geq \sqrt{N/y} \right\} \\ &\subseteq \left\{ \left\lfloor \frac{N/y}{\ell} \right\rfloor : \ell \in \mathbb{N} \text{ and } \ell < \sqrt{N/y} \right\} \cup \left\{ m \in \mathbb{Z} : 0 \leq m \leq \sqrt{N/y} \right\}, \end{aligned}$$

and so:

$$|\mathcal{V}(y)| \leq 1 + 2 \left\lfloor \sqrt{N/y} \right\rfloor \leq 3(N/y)^{1/2}. \quad (5.17)$$

One has also

$$\frac{N/y}{\ell} - \frac{N/y}{\ell+1} = \frac{N/y}{(\ell+1)\ell} < 1 \quad \text{when } \ell \geq \sqrt{N/y},$$

and so, for $0 < y < 1$, the set $\mathcal{V}(y)$ will contain all non-negative integers m that are less than or equal to $(N/y)/\lceil \sqrt{N/y} \rceil$. Noting that, for $N \geq 1 > y > 0$, one has

$$\frac{N/y}{\lceil \sqrt{N/y} \rceil} > \frac{N/y}{\sqrt{N/y} + 1} = \frac{\sqrt{N/y}}{1 + \sqrt{y/N}} > \left(1 - \sqrt{\frac{y}{N}}\right) \sqrt{\frac{N}{y}} = \sqrt{\frac{N}{y}} - 1 > 0,$$

we may conclude that

$$r \geq \max \left\{ \sqrt{\frac{N}{y}} - 1, 1 \right\} \geq \left(\frac{1}{2}\right) \sqrt{\frac{N}{y}} \quad \text{if } 0 < y < 1, r \in \mathbb{N} \cup \{0\} \text{ and } r \notin \mathcal{V}(y). \quad (5.18)$$

By (5.14), (5.17) and (5.18), we are able to conclude that, for $0 < y < 1$, one has:

$$\begin{aligned} \int_0^1 |K^\%(x, y)| dx &= \sum_{r=0}^{N-1} \int_{r/N}^{(r+1)/N} |K^\%(x, y)| dx \\ &\leq \sum_{\substack{0 \leq r < N \\ r \in \mathcal{V}(y)}} \frac{1}{N} + \sum_{\frac{1}{2}\sqrt{N/y} \leq r < N} \frac{1}{(r+1)ry} \\ &\leq N^{-1} \min \{N, |\mathcal{V}(y)|\} + \begin{cases} \frac{1}{\lceil \frac{1}{2}\sqrt{N/y} \rceil y} & \text{if } y > \frac{1}{4N}, \\ 0 & \text{otherwise,} \end{cases} \\ &\leq \min \left\{ 1, \frac{3}{\sqrt{Ny}} \right\} + \min \left\{ 4, \frac{2}{\sqrt{Ny}} \right\} \leq 7 \min \left\{ 1, \frac{1}{\sqrt{Ny}} \right\}. \end{aligned}$$

Using this we find, by (5.10), that one has

$$0 \leq D \leq \int_0^{1/N} 7^2 dy + \left(\frac{7}{\sqrt{N}} \right)^2 \int_{1/N}^1 \frac{dy}{y} = \frac{49(1 + \log N)}{N},$$

as the lemma states. ■

By (5.9) and Lemmas 5.3 and 5.4, we obtain:

$$\begin{aligned} J(\theta, \theta) &= O(\delta) + F_+ + F_- + O(\sqrt{D}) + O(\delta^{-1}D) \\ &= F_+ + F_- + O(\max\{\delta, \delta^{-1}D\}) = F_+ + F_- + O\left(\max\left\{\delta, \frac{1 + \log N}{\delta N}\right\}\right). \end{aligned}$$

By this and Lemma 5.1 it follows that

$$J^\dagger = J^\dagger(\theta, \theta) = F_+ + F_- + O(\delta), \quad (5.19)$$

provided that we have $\delta > \sqrt{(1 + \log N)/N}$ (which is the hypothesis (3.7) of Theorem 3).

Assuming that (3.7) holds (and given that $N \geq 1$), it is trivially the case that the absolute value of the term $2M(N)$ occurring in (5.1) will be less than $2\delta N^2$. We therefore obtain, by (5.1), (5.19) and the relevant definitions (for which, see below (5.7), and below (5.9)), the result that is stated in Theorem 3. ■

§6. Proof of Theorem 4.

We assume throughout this section that N is an integer satisfying $N \geq 3$, and that ε and δ satisfy both (1.8) and (1.9) (and so are both positive). We put $\Delta = e^\varepsilon > 1$, so that, by (1.9),

$$\Delta = \frac{1}{\delta}. \quad (6.1)$$

We shall also use the notations x_j ($j = 1, \dots, N+2$), $\mathcal{R}_{i,j}$ ($i, j \in \{1, \dots, N+1\}$) and $\mu_{i,j}$ ($i, j \in \mathbb{N}$ and $i+j \leq N+1$) that were defined in Section 1; we shall, in particular, take the kernel $K^\triangleleft(x, y) = K^{\triangleleft, N}(x, y)$ to be as defined in the paragraph containing (1.8)-(1.14).

Lemma 6.1. *Subject to the assumptions just above, one has:*

$$\sqrt{3} < e^{N\varepsilon} < \frac{1}{2\varepsilon}, \quad (6.2)$$

$$0 < \left(\frac{63}{256} \right) \left(\frac{\log N}{N} \right) < \varepsilon < \min \left\{ \frac{1}{4}, \frac{\log N}{N} \right\} \quad (6.3)$$

and

$$\frac{1}{1 + \varepsilon} < \frac{1 - \delta}{\varepsilon} < 1 < \frac{\Delta - 1}{\varepsilon} < \frac{1}{1 - \varepsilon}. \quad (6.4)$$

One has, in particular, $0 < \varepsilon < 1/4$,

$$\left(\frac{4}{5}\right)\varepsilon < 1 - \delta < \varepsilon < \Delta - 1 < \left(\frac{4}{3}\right)\varepsilon \quad \text{and} \quad \frac{3}{4} < \delta < 1 < \Delta < \frac{4}{3}. \quad (6.5)$$

Proof. By (1.8) $\sinh(\varepsilon) > 0$, and so $\varepsilon > 0$. The existence and uniqueness of ε (given N) follows from the facts that the left and right sides of (1.8) are continuous and strictly monotonic functions of ε (for $\varepsilon > 0$, that is); since $2\sinh(0) = 0 < e^{-0N} = 1$, while $2\sinh(1/4) > 2/4 > e^{-3/4} \geq e^{-N/4}$, the continuity of the two functions therefore guarantees that (1.8) holds for some $\varepsilon \in (0, 1/4)$; the strict (and opposite) monotonicities of the two functions ensure that there cannot be two distinct choices of $\varepsilon > 0$ that both satisfy (1.8), so we may conclude that ε (satisfying (1.8)) is a function of $N \geq 3$, and satisfies $0 < \varepsilon < 1/4$ (for all such N).

Since $\varepsilon > 0$, we have $\sinh(\varepsilon) = \varepsilon + \varepsilon^3/3! + \dots > \varepsilon$; by this and (1.8), we obtain the rightmost inequality of (6.2), which is equivalent to the relation:

$$N\varepsilon < \log\left(\frac{1}{2\varepsilon}\right). \quad (6.6)$$

For the other inequality in (6.2), we argue that, since $N > 2$ and $\varepsilon > 0$, the equation (1.8) implies that we have $e^{-N\varepsilon} = 2\sinh(\varepsilon) < 2\sinh(\varepsilon)\cosh(\varepsilon) = \sinh(2\varepsilon) < \sinh(N\varepsilon)$, and so $2e^{-N\varepsilon} < e^{N\varepsilon} - e^{-N\varepsilon}$, from which the desired result follows.

The fact that $\varepsilon > 0$ also ensures that we have $\sinh(\varepsilon)/\varepsilon = 1 + \varepsilon^2/3! + \dots < \exp(\varepsilon^2/6) = 1 + (\varepsilon^2/6)/1! + (\varepsilon^2/6)^2/2! + \dots$, and so it follows from (1.8) that $e^{-N\varepsilon} < 2\varepsilon \exp(\varepsilon^2/6)$. We therefore have $N\varepsilon > \log(1/(2\varepsilon)) - (\varepsilon^2/6)$, which (as one has $0 < \varepsilon < 1/4$, and consequently $\log(1/(2\varepsilon)) > \log 2 > 2/3$) implies that

$$N\varepsilon > \left(1 - \frac{\varepsilon^2}{4}\right) \log\left(\frac{1}{2\varepsilon}\right) > \left(\frac{63}{64}\right) \log\left(\frac{1}{2\varepsilon}\right) > \left(\frac{63}{64}\right) \log 2 > \frac{1}{2}. \quad (6.7)$$

It follows that we have $1/(2\varepsilon) < N$, and so may deduce from (6.6) that $\varepsilon < (\log N)/N$. Thus, the rightmost inequality of (6.3) holds. An immediate consequence is that one has $1/(2\varepsilon) > N/(2 \log N)$, and so

$$\log\left(\frac{1}{2\varepsilon}\right) > \left(1 - \frac{\log(2) + \log \log(N)}{\log N}\right) \log N. \quad (6.8)$$

By elementary calculus, the maximum value attained by the function $x^{-1} \log(2x)$ ($x > 0$) is the number $2/e < 3/4$, and so (as $\log N > 0$) it follows from (6.8) that we have $\log(1/(2\varepsilon)) > (\log N)/4$; by this and the first two inequalities of (6.7), we obtain the middle inequality of (6.3). With the leftmost inequality of (6.3) being a consequence of having $N \geq 3 > e$ (and so $\log N > 1 > 0$), our proof of (6.3) is complete.

The results (6.2) and (6.3) have been proved. Since it has been shown that $0 < \varepsilon < \frac{1}{4}$, each relation in (6.5) follows directly from what is asserted in (6.4). The proof of the lemma will therefore be complete once we have demonstrated that (6.4) holds.

For (6.4), we note firstly that, by (1.9), one has $1 - \delta = 1 - e^{-\varepsilon} = \int_{-\varepsilon}^0 e^y dy$, and so $(0 - (-\varepsilon))e^{-\varepsilon} < 1 - \delta < (0 - (-\varepsilon))e^0$. Thus we have $\varepsilon > 1 - \delta$ and $1 - \delta > e^{-\varepsilon}\varepsilon = \delta\varepsilon = (1 - (1 - \delta))\varepsilon$, which implies that $(1 + \varepsilon)(1 - \delta) > \varepsilon$: we are therefore able to conclude that the first two inequalities in (6.4) are satisfied. By similar reasoning, the number defined in (6.1) (that is, $\Delta = 1/\delta = e^\varepsilon$) must satisfy $\varepsilon < \Delta - 1 < \varepsilon e^\varepsilon = \Delta\varepsilon = (1 + (\Delta - 1))\varepsilon$. It follows that we have both $\Delta - 1 > \varepsilon$ and $(1 - \varepsilon)(\Delta - 1) < \varepsilon$, and so we find (given that $0 < \varepsilon < 1/4 < 1$) that the final two inequalities of (6.4) are satisfied: this completes the proof of (6.4) and, with that, the proof of the lemma \blacksquare

Lemma 6.1 is useful for simplifying some of the estimates that we shall obtain, later in this section. We take a first step towards the proof of Theorem 4 by noting that

$$\begin{aligned} \|K^{\boxtimes}\|^2 &= \|K - K^\triangleleft\|^2 = \int_0^1 \int_0^1 (K(x, y) - K^\triangleleft(x, y))^2 dx dy \\ &= \sum_{i=1}^{N+1} \sum_{j=1}^{N+1} \iint_{\mathcal{R}_{i,j}} (K(x, y) - K^\triangleleft(x, y))^2 dx dy = V_0 + V_1, \end{aligned} \quad (6.9)$$

where, with $I_{i,j}(\kappa)$ defined, for $\kappa \in \mathbb{R}$ and $i, j \in \{1, \dots, N+1\}$, by

$$I_{i,j}(\kappa) := \iint_{\mathcal{R}_{i,j}} (K(x, y) - \kappa)^2 dx dy = \int_{x_{i+1}}^{x_i} \left(\int_{x_{j+1}}^{x_j} (K(x, y) - \kappa)^2 dy \right) dx, \quad (6.10)$$

we have:

$$V_0 = \sum_{\substack{i,j \geq 1 \\ i+j \leq N+1}} I_{i,j}(\mu_{i,j}) \quad \text{and} \quad V_1 = \sum_{\substack{i,j \leq N+1 \\ i+j > N+1}} I_{i,j}(0). \quad (6.11)$$

The next two lemmas take care of the estimation of the sum V_1 . Before stating them, we remark that, by its definition in (1.2), the kernel K is such that

$$K(x, y) = -\tilde{B}_1\left(\frac{1}{xy}\right) \quad (0 < x, y \leq 1),$$

where $\tilde{B}_1(t) := \{t\} - \frac{1}{2}$ is the ‘periodic Bernoulli function’ associated with the Bernoulli polynomial of degree 1. For later reference, we recall here (from [OLBC, Section24.2]) that

$$\tilde{B}_n(t) = B_n(\{t\}) \quad (n \in \mathbb{N} \text{ and } t \in \mathbb{R}),$$

and that the Bernoulli polynomials of degrees 1 and 4 are:

$$B_1(X) = X - \frac{1}{2} \quad \text{and} \quad B_4(X) = X^4 - 2X^3 + X^2 - \frac{1}{30}.$$

Lemma 6.2. *Let $s \geq 0$. Then, for some $\theta = \theta(s)$ satisfying $-1 \leq 36\sqrt{3}\theta \leq 1$, one has:*

$$\int_0^s \left(\tilde{B}_1(u)\right)^2 du = \frac{s}{12} + \theta.$$

Proof. Let $n = \lfloor s \rfloor$ and $\sigma = \{s\}$, so that $n \in \mathbb{Z}$, $n \geq 0$, $0 \leq \sigma < 1$ and $s = n + \sigma$. Then, by the periodicity of $\tilde{B}_1(u)$,

$$\int_0^s \left(\tilde{B}_1(u)\right)^2 du = n \int_0^1 \left(\tilde{B}_1(u)\right)^2 du + \int_0^\sigma \left(\tilde{B}_1(u)\right)^2 du .$$

For $0 \leq \tau \leq 1$, we have

$$\int_0^\tau \left(\tilde{B}_1(u)\right)^2 du = \int_0^\tau \left(u - \frac{1}{2}\right)^2 du = \frac{\left(\tau - \frac{1}{2}\right)^3 + \frac{1}{8}}{3} .$$

By this and the previous equation, one has

$$\int_0^s \left(\tilde{B}_1(u)\right)^2 du = \frac{1}{12}(s - \sigma) + \frac{\left(\sigma - \frac{1}{2}\right)^3 + \frac{1}{8}}{3} = \frac{s}{12} + \frac{P(\sigma)}{3} ,$$

where $P(\tau) = \left(\tau - \frac{1}{2}\right)^3 - \frac{1}{4}\tau + \frac{1}{8}$. One has $P'(\tau) = 0$ only if $\left(\tau - \frac{1}{2}\right)^2 = \frac{1}{12}$, and in this case $P(\tau) = \left(\frac{1}{2} - \tau\right)/6 = \pm \frac{1}{12\sqrt{3}}$; since we have also $P(0) = P(1) = 0$, it therefore follows (by elementary principles of calculus) that the maximum and minimum values attained by $P(\tau)$ on the interval $0 \leq \tau \leq 1$ are $\frac{1}{12\sqrt{3}}$ and $-\frac{1}{12\sqrt{3}}$, respectively, and so we obtain what the lemma states (with $\theta(s) = \frac{1}{3}P(\{s\}) \in \left[\frac{-1}{36\sqrt{3}}, \frac{1}{36\sqrt{3}}\right]$) ■

Lemma 6.3. The sum V_1 given by (6.11), (6.10) and (1.2) satisfies

$$V_1 \leq \left(\frac{N\varepsilon + 1}{12}\right) \Delta^{-N} + \left(\frac{N\varepsilon}{18\sqrt{3}}\right) \Delta^{-2N} .$$

Proof. By (1.11) and (1.12), the sets $\mathcal{R}_{i,j}$ ($i, j \in \{1, \dots, N+1\}$) are pairwise disjoint open subsets of the square $[0, 1] \times [0, 1]$. It therefore follows from (6.10), (6.11), (1.10) and (1.12) that

$$V_1 = \iint_{\mathcal{T}} (K(x, y))^2 dx dy ,$$

where

$$\mathcal{T} = \bigcup_{i=1}^{N+1} \bigcup_{j=N+2-i}^{N+1} \mathcal{R}_{i,j} \subset \bigcup_{i=1}^{N+1} \bigcup_{j=N+2-i}^{N+1} (0, \delta^{i-1}) \times (0, \delta^{j-1}) \subset (0, 1) \times (0, 1) .$$

We have here $(x, y) \in \mathcal{T}$ only if $(x, y) \in (0, 1) \times (0, 1)$ and there exists some pair of positive integers i, j such that $i + j \geq N + 2$ and $\delta^{i+j-2} = \delta^{i-1}\delta^{j-1} > xy$. It follows (given that $0 < \delta < 1$, as was shown in Lemma 6.1) that each point $(x, y) \in \mathcal{T}$ must satisfy

$xy < \delta^{N+2-2} = \delta^N$ (as well as $0 < x, y < 1$). This, together with the remarks preceding Lemma 6.2, leads us to the conclusion that one has:

$$V_1 \leq \int_0^1 \int_0^{\min\{1, \delta^N/x\}} \left(\tilde{B}_1\left(\frac{1}{xy}\right) \right)^2 dy dx .$$

By making the change of variables $1/(xy) = s$, $y/x = t$ and noting that $\delta^N = \Delta^{-N}$, we find that this last upper bound on V_1 becomes:

$$V_1 \leq \int_{\Delta^N}^{\infty} \int_{1/s}^s \left(\tilde{B}_1(s) \right)^2 (2t)^{-1} dt s^{-2} ds = \int_{\Delta^N}^{\infty} \left(\tilde{B}_1(s) \right)^2 s^{-2} \log(s) ds . \quad (6.12)$$

By Lemma 6.2, we find that the application of integration by parts gives:

$$\int_{\Delta^N}^{\infty} \left(\left(\tilde{B}_1(s) \right)^2 - \frac{1}{12} \right) s^{-2} \log(s) ds = - \left(\frac{\log(\Delta^N)}{\Delta^{2N}} \right) \theta(\Delta^N) - \int_{\Delta^N}^{\infty} \left(\frac{1 - 2 \log s}{s^3} \right) \theta(s) ds ,$$

where $\theta(s)$ is the (periodic) function of s that appears in Lemma 6.2. Therefore it follows by Lemma 6.2 (again), together with the first inequality of (6.2) (which implies that $2 \log s > \log 3 > 1$ if $s > \Delta^N$), that we have:

$$\int_{\Delta^N}^{\infty} \left(\left(\tilde{B}_1(s) \right)^2 - \frac{1}{12} \right) s^{-2} \log(s) ds \leq \frac{\log(\Delta^N)}{18\sqrt{3}\Delta^{2N}} .$$

By this, combined with (6.12) and the observations that

$$\int_{\Delta^N}^{\infty} s^{-2} \log(s) ds = \frac{1 + \log(\Delta^N)}{\Delta^N} \quad \text{and} \quad \log(\Delta^N) = N \log \Delta = N\varepsilon ,$$

we obtain (after a very short calculation) just what the lemma states \blacksquare

We shall next consider the terms $I_{i,j}(\mu_{i,j})$ occurring in the sum V_0 , in (6.11): we assume (accordingly) that i and j are positive integers satisfying the condition

$$i + j \leq N + 1 . \quad (6.13)$$

We therefore have $i, j \in \{1, \dots, N\}$, and so (by (1.10)-(1.12)) have also: $\mathcal{R}_{i,j} = (\delta^i, \delta^{i-1}) \times (\delta^j, \delta^{j-1}) \subset (0, 1) \times (0, 1)$. It is a special property of the mean value $\mu_{i,j}$ (defined in (1.13)) that, with $I_{i,j}(\kappa)$ defined by (6.10), one has:

$$I_{i,j}(\kappa) = I_{i,j}(\mu_{i,j}) + (\mu_{i,j} - \kappa)^2 \iint_{\mathcal{R}_{i,j}} dx dy \quad (\kappa \in \mathbb{R}). \quad (6.14)$$

We omit the (very straightforward) proof of this, but note that (6.14) is an instance of the ‘parallel axis theorem’: see [Li, Section 2.3.1] for a relevant discussion concerning the

variance of a random variable. As $\mu_{i,j}$ is a real number, while the double integral appearing in (6.14) is positive (being, in fact, equal to $(1 - \delta)^2 \delta^{i+j-2}$), it is therefore an immediate corollary of (6.14) that one has

$$I_{i,j}(\mu_{i,j}) \leq I_{i,j}(\kappa) \quad (\kappa \in \mathbb{R}), \quad (6.15)$$

so that $I_{i,j}(\mu_{i,j})$ is the minimum of the function $I_{i,j}(\kappa)$.

Putting $\kappa = 0$ in (6.14), and then recalling (6.10), we obtain the identity

$$I_{i,j}(\mu_{i,j}) = \iint_{\mathcal{R}_{i,j}} (K(x,y))^2 dx dy - \mu_{i,j}^2 \iint_{\mathcal{R}_{i,j}} dx dy, \quad (6.16)$$

which will appear somewhat familiar to statisticians: see [Li, Page 118], for example. Note that $\mu_{i,j}$ and $I_{i,j}(\mu_{i,j})$ are completely determined by the values of the three double integrals that occur in the pair of equations (1.13), (6.16). By (6.14), those three values also determine the real function $\kappa \mapsto I_{i,j}(\kappa)$. It is therefore significant that, if p is a non-negative integer, then, given the definition (1.2), it follows by the change of variables $x = \delta^{i-1}u$, $y = \delta^{j-1}v$, that one has

$$\iint_{\mathcal{R}_{i,j}} (K(x,y))^p dx dy = \delta^{n-1} \int_{\delta}^1 \int_{\delta}^1 (K(\delta^{n-1}uv, 1))^p dudv,$$

where

$$n = i + j - 1 \in \{1, \dots, N\}. \quad (6.17)$$

Note that the last double integral above depends only on δ (itself determined by N), p and n . Bearing in mind what was deduced from (6.16), we conclude that, since the above holds for $p = 0, 1, 2$ (in particular), the real function $\kappa \mapsto I_{i,j}(\kappa)$, and numbers $I_{i,j}(\mu_{i,j})$ and $\mu_{i,j}$, must each be completely determined by just N and the integer n in (6.17). We therefore have (for n as defined in (6.17)):

$$\mu_{i,j} = \tilde{\mu}_n := \mu_{n,1} \in \mathbb{R}, \quad (6.18)$$

$$I_{i,j}(\kappa) = \tilde{I}_n(\kappa) := I_{n,1}(\kappa) \quad (\kappa \in \mathbb{R}) \quad (6.19)$$

and (consequently) $I_{i,j}(\mu_{i,j}) = \tilde{I}_n(\tilde{\mu}_n)$. By (6.15), (6.18) and (6.19), it follows that

$$\tilde{I}_n(\tilde{\mu}_n) \leq \tilde{I}_n(\kappa) \quad (\kappa \in \mathbb{R} \text{ and } n \in \{1, \dots, N\}). \quad (6.20)$$

Remark. By (1.18), (1.10) and (6.17)-(6.19), the elements h_{ij} of the $N \times N$ matrix $H(N)$ satisfy $h_{ij} = \eta_{i+j-1}$ ($i, j \in \{1, \dots, N\}$), where, for $n = 1, \dots, 2N - 1$, one has:

$$\eta_n = \begin{cases} (\delta^{\frac{n-1}{2}} - \delta^{\frac{n+1}{2}}) \tilde{\mu}_n & \text{if } n \leq N, \\ 0 & \text{otherwise.} \end{cases}$$

Thus $H(N)$ is indeed a Hankel matrix, as was mentioned in our Introduction.

In the preceding we have assumed only that i and j are integers satisfying (6.13) and $i, j \geq 1$, which are precisely the conditions of summation for the sum V_0 , defined in (6.11). We therefore deduce from (6.11), (6.17), (6.18) and (6.19) that one has

$$V_0 = \sum_{1 \leq i \leq n \leq N} \sum_{n=1}^N \tilde{I}_n(\tilde{\mu}_n) = \sum_{n=1}^N n \cdot \tilde{I}_n(\tilde{\mu}_n). \quad (6.21)$$

In order to estimate V_0 we require upper bounds for the numbers $\tilde{I}_1(\tilde{\mu}_1), \dots, \tilde{I}_N(\tilde{\mu}_N)$. The next lemma is a preliminary step towards obtaining such bounds. Following it we have another lemma (really a corollary) stating a formula that, for $N \in \mathbb{N}$, facilitates the numerical evaluation of $\tilde{\mu}_1, \dots, \tilde{\mu}_N$. In relation to our proof of Theorem 4, the latter lemma (i.e. Lemma 6.5) is merely incidental: see the Remark following its proof for some observations justifying its inclusion in this paper.

Lemma 6.4. *Let $n \leq N$ be a positive integer. Then, for $\kappa \in \mathbb{R}$, one has*

$$\begin{aligned} \tilde{I}_n(\kappa) = \varepsilon^2 e^{-n\varepsilon} & \left(\int_{-1}^{\sigma_n} \left(\kappa + e^{(\sigma+n)\varepsilon} - \left(k_n - \frac{1}{2}\right) \right)^2 (1 - |\sigma|) e^{-\sigma\varepsilon} d\sigma \right. \\ & \left. + \int_{\sigma_n}^1 \left(\kappa + e^{(\sigma+n)\varepsilon} - \left(k_n + \frac{1}{2}\right) \right)^2 (1 - |\sigma|) e^{-\sigma\varepsilon} d\sigma \right), \end{aligned}$$

where

$$k_n := \lfloor e^{(n-1)\varepsilon} \rfloor + 1 \quad \text{and} \quad \sigma_n := \min \{1, \varepsilon^{-1} \log(k_n) - n\} \in (-1, 1]. \quad (6.22)$$

Proof. Let $\kappa \in \mathbb{R}$. By (6.19), (6.10) and (1.2), we find that

$$\tilde{I}_n(\kappa) = I_{n,1}(\kappa) = \int_{x_{n+1}}^{x_n} \left(\int_{x_2}^{x_1} (\kappa - K(xy, 1))^2 dy \right) dx.$$

By (1.9) and (1.10), we have also $x_i = e^{(1-i)\varepsilon}$ for $i = 1, \dots, N+1$. Therefore, upon applying the substitutions $x = e^{(\theta-n)\varepsilon}$, $y = e^{(\phi-1)\varepsilon}$ to the last integral above, we obtain:

$$\tilde{I}_n(\kappa) = \varepsilon^2 e^{-(n+1)\varepsilon} \int_0^1 \int_0^1 \left(\kappa - K(e^{(\theta+\phi-(n+1))\varepsilon}, 1) \right)^2 e^{(\theta+\phi)\varepsilon} d\theta d\phi.$$

Then, by means of another pair of substitutions, $\phi = \frac{1}{2}((1-\sigma) + \tau)$, $\theta = \frac{1}{2}((1-\sigma) - \tau)$, we find that

$$\begin{aligned} \tilde{I}_n(\kappa) &= \varepsilon^2 e^{-(n+1)\varepsilon} \int_{-1}^1 \int_{|\sigma|-1}^{1-|\sigma|} \left(\kappa - K(e^{-(\sigma+n)\varepsilon}, 1) \right)^2 e^{(1-\sigma)\varepsilon} \cdot \frac{1}{2} d\tau d\sigma \\ &= \varepsilon^2 e^{-n\varepsilon} \int_{-1}^1 \left(\kappa - K(e^{-(\sigma+n)\varepsilon}, 1) \right)^2 (1 - |\sigma|) e^{-\sigma\varepsilon} d\sigma. \end{aligned} \quad (6.23)$$

By (1.2) we have here:

$$-K(e^{-(\sigma+n)\varepsilon}, 1) = e^{(\sigma+n)\varepsilon} - \left(\lfloor e^{(\sigma+n)\varepsilon} \rfloor + \frac{1}{2} \right) .$$

The stated result of the lemma will therefore follow from (6.23) if it can be shown that one has both

$$\lfloor e^{(\sigma+n)\varepsilon} \rfloor = k_n - 1 \quad \text{for } -1 < \sigma < \sigma_n , \quad (6.24)$$

$$\lfloor e^{(\sigma+n)\varepsilon} \rfloor = k_n \quad \text{for } \sigma_n < \sigma < 1 , \quad (6.25)$$

and also $-1 \leq \sigma_n \leq 1$.

Regarding the last mentioned requirement (proof that $-1 \leq \sigma_n \leq 1$), we observe that, by the first part of (6.22), one has $k_n > e^{(n-1)\varepsilon}$, and so $\varepsilon^{-1} \log(k_n) - n > (n-1) - n = -1$; it therefore follows by the second part of (6.22) that $\sigma_n \leq 1$ and $\sigma_n \in \{1, \varepsilon^{-1} \log(k_n) - n\} \subset (-1, \infty)$, so that one has, in fact, $-1 < \sigma_n \leq 1$.

For (6.24) we observe that, when $\sigma \in (-1, \sigma_n)$, one has

$$e^{(n-1)\varepsilon} < e^{(n+\sigma)\varepsilon} < e^{(n+\sigma_n)\varepsilon} \leq k_n = \left(\lfloor e^{(n-1)\varepsilon} \rfloor + 1 \right) \in \mathbb{Z}$$

(the last two relations holding by virtue of (6.22)), and so

$$k_n - 1 = \lfloor e^{(n-1)\varepsilon} \rfloor \leq \lfloor e^{(n+\sigma)\varepsilon} \rfloor \leq k_n - 1 ,$$

by which we must have $\lfloor e^{(n+\sigma)\varepsilon} \rfloor = k_n - 1$ (as was required to be shown).

Finally, for (6.25) we begin by noting that, when $\sigma_n < \sigma < 1$, one must have $\sigma_n < 1$, and so (by (6.22)) $\sigma_n = \varepsilon^{-1} \log(k_n) - n$, which implies that $k_n = e^{(n+\sigma_n)\varepsilon}$. Therefore, assuming that $\sigma_n < \sigma < 1$, we will have

$$e^{(n+1)\varepsilon} > e^{(\sigma+n)\varepsilon} > e^{(\sigma_n+n)\varepsilon} = k_n ,$$

and so (given that we have $\mathbb{Z} \ni k > e^{(n-1)\varepsilon}$, as has already been noted) may deduce that

$$e^{(n+1)\varepsilon} > \lfloor e^{(\sigma+n)\varepsilon} \rfloor \geq k_n > e^{(n-1)\varepsilon} .$$

Since $n \leq N$, it follows by this last sequence of three inequalities and (1.8) that

$$0 \leq \lfloor e^{(\sigma+n)\varepsilon} \rfloor - k_n < 2 \sinh(\varepsilon) e^{n\varepsilon} \leq 2 \sinh(\varepsilon) e^{N\varepsilon} = 1 ,$$

when $\sigma_n < \sigma < 1$: as two integers that differ by less than 1 must be equal, this completes the proof of (6.25), and, with that, the proof of the lemma also \blacksquare

Lemma 6.5. *Let $n \leq N$ be a positive integer. Let $\tilde{\mu}_n$ be the real number defined by (6.18) and (1.13). Then*

$$\tilde{\mu}_n = k_n - \frac{\varepsilon^2 e^{\varepsilon n} + \sinh(\varepsilon) - (|1 - e^{-\varepsilon \sigma_n}| + \varepsilon e^{-\varepsilon \sigma_n} (1 - |\sigma_n|))}{2 (\cosh(\varepsilon) - 1)} ,$$

where k_n and σ_n are as in (6.22).

Proof. By (6.20), the real function $\kappa \mapsto \tilde{I}_n(\kappa)$ attains its global minimum at the point $\kappa = \tilde{\mu}_n$, and so (bearing in mind that $\tilde{I}_n(\kappa)$ is a polynomial in κ) we must have $\tilde{I}'_n(\tilde{\mu}_n) = 0$. By this and Lemma 6.4, we find that one has

$$0 = \int_{-1}^{\sigma_n} \left(\tilde{\mu}_n + e^{(\sigma+n)\varepsilon} - \left(k_n - \frac{1}{2}\right) \right) (1 - |\sigma|) e^{-\sigma\varepsilon} d\sigma \\ + \int_{\sigma_n}^1 \left(\tilde{\mu}_n + e^{(\sigma+n)\varepsilon} - \left(k_n + \frac{1}{2}\right) \right) (1 - |\sigma|) e^{-\sigma\varepsilon} d\sigma ,$$

and so:

$$(\tilde{\mu}_n - k_n) X(\varepsilon, -1) + e^{n\varepsilon} X(0, -1) = \frac{1}{2} X(\varepsilon, \sigma_n) , \quad (6.26)$$

where

$$X(\omega, \alpha) := \int_{\alpha}^1 (1 - |\sigma|) e^{-\sigma\omega} d\sigma - \int_{-1}^{\alpha} (1 - |\sigma|) e^{-\sigma\omega} d\sigma \quad (6.27)$$

for $\omega \in \mathbb{R}$, $\alpha \in [-1, 1]$. Here $X(0, -1)$ is the area of the triangle with vertices at $(-1, 0)$, $(0, 1)$ and $(0, 1)$, which is, of course, equal to 1. By elementary calculus, we have also:

$$X(\omega, -1) = \frac{2(\cosh(\omega) - 1)}{\omega^2} = \left(\frac{\sinh(\frac{1}{2}\omega)}{\frac{1}{2}\omega} \right)^2 \quad (\mathbb{R} \ni \omega \neq 0). \quad (6.28)$$

By (6.26), (6.27), (6.28) and the equality $X(0, -1) = 1$, we have:

$$\tilde{\mu}_n - k_n = \frac{\frac{1}{2}\varepsilon^2 X(\varepsilon, \sigma_n) - \varepsilon^2 e^{n\varepsilon}}{2(\cosh(\varepsilon) - 1)} .$$

Therefore the stated result of the lemma follows if it can be shown that one has:

$$\frac{1}{2} X(\varepsilon, \sigma_n) = \frac{e^{-\varepsilon\sigma_n} (1 - |\sigma_n|)}{\varepsilon} + \frac{|1 - e^{-\varepsilon\sigma_n}| - \sinh(\varepsilon)}{\varepsilon^2} . \quad (6.29)$$

We begin the verification of (6.29) with the observation that, if $0 \leq \alpha \leq 1$ and $\mathbb{R} \ni \omega \neq 0$, then, by (6.27) and (6.28), one has:

$$\frac{1}{2} X(\omega, \alpha) = \frac{1}{2} \left(2 \int_{\alpha}^1 (1 - \sigma) e^{-\sigma\omega} d\sigma - X(\omega, -1) \right) \\ = \frac{(1 - \alpha)e^{-\alpha\omega}}{\omega} + \frac{e^{-\omega} - e^{-\alpha\omega}}{\omega^2} - \frac{\cosh(\omega) - 1}{\omega^2} \\ = \frac{e^{-\alpha\omega}(1 - \alpha)}{\omega} + \frac{(1 - e^{-\alpha\omega}) - \sinh(\omega)}{\omega^2} . \quad (6.30)$$

Given that we have $\varepsilon > 0$, it follows immediately from (6.30) that (6.29) is valid when $0 \leq \sigma_n \leq 1$. In dealing with the remaining cases (where $-1 < \sigma_n < 0$) we find that

the substitution $\sigma = -\tau$ (in (6.27)) gives $X(\varepsilon, \sigma_0) = -X(-\varepsilon, -\sigma_0)$, and so, by (6.30) for $\omega = -\varepsilon \in \mathbb{R} \setminus \{0\}$, $\alpha = -\sigma_0 \in (0, 1)$, we obtain

$$\begin{aligned} \frac{1}{2}X(\varepsilon, \sigma_n) &= - \left(\frac{e^{-\varepsilon\sigma_0}(1 + \sigma_0)}{(-\varepsilon)} + \frac{(1 - e^{-\varepsilon\sigma_0}) + \sinh(\varepsilon)}{\varepsilon^2} \right), \\ &= \frac{e^{-\varepsilon\sigma_0}(1 + \sigma_0)}{\varepsilon} + \frac{(e^{-\varepsilon\sigma_0} - 1) - \sinh(\varepsilon)}{\varepsilon^2}, \end{aligned}$$

which is in exact agreement with (6.29), given we are supposing here that $\sigma_0 < 0$, and $\varepsilon > 0$. This completes the proof of the lemma \blacksquare

Remark. Lemma 6.5 is not part of our proof of Theorem 4, but is of use in computing eigenvalues of the kernel $K^{\mathcal{A}, N}$: see Section 7 for some details of this.

It can be shown that, when σ_n (as defined in (6.22)) is equal to 1, one has

$$\tilde{I}_n(\tilde{\mu}_n) = \varepsilon^2 e^{n\varepsilon} \left(\frac{\sinh^2(\frac{1}{2}\varepsilon)}{(\frac{1}{2}\varepsilon)^2} - \frac{(\frac{1}{2}\varepsilon)^2}{\sinh^2(\frac{1}{2}\varepsilon)} \right), \quad (6.31)$$

which (by virtue of (1.9) and the identity $4\sinh^2 x = e^{2x} + e^{-2x} - 2$) is a certain rational function of δ , δ^n and ε^2 . With more work one can compute a rational function of δ , δ^n , ε , n , k_n and σ_n (the last two being given by (6.22)) that is equal to $\tilde{I}_n(\tilde{\mu}_n)$ if $0 \leq \sigma_n < 1$, as well as another such function that is equal to $\tilde{I}_n(\tilde{\mu}_n)$ if $-1 < \sigma_n \leq 0$. The latter two exact results (pertaining to cases where $-1 < \sigma_n < 1$) are complicated enough to dissuade us from using them in our proof of Theorem 4, and so it would serve no purpose to provide more details of those results in this paper. We seek instead to work with simpler approximate results, and (what is more) do not believe that the complicated exact results are an ideal starting point from which to make our way to a satisfactory ‘simple’ approximation to $\tilde{I}_n(\tilde{\mu}_n)$. For these reasons we take a different approach: our goal becomes a reasonably sharp upper bound on $\tilde{I}_n(\tilde{\mu}_n)$, to be obtained by the application of the inequality (6.20) for a suitably chosen $\kappa \in \mathbb{R}$ (which, in light of (6.14), must necessarily be an approximation to $\tilde{\mu}_n$).

In order to arrive at a suitable choice for κ , in applying (6.20), we revisit the equations (6.26)-(6.28) that occur in the proof of Lemma 6.5. By (6.26) and (6.27), we have

$$\begin{aligned} &\left(\tilde{\mu}_n + e^{(n+\sigma_n)\varepsilon} - k_n \right) X(\varepsilon, -1) \\ &= \int_{-1}^1 \left(e^{(n+\sigma_n)\varepsilon} - e^{(n+\sigma)\varepsilon} \right) (1 - |\sigma|) e^{-\sigma\varepsilon} d\sigma + \frac{1}{2}X(\varepsilon, \sigma_n) \\ &= e^{(n+\sigma_n)\varepsilon} \int_{-1}^1 \left(1 - e^{(\sigma-\sigma_n)\varepsilon} \right) (1 - |\sigma|) e^{-\sigma\varepsilon} d\sigma + \frac{1}{2}X(\varepsilon, \sigma_n) \end{aligned}$$

We next modify the above equation, replacing each term of form $X(\varepsilon, \alpha)$ by $X(0, \alpha)$, while replacing $\tilde{\mu}_n$ by some μ'_n . Note that $X(0, -1) = 1$, and that by (6.28) we have

$X(\varepsilon, -1) = 1 + O(\varepsilon^2) = (1 + O(\varepsilon^2))X(0, -1)$. By (6.22) and (6.27), we have also $X(\varepsilon, \sigma_n) = X(0, \sigma_n) + O(\varepsilon)$. Thus, given the result (6.3) of Lemma 6.1, we see that if N is large enough then $X(0, -1)$ and $X(0, \sigma_n)$ will be quite good approximations for $X(\varepsilon, -1)$ and $X(\varepsilon, \sigma_n)$, respectively, and so the number μ'_n satisfying

$$\mu'_n + e^{(n+\sigma_n)\varepsilon} - k_n = e^{(n+\sigma_n)\varepsilon} \int_{-1}^1 \left(1 - e^{(\sigma-\sigma_n)\varepsilon}\right) (1 - |\sigma|) e^{-\sigma\varepsilon} d\sigma + \frac{1}{2}X(0, \sigma_n) \quad (6.32)$$

will then be a reasonable approximation for $\tilde{\mu}_n$.

We note that the integrand occurring in the last equation has, as a factor, the product:

$$\left(1 - e^{(\sigma-\sigma_n)\varepsilon}\right) e^{-\sigma\varepsilon} = \left(-(\sigma - \sigma_n)\varepsilon + O(\varepsilon^2)\right) (1 + O(\varepsilon)) = -(\sigma - \sigma_n)\varepsilon + O(\varepsilon^2) .$$

By this, (6.32), (6.27), and a short calculation, we arrive at a second, simpler, approximation to $\tilde{\mu}_n$:

$$\begin{aligned} \mu''_n &:= \left(k_n - e^{(n+\sigma_n)\varepsilon}\right) - \varepsilon e^{(n+\sigma_n)\varepsilon} \int_{-1}^1 (1 - |\sigma|) (\sigma - \sigma_n) d\sigma + \frac{1}{2}X(0, \sigma_n) \\ &= \left(k_n - e^{(n+\sigma_n)\varepsilon}\right) + \varepsilon \sigma_n e^{(n+\sigma_n)\varepsilon} X(0, -1) + \frac{1}{2}X(0, \sigma_n) \\ &= k_n + (\varepsilon \sigma_n - 1) e^{(n+\sigma_n)\varepsilon} - \frac{1}{2}\chi(\sigma_n) \in \mathbb{R} , \end{aligned} \quad (6.33)$$

where

$$\chi(\alpha) := -X(0, \alpha) = (2 - |\alpha|)\alpha \quad (-1 \leq \alpha \leq 1). \quad (6.34)$$

Our application of the inequality (6.20) will, in all cases, be made with $\kappa = \mu''_n$. The following lemma is therefore central to the completion of our proof of Theorem 4.

Lemma 6.6. *Let $n \leq N$ be a positive integer. Then*

$$\tilde{I}_n(\mu''_n) \leq \frac{(1-\chi^2(\sigma_n))}{4} \varepsilon^2 e^{(1-n)\varepsilon} + \psi(\sigma_n) \varepsilon^3 e^{(1+\sigma_n)\varepsilon} + \frac{1}{6} \varepsilon^4 e^{(n+1+2\sigma_n)\varepsilon} + \left(\frac{1}{6} + \sigma_n^2\right) e^{2\varepsilon} \varepsilon^4 ,$$

where

$$\psi(\alpha) := -\frac{2}{3}|\alpha|^3 + \alpha^2 - \frac{1}{3} \quad (-1 \leq \alpha \leq 1), \quad (6.35)$$

while k_n , σ_n , μ''_n and $\chi(\sigma)$ are as defined in (6.22), (6.33) and (6.34).

Proof. By (6.33) and Lemma 6.4, one has

$$\begin{aligned} \tilde{I}_n(\mu''_n) &= \varepsilon^2 \int_{-1}^1 \left((\varepsilon \sigma_n - 1) e^{(n+\sigma_n)\varepsilon} + e^{(\sigma+n)\varepsilon} - \phi(\sigma_n, \sigma) \right)^2 \left(\frac{1 - |\sigma|}{e^{(n+\sigma)\varepsilon}} \right) d\sigma \\ &= \varepsilon^2 \int_{-1}^1 \left(\phi(\sigma_n, \sigma) + e^{(n+\sigma_n)\varepsilon} \varrho((\sigma - \sigma_n)\varepsilon) - e^{(n+\sigma_n)\varepsilon} \varepsilon \sigma \right)^2 \left(\frac{1 - |\sigma|}{e^{(n+\sigma)\varepsilon}} \right) d\sigma , \end{aligned} \quad (6.36)$$

where:

$$\phi(x, y) := \frac{1}{2}\chi(x) + \left\lfloor \frac{1}{2}(y - x) \right\rfloor + \frac{1}{2} \quad (-1 < x, y \leq 1) \quad (6.37)$$

and

$$\varrho(t) := 1 + t - e^t \quad (t \in \mathbb{R}). \quad (6.38)$$

By ‘Taylor’s formula with remainder’, one has here $\rho(t) = -\frac{1}{2}t^2 e^{\vartheta(t)t}$, where the range of the function $\vartheta(t)$ is contained in the interval $(0, 1)$. It follows that

$$0 \geq \varrho(t) \geq -\frac{1}{2}t^2 e^{\max\{0,t\}} \quad (t \in \mathbb{R}). \quad (6.39)$$

As a preliminary to the application of (6.39), we consider now the range of the function

$$f(\sigma) := \phi(\sigma_n, \sigma) + e^{(n+\sigma_n)\varepsilon} \varrho((\sigma - \sigma_n)\varepsilon) - e^{(n+\sigma_n)\varepsilon} \varepsilon \sigma \quad (-1 < \sigma < 1),$$

the square of which is a factor of the integrand seen in (6.36). We note firstly that, when $-1 < \sigma < \sigma_n \leq 1$, it follows by (6.37) and (6.38) that one has both

$$\phi(\sigma_n, \sigma) = \frac{1}{2}\chi(\sigma_n) - \frac{1}{2} \quad \text{and} \quad \varrho((\sigma - \sigma_n)\varepsilon) - \varepsilon\sigma = 1 - \sigma_n\varepsilon - e^{(\sigma - \sigma_n)\varepsilon} > -\sigma_n\varepsilon,$$

and so

$$f(\sigma) > -\left(e^{(n+\sigma_n)\varepsilon} \sigma_n\varepsilon + \frac{1}{2} - \frac{1}{2}\chi(\sigma_n)\right). \quad (6.40)$$

When $-1 < \sigma_n \leq \sigma < 1$, one finds instead that

$$\phi(\sigma_n, \sigma) = \frac{1}{2}\chi(\sigma_n) + \frac{1}{2} \quad \text{and} \quad \varrho((\sigma - \sigma_n)\varepsilon) - \varepsilon\sigma > -(\sigma_n\varepsilon + e^{(1-\sigma_n)\varepsilon} - 1).$$

Therefore, given that one has

$$e^{(n+\sigma_n)\varepsilon} (e^{(1-\sigma_n)\varepsilon} - 1) = (e^\varepsilon - e^{\sigma_n\varepsilon}) e^{n\varepsilon} < 2 \sinh(\varepsilon) e^{n\varepsilon} = e^{(n-N)\varepsilon} \leq 1$$

(the penultimate relation here following by (1.8)), we may conclude that the inequality (6.40), shown (earlier) to hold for all $\sigma \in (-1, \sigma_n)$, does in fact hold for all $\sigma \in (-1, 1)$.

We observe, secondly, that $-1 < \sigma_n \leq 1$, and that if $\sigma_n < 0$, so that $\sigma_n \in (-1, 0)$, then it follows by (6.34) that

$$e^{(n+\sigma_n)\varepsilon} \sigma_n\varepsilon + \frac{1}{2} - \frac{1}{2}\chi(\sigma_n) < \frac{1}{2} - \frac{1}{2}\chi(\sigma_n) < \frac{1}{2} - \frac{1}{2}\chi(-1) = 1;$$

if instead $0 \leq \sigma_n \leq 1$, then, by (1.8), (1.9), (6.1) and the result (6.5) of Lemma 6.1, one has

$$\begin{aligned} e^{(n+\sigma_n)\varepsilon} \sigma_n\varepsilon + \frac{1}{2} - \frac{1}{2}\chi(\sigma_n) &\leq e^{(N+1)\varepsilon} \sigma_n\varepsilon + \frac{1}{2} - \frac{1}{2}\sigma_n \\ &= \left(\frac{\Delta\varepsilon}{\sinh(2\varepsilon)} - \frac{1}{2}\right) \sigma_n + \frac{1}{2} \\ &< \left(\frac{\Delta - 1}{2}\right) \sigma_n + \frac{1}{2} \leq \left(\frac{\Delta - 1}{2}\right) (1) + \frac{1}{2} = \frac{\Delta}{2} < \frac{2}{3}. \end{aligned}$$

By these observations, together with (6.40), we deduce that

$$f(\sigma) > -1 \quad (-1 < \sigma < 1). \quad (6.41)$$

By (6.39) and the definition of $f(\sigma)$, we have

$$f(\sigma) \leq g(\sigma) := \phi(\sigma_n, \sigma) - e^{(n+\sigma_n)\varepsilon} \varepsilon \sigma \quad (-1 < \sigma < 1),$$

and so it follows by (6.41) that, for $-1 < \sigma < 1$, we have:

$$f^2 \leq f^2 + (f - g)^2 = g^2 - 2(g - f)f < g^2 + 2(g - f) \quad (f = f(\sigma), g = g(\sigma)).$$

By (6.39) (again), we have here

$$g(\sigma) - f(\sigma) = -e^{(n+\sigma_n)\varepsilon} \varrho((\sigma - \sigma_n)\varepsilon) \leq \frac{1}{2}\varepsilon^2 (\sigma - \sigma_n)^2 e^{(n+\max\{\sigma_n, \sigma\})\varepsilon} \quad (-1 < \sigma < 1),$$

and so may deduce that

$$f^2(\sigma) \leq g^2(\sigma) + \varepsilon^2 (\sigma - \sigma_n)^2 e^{(n+1)\varepsilon} \quad (-1 < \sigma < 1). \quad (6.42)$$

Since we have

$$\begin{aligned} \int_{-1}^1 (\sigma - \sigma_n)^2 e^{(n+1)\varepsilon} \left(\frac{1 - |\sigma|}{e^{(n+\sigma)\varepsilon}} \right) d\sigma &\leq e^{2\varepsilon} \int_{-1}^1 (\sigma - \sigma_n)^2 (1 - |\sigma|) d\sigma \\ &= 2e^{2\varepsilon} \int_0^1 (\sigma^2 + \sigma_n^2) (1 - \sigma) d\sigma = \left(\frac{1}{6} + \sigma_n^2 \right) e^{2\varepsilon}, \end{aligned}$$

it follows from (6.36) and (6.42) that

$$\begin{aligned} \tilde{I}_n(\mu_n'') &\leq \varepsilon^2 \int_{-1}^1 g^2(\sigma) \left(\frac{1 - |\sigma|}{e^{(n+\sigma)\varepsilon}} \right) d\sigma + \left(\frac{1}{6} + \sigma_n^2 \right) e^{2\varepsilon} \varepsilon^4 \\ &= \left(\frac{1}{6} + \sigma_n^2 \right) e^{2\varepsilon} \varepsilon^4 + \varepsilon^2 \int_{-1}^1 \left(\phi(\sigma_n, \sigma) - e^{(n+\sigma_n)\varepsilon} \varepsilon \sigma \right)^2 \left(\frac{1 - |\sigma|}{e^{(n+\sigma)\varepsilon}} \right) d\sigma \\ &\leq \left(\frac{1}{6} + \sigma_n^2 \right) e^{2\varepsilon} \varepsilon^4 + \varepsilon^2 e^{(1-n)\varepsilon} \int_{-1}^1 \left(\phi(\sigma_n, \sigma) - e^{(n+\sigma_n)\varepsilon} \varepsilon \sigma \right)^2 (1 - |\sigma|) d\sigma. \quad (6.43) \end{aligned}$$

By (6.37), the value of the integral occurring in the last inequality above is:

$$\begin{aligned} &\int_{-1}^{\sigma_n} \left(\frac{1}{2}\chi(\sigma_n) - \frac{1}{2} - e^{(n+\sigma_n)\varepsilon} \varepsilon \sigma \right)^2 (1 - |\sigma|) d\sigma \\ &+ \int_{\sigma_n}^1 \left(\frac{1}{2}\chi(\sigma_n) + \frac{1}{2} - e^{(n+\sigma_n)\varepsilon} \varepsilon \sigma \right)^2 (1 - |\sigma|) d\sigma \\ &= \int_{-1}^1 \left(\left(\frac{1}{2}\chi(\sigma_n) - e^{(n+\sigma_n)\varepsilon} \varepsilon \sigma \right)^2 + \frac{1}{4} \right) (1 - |\sigma|) d\sigma \\ &+ \int_{\sigma_n}^1 \left(\frac{1}{2}\chi(\sigma_n) - e^{(n+\sigma_n)\varepsilon} \varepsilon \sigma \right) (1 - |\sigma|) d\sigma \\ &- \int_{-1}^{\sigma_n} \left(\frac{1}{2}\chi(\sigma_n) - e^{(n+\sigma_n)\varepsilon} \varepsilon \sigma \right) (1 - |\sigma|) d\sigma = J_1 + J_2 - J_3 \quad (\text{say}). \quad (6.44) \end{aligned}$$

Since $\int_{-1}^1 \sigma(1 - |\sigma|)d\sigma = 0$, we have here

$$\begin{aligned} J_1 &= \frac{1}{4} (\chi^2(\sigma_n) + 1) \int_{-1}^1 (1 - |\sigma|) d\sigma + \varepsilon^2 e^{2(n+\sigma_n)\varepsilon} \int_{-1}^1 \sigma^2 (1 - |\sigma|) d\sigma \\ &= \frac{1}{4} (\chi^2(\sigma_n) + 1) (1) + \frac{1}{6} \varepsilon^2 e^{2(n+\sigma_n)\varepsilon} . \end{aligned} \quad (6.45)$$

We have, moreover,

$$J_2 - J_3 = \frac{1}{2} \chi(\sigma_n) X(0, \sigma_n) - \varepsilon e^{(n+\sigma_n)\varepsilon} Y(\sigma_n) ,$$

where $X(\omega, \alpha)$ is given by (6.27), while

$$Y(\alpha) := \int_{\alpha}^1 \sigma(1 - |\sigma|) d\sigma - \int_{-1}^{\alpha} \sigma(1 - |\sigma|) d\sigma \quad (-1 \leq \alpha \leq 1).$$

Use of the substitution $\sigma = -\tau$ reveals that the function $Y(\alpha)$ is even: we find that, for $-1 \leq \alpha \leq \alpha$, one has

$$Y(\alpha) = Y(|\alpha|) = 2 \int_{|\alpha|}^1 \sigma(1 - |\sigma|) d\sigma - \int_{-1}^1 \sigma(1 - |\sigma|) d\sigma = 2 \int_{|\alpha|}^1 \sigma(1 - |\sigma|) d\sigma = -\psi(\alpha) ,$$

with $\psi(\alpha)$ as defined in (6.35). By these observations and (6.34), it follows that

$$J_2 - J_3 = \psi(\sigma_n) \varepsilon e^{(n+\sigma_n)\varepsilon} - \frac{1}{2} \chi^2(\sigma_n) .$$

This, combined with (6.43), (6.44) and (6.45), yields the stated result of the lemma \blacksquare

Lemma 6.6 will (shortly) be used to bring us a step nearer to completing our proof of Theorem 4. Another, different, use for Lemma 6.6 is its application to the problem of computing, for a chosen (sufficiently large) positive integer N , a useful numerical upper bound on the real number $\|K^{\boxtimes}\|^2 = \|K^{\boxtimes, N}\|^2$. Indeed, one solution of this problem of numerical computation is to appeal to (6.9), Lemma 6.3 and (6.21), and then bound each term of the sum over n in (6.21) through separate applications of Lemma 6.6 (combined with the case $\kappa = \mu_n''$ of (6.20)): the bound obtained for that sum over n is then just the sum of the N separate numerical bounds computed for its terms. In the context of such ‘computational’ applications, the result in Lemma 6.6 has the practical advantage of it being (essentially) the strongest upper bound on $\tilde{I}_n(\mu_n'')$ that we are able to prove, and this outweighs the minor drawback of its being a somewhat complicated result to state. In our approach to the proof of Theorem 4, however, we are happy to give up some degree of the precision offered by Lemma 6.6 in exchange for a simpler upper bound on $\tilde{I}_n(\mu_n'')$. At the same time, we aim to be precise enough to ensure that the ‘leading term’ in the resulting upper bound for $\|K^{\boxtimes}\|^2$ is essentially optimal, subject to the constraints imposed by our method of obtaining this bound.

Our first simplification of the result of Lemma 6.6 is based on the observation that the definition (6.35) implies that the range of the function $\psi(\alpha)$ ($-1 \leq \alpha \leq 1$) is the interval

$[\psi(0), \psi(1)] = [-\frac{1}{3}, 0]$. It follows that the second term in the bound of Lemma 6.6 for $\tilde{I}_n(\mu_n'')$, which is the term $\psi(\sigma_n) \varepsilon^3 e^{(1+\sigma_n)\varepsilon}$, is always less than or equal to 0, so that one may omit this term from the inequality stated in Lemma 6.6: the inequality will remain valid. To simplify the result further, we observe that (since $\varepsilon > 0$) it follows by (6.22) that one has $\frac{1}{6} + \sigma_n^2 \leq \frac{7}{6}$ and $\max\{e^{(n+1+2\sigma_n)\varepsilon}, e^{2\varepsilon}\} \leq e^{(n+3)\varepsilon}$, for $n = 1, \dots, N$: by (1.9), (6.1) and (6.5), we have here $1 < e^\varepsilon = \Delta < \frac{4}{3}$, and so may conclude that, as a consequence of Lemma 6.6, (6.33) and (6.20) (for $\kappa = \mu_n''$), one has

$$\tilde{I}_n(\tilde{\mu}_n) \leq \tilde{I}_n(\mu_n'') \leq \frac{(1 - \chi^2(\sigma_n)) \varepsilon^2}{4e^{(n-1)\varepsilon}} + O(\varepsilon^4 e^{n\varepsilon}), \quad (6.46)$$

for $n = 1, \dots, N$.

By (6.34), $1 - \chi^2(\alpha) = 0$ if $\alpha = \pm 1$: in particular, $1 - \chi^2(\sigma_n) = 0$ if $\sigma_n = 1$. It follows, by virtue of the definitions (6.22), that if $k_n \geq e^{(n+1)\varepsilon}$ then one may omit the first term on the right-hand side of the rightmost inequality in (6.46) (for that term will then be equal to 0); if instead $k_n < e^{(n+1)\varepsilon}$, then, by (6.22), k_n is a positive integer lying in the open interval $(e^{(n-1)\varepsilon}, e^{(n+1)\varepsilon}) = (\Delta^{n-1}, \Delta^{n+1})$, so that $e^{(n-1)\varepsilon} > \Delta^{-2}k_n$, while $\sigma_n = \varepsilon^{-1} \log(k_n) - n$. It therefore follows from (6.46) that, for $n = 1, \dots, N$, one has

$$\tilde{I}_n(\tilde{\mu}_n) \leq O(\varepsilon^4 e^{n\varepsilon}) + \frac{1}{4} \Delta^2 \varepsilon^2 \sum_{\Delta^{n-1} < k < \Delta^{n+1}} \frac{1 - \chi^2\left(\frac{\log k}{\varepsilon} - n\right)}{k} \quad (6.47)$$

(we use here the fact that, since $\Delta = e^\varepsilon > 1$, the definition (6.34) ensures that every term of the last sum is defined and greater than or equal to 0). We remark that, when $n \leq N$, it follows by (1.8), (1.9), (6.1) and (6.5), that one has $\Delta^{n+1} - \Delta^{n-1} \leq \Delta^{n-N} \leq 1$, so that the sum over k in (6.47) contains at most one term. When there exists a unique integer k satisfying $\Delta^{n-1} < k < \Delta^{n+1}$, that integer must then be the least integer greater than Δ^{n-1} , and this (given (6.22)) implies that $k = k_n$. We conclude that, for $n = 1, \dots, N$, the bound in (6.47) is never more than a factor of Δ^2 greater than the upper bound on $\tilde{I}_n(\mu_n'')$ in (6.46): since (6.3) and (6.5) give us $\Delta = 1 + O((\log N)/N)$ (as $N \rightarrow \infty$), what we lose by using (6.47) instead of (6.46) is so little as to be insignificant.

The next lemma is an upper bound for the term V_0 that occurs in (6.9). To prove it we shall use of (6.47), combined with the observation in (6.21).

Lemma 6.7 *Let $\tilde{B}_4(t)$ be the periodic Bernoulli function associated with the Bernoulli polynomial $B_4(X)$ of degree 4. Then one has*

$$V_0 \leq \left(\frac{7}{60} + O\left(\frac{1}{\log N}\right) \right) N^{-1} \log^3(N) + \frac{1}{2} \varepsilon R,$$

where

$$R := - \sum_{1 < k < \Delta^{N+1}} (\log k) k^{-1} \tilde{B}_4\left(\frac{\log k}{\varepsilon}\right). \quad (6.48)$$

Proof. By (6.21) and (6.47), one has

$$V_0 = P + \frac{1}{4}\Delta^2\varepsilon^2Q, \quad (6.49)$$

where

$$P = \sum_{n=1}^N n \cdot O(\varepsilon^4 e^{n\varepsilon}) \ll \frac{N\varepsilon^4 e^{N\varepsilon}}{1 - e^{-\varepsilon}} \ll N\varepsilon^2 \quad (6.50)$$

(the last two relation following by virtue of (6.2) and the inequalities $0 < \varepsilon < \frac{1}{4}$), while

$$\begin{aligned} Q &= \sum_{n=1}^N n \sum_{\Delta^{n-1} < k < \Delta^{n+1}} \frac{1 - \chi^2\left(\frac{\log k}{\varepsilon} - n\right)}{k} \\ &= \sum_{1 < k < \Delta^{N+1}} \frac{1}{k} \sum_{\frac{\log k}{\varepsilon} - 1 < n \leq \frac{\log k}{\varepsilon} + 1} \left(1 - \chi^2\left(\frac{\log k}{\varepsilon} - n\right)\right) n \end{aligned} \quad (6.51)$$

(the last of these equalities following by virtue of (6.1), (6.5), (1.9) and (6.34), the last of which ensures that summand in the last sum over n occurring in (6.51) is equal to 0 in any cases where $n = \varepsilon^{-1} \log(k) + 1$).

Now a given $n \in \mathbb{N}$ satisfies the conditions of summation seen in the final line of (6.51) if and only if n lies in one of the two disjoint intervals $(\varepsilon^{-1} \log(k) - 1, \varepsilon^{-1} \log(k)]$, $(\varepsilon^{-1} \log(k), \varepsilon^{-1} \log(k) + 1]$. The integer n lies in the first of these two intervals if and only if $n = \lfloor \varepsilon^{-1} \log(k) \rfloor$; it lies in the second interval if and only if $n = \lfloor \varepsilon^{-1} \log(k) \rfloor + 1$. We conclude that, in the final line of (6.51), the sum over n may be expressed as the sum

$$(1 - \chi^2(\beta))m + (1 - \chi^2(\beta - 1))(m + 1) = (2 - \chi^2(\beta) - \chi^2(\beta - 1)) \varepsilon^{-1} \log k + O(1),$$

where $\beta = \{\varepsilon^{-1} \log(k)\} \in [0, 1)$ and $m = \varepsilon^{-1} \log(k) - \beta = \lfloor \varepsilon^{-1} \log(k) \rfloor$: note that we have here used the fact that (6.34) implies that $|\chi(\alpha)| \leq 1$ for $-1 \leq \alpha \leq 1$. By (6.34) (again), we find that when $0 \leq \beta < 1$ one has: $\chi^2(\beta) + \chi^2(\beta - 1) = (2 - \beta)^2 \beta^2 + (1 + \beta)^2 (\beta - 1)^2 = 2\beta^4 - 4\beta^3 + 2\beta^2 + 1 = 2B_4(\beta) + \frac{16}{15}$.

By (6.51), (1.9), (6.1), (6.2) and the points noted in the preceding paragraph, it follows that

$$\begin{aligned} Q &= O(\log(\Delta^{N+1})) + \frac{1}{\varepsilon} \sum_{1 < k < \Delta^{N+1}} \left(\frac{14}{15} - 2B_4\left(\left\{\frac{\log k}{\varepsilon}\right\}\right) \right) k^{-1} \log k \\ &= O((N+1)\varepsilon) + \frac{14}{15}\varepsilon^{-1} (O(1) + \frac{1}{2}(N+1)^2\varepsilon^2) - \frac{2}{\varepsilon} \sum_{1 < k < \Delta^{N+1}} \tilde{B}_4\left(\frac{\log k}{\varepsilon}\right) k^{-1} \log k \\ &= \frac{7}{15}\varepsilon N^2 + 2\varepsilon^{-1}R + O(\varepsilon^{-1}), \end{aligned}$$

with R as defined in (6.48). By this, together with (6.49) and (6.50), we find (given (6.2) and (6.5)) that

$$V_0 = \left(\frac{7}{60}\varepsilon^3 N^2 + \frac{1}{2}\varepsilon R\right) \Delta^2 + O(\varepsilon^2 N). \quad (6.52)$$

By (6.3) and (6.5), we have here

$$\varepsilon^3 N^2 < N^{-1} \log^3 N, \quad \varepsilon^2 N < N^{-1} \log^2 N = (\log N)^{-1} N^{-1} \log^3 N$$

and

$$\Delta^2 = (1 + O(\varepsilon))^2 = 1 + O(\varepsilon) = 1 + O\left(\frac{\log N}{N}\right) = 1 + O\left(\frac{1}{\log N}\right),$$

and so also:

$$(\Delta^2 - 1) \varepsilon R \ll \varepsilon^2 R \ll \varepsilon^2 \log^2 (\Delta^{N+1}) = \varepsilon^4 (N+1)^2 \ll N^{-2} \log^4 N \ll N^{-1}$$

(the bound $R \ll \log^2 (\Delta^{N+1})$ being an essentially trivial consequence of (6.48)). With the help of these observations we find that (6.52) implies what is stated in the lemma \blacksquare

As noted in the above proof, the term R defined by (6.48) certainly satisfies $R \ll \log^2 (\Delta^{N+1})$, and so (given (1.9), (6.1) and (6.3)) one has: $\frac{1}{2}\varepsilon R \ll \varepsilon^3 N^2 < N^{-1} \log^3 N$. Thus Lemma 6.7 implies the (weaker) result that V_0 satisfies $V_0 \ll N^{-1} \log^3 N$. In completing the proof of Theorem 4 we shall need to sharpen the above mentioned bound on R (which is essentially a trivial estimate). We seek a bound on R that will, at the very least, enable us to deduce from Lemma 6.1 that

$$V_0 \leq \left(\frac{7}{60} + o(1)\right) N^{-1} \log^3 N \quad (N \rightarrow \infty). \quad (6.53)$$

A first step towards (6.53) is the observation that the periodic function $\tilde{B}_4(x)$ has the Fourier expansion

$$\tilde{B}_4(x) = -\frac{4!}{(2\pi i)^4} \sum_{\substack{\ell=-\infty \\ \ell \neq 0}}^{\infty} \frac{e^{2\pi i \ell x}}{\ell^4} = -\frac{3}{\pi^4} \cdot \operatorname{Re} \left(\sum_{\ell=1}^{\infty} \frac{e^{-2\pi i \ell x}}{\ell^4} \right),$$

which (see [OLBC, 24.2(iii) and 24.8.3]) is valid for all real x . By application of this observation in respect of each term $\log(k)k^{-1}\tilde{B}_4(\varepsilon^{-1} \log k)$ that occurs in the sum in (6.48), we find that

$$\begin{aligned} R &= \frac{3}{\pi^4} \sum_{1 < k < \Delta^{N+1}} (\log k)k^{-1} \cdot \operatorname{Re} \left(\sum_{\ell=1}^{\infty} \frac{e^{-2\pi i \ell \varepsilon^{-1} \log k}}{\ell^4} \right) \\ &= \frac{3}{\pi^4} \sum_{\ell=1}^{\infty} \ell^{-4} \cdot \operatorname{Re} \left(\sum_{1 < k < \Delta^{N+1}} (\log k)k^{-1-2\pi i \ell \varepsilon^{-1} i} \right). \end{aligned} \quad (6.54)$$

We recall that $\zeta'(s)$, the first derivative of the Riemann zeta-function, has the Dirichlet series expansion

$$\zeta'(s) = -\sum_{k=2}^{\infty} \frac{\log k}{k^s} \quad (\operatorname{Re}(s) > 1),$$

which suggests that $\zeta'(1 + 2\pi\ell\varepsilon^{-1}i)$ might be a useful approximation for the value of the sum over k that occurs in the last line of (6.54): that this is indeed the case is shown by the following lemma.

Lemma 6.8. *Let $\ell \in \mathbb{N}$. Then, given what we assume in respect of N , ε and Δ , one has*

$$- \sum_{1 < k < \Delta^{N+1}} (\log k) k^{-1-2\pi\ell\varepsilon^{-1}i} = \zeta'(1 + 2\pi\ell\varepsilon^{-1}i) + O\left(\ell^{1/2} N^{-1/2} \log^{3/2} N\right),$$

where the implicit constant is absolute (and so is, in particular, independent of ℓ).

Proof. We let $x = \Delta^{N+1}$ and define $f(s)$ to be the function defined, for $s \in \mathbb{C} \setminus \{1\}$, by:

$$f(s) = \zeta(s) - \sum_{k < x} k^{-s}.$$

We put also $\sigma_0 = 1$, $t_0 = 2\pi\ell\varepsilon^{-1}$ and $s_0 = \sigma_0 + it_0$. It suffices for proof of the lemma that we show that one has $f'(s_0) \ll \ell^{1/2} N^{-1/2} \log^{3/2} N$ (with an implicit constant that is absolute). This will follow if it is shown that, with

$$r := \frac{1}{4 \log x},$$

one has

$$f(s) \ll \ell^{1/2} N^{-1/2} \log^{1/2} N \quad (|s - s_0| \leq r), \quad (6.55)$$

with an implicit constant that is absolute: for, by the residue calculus, one will then have

$$f'(s_0) = \frac{1!}{2\pi i} \int_C \frac{f(s) ds}{(s - s_0)^2}$$

when C is the (positively oriented) circle with radius r and centre at s_0 , and so, by (6.55) (again), may deduce that

$$\begin{aligned} |f'(s_0)| &\leq \frac{1}{2\pi} \int_C |ds| \cdot \max \left\{ \left| f(s) (s - s_0)^{-2} \right| : s \in \mathbb{C} \text{ and } |s - s_0| = r \right\} \\ &= r \cdot r^{-2} \cdot \max \{ |f(s)| : s \in \mathbb{C} \text{ and } |s - s_0| = r \} \\ &\ll r^{-1} \ell^{1/2} N^{-1/2} \log^{1/2} N, \end{aligned}$$

with the factor r^{-1} satisfying $r^{-1} = 4 \log x = 4(N+1)\varepsilon < 6N\varepsilon < \log N$ (the last three relations following by virtue of (1.9), (6.1), (6.3) and our assumption that $N \geq 3$).

We now work to establish (6.55) (and so complete the proof of the lemma). The bounds

$$t_0 > 8\pi\ell \geq 8\pi \quad (6.56)$$

and

$$r < \frac{1}{4N\varepsilon} < \frac{1}{4 \log \sqrt{3}} < \frac{1}{2} \quad (6.57)$$

will be helpful: they are consequences of (6.2) and (6.3). We suppose that $s \in \mathbb{C}$ satisfies the condition in (6.55). It follows that $s = \sigma + it$ for some real numbers σ, t such that one has both $r \geq |\sigma - \sigma_0| = |\sigma - 1|$ and $r \geq |t - t_0|$. Upon combining this with (6.56), (6.57) and the definition of r , we find that

$$\frac{1}{2} < 1 - \frac{1}{4 \log x} \leq \sigma \leq 1 + \frac{1}{4 \log x} < \frac{3}{2} \quad (6.58)$$

and (similarly)

$$2t_0 > t > \frac{t_0}{2} > 4\pi . \quad (6.59)$$

By (6.59), (1.9), (6.1), (1.8) and (6.5), the number $y := (2\pi x)^{-1}t = (2\pi)^{-1}\Delta^{-N-1}t$ satisfies

$$y > \frac{t_0}{4\pi x} = \frac{\ell}{2\varepsilon\Delta^{N+1}} = \left(\frac{\sinh(\varepsilon)}{\varepsilon}\right) \left(\frac{\ell}{\Delta}\right) > \frac{1}{\Delta} > \frac{3}{4} , \quad (6.60)$$

while, by (6.5), we have also $x = \Delta^{N+1} > 1$. It therefore follows, by application of the case $(H, K) = (\frac{3}{2}, \frac{3}{4})$ of the approximate functional equation obtained in [H-L], that

$$f(s) = \chi(s) \sum_{n < y} n^{s-1} + O(x^{-\sigma}) + O(y^{\sigma-1}t^{\frac{1}{2}-\sigma}) , \quad (6.61)$$

where $\chi(s) = \zeta(s)/\zeta(1-s) = 2^s\pi^{s-1} \sin(\frac{\pi}{2}s) \Gamma(1-s)$ ($\Gamma(z)$ being Euler's Gamma-function) and both implicit constants are absolute. By (6.58) we have

$$\left| \sum_{n < y} n^{s-1} \right| \leq \sum_{n < y} n^{\sigma-1} \leq 2y^\sigma$$

and may also deduce (given (6.59)) that the estimate in [Ti, (4.12.3)] implies that we have $\chi(s) \ll t^{\frac{1}{2}-\sigma}$. By these observations, combined with (6.60) and (6.61), we may conclude that $f(s) \ll x^{-\sigma} + y^\sigma t^{\frac{1}{2}-\sigma}$. Here, by (6.58), (6.5), (6.3) and (6.59), we have both

$$x^{-\sigma} \ll x^{-1} = \Delta^{-N-1} < \Delta^{-N} = 2 \sinh(\varepsilon) \ll \varepsilon$$

and

$$y^\sigma t^{\frac{1}{2}-\sigma} = t^{\frac{1}{2}}(2\pi x)^{-\sigma} \ll t_0^{\frac{1}{2}}\varepsilon = (2\pi\ell\varepsilon)^{\frac{1}{2}} ,$$

and so, given (6.3), and given that $\ell \geq 1$, we are able to verify that (6.55) does indeed hold (and thereby complete this proof) \blacksquare

Since $\sum_{\ell=1}^{\infty} \ell^{-\frac{7}{2}} = \zeta(\frac{7}{2}) < \infty$, we may deduce from the results in (6.54) and Lemma 6.8 that the term R (given by (6.48)) satisfies:

$$R = -\frac{3}{\pi^4} \sum_{\ell=1}^{\infty} \ell^{-4} \cdot \operatorname{Re}(\zeta'(1 + 2\pi\ell\varepsilon^{-1}i)) + O\left(N^{-\frac{1}{2}} \log^{\frac{3}{2}} N\right) . \quad (6.62)$$

It is known that one has $\zeta(1+it) \ll (\log t)^{\frac{2}{3}}$ and $\zeta'(1+it)/\zeta(1+it) \ll (\log t)^{\frac{2}{3}}(\log \log t)^{\frac{1}{3}}$ for $t \geq 2$ (see [Ti, Section 6.19] for a relevant discussion, including references to the papers in which these bounds were obtained). An obvious consequence of these estimates is that one has $\zeta'(1+it) \ll (\log t)^{\frac{4}{3}}(\log \log t)^{\frac{1}{3}}$ for $t \geq 2$. Although this last bound might well be considered adequate, in respect of its application in bounding the sum in (6.62), we shall nevertheless take the trouble to sharpen it very slightly, by removal of the factor $(\log \log t)^{\frac{1}{3}}$. The key to this small improvement is the estimate

$$\zeta(\sigma+it) \ll t^{\sqrt{10^4(1-\sigma)^3}} \log^{\frac{2}{3}} t \quad \left(\frac{1}{2} \leq \sigma \leq 1, t \geq 2\right), \quad (6.63)$$

which is a special case of bounds for the Hurwitz zeta-function obtained in a paper [Ri] of Richert. Note that (6.63) features in the above mentioned discussion in [Ti, Section 6.19]. The implicit constant in (6.63) is absolute (i.e. independent of both t and σ). The following is a corollary of (6.63): we state it as a lemma (requiring proof), since we were unable to find a statement of this result in the existing literature.

Lemma 6.9. *One has $\zeta'(1+it) \ll \log^{\frac{4}{3}} t$ for $t \geq 17$.*

Proof. By [Iv, Lemma 1.3], it follows that, for $1 < \sigma \leq \frac{3}{2}$ and $t \geq 1$, one has $\zeta(\sigma+it) \ll 1 + |\zeta(1+i(t+v))|$ (with an implicit constant that is absolute), for some real $v = v(\sigma, t)$ satisfying $|v| \leq \log^2 t$. Therefore, given that $0 \leq \log^2 t \leq (2/e)^2 t < \frac{3}{5}t$ for $t \geq 1$, it may be deduced from the case $\sigma = 1$ of (6.63) that one has:

$$\zeta(\sigma+it) \ll \log^{\frac{2}{3}} t \quad \left(1 < \sigma \leq \frac{3}{2}, t \geq 5\right). \quad (6.64)$$

Let $t \geq e^{\sqrt{8}}$. By an application of the residue calculus similar to that seen in the proof of Lemma 6.8, we find that $|\zeta'(1+it)| \leq \varrho^{-1} \cdot \max\{|\zeta(s)| : s \in \mathbb{C} \text{ and } |s - (1+it)| = \varrho\}$, provided only that $0 < \varrho < t$. One may, in particular, specify that $\varrho = (\log t)^{-\frac{2}{3}}$ here: note that this choice ensures that $0 < \varrho \leq \frac{1}{2}$. Then, when $s = 1+it + \varrho e^{i\theta}$ for some $\theta \in (-\frac{\pi}{2}, \frac{\pi}{2})$, it follows by (6.64) that $\zeta(s) \ll \log^{\frac{2}{3}} t = \varrho^{-1}$. When $s = 1+it + \varrho e^{i\theta}$ for some $\theta \in [\frac{\pi}{2}, \frac{3\pi}{2}]$, one finds, by (6.63), that

$$\zeta(s) \ll (2t)^{\sqrt{10^4 \varrho^3}} \log^{\frac{2}{3}}(2t) = (2t)^{100/\log t} \log^{\frac{2}{3}}(2t) \leq 2^{25\sqrt{2}} e^{100} (2 \log t)^{\frac{2}{3}}.$$

The last two observations show that $\max\{|\zeta(s)| : s \in \mathbb{C} \text{ and } |s - (1+it)| = \varrho\} \ll \varrho^{-1}$, so that (by the application of the residue calculus mentioned earlier) one must have $|\zeta'(1+it)| \ll \varrho^{-2} = \log^{\frac{4}{3}} t$: as $e^{\sqrt{8}} < 17$, the lemma has been proved \blacksquare

Remark. If the Riemann Hypothesis is true then the upper bound in Lemma 6.9 can be improved to $O((\log \log t)^2)$: this would follow from certain conditional estimates for $\log \zeta(s)$ that are mentioned in [Ti, Section 14.33]. Based on preliminary calculations, we believe it therefore follows that the result stated in Theorem 4 could, on the Riemann Hypothesis, be refined to:

$$\begin{aligned} \|K^{\triangleleft, N}\|^2 &\leq \left(\frac{7}{60}(\varepsilon N)^3 + \frac{1}{12}(\varepsilon N)^2 + O((\log \log N)^2 \varepsilon N)\right) N^{-1} \\ &\leq \frac{7}{60} \left(1 - \frac{3 \log \log (N^\beta)}{\log N} + O\left(\left(\frac{\log \log N}{\log N}\right)^2\right)\right) N^{-1} \log^3 N, \end{aligned}$$

with $\beta = 2e^{-\frac{5}{21}}$. We omit the details of those calculations, except to mention that they involve asymptotic estimation of the contribution of the term $\psi(\sigma_n)\varepsilon^3 e^{(1+\sigma_n)\varepsilon}$ occurring in the statement of Lemma 6.6, rather than the use of the bound $\psi(\sigma_n)\varepsilon^3 e^{(1+\sigma_n)\varepsilon} \leq 0$.

The proof of Theorem 4. By (6.9), we have $\|K^\infty\|^2 = V_0 + V_1$, where, by Lemmas 6.3 and 6.7, and (1.8), (1.9), (6.1), (6.2) and (6.3), one has: $V_1 \leq \frac{1}{6}(N\varepsilon + 1) \sinh(\varepsilon) + \frac{2}{9\sqrt{3}}N\varepsilon \sinh^2(\varepsilon) \ll N\varepsilon^2 < N^{-1} \log^2 N$ and $V_0 \leq \frac{7}{60}N^{-1} \log^3 N + O(N^{-1} \log^2 N) + \frac{1}{2}\varepsilon R$, with R as defined in (6.48). Thus

$$\|K^\infty\|^2 \leq \frac{7}{60}N^{-1} \log^3 N + O(N^{-1} \log^2 N) + O(\varepsilon|R|) . \quad (6.65)$$

We have shown (above) that R satisfies (6.62). In each term of the sum that occurs in (6.62) one has $2\pi\ell\varepsilon^{-1} > 8\pi\ell \geq 8\pi$ (by virtue of (6.3) and the fact that the summation in (6.62) is over $\ell \in \mathbb{N}$). We are therefore able to apply the result of Lemma 6.9 to each term of the sum in (6.62), and find (thereby) that

$$\begin{aligned} |R| &\leq \frac{3}{\pi^4} \sum_{\ell=1}^{\infty} \ell^{-4} \cdot O\left(\log^{\frac{4}{3}}(2\pi\ell\varepsilon^{-1})\right) + O\left(N^{-\frac{1}{2}} \log^{\frac{3}{2}} N\right) \\ &\ll 1 + \sum_{\ell \leq 2\pi/\varepsilon} \ell^{-4} \log^{\frac{4}{3}}(4\pi^2\varepsilon^{-2}) + \sum_{\ell > 2\pi/\varepsilon} \ell^{-4} \log^{\frac{4}{3}}(\ell^2) \\ &\leq 1 + 2^{\frac{4}{3}}\zeta(4) \cdot \log^{\frac{4}{3}}(2\pi\varepsilon^{-1}) + 2^{\frac{4}{3}} \sum_{\ell=2}^{\infty} \ell^{-4} \log^{\frac{4}{3}} \ell \ll \log^{\frac{4}{3}}(\varepsilon^{-1}) . \end{aligned}$$

By this and (6.3) we have, in the second O -term of (6.65),

$$\varepsilon|R| \ll \varepsilon \log^{\frac{4}{3}}(\varepsilon^{-1}) \ll \left(\frac{\log N}{N}\right) \cdot \log^{\frac{4}{3}}\left(\frac{N}{\log N}\right) \ll N^{-1} \log^{\frac{7}{3}} N$$

(using that $N \geq 3$), and so we obtain what the theorem states \blacksquare

§7. Numerical Results.

As already noted (in the paragraph below the proof of Lemma 6.6) Lemmas 6.3 and 6.6 may be used in computing, for a chosen integer N , a numerical upper bound on $\|K^{\infty, N}\|$. Lemma 6.5, on the other hand, enables one to compute the numerical values of the elements of the corresponding Hankel matrix $H(N)$, which (by (1.19) and (1.20)) has as its non-zero eigenvalues the reciprocal eigenvalues of the kernel $K^{\leftarrow, N} = K - K^{\infty, N}$. Thus, given a suitable machine (and software including the necessary algorithms), one can compute good numerical approximations to the reciprocal eigenvalues of $K^{\leftarrow, N}$: such approximations, when combined with a numerical upper bound on $\|K^{\infty, N}\|$, lead (via (3.6)) to numerical bounds (both upper and lower) for the corresponding reciprocal eigenvalues of K .

In putting the above observations into practice, we have made use of the ‘GNU Octave’ software, installed on a desktop computer. We found that, when $N = 10^6$ and $K^\infty = K^{\infty, N}$, one has $V_0 < 0.000135821302$ and $V_1 < 0.000021043522$, so that it follows by (6.9) that

$$\|K^\infty\| = \sqrt{V_0 + V_1} < 0.012524569 . \quad (7.1)$$

We also computed (again, for $N = 10^6$) the following table of approximations to the twelve largest (in absolute value) of the reciprocal eigenvalues of the kernel $K^\triangleleft = K^{\triangleleft, N}$.

k	ν_{k-}^\triangleleft	ν_{k+}^\triangleleft
1	-0.0690165028	0.0796889109
2	-0.0566761531	0.0690322925
3	-0.0506170103	0.0552247168
4	-0.0441382847	0.0458979695
5	-0.0417049532	0.0440302591
6	-0.0370487294	0.0377564244

Although we think it likely that the entries in this table are accurate to at least 8 decimal places, we admit to some uncertainty on this point, as we have not given due consideration to, for example, the possible effect of accumulated rounding errors (not to mention other potential sources of error). Assuming that these approximations are indeed accurate to 8 decimal places, it follows by (3.6), (7.1) and the first row of data in the above table that one has $0.0815411 > -\nu_{1-} > 0.0564919$ and $0.0922135 > \nu_{1+} > 0.0671643$. Upon comparing these bounds with the second row of data in the above table, we are forced to concede that, even if it were reliable, the numerical data presented here would seem to be insufficient to establish (for certain) that one does not have both $\nu_{2+} = \nu_{1+}$ and $\nu_{2-} = \nu_{1-}$.

We can, at least, make slightly more efficient use of our data by observing that it follows by Lemmas 3.1, the Cauchy-Schwarz inequality and Lemma 2.1 that one has:

$$\begin{aligned} \nu_{1+} - \nu_{3+} &\geq (\nu_{1+}^\triangleleft + \nu_{1-}^\boxtimes) - (\nu_{3+}^\triangleleft + \nu_{1+}^\boxtimes) \\ &\geq \nu_{1+}^\triangleleft - \nu_{3+}^\triangleleft - 2^{\frac{1}{2}} \left((\nu_{1-}^\boxtimes)^2 + (\nu_{1+}^\boxtimes)^2 \right)^{\frac{1}{2}} \\ &\geq \nu_{1+}^\triangleleft - \nu_{3+}^\triangleleft - 2^{\frac{1}{2}} \|K^\boxtimes\| . \end{aligned}$$

By this, along with (7.1) and the data from the above table, we arrive at the bound $\nu_{1+} - \nu_{3+} > 0.0067517$. We find (similarly) that $|\nu_{1-}| - |\nu_{3-}| > 0.0006870$. It therefore would appear that each one of the numbers $\lambda_{1+} = 1/\nu_{1+}$, $\lambda_{1-} = 1/\nu_{1-}$ is an eigenvalue (of K) that has index less than or equal to 2: if this is not so, then our computed approximations to the relevant eigenvalues of K^\triangleleft must be disappointingly inaccurate.

The results of the numerical computations just described could certainly be improved by means of similar computations in respect of a significantly increased value of N . However, our attempts to carry out such computations have essentially failed. This is due partly to limitations on the precision with which arithmetic operations are carried out in GNU Octave, and partly to limitations on working memory (when running Octave under a Windows operating system). Taking a different approach, one may instead seek to utilise the information on eigenvectors of the matrix $H(N)$ that one obtains as a by-product of the computation of the eigenvalues of that matrix. We have had some success with this second approach (without needing to consider any $N > 10^6$). The present paper, however, is not the right place for a full discussion of this matter, as such a discussion would (ideally)

have to extend to questions concerning the degree of accuracy of certain computed approximations to eigenfunctions of K : rather than lengthen the present paper any further, we reserve that discussion for another occasion.

References.

- [C-R] R.R. Coifman and R. Rochberg, ‘Representation theorems for holomorphic and harmonic functions in L^p ’, *Asterisque* **77** (1980), 11-66.
- [H-L] G.H. Hardy and J.E. Littlewood, ‘The approximate functional equation in the theory of the zeta-function, with applications to the divisor problems of Dirichlet and Piltz’, *Proc. London Math. Soc.* (2), **21** (1922), 39-74.
- [HW] M.N. Huxley and N. Watt, ‘Mertens sums requiring fewer values of the Möbius function’, preprint (to appear).
- [Iv] A. Ivić, *The Riemann zeta-function: theory and applications*, Dover Publications, Inc., Mineola, New York (2003); originally published by Wiley, New York (1985).
- [Li] B.W. Lindgren, *Statistical Theory* (2nd edition), The Macmillan Company, New York (1968).
- [OLBC] F.W.J. Olver, D.W.L. Lozier, R.F. Boisvert, C.W. Clark (editors), *NIST Handbook of Mathematical Functions*, Cambridge University Press, New York 2010.
- [Pa] J.R. Partington, *An Introduction to Hankel Operators*, London Math. Soc. Student Texts **13**, Cambridge University Press, Cambridge (1988).
- [Pe] V.V. Peller, ‘Hankel operators of class C_p and their applications (rational approximation, Gaussian processes, the problem of majorizing operators)’, *Math. USSR Sbornik* **41** (1982), No. 4, 443-479.
- [Ri] H.-E. Richert, ‘Zur Abschätzung der Riemannschen Zetafunktion in der Nähe der Vertikalen $\sigma = 1$ ’, *Math. Ann.* **169** (1967), 97-101.
- [Sch] E. Schmidt, ‘Zur Theorie der linearen und nicht linearen Integralgleichungen. I. Teil: Entwicklung willkürlicher Funktionen nach Systemen vorgeschriebener’, *Math. Ann.* **63** (1907), 433-476.
- [Ti] E.C. Titchmarsh (revised by D.R. Heath-Brown), *The Theory of the Riemann Zeta-function*, Oxford Univ. Press, 1986
- [Tr] F.G. Tricomi, *Integral Equations*, Dover Publications, New York (1985).
- [Wa] N. Watt, ‘On eigenvalues of the kernel $\frac{1}{2} + \lfloor \frac{1}{xy} \rfloor - \frac{1}{xy}$ ($0 < x, y \leq 1$)’, preprint; arXiv:1811:05246v1 [math.NT].
- [Wa2] N. Watt, ‘On eigenfunctions of the kernel $\frac{1}{2} + \lfloor \frac{1}{xy} \rfloor - \frac{1}{xy}$ ($0 < x, y \leq 1$)’, preprint (in preparation).
- [We] H. Weyl, ‘Ueber die asymptotische Verteilung der Eigenwerte’, *Nachrichten von der Gesellschaft der Wissenschaften zu Göttingen, Mathematisch-Physikalische Klasse* (1911), pages 110-117