

# CLASSIFYING RATIONAL PARALLELEPIPEDS

RANDALL L. RATHBUN

ABSTRACT. It is proposed that the name ‘*Diophantine parallelepiped*’ be applied to rational parallelepipeds. By examining the count of possible right angles  $\perp$  at any vertice, the parallelepipeds become classified into 5 types: *acute anorthogonal triclinic*; *obtuse anorthogonal triclinic*; *monorthogonal bi-clinic*; *bi-orthogonal monoclinic*; and *rectangular cuboid*; There is sufficient confusion in the present literature to warrant an attempt at clarity for these types of pipeds and to offer a classification scheme.

The possible rational components of the Diophantine parallelepiped are examined and some results from computer searches are also presented.

## 1. INTRODUCTION TO DIOPHANTINE PARALLELEPIPEDS

Parallelepipeds have been examined, in Diophantine analysis of Number Theory, but a lack of clarity still seems to exist.

We define a rational parallelepiped in  $n$ -dimensions as a polytope spanned by  $n$  vectors  $\vec{v}_1, \dots, \vec{v}_n$  in a vector space over the rationals,  $\mathbb{Q}$ , or integers,  $\mathbb{Z}$

$$\text{span}(\vec{v}_1, \dots, \vec{v}_n) = t_1\vec{v}_1 + \dots + t_n\vec{v}_n \text{ for } t_i \in \mathbb{Q} \text{ for } i = 1 \rightarrow n$$

Here we are interested in three dimensions,  $n = 3$ , so a rational parallelepiped is a prism determined by the 3 rational basis vectors  $\vec{a}$ ,  $\vec{b}$ ,  $\vec{c}$ . The prism has 8 vertices, 3 pairs of parallel faces which are all parallelograms, and 12 edges (3 distinct)  $\in \mathbb{Q}$ .

$$(1) \quad \text{span}(\vec{a}, \vec{b}, \vec{c}) = t_1\vec{a} + t_2\vec{b} + t_3\vec{c} \text{ for } t_{i=1 \rightarrow 3} \in \mathbb{Q}$$




FIGURE 1. The rational parallelepiped with 3 rational basis vectors  $\vec{a}$ ,  $\vec{b}$ , and  $\vec{c}$

In Figure 1, the lengths of the basis vectors,  $\vec{a}$ ,  $\vec{b}$ , and  $\vec{c}$ , are rational  $\in \mathbb{Q}$ , and they determine the parallelepiped.

## 2. CLASSIFYING THE PARALLELEPIPEDS BY ORTHOGONALITY

Literature abounds with examples confusing rectangular parallelepipeds with monoclinic pipeds. The bi-clinic or mon-orthogonal case is hardly even known. The rectangular parallelepiped is called the ‘*cuboid*’,

---

*Date:* November 27, 2018.

*2010 Mathematics Subject Classification.* 11D09.

*Key words and phrases.* integer cuboid, rectangular cuboid, perfect parallelepiped.

or ‘*brick*’, or ‘*Euler brick*’, or even the ‘*orthotope*’, engendering confusion over the preferred name for the same object.

In an attempt at clarity for Diophantine analysis, this author proposes a classification scheme based upon the maximum count of right angles  $\perp$  which might occur at any of the 8 vertices in the Diophantine parallelepiped.

Consider the 3 surface angles from Figure 1.,  $\alpha$  between  $\vec{a}$  and  $\vec{b}$ ;  $\beta$  between  $\vec{a}$  and  $\vec{c}$ ; and  $\gamma$  between  $\vec{b}$  and  $\vec{c}$ . There may be none, or 1, or 2, or 3  $\perp$  right angles at a vertex of the Diophantine parallelepiped. All 8 vertices need to be considered.

### 3. FIVE CLASSES OF RATIONAL DIOPHANTINE PARALLELEPIPEDS

There are 5 unique classes of parallelepipeds which do exist, depending upon the count of right angles at the origin [the intersection of the basis vectors], as shown and proved in Appendix A.

TABLE 1. The five classes of rational parallelepipeds by surface angles at a vertice.

Name	Comment	Vertex Group(s)
acute	(anorthic) [no $\perp$ ]	$[-1 \ -1 \ 1]$ [ <b>1 1 1</b> ]
obtuse	(anorthic) [no $\perp$ ]	$[-1 \ -1 \ -1]$ $[-1 \ 1 \ 1]$
1-ortho	(bi-clinic)	$[-1 \ -1 \ 0]$ $[-1 \ 0 \ 1]$ [ <b>0 1 1</b> ]
2-ortho	(monoclinic)	$[-1 \ 0 \ 0]$ [ <b>0 0 1</b> ]
3-ortho	(rectangular)	[ <b>0 0 0</b> ]

In the table above, the classifier for the vertice group is shown in the Vertex Group(s) column. The meaning is as follows: 1 = acute  $\angle$  angle; 0 = right  $\perp$  angle; and  $-1$  = obtuse  $\sphericalangle$  angle. The classifier is color highlighted for the right  $\perp$  angle(s).

NOTE: The vertice group gives the actual types of vertices which actually occur in the class.

This suggests a possible classification scheme with 5 categories or class names:

- the Diophantine acute anorthic parallelepiped (*acute triclinic*)
- the Diophantine obtuse anorthic parallelepiped (*obtuse triclinic*)
- the Diophantine 1-ortho parallelepiped (*biclinic*)
- the Diophantine 2-ortho parallelepiped (*monoclinic*)
- the Diophantine 3-ortho parallelepiped (*rectangular*)

Please note that for the first two classes, the *acute anorthic*, and the *obtuse anorthic*, that no right angle  $\perp$  exists at any vertex of the parallelepiped.

The labeling of the pipeds using the terms anorthic, or triclinic, the relatively unknown biclinic and the well known monoclinic is deliberate, and comes from the mineral crystallography classification scheme, since the crystals systems share some of the same morphology as the Diophantine pipeds.

Thus the proposed classification scheme then depends upon whether or not (0,1,2,3) right angles exist for at least one of the eight vertices of the piped (actually at the origin or intersection of the basis vectors creating the piped).

### 4. RATIONAL PARALLELEPIPEDS AND THEIR COMPONENTS

We need to take a careful look at the rational components of the Diophantine piped for Diophantine analysis. Due to the extensive symmetry and parallelism of the parallelepiped figure, many of its components are congruent, thus insuring rationality for all elements in the same congruent group, such as the edges, the face diagonals, the skew triangles, the face or body triangles, and the parallelogram areas.

TABLE 2. Diophantine parallelepiped classification scheme into 5 classes.

proposed name	<i>acute</i>	<i>obtuse</i>	<i>1-ortho</i>	<i>2-ortho</i>	<i>rectangular</i>
common name	triclinic	triclinic	parallelepiped	monoclinic	cuboid
right angle(s) $\perp$	0	0	1	2	3

4.1. **Derivation from 3 rational or integer basis-vectors.** Again, as in eq(1) we first start with 3 finite vectors which span  $\mathbb{R}^3$  as shown in figure 2. Let their magnitudes be rational  $\in \mathbb{Q}$  or preferably as integer  $\in (\mathbb{Z})$ .




FIGURE 2. The three basis vectors,  $\vec{v}_1, \vec{v}_2, \vec{v}_3$ . with magnitudes  $a, b, c \in \mathbb{Z}$

Call the integer lengths of these vectors  $a, b$ , and  $c$  respectively. Using these 3 vectors we can create 3 unique parallelograms, using a pair of the 3 vectors, and then combine the 3 parallelograms into a parallelepiped.

Where 3 parallelograms exist, with their sides of length  $[a, b]$ , and  $[a, c]$ , and  $[b, c]$ , then a parallelepiped solution exists, where  $a, b$ , and  $c$  are the magnitudes of the 3 basis vectors. We label the 3 pairs of the diagonals of the 3 parallelograms as  $d, e; f, g$ ; and  $h, j$ .

parallelogram	sides	diagonals
1	$a, b$	$d, e$
2	$a, c$	$f, g$
3	$b, c$	$h, j$

TABLE 3. 3-basis vector solution  $a, b, c$  for 3 parallelograms with matching sides

4.2. **The Parallelogram equation and rational parallelograms.** The parallelogram equation (2) shows that we can pick 3 values of a parallelogram to be rational  $\in \mathbb{Q}$ .

$$(2) \quad 2(a^2 + b^2) = c^2 + d^2$$

where  $a, b$  are the sides and  $c, d$  are the diagonals of the parallelogram. We can pick 3 rational values for a parallelogram, satisfying equation (2), say sides  $a, b$  and the diagonal  $c$ , then the other diagonal  $d$  may or may not be rational. If the second diagonal is rational, we call the parallelogram the *rational* or *integer parallelogram* (upon scaling).

It is noted that these integer parallelograms are created from integer triangles, so every parallelogram has at least 1 rational diagonal, and if a parallelepiped is constructed from 3 such triangles, then at least 3 diagonals, say  $d, f$ , and  $h$ , from Table 3 are rational, while the remaining three  $e, g$ , and  $j$  may or may not be rational.

4.3. **Locating coordinates in  $\mathbb{R}^3$  for permuted tetrahedrons of a piped.** Using these 3 parallelograms, which have at least 1 rational diagonal each, 48 tetrahedrons can be created from the permutations, as 6 sets of 8 tetrahedrons. However only 1 set of the 8 tetrahedrons is necessary to completely specify all 6 sets.

A convenient and unique collection covering the 8 possible arrangements of a convenient set is shown in Table 4 below.

TABLE 4. Combinations of the sides/diagonals of 3 Parallelograms for a Tetrahedron

Tetrahedron Sides			
a d b f c h	a d b f c j	a d b g c h	a d b g c j
a e b f c h	a e b f c j	a e b g c h	a e b g c j

Depending upon whether or not the second diagonal(s) is rational or not, we can create an *integer tetrahedron*, in at least 1 case of the 8 given above.

The reason we deal with *integer tetrahedrons* is that they provide us with the 6 key lengths:  $a, b, c, d, e, f$ ; which enable us to fix the vertices of the Parallelepiped into  $\mathbb{R}^3$  Cartesian space.

An *integer tetrahedron* is shown below in figure 3 with 6 integer (or rational) sides,  $a, b, c, d, e, \& f$ . These sides are slightly relabeled from the set of 8 previously given so that the equations (3, 4) below giving the lattice locations are correct.

You can quickly recognize the 3 vectors  $\vec{v}_1, \vec{v}_2$  and  $\vec{v}_3$  creating edges  $a, c$ , and  $e$  in the figure:




FIGURE 3. Integer tetrahedron.

We have to locate the vertices of this tetrahedron, given the lengths, into  $\mathbb{R}^3$ , and I choose the following method to fix the vertices  $v_1, v_2, v_3$ , and  $v_4$  into Cartesian 3D space. We find the 3 vertices first.

$$\begin{aligned}
 &\text{let tetrahedron edges be } a, b, c, d, e, f \in \mathbb{Z} \\
 &v_1 = [0, 0, 0] \\
 &v_2 = [a, 0, 0] \\
 (3) \quad &\text{let } \cos(B) = \frac{a^2 + c^2 - b^2}{2ac} \\
 &\text{and } \sin(B) = \frac{\sqrt{(a+b+c)(a+b-c)(a+c-b)(b+c-a)}}{2ac} \\
 &v_3 = c[\cos(B), \sin(B), 0]
 \end{aligned}$$

By using the intersections of three  $S^2$  spheres from the 3 known vertices  $[0, 0, 0]$ ,  $v_1$ , and  $v_2$  with the fourth vertex  $v_4$ , that vertex can be located.

$$\begin{aligned}
 x &= \frac{a^2 + e^2 - d^2}{2a} \\
 y &= \frac{c^2 + e^2 - f^2 - 2rx}{2s} \\
 \text{where } r &= c \cdot \cos(B) \\
 \text{and } s &= c \cdot \sin(B) \text{ both from } v_3 \\
 \text{then } z &= \pm \sqrt{e^2 - x^2 - y^2} \\
 v_4 &= [x, y, z]
 \end{aligned}
 \tag{4}$$

The  $+z$  value was used (although  $-z$  is another solution).

Most of the time, the vertices are located in a quadratic field  $K = \mathbf{Q}(\sqrt{D})$ , occasionally they can be on a rational lattice.

We have to follow the same process in equations (3),(4) for all 8 *integer tetrahedrons* in Table 4. The parallelepiped (1 of 8) is derived from these 3 vectors and we strictly label the vertices as shown and follow this order for the rest of this paper.




FIGURE 4. The labeled Diophantine parallelepiped.

This labeled parallelepiped contains several (possibly rational) subcomponents, and they are examined in the following §§.

**4.4. The 28 Diophantine Piped Diagonals.** The first subcomponent to be considered are the diagonals, a line segment between 2 vertices. There are 28 diagonals in the Diophantine piped, the parallelogram in which they reside is given also, if it exists.

TABLE 5. 28 Diagonals in the Diophantine piped with parallelograms

12 Face Diagonals				$\square$	$\square$
1-4	2-3	5-8	6-7	F1234	F5678
1-6	2-5	3-8	4-7	F1256	F3478
1-7	3-5	2-8	4-6	F1357	F2468
4 Body Diagonals				$\square$	$\square$
1-8	3-6	2-7	4-5	B1368	B2457
				B1278	B3456
				B1458	B2367

12 edges				$\square$	$\square$
1-2	3-4	5-6	7-8	F1234	F5678
1-3	2-4	5-7	6-8	F1234	F5678
1-5	2-6	3-7	4-8	F1256	F3478

There are 6 face parallelograms containing 12 diagonals, and 6 body parallelograms with 4 unique diagonals, since they share the same set of 4 diagonals in a permuted manner.

**4.5. The 56 Diophantine Piped Triangles.** Triangles are composed of 3 connected diagonals, or any 3 vertices. There are 56 triangles to consider in the Diophantine piped. However only 48 reside on the face or body parallelograms.

TABLE 6. 48 Diophantine Piped Triangles on Parallelograms.

$\triangle$	$\triangle$	$\square$	$\triangle$	$\triangle$	$\square$
123	234	F1234	567	678	F5678
124	134	F1234	568	578	F5678
125	256	F1256	347	478	F3478
126	156	F1256	378	348	F3478
135	357	F1357	246	468	F2468
137	157	F1357	248	268	F2468
127	278	B1278	345	456	B3456
128	178	B1278	346	356	B3456
136	368	B1368	245	457	B2457
138	168	B1368	247	257	B2457
148	158	B1458	236	367	B2367
145	458	B1458	237	267	B2367

There are also 8 triangles unaccounted for. These are not found on the parallelogram components of the Diophantine piped. I call them the *'skew triangles'* as they are not located on any of the 12 parallelograms.

The 8 skew non-parallelogram triangles are

TABLE 7. 8 Skew Triangles

$\triangle$	$\triangle$	$\triangle$	$\triangle$
146	147	167	467
853	852	832	532

The matching congruent triangles listed above are given in a column, preserving the isometry of the lengths of any two vertices of the triangle, following the order of the vertice labeling.

**4.6. The 12 Diophantine Piped Parallelograms.** The Diophantine piped also contains 12 parallelograms.

TABLE 8. 12 Parallelograms in the Piped.

face parallelograms						
$\square$			$\square$			$\square$
F1234			F1357			F1256
F5678			F2468			F3478
body parallelograms						
$\square$	$\square$	$\square$	$\square$	$\square$	$\square$	
B1278	B1368	B1458	B2367	B2457	B3456	

The face parallelograms given above in a column in Table 8. are congruent, and have the same area and diagonal lengths. However please note that the 6 body parallelograms have distinct areas.

**4.7. The Diophantine Parallelepiped Itself.** Finally there is the parallelepiped itself, which has 8 ordered vertices such that there is a pair of 4 vertices which are co-planar and parallel. This figure is conventionally called the prism.

There is one component to consider for rationality, the volume:

$$\frac{\text{parallelepiped volume}}{P12345678}$$

**4.8. Summarizing the Diophantine Analysis for all 83 Components.** Considering all the components and the duplications of lengths which occur in the piped, there is a unique signature which can be checked for rationality:

**Rational Signature Considerations**

- 3 edge lengths (already rational)
- 6 face diagonals (3 pairs matches lengths)
- 4 body diagonals
- 56 triangle areas\* (48 subsumed in parallelograms, 8 are not)
- 4 skew triangle areas
- 3 face parallelograms diagonals and area
- 6 body parallelograms diagonals and area
- 1 piped volume

We don't actually check all 56 triangles\*, but let the 48 triangles on the parallelograms be subsumed by the rationality checks upon the body or face parallelograms. However, we must check 4 of the 8 skew triangles for rational area.

Additionally, the volume definitely needs to be checked, as it was discovered by computer testing that certain combinations of  $a, b, c, d, e,$  and  $f$  for the integer tetrahedron edges actually led to degenerate 'flat' tetrahedrons where all three vectors  $v_1, v_2, v_3$  were co-planer, or spanning only  $\mathbb{R}^2$ .

Putting all this together results in the unique rational check signature for a given Diophantine parallelepiped.

**4.9. Unique Rationality Check of the Diophantine Piped.** We discover that not all 83 components need to be checked for rationality. Since there are congruences between the lengths of the edges and diagonals, and congruences between the areas of the triangles, face parallelograms, and body parallelograms then not all need to be checked. In fact:

- 27 components need to be checked for rationality

We provide an example of this signature for three sample pipeds, where 0 denotes an irrational value and 1 a rational value.

TABLE 9. Three sample Diophantine parallelepipeds

$v_1$	Basis Vectors			Tetrahedron Sides					
	$v_2$	$v_3$		a	b	c	d	e	f
(103,0,0)	$(\frac{-3179}{103}, \frac{120}{103}\sqrt{17458}, 0)$	$(\frac{-3179}{103}, \frac{-2464734}{4495435}\sqrt{17458}, \frac{1122}{43645}\sqrt{27915342})$		103	204	157	204	157	264
(10,0,0)	$(-5, 6\sqrt{14}, 0)$	$(0, \frac{-24}{7}\sqrt{14}, \frac{24}{7}\sqrt{35})$		10	27	23	26	24	41
(44,0,0)	$(0, 117, 0)$	$(0, 0, 240)$		44	125	117	244	240	267

TABLE 10. 27 rational component checks for the three sample Diophantine pipeds

edges	skew triangles	face diagonals	body diagonals	face area	body area	vol
$\begin{vmatrix} 1 & 1 & 1 \\ 1 & 1 & 1 \\ 1 & 1 & 1 \end{vmatrix}$	$\begin{vmatrix} 1 & 0 & 1 & 0 \\ 1 & 0 & 0 & 0 \\ 1 & 0 & 0 & 0 \end{vmatrix}$	$\begin{vmatrix} 1 & 1 & 1 & 1 & 1 & 1 \\ 1 & 1 & 1 & 1 & 1 & 1 \\ 1 & 1 & 1 & 1 & 1 & 1 \end{vmatrix}$	$\begin{vmatrix} 0 & 0 & 0 & 0 \\ 1 & 1 & 1 & 0 \\ 0 & 0 & 0 & 0 \end{vmatrix}$	$\begin{vmatrix} 0 & 0 & 1 \\ 1 & 0 & 0 \\ 1 & 1 & 1 \end{vmatrix}$	$\begin{vmatrix} 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 \\ 1 & 1 & 1 & 1 & 1 & 1 \end{vmatrix}$	$\begin{vmatrix} 0 \\ 0 \\ 1 \end{vmatrix}$

## 5. RATIONAL CONSTRAINTS FOR DIOPHANTINE OBJECTS

There is an geometric increase in the rationality requirements as the embedding dimension of the geometric object increases. For this reason, it is necessary to examine both rational triangles and parallelograms, before considering *Diophantine parallelepipeds*, in order to better understand those requirements.

**5.1. Rational Triangles.** In Diophantine analysis of Number Theory, we can study triangles, for example, for rational components. Since the triangle is 3 sided polygon, (it is actually determined by 2 non-linear vectors), the following properties need to be checked for rationality.

### Triangle Constraints

- rational sides (3 sides)
- rational area (1 area)

This means that  $2^4 = 16$  possibilities exist, when considering the rationality of the triangle. But collectively all 3 sides are considered rational and we definitely are interested if the area is rational. In fact triangles with this property have been given a special name: *'Heron triangles'*.

We won't consider internal parts of the triangle, such as the medians, or the angle bisectors, or the perpendicular bisectors, or other civians, for example, although their study in Diophantine analysis is important. We are simply making the point here that we have 2 components to consider for rationality: the *sides* and the *area*.

If we force the triangle to have 3 integer sides,  $a, b, c \in \mathbb{Q}$ , then we only have to consider the area  $\in \mathbb{Q}$ .

The answer to the question, *can we find a triangle with integer sides and rational area?* has been known in antiquity, the 3,4,5 right triangle with area 6 is the classic answer.

Less well known is the fact that the primitive 5,5,6 isosceles Heron triangle, and the primitive 5,5,8 isosceles Heron triangle, both with area 12, are the next smallest integer triangles with rational area. In figure 5 below, note that the 5-5-6 Heron triangle was created from joining two 3-4-5 right triangles along the side length 4.

The next 2 triangles are the composite 6,8,10 right triangle with area 24, and the primitive Heron triangle 9,10,17 with area 36 which is not isosceles. It should be pointed out that if the triangle sides  $\in \mathbb{Z}$ , and the






FIGURE 5. The 3,4,5 Right Triangle and 5,5,6 isosceles Heron Triangle

triangle has rational area, then its area  $\in \mathbb{Z}$  also. If the sides are fractional  $\in \mathbb{Q}$ , and if the area is rational, then the area  $\in \mathbb{Q}$ .

Of note is the Heron formula for area of a triangles with sides  $a$ ,  $b$ , and  $c$

$$(5) \quad A = \frac{1}{4} \sqrt{(a+b+c)(-a+b+c)(a-b+c)(a+b-c)}$$

which is used to help discover rational area for the triangle in question. These triangles can be parameterized, see §5.7.

**5.2. Rational Parallelograms.** We now consider parallelograms. They are 4 sided planar figures, with 2 pairs of congruent parallel edges;  $(a, c)$ ,  $a = c$  and  $a \parallel c$  and  $(b, d)$ ,  $b = d$  and  $b \parallel d$ . They also can be determined by two non-linear vectors but there are 2 additional considerations to be added to the Diophantine analysis.

The parallelogram has the following properties to check for rationality.

**Parallelogram Constraints**

- rational sides (4 sides)
- rational diagonals (2 diagonals)
- rational area (1 area)

This means that we have to consider any one of the  $2^7 = 128$  permutations of the parallelograms properties for rationality. Let the parallelogram have sides  $(a,c)$ ,  $(b,d)$ , with diagonals  $d_1, d_2$  and area  $A$ . Considering the pairs of sides as a unit, we then have only 32 ( $2^5$ ) combinations to consider. But forcing both pairs of sides to be rational further shortens down the 128 possibilities to just 8 cases.

TABLE 11. Possible rationality checks for a parallelogram

parallelogram	component	$(a, c)$	$(b, d)$	$d_1$	$d_2$	$A$
case 1	rational ?	✓	✓	-	-	-
case 2	rational ?	✓	✓	✓	-	-
case 3	rational ?	✓	✓	-	✓	-
case 4	rational ?	✓	✓	✓	✓	-
case 5	rational ?	✓	✓	-	-	✓
case 6	rational ?	✓	✓	✓	-	✓
case 7	rational ?	✓	✓	-	✓	✓
case 8	rational ?	✓	✓	✓	✓	✓

From examination, it can be seen that case 3 is identical to case 2 and case 7 is identical to case 6, because order is not important, so this narrows our possibilities to just 6 actual cases:

TABLE 12. Actual rationality checks for the 6 cases of a parallelogram

parallelogram	component	$(a, c)$	$(b, d)$	$d_1$	$d_2$	$A$
case 1	rational ?	✓	✓	–	–	–
case 2	rational ?	✓	✓	✓	–	–
case 3	rational ?	✓	✓	✓	✓	–
case 4	rational ?	✓	✓	–	–	✓
case 5	rational ?	✓	✓	✓	–	✓
case 6	rational ?	✓	✓	✓	✓	✓

Mathematicians would be interested in case 2 and definitely in case 3. In fact case 3 has been recently parameterized[8, 9] (see §5.7). They might consider case 4 an annoyance, to be avoided. Case 5 might be momentarily considered, but then abandoned because it has 1 irrational diagonal. So this leaves case 6 as the one which would definitely be considered, since it has all components rational.

I have purposefully gone through this exercise for both triangles and parallelograms, to emphasize that the possible rational cases have to be carefully delineated, when studying the Diophantine object in question. Sorry to say, sometimes this careful analysis is often overlooked, particularly when considering rational parallelepipeds.

As you can imagine, the rational components increase geometrically, when the embedding dimension containing the object increments, because the figure becomes more complex, and it become important to account for all the possibilities.

**5.3. Some examples of the smallest integer parallelograms.** The following parallelograms are the smallest integer parallelograms that exist (the next 2 tables list the parallelograms two to a row).

TABLE 13. Case 3. First 10 parallelograms with 2 rational diagonals.

sides		diags		sides		diags	
3	4	5	5	4	7	7	9
5	5	6	8	5	10	9	13
5	12	13	13	6	7	7	11
6	8	10	10	6	13	11	17
6	17	17	19	7	9	8	14

TABLE 14. Case 6. First 10 parallelograms with 2 diagonals rational and area rational.

sides		diags		sides		diags	
3	4	5	5	5	5	6	8
5	12	13	13	6	8	10	10
7	24	25	25	8	15	17	17
9	12	15	15	9	40	41	41
10	10	12	16	10	24	26	26

Please note that if the 2 diagonals are equal, the parallelogram is a rectangle or square. There are some parallelograms in common with both lists.

TABLE 15. Parallelograms in common - 4 right triangles - 1 Heron triangle.

sides		diags		sides		diags	
3	4	5	5	5	12	13	13
5	5	6	8	6	8	10	10
7	24	25	25				

5.4. **Some examples of rational tetrahedrons.** From these smallest integer parallelograms, come the smallest integer tetrahedrons. A computer search quickly discovers the following integer tetrahedrons.

TABLE 16. The first 40 smallest integer tetrahedrons.

tetrahedron edges						tetrahedrons edges					
5	6	5	9	10	13	5	6	5	9	10	9
5	6	5	13	10	13	5	6	5	13	10	9
5	6	5	13	12	13	5	8	5	9	10	13
5	8	5	9	10	9	5	8	5	13	10	13
5	8	5	13	10	9	5	8	5	13	12	13
7	8	9	21	22	17	7	8	9	21	22	29
7	8	9	25	22	17	7	8	9	25	22	29
7	12	11	13	16	15	7	12	11	13	16	23
7	12	11	21	16	15	7	12	11	21	16	23
7	13	16	21	22	18	7	13	16	21	22	34
7	13	16	25	22	18	7	13	16	25	22	34
7	14	9	21	22	17	7	14	9	21	22	29
7	14	9	25	22	17	7	14	9	25	22	29
7	14	11	13	16	15	7	14	11	13	16	23
7	14	11	21	16	15	7	14	11	21	16	23
7	21	16	21	22	18	7	21	16	21	22	34
7	21	16	25	22	18	7	21	16	25	22	34
7	21	22	25	24	26	7	21	22	25	24	38
7	25	22	25	24	26	7	25	22	25	24	38
13	10	13	85	84	85	13	24	13	85	84	85

5.5. **The Tetrahedron Permutation Group.** It is important to note that for integer tetrahedrons, using just the six side lengths  $a, b, c, d, e, & f$ , that up to 24 tetrahedrons can be found in a family. The family is found by considering all permutations of sides, and accounting for those which create a valid tetrahedron. This permutation group accounts for the appearance of the similar tetrahedrons seen in Table 16 above.

TABLE 17. The Tetredron Permutation Group

$a$	$b$	$c$	$d$	$e$	$f$
$a$	$b$	$c$	$\sqrt{2(a^2 + e^2) - d^2}$	$e$	$\sqrt{2(c^2 + e^2) - f^2}$
$a$	$\sqrt{2(a^2 + c^2) - b^2}$	$c$	$d$	$e$	$\sqrt{2(c^2 + e^2) - f^2}$
$a$	$\sqrt{2(a^2 + c^2) - b^2}$	$c$	$\sqrt{2(a^2 + e^2) - d^2}$	$e$	$f$
$a$	$d$	$e$	$b$	$c$	$f$
$a$	$d$	$e$	$b$	$c$	$\sqrt{2(c^2 + e^2) - f^2}$
$a$	$\sqrt{2(a^2 + e^2) - d^2}$	$e$	$\sqrt{2(a^2 + c^2) - b^2}$	$c$	$\sqrt{2(c^2 + e^2) - f^2}$

*continued on next page*

Table 17 – The Tetradron Permutation Group - *continued*

$a$	$\sqrt{2(a^2 + e^2) - d^2}$	$e$	$\sqrt{2(a^2 + c^2) - b^2}$	$c$	$f$
$c$	$b$	$a$	$f$	$e$	$d$
$c$	$b$	$a$	$\sqrt{2(c^2 + e^2) - f^2}$	$e$	$\sqrt{2(a^2 + e^2) - d^2}$
$c$	$\sqrt{2(a^2 + c^2) - b^2}$	$a$	$f$	$e$	$\sqrt{2(a^2 + e^2) - d^2}$
$c$	$\sqrt{2(a^2 + c^2) - b^2}$	$a$	$\sqrt{2(c^2 + e^2) - f^2}$	$e$	$d$
$c$	$f$	$e$	$b$	$a$	$d$
$c$	$f$	$e$	$\sqrt{2(a^2 + c^2) - b^2}$	$a$	$\sqrt{2(a^2 + e^2) - d^2}$
$c$	$\sqrt{2(c^2 + e^2) - f^2}$	$e$	$b$	$a$	$\sqrt{2(a^2 + e^2) - d^2}$
$c$	$\sqrt{2(c^2 + e^2) - f^2}$	$e$	$\sqrt{2(a^2 + c^2) - b^2}$	$a$	$d$
$e$	$d$	$a$	$f$	$c$	$b$
$e$	$d$	$a$	$\sqrt{2(c^2 + e^2) - f^2}$	$c$	$\sqrt{2(a^2 + c^2) - b^2}$
$e$	$\sqrt{2(a^2 + e^2) - d^2}$	$a$	$f$	$c$	$\sqrt{2(a^2 + c^2) - b^2}$
$e$	$\sqrt{2(a^2 + e^2) - d^2}$	$a$	$\sqrt{2(c^2 + e^2) - f^2}$	$c$	$b$
$e$	$f$	$c$	$d$	$a$	$b$
$e$	$f$	$c$	$\sqrt{2(a^2 + e^2) - d^2}$	$a$	$\sqrt{2(a^2 + c^2) - b^2}$
$e$	$\sqrt{2(c^2 + e^2) - f^2}$	$c$	$d$	$a$	$\sqrt{2(a^2 + c^2) - b^2}$
$e$	$\sqrt{2(c^2 + e^2) - f^2}$	$c$	$\sqrt{2(a^2 + e^2) - d^2}$	$a$	$b$

The actual group is a set union of the 24 rows given above.

NOTE: While 24 tetrahedrons are usually in the group, sometimes less can occur, due to symmetries of identical tetrahedron sides, which can create identical rows. For *rectangular pipeds*, there are 6 tetrahedrons in the group.

**5.6. Density of rational solutions.** The next thing which must be considered is the density of the rational solutions in  $\mathbb{Q}$  as compared to the irrational solutions in  $\mathbb{R}$  real space for a Diophantine geometrical object, when considering the set of desired rational components. Sometimes this density is very sparse and clever algorithms must be utilized to even find the rational solutions that match the desired set.

For example, while running computer studies of random parallelepipeds, it became necessary to create all possible parallelepipeds using 2 integer sides and a 3rd integer diagonal, thus satisfying case 2, and case 5, for the parallelogram, automatically.

A record was kept of one such run using 3 integers,  $a, b, c$ , two  $(a, b)$  for the 4 sides and  $c$  for 1 rational diagonal and ran the integers for  $0 < a \leq 100$  and  $0 < b < a$  and  $a - b < c < a + b$  to create the parallelograms.

TABLE 18. Statistics for a short run of 746,344 integer sided parallelograms.

count	percent	case	rational diagonal ?	rational area ?
737628	98.832%	case 2	no	no
1827	0.2448%	case 5	no	yes
6683	0.8954%	case 3	yes	no
206	0.0276%	case 6	yes	yes
63	0.00844%	ibid	yes	right triangle
143	0.01916%	ibid	yes	scalene triangle
746344	100.000%	–	–	

So it can be seen that rational solutions are sparse, only 206 solutions were found from 746,344 examined, even for a computer algorithm which automatically satisfied four rational sides and 1 rational diagonal  $\in \mathbb{Z}$ .

If we can parameterize the solutions and show that they completely cover the rational solution space in  $\mathbb{Z}$  or  $\mathbb{Q}$ , this is a vast improvement in hunting for complete solutions for our set of properties to satisfy.

**5.7. Parameterizing Heron Triangles and Rational Parallelograms.** In order to efficiently discover rational solutions to *Diophantine parallelepipeds*, it is very convenient to utilize parametric solutions to automatically speed up the search process efficiently.

Such is the case here for both triangles and parallelograms.

The Heron Triangle is efficiently parameterized for the computer as the following solution using 4 integer parameters  $m, n, p$ , and  $q$ , all  $\in \mathbb{Z}$ :

**A parametric solution[6] for Heron Triangles**

$$(6) \quad \begin{aligned} \text{sides } a &= mn(p^2 + q^2) \\ b &= pq(m^2 + n^2) \\ c &= pq(n^2 - m^2) + mn(q^2 - p^2) \\ \triangle \text{ area} &= 4mnpq(mq + np)(nq - mp) \end{aligned}$$

This solution has been proven to fully cover the rational space  $\mathbb{Q}$ .

Another very helpful solution is that for parallelograms with 2 rational diagonals. This has been recently parameterized by Walter Wyss[8][9]. I changed his rational solution of  $u, m, n \in \mathbb{Q}$  to that of  $k, m, n, p, q \in \mathbb{Z}$  since the integer case is easier to handle in computers than rational numbers.

In this parametric solution, the 5 integer parameters are  $k, m, n, p$ , and  $q$ . The scaling  $k$  parameter creates composite solutions (normally  $k = 1$ ). The 4 sides of the parallelogram are  $a, b$  and the 2 diagonals are  $c, d$ .

**A parametric solution[9] for rational parallelograms**

$$(7) \quad \begin{aligned} \text{sides } a &= k(nq - mp) \\ b &= k(mq + np) \\ \text{diagonals } c &= k(p(m - n) + q(m + n)) \\ d &= k(p(n + m) + q(n - m)) \end{aligned}$$

This solution also fully covers rational space  $\mathbb{Q}$

These 2 parameterization enormously speeded up the recovery of rational triangles and parallelograms used to assemble rational component parallelepipeds, since the natural construction sequence is triangles  $\rightarrow$  parallelograms  $\rightarrow$  parallelepipeds.

**5.8. Density of Rational Solutions using Parameterization.** Even with parameterization, the density of rational solutions still has sparsity. In a run using a parallelogram parameterization[§5.7 (7)], satisfying case 3 automatically, 95,974,602 parallelograms with both diagonals rational were created. Only 25,088,615 were unique. Of those 25+ million, only 21,755 had rational area. This indicates that in the search range of  $0 < \textit{side} < 10,001$  for the sides, only 0.0867% satisfied the additional rational constraint for case 6 using the parameterization.

Using the Heron parameterization[§5.7 (6)] which satisfies case 5 automatically, of the 225,523 unique solutions found, with  $0 < \textit{sides} < 10,001$  only 5,302 had the 4th diagonal rational, while 220,221 were irrational. This meant that only 2.351% satisfied the constraint for case 6. This is better than the previous 0.0867% for the parallelogram 2 diagonals rational parameterization, but 2% is still low.

In another run of a search program using the Heron parametrization, 72,329,230 Heron triangles were found but only 704,953 were unique. Of those, 686,264 had one diagonal irrational, only 18,689 satisfied case 6 with both diagonals rational and area rational. This means that only 2.651% of the unique solutions were fully rational, but only 0.975% of the initial solutions satisfied case 6 constraints, for a net result of 0.0258% of the raw solutions, even using an efficient Heron triangle parameterization.

From this, it is seen that even when using parameterizations, the sparsity of rational solutions satisfying constraints is low. It is necessary to use clever algorithms in computer searching for Diophantine objects in embedded in 2d space. The requirements are more severe with objects embedded in 3d space, and these constraints increase geometrically as we shall see.

## 6. COMPUTER SEARCHES OF RATIONAL PARALLELEPIPEDS

We offer some interesting discoveries of pipeds from the computer searches.

To start, the extensive computer runs created 1,981,336,681 unique integer tetrahedrons which required over 441.8 gigabytes of storage space to classify. The computer search also found 79,580 degenerate parallelepipeds, where the volume = 0, the algorithms creating the pipeds did not specifically check for non-zero volume while assembling possible pipeds as this would have significantly increased the search time.

First of all, the following counts for the five classes of Diophantine parallelepipeds was obtained:

class	count	% abundance
<i>acute triclinic</i>	1315235647	66.381 229 379 8
<i>obtuse triclinic</i>	659297660	33.275 397 680 9
<i>1-ortho biclinic</i>	6743400	0.340 345 992 9
<i>2-ortho monoclinic</i>	59930	0.003 024 725 7
<i>rectangular</i>	44	0.000 002 220 7
<i>Total count</i>	1981336681	100.000 000 000 0

TABLE 19. Counts of Diophantine pipeds in each class

This shows that about 1 *obtuse* piped occurred for every 2 *acute* pipeds, while both of them together are 292.8× as numerous as *1-ortho* pipeds. Both *acute* and *obtuse* pipeds together are 32,947.3× more numerous than *2-ortho* pipeds, while *rectangular* pipeds are very scarce, accounting for only 0.0000022207% of the tetrahedrons discovered. Furthermore the ratio between the *1-ortho* and *2-ortho* pipeds is about 113 to 1.

There were 33,516 pipeds found which had rational volume, this is 0.0016915852% or only about 1 in 59,116 pipeds.

**6.1. 115 Unique Categories of Diophantine Pipeds.** After sorting the 1,981,336,681 integer tetrahedrons, it was discovered that 1,923 unique entries existed, grouped by the 27 rationality checks, and 115 unique categories resulted.

In the category table given below, we refer back to the 27 rationality checks previously determined in §4.9 but we only supply the counts in each rational check. For examine the skew triangle can have 16 possibilities, binary 4-place, decimal values 0...15, but we only give the total count of the rational components, if they occur, thus for the skew column, the count of rational occurrences would be 0...4.

NOTE: For the volume check, -1 denotes 0 volume, no 3d object exists, while 0 denotes irrational volume, and 1 denotes rational volume.

TABLE 20. 115 Unique categories of Diophantine Parallelepipeds

cat.	count	edges	skew	face	body	f area	b area	vol.	notes
1	268	3	0	6	0	0	0	-1	
2	1914630558	3	0	6	0	0	0	0	
3	14628	3	0	6	0	0	0	1	

*Continued on next page*

Table 20 – 115 Unique Categories - continued from previous page

cat.	count	edges	skew	face	body	f area	b area	vol.	notes
4	2585672	3	0	6	0	0	1	0	
5	48	3	0	6	0	0	1	1	
6	13252	3	0	6	0	0	2	0	
7	96	3	0	6	0	0	2	1	
8	48	3	0	6	0	0	3	0	
9	32448	3	0	6	0	1	0	-1	
10	19497256	3	0	6	0	1	0	0	
11	912	3	0	6	0	1	0	1	
12	6831972	3	0	6	0	1	1	0	
13	72	3	0	6	0	1	1	1	
14	1412	3	0	6	0	1	2	0	
15	2304	3	0	6	0	1	2	1	
16	12232	3	0	6	0	1	3	0	
17	69136	3	0	6	0	2	0	0	
18	2368	3	0	6	0	2	1	0	
19	20898	3	0	6	0	2	2	0	
20	110	3	0	6	0	2	4	0	
21	4440	3	0	6	0	3	0	0	
22	19628	3	0	6	0	3	1	0	
23	7962	3	0	6	0	3	2	1	
24	3776	3	0	6	0	3	3	0	
25	44	3	0	6	0	3	6	1	<i>rect</i>
26	24	3	0	6	1	0	0	-1	
27	32314073	3	0	6	1	0	0	0	
28	5036	3	0	6	1	0	0	1	
29	145920	3	0	6	1	0	1	0	
30	192	3	0	6	1	0	1	1	
31	2648	3	0	6	1	0	2	0	
32	5232	3	0	6	1	1	0	-1	
33	477828	3	0	6	1	1	0	0	
34	384	3	0	6	1	1	0	1	
35	65516	3	0	6	1	1	1	0	
36	24	3	0	6	1	1	2	0	
37	208	3	0	6	1	1	2	1	
38	196	3	0	6	1	1	3	0	
39	1892	3	0	6	1	2	0	0	
40	216	3	0	6	1	2	1	0	
41	264	3	0	6	1	3	1	0	
42	834952	3	0	6	2	0	0	0	
43	264	3	0	6	2	0	0	1	
44	24	3	0	6	2	0	1	-1	
45	12240	3	0	6	2	0	1	0	
46	504	3	0	6	2	0	2	0	
47	3056	3	0	6	2	1	0	-1	
48	102768	3	0	6	2	1	0	0	

Continued on next page

Table 20 – 115 Unique Categories - continued from previous page

cat.	count	edges	skew	face	body	f area	b area	vol.	notes
49	53252	3	0	6	2	1	1	0	
50	1320	3	0	6	2	1	3	0	
51	24	3	0	6	2	2	0	0	
52	718	3	0	6	2	2	2	0	
53	72	3	0	6	2	3	1	0	
54	182	3	0	6	2	3	2	1	
55	19416	3	0	6	3	0	0	0	
56	192	3	0	6	3	0	1	0	
57	64	3	0	6	3	1	0	-1	
58	5160	3	0	6	3	1	0	0	
59	1912	3	0	6	3	1	1	0	
60	124	3	0	6	4	0	0	0	<i>perfect</i>
61	600	3	0	6	4	0	3	-1	
62	4	3	0	6	4	1	0	0	<i>perfect</i>
63	1324	3	1	6	0	0	0	-1	
64	3126152	3	1	6	0	0	0	0	
65	96	3	1	6	0	0	0	1	
66	29040	3	1	6	0	0	1	0	
67	648	3	1	6	0	0	2	0	
68	74084	3	1	6	0	1	0	0	
69	336	3	1	6	0	1	0	1	
70	83088	3	1	6	0	1	1	0	
71	144	3	1	6	0	1	1	1	
72	24	3	1	6	0	1	2	0	
73	248	3	1	6	0	1	2	1	
74	300	3	1	6	0	1	3	0	
75	312	3	1	6	0	2	0	0	
76	108	3	1	6	0	3	1	0	
77	120	3	1	6	1	0	0	-1	
78	128176	3	1	6	1	0	0	0	
79	2988	3	1	6	1	0	1	0	
80	5160	3	1	6	1	1	0	0	
81	24	3	1	6	1	1	0	1	
82	2880	3	1	6	1	1	1	0	
83	24	3	1	6	1	1	2	1	
84	72	3	1	6	1	2	0	0	
85	5136	3	1	6	2	0	0	0	
86	48	3	1	6	2	0	0	1	
87	1304	3	1	6	2	0	1	-1	
88	360	3	1	6	2	1	0	0	
89	2460	3	1	6	2	1	1	0	
90	432	3	1	6	2	1	2	0	
91	192	3	1	6	2	1	3	0	
92	72	3	1	6	3	0	0	0	
93	12	3	1	6	3	1	1	0	

*Continued on next page*



Table 20 – 115 Unique Categories - continued from previous page

cat.	count	edges	skew	face	body	f area	b area	vol.	notes
94	16352	3	2	6	0	0	0	0	
95	24	3	2	6	0	0	0	1	
96	916	3	2	6	0	0	1	0	
97	1476	3	2	6	0	1	0	0	
98	3388	3	2	6	0	1	1	0	
99	36	3	2	6	0	1	2	1	
100	12	3	2	6	0	1	3	0	
101	96	3	2	6	0	2	0	0	
102	12	3	2	6	0	2	2	0	
103	36	3	2	6	0	3	1	0	
104	12	3	2	6	0	3	2	1	
105	808	3	2	6	1	0	0	0	
106	3024	3	2	6	1	1	0	0	
107	1404	3	2	6	1	1	1	0	
108	192	3	2	6	1	1	2	1	
109	48	3	2	6	2	1	0	0	
110	144	3	2	6	2	1	1	0	
111	180	3	3	6	0	1	1	0	
112	32744	3	4	6	0	3	6	-1	
113	48	3	4	6	1	3	6	-1	
114	2300	3	4	6	2	3	6	-1	
115	24	3	4	6	4	3	6	-1	planar

*1,981,336,681 Diophantine pipeds in 1923 entries in 115 categories*

We will select interesting examples from these 115 categories of pipeds in the next sections.

**6.2. Perfect Diophantine parallelepipeds.** *Perfect Diophantine parallelepipeds* are those where both the 6 face diagonals and the 4 body diagonals are rational.

The first perfect parallelepiped discovered was the Diophantine *acute piped* created by these 3 vectors:

vector	direction vector	length
$v_1$	$[106, 0, 0]$	106
$v_2$	$[\frac{4913}{53}, \frac{30\sqrt{202398}}{53}, 0]$	271
$v_3$	$[\frac{2911}{53}, \frac{3468\sqrt{202398}}{255407}, \frac{66\sqrt{40277202}}{4819}]$	103




FIGURE 6. The Sawyer-Reiter Acute Diophantine Piped  
 $\alpha = \arccos\left(\frac{4913}{14363}\right)$   $\beta = \arccos\left(\frac{2911}{5459}\right)$   $\gamma = \arccos\left(\frac{6647}{27913}\right)$  and  $a \neq b \neq c$

This *acute* piped discovered by Jorge F. Sawyer and Clifford A. Reiter[5] in 2008 or 2009 was the first known that had all 4 body diagonals rational as well as all 12 face diagonals rational.

The computer searches found 6 *perfect Diophantine parallelepipeds* including the Sawyer-Reiter perfect piped shown in Fig. 6. Further computer searches have revealed that the *2-ortho monoclinic piped* can have all

TABLE 21. Six perfect Diophantine parallelepipeds

	edges	skew triangles	face diagonals	body diagonals	face area	body area	vol		
	1 1 1	0 0 0 0	1 1 1 1 1 1	1 1 1 1	0 0 0	0 0 0 0 0 0	0		
Class	Basis Vectors			Tetrahedron Sides					
	$v_1$	$v_2$	$v_3$	a	b	c	d	e	f
acute	(103, 0, 0)	$\left(\frac{5822}{103}, \frac{84\sqrt{12090}}{103}, 0\right)$	$\left(\frac{6647}{103}, \frac{26010\sqrt{12090}}{41509}, \frac{66\sqrt{2405910}}{403}\right)$	103	101	106	266	271	255
acute	(335, 0, 0)	$\left(\frac{24157}{335}, \frac{96\sqrt{1558986}}{335}, 0\right)$	$\left(\frac{15510}{67}, \frac{2089315\sqrt{1558986}}{5356516}, \frac{65\sqrt{160551933366}}{79948}\right)$	335	444	365	595	630	385
1-ortho	(340, 0, 0)	(0, 357, 0)	$\left(\frac{7400}{17}, \frac{6384}{17}, \frac{720\sqrt{321}}{17}\right)$	340	493	357	852	952	875
acute	(342, 0, 0)	$\left(\frac{21385}{57}, \frac{70\sqrt{141410}}{57}, 0\right)$	$\left(\frac{18847}{57}, \frac{614608\sqrt{141410}}{4030185}, \frac{18\sqrt{6688793542510}}{70705}\right)$	342	463	595	661	739	774
acute	(375, 0, 0)	$(468, 72\sqrt{14}, 0)$	$\left(\frac{11951}{25}, -\frac{6088\sqrt{14}}{75}, \frac{8\sqrt{343966}}{15}\right)$	375	285	540	448	647	653
acute	(422, 0, 0)	$\left(\frac{81891}{211}, \frac{42\sqrt{4659370}}{211}, 0\right)$	$\left(\frac{175195}{211}, -\frac{2958000\sqrt{4659370}}{98312707}, \frac{30\sqrt{39076439247446}}{465937}\right)$	422	431	579	577	925	776

diagonals rational also. Rathbun found parametric formulii for such pipeds[3],[4].

**6.3. Interesting examples of Diophantine pipeds found by searches.** In the 14 examples given below, in general, the first occurrences of the designated piped is given.

6.3.1. *Rational volume pipeds.* Searching found that Diophantine parallelepipeds can have a rational volume, although none of the body diagonals have a rational length, nor does any of the 6 body or 3 face parallelograms have a rational area. Interesting enough, the 2<sup>nd</sup> and the 3<sup>rd</sup> pipeds in Table 22. have the same volume,

TABLE 22. Rational volume Diophantine parallelepipeds

edges	skew triangles	face diagonals	body diagonals	face area	body area	vol				
1 1 1	0 0 0 0	1 1 1 1 1 1	0 0 0 0	0 0 0	0 0 0 0 0 0	1				
Class	Basis Vectors			Tetrahedron Sides			Volume			
	$v_1$	$v_2$	$v_3$	a	b	c	d	e	f	
acute	(17, 0, 0)	$(\frac{473}{17}, \frac{24\sqrt{455}}{17}, 0)$	$(\frac{876}{17}, \frac{1848\sqrt{455}}{1105}, \frac{108\sqrt{455}}{65})$	17	32	41	61	72	43	18144
obtuse	(19, 0, 0)	$(\frac{122}{19}, \frac{84\sqrt{170}}{19}, 0)$	$(\frac{203}{19}, -\frac{216\sqrt{170}}{323}, \frac{24\sqrt{170}}{17})$	19	59	58	22	23	69	20160
acute	(25, 0, 0)	$(\frac{342}{25}, \frac{12\sqrt{3094}}{25}, 0)$	$(22, \frac{132\sqrt{3094}}{221}, \frac{120\sqrt{3094}}{221})$	25	29	30	45	50	32	20160
acute	(26, 0, 0)	$(\frac{207}{13}, \frac{30\sqrt{238}}{13}, 0)$	$(\frac{216}{13}, \frac{444\sqrt{238}}{221}, \frac{36\sqrt{238}}{17})$	26	37	39	46	48	33	30240
acute	(28, 0, 0)	$(14, 21\sqrt{5}, 0)$	$(18, \frac{111\sqrt{5}}{5}, \frac{48\sqrt{5}}{5})$	28	49	49	55	57	22	28224

20160.

6.3.2. *Rational pipeds with one body parallelogram area rational and rational volume.* Diophantine pipeds were found that had one body parallelogram with rational area, while the other 5 are irrational. None of the face parallelograms had rational area. The volume was rational also.

TABLE 23. Rational pipeds with one rational body parallelogram area and rational volume

Class	Basis Vectors			Tetrahedron Sides					
	$v_1$	$v_2$	$v_3$	a	b	c	d	e	f
obtuse	(99, 0, 0)	$(\frac{301}{3}, \frac{104\sqrt{170}}{3}, 0)$	$(\frac{77}{3}, -\frac{17072\sqrt{170}}{3315}, \frac{16632\sqrt{170}}{1105})$	99	452	463	220	209	560
obtuse	(99, 0, 0)	$(-\frac{301}{3}, \frac{104\sqrt{170}}{3}, 0)$	$(-\frac{77}{3}, -\frac{17072\sqrt{170}}{3315}, \frac{16632\sqrt{170}}{1105})$	99	494	463	242	209	560
acute	(274, 0, 0)	$(\frac{42364}{137}, \frac{120\sqrt{114130}}{137}, 0)$	$(\frac{49679}{137}, \frac{1829856\sqrt{114130}}{1563581}, \frac{10296\sqrt{114130}}{11413})$	274	298	428	507	617	325
obtuse	(209, 0, 0)	$(\frac{2525}{19}, \frac{336\sqrt{629}}{19}, 0)$	$(\frac{231}{19}, -\frac{12408\sqrt{629}}{11951}, \frac{2376\sqrt{629}}{629})$	209	450	463	220	99	494
obtuse	(209, 0, 0)	$(\frac{2525}{19}, \frac{336\sqrt{629}}{19}, 0)$	$(-\frac{231}{19}, \frac{12408\sqrt{629}}{11951}, \frac{2376\sqrt{629}}{629})$	209	450	463	242	99	452

TABLE 24. Rational component checks for the rational body area pipeds

edges	skew triangles	face diagonals	body diagonals	face area	body area	vol
1 1 1	0 0 0 0	1 1 1 1 1 1	0 0 0 0	0 0 0	1 0 0 0 0 0	1
1 1 1	0 0 0 0	1 1 1 1 1 1	0 0 0 0	0 0 0	1 0 0 0 0 0	1
1 1 1	0 0 0 0	1 1 1 1 1 1	0 0 0 0	0 0 0	0 0 0 1 0 0	1
1 1 1	0 0 0 0	1 1 1 1 1 1	0 0 0 0	0 0 0	0 0 0 0 0 1	1
1 1 1	0 0 0 0	1 1 1 1 1 1	0 0 0 0	0 0 0	0 0 0 0 0 1	1

The volume of these 5 pipeds are 8781696, 8781696, 24710400, 8781696, 8781696. As shown by these identical volumes the 1<sup>st</sup>, 2<sup>nd</sup>, 4<sup>th</sup>, and 5<sup>th</sup> tetrahedrons are in the same family, while the 3<sup>rd</sup> is in another family.

6.3.3. *Rational pipeds with all 9 parallelograms of rational area and rational volume.* It was discovered that the *rectangular* class of Diophantine pipeds, has all 3 face parallelograms and all 6 body parallelograms with rational area, as well as the volume rational. Unfortunately none of the 4 body diagonals were rational. This class of piped is related to the question of whether or not *perfect cuboids* exist, still unanswered.

TABLE 25. Parellepipeds with all face and body parallelograms of rational area

edges	skew triangles				face diagonals				body diagonals				face area			body area			vol									
1	1	1	0	0	0	0	1	1	1	1	1	1	0	0	0	0	1	1	1	1	1	1	1	1	1	1	1	1
Class	Basis Vectors			Tetrahedron Sides						Volume																		
	$v_1$	$v_2$	$v_3$	a	b	c	d	e	f																			
rectangular	(44, 0, 0)	(0, 117, 0)	(0, 0, 240)	44	125	117	244	240	267	1235520																		
rectangular	(85, 0, 0)	(0, 132, 0)	(0, 0, 720)	85	157	132	725	720	732	8078400																		
rectangular	(140, 0, 0)	(0, 480, 0)	(0, 0, 693)	140	500	480	707	693	843	46569600																		
rectangular	(160, 0, 0)	(0, 231, 0)	(0, 0, 792)	160	281	231	808	792	825	29272320																		
rectangular	(187, 0, 0)	(0, 1020, 0)	(0, 0, 1584)	187	1037	1020	1595	1584	1884	302132160																		
rectangular	(240, 0, 0)	(0, 252, 0)	(0, 0, 275)	240	348	252	365	275	373	16632000																		

6.3.4. *Rational piped with 1 rational body diagonal, and some rational area parallelograms.* Computer searches led to the discovery of one piped which had 1 body diagonal rational, and 2 body parallelograms with rational area, as well as one face parallelogram with rational area.

TABLE 26. Diophantine piped with a rational body diagonal and some parallelograms of rational area

edges	skew triangles				face diagonals				body diagonals				face area			body area			vol								
1	1	1	0	0	0	0	1	1	1	1	1	1	0	0	0	1	1	0	0	0	0	0	1	0	0	0	0
Class	Basis Vectors			Tetrahedron Sides																							
	$v_1$	$v_2$	$v_3$	a	b	c	d	e	f																		
acute	(385, 0, 0)	$\left(\frac{1211}{5}, \frac{24\sqrt{4929}}{5}, 0\right)$	$\left(\frac{539}{5}, \frac{14168\sqrt{4929}}{8215}, \frac{2464\sqrt{54219}}{1643}\right)$	385	366	415	462	385	432																		

6.3.5. *Rational piped with 2 rational body diagonals and 2 rational area face parallelograms.* One Diophantine parallelepiped was found that had 2 rational body diagonals (of 4 possible) and 2 rational area face parallelograms (of 3 possible).

TABLE 27. Diophantine piped with 2 rational body diagonals and 2 rational area face parallelograms

edges	skew triangles				face diagonals				body diagonals				face area			body area			vol								
1	1	1	0	0	0	0	1	1	1	1	1	1	0	1	0	1	0	0	0	0	0	0	0	0	0	0	0
Class	Basis Vectors			Tetrahedron Sides																							
	$v_1$	$v_2$	$v_3$	a	b	c	d	e	f																		
1-ortho	(175, 0, 0)	$\left(\frac{4017}{175}, \frac{792\sqrt{3149}}{175}, 0\right)$	$\left(0, \frac{122500\sqrt{3149}}{34639}, \frac{280\sqrt{2096947441}}{34639}\right)$	175	296	255	455	420	375																		

6.3.6. *Rational piped with 4 rational body diagonals and 1 rational area face parallelogram.* One Diophantine piped was found that had all 4 body diagonals rational, and had one face  $\square$  parallelogram with rational area.

TABLE 28. Diophantine piped with 4 rational body diagonals and 1 rational area face parallelogram

edges	skew triangles	face diagonals	body diagonals	face area	body area	vol			
1 1 1	0 0 0 0	1 1 1 1 1 1	1 1 1 1	0 1 0	0 0 0 0 0 0	0			
Class	Basis Vectors			Tetrahedron Sides					
	$v_1$	$v_2$	$v_3$	a	b	c	d	e	f
1-ortho	(340, 0, 0)	(0, 357, 0)	$\left(\frac{7400}{17}, \frac{6384}{17}, \frac{720\sqrt{321}}{17}\right)$	340	493	357	852	952	875

6.3.7. *Rational piped with 1 rational area face and 2 rational area body parallelograms.* One piped was found that had a rational face area parallelogram and 2 rational area body parallelograms.

TABLE 29. Diophantine piped with 1 rational area face and 2 rational area body parallelograms

edges	skew triangles	face diagonals	body diagonals	face area	body area	vol			
1 1 1	0 0 0 1	1 1 1 1 1 1	0 0 0 0	1 0 0	1 0 0 0 1 0	0			
Class	Basis Vectors			Tetrahedron Sides					
	$v_1$	$v_2$	$v_3$	a	b	c	d	e	f
1-ortho	(204, 0, 0)	$\left(\frac{2400}{17}, \frac{720\sqrt{21}}{17}, 0\right)$	$\left(0, \frac{680\sqrt{21}}{21}, \frac{85\sqrt{20265}}{21}\right)$	204	204	240	629	595	595

6.3.8. *Rational piped with 1 rational body diagonal, 1 rational face parallelogram and 2 rational body parallelograms and rational volume.* One Diophantine piped was found which has 1 rational body diagonal, 1 face parallelogram with rational area, and 2 body parallelograms with rational area, and rational volume.

TABLE 30. Diophantine piped with 1 rational body diagonal and rational body/face parallelograms

edges	skew triangles	face diagonals	body diagonals	face area	body area	vol				
1 1 1	0 0 0 1	1 1 1 1 1 1	1 0 0 0	0 1 0	0 0 1 0 0 1	1				
Class	Basis Vectors			Tetrahedron Sides						
	$v_1$	$v_2$	$v_3$	a	b	c	d	e	f	Volume
acute	(137, 0, 0)	$\left(\frac{3281}{137}, \frac{18480}{137}, 0\right)$	$\left(\frac{20475}{137}, \frac{17160}{137}, 216\right)$	137	176	137	250	291	250	3991680

6.3.9. *Rational piped with 3 rational body diagonals, 1 rational face parallelogram and 1 rational body parallelogram.* One parallelepiped was found that had 3 rational body diagonals, 1 rational area face parallelogram and 1 rational area body parallelogram.

TABLE 31. Diophantine piped with 3 rational body diagonals and 1 rational face and 1 rational body parallelograms

	edges	skew triangles	face diagonals	body diagonals	face area	body area	vol		
	1 1 1	0 1 0 0	1 1 1 1 1 1	1 0 1 1	0 0 1	0 0 0 1 0 0	0		
Class	Basis Vectors			Tetrahedron Sides					
	$v_1$	$v_2$	$v_3$	a	b	c	d	e	f
acute	(273, 0, 0)	$(\frac{14022}{91}, \frac{20\sqrt{3552982}}{91}, 0)$	$(\frac{14022}{91}, \frac{3068300\sqrt{3552982}}{161660681}, \frac{95760\sqrt{58624203}}{1776491})$	273	431	442	431	442	560

6.3.10. *Rational piped with 1 rational area face parallelogram and 3 rational area body parallelograms.* Computer searching found a Diophantine piped which had 1 face parallelogram with rational area, and 3 body parallelograms with rational area.

TABLE 32. Rational piped with 1 rational area face parallelogram and 3 rational area body parallelograms

	edges	skew triangles	face diagonals	body diagonals	face area	body area	vol		
	1 1 1	0 1 0 1	1 1 1 1 1 1	0 0 0 0	0 0 1	1 0 1 0 1 0	0		
Class	Basis Vectors			Tetrahedron Sides					
	$v_1$	$v_2$	$v_3$	a	b	c	d	e	f
obtuse	(361, 0, 0)	$(-\frac{64736}{361}, \frac{1320\sqrt{11041}}{361}, 0)$	$(\frac{64736}{361}, \frac{121178040\sqrt{11041}}{43843811}, \frac{80640\sqrt{143533}}{121451})$	361	663	424	425	424	448

6.3.11. *Rational pipeds with 3 rational face parallelograms and 2 rational body parallelograms.* Computer searching found two examples of *2-ortho class* pipeds, which have all the face parallelograms rational and 2 body parallelograms rational (out of 4). These pipeds are closely related to the *face cuboids* found in the rational cuboid table solutions.

TABLE 33. Diophantine cuboids related to *face cuboids*

	edges	skew triangles	face diagonals	body diagonals	face area	body area	vol			
	1 1 1	1 0 1 0	1 1 1 1 1 1	0 0 0 0	1 1 1	1 0 0 1 0 0	1			
	1 1 1	0 1 1 0	1 1 1 1 1 1	0 0 0 0	1 1 1	0 0 1 0 0 1	1			
Class	Basis Vectors			Tetrahedron Sides			Volume			
	$v_1$	$v_2$	$v_3$	a	b	c	d	e	f	
2-ortho	(153, 0, 0)	(0, 680, 0)	$(0, \frac{55096}{85}, \frac{17472}{85})$	153	697	680	697	680	208	21385728
2-ortho	(185, 0, 0)	$(\frac{12593}{185}, \frac{31824}{185}, 0)$	(0, 0, 672)	185	208	185	697	672	697	21385728

6.3.12. *Rational pipeds with 2 rational area skew triangles and 2 rational body diagonals and 1 rational area face parallelogram and 1 rational body parallelogram.* The computer searches found some interesting Diophantine pipeds which had 2 skew triangles with rational area and 2 rational body diagonals, with 1 face parallelogram rational and 1 body parallelogram rational.

Pipeds with rational skew area triangles were quite infrequent.

TABLE 34. Diophantine pipeds with 2 rational area skew triangles

	edges	skew triangles	face diagonals	body diagonals	face area	body area	vol		
	1 1 1	1 0 1 0	1 1 1 1 1 1	0 0 1 1	0 0 1	0 0 0 1 0 0	0		
	1 1 1	0 1 0 1	1 1 1 1 1 1	0 0 1 1	0 0 1	0 0 0 1 0 0	0		
	1 1 1	0 0 1 1	1 1 1 1 1 1	1 0 0 1	1 0 0	0 1 0 0 0 0	0		
Class	Basis Vectors			Tetrahedron Sides					
	$v_1$	$v_2$	$v_3$	a	b	c	d	e	f
obtuse	(36, 0, 0)	$(\frac{8}{3}, \frac{5\sqrt{2849}}{3}, 0)$	$(\frac{8}{3}, -\frac{8795\sqrt{2849}}{8547}, \frac{160\sqrt{1552705}}{2849})$	36	95	89	95	89	160
acute	(96, 0, 0)	$(\frac{825}{8}, \frac{1\sqrt{1956751}}{8}, 0)$	$(\frac{825}{8}, -\frac{552049\sqrt{1956751}}{15654008}, \frac{2520\sqrt{16966987921}}{1956751})$	96	175	203	175	203	280
obtuse	(593, 0, 0)	$(\frac{60352}{593}, \frac{11040\sqrt{1533}}{593}, 0)$	$(\frac{80801}{593}, -\frac{788020\sqrt{1533}}{303023}, \frac{620\sqrt{219219}}{511})$	593	879	736	736	593	1007

6.3.13. *Closest rational Diophantine (degenerate) piped to a superperfect piped.* The computer searches found some interesting pipeds in which all the rational checks were rational, except the volume was 0, indicating that these pipeds were degenerate planar figures.

What is interesting is that these are *2-ortho pipeds*.

TABLE 35. Diophantine piped with all rational components except 0 volume

	edges	skew triangles	face diagonals	body diagonals	face area	body area	vol		
	1 1 1	1 1 1 1	1 1 1 1 1 1	1 1 1 1	1 1 1	1 1 1 1 1 1	-1		
Class	Basis Vectors			Tetrahedron Sides					
	$v_1$	$v_2$	$v_3$	a	b	c	d	e	f
2-ortho	(120, 0, 0)	(0, 182, 0)	(0, 209, 0)	120	218	182	241	209	27

6.3.14. *Rectangular Diophantine pipeds*. There were 46 *rectangular* Diophantine pipeds found during the computer searches. These are the *body type* of cuboid solutions. Unfortunately none of them has a rational body diagonal leading to a *perfect* cuboid.

These pipeds all have the same signature for their rationality checks.

TABLE 36. Rational Component checks for the *Rectangular* Diophantine pipeds

edges	skew triangles	face diagonals	body diagonals	face area	body area	vol
1 1 1	0 0 0 0	1 1 1 1 1 1	0 0 0 0	1 1 1	1 1 1 1 1 1	1

TABLE 37. *Rectangular* Diophantine Parallelepipeds

Class	$v_1$	$v_2$	$v_3$	a	b	c	d	e	f
rectangular	(44, 0, 0)	(0, 117, 0)	(0, 0, 240)	44	125	117	244	240	267
rectangular	(44, 0, 0)	(0, 240, 0)	(0, 0, 117)	44	244	240	125	117	267
rectangular	(85, 0, 0)	(0, 132, 0)	(0, 0, 720)	85	157	132	725	720	732
rectangular	(85, 0, 0)	(0, 720, 0)	(0, 0, 132)	85	725	720	157	132	732
rectangular	(88, 0, 0)	(0, 234, 0)	(0, 0, 480)	88	250	234	488	480	534
rectangular	(88, 0, 0)	(0, 480, 0)	(0, 0, 234)	88	488	480	250	234	534
rectangular	(117, 0, 0)	(0, 44, 0)	(0, 0, 240)	117	125	44	267	240	244
rectangular	(117, 0, 0)	(0, 240, 0)	(0, 0, 44)	117	267	240	125	44	244
rectangular	(132, 0, 0)	(0, 85, 0)	(0, 0, 720)	132	157	85	732	720	725
rectangular	(132, 0, 0)	(0, 351, 0)	(0, 0, 720)	132	375	351	732	720	801
rectangular	(132, 0, 0)	(0, 720, 0)	(0, 0, 85)	132	732	720	157	85	725
rectangular	(132, 0, 0)	(0, 720, 0)	(0, 0, 351)	132	732	720	375	351	801
rectangular	(140, 0, 0)	(0, 480, 0)	(0, 0, 693)	140	500	480	707	693	843
rectangular	(140, 0, 0)	(0, 693, 0)	(0, 0, 480)	140	707	693	500	480	843
rectangular	(160, 0, 0)	(0, 231, 0)	(0, 0, 792)	160	281	231	808	792	825
rectangular	(187, 0, 0)	(0, 1020, 0)	(0, 0, 1584)	187	1037	1020	1595	1584	1884
rectangular	(234, 0, 0)	(0, 88, 0)	(0, 0, 480)	234	250	88	534	480	488
rectangular	(234, 0, 0)	(0, 480, 0)	(0, 0, 88)	234	534	480	250	88	488
rectangular	(240, 0, 0)	(0, 44, 0)	(0, 0, 117)	240	244	44	267	117	125
rectangular	(240, 0, 0)	(0, 117, 0)	(0, 0, 44)	240	267	117	244	44	125
rectangular	(240, 0, 0)	(0, 252, 0)	(0, 0, 275)	240	348	252	365	275	373
rectangular	(240, 0, 0)	(0, 275, 0)	(0, 0, 252)	240	365	275	348	252	373
rectangular	(252, 0, 0)	(0, 240, 0)	(0, 0, 275)	252	348	240	373	275	365

*Continued on next page*



Table 37 – *Rectangular Piped*s - continued from previous page

Class	$v_1$	$v_2$	$v_3$	a	b	c	d	e	f
rectangular	(252, 0, 0)	(0, 275, 0)	(0, 0, 240)	252	373	275	348	240	365
rectangular	(275, 0, 0)	(0, 240, 0)	(0, 0, 252)	275	365	240	373	252	348
rectangular	(275, 0, 0)	(0, 252, 0)	(0, 0, 240)	275	373	252	365	240	348
rectangular	(351, 0, 0)	(0, 132, 0)	(0, 0, 720)	351	375	132	801	720	732
rectangular	(351, 0, 0)	(0, 720, 0)	(0, 0, 132)	351	801	720	375	132	732
rectangular	(480, 0, 0)	(0, 88, 0)	(0, 0, 234)	480	488	88	534	234	250
rectangular	(480, 0, 0)	(0, 140, 0)	(0, 0, 693)	480	500	140	843	693	707
rectangular	(480, 0, 0)	(0, 234, 0)	(0, 0, 88)	480	534	234	488	88	250
rectangular	(480, 0, 0)	(0, 504, 0)	(0, 0, 550)	480	696	504	730	550	746
rectangular	(480, 0, 0)	(0, 550, 0)	(0, 0, 504)	480	730	550	696	504	746
rectangular	(480, 0, 0)	(0, 693, 0)	(0, 0, 140)	480	843	693	500	140	707
rectangular	(504, 0, 0)	(0, 480, 0)	(0, 0, 550)	504	696	480	746	550	730
rectangular	(504, 0, 0)	(0, 550, 0)	(0, 0, 480)	504	746	550	696	480	730
rectangular	(550, 0, 0)	(0, 480, 0)	(0, 0, 504)	550	730	480	746	504	696
rectangular	(550, 0, 0)	(0, 504, 0)	(0, 0, 480)	550	746	504	730	480	696
rectangular	(693, 0, 0)	(0, 140, 0)	(0, 0, 480)	693	707	140	843	480	500
rectangular	(693, 0, 0)	(0, 480, 0)	(0, 0, 140)	693	843	480	707	140	500
rectangular	(720, 0, 0)	(0, 85, 0)	(0, 0, 132)	720	725	85	732	132	157
rectangular	(720, 0, 0)	(0, 132, 0)	(0, 0, 85)	720	732	132	725	85	157
rectangular	(720, 0, 0)	(0, 132, 0)	(0, 0, 351)	720	732	132	801	351	375
rectangular	(720, 0, 0)	(0, 351, 0)	(0, 0, 132)	720	801	351	732	132	375

## 7. THE RECTANGULAR DIOPHANTINE PIPED AND THE PERFECT CUBOID

The most commonly known Diophantine parallelepiped known today is the *rectangular piped*, often called "the brick" or "box" or "Euler brick" also known as the *cuboid* or *integer cuboid* in math literature. There are 7 lengths of interest in this solid, the 3 edges, the 3 face diagonals and the body diagonal.

It has been found by search, that there are 3 cases which exist, called the "body cuboid", the "face cuboid", and the "edge cuboid"; where the 7th length is irrational. This is problem D18 in *Unsolved Problems in Number Theory* (UPINT) by Richard K. Guy[2].

Walter Wyss recently attempted to show that the 7th value cannot be rational by utilizing the proof of Leonard Euler that the sum or difference of 2 quartics cannot be a square[10]. He was able to parametrize the monoclinic piped and showed that approaching a right angle  $\perp$  sets up the impossible quartic condition which Euler disproved. But he failed to prove that no perfect integer cuboid exists, hence question is still open today.

Recently Renyxa D'Arox recently reached a milestone[1] with computer searching extending to  $2^{53} = 9007199254740992$  for the body diagonal, if a *perfect cuboid* exists, its body diagonal must exceed this value.

## 8. POSSIBLE FUTURE STUDIES FOR DIOPHANTINE ANALYSIS

There are some conjectures and questions created by the computer study of the 1,981,336,681 Diophantine pipeds.

Some questions:

- *Is there a rational piped with 4 rational area skew triangles?*
- *Is there a perfect parallelepiped with 2 or 3 rational area face parallelograms?*
- *Is there any perfect parallelepiped with 1..6 rational area body parallelograms?*
- *Is there any perfect parallelepipeds with a rational area skew triangle? 2? 3? or 4?*
- *Is there a perfect parallelepiped with rational volume? (asked by Sawyer, Reiter)[5]*
- *Is there a perfect parallelepiped on the rational lattice? (asked by Sawyer, Reiter)[5]*

Two conjectures:

- *Conjecture 1. Other Diophantine pipeds do exist with less than 6 rational face diagonals as covered by the computer studies done here*
- *Conjecture 2. No Diophantine pipeds exist with all 27 components rational*

These are just some questions and two conjectures relating to the computer search results.

Some interesting problems for further analysis are the following:

- Parametrize the *acute* or *obtuse triclinic* piped as an infinite family
- Parametrize the *mono-orthogonal* or *biclinic* piped as an infinite family

## APPENDIX A - PROOF OF 5 PARALLELEPIPED CLASSES

Let the parallelepiped be situated so the the main vertice  $v_1$  determined by the 3 basis vectors is at the origin  $[0,0,0]$ . Figure 7 shows the parallelepiped and its 8 labeled vertices. Previously, Figure 1 showed the 3 basis vectors, the same piped is shown.




FIGURE 7. Labeled vertices of the parallelepiped.

The origin  $[0,0,0]$  is labeled  $v_1$ . The  $\vec{a}$  basis vector lies along the positive x-axis,  $v_1$  to  $v_2$ ; the  $\vec{b}$  basis vector is in the x-y plane, along  $v_1$  to  $v_3$ ; and the  $\vec{c}$  basis vector is along the line from  $v_1$  to  $v_5$ , with a non-zero z component.

We label the surface angles, as found by the angle traced by the 3 ordered vertices.

TABLE 38. The 3 surface angles at each vertice of parallelepiped

vertice	$angle_1$	$angle_2$	$angle_3$
1	$v_2v_1v_3$	$v_2v_1v_5$	$v_3v_1v_5$
2	$v_1v_2v_4$	$v_1v_2v_6$	$v_4v_2v_6$
3	$v_1v_3v_4$	$v_1v_3v_7$	$v_4v_3v_7$
4	$v_2v_4v_3$	$v_2v_4v_8$	$v_3v_4v_8$
5	$v_1v_5v_6$	$v_1v_5v_7$	$v_6v_5v_7$
6	$v_2v_6v_5$	$v_2v_6v_8$	$v_5v_6v_8$
7	$v_3v_7v_5$	$v_3v_7v_8$	$v_5v_7v_8$
8	$v_4v_8v_6$	$v_4v_8v_7$	$v_6v_8v_7$

We know that the interior angles on the diagonals of a parallelogram are equal, and that 2 adjacent angles are supplementary. This fact enables us to substitute the equivalent angles as necessary for the piped, in order to redefine all surface angles as a subset of the 3 surface angles at vertice #1.

Equivalent angles for the parallelepiped vertices occurring on a basis plane.

$a - b$ plane	$a - c$ plane	$b - c$ plane
$v_2v_1v_5$	$v_2v_1v_3$	$v_3v_1v_5$
$v_2v_6v_5 = v_2v_1v_5$	$v_1v_2v_4 = 180^\circ - v_2v_1v_3$	$v_3v_7v_5 = v_3v_1v_5$
$v_1v_5v_6 = 180^\circ - v_2v_1v_5$	$v_1v_3v_4 = 180^\circ - v_2v_1v_3$	$v_1v_3v_7 = 180^\circ - v_3v_1v_5$
$v_1v_2v_6 = 180^\circ - v_2v_1v_5$	$v_2v_4v_3 = v_2v_1v_3$	$v_1v_5v_7 = 180^\circ - v_3v_1v_5$
$v_4v_3v_7 = v_2v_1v_5$	$v_6v_5v_7 = v_2v_1v_3$	$v_4v_2v_6 = v_3v_1v_5$
$v_4v_8v_7 = v_2v_1v_5$	$v_5v_6v_8 = 180^\circ - v_2v_1v_3$	$v_4v_8v_6 = v_3v_1v_5$
$v_3v_7v_8 = 180^\circ - v_2v_1v_5$	$v_5v_7v_8 = 180^\circ - v_2v_1v_3$	$v_2v_4v_8 = 180^\circ - v_3v_1v_5$
$v_3v_4v_8 = 180^\circ - v_2v_1v_5$	$v_6v_8v_7 = v_2v_1v_3$	$v_2v_6v_8 = 180^\circ - v_3v_1v_5$

We substitute the 3 surface angles or their supplements found at vertice #1, for all 8 vertices, such that all surface angles are derived from just the angles at the origin, or vertice #1.

After substitutions, we find the surface angles.

Equivalent 3 surface angles at each vertice of parallelepiped

vertice	$angle_1$	$angle_2$	$angle_3$
1	$v_2v_1v_3$	$v_2v_1v_5$	$v_3v_1v_5$
2	$180^\circ - v_2v_1v_3$	$180^\circ - v_2v_1v_5$	$v_3v_1v_5$
3	$180^\circ - v_2v_1v_3$	$180^\circ - v_3v_1v_5$	$v_2v_1v_5$
4	$v_2v_1v_3$	$180^\circ - v_3v_1v_5$	$180^\circ - v_2v_1v_5$
5	$180^\circ - v_2v_1v_5$	$180^\circ - v_3v_1v_5$	$v_2v_1v_3$
6	$v_2v_1v_5$	$180^\circ - v_3v_1v_5$	$180^\circ - v_2v_1v_3$
7	$v_3v_1v_5$	$180^\circ - v_2v_1v_5$	$180^\circ - v_2v_1v_3$
8	$v_3v_1v_5$	$v_2v_1v_5$	$v_2v_1v_3$

In order to classify the angles, we consider the  $\sphericalangle$  such that

$$\begin{aligned}
 0^\circ < \sphericalangle < 90^\circ &\rightarrow \text{sign}(\cos(\sphericalangle)) = 1 \\
 \sphericalangle = 90^\circ &\rightarrow \text{sign}(\cos(\sphericalangle)) = 0 \\
 180^\circ > \sphericalangle > 90^\circ &\rightarrow \text{sign}(\cos(\sphericalangle)) = -1
 \end{aligned}$$

We substitute these values for the surface angles in the labeling above. For example, vertice 1 could have the signs: [-1, 0, 1], if its angles were  $108^\circ$ ,  $90^\circ$ , and  $60^\circ$  respectively.

We notice that if the sign of an angle = 1, then its supplement = -1, or vice versa. If the sign of an angle = 0, then the supplement is also = 0.

Since all other 7 vertices are given as some combination of the 3 surface angles of vertice #1, we only have to consider the 27 permutations of -1,0,1 taken 3 at a time for vertice #1, in order to enumerate all possible states of the parallelepiped, and obtain an exhaustive list.

For each permutation, we create a set union of all 8 vertices to distinguish the actual unique signature of each possible permutation group.

TABLE 39. The set union set of the 27 permutations for all 8 vertices of the parallelepiped

#	group	#	group	#	group
1	[-1, -1, -1], [-1, 1, 1]	10	[-1, -1, 0], [-1, 0, 1], [0, 1, 1]	19	[-1, -1, 1], [1, 1, 1]
2	[-1, -1, 0], [-1, 0, 1], [0, 1, 1]	11	[-1, 0, 0], [0, 0, 1]	20	[-1, -1, 0], [-1, 0, 1], [0, 1, 1]
3	[-1, -1, 1], [1, 1, 1]	12	[-1, -1, 0], [-1, 0, 1], [0, 1, 1]	21	[-1, -1, -1], [-1, 1, 1]
4	[-1, -1, 0], [-1, 0, 1], [0, 1, 1]	13	[-1, 0, 0], [0, 0, 1]	22	[-1, -1, 0], [-1, 0, 1], [0, 1, 1]
5	[-1, 0, 0], [0, 0, 1]	14	[0, 0, 0]	23	[-1, 0, 0], [0, 0, 1]
6	[-1, -1, 0], [-1, 0, 1], [0, 1, 1]	15	[-1, 0, 0], [0, 0, 1]	24	[-1, -1, 0], [-1, 0, 1], [0, 1, 1]
7	[-1, -1, 1], [1, 1, 1]	16	[-1, -1, 0], [-1, 0, 1], [0, 1, 1]	25	[-1, -1, -1], [-1, 1, 1]
8	[-1, -1, 0], [-1, 0, 1], [0, 1, 1]	17	[-1, 0, 0], [0, 0, 1]	26	[-1, -1, 0], [-1, 0, 1], [0, 1, 1]
9	[-1, -1, -1], [-1, 1, 1]	18	[-1, -1, 0], [-1, 0, 1], [0, 1, 1]	27	[-1, -1, 1], [1, 1, 1]

We discover that 5 unique classes result, after creating a set union of the 27 permutation set union groups shown in Table 39.

TABLE 40. Five classes of parallelepipeds

name	class
rectangular	[0, 0, 0]
monoclinic	[-1, 0, 0], [0, 0, 1]
bi-clinic	[-1, -1, 0], [-1, 0, 1], [0, 1, 1]
obtuse triclinic	[-1, -1, -1], [-1, 1, 1]
acute triclinic	[-1, -1, 1], [1, 1, 1]

□

#### REFERENCES

- [1] Renyxa D'Arox, *Body diagonal of the perfect cuboid must exceed  $2^{53}$* , private communication, November 25, 2018 11:04 PM PST
- [2] Richard K. Guy, *Unsolved Problems in Number Theory*, ver. 3, Springer-Verlag, New York, ISBN 978-0-387-20860-2 (hardcover), 2004, 438 pages
- [3] Randall Rathbun, *Some new parameterizations for the Diophantine bi-orthogonal monoclinic piped*, <https://arxiv.org/abs/1709.01138> version [v1] Mon, 4 Sep 2017 20:02:15 UTC
- [4] Randall Rathbun, *Four integer parametrizations for the monoclinic Diophantine piped*, <https://arxiv.org/abs/1705.07734> version [v2] Fri, 2 Jun 2017 17:12:44 UTC
- [5] J. Sawyer, C.A. Reiter, *Perfect parallelepipeds exist*, Math Comp., 80, (2011), 1037-1040 Also Randall Rathbun discovered some earlier.
- [6] Hermann Schubert, §2 *Heron Triangles, formula, page 10*, Integrality in Algebraic Geometry 1905, translation of Die Ganzzahligkeit in der Algebraischen Geometrie" by Ralph Buchholz 2005
- [7] Benjamin D. Sokolowsky, Amy G. VanHooft, Rachel M. Volkert, Clifford A. Reiter, *An Infinite Family of Perfect Parallelepipeds*, Mathematics of Computation, Volume 83, Number 289, September 2014, pp 2441-2454
- [8] Walter Wyss, *Perfect Parallelograms*, American Math Monthly, 119 (6) (2012), pp. 513-515

- [9] Walter Wyss, *Sums of Squares, Bijective Parameter Representation*,  
<https://arxiv.org/abs/1402.0102> version #1 Sat, 1 Feb 2014 16:49:20 GMT
- [10] Walter Wyss, *No Perfect Cuboid*, <http://arxiv.org/abs/1506.02215v4> version #4 Tue 14 Jun 2016  
18:30:50 GMT

---

*E-mail address:* [randallrathbun@gmail.com](mailto:randallrathbun@gmail.com)