

# Ramanujan's theta functions and linear combinations of four triangular numbers

Zhi-Hong Sun

School of Mathematics and Statistics  
 Huaiyin Normal University  
 Huaian, Jiangsu 223300, P.R. China  
 Email: zhsun@hytc.edu.cn  
 Homepage: <http://maths.hytc.edu.cn/szh1.htm>

## Abstract

Let  $\mathbb{Z}$  and  $\mathbb{Z}^+$  be the set of integers and the set of positive integers, respectively. For  $a, b, c, d, n \in \mathbb{Z}^+$  let  $t(a, b, c, d; n)$  be the number of representations of  $n$  by  $ax(x+1)/2 + by(y+1)/2 + cz(z+1)/2 + dw(w+1)/2$  ( $x, y, z, w \in \mathbb{Z}$ ). In this paper, by using Ramanujan's theta functions  $\varphi(q)$  and  $\psi(q)$  we present many formulas and conjectures on  $t(a, b, c, d; n)$ .

Keywords: theta function; triangular number; quadratic form

Mathematics Subject Classification 2010: 11D85, 11E25, 30B10, 33E20

## 1. Introduction

Let  $\mathbb{Z}$ ,  $\mathbb{Z}^+$  and  $\mathbb{N}$  be the set of integers, the set of positive integers and the set of nonnegative integers, respectively, and let  $\mathbb{Z}^k = \underbrace{\mathbb{Z} \times \mathbb{Z} \times \cdots \times \mathbb{Z}}_{k \text{ times}}$ . For  $a_1, a_2, \dots, a_k \in \mathbb{Z}^+$  ( $k \geq 2$ )

and  $n \in \mathbb{N}$  set

$$N(a_1, a_2, \dots, a_k; n) = \left| \{(x_1, \dots, x_k) \in \mathbb{Z}^k \mid n = a_1x_1^2 + a_2x_2^2 + \cdots + a_kx_k^2\} \right|,$$

$$t(a_1, a_2, \dots, a_k; n) = \left| \left\{ (x_1, \dots, x_k) \in \mathbb{Z}^k \mid n = a_1 \frac{x_1(x_1-1)}{2} + a_2 \frac{x_2(x_2-1)}{2} + \cdots + a_k \frac{x_k(x_k-1)}{2} \right\} \right|$$

and

$$C(a_1, \dots, a_k) = \frac{i_1(i_1-1)(i_1-2)(i_1-3)}{4!} + \frac{i_1(i_1-1)i_2}{2} + i_1i_3,$$

where  $i_j$  denotes the number of elements in  $\{a_1, \dots, a_k\}$  which are equal to  $j$ . For convenience we also define

$$t(a_1, a_2, \dots, a_k; n) = N(a_1, a_2, \dots, a_k; n) = 0 \quad \text{for } n \notin \mathbb{N}.$$

In 2005 Adiga, Cooper and Han [ACH] showed that

$$(1.1) \quad \begin{aligned} & t(a_1, a_2, \dots, a_k; n) \\ &= \frac{2}{2 + C(a_1, \dots, a_k)} N(a_1, \dots, a_k; 8n + a_1 + \cdots + a_k) \quad \text{for } a_1 + \cdots + a_k \leq 7. \end{aligned}$$

In 2008 Baruah, Cooper and Hirschhorn [BCH] proved that

$$(1.2) \quad \begin{aligned} & t(a_1, a_2, \dots, a_k; n) \\ &= \frac{2}{2 + C(a_1, \dots, a_k)} (N(a_1, \dots, a_k; 8n + 8) - N(a_1, \dots, a_k; 2n + 2)) \\ & \quad \text{for } a_1 + \dots + a_k = 8. \end{aligned}$$

Ramanujan's theta functions  $\varphi(q)$  and  $\psi(q)$  are defined by

$$\varphi(q) = \sum_{n=-\infty}^{\infty} q^{n^2} = 1 + 2 \sum_{n=1}^{\infty} q^{n^2} \quad \text{and} \quad \psi(q) = \sum_{n=0}^{\infty} q^{n(n+1)/2} \quad (|q| < 1).$$

It is evident that for positive integers  $a_1, \dots, a_k$  and  $|q| < 1$ ,

$$(1.3) \quad \sum_{n=0}^{\infty} N(a_1, \dots, a_k; n) q^n = \varphi(q^{a_1}) \cdots \varphi(q^{a_k}),$$

$$(1.4) \quad \sum_{n=0}^{\infty} t(a_1, \dots, a_k; n) q^n = 2^k \psi(q^{a_1}) \cdots \psi(q^{a_k}).$$

There are many identities involving  $\varphi(q)$  and  $\psi(q)$ . From [BCH, Lemma 4.1] or [Be] we know that for  $|q| < 1$ ,

$$(1.5) \quad \psi(q)^2 = \varphi(q)\psi(q^2),$$

$$(1.6) \quad \varphi(q) = \varphi(q^4) + 2q\psi(q^8) = \varphi(q^{16}) + 2q^4\psi(q^{32}) + 2q\psi(q^8),$$

$$(1.7) \quad \varphi(q)^2 = \varphi(q^2)^2 + 4q\psi(q^4)^2 = \varphi(q^4)^2 + 4q^2\psi(q^8)^2 + 4q\psi(q^4)^2,$$

$$(1.8) \quad \psi(q)\psi(q^3) = \varphi(q^6)\psi(q^4) + q\varphi(q^2)\psi(q^{12}).$$

By [S1, Lemma 2.4],

$$(1.9) \quad \varphi(q)^2 = \varphi(q^8)^2 + 4q^4\psi(q^{16})^2 + 4q^2\psi(q^8)^2 + 4q\varphi(q^{16})\psi(q^8) + 8q^5\psi(q^8)\psi(q^{32}).$$

By [S1, Lemma 2.3], for  $|q| < 1$  we have

$$(1.10) \quad \begin{aligned} \varphi(q)\varphi(q^3) &= \varphi(q^{16})\varphi(q^{48}) + 4q^{16}\psi(q^{32})\psi(q^{96}) + 2q\varphi(q^{48})\psi(q^8) + 2q^3\varphi(q^{16})\psi(q^{24}) \\ & \quad + 6q^4\psi(q^8)\psi(q^{24}) + 4q^{13}\psi(q^8)\psi(q^{96}) + 4q^7\psi(q^{24})\psi(q^{32}). \end{aligned}$$

Let  $a, b, c, d, n \in \mathbb{Z}^+$ . From 1859 to 1866 Liouville made about 90 conjectures on  $N(a, b, c, d; n)$  in a series of papers. Most conjectures of Liouville have been proved. See Cooper's survey paper [C], Dickson's historical comments [D] and Williams' book [W]. Recently, some connections between  $t(a, b, c, d; n)$  and  $N(a, b, c, d; 8n + a + b + c + d)$  have been found. See [ACH, BCH, S1, S3, WS2]. More recently Yao [Y] confirmed some conjectures posed by the author in [S1]. We also note that the evaluations of  $t(a, b, c, d; n)$  ( $a + b + c + d \geq 8$ ) have been given for some special values of  $(a, b, c, d)$ . In [C] Cooper determined  $t(a, b, c, d; n)$  for  $(a, b, c, d) = (1, 3, 3, 3), (1, 2, 2, 3), (1, 3, 6, 6), (1, 3, 4, 4), (1, 1, 2, 6), (1, 3, 12, 12)$ , in [WS1] Wang and Sun determined  $t(a, b, c, d; n)$  for  $(a, b, c, d) = (1, 2, 2, 4), (1, 2, 4, 4), (1, 1, 4, 4), (1, 4, 4, 4), (1, 3, 3, 9), (1, 1, 9, 9), (1, 9, 9, 9), (1, 1, 1, 9), (1, 3, 9, 9), (1, 1, 3, 9)$ , in [WS2] Wang and Sun determined  $t(a, b, c, d; n)$  for  $(a, b, c, d) = (1, 1, 2, 8)$ ,

$(1, 1, 2, 16), (1, 2, 3, 6), (1, 3, 4, 12), (1, 1, 3, 4), (1, 1, 5, 5), (1, 5, 5, 5), (1, 3, 3, 12), (1, 1, 1, 12), (1, 1, 3, 12), (1, 3, 3, 4)$ , in [S1] Sun determined  $t(a, b, c, d; n)$  for  $(a, b, c, d) = (1, 3, 3, 6), (1, 1, 8, 8), (1, 1, 4, 8)$ . In [S1] the author stated that for odd  $a$ ,

$$\begin{aligned} t(a, a, 2a, 4b; 4n + 3a) &= 4t(a, 2a, 4a, b; n), & t(a, a, 6a, 4b; 4n + 3a) &= 2t(a, a, 6a, b; n), \\ t(a, a, 8a, 2b; 2n) &= t(a, 2a, 2a, b; n), & t(a, a, 8a, 2b; 2n + a) &= 2t(a, 4a, 4a, b; n). \end{aligned}$$

In Section 2, using theta function identities we establish new general results for  $t(a, b, c, d; n)$ . Let  $a, b, n \in \mathbb{Z}^+$  and  $k \in \mathbb{N}$ . We show that for odd integers  $a$  and  $b$ ,

$$\begin{aligned} t(a, 2a, 2a, 2b; n) &= \frac{1}{2}N(a, a, 4a, 2b; 8n + 5a + 2b), \\ t(a, 3a, 3a, 2b; n) &= N(3a, 3a, 4a, 2b; 8n + 7a + 2b), \\ t(a, a, 2a, b; n) &= 2N(a, 4a, 8a, b; 8n + 4a + b) \quad \text{for } a \equiv -b \pmod{4}, \\ t(2a, 2a, 3a, b; n) &= \frac{1}{3}N(a, 3a, 16a, 4b; 32n + 28a + 4b) \quad \text{for } a \equiv -b \pmod{4}, \\ t(a, 6a, 6a, b; n) &= \frac{1}{3}N(a, 3a, 48a, 4b; 32n + 52a + 4b) \quad \text{for } a \equiv b \pmod{4}, \\ t(a, 3a, b, 3b; n) &= 4N(a, 3a, b, 3b; 2n + a + b) \quad \text{for } n \equiv (a - b)/2 \pmod{2}. \end{aligned}$$

For  $a, b \in \mathbb{Z}^+$  let  $(a, b)$  be the greatest common divisor of  $a$  and  $b$ . For an odd prime  $p$  and  $a \in \mathbb{Z}$  let  $\left(\frac{a}{p}\right)$  be the Legendre symbol. Suppose that  $a, b, n \in \mathbb{Z}^+$ ,  $(a, b) = 1$ ,  $b \not\equiv 0, -a \pmod{4}$  and there is an odd prime divisor  $p$  of  $b$  such that  $\left(\frac{a(8n+9a)}{p}\right) = -1$ . We prove that

$$t(a, 4a, 4a, b; n) = \frac{1}{2}N(a, 4a, 4a, b; 8n + 9a + b).$$

In Section 3, using theta function identities we establish 31 transformation formulas for  $t(a, b, c, d; n)$ . As typical examples, for  $a, b, c, n \in \mathbb{Z}^+$  with  $2 \nmid a$  we have

$$\begin{aligned} t(a, a, 2b, 2c; 2n + a) &= 2t(a, 4a, b, c; n), \\ t(a, 3a, 4b, 4c; 4n + 3a) &= 2t(3a, 4a, b, c; n), \\ t(a, 3a, 4b, 4c; 4n + 6a) &= 2t(a, 12a, b, c; n), \\ t(a, 7a, 2b, 2c; 2n + a) &= t(a, 7a, b, c; n), \\ t(3a, 5a, 2b, 2c; 2n + 3a) &= t(a, 15a, b, c; n), \\ t(a, 15a, 2b, 2c; 2n) &= t(3a, 5a, b, c; n), \\ t(a, a, 6a, 8b; 8n + 6a) &= 4t(2a, 2a, 3a, b; n), \\ t(2a, 3a, 3a, 8b; 8n + 12a) &= 4t(a, 6a, 6a, b; n). \end{aligned}$$

In Section 4, we completely determine  $t(2, 3, 3, 8; n)$ ,  $t(1, 1, 6, 24; n)$  and  $t(1, 1, 6, 8; n)$  for any positive integer  $n$ . In Section 5 we prove some special relations between  $t(a, b, c, d; n)$  and  $N(a, b, c, d; n)$ , and pose many challenging conjectures based on calculations on Maple.

## 2. New general formulas for $t(a, b, c, d; n)$

In this section we present several general formulas for  $t(a, b, c, d; n)$ , which were found by calculations on Maple and proved by using Ramanujan's theta functions.

**Theorem 2.1.** Let  $a, b \in \mathbb{Z}^+$  with  $2 \nmid ab$ . For  $n = 0, 1, 2, \dots$  we have

$$t(a, 2a, 2a, 2b; n) = \frac{1}{2}N(a, a, 4a, 2b; 8n + 5a + 2b).$$

Proof. By (1.6) and (1.9),

$$\begin{aligned} & \sum_{n=0}^{\infty} N(a, a, 4a, 2b; n)q^n \\ &= \varphi(q^a)^2 \varphi(q^{4a}) \varphi(q^{2b}) \\ &= (\varphi(q^{8a})^2 + 4q^{4a} \psi(q^{16a})^2 + 4q^{2a} \psi(q^{8a})^2 + 4q^a \varphi(q^{16a}) \psi(q^{8a}) + 8q^{5a} \psi(q^{8a}) \psi(q^{32a})) \\ & \quad \times (\varphi(q^{16a}) + 2q^{4a} \psi(q^{32a})) (\varphi(q^{8b}) + 2q^{2b} \psi(q^{16b})). \end{aligned}$$

Collecting the terms of the form  $q^{8n+5a+2b}$  we get

$$\begin{aligned} & \sum_{n=0}^{\infty} N(a, a, 4a, 2b; 8n + 5a + 2b)q^{8n+5a+2b} \\ &= 8q^{5a} \psi(q^{8a}) \psi(q^{32a}) \cdot \varphi(q^{16a}) \cdot 2q^{2b} \psi(q^{16b}) \\ & \quad + 4q^a \varphi(q^{16a}) \psi(q^{8a}) \cdot 2q^{4a} \psi(q^{32a}) \cdot 2q^{2b} \psi(q^{16b}) \\ &= 32q^{5a+2b} \psi(q^{8a}) \psi(q^{16b}) \varphi(q^{16a}) \psi(q^{32a}) = 32q^{5a+2b} \psi(q^{8a}) \psi(q^{16a})^2 \psi(q^{16b}). \end{aligned}$$

Therefore,

$$\begin{aligned} & \sum_{n=0}^{\infty} N(a, a, 4a, 2b; 8n + 5a + 2b)q^n \\ &= 32 \psi(q^a) \psi(q^{2b}) \varphi(q^{2a}) \psi(q^{4a}) = 2 \sum_{n=0}^{\infty} t(a, 2a, 4a, 2b; n)q^n. \end{aligned}$$

Hence the result follows.

**Theorem 2.2.** Let  $a, b \in \mathbb{Z}^+$  with  $ab \equiv -1 \pmod{4}$ . For  $n \in \mathbb{Z}^+$  we have

$$t(a, a, 2a, b; n) = 2N(a, 4a, 8a, b; 8n + 4a + b).$$

Proof. It is clear that

$$\begin{aligned} & \sum_{n=0}^{\infty} N(a, 4a, 8a, b; n)q^n = \varphi(q^a) \varphi(q^b) \varphi(q^{4a}) \varphi(q^{8a}) \\ &= (\varphi(q^{16a}) + 2q^{4a} \psi(q^{32a}) + 2q^a \psi(q^{8a})) (\varphi(q^{16b}) + 2q^{4b} \psi(q^{32b}) + 2q^b \psi(q^{8b})) \\ & \quad \times (\varphi(q^{16a}) + 2q^{4a} \psi(q^{32a})) \varphi(q^{8a}). \end{aligned}$$

Thus,

$$\begin{aligned} & \sum_{n=0}^{\infty} N(a, 4a, 8a, b; 8n + 4a + b)q^{8n+4a+b} \\ &= 2q^b \psi(q^{8b}) (\varphi(q^{16a}) \cdot 2q^{4a} \psi(q^{32a}) + 2q^{4a} \psi(q^{32a}) \cdot \varphi(q^{16a})) \varphi(q^{8a}) \end{aligned}$$

and so

$$\begin{aligned}
& \sum_{n=0}^{\infty} N(a, 4a, 8a, b; 8n + 4a + b)q^n \\
&= 8\psi(q^b)\varphi(q^{2a})\psi(q^{4a})\varphi(q^a) = 8\psi(q^b)\psi(q^{2a})^2\varphi(q^a) = 8\psi(q^a)^2\psi(q^{2a})\psi(q^b) \\
&= \frac{1}{2} \sum_{n=0}^{\infty} t(a, a, 2a, b; n)q^n.
\end{aligned}$$

This yields the result.

**Corollary 2.1.** *Suppose  $a, b, n \in \mathbb{Z}^+$  and  $ab \equiv -1 \pmod{4}$ . Then*

$$3N(a, 4a, 8a, b; 8n + 4a + b) = N(a, a, 2a, b; 8n + 4a + b) - N(a, a, 2a, 4b; 8n + 4a + b).$$

Proof. From [S3, Theorem 3.1] we know that

$$t(a, a, 2a, b; n) = \frac{2}{3}(N(a, a, 2a, b; 8n + 4a + b) - N(a, a, 2a, 4b; 8n + 4a + b)).$$

This together with Theorem 2.2 yields the result.

**Corollary 2.2.** *Suppose that  $n \in \{0, 1, 2, \dots\}$ . Then*

$$N(1, 3, 4, 8; 8n + 7) = \frac{1}{5}N(1, 1, 2, 3; 8n + 7).$$

Proof. By Theorem 2.2,  $t(1, 1, 2, 3; n) = 2N(1, 4, 8, 3; 8n+7)$ . By [ACH],  $t(1, 1, 2, 3; n) = \frac{2}{5}N(1, 1, 2, 3; 8n + 7)$ . Thus the result follows.

**Theorem 2.3.** *Let  $a, b \in \mathbb{Z}^+$  with  $2 \nmid ab$ . For  $n \in \mathbb{Z}^+$  we have*

$$t(a, 3a, 3a, 2b; n) = N(3a, 3a, 4a, 2b; 8n + 7a + 2b).$$

Proof. Clearly,

$$\begin{aligned}
& \sum_{n=0}^{\infty} N(3a, 3a, 4a, 2b; n)q^n = \varphi(q^{3a})^2\varphi(q^{4a})\varphi(q^{2b}) \\
&= (\varphi(q^{12a})^2 + 4q^{6a}\psi(q^{24a})^2 + 4q^{3a}\psi(q^{12a})^2)(\varphi(q^{8b}) + 2q^{2b}\psi(q^{16b}))\varphi(q^{4a}).
\end{aligned}$$

Thus,

$$\begin{aligned}
& \sum_{n=0}^{\infty} N(3a, 3a, 4a, 2b; 4n + 3a + 2b)q^{4n+3a+2b} \\
&= 4q^{3a}\psi(q^{12a})^2 \cdot 2q^{2b}\psi(q^{16b}) \cdot \varphi(q^{4a}) = 8q^{3a+2b}\varphi(q^{4a})\psi(q^{12a})^2\psi(q^{16b})
\end{aligned}$$

and so

$$\sum_{n=0}^{\infty} N(3a, 3a, 4a, 2b; 4n + 3a + 2b)q^n = 8\varphi(q^a)\psi(q^{3a})^2\psi(q^{4b}) = 8\varphi(q^a)\varphi(q^{3a})\psi(q^{6a})\psi(q^{4b}).$$

Applying (1.6) and (1.8) we see that

$$\begin{aligned}
& \sum_{n=0}^{\infty} N(3a, 3a, 4a, 2b; 4n + 3a + 2b)q^n \\
&= 8(\varphi(q^{4a}) + 2q^a\psi(q^{8a}))(\varphi(q^{12a}) + 2q^{3a}\psi(q^{24a}))\psi(q^{6a})\psi(q^{4b}) \\
&= 8(\varphi(q^{4a})\varphi(q^{12a}) + 4q^{4a}\psi(q^{8a})\psi(q^{24a}) + 2q^a\psi(q^{2a})\psi(q^{6a}))\psi(q^{6a})\psi(q^{4b}).
\end{aligned}$$

Hence

$$\sum_{n=0}^{\infty} N(3a, 3a, 4a, 2b; 4(2n + a) + 3a + 2b)q^{2n+a} = 16q^a\psi(q^{2a})\psi(q^{6a})^2\psi(q^{4b})$$

and so

$$\sum_{n=0}^{\infty} N(3a, 3a, 4a, 2b; 8n + 7a + 2b)q^n = 16\psi(q^a)\psi(q^{3a})^2\psi(q^{2b}) = \sum_{n=0}^{\infty} t(a, 3a, 3a, 2b; n)q^n,$$

which gives the result.

**Corollary 2.3.** *Suppose  $n \in \mathbb{Z}^+$ . Then*

$$N(3, 3, 4, 6; 8n + 13) = 2N(1, 3, 12, 24; 8n + 13) = \frac{2}{5}N(1, 3, 3, 6; 8n + 13).$$

Proof. By Theorem 2.3,  $t(1, 3, 3, 6; n) = N(3, 3, 4, 6; 8n + 13)$ . By Theorem 2.2,  $t(1, 3, 3, 6; n) = 2N(1, 3, 12, 24; 8n + 13)$ . By [S1, Theorem 2.10],  $t(1, 3, 3, 6; n) = \frac{2}{5}N(1, 3, 3, 6; 8n + 13)$ . Thus, the result follows.

**Corollary 2.4.** *Suppose  $a, b, n \in \mathbb{Z}^+$  and  $2 \nmid ab$ . Then*

$$\begin{aligned}
\text{(i)} \quad & N(3a, 4a, 12a, 2b; 8n + 7a + 2b) = \frac{1}{2}N(3a, 3a, 4a, 2b; 8n + 7a + 2b), \\
\text{(ii)} \quad & N(a, 3a, 3a, 8b; 8n + 7a + 2b) \\
&= N(a, 3a, 3a, 2b; 8n + 7a + 2b) - 2N(3a, 3a, 4a, 2b; 8n + 7a + 2b), \\
\text{(iii)} \quad & N(a, 3a, 12a, 2b; 8n + 7a + 2b) \\
&= N(a, 3a, 3a, 2b; 8n + 7a + 2b) - \frac{3}{2}N(3a, 3a, 4a, 2b; 8n + 7a + 2b).
\end{aligned}$$

Proof. By [S3, Corollary 4.2],  $t(a, 3a, 3a, 2b; n) = 2N(4a, 12a, 3a, 2b; 8n + 7a + 2b)$ . This together with Theorem 2.3 proves (i). By [S3, Theorem 3.1] and Theorem 2.3,

$$\begin{aligned}
& N(a, 3a, 3a, 2b; 8n + 7a + 2b) - N(a, 3a, 3a, 8b; 8n + 7a + 2b) \\
&= 2t(a, 3a, 3a, 2b; n) = 2N(3a, 3a, 4a, 2b; 8n + 7a + 2b).
\end{aligned}$$

This yields (ii). By [S3, Theorem 4.5] and Theorem 2.3,

$$\begin{aligned}
& \frac{2}{3}(N(a, 3a, 3a, 2b; 8n + 7a + 2b) - N(a, 3a, 12a, 2b; 8n + 7a + 2b)) \\
&= t(a, 3a, 3a, 2b; n) = N(3a, 3a, 4a, 2b; 8n + 7a + 2b),
\end{aligned}$$

which yields part(iii). Hence the theorem is proved.

**Theorem 2.4.** *Let  $a, b, n \in \mathbb{Z}^+$  with  $2 \nmid ab$ . For  $n \in \mathbb{Z}^+$  with  $n \equiv \frac{a-b}{2} \pmod{2}$  we have*

$$t(a, 3a, b, 3b; n) = 4N(a, 3a, b, 3b; 2n + a + b).$$

Proof. Clearly,

$$\begin{aligned} \sum_{n=0}^{\infty} t(a, 3a, b, 3b; n)q^n &= 16\psi(q^a)\psi(q^{3a})\psi(q^b)\psi(q^{3b}) \\ &= 16(\varphi(q^{6a})\psi(q^{4a}) + q^a\varphi(q^{2a})\psi(q^{12a}))(\varphi(q^{6b})\psi(q^{4b}) + q^b\varphi(q^{2b})\psi(q^{12b})). \end{aligned}$$

Thus,

$$\begin{aligned} \sum_{n=0}^{\infty} t(a, 3a, b, 3b; 2n)q^{2n} \\ = 16\varphi(q^{6a})\psi(q^{4a})\varphi(q^{6b})\psi(q^{4b}) + 16q^{a+b}\varphi(q^{2a})\psi(q^{12a})\varphi(q^{2b})\psi(q^{12b}) \end{aligned}$$

and

$$\begin{aligned} \sum_{n=0}^{\infty} t(a, 3a, b, 3b; 2n+1)q^{2n+1} \\ = 16q^a\varphi(q^{2a})\psi(q^{12a})\varphi(q^{6b})\psi(q^{4b}) + 16q^b\varphi(q^{6a})\psi(q^{4a})\varphi(q^{2b})\psi(q^{12b}). \end{aligned}$$

Therefore,

$$(2.1) \quad \begin{aligned} \sum_{n=0}^{\infty} t(a, 3a, b, 3b; 2n)q^n \\ = 16\varphi(q^{3a})\psi(q^{2a})\varphi(q^{3b})\psi(q^{2b}) + 16q^{\frac{a+b}{2}}\varphi(q^a)\psi(q^{6a})\varphi(q^b)\psi(q^{6b}) \end{aligned}$$

and

$$(2.2) \quad \begin{aligned} \sum_{n=0}^{\infty} t(a, 3a, b, 3b; 2n+1)q^n \\ = 16q^{\frac{a-1}{2}}\varphi(q^a)\psi(q^{6a})\varphi(q^{3b})\psi(q^{2b}) + 16q^{\frac{b-1}{2}}\varphi(q^{3a})\psi(q^{2a})\varphi(q^b)\psi(q^{6b}). \end{aligned}$$

On the other hand,

$$\begin{aligned} \sum_{n=0}^{\infty} N(a, 3a, b, 3b; n)q^n &= \varphi(q^a)\varphi(q^{3a})\varphi(q^b)\varphi(q^{3b}) \\ &= (\varphi(q^{4a}) + 2q^a\psi(q^{8a}))(\varphi(q^{12a}) + 2q^{3a}\psi(q^{24a})) \\ &\quad \times (\varphi(q^{4b}) + 2q^b\psi(q^{8b}))(\varphi(q^{12b}) + 2q^{3b}\psi(q^{24b})). \end{aligned}$$

Collecting the terms of the form  $q^{4n+2}$  yields for  $a \equiv -b \pmod{4}$ ,

$$\sum_{n=0}^{\infty} N(a, 3a, b, 3b; 4n+2)q^{4n+2}$$

$$\begin{aligned}
&= 2q^a \psi(q^{8a}) \cdot \varphi(q^{12a}) \cdot \varphi(q^{4b}) \cdot 2q^{3b} \psi(q^{24b}) \\
&\quad + 2q^b \psi(q^{8b}) \cdot \varphi(q^{12b}) \cdot \varphi(q^{4a}) \cdot 2q^{3a} \psi(q^{24a}) \\
&= 4q^{a+b+2} (q^{2b-2} \varphi(q^{4b}) \psi(q^{24b}) \psi(q^{8a}) \varphi(q^{12a}) + q^{2a-2} \varphi(q^{4a}) \psi(q^{24a}) \psi(q^{8b}) \varphi(q^{12b})).
\end{aligned}$$

Hence

$$\begin{aligned}
(2.3) \quad &\sum_{n=0}^{\infty} N(a, 3a, b, 3b; 4n+2) q^n \\
&= 4q^{\frac{a+b}{4}} (q^{\frac{b-1}{2}} \varphi(q^b) \psi(q^{6b}) \psi(q^{2a}) \varphi(q^{3a}) + q^{\frac{a-1}{2}} \varphi(q^a) \psi(q^{6a}) \psi(q^{2b}) \varphi(q^{3b})).
\end{aligned}$$

Similarly, for  $a \equiv b \pmod{4}$  we have

$$\begin{aligned}
&\sum_{n=0}^{\infty} N(a, 3a, b, 3b; 4n+2) q^{4n+2} \\
&= 2q^a \psi(q^{8a}) \cdot \varphi(q^{12a}) \cdot 2q^b \psi(q^{8b}) \cdot \varphi(q^{12b}) + \varphi(q^{4a}) \cdot 2q^{3a} \psi(q^{24a}) \cdot \varphi(q^{4b}) \cdot 2q^{3b} \psi(q^{24b}) \\
&= 4q^{a+b} (\varphi(q^{12a}) \varphi(q^{12b}) \psi(q^{8a}) \psi(q^{8b}) + q^{2a+2b} \varphi(q^{4a}) \varphi(q^{4b}) \psi(q^{24a}) \psi(q^{24b}))
\end{aligned}$$

and so

$$\begin{aligned}
(2.4) \quad &\sum_{n=0}^{\infty} N(a, 3a, b, 3b; 4n+2) q^n \\
&= 4q^{\frac{a+b-2}{4}} (\varphi(q^{3a}) \varphi(q^{3b}) \psi(q^{2a}) \psi(q^{2b}) + q^{\frac{a+b}{2}} \varphi(q^a) \varphi(q^b) \psi(q^{6a}) \psi(q^{6b})).
\end{aligned}$$

For  $a \equiv -b \pmod{4}$  combining (2.2) with (2.3) gives

$$\sum_{n=0}^{\infty} N(a, 3a, b, 3b; 4n+2) q^n = \frac{4}{16} q^{\frac{a+b}{4}} \sum_{n=0}^{\infty} t(a, 3a, b, 3b; 2n+1) q^n$$

and so

$$t(a, 3a, b, 3b; 2n+1) = 4N(a, 3a, b, 3b; 4n+2+a+b).$$

For  $a \equiv b \pmod{4}$  combining (2.1) with (2.4) gives

$$\sum_{n=0}^{\infty} N(a, 3a, b, 3b; 4n+2) q^n = \frac{4}{16} q^{\frac{a+b-2}{4}} \sum_{n=0}^{\infty} t(a, 3a, b, 3b; 2n) q^n$$

and so

$$t(a, 3a, b, 3b; 2n) = 4N(a, 3a, b, 3b; 4n+a+b).$$

This completes the proof.

**Theorem 2.5.** *Let  $a, b \in \mathbb{Z}^+$  with  $ab \equiv -1 \pmod{4}$ . For  $n \in \mathbb{Z}^+$  we have*

$$t(2a, 2a, 3a, b; n) = \frac{1}{3} N(a, 3a, 16a, 4b; 32n + 28a + 4b).$$

Proof. By (1.10),

$$\sum_{n=0}^{\infty} N(a, 3a, 16a, 4b; n) q^n = \varphi(q^a) \varphi(q^{3a}) \varphi(q^{16a}) \varphi(q^{4b})$$



$$\begin{aligned}
&= (\varphi(q^{16a})\varphi(q^{48a}) + 4q^{16a}\psi(q^{32a})\psi(q^{96a}) + 2q^a\varphi(q^{48a})\psi(q^{8a}) + 2q^{3a}\varphi(q^{16a})\psi(q^{24a}) \\
&\quad + 6q^{4a}\psi(q^{8a})\psi(q^{24a}) + 4q^{13a}\psi(q^{8a})\psi(q^{96a}) + 4q^{7a}\psi(q^{24a})\psi(q^{32a})) \\
&\quad \times \varphi(q^{16a})(\varphi(q^{16b}) + 2q^{4b}\psi(q^{32b})).
\end{aligned}$$

Collecting the terms of the form  $q^{8n}$  yields

$$\begin{aligned}
&\sum_{n=0}^{\infty} N(a, 3a, 16a, 4b; 8n)q^{8n} \\
&= \varphi(q^{16a})^2\varphi(q^{48a})\varphi(q^{16b}) + 4q^{16a}\varphi(q^{16a})\psi(q^{32a})\psi(q^{96a})\varphi(q^{16b}) \\
&\quad + 12q^{4a+4b}\varphi(q^{16a})\psi(q^{8a})\psi(q^{24a})\psi(q^{32b})
\end{aligned}$$

and so

$$\begin{aligned}
&\sum_{n=0}^{\infty} N(a, 3a, 16a, 4b; 8n)q^n \\
&= \varphi(q^{2a})^2\varphi(q^{6a})\varphi(q^{2b}) + 4q^{2a}\varphi(q^{2a})\psi(q^{4a})\psi(q^{12a})\varphi(q^{2b}) \\
&\quad + 12q^{(a+b)/2}\varphi(q^{2a})\psi(q^a)\psi(q^{3a})\psi(q^{4b}) \\
&= \varphi(q^{2a})^2\varphi(q^{6a})\varphi(q^{2b}) + 4q^{2a}\varphi(q^{2a})\psi(q^{4a})\psi(q^{12a})\varphi(q^{2b}) \\
&\quad + 12q^{(a+b)/2}\varphi(q^{2a})\psi(q^{4b})(\varphi(q^{6a})\psi(q^{4a}) + q^a\varphi(q^{2a})\psi(q^{12a})).
\end{aligned}$$

Therefore

$$\sum_{n=0}^{\infty} N(a, 3a, 16a, 4b; 8(2n+1))q^{2n+1} = 12q^{(a+b)/2+a}\varphi(q^{2a})^2\psi(q^{12a})\psi(q^{4b})$$

and so

$$\begin{aligned}
&\sum_{n=0}^{\infty} N(a, 3a, 16a, 4b; 8(2n+1))q^n \\
&= 12q^{(a+b)/4+(a-1)/2}\varphi(q^a)^2\psi(q^{6a})\psi(q^{2b}) \\
&= 12q^{(a+b)/4+(a-1)/2}(\varphi(q^{2a})^2 + 4q^a\psi(q^{4a})^2)\psi(q^{6a})\psi(q^{2b}).
\end{aligned}$$

This yields

$$\sum_{n=0}^{\infty} N(a, 3a, 16a, 4b; 16n+12a+4b)q^n = 12\varphi(q^{2a})^2\psi(q^{6a})\psi(q^{2b}) + 48q^a\psi(q^{4a})^2\psi(q^{6a})\psi(q^{2b}).$$

Hence

$$\begin{aligned}
&\sum_{n=0}^{\infty} N(a, 3a, 16a, 4b; 32n+12a+b)q^{2n} = 12\varphi(q^{2a})^2\psi(q^{6a})\psi(q^{2b}), \\
&\sum_{n=0}^{\infty} N(a, 3a, 16a, 4b; 16(2n+1)+12a+4b)q^{2n+1} = 48q^a\psi(q^{4a})^2\psi(q^{6a})\psi(q^{2b})
\end{aligned}$$

and so

$$\sum_{n=0}^{\infty} N(a, 3a, 16a, 4b; 32n+12a+4b)q^n = 12\varphi(q^a)^2\psi(q^{3a})\psi(q^b),$$

$$\sum_{n=0}^{\infty} N(a, 3a, 16a, 4b; 16(2n+1) + 12a + 4b)q^n = 48q^{(a-1)/2}\psi(q^{2a})^2\psi(q^{3a})\psi(q^b).$$

Thus,

$$\sum_{n=0}^{\infty} N(a, 3a, 16a, 4b; 32n + 28a + 4b)q^n = 48\psi(q^{2a})^2\psi(q^{3a})\psi(q^b).$$

This yields

$$N(a, 3a, 16a, 4b; 32n + 28a + 4b) = 3t(2a, 2a, 3a, b; n).$$

Putting all the above together proves the theorem.

**Theorem 2.6.** *Let  $a, b \in \mathbb{Z}^+$  with  $ab \equiv 1 \pmod{4}$ . For  $n \in \mathbb{Z}^+$  we have*

$$t(a, 6a, 6a, b; n) = \frac{1}{3}N(a, 3a, 48a, 4b; 32n + 52a + 4b).$$

Proof. By (1.6),

$$\begin{aligned} \sum_{n=0}^{\infty} N(a, 3a, 48a, 4b; n)q^n &= \varphi(q^a)\varphi(q^{3a})\varphi(q^{48a})\varphi(q^{4b}) \\ &= (\varphi(q^{4a}) + 2q^{4a}\psi(q^{8a}))(\varphi(q^{12a}) + 2q^{12a}\psi(q^{24a}))\varphi(q^{48a})\varphi(q^{4b}) \\ &= (\varphi(q^{4a})\varphi(q^{12a}) + 4q^{4a}\psi(q^{8a})\psi(q^{24a}) + 2q^a(\varphi(q^{12a})\psi(q^{8a}) + q^{2a}\varphi(q^{4a})\psi(q^{24a}))) \\ &\quad \times \varphi(q^{48a})\varphi(q^{4b}). \end{aligned}$$

Thus,

$$\sum_{n=0}^{\infty} N(a, 3a, 48a, 4b; 4n)q^{4n} = (\varphi(q^{4a})\varphi(q^{12a}) + 4q^{4a}\psi(q^{8a})\psi(q^{24a}))\varphi(q^{48a})\varphi(q^{4b}).$$

Appealing to (1.6) and (1.8) we see that

$$\begin{aligned} \sum_{n=0}^{\infty} N(a, 3a, 48a, 4b; 4n)q^n &= (\varphi(q^a)\varphi(q^{3a}) + 4q^a\psi(q^{2a})\psi(q^{6a}))\varphi(q^{12a})\varphi(q^b) \\ &= (\varphi(q^{4a})\varphi(q^{12a}) + 4q^{4a}\psi(q^{8a})\psi(q^{24a}) + 6q^a(\varphi(q^{12a})\psi(q^{8a}) + q^{2a}\varphi(q^{4a})\psi(q^{24a}))) \\ &\quad \times \varphi(q^{12a})(\varphi(q^{4b}) + 2q^b\psi(q^{8b})). \end{aligned}$$

Therefore,

$$\sum_{n=0}^{\infty} N(a, 3a, 48a, 4b; 4(4n + a + b))q^{4n+a+b} = 6q^a\varphi(q^{12a})\psi(q^{8a}) \cdot \varphi(q^{12a}) \cdot 2q^b\psi(q^{8b})$$

and so

$$\begin{aligned} \sum_{n=0}^{\infty} N(a, 3a, 48a, 4b; 4(4n + a + b))q^n \\ = 12\varphi(q^{3a})^2\psi(q^{2a})\psi(q^{2b}) = 12(\varphi(q^{6a})^2 + 4q^{3a}\psi(q^{12a})^2)\psi(q^{2a})\psi(q^{2b}). \end{aligned}$$

It then follows that

$$\sum_{n=0}^{\infty} N(a, 3a, 48a, 4b; 4(4(2n+3a)+a+b))q^{2n+3a} = 48q^{3a}\psi(q^{12a})^2\psi(q^{2a})\psi(q^{2b})$$

and therefore

$$\sum_{n=0}^{\infty} N(a, 3a, 48a, 4b; 4(4(2n+3a)+a+b))q^n = 48\psi(q^a)\psi(q^{6a})^2\psi(q^b) = 3 \sum_{n=0}^{\infty} t(a, 6a, 6a, b; n)q^n.$$

This yields the result.

For  $n \in \{0, 1, 2, \dots\}$  let

$$r_3(n) = N(1, 1, 1; n) = |\{(x, y, z) \in \mathbb{Z}^3 \mid n = x^2 + y^2 + z^2\}|.$$

**Theorem 2.7.** *For  $n \in \mathbb{Z}^+$  we have*

$$\begin{aligned} & t(1, 4, 4; n) - \frac{1}{2}N(1, 4, 4; 8n+9) \\ &= t(1, 4, 4; n) - \frac{1}{6}r_3(8n+9) \\ &= \begin{cases} (-1)^{\frac{m+1}{2}}m & \text{if } 8n+9 = m^2 \text{ for some } m \in \mathbb{Z}^+, \\ 0 & \text{otherwise.} \end{cases} \end{aligned}$$

Proof. Since

$$\begin{aligned} \sum_{n=0}^{\infty} t(1, 1, 8; n)q^n &= 8\psi(q)^2\psi(q^8) = 8\varphi(q)\psi(q^2)\psi(q^8) \\ &= 8(\varphi(q^4) + 2q\psi(q^8))\psi(q^2)\psi(q^8), \end{aligned}$$

we see that

$$\sum_{n=0}^{\infty} t(1, 1, 8; 2n+1)q^{2n+1} = 16q\psi(q^2)\psi(q^8)^2$$

and so

$$\sum_{n=0}^{\infty} t(1, 1, 8; 2n+1)q^n = 16\psi(q)\psi(q^4)^2 = 2 \sum_{n=0}^{\infty} t(1, 4, 4; n)q^n.$$

Thus,

$$t(1, 4, 4; n) = \frac{1}{2}t(1, 1, 8; 2n+1).$$

By [S2, Theorem 3.1],

$$t(1, 1, 8; 2n+1) - \frac{1}{3}r_3(8n+9) = \begin{cases} 2(-1)^{\frac{m+1}{2}}m & \text{if } 8n+9 = m^2 \text{ for some } m \in \mathbb{Z}^+, \\ 0 & \text{otherwise.} \end{cases}$$

Thus,

$$(2.5) \quad t(1, 4, 4; n) - \frac{1}{6}r_3(8n+9) = \begin{cases} (-1)^{\frac{m+1}{2}}m & \text{if } 8n+9 = m^2 \text{ for some } m \in \mathbb{Z}^+, \\ 0 & \text{otherwise.} \end{cases}$$

On the other hand,

$$\sum_{n=0}^{\infty} N(1, 4, 4; n)q^n = \varphi(q)\varphi(q^4)^2 = (\varphi(q^{16}) + 2q^4\psi(q^{32}) + 2q\psi(q^8))(\varphi(q^8)^2 + 4q^4\psi(q^{16})^2).$$

Thus,

$$\sum_{n=0}^{\infty} N(1, 4, 4; 8n + 1)q^{8n+1} = 2q\varphi(q^8)^2\psi(q^8)$$

and so

$$\sum_{n=0}^{\infty} N(1, 4, 4; 8n + 1)q^n = 2\varphi(q)^2\psi(q).$$

Hence

$$(2.6) \quad \sum_{n=0}^{\infty} N(1, 4, 4; 8n + 9)q^n = \frac{2\varphi(q)^2\psi(q) - 2}{q}.$$

By [S2, (2.13)],

$$\sum_{n=0}^{\infty} r_3(4n + 1)q^n = 6\varphi(q)\psi(q)^2 = 6\varphi(q)^2\psi(q^2) = 6(\varphi(q^2)^2 + 4q\psi(q^4)^2)\psi(q^2).$$

Hence

$$\sum_{n=0}^{\infty} r_3(8n + 1)q^{2n} = 6\varphi(q^2)^2\psi(q^2)$$

and so

$$\sum_{n=0}^{\infty} r_3(8n + 1)q^n = 6\varphi(q)^2\psi(q).$$

It then follows that

$$(2.7) \quad \sum_{n=0}^{\infty} r_3(8n + 9)q^n = \frac{6\varphi(q)^2\psi(q) - 6}{q}.$$

Combining (2.6) with (2.7) gives  $r_3(8n + 9) = 3N(1, 4, 4; 8n + 9)$ . Now putting all the above together proves the theorem.

**Corollary 2.5.** *Let  $n \in \mathbb{Z}^+$  with  $n \equiv 1 \pmod{3}$ . Then*

$$t(1, 4, 4; n) = \frac{1}{2}N(1, 4, 4; 8n + 9).$$

**Theorem 2.8.** *Suppose  $a, b, n \in \mathbb{Z}^+$ ,  $(a, b) = 1$  and there is an odd prime divisor  $p$  of  $b$  such that  $(\frac{a(8n+9a)}{p}) = -1$ . Then*

$$t(a, 4a, 4a, b; n) = \frac{1}{2}(N(a, 4a, 4a, b; 8n + 9a + b) - N(a, 4a, 4a, 4b; 8n + 9a + b)).$$

Proof. By Theorem 2.7,

$$t(a, 4a, 4a, b; n)$$

$$\begin{aligned}
&= \sum_{w \in \mathbb{Z}} t(a, 4a, 4a; n - bw(w-1)/2) = \sum_{w \in \mathbb{Z}} t\left(1, 4, 4; \frac{n - bw(w-1)/2}{a}\right) \\
&= \frac{1}{2} \sum_{w \in \mathbb{Z}} N\left(1, 4, 4; 8 \frac{n - bw(w-1)/2}{a} + 9\right) \\
&= \frac{1}{2} \sum_{w \in \mathbb{Z}} N(a, 4a, 4a; 8n - 4bw(w-1) + 9a) \\
&= \frac{1}{2} \sum_{w \in \mathbb{Z}} N(a, 4a, 4a; 8n + 9a + b - b(2w-1)^2) \\
&= \frac{1}{2} \sum_{w \in \mathbb{Z}} (N(a, 4a, 4a; 8n + 9a + b - bw^2) - N(a, 4a, 4a; 8n + 9a + b - b(2w)^2)) \\
&= \frac{1}{2} (N(a, 4a, 4a, b; 8n + 9a + b) - N(a, 4a, 4a, 4b; 8n + 9a + b)).
\end{aligned}$$

This proves the theorem.

**Theorem 2.9.** *Suppose  $a, b, n \in \mathbb{Z}^+$ ,  $(a, b) = 1$ ,  $b \not\equiv 0, -a \pmod{4}$  and there is an odd prime divisor  $p$  of  $b$  such that  $\left(\frac{a(8n+9a)}{p}\right) = -1$ . Then*

$$t(a, 4a, 4a, b; n) = \frac{1}{2} N(a, 4a, 4a, b; 8n + 9a + b).$$

*Proof.* Suppose  $8n + 9a + b = ax^2 + 4ay^2 + 4az^2 + 4bw^2$  for some  $x, y, z, w \in \mathbb{Z}$ . Then  $ax^2 \equiv 9a + b \equiv a + b \pmod{4}$  and so  $b \equiv a(x^2 - 1) \equiv 0, -a \pmod{4}$ . But,  $b \not\equiv 0, -a \pmod{4}$ . Hence,  $N(a, 4a, 4a, 4b; 8n + 9a + b) = 0$ . Now the result follows from Theorem 2.8.

**Corollary 2.6.** *Suppose  $a, b, n \in \mathbb{Z}^+$ ,  $3 \nmid a$ ,  $3 \mid b$ ,  $b \not\equiv 0, 3a \pmod{4}$  and  $3 \mid n - a$ . Then*

$$t(a, 4a, 4a, b; n) = \frac{1}{2} N(a, 4a, 4a, b; 8n + 9a + b).$$

**Corollary 2.7.** *Suppose  $m, n \in \mathbb{Z}^+$ ,  $m \equiv 1, 2 \pmod{4}$  and  $n \equiv 1, 3 \pmod{5}$ . Then*

$$t(1, 4, 4, 5m; n) = \frac{1}{2} N(1, 4, 4, 5m; 8n + 5m + 9).$$

**Theorem 2.10.** *Suppose  $a, b, n \in \mathbb{Z}^+$ ,  $(a, b) = 1$  and there is an odd prime divisor  $p$  of  $b$  such that  $\left(\frac{a(4n+5a)}{p}\right) = -1$ . Then*

$$t(a, a, 8a, b; n) = \frac{1}{2} (N(a, a, 8a, b; 8n + 10a + b) - N(a, a, 8a, 4b; 8n + 10a + b)).$$

*Proof.* By [S2, Theorem 3.1],

$$\begin{aligned}
&t(a, a, 8a, b; n) \\
&= \sum_{w \in \mathbb{Z}} t(a, a, 8a; n - bw(w-1)/2) = \sum_{w \in \mathbb{Z}} t\left(1, 1, 8; \frac{n - bw(w-1)/2}{a}\right) \\
&= \frac{1}{2} \sum_{w \in \mathbb{Z}} N\left(1, 1, 8; 8 \frac{n - bw(w-1)/2}{a} + 10\right)
\end{aligned}$$

$$\begin{aligned}
&= \frac{1}{2} \sum_{w \in \mathbb{Z}} N(a, a, 8a; 8n - 4bw(w-1) + 10a) \\
&= \frac{1}{2} \sum_{w \in \mathbb{Z}} N(a, a, 8a; 8n + 10a + b - b(2w-1)^2) \\
&= \frac{1}{2} \sum_{w \in \mathbb{Z}} (N(a, a, 8a; 8n + 10a + b - bw^2) - N(a, a, 8a; 8n + 10a + b - b(2w)^2)) \\
&= \frac{1}{2} (N(a, a, 8a, b; 8n + 10a + b) - N(a, a, 8a, 4b; 8n + 10a + b)).
\end{aligned}$$

This proves the theorem.

**Theorem 2.11.** *Suppose  $a, b, n \in \mathbb{Z}^+$ ,  $(a, b) = 1$  and there is an odd prime divisor  $p$  of  $b$  such that  $(\frac{a(4n+5a)}{p}) = -1$ . Assume that  $a$  is even or  $ab \equiv 1, 4, 5 \pmod{8}$  for odd  $a$ . Then*

$$t(a, a, 8a, b; n) = \frac{1}{2} N(a, a, 8a, b; 8n + 10a + b).$$

*Proof.* Suppose  $8n + 10a + b = ax^2 + ay^2 + 8az^2 + 4bw^2$  for some  $x, y, z, w \in \mathbb{Z}$ . Then  $a(x^2 + y^2) \equiv 2a + b \pmod{4}$ . If  $2 \mid a$ , then  $2 \nmid b$  and so  $a(x^2 + y^2) \not\equiv 2a + b \pmod{4}$ . We get a contradiction. If  $4 \mid ab - 1$ , then  $a \equiv b \pmod{4}$  and so  $a(x^2 + y^2) \equiv 3a \pmod{4}$ . Hence  $x^2 + y^2 \equiv 3 \pmod{4}$ . This is impossible since  $t^2 \equiv 0, 1 \pmod{4}$  for  $t \in \mathbb{Z}$ . If  $2 \nmid a$  and  $ab \equiv 4 \pmod{8}$ , then  $2 \mid b$  and so  $a(x^2 + y^2) \equiv 2a + b \pmod{8}$ . This yields  $x^2 + y^2 \equiv 2a^2 + ab \equiv ab + 2 \equiv 6 \pmod{8}$ . Since  $t^2 \equiv 0, 1, 4 \pmod{8}$  for  $t \in \mathbb{Z}$  we see that  $x^2 + y^2 \not\equiv 6 \pmod{8}$  and get a contradiction. By the above,  $N(a, a, 8a, 4b; 8n + 10a + b) = 0$ . Now the result follows from Theorem 2.10.

**Remark 2.1** Theorem 2.11 is a generalization of [S2, Theorem 3.2].

### 3. New transformation formulas for $t(a, b, c, d; n)$

In this section we present 31 transformation formulas for  $t(a, b, c, d; n)$ .

**Theorem 3.1.** *Let  $a, b, c \in \mathbb{Z}^+$  and  $2 \nmid a$ . For  $n \in \mathbb{Z}^+$  we have*

$$(3.1) \quad t(a, a, 2b, 2c; 2n + a) = 2t(a, 4a, b, c; n),$$

$$(3.2) \quad t(a, 3a, 4a, 2b; 2n + a) = t(a, a, 6a, b; n),$$

$$(3.3) \quad t(a, 3a, 12a, 2b; 2n) = t(2a, 3a, 3a, b; n).$$

*Proof.* Since

$$\begin{aligned}
\sum_{n=0}^{\infty} t(a, a, 2b, 2c; n) q^n &= 16\psi(q^a)^2 \psi(q^{2b}) \psi(q^{2c}) \\
&= 16\varphi(q^a) \psi(q^{2a}) \psi(q^{2b}) \psi(q^{2c}) = 16(\varphi(q^{4a}) + 2q^a \psi(q^{8a})) \psi(q^{2a}) \psi(q^{2b}) \psi(q^{2c})
\end{aligned}$$

we see that

$$\sum_{n=0}^{\infty} t(a, a, 2b, 2c; 2n + a) q^{2n+a} = 32q^a \psi(q^{8a}) \psi(q^{2a}) \psi(q^{2b}) \psi(q^{2c})$$

and so

$$\sum_{n=0}^{\infty} t(a, a, 2b, 2c; 2n + a)q^n = 32\psi(q^a)\psi(q^{4a})\psi(q^b)\psi(q^c) = 2\sum_{n=0}^{\infty} t(a, 4a, b, c; n)q^n,$$

which yields  $t(a, a, 2b, 2c; 2n + a) = 2t(a, 4a, b, c; n)$ .

It is clearly that

$$\begin{aligned} & \sum_{n=0}^{\infty} t(a, 3a, 4a, 2b; n)q^n \\ &= 16\psi(q^a)\psi(q^{3a})\psi(q^{4a})\psi(q^{2b}) = 16(\varphi(q^{6a})\psi(q^{4a}) + q^a\varphi(q^{2a})\psi(q^{12a}))\psi(q^{4a})\psi(q^{2b}). \end{aligned}$$

Thus,

$$\sum_{n=0}^{\infty} t(a, 3a, 4a, 2b; 2n+a)q^{2n+a} = 16q^a\varphi(q^{2a})\psi(q^{4a})\psi(q^{12a})\psi(q^{2b}) = 16q^a\psi(q^{2a})^2\psi(q^{12a})\psi(q^{2b})$$

and so

$$\sum_{n=0}^{\infty} t(a, 3a, 4a, 2b; 2n + a)q^n = 16\psi(q^a)^2\psi(q^{6a})\psi(q^b) = \sum_{n=0}^{\infty} t(a, a, 6a, b; n)q^n.$$

This yields  $t(a, 3a, 4a, 2b; 2n + a) = t(a, a, 6a, b; n)$ .

Note that

$$\begin{aligned} & \sum_{n=0}^{\infty} t(a, 3a, 12a, 2b; n)q^n \\ &= 16\psi(q^a)\psi(q^{3a})\psi(q^{12a})\psi(q^{2b}) = 16(\varphi(q^{6a})\psi(q^{4a}) + q^a\varphi(q^{2a})\psi(q^{12a}))\psi(q^{12a})\psi(q^{2b}). \end{aligned}$$

We have

$$\sum_{n=0}^{\infty} t(a, 3a, 12a, 2b; 2n)q^{2n} = 16\varphi(q^{6a})\psi(q^{4a})\psi(q^{12a})\psi(q^{2b}) = 16\psi(q^{6a})^2\psi(q^{4a})\psi(q^{2b})$$

and so

$$\sum_{n=0}^{\infty} t(a, 3a, 12a, 2b; 2n)q^n = 16\psi(q^{2a})\psi(q^{3a})^2\psi(q^b) = \sum_{n=0}^{\infty} t(2a, 3a, 3a, b; n)q^n,$$

which gives  $t(a, 3a, 12a, 2b; 2n) = t(2a, 3a, 3a, b; n)$ . This completes the proof.

**Theorem 3.2.** *Suppose  $a, b, c \in \mathbb{Z}^+$  and  $2 \nmid a$ . For  $n \in \mathbb{Z}^+$  we have*

$$(3.4) \quad t(a, 3a, 4b, 4c; 4n + 3a) = 2t(3a, 4a, b, c; n),$$

$$(3.5) \quad t(a, 3a, 4b, 4c; 4n + 6a) = 2t(a, 12a, b, c; n),$$

$$(3.6) \quad t(a, 3a, 48a, 4b; 4n) = t(a, 6a, 6a, b; n),$$

$$(3.7) \quad t(a, 3a, 16a, 4b; 4n + a) = t(2a, 2a, 3a, b; n),$$

$$(3.8) \quad t(2a, 3a, 3a, 4b; 4n + 3a) = 2t(2a, 3a, 3a, b; n).$$

Proof. It is clear that for  $|q| < 1$ ,

$$\begin{aligned} & \sum_{n=0}^{\infty} t(a, 3a, 4b, 4c; n)q^n \\ &= 16\psi(q^a)\psi(q^{3a})\psi(q^{4b})\psi(q^{4c}) = 16(\varphi(q^{6a})\psi(q^{4a}) + q^a\varphi(q^{2a})\psi(q^{12a}))\psi(q^{4b})\psi(q^{4c}) \\ &= 16((\varphi(q^{24a}) + 2q^{6a}\psi(q^{48a}))\psi(q^{4a}) + q^a(\varphi(q^{8a}) + 2q^{2a}\psi(q^{16a}))\psi(q^{12a}))\psi(q^{4b})\psi(q^{4c}). \end{aligned}$$

Thus,

$$\begin{aligned} & \sum_{n=0}^{\infty} t(a, 3a, 4b, 4c; 4n + 6a)q^{4n+6a} = 32q^{6a}\psi(q^{48a})\psi(q^{4a})\psi(q^{4b})\psi(q^{4c}), \\ & \sum_{n=0}^{\infty} t(a, 3a, 4b, 4c; 4n + 3a)q^{4n+3a} = 32q^{3a}\psi(q^{16a})\psi(q^{12a})\psi(q^{4b})\psi(q^{4c}), \\ & \sum_{n=0}^{\infty} t(a, 3a, 4b, 48a; 4n)q^{4n} = 16\varphi(q^{24a})\psi(q^{4a})\psi(q^{4b})\psi(q^{48a}), \\ & \sum_{n=0}^{\infty} t(a, 3a, 4b, 16a; 4n + a)q^{4n+a} = 16q^a\varphi(q^{8a})\psi(q^{12a})\psi(q^{4b})\psi(q^{16a}) \end{aligned}$$

and so

$$\begin{aligned} & \sum_{n=0}^{\infty} t(a, 3a, 4b, 4c; 4n + 6a)q^n = 32\psi(q^a)\psi(q^{12a})\psi(q^b)\psi(q^c) = 2 \sum_{n=0}^{\infty} t(a, 12a, b, c; n)q^n, \\ & \sum_{n=0}^{\infty} t(a, 3a, 4b, 4c; 4n + 3a)q^n = 32\psi(q^{4a})\psi(q^{3a})\psi(q^b)\psi(q^c) = 2 \sum_{n=0}^{\infty} t(3a, 4a, b, c; n)q^n, \\ & \sum_{n=0}^{\infty} t(a, 3a, 4b, 48a; 4n)q^n = 16\varphi(q^{6a})\psi(q^{12a})\psi(q^a)\psi(q^b) = 16\psi(q^a)\psi(q^{6a})^2\psi(q^b), \\ & \sum_{n=0}^{\infty} t(a, 3a, 4b, 16a; 4n + a)q^n = 16\varphi(q^{2a})\psi(q^{3a})\psi(q^b)\psi(q^{4a}) = 16\psi(q^{2a})^2\psi(q^{3a})\psi(q^b). \end{aligned}$$

This yields (3.4)-(3.7). To prove (3.8), appealing to Theorem 3.1 we see that

$$t(2a, 3a, 3a, 4b; 4n + 3a) = 2t(a, 3a, 12a, 2b; 2n) = 2t(2a, 3a, 3a, b; n).$$

**Theorem 3.3.** *Suppose  $a, b \in \mathbb{Z}^+$  with  $2 \nmid a$ . For  $n \in \mathbb{Z}^+$  we have*

$$(3.9) \quad t(2a, 3a, 3a, 8b; 8n + 9a) = 4t(3a, 3a, 4a, b; n),$$

$$(3.10) \quad t(a, a, 6a, 8b; 8n + 13a) = 4t(a, a, 12a, b; n),$$

$$(3.11) \quad t(a, a, 6a, 8b; 8n + 4a) = 2t(a, a, 3a, b; n),$$

$$(3.12) \quad t(a, a, 6a, 8b; 8n + 6a) = 4t(2a, 2a, 3a, b; n),$$

$$(3.13) \quad t(2a, 3a, 3a, 8b; 8n + 12a) = 4t(a, 6a, 6a, b; n),$$

$$(3.14) \quad t(2a, 3a, 3a, 8b; 8n + 6a) = 2t(a, 3a, 3a, b; n).$$



Proof. By Theorems 3.1 and 3.2,

$$\begin{aligned} t(2a, 3a, 3a, 8b; 8n + 9a) &= 2t(a, 3a, 12a, 4b; 4n + 3a) = 4t(3a, 3a, 4a, b; n), \\ t(a, a, 6a, 8b; 8n + 13a) &= 2t(a, 3a, 4a, 4b; 4n + 6a) = 4t(a, a, 12a, b; n). \end{aligned}$$

This proves (3.9) and (3.10). Since  $\psi(q)^2 = \varphi(q)\psi(q^2)$  and  $\varphi(q) = \varphi(q^4) + 2q\psi(q^8)$  we see that

$$\begin{aligned} &\sum_{n=0}^{\infty} t(a, a, 6a, 8b; n)q^n \\ &= 16\psi(q^a)^2\psi(q^{6a})\psi(q^{8b}) = 16\varphi(q^a)\psi(q^{2a})\psi(q^{6a})\psi(q^{8b}) \\ &= 16(\varphi(q^{4a}) + 2q^a\psi(q^{8a}))(\varphi(q^{12a})\psi(q^{8a}) + q^{2a}\varphi(q^{4a})\psi(q^{24a}))\psi(q^{8b}) \\ &= 16(\varphi(q^{16a}) + 2q^{4a}\psi(q^{32a}) + 2q^a\psi(q^{8a})) \\ &\quad \times (\varphi(q^{48a})\psi(q^{8a}) + 2q^{12a}\psi(q^{96a})\psi(q^{8a}) + q^{2a}\varphi(q^{16a})\psi(q^{24a}) + 2q^{6a}\psi(q^{24a})\psi(q^{32a}))\psi(q^{8b}). \end{aligned}$$

Thus,

$$\begin{aligned} &\sum_{n=0}^{\infty} t(a, a, 6a, 8b; 8n + 6a)q^{8n+6a} \\ &= 16\psi(q^{8b})(\varphi(q^{16a}) \cdot 2q^{6a}\psi(q^{24a})\psi(q^{32a}) + 2q^{4a}\psi(q^{32a}) \cdot q^{2a}\varphi(q^{16a})\psi(q^{24a})) \\ &= 64q^{6a}\varphi(q^{16a})\psi(q^{32a})\psi(q^{24a})\psi(q^{8b}) = 64q^{6a}\psi(q^{16a})^2\psi(q^{24a})\psi(q^{8b}) \end{aligned}$$

and

$$\begin{aligned} &\sum_{n=0}^{\infty} t(a, a, 6a, 8b; 8n + 4a)q^{8n+4a} \\ &= 16 \cdot 2q^{4a}(\varphi(q^{48a})\psi(q^{32a}) + q^{8a}\varphi(q^{16a})\psi(q^{96a}))\psi(q^{8a})\psi(q^{8b}) \\ &= 32q^{4a}\psi(q^{8a})^2\psi(q^{24a})\psi(q^{8b}). \end{aligned}$$

It then follows that

$$\begin{aligned} \sum_{n=0}^{\infty} t(a, a, 6a, 8b; 8n + 6a)q^n &= 64\psi(q^{2a})^2\psi(q^{3a})\psi(q^b) = 4 \sum_{n=0}^{\infty} t(2a, 2a, 3a, b; n)q^n, \\ \sum_{n=0}^{\infty} t(a, a, 6a, 8b; 8n + 4a)q^n &= 32\psi(q^a)^2\psi(q^{3a})\psi(q^b) = 2 \sum_{n=0}^{\infty} t(a, a, 3a, b; n)q^n, \end{aligned}$$

which yields (3.11) and (3.12). By (1.5), (1.6) and (1.8),

$$\begin{aligned} &\sum_{n=0}^{\infty} t(2a, 3a, 3a, 8b; n)q^n \\ &= 16\psi(q^{2a})\psi(q^{3a})^2\psi(q^{8b}) = 16\varphi(q^{3a})\psi(q^{2a})\psi(q^{6a})\psi(q^{8b}) \\ &= 16(\varphi(q^{12a}) + 2q^{3a}\psi(q^{24a}))(\varphi(q^{12a})\psi(q^{8a}) + q^{2a}\varphi(q^{4a})\psi(q^{24a}))\psi(q^{8b}). \end{aligned}$$

Thus

$$\sum_{n=0}^{\infty} t(2a, 3a, 3a, 8b; 4n)q^{4n} = 16\varphi(q^{12a})^2\psi(q^{8a})\psi(q^{8b}),$$

$$\sum_{n=0}^{\infty} t(2a, 3a, 3a, 8b; 4n + 2a)q^{4n+2a} = 16q^{2a}\varphi(q^{4a})\psi(q^{8b})\varphi(q^{12a})\psi(q^{24a})$$

and so

$$\begin{aligned} \sum_{n=0}^{\infty} t(2a, 3a, 3a, 8b; 4n)q^n &= 16\varphi(q^{3a})^2\psi(q^{2a})\psi(q^{2b}) \\ &= 16(\varphi(q^{6a})^2 + 4q^{3a}\psi(q^{12a})^2)\psi(q^{2a})\psi(q^{2b}), \\ \sum_{n=0}^{\infty} t(2a, 3a, 3a, 8b; 4n + 2a)q^n &= 16\varphi(q^a)\psi(q^{2b})\varphi(q^{3a})\psi(q^{6a}) \\ &= 16(\varphi(q^{4a}) + 2q^a\psi(q^{8a}))(\varphi(q^{12a}) + 2q^{3a}\psi(q^{24a}))\psi(q^{6a})\psi(q^{2b}). \end{aligned}$$

Hence,

$$\begin{aligned} \sum_{n=0}^{\infty} t(2a, 3a, 3a, 8b; 4(2n + 3a))q^{2n+3a} &= 64q^{3a}\psi(q^{2a})\psi(q^{12a})^2\psi(q^{2b}), \\ \sum_{n=0}^{\infty} t(2a, 3a, 3a, 8b; 4(2n + a) + 2a)q^{2n+a} \\ &= 32q^a(\varphi(q^{12a})\psi(q^{8a}) + q^{2a}\varphi(q^{4a})\psi(q^{24a}))\psi(q^{6a})\psi(q^{2b}) = 32q^a\psi(q^{2a})\psi(q^{6a})^2\psi(q^{2b}). \end{aligned}$$

This yields

$$\begin{aligned} \sum_{n=0}^{\infty} t(2a, 3a, 3a, 8b; 8n + 12a)q^n &= 64\psi(q^a)\psi(q^{6a})^2\psi(q^b) = 4 \sum_{n=0}^{\infty} t(a, 6a, 6a, b; n)q^n \\ \sum_{n=0}^{\infty} t(2a, 3a, 3a, 8b; 8n + 6a)q^n &= 32\psi(q^a)\psi(q^{3a})^2\psi(q^b) = 2 \sum_{n=0}^{\infty} t(a, 3a, 3a, b; n)q^n, \end{aligned}$$

which yields (3.13) and (3.14). The proof is now complete.

**Theorem 3.4.** *Suppose  $a, b, c \in \mathbb{Z}^+$  with  $2 \nmid a$ . For  $n \in \mathbb{Z}^+$  we have*

$$(3.15) \quad t(a, 7a, 2b, 2c; 2n + a) = t(a, 7a, b, c; n),$$

$$(3.16) \quad t(a, 7a, 8b, 8c; 8n + 10a) = 2t(4a, 7a, b, c; n),$$

$$(3.17) \quad t(a, 7a, 8b, 8c; 8n + 28a) = 2t(a, 28a, b, c; n),$$

$$(3.18) \quad t(a, 7a, 8a, 4b; 4n + 6a) = t(a, a, 14a, b; n),$$

$$(3.19) \quad t(a, 7a, 56a, 4b; 4n) = t(2a, 7a, 7a, b; n).$$

Proof. By [Be, p.315],

$$(3.20) \quad \psi(q)\psi(q^7) = \psi(q^8)\varphi(q^{28}) + q\psi(q^2)\psi(q^{14}) + q^6\varphi(q^4)\psi(q^{56}).$$

Thus,

$$(3.21) \quad \psi(q^2)\psi(q^{14}) = \psi(q^{16})\varphi(q^{56}) + q^2\psi(q^4)\psi(q^{28}) + q^{12}\varphi(q^8)\psi(q^{112}).$$

Combining (3.20) with (3.21) gives

$$(3.22) \quad \begin{aligned} \psi(q)\psi(q^7) &= \psi(q^8)\varphi(q^{28}) + q^6\varphi(q^4)\psi(q^{56}) + q\psi(q^{16})\varphi(q^{56}) \\ &\quad + q^3\psi(q^4)\psi(q^{28}) + q^{13}\varphi(q^8)\psi(q^{112}). \end{aligned}$$

Using (3.20) we see that

$$\begin{aligned} \sum_{n=0}^{\infty} t(a, 7a, 2b, 2c; n)q^n &= 16\psi(q^a)\psi(q^{7a})\psi(q^{2b})\psi(q^{2c}) \\ &= 16(\psi(q^{8a})\varphi(q^{28a}) + q^{6a}\varphi(q^{4a})\psi(q^{56a}) + q^a\psi(q^{2a})\psi(q^{14a}))\psi(q^{2b})\psi(q^{2c}) \end{aligned}$$

and so

$$\sum_{n=0}^{\infty} t(a, 7a, 2b, 2c; 2n+a)q^{2n+a} = 16q^a\psi(q^{2a})\psi(q^{14a})\psi(q^{2b})\psi(q^{2c}).$$

This yields

$$\sum_{n=0}^{\infty} t(a, 7a, 2b, 2c; 2n+a)q^n = 16\psi(q^a)\psi(q^{7a})\psi(q^b)\psi(q^c).$$

Hence (3.15) is true. Applying (3.22) we see that

$$\begin{aligned} \sum_{n=0}^{\infty} t(a, 7a, 4b, 4c; n)q^n &= 16\psi(q^a)\psi(q^{7a})\psi(q^{4b})\psi(q^{4c}) \\ &= 16(\psi(q^{8a})\varphi(q^{28a}) + q^{6a}\varphi(q^{4a})\psi(q^{56a}) + q^a\psi(q^{16a})\varphi(q^{56a}) \\ &\quad + q^{3a}\psi(q^{4a})\psi(q^{28a}) + q^{13a}\varphi(q^{8a})\psi(q^{112a}))\psi(q^{4b})\psi(q^{4c}). \end{aligned}$$

Thus,

$$\begin{aligned} \sum_{n=0}^{\infty} t(a, 7a, 4b, 4c; 4n)q^{4n} &= 16\psi(q^{8a})\varphi(q^{28a})\psi(q^{4b})\psi(q^{4c}), \\ \sum_{n=0}^{\infty} t(a, 7a, 4b, 4c; 4n+6a)q^{4n+6a} &= 16q^{6a}\varphi(q^{4a})\psi(q^{56a})\psi(q^{4b})\psi(q^{4c}) \end{aligned}$$

and so

$$(3.23) \quad \sum_{n=0}^{\infty} t(a, 7a, 4b, 4c; 4n)q^n = 16\psi(q^{2a})\varphi(q^{7a})\psi(q^b)\psi(q^c),$$

$$(3.24) \quad \sum_{n=0}^{\infty} t(a, 7a, 4b, 4c; 4n+6a)q^n = 16\varphi(q^a)\psi(q^{14a})\psi(q^b)\psi(q^c).$$

By (3.24) and (1.6),

$$\sum_{n=0}^{\infty} t(a, 7a, 8b, 8c; 4n+6a)q^n = 16(\varphi(q^{4a}) + 2q^a\psi(q^{8a}))\psi(q^{14a})\psi(q^{2b})\psi(q^{2c}).$$

Therefore

$$\sum_{n=0}^{\infty} t(a, 7a, 8b, 8c; 4(2n+a)+6a)q^{2n+a} = 32q^a\psi(q^{8a})\psi(q^{14a})\psi(q^{2b})\psi(q^{2c})$$

and so

$$\sum_{n=0}^{\infty} t(a, 7a, 8b, 8c; 4(2n + a) + 6a)q^n = 32\psi(q^{4a})\psi(q^{7a})\psi(q^b)\psi(q^c).$$

This yields (3.16). Similarly, from (3.23) and (1.6),

$$\sum_{n=0}^{\infty} t(a, 7a, 8b, 8c; 4n)q^n = 16(\varphi(q^{28a}) + 2q^{7a}\psi(q^{56a}))\psi(q^{2a})\psi(q^{2b})\psi(q^{2c}).$$

Therefore,

$$\sum_{n=0}^{\infty} t(a, 7a, 8b, 8c; 4(2n + 7a))q^{2n+7a} = 32q^{7a}\psi(q^{56a})\psi(q^{2a})\psi(q^{2b})\psi(q^{2c})$$

and so

$$\sum_{n=0}^{\infty} t(a, 7a, 8b, 8c; 4(2n + 7a))q^n = 32\psi(q^a)\psi(q^{28a})\psi(q^b)\psi(q^c),$$

which gives (3.17). Recall that  $\varphi(q)\psi(q^2) = \psi(q)^2$ . From (3.23) and (3.24) we see that

$$\begin{aligned} \sum_{n=0}^{\infty} t(a, 7a, 56a, 4b; 4n)q^n &= 16\psi(q^{2a})\varphi(q^{7a})\psi(q^{14a})\psi(q^b) = 16\psi(q^{2a})\psi(q^{7a})^2\psi(q^b), \\ \sum_{n=0}^{\infty} t(a, 7a, 8a, 4b; 4n + 6a)q^n &= 16\varphi(q^a)\psi(q^{2a})\psi(q^{14a})\psi(q^b) = 16\psi(q^a)^2\psi(q^{14a})\psi(q^b), \end{aligned}$$

which yields (3.18) and (3.19). The proof is now complete.

From [Be, p.377] we know that for  $|q| < 1$ ,

$$(3.25) \quad \psi(q^3)\psi(q^5) = \varphi(q^{60})\psi(q^8) + q^3\psi(q^2)\psi(q^{30}) + q^{14}\varphi(q^4)\psi(q^{120}).$$

From [XZ, (2.10)] we know that

$$(3.26) \quad \psi(q)\psi(q^{15}) = \psi(q^6)\psi(q^{10}) + q\varphi(q^{20})\psi(q^{24}) + q^3\varphi(q^{12})\psi(q^{40}).$$

Thus,

$$\psi(q^2)\psi(q^{30}) = \psi(q^{12})\psi(q^{20}) + q^2\varphi(q^{40})\psi(q^{48}) + q^6\varphi(q^{24})\psi(q^{80}).$$

This together with (3.25) yields

$$(3.27) \quad \begin{aligned} \psi(q^3)\psi(q^5) &= \varphi(q^{60})\psi(q^8) + q^{14}\varphi(q^4)\psi(q^{120}) + q^3\psi(q^{12})\psi(q^{20}) \\ &\quad + q^5\varphi(q^{40})\psi(q^{48}) + q^9\varphi(q^{24})\psi(q^{80}). \end{aligned}$$

By (3.25),

$$\psi(q^6)\psi(q^{10}) = \varphi(q^{120})\psi(q^{16}) + q^6\psi(q^4)\psi(q^{60}) + q^{28}\varphi(q^8)\psi(q^{240}).$$

Combining this with (3.26) gives

$$(3.28) \quad \begin{aligned} \psi(q)\psi(q^{15}) &= \varphi(q^{120})\psi(q^{16}) + q^{28}\varphi(q^8)\psi(q^{240}) + q^6\psi(q^4)\psi(q^{60}) \\ &\quad + q\varphi(q^{20})\psi(q^{24}) + q^3\varphi(q^{12})\psi(q^{40}). \end{aligned}$$

**Theorem 3.5.** *Suppose  $a, b, c \in \mathbb{Z}^+$  with  $2 \nmid a$ . For  $n \in \mathbb{Z}^+$  we have*

$$(3.29) \quad t(3a, 5a, 2b, 2c; 2n + 3a) = t(a, 15a, b, c; n),$$

$$(3.30) \quad t(a, 15a, 2b, 2c; 2n) = t(3a, 5a, b, c; n),$$

$$(3.31) \quad t(3a, 5a, 4b, 4c; 4n + 3a) = t(3a, 5a, b, c; n),$$

$$(3.32) \quad t(a, 15a, 4b, 4c; 4n + 6a) = t(a, 15a, b, c; n),$$

$$(3.33) \quad t(3a, 5a, 8b, 8c; 8n + 18a) = 2t(4a, 15a, b, c; n),$$

$$(3.34) \quad t(3a, 5a, 8b, 8c; 8n + 60a) = 2t(a, 60a, b, c; n),$$

$$(3.35) \quad t(3a, 5a, 8a, 4b; 4n + 14a) = t(a, a, 30a, b; n),$$

$$(3.36) \quad t(3a, 5a, 120a, 4b; 4n) = t(2a, 15a, 15a, b; n),$$

$$(3.37) \quad t(a, 15a, 8b, 8c; 8n + 15a) = 2t(5a, 12a, b, c; n),$$

$$(3.38) \quad t(a, 15a, 8b, 8c; 8n + 21a) = 2t(3a, 20a, b, c; n),$$

$$(3.39) \quad t(a, 15a, 24a, 4b; 4n + 3a) = t(3a, 3a, 10a, b; n),$$

$$(3.40) \quad t(a, 15a, 40a, 4b; 4n + a) = t(5a, 5a, 6a, b; n).$$

Proof. Using (3.25) and (3.26) we see that

$$\begin{aligned} \sum_{n=0}^{\infty} t(3a, 5a, 2b, 2c; n)q^n &= 16\psi(q^{3a})\psi(q^{5a})\psi(q^{2b})\psi(q^{2c}) \\ &= 16(\varphi(q^{60a})\psi(q^{8a}) + q^{14a}\varphi(q^{4a})\psi(q^{120a}) + q^{3a}\psi(q^{2a})\psi(q^{30a}))\psi(q^{2b})\psi(q^{2c}), \\ \sum_{n=0}^{\infty} t(a, 15a, 2b, 2c; n)q^n &= 16\psi(q^a)\psi(q^{15a})\psi(q^{2b})\psi(q^{2c}) \\ &= 16(\psi(q^{6a})\psi(q^{10a}) + q^a\varphi(q^{20a})\psi(q^{24a}) + q^{3a}\varphi(q^{12a})\psi(q^{40a}))\psi(q^{2b})\psi(q^{2c}). \end{aligned}$$

Hence

$$\begin{aligned} \sum_{n=0}^{\infty} t(3a, 5a, 2b, 2c; 2n + 3a)q^{2n+3a} &= 16q^{3a}\psi(q^{2a})\psi(q^{30a})\psi(q^{2b})\psi(q^{2c}), \\ \sum_{n=0}^{\infty} t(a, 15a, 2b, 2c; 2n)q^{2n} &= 16\psi(q^{6a})\psi(q^{10a})\psi(q^{2b})\psi(q^{2c}). \end{aligned}$$

It then follows that

$$\begin{aligned} \sum_{n=0}^{\infty} t(3a, 5a, 2b, 2c; 2n + 3a)q^n &= 16\psi(q^a)\psi(q^{15a})\psi(q^b)\psi(q^c), \\ \sum_{n=0}^{\infty} t(a, 15a, 2b, 2c; 2n)q^n &= 16\psi(q^{3a})\psi(q^{5a})\psi(q^b)\psi(q^c), \end{aligned}$$

which implies (3.29) and (3.30). By (3.29) and (3.30),

$$\begin{aligned} t(3a, 5a, 4b, 4c; 4n + 3a) &= t(a, 15a, 2b, 2c; 2n) = t(3a, 5a, b, c; n), \\ t(a, 15a, 4b, 4c; 4n + 6a) &= t(3a, 5a, 2b, 2c; 2n + 3a) = t(a, 15a, b, c; n). \end{aligned}$$

Thus, (3.31) and (3.32) hold. Appealing to (3.27),

$$\begin{aligned} \sum_{n=0}^{\infty} t(3a, 5a, 4b, 4c; n)q^n &= 16\psi(q^{3a})\psi(q^{5a})\psi(q^{4b})\psi(q^{4c}) \\ &= 16(\varphi(q^{60a})\psi(q^{8a}) + q^{14a}\varphi(q^{4a})\psi(q^{120a}) + q^{3a}\psi(q^{12a})\psi(q^{20a}) \\ &\quad + q^{5a}\varphi(q^{40a})\psi(q^{48a}) + q^{9a}\varphi(q^{24a})\psi(q^{80a}))\psi(q^{4b})\psi(q^{4c}). \end{aligned}$$

Thus,

$$\begin{aligned} \sum_{n=0}^{\infty} t(3a, 5a, 4b, 4c; 4n)q^{4n} &= 16\varphi(q^{60a})\psi(q^{8a})\psi(q^{4b})\psi(q^{4c}), \\ \sum_{n=0}^{\infty} t(3a, 5a, 4b, 4c; 4n + 14a)q^{4n+14a} &= 16q^{14a}\varphi(q^{4a})\psi(q^{120a})\psi(q^{4b})\psi(q^{4c}), \end{aligned}$$

which yields

$$(3.41) \quad \sum_{n=0}^{\infty} t(3a, 5a, 4b, 4c; 4n)q^n = 16\varphi(q^{15a})\psi(q^{2a})\psi(q^b)\psi(q^c),$$

$$(3.42) \quad \sum_{n=0}^{\infty} t(3a, 5a, 4b, 4c; 4n + 14a)q^n = 16\varphi(q^a)\psi(q^{30a})\psi(q^b)\psi(q^c).$$

From (3.42), (3.41) and (1.6) we see that

$$\begin{aligned} \sum_{n=0}^{\infty} t(3a, 5a, 8b, 8c; 4n + 14a)q^n &= 16(\varphi(q^{4a}) + 2q^a\psi(q^{8a}))\psi(q^{30a})\psi(q^{2b})\psi(q^{2c}), \\ \sum_{n=0}^{\infty} t(3a, 5a, 8b, 8c; 4n)q^n &= 16(\varphi(q^{60a}) + 2q^{15a}\psi(q^{120a}))\psi(q^{2a})\psi(q^{2b})\psi(q^{2c}). \end{aligned}$$

Thus,

$$\begin{aligned} \sum_{n=0}^{\infty} t(3a, 5a, 8b, 8c; 4(2n + a) + 14a)q^{2n+a} &= 32q^a\psi(q^{8a})\psi(q^{30a})\psi(q^{2b})\psi(q^{2c}), \\ \sum_{n=0}^{\infty} t(3a, 5a, 8b, 8c; 4(2n + 15a))q^{2n+15a} &= 32q^{15a}\psi(q^{120a})\psi(q^{2a})\psi(q^{2b})\psi(q^{2c}) \end{aligned}$$

and so

$$\begin{aligned} \sum_{n=0}^{\infty} t(3a, 5a, 8b, 8c; 4(2n + a) + 14a)q^n &= 32\psi(q^{4a})\psi(q^{15a})\psi(q^b)\psi(q^c), \\ \sum_{n=0}^{\infty} t(3a, 5a, 8b, 8c; 4(2n + 15a))q^n &= 32\psi(q^a)\psi(q^{60a})\psi(q^b)\psi(q^c), \end{aligned}$$

which yields (3.33) and (3.34). Note that  $\varphi(q)\psi(q^2) = \psi(q)^2$ . Taking  $c = 2a$  in (3.42) and  $c = 30a$  in (3.41) yields (3.35) and (3.36). Using (3.28) we see that

$$\sum_{n=0}^{\infty} t(a, 15a, 4b, 4c; n)q^n = 16\psi(q^a)\psi(q^{15a})\psi(q^{4b})\psi(q^{4c})$$

$$= 16(\varphi(q^{120a})\psi(q^{16a}) + q^{28a}\varphi(q^{8a})\psi(q^{240a}) + q^{6a}\psi(q^{4a})\psi(q^{60a}) \\ + q^a\varphi(q^{20a})\psi(q^{24a}) + q^{3a}\varphi(q^{12a})\psi(q^{40a}))\psi(q^{4b})\psi(q^{4c}).$$

From this it follows that

$$\sum_{n=0}^{\infty} t(a, 15a, 4b, 4c; 4n + a)q^{4n+a} = 16q^a\varphi(q^{20a})\psi(q^{24a})\psi(q^{4b})\psi(q^{4c}), \\ \sum_{n=0}^{\infty} t(a, 15a, 4b, 4c; 4n + 3a)q^{4n+3a} = 16q^{3a}\varphi(q^{12a})\psi(q^{40a})\psi(q^{4b})\psi(q^{4c})$$

and so

$$(3.43) \quad \sum_{n=0}^{\infty} t(a, 15a, 4b, 4c; 4n + a)q^n = 16\varphi(q^{5a})\psi(q^{6a})\psi(q^b)\psi(q^c),$$

$$(3.44) \quad \sum_{n=0}^{\infty} t(a, 15a, 4b, 4c; 4n + 3a)q^n = 16\varphi(q^{3a})\psi(q^{10a})\psi(q^b)\psi(q^c).$$

Hence, applying (1.6) we get

$$\sum_{n=0}^{\infty} t(a, 15a, 8b, 8c; 4n + a)q^n = 16(\varphi(q^{20a}) + 2q^{5a}\psi(q^{40a}))\psi(q^{6a})\psi(q^{2b})\psi(q^{2c}), \\ \sum_{n=0}^{\infty} t(a, 15a, 8b, 8c; 4n + 3a)q^n = 16(\varphi(q^{12a}) + 2q^{3a}\psi(q^{24a}))\psi(q^{10a})\psi(q^{2b})\psi(q^{2c})$$

and so

$$\sum_{n=0}^{\infty} t(a, 15a, 8b, 8c; 4(2n + 5a) + a)q^{2n+5a} = 32q^{5a}\psi(q^{40a})\psi(q^{6a})\psi(q^{2b})\psi(q^{2c}), \\ \sum_{n=0}^{\infty} t(a, 15a, 8b, 8c; 4(2n + 3a) + 3a)q^{2n+3a} = 32q^{3a}\psi(q^{24a})\psi(q^{10a})\psi(q^{2b})\psi(q^{2c}).$$

It then follows that

$$\sum_{n=0}^{\infty} t(a, 15a, 8b, 8c; 4(2n + 5a) + a)q^n = 32\psi(q^{20a})\psi(q^{3a})\psi(q^b)\psi(q^c), \\ \sum_{n=0}^{\infty} t(a, 15a, 8b, 8c; 4(2n + 3a) + 3a)q^n = 32\psi(q^{12a})\psi(q^{5a})\psi(q^b)\psi(q^c),$$

which yields (3.37) and (3.38). Finally, taking  $c = 6a$  in (3.44) and  $c = 10a$  in (3.43) and then applying (1.5) we deduce (3.39) and (3.40). The proof is now complete.

## 4. Evaluation of $t(2, 3, 3, 8; n)$ , $t(1, 1, 6, 24; n)$ and $t(1, 1, 6, 8; n)$

In this section we determine  $t(2, 3, 3, 8; n)$ ,  $t(1, 1, 6, 24; n)$  and  $t(1, 1, 6, 8; n)$  for any positive integer  $n$ .

**Theorem 4.1.** *Let  $n$  be a positive integer.*

(i) *If  $2n + 5 = 3^\beta n_1$  ( $3 \nmid n_1$ ), then*

$$t(1, 1, 6, 24; 2n + 1) = t(2, 3, 3, 8; 2n + 3) = 4(\sigma(n_1) - (-1)^n a(2n + 5)),$$

where  $\{a(n)\}$  is given by

$$q \prod_{k=0}^{\infty} (1 - q^{2k})(1 - q^{4k})(1 - q^{6k})(1 - q^{12k}) = \sum_{n=1}^{\infty} a(n)q^n \quad (|q| < 1).$$

(ii) *If  $n + 1 = 2^\alpha 3^\beta n_1$  with  $2 \nmid n_1$  and  $3 \nmid n_1$ , then*

$$t(1, 1, 6, 24; 2n - 2) + t(2, 3, 3, 8; 2n) = 2^{\alpha+4} \sigma(n_1).$$

Proof. By (3.1),  $t(1, 1, 6, 24; 2n + 1) = 2t(1, 3, 4, 12; n) = t(2, 3, 3, 8; 2n + 3)$ . By [WS2, Theorem 3.4], if  $2n+5 = 3^\beta n_1$  with  $3 \nmid n_1$ , then  $t(1, 3, 4, 12; n) = 2(\sigma(n_1) - (-1)^n a(2n+5))$ . Thus (i) is true.

Now we prove (ii). Since  $\psi(q)^2 = \varphi(q)\psi(q^2) = (\varphi(q^4) + 2q\psi(q^8))\psi(q^2)$  we see that

$$\sum_{n=0}^{\infty} t(1, 1, 6, 24; n)q^n = 16\psi(q)^2\psi(q^6)\psi(q^{24}) = (16\varphi(q^4) + 32q\psi(q^8))\psi(q^2)\psi(q^6)\psi(q^{24}).$$

Thus,

$$\sum_{n=0}^{\infty} t(1, 1, 6, 24; 2n)q^{2n} = 16\varphi(q^4)\psi(q^2)\psi(q^6)\psi(q^{24})$$

and so

$$(4.1) \quad \sum_{n=0}^{\infty} t(1, 1, 6, 24; 2n)q^n = 16\varphi(q^2)\psi(q)\psi(q^3)\psi(q^{12}).$$

Similarly,

$$\sum_{n=0}^{\infty} t(2, 3, 3, 8; n)q^n = 16\psi(q^2)\psi(q^3)^2\psi(q^8) = (16\varphi(q^{12}) + 32q^3\psi(q^{24}))\psi(q^6)\psi(q^2)\psi(q^8).$$

Thus,

$$\sum_{n=0}^{\infty} t(2, 3, 3, 8; 2n)q^{2n} = 16\psi(q^2)\psi(q^6)\psi(q^8)\varphi(q^{12})$$

and so

$$(4.2) \quad \sum_{n=0}^{\infty} t(2, 3, 3, 8; 2n)q^n = 16\psi(q)\psi(q^3)\psi(q^4)\varphi(q^6).$$

From (4.1) and (4.2) we see that

$$\sum_{n=0}^{\infty} (t(2, 3, 3, 8; 2n) + t(1, 1, 6, 24; 2n - 2))q^n$$



$$= 16\psi(q)\psi(q^3)(\varphi(q^6)\psi(q^4) + q\varphi(q^2)\psi(q^{12})) = 16\psi(q)^2\psi(q^3)^2.$$

Hence

$$(4.3) \quad t(2, 3, 3, 8; 2n) + t(1, 1, 6, 24; 2n - 2) = t(1, 1, 3, 3; n).$$

By [WS1, Lemma 4.1], if  $n + 1 = 2^\alpha 3^\beta n_1$  with  $2 \nmid n_1$  and  $3 \nmid n_1$ , then

$$(4.4) \quad t(1, 1, 3, 3; n) = 2^{\alpha+4}\sigma(n_1).$$

Thus (ii) holds and the proof is complete.

**Theorem 4.2.** *Let  $n$  be a positive integer.*

(i) *If  $n + 1 = 2^\alpha 3^\beta n_1$  with  $2 \nmid n_1$  and  $3 \nmid n_1$ , then*

$$t(2, 3, 3, 8; 4n + 2) = t(1, 1, 6, 24; 4n) = 2^{\alpha+4}\sigma(n_1).$$

(ii) *If  $2n + 1 = 3^\beta n_1$  ( $3 \nmid n_1$ ), then*

$$\begin{aligned} t(2, 3, 3, 8; 4n) &= 8(\sigma(n_1) + a(2n + 1)), \\ t(1, 1, 6, 24; 4n - 2) &= 8(\sigma(n_1) - a(2n + 1)). \end{aligned}$$

Proof. If  $n + 1 = 2^\alpha 3^\beta n_1$  with  $2 \nmid n_1$  and  $3 \nmid n_1$ , from Theorem 4.1 we have

$$t(1, 1, 6, 24; 4n) + t(2, 3, 3, 8; 4n + 2) = 2^{\alpha+5}\sigma(n_1).$$

By [S1, Theorem 2.14],  $t(1, 1, 6, 24; 4n) = 2^{\alpha+4}\sigma(n_1)$ . Thus,  $t(2, 3, 3, 8; 4n+2) = 2^{\alpha+4}\sigma(n_1)$ . This proves (i).

Now we consider (ii). Suppose  $2n + 1 = 3^\beta n_1$  ( $3 \nmid n_1$ ). We first assume that  $n$  is odd and  $n = 2m + 1$ . By (3.13),  $t(2, 3, 3, 8; 8m + 12) = 4t(1, 1, 6, 6; m)$ . This together with [S3, Theorem 4.15] yields

$$t(2, 3, 3, 8; 8m + 4) = 4t(1, 1, 6, 6; m - 1) = 8(\sigma(n_1) + a(4m + 3)).$$

Now, appealing to Theorem 4.1(ii) we get

$$t(1, 1, 6, 24; 8m + 2) = 16\sigma(n_1) - t(2, 3, 3, 8; 8m + 4) = 8(\sigma(n_1) - a(4m + 3)).$$

From now on suppose that  $n$  is even and  $n = 2m$ . From (3.12) and [S3, Theorem 4.15] we see that

$$t(1, 1, 6, 24; 8m - 2) = 4t(2, 2, 3, 3; m - 1) = 8(\sigma(n_1) - a(4m + 1)).$$

Now applying Theorem 4.1(ii) gives

$$t(2, 3, 3, 8; 8m) = 16\sigma(n_1) - t(1, 1, 6, 24; 8m - 2) = 8(\sigma(n_1) + a(4m + 1)).$$

Putting all the above together proves the theorem.

**Theorem 4.3.** *Let  $n \in \mathbb{Z}^+$ .*

(i) If  $n + 1 = 2^\alpha 3^\beta n_1$  with  $2 \nmid n_1$  and  $3 \nmid n_1$ , then

$$t(1, 1, 6, 8; 2n) = 2^{\alpha+2} \left( 3^{\beta+1} \left( \frac{3}{n_1} \right) + (-1)^{\alpha+\beta+\frac{n_1-1}{2}} \right) \sum_{d|n_1} d \left( \frac{3}{d} \right).$$

(ii) If  $2n + 3 = 3^\beta n_1$  with  $3 \nmid n_1$ , then

$$t(1, 1, 6, 8; 2n + 1) = 2 \left( 3^{\beta+1} \left( \frac{3}{n_1} \right) + (-1)^n \right) \sum_{d|n_1} d \left( \frac{3}{d} \right) - 4 \sum_{\substack{a, b \in \mathbb{Z}^+, 2 \nmid a \\ 8n+12=a^2+3b^2}} (-1)^{(a-1)/2} a.$$

Proof. From [S1, Theorem 2.7],

$$t(1, 1, 6, 8; 2n) = t(1, 2, 2, 3; n) \quad \text{and} \quad t(1, 1, 6, 8; 2n + 1) = 2t(1, 3, 4, 4; n).$$

By [C, (5.6) and Theorem 5.4], if  $n + 1 = 2^\alpha 3^\beta n_1$  with  $2 \nmid n_1$  and  $3 \nmid n_1$ , then

$$\begin{aligned} t(1, 2, 2, 3; n) &= 2^{\alpha+2} \left( 3^{\beta+1} + (-1)^{\alpha+\beta} \left( \frac{-3}{n_1} \right) \right) \sum_{d|n_1} \frac{n_1}{d} \left( \frac{3}{d} \right) \\ &= 2^{\alpha+2} \left( 3^{\beta+1} + (-1)^{\alpha+\beta} \left( \frac{-3}{n_1} \right) \right) \sum_{d|n_1} d \left( \frac{3}{n_1/d} \right) \\ &= 2^{\alpha+2} \left( 3^{\beta+1} \left( \frac{3}{n_1} \right) + (-1)^{\alpha+\beta+(n_1-1)/2} \right) \sum_{d|n_1} d \left( \frac{3}{d} \right). \end{aligned}$$

Thus part(i) is true.

Let us consider part(ii). By [C, (5.6) and Theorem 5.6], if  $2n + 3 = 3^\beta n_1$  with  $3 \nmid n_1$ , then

$$t(1, 3, 4, 4; n) = \left( 3^{\beta+1} - (-1)^\beta \left( \frac{-3}{n_1} \right) \right) \sum_{d|n_1} \frac{n_1}{d} \left( \frac{3}{d} \right) - 2g_1(2n + 3),$$

where  $\{g_1(n)\}$  is given by

$$q\psi(q^6) \prod_{k=1}^{\infty} (1 - q^{2k})^3 = \sum_{n=1}^{\infty} g_1(n) q^n.$$

Using Jacobi's identity,

$$\begin{aligned} q\psi(q^6) \prod_{k=1}^{\infty} (1 - q^{2k})^3 &= q \left( \sum_{m=0}^{\infty} q^{3m(m+1)} \right) \left( \sum_{k=0}^{\infty} (-1)^k (2k + 1) q^{k(k+1)} \right) \\ &= \left( \sum_{m=0}^{\infty} q^{3(2m+1)^2/4} \right) \left( \sum_{k=0}^{\infty} (-1)^k (2k + 1) q^{(2k+1)^2/4} \right) \\ &= \sum_{n=0}^{\infty} \left( \sum_{\substack{k, m \in \{0, 1, 2, \dots\} \\ 4n = (2k+1)^2 + 3(2m+1)^2}} (-1)^k (2k + 1) \right) q^n \\ &= \sum_{n=0}^{\infty} \left( \sum_{\substack{a, b \in \mathbb{Z}^+, 2 \nmid a \\ 4n = a^2 + 3b^2}} (-1)^{(a-1)/2} a \right) q^n. \end{aligned}$$

Hence

$$g_1(n) = \sum_{\substack{a,b \in \mathbb{Z}^+, 2 \nmid a \\ 4n = a^2 + 3b^2}} (-1)^{(a-1)/2} a.$$

It then follows that

$$\begin{aligned} t(1, 1, 6, 8; 2n+1) &= 2t(1, 3, 4, 4; n) = 2 \left( 3^{\beta+1} - (-1)^\beta \left( \frac{-3}{n_1} \right) \right) \sum_{d|n_1} \frac{n_1}{d} \left( \frac{3}{d} \right) - 4g_1(2n+3) \\ &= 2 \left( 3^{\beta+1} \left( \frac{3}{n_1} \right) - (-1)^\beta \left( \frac{-1}{n_1} \right) \right) \sum_{d|n_1} d \left( \frac{3}{d} \right) - 4 \sum_{\substack{a,b \in \mathbb{Z}^+, 2 \nmid a \\ 8n+12 = a^2 + 3b^2}} (-1)^{(a-1)/2} a. \end{aligned}$$

Observe that  $(-1)^\beta \left( \frac{-1}{n_1} \right) \equiv 3^\beta n_1 = 2n+3 \equiv (-1)^{n-1} \pmod{4}$ . We then obtain part(ii). Now the proof is complete.

## 5. Special relations between $t(a, b, c, d; n)$ and $N(a, b, c, d; n)$

In this section, we present some special relations between  $t(a, b, c, d; n)$  and  $N(a, b, c, d; n)$  and pose a lot of related conjectures.

**Theorem 5.1.** *Let  $n \in \mathbb{Z}^+$  Then*

$$\begin{aligned} t(1, 4, 7, 8; n) &= 2N(1, 4, 7, 8; 2n+5) \quad \text{for } n \equiv 3 \pmod{4}, \\ t(1, 4, 8, 15; n) &= 2N(1, 4, 8, 15; 2n+7) \quad \text{for } n \equiv 2 \pmod{4}, \\ t(3, 5, 12, 24; n) &= 2N(3, 5, 12, 24; 2n+11) \quad \text{for } n \equiv 3 \pmod{4}, \\ t(3, 5, 20, 40; n) &= 2N(3, 5, 20, 40; 2n+17) \quad \text{for } n \equiv 3 \pmod{4}. \end{aligned}$$

*Proof.* From (3.15), (3.31) and (3.32) we see that

$$\begin{aligned} t(1, 4, 7, 8; 4n+3) &= t(1, 2, 7, 4; 2n+1) = t(1, 1, 2, 7; n), \\ t(1, 4, 8, 15; 4n+6) &= t(1, 1, 2, 15; n), \\ t(3, 5, 12, 24; 4n+3) &= t(3, 3, 5, 6; n), \\ t(3, 5, 20, 40; 4n+3) &= t(3, 5, 5, 10; n). \end{aligned}$$

Now applying Theorem 2.2 yields the result.

**Theorem 5.2.** *For  $n \in \mathbb{Z}^+$  we have*

$$\begin{aligned} t(2, 3, 3, 4; n) &= 2N(2, 3, 3, 4; 2n+3) \quad \text{for } n \equiv 2, 3 \pmod{4}, \\ t(2, 3, 3, 12; n) &= 2N(2, 3, 3, 12; 2n+5) \quad \text{for } n \equiv 0, 1 \pmod{4}, \\ t(2, 3, 3, 24; n) &= 4N(2, 3, 3, 24; 2n+8) \quad \text{for } n \equiv 2 \pmod{4}, \\ t(2, 3, 3, 36; n) &= 2N(2, 3, 3, 36; 2n+11) \quad \text{for } n \equiv 2, 3 \pmod{4}, \\ t(1, 1, 6, 12; n) &= 2N(1, 1, 6, 12; 2n+5) \quad \text{for } n \equiv 0, 3 \pmod{4}, \\ t(1, 1, 6, 16; n) &= \begin{cases} N(1, 1, 3, 8; n+3) & \text{if } n \equiv 2 \pmod{8}, \\ 4N(1, 1, 3, 8; n+3) & \text{if } n \equiv 4 \pmod{8}. \end{cases} \end{aligned}$$

Proof. By Theorem 2.3 and (3.8),

$$N(2, 3, 3, 4; 8n + 9) = t(1, 2, 3, 3; n) = \frac{1}{2}t(2, 3, 3, 4; 4n + 3).$$

By (1.6) and (1.9),

$$\begin{aligned} \sum_{n=0}^{\infty} N(2, 3, 3, 4; n)q^n &= \varphi(q^2)\varphi(q^4)\varphi(q^3)^2 \\ &= (\varphi(q^8) + 2q^2\psi(q^{16}))(\varphi(q^{16}) + 2q^4\psi(q^{32})) \\ &\quad \times (\varphi(q^{24})^2 + 4q^{12}\psi(q^{48})^2 + 4q^6\psi(q^{24})^2 + 4q^3\varphi(q^{48})\psi(q^{24}) + 8q^{15}\psi(q^{24})\psi(q^{96})). \end{aligned}$$

Thus

$$\begin{aligned} \sum_{n=0}^{\infty} N(2, 3, 3, 4; 8n + 7)q^{8n+7} \\ &= \varphi(q^8)\varphi(q^{16}) \cdot 8q^{15}\psi(q^{24})\psi(q^{96}) + 2q^4\varphi(q^8)\psi(q^{32}) \cdot 4q^3\varphi(q^{48})\psi(q^{24}) \\ &= 8q^7\varphi(q^8)\psi(q^{24})(\varphi(q^{48})\psi(q^{32}) + q^8\varphi(q^{16})\psi(q^{96})) \\ &= 8q^7\varphi(q^8)\psi(q^{24})\psi(q^8)\psi(q^{24}) \end{aligned}$$

and so

$$(5.1) \quad \sum_{n=0}^{\infty} N(2, 3, 3, 4; 8n + 7)q^n = 8\varphi(q)\psi(q)\psi(q^3)^2.$$

On the other hand, using (1.5), (1.6) and (1.8) we see that for  $b \in \mathbb{Z}^+$ ,

$$\begin{aligned} (5.2) \quad \sum_{n=0}^{\infty} t(2, 3, 3, 4b; n)q^n \\ &= 16\psi(q^2)\psi(q^3)^2\psi(q^{4b}) = 16\varphi(q^3)\psi(q^2)\psi(q^6)\psi(q^{4b}) \\ &= 16(\varphi(q^{12}) + 2q^3\psi(q^{24}))(\varphi(q^{12})\psi(q^8) + q^2\varphi(q^4)\psi(q^{24}))\psi(q^{4b}). \end{aligned}$$

Thus

$$\sum_{n=0}^{\infty} t(2, 3, 3, 4b; 4n + 2)q^{4n+2} = 16q^2\varphi(q^4)\psi(q^{4b})\varphi(q^{12})\psi(q^{24})$$

and so

$$\sum_{n=0}^{\infty} t(2, 3, 3, 4b; 4n + 2)q^n = 16\varphi(q)\psi(q^b)\varphi(q^3)\psi(q^6) = 16\varphi(q)\psi(q^3)^2\psi(q^b).$$

This together with (5.1) yields  $t(2, 3, 3, 4; 4n + 2) = 2N(2, 3, 3, 4; 8n + 7)$ . Therefore  $t(2, 3, 3, 4; n) = 2N(2, 3, 3, 4; 2n + 3)$  for  $n \equiv 2, 3 \pmod{4}$ . The remaining results can be proved similarly.

**Lemma 5.1.** *Suppose  $a, b \in \mathbb{Z}^+$  and  $ab \equiv -1 \pmod{4}$ . Then*

$$\sum_{n=0}^{\infty} N(a, a, a, 2b; 8n + 5a)q^n = 24\varphi(q^b)\psi(q^a)\psi(q^{2a})^2$$

$$\sum_{n=0}^{\infty} N(a, a, a, 2b; 8n + a + 2b)q^n = 12\varphi(q^a)^2\psi(q^a)\psi(q^{2b}).$$

Proof. It is evident that

$$\begin{aligned} & \sum_{n=0}^{\infty} N(a, a, a, 2b; n)q^n \\ &= \varphi(q^a)^3\varphi(q^{2b}) = (\varphi(q^{16a}) + 2q^{4a}\psi(q^{32a}) + 2q^a\psi(q^{8a}))^3(\varphi(q^{8b}) + 2q^{2b}\psi(q^{16b})) \\ &= ((\varphi(q^{16a}) + 2q^{4a}\psi(q^{32a}))^3 + 6q^a(\varphi(q^{16a}) + 2q^{4a}\psi(q^{32a}))^2\psi(q^{8a}) + 12q^{2a}(\varphi(q^{16a}) \\ &\quad + 2q^{4a}\psi(q^{32a}))\psi(q^{8a})^2 + 8q^{3a}\psi(q^{8a})^3)(\varphi(q^{8b}) + 2q^{2b}\psi(q^{16b})). \end{aligned}$$

Thus,

$$\begin{aligned} & \sum_{n=0}^{\infty} N(a, a, a, 2b; 8n + 5a)q^{8n+5a} = 6q^a\varphi(q^{16a}) \cdot 4q^{4a}\psi(q^{32a})\psi(q^{8a})\varphi(q^{8b}), \\ & \sum_{n=0}^{\infty} N(a, a, a, 2b; 8n + a + 2b)q^{8n+a+2b} \\ &= 3(\varphi(q^{16a})^2 + 4q^{8a}\psi(q^{32a})^2) \cdot 2q^a\psi(q^{8a}) \cdot 2q^{2b}\psi(q^{16b}) = 12q^{a+2b}\varphi(q^{8a})^2\psi(q^{8a})\psi(q^{16b}) \end{aligned}$$

and so the result follows.

**Theorem 5.3.** For  $n \in \mathbb{Z}^+$  with  $n \equiv 3, 5 \pmod{8}$ ,

$$\begin{aligned} t(1, 1, 2, 12; n) &= 4N(1, 1, 4, 6; n + 2) = \frac{8}{3}N(1, 1, 1, 6; n + 2) \\ t(3, 3, 4, 6; n) &= \frac{8}{3}N(2, 3, 3, 3; n + 2). \end{aligned}$$

Proof. By (3.1) and [S1, Theorem 2.11],

$$\begin{aligned} t(1, 1, 2, 12; 8n + 3) &= 2t(1, 1, 4, 6; 4n + 1) = 4t(1, 2, 3, 4; 2n) = 4N(1, 1, 4, 6; 8n + 5), \\ t(1, 1, 2, 12; 8n + 5) &= 2t(1, 1, 4, 6; 4n + 2) = 4N(1, 1, 4, 6; 8n + 7). \end{aligned}$$

From [S1, p.283] and Lemma 5.1 we know that

$$\begin{aligned} \sum_{n=0}^{\infty} t(1, 1, 4, 6; 4n + 1)q^n &= 32\varphi(q^3)\psi(q)\psi(q^2)^2 = \frac{4}{3} \sum_{n=0}^{\infty} N(1, 1, 1, 6; 8n + 5)q^n, \\ \sum_{n=0}^{\infty} t(1, 1, 4, 6; 4n + 2)q^n &= 16\varphi(q)^2\psi(q)\psi(q^6) = \frac{4}{3} \sum_{n=0}^{\infty} N(1, 1, 1, 6; 8n + 7)q^n. \end{aligned}$$

Hence the formula for  $t(1, 1, 2, 12; n)$  is true. From (5.2) we see that

$$\begin{aligned} \sum_{n=0}^{\infty} t(2, 3, 3, 12; 4n)q^{4n} &= 16\varphi(q^{12})^2\psi(q^8)\psi(q^{12}), \\ \sum_{n=0}^{\infty} t(2, 3, 3, 12; 4n + 5)q^{4n+5} &= 32q^5\psi(q^{24})\varphi(q^4)\psi(q^{24})\psi(q^{12}). \end{aligned}$$

Thus, appealing to Lemma 5.1 we get

$$\begin{aligned}\sum_{n=0}^{\infty} t(2, 3, 3, 12; 4n)q^n &= 16\varphi(q^3)^2\psi(q^2)\psi(q^3) = \frac{4}{3}\sum_{n=0}^{\infty} N(2, 3, 3, 3; 8n + 5)q^n, \\ \sum_{n=0}^{\infty} t(2, 3, 3, 12; 4n + 5)q^n &= 32\varphi(q)\psi(q^3)\psi(q^6)^2 = \frac{4}{3}\sum_{n=0}^{\infty} N(2, 3, 3, 3; 8n + 15)q^n.\end{aligned}$$

This together with (3.1) gives

$$\begin{aligned}t(3, 3, 4, 6; 8n + 3) &= 2t(2, 3, 3, 12; 4n) = \frac{8}{3}\sum_{n=0}^{\infty} N(2, 3, 3, 3; 8n + 5), \\ t(3, 3, 4, 6; 8n + 5) &= 2t(2, 3, 3, 12; 4n + 1) = 4t(1, 3, 6, 12; 2n - 1) \\ &= \frac{8}{3}\sum_{n=0}^{\infty} N(2, 3, 3, 3; 8n + 15),\end{aligned}$$

which completes the proof.

**Corollary 5.1.** *For  $n \in \mathbb{Z}^+$  with  $n \equiv 5, 7 \pmod{8}$ ,*

$$N(1, 1, 1, 6; 4n) = 5N(1, 1, 1, 6; n) \quad \text{and} \quad N(2, 3, 3, 3; 4n) = 5N(2, 3, 3, 3; n).$$

*Proof.* From Theorem 5.3 and [S1, Theorem 2.1],

$$\begin{aligned}\frac{8}{3}N(1, 1, 1, 6; n) &= t(1, 1, 2, 12; n - 2) = \frac{2}{3}(N(1, 1, 1, 6; 4n) - N(1, 1, 1, 6; n)), \\ \frac{8}{3}N(2, 3, 3, 3; n) &= t(3, 3, 4, 6; n - 2) = \frac{2}{3}(N(2, 3, 3, 3; 4n) - N(2, 3, 3, 3; n)).\end{aligned}$$

This yields the result.

Using similar method one can prove the following results:

$$(5.3) \quad t(3, 3, 4, 18; n) = 2N(3, 3, 4, 18; 2n + 7) \quad \text{for } n \equiv 0, 1 \pmod{4},$$

$$(5.4) \quad t(1, 3, 8, 12; n) = 4N(1, 3, 8, 12; n + 3) \quad \text{for } n \equiv 2, 4 \pmod{8},$$

$$(5.5) \quad t(1, 1, 2, 28; n) = 4N(1, 1, 2, 28; n + 4) \quad \text{for } n \equiv 1, 3 \pmod{8},$$

$$(5.6) \quad t(1, 3, 4, 24; n) = 4N(1, 3, 4, 24; n + 4) \quad \text{for } n \equiv 1, 3 \pmod{8},$$

$$(5.7) \quad t(2, 3, 3, 48; n) = N(2, 3, 3, 48; 2n + 14) \quad \text{for } n \equiv 0 \pmod{8},$$

$$(5.8) \quad t(1, 1, 8, 14; n) = 8N(1, 1, 8, 14; n + 3) \quad \text{for } n \equiv 1 \pmod{8},$$

$$(5.9) \quad t(1, 1, 10, 20; n) = 4N(1, 1, 10, 20; n + 4) \quad \text{for } n \equiv 1 \pmod{8},$$

$$(5.10) \quad t(1, 1, 14, 16; n) = 4N(1, 1, 14, 16; n + 4) \quad \text{for } n \equiv 1 \pmod{8},$$

$$(5.11) \quad t(1, 2, 7, 14; n) = 8N(1, 2, 7, 14; n + 3) \quad \text{for } n \equiv 1 \pmod{8},$$

$$(5.12) \quad t(1, 1, 8, 30; n) = 4N(1, 1, 8, 30; 2n + 10) \quad \text{for } n \equiv 1 \pmod{8},$$

$$(5.13) \quad t(1, 3, 4, 16; n) = \frac{4}{3}N(1, 3, 4, 16; 2n + 6) \quad \text{for } n \equiv 1 \pmod{8},$$

$$(5.14) \quad t(3, 3, 10, 48; n) = 4N(3, 3, 10, 48; 2n + 16) \quad \text{for } n \equiv 1 \pmod{8},$$

$$(5.15) \quad t(1, 1, 8, 14; n) = 4N(1, 1, 8, 14; 2n + 6) \quad \text{for } n \equiv 3 \pmod{8},$$

- (5.16)  $t(2, 15, 15, 24; n) = 4N(2, 15, 15, 24; 2n + 14)$  for  $n \equiv 3 \pmod{8}$ ,
- (5.17)  $t(5, 5, 6, 8; n) = 4N(5, 5, 6, 8; 2n + 6)$  for  $n \equiv 3 \pmod{8}$ ,
- (5.18)  $t(1, 3, 12, 48; n) = \frac{4}{3}N(1, 3, 12, 48; 2n + 16)$  for  $n \equiv 4 \pmod{8}$ ,
- (5.19)  $t(2, 4, 5, 5; n) = 4N(2, 4, 5, 5; n + 2)$  for  $n \equiv 5 \pmod{8}$ ,
- (5.20)  $t(4, 7, 7, 14; n) = 4N(4, 7, 7, 14; n + 4)$  for  $n \equiv 5 \pmod{8}$ ,
- (5.21)  $t(1, 1, 16, 30; n) = 4N(1, 1, 16, 30; 2n + 12)$  for  $n \equiv 5 \pmod{8}$ ,
- (5.22)  $t(1, 1, 30, 40; n) = 4N(1, 1, 30, 40; 2n + 18)$  for  $n \equiv 5 \pmod{8}$ ,
- (5.23)  $t(1, 3, 16, 36; n) = \frac{4}{3}N(1, 3, 16, 36; 2n + 14)$  for  $n \equiv 5 \pmod{8}$ ,
- (5.24)  $t(2, 3, 3, 32; n) = 4N(2, 3, 3, 32; 2n + 10)$  for  $n \equiv 5 \pmod{8}$ ,
- (5.25)  $t(2, 7, 7, 24; n) = 4N(2, 7, 7, 24; 2n + 10)$  for  $n \equiv 5 \pmod{8}$ ,
- (5.26)  $t(3, 3, 10, 24; n) = 4N(3, 3, 10, 24; 2n + 10)$  for  $n \equiv 5 \pmod{8}$ ,
- (5.27)  $t(1, 7, 16, 16; n) = 4N(1, 7, 16, 16; n + 5)$  for  $n \equiv 6 \pmod{8}$ ,
- (5.28)  $t(2, 3, 3, 48; n) = 4N(2, 3, 3, 48; 2n + 14)$  for  $n \equiv 6 \pmod{8}$ ,
- (5.29)  $t(1, 1, 10, 20; n) = 4N(1, 1, 10, 20; n + 4)$  for  $n \equiv 7 \pmod{8}$ ,
- (5.30)  $t(2, 4, 5, 5; n) = 4N(2, 4, 5, 5; n + 2)$  for  $n \equiv 7 \pmod{8}$ ,
- (5.31)  $t(4, 7, 7, 14; n) = 4N(4, 7, 7, 14; n + 4)$  for  $n \equiv 7 \pmod{8}$ ,
- (5.32)  $t(1, 1, 14, 16; n) = 4N(1, 1, 14, 16; 2n + 8)$  for  $n \equiv 7 \pmod{8}$ ,
- (5.33)  $t(5, 5, 6, 40; n) = 4N(5, 5, 6, 40; 2n + 14)$  for  $n \equiv 7 \pmod{8}$ .

By doing calculations on Maple, we pose the following conjectures.

**Conjecture 5.1.** *Suppose  $n \in \mathbb{Z}^+$ . Then*

$$\begin{aligned} t(1, 3, 5, 15; n) &= 8N(1, 3, 5, 15; n + 3) \quad \text{for } n \equiv 3 \pmod{4}, \\ t(1, 3, 7, 21; n) &= 8N(1, 3, 7, 21; n + 4) \quad \text{for } n \equiv 2 \pmod{4}, \\ t(1, 3, 9, 27; n) &= \frac{16}{13}N(1, 3, 9, 27; 2n + 10) \quad \text{for } n \equiv 7 \pmod{8}. \end{aligned}$$

We remark that Dongxi Ye informed the author he proved the formula for  $t(1, 3, 5, 15; n)$  and  $t(1, 3, 7, 21; n)$  by using theta function identities and weight 2 modular forms.

**Conjecture 5.2.** *Let  $n \in \mathbb{Z}^+$ . Then*

$$t(1, 2, 3, 10; n) = \begin{cases} \frac{4}{3}N(1, 2, 3, 10; 2n + 4) = \frac{4}{9}N(1, 2, 3, 10; 8n + 16) & \text{if } 4 \mid n - 1, \\ \frac{8}{3}N(1, 2, 3, 10; 2n + 4) & \text{if } 8 \mid n, \\ \frac{16}{9}N(1, 2, 3, 10; 2n + 4) & \text{if } n \equiv 10 \pmod{16}, \\ 4N(1, 2, 3, 10; 2n + 4) & \text{if } n \equiv 11, 15 \pmod{20}. \end{cases}$$

**Conjecture 5.3.** *Let  $n \in \mathbb{Z}^+$ . Then*

$$t(1, 2, 3, 18; n) = \begin{cases} \frac{4}{3}N(1, 2, 3, 18; 2n + 6) = \frac{4}{9}N(1, 2, 3, 18; 8n + 24) & \text{if } 4 \mid n, \\ \frac{8}{3}N(1, 2, 3, 18; 2n + 6) & \text{if } 8 \mid n - 3, \\ 4N(1, 2, 3, 18; 2n + 6) & \text{if } 12 \mid n - 6, \\ \frac{8}{5}N(1, 2, 3, 18; 2n + 6) & \text{if } 24 \mid n - 15, \end{cases}$$

**Conjecture 5.4.** *Let  $n \in \mathbb{Z}^+$ . Then*

$$t(1, 3, 6, 30; n) = \begin{cases} \frac{4}{3}N(1, 3, 6, 30; 2n + 10) = \frac{4}{9}N(1, 3, 6, 30; 8n + 40) & \text{if } 4 \mid n, \\ \frac{8}{3}N(1, 3, 6, 30; 2n + 10) = \frac{8}{15}N(1, 3, 6, 30; 8n + 40) & \text{if } 8 \mid n - 1, \\ \frac{16}{9}N(1, 3, 6, 30; 2n + 10) = \frac{16}{33}N(1, 3, 6, 30; 8n + 40) & \text{if } 16 \mid n + 1. \end{cases}$$

**Conjecture 5.5.** *Let  $n \in \mathbb{Z}^+$ . Then*

$$t(1, 3, 18, 18; n) = \begin{cases} \frac{4}{3}N(1, 3, 18, 18; 2n + 10) = \frac{4}{9}N(1, 3, 18, 18; 8n + 40) & \text{if } 4 \mid n - 2, \\ \frac{8}{3}N(1, 3, 18, 18; 2n + 10) = \frac{8}{15}N(1, 3, 18, 18; 8n + 40) & \text{if } 8 \mid n - 1, \\ \frac{16}{9}N(1, 3, 18, 18; 2n + 10) = \frac{16}{33}N(1, 3, 18, 18; 8n + 40) & \text{if } 16 \mid n - 7, \\ 4N(1, 3, 18, 18; 2n + 10) = \frac{4}{7}N(1, 3, 18, 18; 8n + 40) & \text{if } 12 \mid n - 4, \\ \frac{8}{5}N(1, 3, 18, 18; 2n + 10) = \frac{8}{17}N(1, 3, 18, 18; 8n + 40) & \text{if } 24 \mid n - 13. \end{cases}$$

**Conjecture 5.6.** *Let  $n \in \mathbb{Z}^+$ . Then*

$$t(2, 3, 9, 18; n) = \begin{cases} \frac{4}{3}N(2, 3, 9, 18; 2n + 8) = \frac{4}{9}N(2, 3, 9, 18; 8n + 32) & \text{if } 4 \mid n - 3, \\ \frac{8}{3}N(2, 3, 9, 18; 2n + 8) & \text{if } 8 \mid n - 2, \\ \frac{16}{9}N(2, 3, 9, 18; 2n + 8) & \text{if } 16 \mid n - 8, \\ \frac{32}{15}N(2, 3, 9, 18; 2n + 8) & \text{if } 32 \mid n - 20, \\ \frac{8}{5}N(2, 3, 9, 18; 2n + 8) & \text{if } 24 \mid n - 14, \\ 4N(2, 3, 9, 18; 2n + 8) & \text{if } 12 \mid n - 5. \end{cases}$$

**Conjecture 5.7.** *Let  $n \in \mathbb{Z}^+$ . Then*

$$t(2, 5, 10, 15; n)$$



$$= \begin{cases} \frac{4}{3}N(2, 5, 10, 15; 2n + 8) = \frac{4}{9}N(2, 5, 10, 15; 8n + 32) & \text{if } 4 \mid n - 3, \\ \frac{8}{3}N(2, 5, 10, 15; 2n + 8) & \text{if } 8 \mid n - 6, \\ \frac{16}{9}N(2, 5, 10, 15; 2n + 8) & \text{if } 16 \mid n - 8, \\ 4N(2, 5, 10, 15; 2n + 8) & \text{if } n \equiv 61, 81 \pmod{100} \end{cases}$$

**Conjecture 5.8.** *Let  $n \in \mathbb{Z}^+$ . Then*

$$t(5, 6, 15, 30; n) = \begin{cases} \frac{4}{3}N(5, 6, 15, 30; 2n + 14) & \text{if } 4 \mid n - 2, \\ \frac{8}{3}N(5, 6, 15, 30; 2n + 14) & \text{if } 8 \mid n - 7, \\ \frac{16}{9}N(5, 6, 15, 30; 2n + 14) & \text{if } 16 \mid n - 13. \end{cases}$$

**Conjecture 5.9.** *Let  $n \in \mathbb{Z}^+$ . Then*

$$t(1, 6, 9, 12; n) = 2N(1, 6, 9, 12; 2n + 7) \quad \text{for } n \equiv 2, 3 \pmod{4}.$$

**Conjecture 5.10.** *Let  $n \in \mathbb{Z}^+$  with  $n \equiv 0 \pmod{8}$ . Then*

$$\begin{aligned} t(1, 3, 24, 28; n) &= 4N(1, 3, 24, 28; n + 7), \\ t(1, 6, 10, 15; n) &= 8N(1, 6, 10, 15; n + 4), \\ t(1, 2, 21, 24; n) &= 4N(1, 2, 21, 24; 2n + 12), \\ t(1, 6, 8, 33; n) &= 4N(1, 6, 8, 33; 2n + 12). \end{aligned}$$

**Conjecture 5.11.** *Let  $n \in \mathbb{Z}^+$  with  $n \equiv 1 \pmod{8}$ . Then*

$$\begin{aligned} t(1, 2, 5, 32; n) &= 4N(1, 2, 5, 32; 2n + 10), \\ t(1, 2, 13, 24; n) &= 4N(1, 2, 13, 24; 2n + 10), \\ t(2, 5, 9, 24; n) &= 4N(2, 5, 9, 24; 2n + 10). \end{aligned}$$

**Conjecture 5.12.** *Let  $n \in \mathbb{Z}^+$  with  $n \equiv 2 \pmod{8}$ . Then*

$$\begin{aligned} t(1, 2, 5, 8; n) &= \frac{4}{3}N(1, 2, 5, 8; 2n + 4), \\ t(1, 2, 5, 24; n) &= 4N(1, 2, 5, 24; 2n + 8). \end{aligned}$$

**Conjecture 5.13.** *Let  $n \in \mathbb{Z}^+$  with  $n \equiv 3 \pmod{8}$ . Then*

$$\begin{aligned} t(1, 6, 8, 9; n) &= 4N(1, 6, 8, 9; 2n + 6), \\ t(1, 18, 24, 45; n) &= 4N(1, 18, 24, 45; 2n + 22), \\ t(2, 3, 24, 27; n) &= 4N(2, 3, 24, 27; 2n + 14), \\ t(2, 5, 10, 15; n) &= \frac{4}{3}N(2, 5, 10, 15; 2n + 8), \\ t(3, 6, 8, 39; n) &= 4N(3, 6, 8, 39; 2n + 14). \end{aligned}$$

**Conjecture 5.14.** *Let  $n \in \mathbb{Z}^+$  with  $n \equiv 4 \pmod{8}$ . Then*

$$t(1, 8, 16, 31; n) = 4N(1, 8, 16, 31; n + 7),$$

$$t(1, 9, 30, 40; n) = 4N(1, 9, 30, 40; 2n + 20),$$

$$t(2, 2, 3, 9; n) = \frac{8}{3}N(2, 2, 3, 9; 2n + 4).$$

**Conjecture 5.15.** *Let  $n \in \mathbb{Z}^+$  with  $n \equiv 5 \pmod{8}$ . Then*

$$t(1, 8, 16, 23; n) = 4N(1, 8, 16, 23; n + 6),$$

$$t(1, 5, 10, 40; n) = \frac{4}{3}N(1, 5, 10, 40; 2n + 14),$$

$$t(2, 3, 11, 24; n) = 4N(2, 3, 11, 24; 2n + 10).$$

**Conjecture 5.16.** *Let  $n \in \mathbb{Z}^+$  with  $n \equiv 6 \pmod{8}$ . Then*

$$t(1, 18, 21, 24; n) = 4N(1, 18, 21, 24; 2n + 16),$$

$$t(3, 6, 8, 15; n) = 4N(3, 6, 8, 15; 2n + 8),$$

$$t(5, 6, 8, 45; n) = 4N(5, 6, 8, 45; 2n + 16).$$

**Conjecture 5.17.** *Let  $n \in \mathbb{Z}^+$  with  $n \equiv 7 \pmod{8}$ . Then*

$$t(2, 3, 5, 30; n) = 8N(2, 3, 5, 30; n + 5),$$

$$t(1, 6, 9, 24; n) = \frac{4}{3}N(1, 6, 9, 24; 2n + 10),$$

$$t(2, 5, 24, 25; n) = 4N(2, 5, 24, 25; 2n + 14),$$

$$t(2, 9, 21, 24; n) = 4N(2, 9, 21, 24; 2n + 14),$$

$$t(3, 6, 7, 8; n) = 4N(3, 6, 7, 8; 2n + 6),$$

$$t(3, 15, 30, 40; n) = 4N(3, 15, 30, 40; 2n + 22).$$

**Conjecture 5.18.** *Let  $n \in \mathbb{Z}^+$ . Then*

$$t(1, 1, 25, 25; n) = 2N(1, 1, 25, 25; 2n + 13) \quad \text{for } n \equiv 0, 2 \pmod{5},$$

$$t(1, 1, 5, 21; n) = 2N(1, 1, 5, 21; 2n + 7) \quad \text{for } n \equiv 0, 3 \pmod{5},$$

$$t(1, 6, 6, 15; n) = 2N(1, 6, 6, 15; 2n + 7) \quad \text{for } n \equiv 0, 3 \pmod{5},$$

$$t(2, 2, 3, 5; n) = 2N(2, 2, 3, 5; 2n + 3) \quad \text{for } n \equiv 3, 4 \pmod{5},$$

$$t(3, 7, 7, 35; n) = 2N(3, 7, 7, 35; 2n + 13) \quad \text{for } n \equiv 3, 4 \pmod{5},$$

$$t(3, 3, 7, 15; n) = 2N(3, 3, 7, 15; 2n + 7) \quad \text{for } n \equiv 1, 2 \pmod{5}.$$

$$t(1, 1, 3, 5; n) = \frac{2}{5}N(1, 1, 3, 5; 8n + 10) \quad \text{for } n \equiv 2, 3 \pmod{5},$$

$$t(1, 3, 3, 15; n) = \frac{2}{5}N(1, 3, 3, 15; 8n + 22) \quad \text{for } n \equiv 0, 2 \pmod{5}.$$

**Conjecture 5.19.** *Let  $n \in \mathbb{Z}^+$  with  $n \equiv 0, 1, 5 \pmod{7}$ . Then*

$$t(1, 2, 2, 7; n) = 2N(1, 2, 2, 7; 2n + 3).$$

**Conjecture 5.20.** *Let  $n \in \mathbb{Z}^+$  with  $n \equiv 0, 2, 3 \pmod{7}$ . Then*

$$t(1, 1, 1, 7; n) = \frac{2}{5}N(1, 1, 1, 7; 8n + 10).$$

**Conjecture 5.21.** *Let  $n \in \mathbb{Z}^+$ . Then*

$$t(1, 1, 1, 33; n) = 2N(1, 1, 1, 33; 2n + 9) \quad \text{for } n \equiv 2, 4, 5, 6, 10 \pmod{11},$$

$$t(1, 1, 9, 33; n) = 2N(1, 1, 9, 33; 2n + 11) \quad \text{for } n \equiv 1, 3, 4, 5, 9 \pmod{11},$$

$$t(1, 9, 9, 33; n) = 2N(1, 9, 9, 33; 2n + 13) \quad \text{for } n \equiv 0, 2, 3, 4, 8 \pmod{11}.$$

**Remark 5.1** From [S3, Theorem 4.2] we know that if  $a, b, c, d, n \in \mathbb{Z}^+$  and  $a \equiv b \equiv c \equiv d \equiv \pm 1 \pmod{4}$ , then

$$t(a, b, c, d; n) = N(a, b, c, d; 8n + a + b + c + d) - N(a, b, c, d; 2n + (a + b + c + d)/4).$$

## References

- [ACH] C. Adiga, S. Cooper and J. H. Han, *A general relation between sums of squares and sums of triangular numbers*, Int. J. Number Theory **1**(2005), 175-182.
- [BCH] N. D. Baruah, S. Cooper and M. Hirschhorn, *Sums of squares and sums of triangular numbers induced by partitions of 8*, Int. J. Number Theory **4**(2008), 525-538.
- [Be] B.C. Berndt, *Ramanujan's Notebooks*, Part III, Springer, New York, 1991.
- [C] S. Cooper, *On the number of representations of integers by certain quadratic forms, II*, J. Combin. Number Theory **1**(2009), 153-182.
- [D] L.E. Dickson, *History of the Theory of Numbers*, Vol. II, Carnegie Institute of Washington, Washington D.C., 1923. Reprinted by AMS Chelsea, 1999.
- [S1] Z.H. Sun, *Some relations between  $t(a, b, c, d; n)$  and  $N(a, b, c, d; n)$* , Acta Arith. **175**(2016), 269-289.
- [S2] Z.H. Sun, *Ramanujan's theta functions and sums of triangular numbers*, Int. J. Number Theory **15**(2019), 969-989.
- [S3] Z.H. Sun, *The number of representations of  $n$  as a linear combination of triangular numbers*, Int. J. Number Theory **15**(2019), 1191-1218.
- [XZ] E.X.W. Xia and Y. Zhang, *Proofs of some conjectures of Sun on the relations between sums of squares and sums of triangular numbers*, Int. J. Number Theory **15**(2019), 189-212.
- [Y] X.M. Yao, *The relations between  $N(a, b, c, d; n)$  and  $t(a, b, c, d; n)$  and  $(p, k)$ -parametrization of theta functions*, J. Math. Anal. Appl. **453**(2017), 125-143.
- [WS1] M. Wang and Z.H. Sun, *On the number of representations of  $n$  as a linear combination of four triangular numbers*, Int. J. Number Theory **12**(2016), 1641-1662.
- [WS2] M. Wang and Z.H. Sun, *On the number of representations of  $n$  as a linear combination of four triangular numbers II*, Int. J. Number Theory **13**(2017), 593-617.
- [W] K.S. Williams, *Number Theory in the Spirit of Liouville*, Cambridge Univ. Press, New York, 2011.